Decomposable Neuro Symbolic Regression

Abstract

Symbolic regression (SR) models complex systems by discovering mathematical expressions that capture underlying relationships in observed data. However, most SR methods prioritize minimizing prediction error over identifying the governing equations, often producing overly complex or inaccurate expressions. To address this, we present a decomposable SR method that generates interpretable multivariate expressions leveraging transformer models, genetic algorithms (GAs), and genetic programming (GP). In particular, our explainable SR method distills a trained "opaque" regression model into mathematical expressions that serve as explanations of its computed function. Our method employs a Multi-Set Transformer to generate multiple univariate symbolic skeletons that characterize how each variable influences the opaque model's response. We then evaluate the generated skeletons' performance using a GA-based approach to select a subset of high-quality candidates before incrementally merging them via a GP-based cascade procedure that preserves their original skeleton structure. The final multivariate skeletons undergo coefficient optimization via a GA. We evaluated our method on problems with controlled and varying degrees of noise, demonstrating lower or comparable interpolation and extrapolation errors compared to two GP-based and three neural SR methods. Unlike these methods, our approach consistently learned expressions that matched the original mathematical structure.

1 Introduction

Deep learning-based systems have achieved remarkable success across various domains due to their ability to model complex, nonlinear functions. However, these systems are often described as "opaque" models[1] in the literature, emphasizing the high complexity of the functions they learn and their many required parameters. Their lack of interpretability and traceability poses challenges in applications where understanding the underlying mechanisms is crucial (Linardatos et al., 2021).

In scientific research, particularly in the physical sciences, interpretability is essential for uncovering governing equations that describe observed phenomena (Camps-Valls et al., 2023; Lee & Kumar, 2023). Data-driven methods play a crucial role in this process by identifying patterns and relationships directly from empirical data. Symbolic regression (SR) emerges as a powerful technique, offering an alternative to opaque models for discovering interpretable mathematical representations from data (Filho et al., 2020). Unlike conventional regression techniques that assume predefined model structures, SR searches the space of mathematical expressions to identify compact and accurate equations describing the observed data (Kronberger et al., 2024).

Most existing SR approaches focus primarily on minimizing prediction error rather than extracting the governing equations (Bertschinger et al., 2024). This often leads to expressions that fit the data well but are overly complex and difficult to interpret (La Cava et al., 2021). Thus, recent neural SR techniques (Biggio et al., 2021; Kamienny et al., 2022; Bertschinger et al., 2024) leverage transformer-based models to generate symbolic skeletons or full mathematical expressions. However, they process all variables simultaneously and often fail to capture the functional form between each variable and the system's response correctly [Ref. 1].[2]

To tackle these limitations, we introduce a novel multivariate SR method called **S**ymbolic **R**egression using **T**ransformers, **G**enetic **A**lgorithms, and genetic **P**rogramming (SeTGAP), illustrated in Fig. 1. SeTGAP is

---

[1]This term is preferred over "black-box": acm.org/diversity-inclusion/words-matter
[2]These references are hidden for double-blind review purposes.

Figure 1: An example of the symbolic discovery process using SeTGAP.

an explainable SR approach that extracts mathematical expressions that explain a trained opaque regression model's learned function. It employs a Multi-Set Transformer model to generate multiple univariate symbolic skeletons, characterizing the functional relationships between each independent variable and the opaque model's response. A genetic algorithm (GA)-based selection process then filters out low-quality skeletons, retaining only the most informative ones. Then, they are merged through an incremental genetic programming (GP)-based procedure, which ensures that the merged expressions remain interpretable and aligned with the original structures. Finally, a GA refines the numerical coefficients of the resulting multivariate expression. Our experiments demonstrate that SeTGAP consistently recovers the correct mathematical form of the underlying equations, unlike existing GP-based and neural SR methods, which often fail to do so. In addition, SeTGAP achieves lower or comparable interpolation and extrapolation error rates across various problem settings, including those with controlled and varying noise levels.

2 Related Work

SR constitutes an NP-hard problem that becomes increasingly complex as the number of observations, operators, and variables increases (Virgolin & Pissis, 2022). As such, brute-force approaches become infeasible. Most SR approaches are based on GP (Kronberger et al., 2024). They evolve candidate solutions, or programs, to identify suitable functional forms and optimize their parameters. This process involves applying genetic operators such as mutation, crossover, and selection to populations of expressions (Cranmer, 2023; Makke & Chawla, 2022; Orzechowski et al., 2018; Schmidt & Lipson, 2009). A candidate's fitness is evaluated based on its ability to minimize prediction error, guiding the evolution toward more accurate models.

GP's flexibility in generating programs of varying lengths allows it to explore a vast search space. One significant drawback is called "code bloat," the uncontrolled growth of program size during evolution. This phenomenon leads to increasingly complex solutions, inflating computational costs and reducing model interpretability (Dignum & Poli, 2007; Poli et al., 2007). The redundancy in the search space further exacerbates this issue, as many solutions represent nearly equivalent functions within the studied domain but differ syntactically (Ebner, 1999; Amir Haeri et al., 2017). This redundancy increases the difficulty of identifying optimal solutions and contributes to GP's bias toward generating larger expressions (Langdon et al., 1999).

Another limitation is that fitness evaluation typically focuses on the overall output of a program, neglecting evaluation of the intermediate components that contribute to the final result. To address this, Arnaldo et al. (2014) proposed to optimize the generated programs before the selection step by optimizing the fitness contributions of its subexpressions. Conversely, Jiang & Xue (2023) proposed Control Variable Genetic Programming (CVGP), which introduces independent variables into the analysis sequentially through a series of controlled experiments. In each stage, a GP learns simplified expressions by holding all but one variable constant, progressively refining candidate equations as more variables are incorporated. This iterative procedure enables CVGP to recover complex symbolic expressions more reliably than end-to-end approaches.

GP has been combined with deep learning techniques for SR. Mundhenk et al. (2021) introduced a recurrent neural network (RNN) to generate initial GP populations. The best-performing individuals are then used to retrain the RNN. Similarly, Petersen et al. (2021) presented a method that utilizes RNNs to generate mathematical expressions in a reinforcement learning framework. In addition, Udrescu & Tegmark (2020) proposed a recursive algorithm called AI Feynmann that uses NNs to identify properties such as symmetry, separability, and compositionally, which are exploited to define sub-problems that are easier to solve.

Several methods have proposed training NNs and pruning irrelevant parts of them until a simple equation can be distilled from the network weights (Martius & Lampert, 2016; Sahoo et al., 2018; Tsoi et al., 2025; Werner et al., 2021). One of their main limitations is that they do not leverage past experiences, and each problem is learned from scratch. Transformer-based methods have been proposed recently as an alternative. Biggio et al. (2021) presented a Set Transformer-based model (Lee et al., 2019) and pre-trained it on a synthetic dataset of multivariate equations to predict symbolic skeletons from input-output data pairs. During inference, the observed data is used to predict the equation's skeleton using the pre-trained transformer. Next, the constants within the skeleton are optimized using the BFGS algorithm (Fletcher, 1987).

Alternatively, Kamienny et al. (2022) proposed an end-to-end (E2E) transformer model that directly predicts complete mathematical expressions, bypassing skeleton prediction. Learned constants can be fine-tuned using a non-convex optimizer such as BFGS. Shojaee et al. (2023) introduced Transformer-based Planning for Symbolic Regression (TPSR), a decoding strategy that uses Monte Carlo Tree Search (MCTS) to guide equation generation via lookahead planning. TPSR is model-agnostic and works with any pre-trained transformer-based SR model, enabling optimization for non-differentiable objectives like reward and complexity. Their experiments show that applying TPSR on the E2E model boosts performance on benchmark datasets.

In addition, Bertschinger et al. (2024) proposed an SR neuro-evolution approach that trains a population of transformer models using two objective functions: prediction error and symbolic loss. Finally, in [Ref. 1][2], we noted that SR methods analyzing all variables simultaneously often fail to identify the functional relationships between each variable and the system's response. To address this, we introduced the Multi-Set Transformer, trained on synthetic data to identify symbolic skeletons shared across multiple input-response sets. Then, we proposed a *post-hoc* explainability method that extracts univariate skeletons, providing interpretable approximations of the function learned by an opaque regression model.

3 Deep Evolutionary Symbolic Regression

We consider a system with response $y \in \mathbb{R}$ and t explanatory variables $\mathbf{x} = \{x_1, \ldots, x_t\}$. We assume that its underlying function $f(\mathbf{x}) = f(x_1, \ldots, x_t)$ can be constructed using a finite number of unary and binary operations. The response is expressed as $y = f(\mathbf{x}) + \varepsilon_a$, where ε_a represents the error term due to aleatoric uncertainty. Below, we define the SR problem formally and describe our SeTGAP methodology.

3.1 Problem Definition

Let us define our SR problem formally:

Definition 1. *Given a dataset $(\mathbf{X}, \mathbf{y})$, with inputs $\mathbf{X} = \{\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_{N_R}\} \subseteq \mathbb{R}^t$ and target responses $\mathbf{y} = \{y_1, y_2, \ldots, y_{N_R}\} \subseteq \mathbb{R}$, the SR problem seeks to discover a function $\tilde{f} : \mathbb{R}^t \to \mathbb{R}$ that approximates the unknown underlying function $f(\mathbf{x})$. The function $\tilde{f}$ is represented as a composition of a finite number of unary and binary operators, such that $\tilde{f}(\mathbf{x}) \approx f(\mathbf{x})$ while capturing its functional structure and behavior.*

Thus, $\tilde{f}(\mathbf{x})$ can be expressed as $\tilde{f}(\mathbf{x}) = f(\mathbf{x}) + \varepsilon_a + \varepsilon_e$, where ε_e is the error due to epistemic uncertainty, which is attributable to a lack of knowledge about f and can be reduced by acquiring additional information and improving the predictive model. In this context, solving the SR problem implies minimizing ε_e by selecting a suitable representation for $\tilde{f}(\mathbf{x})$ and identifying an optimal set of parameters $\boldsymbol{\theta}_{\tilde{f}}$ for such representation.

3.2 Univariate Symbolic Skeleton Prediction

We deviate from SR approaches that prioritize the minimization of the prediction error of the learned functions. We argue that a correctly identified functional form $\tilde{f}$ inherently leads to a low estimation error

and emphasizes interpretability and faithfulness to the underlying system's governing principles. Existing works on Symbolic Skeleton Prediction (SSP) (Bertschinger et al., 2024; Biggio et al., 2021; Petersen et al., 2021; Valipour et al., 2022) attempt to address this issue by generating multivariate symbolic skeletons that describe the behavior of f, with numerical coefficients subsequently optimized to minimize prediction error.

Given a mathematical expression, its symbolic skeleton is an expression that replaces the numerical constants with placeholders. For example, if $f(\mathbf{x}) = 5 \log x_1 (\sin(x_2^2) + 1) - 4$, its skeleton is expressed as $\mathbf{e}(\mathbf{x}) = \kappa(f(\mathbf{x})) = c_1 \log x_1 (\sin(x_2^2) + c_2) + c_3$, where $\kappa(\cdot)$ represents the skeleton function and c_i are placeholders. In prior work [Ref. 1][2], we showed that SSP methods struggle to identify the correct functional form of all variables in a system. Thus, we introduced a univariate skeleton prediction method that produces univariate skeletons explaining the relationships between each variable and the system's response. Following the previous example, its univariate skeletons are: $\mathbf{e}(x_1) = \kappa(f(\mathbf{x}); x_1) = c_4(\log x_1) + c_5$, and $\mathbf{e}(x_2) = \kappa(f(\mathbf{x}); x_2) = c_6 \sin(x_2^2) + c_7$. Here, $\kappa(\cdot; x_v)$ considers the remaining variables $\mathbf{x} \setminus x_v$ irrelevant when describing the functional form between x_v ($v \in [1, \ldots, t]$) and the system's response. In this case, the placeholders c_i may represent numeric constants or functions of other variables.

This method uses a trained regression model $\hat{f}$ (e.g., a neural network) that approximates the underlying function f by capturing the association between $\mathbf{X}$ and $\mathbf{y}$. Then, each variable of the system, x_v, is analyzed separately. To do this, N_s artificial sets of input–response pairs $\{(\tilde{\mathbf{X}}^{(1)}, \tilde{\mathbf{y}}^{(1)}), \ldots, (\tilde{\mathbf{X}}^{(N_s)}, \tilde{\mathbf{y}}^{(N_s)})\}$ are generated. To isolate the influence of variable x_v, the set $\tilde{\mathbf{X}}^{(s)}$ ($s \in [1, \ldots, N_s]$) is constructed such that the variable x_v (i.e., the v-th column $\tilde{\mathbf{X}}_v^{(s)}$) is allowed to vary while the other variables are fixed to random values. Thus, $\tilde{\mathbf{y}}^{(s)}$ denotes the estimated response of inputs $\tilde{\mathbf{X}}^{(s)}$ using the trained model $\hat{f}$ as $\tilde{\mathbf{y}}^{(s)} = \hat{f}(\tilde{\mathbf{X}}^{(s)})$.

Fixing the columns corresponding to the variables $\mathbf{x} \setminus x_v$ ensures that $\tilde{\mathbf{y}}^{(s)}$ depends only on the column $\tilde{\mathbf{X}}_v^{(s)}$. However, this process may project the function into a space where its functional form becomes less recognizable. To address this, the influence of x_v on the system's response is analyzed using N_s sets of input–response pairs, each reflecting a different effect of the variables $\mathbf{x} \setminus \{x_v\}$. As such, each set $(\tilde{\mathbf{X}}^{(s)}, \tilde{\mathbf{y}}^{(s)})$ is generated independently by fixing $\mathbf{x} \setminus \{x_v\}$ at different values. The relationship between each $\tilde{\mathbf{X}}_v^{(s)}$ and $\tilde{\mathbf{y}}^{(s)}$ can be described by a univariate function $f_v^{(s)}$. Note that functions $f_v^{(1)}, \ldots, f_v^{(N_S)}$ have been derived from the same function $f(\mathbf{x})$ and should share the same symbolic skeleton $\mathbf{e}(x_v)$, which is unknown. The task of predicting a skeleton $\hat{\mathbf{e}}(x_v)$ that describes the shared function form of input $\tilde{\mathbf{D}}_v = \{\tilde{\mathbf{D}}_v^{(1)}, \ldots, \tilde{\mathbf{D}}_v^{(N_s)}\}$ (i.e., $\hat{\mathbf{e}}(x_v) \approx \mathbf{e}(x_v)$), s.t. $\tilde{\mathbf{D}}_v^{(s)} = (\tilde{\mathbf{X}}_v^{(s)}, \tilde{\mathbf{y}}^{(s)})$, is known as multi-set symbolic skeleton prediction (MSSP).

The MSSP problem has been addressed in [Ref. 1][2] by designing a Multi-Set Transformer. The model's function and parameters are denoted by $g(\cdot)$ and Θ, respectively. It was trained on a large dataset of synthetically generated MSSP problems to produce accurate estimated skeletons. Given an input collection $\tilde{\mathbf{D}}_v$, the estimated skeleton obtained for variable x_v is computed as $\tilde{\mathbf{e}}(x_v) = g(\tilde{\mathbf{D}}_v, \Theta)$.

Unlike previous work, we generate up to n_{cand} distinct candidate skeletons, as described in Algorithm 1. We generate a $\tilde{\mathbf{D}}_v$ collection and feed it into g to obtain n_B skeletons using a diverse beam search (DBS) strategy (Vijayakumar et al., 2018) to promote variability among the n_B generated skeletons. This process is repeated n_{cand} times, yielding a total of $n_{\text{cand}} n_B$ skeletons. Each repetition yields a new $\tilde{\mathbf{D}}_v$ collection using different combinations of fixed values of $\mathbf{x} \setminus \{x_v\}$, increasing input diversity and potentially leading to different skeletons. Then, we discard identical or mathematically equivalent skeletons. For example, we consider the skeleton $c_1 \sin(c_2 x_v + c_3)$ is equivalent to $c_4 \cos(c_5 x_v + c_6)$ if $c_1 = c_4$, $c_2 = c_5$, and $c_3 = c_6 - \pi/2$.

If the generated skeleton list, $\texttt{genSks}_v$, exceeds n_{cand} elements, we evaluate their performance and select the top n_{cand} candidates. To do this, an additional collection $\tilde{\mathbf{D}}_v$ is generated. Since a skeleton expression for variable $\mathbf{x}_v$ is expected to describe the functional form of all sets in $\tilde{\mathbf{D}}_v$, we choose a random set $\tilde{\mathbf{D}}_v^{(\text{test})} = (\tilde{\mathbf{X}}_v^{(\text{test})}, \tilde{\mathbf{y}}^{(\text{test})}) \subset \tilde{\mathbf{D}}_v$ and use it to fit the coefficients of each skeleton $\hat{\mathbf{e}}_k(x_v)$, where $k \in \{1, \ldots, |\texttt{genSks}_v|\}$. The coefficient fitting problem is described as follows. Let $f_{est}(x_v) = \texttt{setConstants}(\hat{\mathbf{e}}_k(x_v), \mathbf{c})$ be the function obtained when replacing the n_c coefficients in $\hat{\mathbf{e}}_k(x_v)$ with the numerical values in a given set $\mathbf{c} \in \mathbb{R}^{n_c}$. Then, the objective is to find an optimal set $\mathbf{c}^*$ that maximizes the following Pearson correlation:

$$\mathbf{c}^* = \underset{\mathbf{c}}{\arg\max} \, \text{corr}(f_{est}(\tilde{\mathbf{X}}_v^{(\text{test})}), \tilde{\mathbf{y}}^{(\text{test})}).$$

Algorithm 1 Univariate Skeleton Generation

1: **function** GENERATEUNIVSKS($v, n, N_s, \hat{f}, g, n_{\text{cand}}, n_B$)
2:     $\text{genSks}_v \leftarrow []$
3:     **for** each $i \in (1, n_{\text{cand}})$ **do**
4:         $\tilde{\mathbf{D}}_v \leftarrow \text{generateCollection}(v, n, N_s, \hat{f})$
5:         $\text{genSks}_v.\text{append}(g(\tilde{\mathbf{D}}_v, \Theta; n_B))$
6:     $\text{genSks}_v \leftarrow \text{removeDuplicates}(\text{genSks}_v)$ $\triangleright \text{genSks}_v = \{\hat{\mathbf{e}}_1(x_v), \ldots, \hat{\mathbf{e}}_{|\text{genSks}_v|}(x_v)\}$
7:     $\tilde{\mathbf{D}}_v^{(\text{test})} \leftarrow \text{generateCollection}(v, n, N_s, \hat{f})$
8:     $\text{corrVals}_v \leftarrow \text{zeros}(|\text{genSks}_v|)$
9:     **for** each $k \in (1, n_{\text{cand}})$ **do**
10:         $\text{corrVals}_v[k] \leftarrow \text{fitCoefficients}(\hat{\mathbf{e}}_k(x_v), \tilde{\mathbf{D}}_v^{(\text{test})})$
11:     $\text{genSks}_v \leftarrow \text{sortSkeletons}(\text{genSks}_v, \text{corrVals}_v)$
12:     **if** $|\text{genSks}_v| > n_{\text{cand}}$ **then**
13:         $\text{genSks}_v, \text{corrVals}_v \leftarrow \text{genSks}_v[1 : n_{\text{cand}}], \text{corrVals}_v[1 : n_{\text{cand}}]$
14:     **return** $\text{genSks}_v, \text{corrVals}_v$

Note that in generating $\tilde{\mathbf{D}}_v$, we fixed the values of $\mathbf{x} \setminus x_v$, so we can assume that all coefficients in $\mathbf{c}$ are numerical values. However, the learned coefficients are then discarded, as they serve only to evaluate the fit of a univariate skeleton to the data. We use the Pearson correlation between $f_{est}(\tilde{\mathbf{X}}_v^{(\text{test})})$ and the estimated response $\tilde{\mathbf{y}}^{(s)}$ instead of mean squared error (MSE) to evaluate the functional suitability of skeleton $\hat{\mathbf{e}}_k(x_v)$. This choice is motivated by the fact that we aim to assess whether the candidate skeleton captures the functional form regardless of scale or vertical shift. Unlike MSE, the correlation coefficient is invariant to linear transformations, thus allowing us to compare skeletons without explicitly optimizing scale and offset parameters, which would otherwise increase computational cost.

This problem is solved using a genetic algorithm (GA) (Holland, 1992). The individuals of our GA are arrays of n_c elements that represent potential $\mathbf{c}$ sets. Then the optimization process is carried out by function $\text{fitCoefficients}(\hat{\mathbf{e}}_k(x_v), \tilde{\mathbf{D}}_v^{(\text{test})})$. This optimization process is repeated for all system variables to derive their univariate skeleton expressions with respect to the system's response.

3.3 Merging Univariate Symbolic Skeletons

The next step is to merge the univariate skeleton candidates to produce multivariate expressions. This process is carried out in a cascade fashion until a final expression incorporating all variables is formed.

3.3.1 Merging Skeleton Expressions

Given two skeleton expressions, multiple mathematically valid ways to combine them may exist. Here, we explore how to generate such combinations. We start with the following proposition:

Proposition 1. *Let $f(\mathbf{x})$ be a scalar-valued function defined by a finite composition of real-valued unary and binary operators applied to scalar sub-expressions. Then, $f(\mathbf{x})$ can always be expressed as: $f(\mathbf{x}) = c_0 + \sum_i c_{i,j} \prod_j \nu_{i,j}(T_{i,j}(\mathbf{x}))$, where $c_0, c_{i,j} \in \mathbb{R}$, $\nu_{i,j}$ is a unary operator (including the identity function $I(f(\mathbf{x})) = f(\mathbf{x})$), and $T_{i,j}(\mathbf{x})$ is a sub-expression. Moreover, each $T_{i,j}(\mathbf{x})$ can be recursively decomposed in the same structure as f, continuing until the decomposition reduces to variables or constants.*

Proposition 1 is proved in Appendix A. Then, let $\mathbf{e}_1(\mathbf{x}_S)$ and $\mathbf{e}_2(x_q)$ be two candidate skeletons to be merged. Here, S is an index set, $S \subset \{1, \ldots, t\}$, specifying the variables that $\mathbf{e}_1$ depends on; i.e., $\mathbf{x}_S = \{x_r | r \in S\}$. In contrast, x_q is a variable distinct from those in $\mathbf{x}_S$ ($q \notin S$). Following from Proposition 1, the skeletons can be expressed as $\mathbf{e}_1(\mathbf{x}_S) = c_0 + \sum_i c_{i,j} \prod_j \nu_{i,j}(T_{i,j}(\mathbf{x}_S))$ and $\mathbf{e}_2(x_q) = c_0' + \sum_i c_{i',j'}' \prod_{j'} \nu_{i',j'}'(T_{i',j'}'(x_q))$. Below, we explain how the subtrees of both expressions can be merged recursively.

The key idea is that a constant placeholder in $\mathbf{e}_1(\mathbf{x}_S)$ may be replaced by a subtree of $\mathbf{e}_2(x_q)$, and vice versa. Given two skeletons, $\mathbf{e}_1(\mathbf{x}_S) = c_1 T_1(\mathbf{x}_S)$ and $\mathbf{e}_2(x_q) = c_2 T_2(x_q)$, a straightforward way to merge them is by recognizing that part of c_1 may be a function of x_q, while part of c_2 may be a function of $\mathbf{x}_S$. Thus, their combination, denoted by the operation $\bowtie$, is given by $\mathbf{e}_3(\mathbf{x}_S \cup x_q) = \mathbf{e}_1(\mathbf{x}_S) \bowtie \mathbf{e}_2(x_q) = c_3(c_4 + T_1(\mathbf{x}_S))(c_5 + T_2(x_q))$. Expanding this expression conforms to the expected functional form stated in Proposition 1. Note that the skeletons with respect to the corresponding initial variable sets remain unchanged; that is, $\kappa(\mathbf{e}_1(\mathbf{x}_S); \mathbf{x}_S) = \kappa(\mathbf{e}_3(\mathbf{x}_S \cup x_q); \mathbf{x}_S) = c_1 T_1(\mathbf{x}_S)$ and $\kappa(\mathbf{e}_2(x_q); x_q) = \kappa(\mathbf{e}_3(\mathbf{x}_S \cup x_q); x_q) = c_q T_2(x_q)$.

Figure 2: Example of a selected subtree of $\mathbf{e}_2(x_q)$ within a sum merging with one or more subtrees from $\mathbf{e}_1(\mathbf{x}_S)$, illustrating four out of the nine possible cases.

Applying the same principle to a more general case in which $\mathbf{e}_1(\mathbf{x}_S) = c_1 + c_2 \prod_j T_{1,j}(\mathbf{x}_S)$ and $\mathbf{e}_2(x_q) = c_3 + c_4 \prod_{j'} T_{2,j'}(x_q)$, we obtain $\mathbf{e}_3(\mathbf{x}_S \cup x_q) = \mathbf{e}_1(\mathbf{x}_S) \bowtie \mathbf{e}_2(x_q) = c_5 + c_6 \prod_j (c_{7,j} + T_{1,j}(\mathbf{x}_S)) \prod_{j'} (c_{8,j'} + T_{2,j'}(\mathbf{x}_S))$. Nevertheless, more combinations are possible if the candidate skeletons share functions with compatible mathematical structures. For example, if $\mathbf{e}_1(x_1, x_2) = c_1 \sin(c_2 \, x_1 x_2 + c_3)$ and $\mathbf{e}_2(x_3) = c_4 \sin(c_5 \, x_3 + c_6)$, we could obtain $\mathbf{e}_3(x_1, x_2, x_3) = \mathbf{e}_1(x_1, x_2) \bowtie \mathbf{e}_2(x_3) = c_7(c_8 + \sin(c_9 \, x_1 x_2 + c_{11}))(c_{12} + \sin(c_{13} \, x_3 + c_{14}))$, as shown before, but also $\mathbf{e}_3(x_1, x_2, x_3) = c_{15} \sin(c_{16} \, x_1 x_2 + c_{17} x_3)$ and $\mathbf{e}_3(x_1, x_2, x_3) = c_{18} \sin(c_{19} \, x_1 x_2 x_3 + c_{20})$, which yield to the same skeletons with respect to the initial variable sets.

Algorithm 2 shows our recursive merging procedure, $\texttt{merge}(\texttt{ex}_1, \texttt{ex}_2)$, which takes as inputs two skeleton expressions $\texttt{ex}_1$ and $\texttt{ex}_2$. The lists of subtrees of both expressions are retrieved by function $\texttt{getSubtreesLists}(\texttt{ex}_1, \texttt{ex}_2)$, returning the one with fewer subtrees first. We shuffle both lists to introduce variability in the process. If $\texttt{exShort}$ lacks a constant term, we add one to conform to the structure in Proposition 1 and ensure that is the last element. If both expressions represent sums, the algorithm iterates over the elements of $\texttt{exShort}$, attempting to merge each with compatible subtrees from $\texttt{exLong}$; i.e., subtrees that share the same unary or binary operator. The compatibility check is performed by $\texttt{findComp}$, and a subset of matching subtrees is randomly selected for merging, as depicted in Fig 2. If only one compatible subtree is selected, it is merged recursively with $\texttt{exShort}[i]$. However, if multiple subtrees are selected, the only way to maintain the original skeleton structure is to treat their sum as a single, indivisible entity, effectively a constant term with respect to x_q, which is then multiplied by $\texttt{exShort}[i]$ (Line 16). Then, the constant term that appears at the end of $\texttt{exShort}$ absorbs any remaining terms from $\texttt{exLong}$.

If neither expression is a sum or a product but they share the same function, the function remains unchanged, and their inner arguments are merged recursively. For products, if $\texttt{exShort}$ consists only of symbols, the expressions are combined as explained before (Line 23). Otherwise, it iterates through the terms, merging them based on compatibility, as was done for the sum case. Unlike in the sum case, each subtree $\texttt{exShort}[i]$ can merge with only one subtree from $\texttt{exLong}$ at a time, with a 0.5 probability of merging to introduce variability. Finally, the function returns the merged expression preserving their initial structures.

3.3.2 Selecting Combined Skeleton Expressions

Algorithm 2 generates a single combination of skeletons. Here, we extend this process to construct a population of candidate skeletons and select the top-performing one. Thus, we employ an evolutionary strategy to assess the performance of each skeleton combination. The population of skeletons, $\texttt{candSks}$, is constructed by repeatedly applying Algorithm 2. Since the merging process is stochastic, each application of the algorithm may yield a different skeleton combination. The process continues until a population size $P_{\max}$ is reached or a patience criterion is met. As such, if no new valid skeleton is found after a certain number of attempts, the generation process halts, assuming that all viable combinations have been explored.

We aim to identify the most promising combination in $\texttt{candSks}$ by evaluating how well each skeleton can fit the test data. Algorithm 3 provides a high-level overview of this process. We generate test data $\tilde{\mathbf{D}}^{(\text{test})} = (\tilde{\mathbf{X}}^{(\text{test})}, \tilde{\mathbf{y}}^{(\text{test})})$. Unlike in Sec. 3.2, here the columns in $S' = \{S \cup q\}$ (i.e., the indices of variables present in the combined skeletons) are allowed to vary, while all other variables are fixed to random values. The set

---

Algorithm 2 Recursive Skeleton Merging

1: **function** MERGE(ex_1, ex_2)
2: exShort, exLong $\leftarrow$ **getSubtreesLists**(ex_1, ex_2) ▷ Return lists of subtrees in sum
3: exShort.**shuffle**(), exLong.**shuffle**()
4: **if** isSum(ex_1) **and** isSum(ex_2) **then**
5:      **for** each $i \in (1, |\text{exShort}|)$ **do**
6:          **if** $i = |\text{exShort}|$ **and** $|\text{exLong}| > 0$ **then**
7:              $\text{exShort}[i] \leftarrow \text{exShort}[i] + \sum_j \text{exLong}[j]$
8:          **else**
9:              args $\leftarrow$ **findComp**(exLong, exShort[i]) ▷ Find exLong args compatible w/exShort[i]
10:              selectedArgs $\leftarrow$ **sample**(args, randInt(0, |args|))
11:              **if** $|\text{selectedArgs}| = 0$ **then continue**
12:              exLong.**remove**(selectedArgs)
13:              **if** $|\text{selectedArgs}| = 1$ **then**
14:                  $\text{exShort}[i] \leftarrow \textbf{merge}(\text{exShort}[i], \text{selectedArgs}[0])$
15:              **else**
16:                  $\text{exShort}[i] \leftarrow \text{exShort}[i] \sum_j \text{selectedArgs}[j]$
17:      mergedEx $\leftarrow \sum_i \text{exShort}[i]$
18: **else**
19:      **if** !isMult(ex_1) **then** ▷ If compatible, merge inner arguments
20:          mergedEx $\leftarrow \text{ex}_1.\textbf{func}(\textbf{merge}(\text{ex}_1.\textbf{args}, \text{ex}_2.\textbf{args}))$
21:      **else**
22:          **if** isAllSymbols(exShort) **then**
23:              mergedEx $\leftarrow c_1 \prod_i (c_{2,i} + \text{exShort}[i]) \prod_j (c_{3,j} + \text{exLong}[j])$
24:          **else**
25:              mergedEx $\leftarrow$ **null**
26:              **for** each $i \in (1, |\text{exShort}|)$ **do**
27:                  **if** $i = |\text{exShort}|$ **and** $|\text{exLong}| > 0$ **then**
28:                      mergedEx $\leftarrow \text{exShort}[i] \prod_{i'=0}^{i-1} (c_{1,i'} + \text{exShort}[i']) \prod_j (c_{2,j} + \text{exLong}[j])$
29:                  **else**
30:                      args $\leftarrow$ **findComp**(exLong, exShort[i])
31:                      **if** $|\text{args}| = 0$ **then continue**
32:                      selectedArg $\leftarrow$ **choice**(args)
33:                      **if** random$(0, 1) < 0.5$ **then continue**
34:                      exLong.**remove**(selectedArg)
35:                      $\text{exShort}[i] \leftarrow \textbf{merge}(\text{exShort}[i], \text{selectedArg})$
36:              **if** mergedEx is **null** **then**
37:                  mergedEx $\leftarrow \prod_i \text{exShort}[i]$
38: **return** mergedEx

Algorithm 3 Skeleton Combination with Genetic Programming

1: **function** SELECTCOMBINATION(candSks, rep, maxG, $\tilde{\mathbf{D}}_{S'}^{(\text{test})}$)
2: Pop $\leftarrow$ [ ]
3: **for** each $s \in (1, |\text{candSks}|)$ **do**
4:      exps $\leftarrow$ [ ]
5:      **for** $i = 1$ to rep **do** exps.**append**(**assignValues**(candSks[s]))
6:      Pop.**append**(exps)
7: **for** each gen $\in (1, \text{maxG})$ **do**
8:      fitnesses, bestFitnesperSk $\leftarrow$ **evalCorr**(Pop, $\tilde{\mathbf{D}}_{S'}^{(\text{test})}$)
9:      Pop $\leftarrow$ **evolveSks**(Pop, fitnesses)
10: **return** candSks[**argmax**(bestFitnesperSk)], **max**(bestFitnesperSk)

$\tilde{\mathbf{D}}_{S'}^{(\text{test})} = (\tilde{\mathbf{X}}_{S'}^{(\text{test})}, \tilde{\mathbf{y}}^{(\text{test})})$ is then used for evaluation. Each skeleton in `candSks` is replicated `rep` times, with randomly assigned coefficient values, to form an initial pool of candidate expressions `Pop`.

The population then undergoes an evolution process for `maxG` generations. In each iteration, the fitness of every expression in `Pop` is evaluated using `evalCorr`, which assesses the Pearson correlation value between $\tilde{\mathbf{y}}^{(\text{test})}$ and the output produced by the expression when evaluated on $\tilde{\mathbf{X}}^{(\text{test})}$. Then, `evolveSks` updates the population by selecting high-performing candidates and introducing variations through crossover and mutation. While coefficient values are subject to modification, the structure of each skeleton remains unchanged. This is ensured by restricting crossover to expressions derived from the same skeleton. Finally, the highest-fitness skeleton across all evolved instances is selected as the best candidate. This approach bears similarities to GP, with the distinction that the search for viable skeletons is decoupled from the evolutionary optimization process. Rather than evolving the expressions dynamically, we first enumerate all admissible skeleton combinations via Algorithm 2 and then apply a constrained form of genetic programming, where evolution occurs with the structure of all mathematical expressions fixed to the precomputed combined skeletons.

3.3.3 Cascade Merging

The construction of multivariate skeletons follows an incremental merging process in which univariate skeleton candidates are combined progressively. For each variable x_v Algorithm 1 generates up to n_{cand} candidate skeletons $\texttt{genSks}_v$ along with their corresponding correlation values $\texttt{corrVals}_v$. To determine the merging order, the variables are ranked according to the maximum correlation value obtained across their generated skeletons. Variables with higher correlation scores are prioritized in the merging order, as their generated skeletons exhibit stronger relationships with the data, reducing the risk of propagating structural uncertainty.

Let S denote the indices of the variables whose skeletons have already been merged. Initially, S contains only the index of the variable that corresponds to the skeleton with the highest correlation value. At each iteration, a new variable x_q is selected according to the ranking, and its univariate skeletons are merged with those corresponding to S. Specifically, Algorithms 2 and 3 are applied to combine each skeleton generated for $\mathbf{x}_S$ with each skeleton generated for x_q, producing at most n_{cand}^2 skeleton combinations. Since Algorithm 3 returns both the selected merged skeleton and its fitness, we apply a greedy selection strategy, retaining only the n_{cand} highest-performing skeletons at each step. This process repeats until skeletons incorporating all t variables have been generated, which serve as interpretations that describe the functional form of model $\hat{f}$.

We argue that our cascade approach ensures a structured integration of variables into the final multivariate skeleton. Directly constructing a multivariate skeleton from all univariate candidates would prioritize minimizing overall prediction error, potentially obscuring individual contributions and leading to overfitting. Instead, incorporating one variable at a time allows for a more interpretable learning process, where each newly added variable's effect is evaluated in the context of the previously analyzed ones.

It is worth pointing out that our cascade-fashion approach shares similarities with the CVGP method proposed by Jiang & Xue (2023). CVGP analyzes the t system variables in a fixed order: $x_1, x_2, \ldots, x_t$. First, multiple datasets (trials) are generated where only x_1 varies while others remain fixed. A GP algorithm extracts a skeleton involving x_1. Then, new trials are generated where x_1 and x_2 vary, and the previously discovered skeleton is fixed while GP extends it to include both variables. This process repeats until all variables are included. Our approach differs in both motivation and implementation. SeTGAP starts by independently identifying univariate skeletons for each variable, focusing on interpreting the individual functional relationships learned by the opaque model $\hat{f}$. Once fixed, it aims to discover mathematically valid ways to merge them into coherent multivariate expressions. In contrast, CVGP builds successive expressions atop a single skeleton for x_1. If this initial skeleton is poorly fitted, subsequent expressions tend to compensate for the misspecification rather than recover accurate functional relationships for the remaining variables.

3.3.4 Underlying Function Estimation

We utilize the N_e multivariate skeletons $\hat{\mathbf{e}}_1(\mathbf{x}), \ldots, \hat{\mathbf{e}}_{N_e}(\mathbf{x})$ produced by our cascade merging procedure, where $N_e \leq n_{\text{cand}}$. The goal is to construct functions $\tilde{f}_i(\mathbf{x})$ that approximate the underlying function $f(\mathbf{x})$ using the corresponding skeleton $\hat{\mathbf{e}}_i(\mathbf{x})$, $\forall i \in (1, \ldots, N_e)$. This defines a coefficient fitting problem similar to the one presented in Sec. 3.2. Unlike earlier sections, we minimize the MSE of the response calculated by evaluating $\tilde{f}_i(\mathbf{x})$ on the original dataset $(\mathbf{X}, \mathbf{y})$. This is feasible because $\hat{\mathbf{e}}(\mathbf{x})$ contains all system variables and there is no need to generate a synthetic set of estimated points using $\hat{f}$. Hence, the new coefficient fitting problem consists of finding an optimal set of coefficient values that minimizes the prediction MSE:

$$\mathbf{c}^* = \operatorname*{argmin}_{\mathbf{c}} \frac{1}{|\mathbf{X}|} \sum_{(\mathbf{x}_j, y_j) \in (\mathbf{X}, \mathbf{y})} \left(\texttt{setConstants}(\hat{\mathbf{e}}_i(\mathbf{x}_j), \mathbf{c}) - y_j \right)^2$$

such that $\tilde{f}_i(\mathbf{x}) = \texttt{setConstants}(\hat{\mathbf{e}}_i(\mathbf{x}), \mathbf{c}^*)$. This optimization is carried out using the GA-based function $\texttt{fitCoefficients}(\hat{\mathbf{e}}_i(\mathbf{x}), (\mathbf{X}, \mathbf{y}))$ introduced in Sec. 3.2 but modified to minimize MSE.

4 Experimental Results

We evaluated SeTGAP on 13 synthetic SR problems derived from equations inspired by previous work. Table 1 lists these equations along with their domain ranges. The datasets generated from these equations consisted of 10,000 points and each variable was sampled according to a uniform distribution.

Table 1: Equations used for experiments

Eq.	Underlying equation	Reference	Domain range
E1	$(3.0375x_0x_1 + 5.5\sin(9/4(x_0 - 2/3)(x_1 - 2/3)))/5$	Jin et al. (2020)	$[-5,5]^2$
E2	$5.5 + (1 - x_0/4)^2 + \sqrt{x_1 + 10}\sin(x_2/5)$	[Ref. 1][2]	$[-10,10]^2$
E3	$(1.5e^{1.5x_0} + 5\cos(3x_1))/10$	Jin et al. (2020)	$[-5,5]^2$
E4	$((1 - x_0)^2 + (1 - x_2)^2 + 100(x_1 - x_0^2)^2 + 100(x_3 - x_2^2)^2)/10000$	Rosenbrock-4D	$[-5,5]^4$
E5	$\sin(x_0 + x_1x_2) + \exp(1.2x_3)$	[Ref. 1][2]	$x_1 \in [-10,10], x_2 \in [-5,5],$ $x_3 \in [-5,5], x_4 \in [-3,3]$
E6	$\tanh(x_0/2) + |x_1|\cos(x_2^2/5)$	[Ref. 1][2]	$[-10,10]^3$
E7	$(1 - x_1^2)/(\sin(2\pi x_0) + 1.5)$	Werner et al. (2021)	$[-5,5]^2$
E8	$x_0^4/(x_0^4 + 1) + x_1^4/(x_1^4 + 1)$	Trujillo et al. (2016)	$[-5,5]^2$
E9	$\log(2x_1 + 1) - \log(4x_0^2 + 1)$	Trujillo et al. (2016)	$[0,5]^2$
E10	$\sin(x_0 \, e^{x_1})$	Bertschinger et al. (2023)	$x_1 \in [-2,2], x_2 \in [-4,4]$
E11	$x_0 \, \log(x_1^4)$	Bertschinger et al. (2023)	$[-5,5]^2$
E12	$1 + x_0 \sin(1/x_1)$	Bertschinger et al. (2023)	$[-10,10]^2$
E13	$\sqrt{x_0} \, \log(x_1^2)$	Bertschinger et al. (2023)	$x_1 \in [0,20], x_2 \in [-5,5]$

Details on the Multi-Set Transformer g architecture and training can be found in our previous work [Ref. 1][2]. In all cases, the generated collections $\tilde{\mathbf{D}}_v$ consist of N_S sets of $n = 3000$ elements. The regression models $\hat{f}$ were feedforward networks trained on 90% of the data and validated on the remaining 10%. They featured three hidden layers for problem E2, five for E1, E4, E5, and E7, and four for all other cases, with each layer containing 500 ReLU-activated nodes. The hyperparameters in Algorithm 1 include the beam size n_B and the number of univariate skeleton candidates n_{cand}. We set $n_B = 3$ and $n_{\mathrm{cand}} = 4$, as higher values did not yield more distinct skeletons across all problems. The GAs in Secs. 3.2 and 3.3 share the same configuration but differ in loss functions: the former maximizes Pearson correlation, while the latter minimizes MSE. The GA runs with a population of 500 individuals and terminates when the objective function change remains below 10^{-6} for 30 consecutive generations. It uses tournament selection, binomial crossover, and generational replacement. Algorithm 3 uses the number of instance expressions per candidate skeleton combination `rep` and the maximum number of generations `maxG`, with values set to `rep` = 150 and `maxG` = 300. In addition, the initial population `candSks` was generated with a maximum size $P_{max} = 5000$, though none of the cases reached this limit. This setup was chosen for its consistent effective optimization results across all problems.

For each problem, SeTGAP generates up to n_{cand} multivariate expressions, but for brevity, we report only the one with the lowest MSE. These results are compared against expressions obtained from two GP-based methods, PySR (Cranmer, 2023) and TaylorGP (He et al., 2022), and three neural SR approaches, NeSym-ReS (Biggio et al., 2021), E2E (Kamienny et al., 2022), and TPSR (Shojaee et al., 2023). The comparison is limited to neural SR methods with publicly available models, as training large transformer-based architectures is computationally prohibitive. NeSymReS could not be applied to E4 and E5 due to its limitation to systems with at most three variables. The only hyperparameters tuned were the beam size and the number of BFGS restarts, both initially set to 5 and 4, respectively. However, it was found that a beam size above 2 and more than 2 BFGS restarts were unnecessary. For E2E, we limited the number of generated candidates to a maximum of $K = 10$, as increasing this value provided no observable improvement. TPSR employed E2E as its backbone, and its regularization parameter λ, which controls the trade-off between accuracy and expression complexity, was set to 1, as smaller values led to unnecessarily complex expressions.

For GP-based methods, we set an iteration limit of 10,000, though all cases converged earlier. Population sizes of 100, 200, 500, and 1000 were tested, with no observed benefits beyond 500. Both methods were configured to use the same set of operators as those present in the vocabularies used by the neural SR methods. Although our approach may appear similar to CVGP, as discussed in Sec. 3.3.3, we did not include it in our experiments. This is because CVGP relies heavily on a data oracle and does not apply to a standard SR setting where a fixed initial dataset is available, making a fair comparison infeasible. Tables 2–4 present the learned expressions, with shaded cells indicating that the learned expression's functional form matches that of the underlying function. To ensure a fair comparison, the evaluation was repeated nine additional times, each with a newly generated dataset using a different random seed (Appendix B).

We evaluated the extrapolation capability of the learned expressions by testing them on an extended domain range. The original domain range, or interpolation range, for a variable x_v is denoted as $[x_v^\ell, x_v^u]$, while its extrapolation range is defined as $[2x_v^\ell, x_v^\ell[\, \cup \,]x_v^u, 2x_v^u]$. This evaluation was repeated for each of the 10 expressions learned by each method. We also evaluated the extrapolation capability of the opaque models

Table 2: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 1

Method	E1	E2	E3	E4
PySR	$0.61x_0x_1$	$0.41x_2 + ||x_0 - 3.51| - 1.95| + 4.11$	$0.34e^{x_0}|\sinh(0.47x_0)|$	$0.21x_0^2 - 0.18x_1 + 0.21x_2^2 - 0.18x_3 - 0.76$
TaylorGP	$0.64x_0x_1$	$-0.5x_0 + 0.001x_1 + 0.39x_2 + 8.62$	$0.23x_0e^{x_0}$	$0.29x_0^2$
NeSymReS	$0.59x_0x_1 + \cos(0.01(x_1 - x_0 - 0.08)^2)$	$-x_0 + 0.40x_2 + e^{e^{-0.001x_1}} + 5.88$	$9.10e^{0.72x_0}\cos(0.15x_1)$	—
E2E	$1.08(0.56x_0x_1 - 0.03x_0 + 0.02x_1 - \sin(0.01x_0^2 + 8.6x_0 + 0.45) - 0.01)$	$0.06x_0^2 - 0.51x_0 - 0.22x_1\cos(0.18x_2 + 1.43) - 0.01x_1 + 0.01x_2 - 3.25\cos(0.18x_2 + 1.43) + 6.56$	$0.14e^{1.52x_0} + 0.52\cos(3.45x_1 + 0.05) + 0.11$	$0.001|8.99(-0.88x_1 + (x_0 + 0.01)^2 + 0.62)^2 + 9.72(-x_3 + 0.98(x_2 + 0.01)^2 + 0.01)^2| + 0.0023$
TPSR	$0.146x_1(4.161x_0 - 0.071) - 0.036\sin(5.687x_1 - 1.848) - 0.002$	$\left(0.014 + \frac{1}{x_0 + 14.596}\right)(0.039x_0 + 3.247)\left(-0.059x_1 - 83.999 + \frac{0.061}{x_2}\right)(0.028x_2 - 0.409\arctan(0.14474x_2 + 0.0015) - 0.334)$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$0.175x_1 - 0.124x_3 + 0.3626 + (8.858 - 5.843x_1)(0.001x_1 + 0.073) + 0.0016\left(x_0 + 0.595(0.944x_0 - 1)^2\right)^3$
SeTGAP	$0.61x_0x_1 + 1.15\sin((2.24x_0 - 1.5)(x_1 - 0.68))$	$0.06x_0^2 - 0.5x_0 + (3.37\sqrt{0.1x_1 + 1} - 0.19)(\sin(0.2x_2) + 0.01) + 6.49$	$0.15e^{1.5x_0} + 0.5\sin(3x_1 - 4.71)$	$0.01x_0^4 - 0.02x_0^2x_1 - 0.001x_0 + 0.01x_1^2 + 0.01x_2^4 + 0.01x_3^2 - (0.02x_2^2 + 0.004)(x_3 - 0.11) - 0.02$

Table 3: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 1

Method	E5	E6	E7	E8
PySR	$e^{1.2x_3} + \sin(x_0 + x_1x_2)$	$\tanh(e^{x_2})$	$(0.56 - 0.59x_0^2)/(\sinh(\sinh(\tanh(e^{\sinh(\sin(6.28x_1))}))))$	$(\tanh(\cosh(x_0) - 1.04) + \tanh(\cosh(x_1) - 1.04))$
TaylorGP	$0.51e^{x_3}e^{\sin(0.87x_3)}$	$-\frac{\sin(0.34x_2^2)}{-\sqrt{|x_2|} + \sin(\sqrt{|x_2|})}$	$\sqrt{|x_1|} - x_1^2$	2
NeSymReS	—	$-0.39x_0 + x_1\sin(\frac{x_0}{x_1} - 0.001x_2)$	$\frac{0.12x_0 + x_1^2}{\cos(3.1(-0.02x_1 - 1)^2) - 0.31}$	$\cos(\sin(1.69x_0)/(x_0x_1)) + 0.71$
E2E	$e^{1.2x_3} - 0.91\cos((2.62x_0 + 0.15)(24.66x_1 + 1.24)) - 0.05$	$0.01x_1(-7.5\cos(15.41x_1 + 0.21) - 0.18) + 0.69\operatorname{atan}(0.75x_0 + 0.05) + 0.47$	$(-0.03x_1 - 0.03)(0.34x_1 - 0.35)(41.59(1 - 0.5\sin(6.74x_0 + 0.23))^2 + 40)$	$2.01 - 1.05e^{-0.06|x_0 2.73 - 0.14||0.59x_1 + 0.1|}$
TPSR	$1.032e^{-0.001x_2 + 1.191x_3} - 0.054 + 0.063\cos(-2.276x_0 + x_1 + 5.52)$	$(0.003 - 0.024(\cos(39.378(1 - 0.005x_2)^2 - 38.823) + 0.56)^2)(-3.997x_0 + 51.732x_2 + 16.608\cos(1.089x_1 - 0.479) - 135.856)$	$0.681 - 0.848(-x_1 - 0.134\cos(x_0 + 2.153) - 0.069)^2$	$2.002 - 1.474e^{-0.009|39.99x_0 - 0.28|(1.08x_1 + 0.03)|}$
SeTGAP	$0.999e^{1.2x_3} - \sin(x_0 + x_1x_2 + 9.42)$	$\cos(0.2x_2^2 + 0.05)|x_1| + \tanh(0.5x_0)$	$\frac{4.53 - 4.54x_1^2}{4.54\sin(6.28x_0 + 6.28) + 6.81}$	$2 - \frac{19.76}{19.31x_0^4 + 0.12x_0^3 + 0.42x_0^2 + 19.72} - \frac{5.33}{5.44x_1^4 - 0.09x_1^2 + 5.34}$

$\hat{f}$ trained for each problem. Each extrapolation set comprised 10,000 points sampled uniformly within this range. Table 5 shows the rounded mean and standard deviation of the extrapolation MSE across these runs. Bold entries indicate the method with the lowest mean error and a statistically significant difference, determined by Tukey's honestly significant difference test at the 0.05 significance level.

Finally, we tested SeTGAP under increasingly noisy conditions. We considered a normal error term, defined as $\varepsilon_a = \mathcal{N}(0, \sigma_a\sigma_\mathbf{y})$, and four noise levels: $\sigma_a = \{0, 0.01, 0.03, 0.05\}$. Here, $\sigma_\mathbf{y}$ denotes the standard deviation of the response variable so that the noise is scaled relative to the dispersion of each problem. The obtained interpolation and extrapolation MSE are shown in Table 6, where shaded cells indicate an incorrectly identified functional form. The corresponding learned expressions are provided in Appendix C.

5 Discussion

SeTGAP can be viewed as a *post-hoc* interpretability method, as it extracts mathematical expressions that align with the functional response learned by a given opaque regression model. Our decomposable approach learns and preserves functional relationships between input variables and the system's response, and increments them progressively, allowing for an interpretable evolution.. This prevents the resulting expressions from focusing solely on error minimization, encouraging alignment with the true functional form instead.

From Tables 2–4, we confirmed that SeTGAP successfully learned mathematical expressions equivalent to the underlying functions in Table 1 across all tested problems and all 10 iterations. In contrast, none of the competing methods correctly identified the underlying functions in more than seven out of the 13 problems in any of the 10 iterations. Notably, some methods only captured the functional form of the most influential variables; i.e., those contributing most to the response value. For instance, E2E recovered correctly the term $0.06x_0^2 - 0.51x_0 + 6.5$ for x_0 in E2 but failed for the other variables. E2E and TPSR were the only methods that produced expressions longer than reported, requiring simplification via a symbolic manipulation library[3].

---

[3]SymPy: `https://www.sympy.org/`

Table 4: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 1

Method	E9	E10	E11	E12	E13
PySR	$\log(\frac{x_1+0.5}{0.5+2x_0^2})$	$\sin(x_0 e^{x_1})$	$x_0\log(x_1^4)$	$\sin(\frac{x_0}{x_1/0.12})|x_0|+0.99$	$\sqrt{x_0}\log(x_1^2)$
TaylorGP	$\log(\frac{0.79}{|x_0|})-$ $2.36e^{-x_1}$	$x_0 e^{-\sqrt{|x_1|}}$	$4x_0\log(|x_1|)$	$(x_0+\sqrt{|x_1|}-$ $0.91)\sin(\frac{0.73}{x_1})$	$\sqrt{e^{\sqrt{x_0}}}\log(|x_1|)+$ $\log(|x_1|)+0.58$
NeSymReS	$1.12\log(|x_1/x_0|)-1.37$	$\sin(x_0 e^{x_1})$	$x_0\log(x_1^4)$	$(x_0+x_1)\sin(1/x_1)$	$0.31x_0+3.19\log(x_1^2)-3.25$
E2E	$2-0.60\log(13.36(0.004-$ $x_0)^2(1-0.13/$ $(-0.06x_1-0.02))^2+0.8)$	$-0.98\sin((0.06-2.86e^{1.03x_1})$ $(0.32x_0+0.002))-0.007$	$-0.74x_0(-5.62$ $\log(0.07|-6.94x_1+0.13|$ $+0.01)-3.74)$	$(0.79x_0-0.04)$ $\sin(4.5/(3.4x_1+0.08))$	$(-90.0+\frac{9}{0.12|3.4x_1+0.12|+0.04})$ $(0.09-0.1\log(0.17x_0+3.29))$
TPSR	$0.401x_1-0.025|8.656x_0-$ $120.665\arctan(-0.611x_0-$ $0.004)+0.098|+0.94$	$1.0\sin(1.002x_0 e^{x_1})$	$(0.022-0.601x_0)$ $(-0.223x_1^2-7.502-$ $\frac{36.162}{-4.305|x_1|-0.971})$	$0.001x_0(17.945x_1-$ $0.666+\frac{1.203}{x_1})-$ $0.008x_0+0.941$	$\left(-0.008+\frac{0.742}{5.792|1.369x_1+0.005|+1.035}\right)$ $\left(7.888-\frac{28.241}{x_0+4.024}\right)(x_1-0.995)$ $(1.462x_0+44.99)(0.112x_1+0.111)$
SeTGAP	$-\log(13.95x_0^2+3.48)+$ $|\log(8.32x_1+4.18)|-0.18$	$\sin\left(x_0 e^{0.999x_1}\right)$	$1.998x_0\log(x_1^2)$	$x_0\sin(1/x_1)+1$	$2\sqrt{x_0}\log|x_1|$

Table 5: Extrapolation MSE Comparison

Eq.	NN	PySR	TaylorGP	NeSymRes	E2E	TPSR	SeTGAP
E1	431.1 ± 0.39	0.37 ± 0.40	1.22 ± 1.31	3.37 ± 3.49	1.22 ± 0.13	0.65 ± 0.61	**0.12 ± 0.09**
E2	30.77 ± 0.37	98.99 ± 64.70	238.50 ± 75.63	8.59e+7 ± 1.91e+8	68.85 ± 68.22	1.38e+7 ± 4.04e+7	**0.08 ± 7.62e-2**
E3	2.14e+4 ± 594.9	2.07e+3 ± 6.19e+3	3.15e+4 ± 3.96e+3	4.65e+4 ± 9.21e+3	1.45e+5 ± 3.68e+5	9.42e+7 ± 2.83e+8	10.28 ± 17.89
E4	2869 ± 15.4	2.21e+5 ± 9.04e+4	6.62e+3 ± 1.06e+3	—	876.20 ± 2.32e+3	2.44e+5 ± 7.26e+5	**1.62 ± 1.89**
E5	9.29e+4 ± 1.19e+3	**5.01e-2 ± 0.15**	2.22e+4 ± 2.99e+4	—	2.05e+3 ± 2.81e+3	5.73e+4 ± 1.72e+5	5.49 ± 12.05
E6	200.8 ± 0.52	18.26 ± 34.46	116.10 ± 3.86	189.20 ± 40.71	130.20 ± 25.14	213.9 ± 191.3	**3.83 ± 5.89**
E7	2330 ± 14.28	570.70 ± 1.11e+3	1.04e+3 ± 6.62	1.62e+3 ± 819.30	1.78e+3 ± 525.90	1.28e+3 ± 656.6	**3.60e-2 ± 4.49e-2**
E8	1.68e-2 ± 4.66e-5	**3.32e-4 ± 4.76e-4**	1.18 ± 2.12	0.12 ± 1.91e-2	0.13 ± 0.38	1.25e-2 ± 1.83e-2	**3.38e-8 ± 6.31e-8**
E9	6.09e-2 ± 3.73e-4	779.40 ± 2.33e+3	0.30 ± 0.15	1.03 ± 4.76e-2	0.43 ± 0.17	0.77 ± 1.11	**2.72e-6 ± 4.09e-6**
E10	2.39 ± 3.38e-2	**7.3e-11 ± 1.5e-10**	0.17 ± 0.21	**0.00 ± 0.00**	0.36 ± 4.76e-2	4.96e-12 ± 1.45e-11	3.72e-4 ± 5.62e-4
E11	240.2 ± 3.09	1.94e-2 ± 5.83e-2	3.08 ± 8.02	**0.00 ± 0.00**	98.80 ± 104.80	200.8 ± 125.0	9.36e-5 ± 2.81e-4
E12	4.35 ± 0.24	0.80 ± 2.41	2.97 ± 1.57	2.57e-2 ± 0.00	5.31 ± 1.66	15.42 ± 20.69	**1.11e-6 ± 2.22e-6**
E13	1.43 ± 5.5e-2	2.72 ± 8.16	9.43 ± 23.11	60.55 ± 15.63	182.90 ± 330.00	26.17 ± 22.79	**8.59e-7 ± 1.78e-6**

Table 5 confirms that SeTGAP achieved lower or comparable extrapolation MSE values across all problems. These results suggest that other methods, which optimize purely for in-domain MSE, tend to overfit and learn expressions lacking the structural flexibility needed for extrapolation. In contrast, SeTGAP's decompositional approach learns functional forms that better capture underlying relationships, enabling superior generalization beyond the training domain. For example, in problem E8, most SR methods effectively minimized prediction MSE. TaylorGP, prioritizing parsimony, evolved the expression $\tilde{f}(\mathbf{x}) = 2$, effectively smoothing the data but offering no meaningful solution. This illustrates that the simplest solution isn't always best, as small data variations may reflect functional components important for generalization. Table 5 also shows cases where NeSymRes achieved zero extrapolation error across all 10 iterations for E10 and E11, while SeTGAP had low but nonzero MSE. This is because some methods learned expressions that exactly matched the underlying form, eliminating the need for coefficient fitting. PySR, for instance, identified $\sqrt{x_0}\log(x_1^2)$ for E13 in its first iteration—an expression with no numerical coefficients. SeTGAP, by contrast, produced $2.000137\sqrt{x_0}\log|x_1|$, where the fitted constant caused minor errors. We also compare SeTGAP's extrapolation with that of the opaque NNs from which its symbolic expressions are derived. Despite being trained on the same in-domain data, the original networks consistently show significantly higher extrapolation errors, as shown in Table 5. This highlights SeTGAP's value not only as a regression method but also as an interpretability tool that extracts functional approximations with superior generalization.

Since SeTGAP was the only method that consistently identified the correct functional form, we evaluated its robustness under different noise levels, as shown in Table 6. As expected, interpolation and extrapolation errors increased with higher noise levels. Interpolation errors remained low across all cases, while extrapolation errors showed a few exceptions due to poor coefficient fitting or incorrect functional form identification. For example, E3 and E5 exhibited high extrapolation errors because their functions contain exponential terms. Even when the learned expressions matched the expected functional form, small coefficient errors in the exponential term led to significant deviations for larger values of x_1. A similar case was observed in E1, where extrapolation errors were high for $\sigma_a \geq 0.03$ despite correctly identifying the functional form. The ability to recover the correct functional form in the presence of noise can be attributed to two factors. First, the Multi-Set Transformer was trained with small noise levels, making it more resilient to perturbations. Second, the regression NN $\hat{f}$ used during inference to generate the multiple input sets $\tilde{\mathbf{D}}_v$

Table 6: MSE comparison using SeTGAP with noisy data

Eq.	Interpolation ($\sigma_a = 0$)	Interpolation ($\sigma_a = 0.01$)	Interpolation ($\sigma_a = 0.03$)	Interpolation ($\sigma = 0.05$)	Extrapolation ($\sigma_a = 0$)	Extrapolation ($\sigma_a = 0.01$)	Extrapolation ($\sigma_a = 0.03$)	Extrapolation ($\sigma_a = 0.05$)
E1	7.159e-03	1.259e-02	1.236e-01	5.455e-01	1.012e-01	4.433e-01	1.645e+00	3.845e+00
E2	7.024e-03	4.459e-03	1.563e-02	4.754e-02	1.300e-01	5.232e-02	1.017e-01	3.504e-01
E3	1.063e-03	1.022e-03	7.219e-03	2.003e-02	2.961e-01	2.055e+02	7.467e+01	1.652e+02
E4	1.158e-04	2.969e-03	2.061e-01	2.228e-02	4.121e-02	1.022e+01	7.420e+01	3.105e+01
E5	7.211e-06	6.807e-03	5.288e-01	1.694e-01	3.069e-01	1.097e+01	8.402e+01	2.312e+02
E6	5.619e-03	1.147e-02	1.726e-02	4.819e-02	4.508e-02	4.651e+00	2.081e-01	5.598e-01
E7	2.655e-06	1.076e-02	6.822e-02	1.903e-01	6.447e-05	2.517e-01	1.247e+00	3.520e+00
E8	1.742e-06	2.413e-05	2.103e-04	5.828e-04	1.817e-10	1.686e-07	7.539e-09	1.672e-07
E9	4.907e-08	2.160e-04	1.943e-03	5.408e-03	2.780e-07	3.063e-04	2.756e-03	7.943e-03
E10	2.282e-06	3.298e-05	2.950e-04	8.192e-04	1.857e-03	2.233e-04	4.793e-04	3.379e-03
E11	1.178e-04	1.838e-02	1.654e-01	4.595e-01	9.308e-04	1.453e-01	1.308e+00	3.632e+00
E12	8.029e-06	4.948e-04	4.436e-03	1.232e-02	7.273e-06	1.031e-03	9.257e-03	2.571e-02
E13	2.583e-07	4.200e-03	3.991e-02	1.047e-01	1.015e-06	9.599e-03	9.348e-02	2.389e-01

smooths the estimated response values, mitigating the impact of noise. In the remaining cases, we observed two outcomes. SeTGAP correctly identified the functional forms of individual variables, but noise hindered the detection of relationships between variables. Otherwise, incorrect but reasonable univariate skeletons were identified, leading to expressions with small errors; e.g., for E9 and $\sigma_a = 0.05$, SeTGAP produced $\tilde{f}(\mathbf{x}) = 5.965\sqrt{0.63\log(9.4x_1 + 6.4) + 1} - \log(11.26x_0^2 + 2.82) - 7.71$.

SeTGAP is limited by the complexity of skeletons and the set of unary and binary operators used to train the Multi-Set Transformer. This limitation, inherent to any neural SR method, can be addressed through transfer learning, where the model is fine-tuned on more complex tasks to expand its expressivity. Another limitation is the computational cost due to multiple intermediate optimization processes, making SeTGAP less efficient than all compared approaches. However, in applications like scientific discovery, where the goal is to derive interpretable and reliable mathematical expressions rather than simply optimizing predictive accuracy, the additional computational effort is warranted. In this work, we do not aim to achieve competitive computational complexity, but rather to demonstrate that a decomposable approach reconstructs the correct functional form of a system more effectively. Future work will focus on developing fast and scalable merging strategies capable of synthesizing high-dimensional expressions efficiently.

6 Conclusion

Symbolic regression aims to find interpretable equations that represent the relationships between input variables and their response. As such, SR represents a promising avenue for building interpretable models, a key aspect of modern machine learning. By distilling opaque regression models into transparent mathematical expressions, SR provides insight into their inner workings and enhances confidence in their reliability.

In this work, we introduced SeTGAP, a decomposable SR method that integrates transformers, genetic algorithms, and genetic programming. It first generates multiple univariate skeletons that capture the functional relationship between each variable and the system's response. These skeletons are systematically merged using evolutionary approaches, ensuring interpretability throughout the process. Experimental results demonstrated that SeTGAP consistently recovered the correct functional forms across all tested problems, unlike the compared methods. In addition, it exhibited robustness against varying noise levels.

Future work will investigate how the order of skeleton merging influences the learned expressions and their performance. We also aim to extend SeTGAP beyond unary and binary operators to identify more complex functions, such as differential operators and integral transforms. This expansion would significantly enlarge the search space, increasing the challenge of exploration. To address this, we plan to refine the Multi-Set Transformer's structure, incorporating prior knowledge to better guide the search process. Finally, while effective in consistently recovering ground-truth functions in low-dimensional settings, the evolutionary merging process becomes computationally infeasible as dimensionality increases. Future work will focus on developing faster, scalable merging strategies, grounded in a formal theoretical framework, to determine when correctly inferred skeletons can be reliably combined into coherent multivariate expressions.

References

Maryam Amir Haeri, Mohammad Mehdi Ebadzadeh, and Gianluigi Folino. Statistical genetic programming for symbolic regression. *Applied Soft Computing*, 60:447–469, 2017.

Ignacio Arnaldo, Krzysztof Krawiec, and Una-May O'Reilly. Multiple regression genetic programming. In *Genetic and Evolutionary Computation Conference*, pp. 879–886, 2014.

Amanda Bertschinger, Q. Tyrell Davis, James Bagrow, and Joshua Bongard. The metric is the message: Benchmarking challenges for neural symbolic regression. In *Machine Learning and Knowledge Discovery in Databases*, pp. 161–177, 2023.

Amanda Bertschinger, James Bagrow, and Joshua Bongard. Evolving Form and Function: Dual-Objective Optimization in Neural Symbolic Regression Networks. In *Genetic and Evolutionary Computation Conference*, pp. 277–285, July 2024. ISBN 979-8-4007-0494-9.

Luca Biggio, Tommaso Bendinelli, Alexander Neitz, Aurelien Lucchi, and Giambattista Parascandolo. Neural symbolic regression that scales. In *International Conference on Machine Learning*, volume 139, pp. 936–945, 2021.

Gustau Camps-Valls, Andreas Gerhardus, Urmi Ninad, Gherardo Varando, Georg Martius, Emili Balaguer-Ballester, Ricardo Vinuesa, Emiliano Diaz, Laure Zanna, and Jakob Runge. Discovering causal relations and equations from data. *ArXiv*, abs/2305.13341, 2023.

Miles Cranmer. Interpretable machine learning for science with PySR and SymbolicRegression.jl. *ArXiv*, abs/2305.01582, 2023.

Stephen Dignum and Riccardo Poli. Generalisation of the limiting distribution of program sizes in tree-based genetic programming and analysis of its effects on bloat. In *Genetic and Evolutionary Computation Conference*, pp. 1588–1595, 2007.

M. Ebner. On the search space of genetic programming and its relation to nature's search space. In *IEEE Congress on Evolutionary Computation*, volume 2, pp. 1357–1361, 1999.

Renato Filho, Anisio Lacerda, and Gisele Pappa. Explaining symbolic regression predictions. In *IEEE Congress on Evolutionary Computation*, pp. 1–8, 2020.

Roger Fletcher. *Practical Methods of Optimization.* John Wiley & Sons, New York, NY, USA, second edition, 1987.

Baihe He, Qiang Lu, Qingyun Yang, Jake Luo, and Zhiguang Wang. Taylor genetic programming for symbolic regression. In *Genetic and Evolutionary Computation Conference*, pp. 946–954, 2022.

John H. Holland. Genetic algorithms. *Scientific American*, July 1992.

Nan Jiang and Yexiang Xue. Symbolic regression via control variable genetic programming. In *Machine Learning and Knowledge Discovery in Databases: Research Track*, pp. 178–195, 2023.

Ying Jin, Weilin Fu, Jian Kang, Jiadong Guo, and Jian Guo. Bayesian symbolic regression. *ArXiv*, abs/1910.08892, 2020.

Pierre-Alexandre Kamienny, Stéphane d'Ascoli, Guillaume Lample, and Francois Charton. End-to-end symbolic regression with transformers. In *Advances in Neural Information Processing Systems*, volume 35, pp. 10269–10281, 2022.

Gabriel Kronberger, Bogdan Burlacu, Michael Kommenda, Stephan M. Winkler, and Michael Affenzeller. *Symbolic Regression.* Chapman and Hall/CRC, New York, August 2024. ISBN 978-1-315-16640-7.

William La Cava, Patryk Orzechowski, Bogdan Burlacu, Fabrício Olivetti de França, Marco Virgolin, Ying Jin, Michael Kommenda, and Jason Moore. Contemporary symbolic regression methods and their relative performance. In *Advances in Neural Information Processing Systems*, 2021.

W. B. Langdon, T. Soule, R. Poli, and J. A. Foster. The Evolution of Size and Shape. In *Advances in Genetic Programming*. The MIT Press, 07 1999.

Juho Lee, Yoonho Lee, Jungtaek Kim, Adam Kosiorek, Seungjin Choi, and Yee Whye Teh. Set Transformer: A framework for attention-based permutation-invariant neural networks. In *International Conference on Machine Learning*, volume 97, pp. 3744–3753, 2019.

Kin Long Kelvin Lee and Nalini Kumar. Artificial intelligence for scientific discovery at high-performance computing scales. *Computer*, 56(4):116–122, 2023.

Pantelis Linardatos, Vasilis Papastefanopoulos, and Sotiris Kotsiantis. Explainable AI: A review of machine learning interpretability methods. *Entropy*, 23(1), 2021.

Nour Makke and Sanjay Chawla. Interpretable scientific discovery with symbolic regression: A review. *ArXiv*, abs/2211.10873, 2022.

Georg Martius and Christoph H. Lampert. Extrapolation and learning equations. *ArXiv*, abs/1610.02995, 2016.

T. Nathan Mundhenk, Mikel Landajuela, Ruben Glatt, Claudio Santiago, Daniel Faissol, and Brenden Petersen. Symbolic regression via neural-guided genetic programming population seeding. In *Advances in Neural Information Processing Systems*, 2021.

Patryk Orzechowski, William La Cava, and Jason Moore. Where are we now? A large benchmark study of recent symbolic regression methods. In *Genetic and Evolutionary Computation Conference*, pp. 1183–1190, 2018.

Brenden K Petersen, Mikel Landajuela, T Nathan Mundhenk, Claudio P Santiago, Soo K Kim, and Joanne T Kim. Recovering mathematical expressions from data via risk-seeking policy gradients. In *International Conference on Learning Representations*, 2021.

Riccardo Poli, William B. Langdon, and Stephen Dignum. On the limiting distribution of program sizes in tree-based genetic programming. In *Genetic Programming*, pp. 193–204, Berlin, Heidelberg, 2007. Springer Berlin Heidelberg.

Subham Sahoo, Christoph Lampert, and Georg Martius. Learning equations for extrapolation and control. In *International Conference on Machine Learning*, pp. 4442–4450, 10–15 Jul 2018.

Michael Schmidt and Hod Lipson. Distilling free-form natural laws from experimental data. *Science*, 324 (5923):81–85, 2009.

Parshin Shojaee, Kazem Meidani, Amir Barati Farimani, and Chandan Reddy. Transformer-based Planning for Symbolic Regression. In *Advances in Neural Information Processing Systems*, volume 36, pp. 45907–45919, December 2023.

Leonardo Trujillo, Luis Muñoz, Edgar Galván-López, and Sara Silva. neat genetic programming: Controlling bloat naturally. *Information Sciences*, 333:21–43, 2016.

Ho Fung Tsoi, Vladimir Loncar, Sridhara Rao Dasu, and Philip Harris. Symbolnet: Neural symbolic regression with adaptive dynamic pruning for compression. *Machine Learning: Science and Technology*, 2025.

Silviu-Marian Udrescu and Max Tegmark. AI Feynman: A physics-inspired method for symbolic regression. *Science Advances*, 6(16):eaay2631, 2020.

Mojtaba Valipour, Bowen You, Maysum Panju, and Ali Ghodsi. SymbolicGPT: A generative transformer model for symbolic regression. In *Advances in Neural Information Processing Systems*, 2022.

Ashwin K Vijayakumar, Michael Cogswell, Ramprasath R. Selvaraju, Qing Sun, Stefan Lee, David Crandall, and Dhruv Batra. Diverse beam search: Decoding diverse solutions from neural sequence models. *ArXiv*, abs/1610.02424, 2018.

Marco Virgolin and Solon P Pissis. Symbolic regression is NP-hard. *Transactions on Machine Learning Research*, 2022. ISSN 2835-8856.

Matthias Werner, Andrej Junginger, Philipp Hennig, and Georg Martius. Informed equation learning. *ArXiv*, abs/2105.06331, 2021.

A Proof of Proposition 1

Proof. We prove the proposition by structural induction on the composition of operations that define $f(\mathbf{x})$. We begin with the base cases. If $f(\mathbf{x})$ is a constant, the required form is satisfied trivially by setting $c_0 = c$ with no additional terms. If $f(\mathbf{x}) = x_i$ is a single-variable function, the structure is preserved by setting $c_0 = 0$, $c_{1,1} = 1$, and $T_{1,1}(\mathbf{x}) = x_i$.

For the inductive hypothesis, assume the decomposition holds for functions $h(\mathbf{x})$ and $u(\mathbf{x})$, which are composed of unary or binary operations. That is, each function can be written in the form $h(\mathbf{x}) = c_0' + \sum_i c_{i,j}' \prod_j \nu_{i,j}'(T_{i,j}'(\mathbf{x}))$ and $u(\mathbf{x}) = c_0'' + \sum_i c_{i,j}'' \prod_j \nu_{i,j}''(T_{i,j}''(\mathbf{x}))$, with terms $T_{i,j}'$ and $T_{i,j}''$ themselves recursively decomposable in the same way.

For the inductive step, we consider the result of applying unary or binary operations to such functions. For a unary operation $f(\mathbf{x}) = \nu(h(\mathbf{x}))$, the expression satisfies the required structure by treating the composition as a single term ($i = 1$, $j = 1$) where $T_{1,1}(\mathbf{x}) = h(\mathbf{x})$, $\nu_{1,1} = \nu$, $c_0 = 0$, and $c_{1,1} = 1$. Since $h(\mathbf{x})$ itself satisfies the recursive form by the inductive hypothesis, $f(\mathbf{x})$ does as well.

Now consider a binary operation $f(\mathbf{x}) = h(\mathbf{x}) \circ u(\mathbf{x})$, where $\circ$ is a binary operator. For the addition operation ($\circ = +$), substituting the expressions and grouping constants and summation terms yields:

$$f(\mathbf{x}) = (c_0' + c_0'') + \sum_i c_{i,j}' \prod_j \nu_{i,j}'(T_{i,j}'(\mathbf{x})) + \sum_i c_{i,j}'' \prod_j \nu_{i,j}''(T_{i,j}''(\mathbf{x})).$$

This expansion matches the desired structure $f(\mathbf{x}) = c_0 + \sum_i c_i \prod_j \nu_{i,j}(T_{i,j}(\mathbf{x}))$, considering that $c_0 = c_0' + c_0''$ and that the summation terms come directly from the sub-expressions of $h(\mathbf{x})$ and $u(\mathbf{x})$.

For the multiplication operation ($\circ = \cdot$), the product expansion gives:

$$f(\mathbf{x}) = c_0' c_0'' + c_0' \sum_i c_{i,j}'' \prod_j \nu_{i,j}''(T_{i,j}''(\mathbf{x})) + c_0'' \sum_i c_{i,j}' \prod_j \nu_{i,j}'(T_{i,j}'(\mathbf{x}))$$
$$+ \sum_{i,j} c_{i,j}' c_{i,j}'' \prod_j \nu_{i,j}'(T_{i,j}'(\mathbf{x})) \prod_j \nu_{i,j}''(T_{i,j}''(\mathbf{x})).$$

Each term fits into the structure $f(\mathbf{x}) = c_0 + \sum_i c_i \prod_j \nu_{i,j}(T_{i,j}(\mathbf{x}))$: the constant term can be expressed as $c_0 = c_0' \cdot c_0''$; the second and third terms are sums over products of unary operator applications, consistent with the required form; and the fourth term comprises products of sub-expressions from $h(\mathbf{x})$ and $u(\mathbf{x})$, which can be grouped as new terms $\prod_j \nu_{i,j}(Ti, j(\mathbf{x}))$. For other scalar-valued binary operations that are algebraically reducible, such as subtraction and division, we use the identities $h(\mathbf{x}) - u(\mathbf{x}) = h(\mathbf{x}) + (-u(\mathbf{x}))$ and $h(\mathbf{x})/u(\mathbf{x}) = h(\mathbf{x}) \cdot u(\mathbf{x})^{-1}$. Negation and inversion are unary operations, and since the unary case has been established, the result follows. Binary operations that involve non-scalar interactions (e.g., convolution or vector operations) are beyond the scope of this structural decomposition.

Thus, any function $f(\mathbf{x})$, defined by finite compositions of unary and binary operations, can always be expressed in the required form. Since the base case holds and the inductive step is proven, the proposition is established by structural induction. □

B Comparison of Predicted Mathematical Expressions

This appendix presents in Tables 7–33 the complete set of mathematical expressions learned by each symbolic regression method across iterations 2 to 9 of dataset generation. Results corresponding to the first iteration were reported in Tables 2–4.

Table 7: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 2

Method	E1	E2	E3	E4
PySR	$0.607x_0x_1$	$0.726x_0 - 0.759x_2 + 12.944\tanh(0.107x_2) - 4.572 + 11.225e^{-0.099x_0}$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$0.091\cosh(x_2) + \cosh(0.064x_0^2 - 0.052x_1 - 0.052x_3 + 1.154) - 1.914$
TaylorGP	$0.597x_0x_1$	$2.95(\sqrt{e^{\sqrt{\log(2x_1)}}} + 0.206 + e^{-0.294x_0} + e^{\tanh(x_2)} + e^{\tanh(\log(x_0))} + \sin^{0.5}(1.821((-e^{\log(x_0\tanh(x_0)+x_1)^{0.5}} - e^{-0.294x_0)^{0.5}} + 0.179)^{0.5} + e^{\tanh(x_2)} + e^{\tanh(\log(x_0))} + \sqrt{\sin(x_2)} + e^{-0.294x_0}) + e^{-0.294x_0})^{0.5}$	$-x_0 - (-x_0 - 0.868(e^{x_0})^{0.5} + e^{x_0})^{0.5} + e^{x_0} - \cos(\cos^{0.5}(\frac{x_1}{x_0})) + 0.092$	$\sqrt{x_0}\log(x_0) + x_2 - e^{\tanh(\log(x_2))}$
NeSymReS	$0.586x_0x_1 + \cos(0.593(x_0 - 0.979x_1)^2)$	$-x_0 + 0.001e^{0.597x_1} + e^{\sin(0.175x_2)} + 7.267$	$0.581e^{x_0} - 0.463\sin(1.104(0.003x_1 + 1)^2)$	—
E2E	$(0.081x_0 - 17.975)(0.051\sqrt{0.008x_1 + 1} + 0.006)\cos1.744(0.045(0.001 \\ |15810201.601(0.292x_0 + 1)^3 + 0.016|+1)^2 \\ +1)^{0.5} - 0.007) + (2.709x_0 - 0.014) \\ (0.223x_1 + 0.002)$	$-0.002x_1 + ((0.016x_0 + 0.004)(1.231x_0 - 9.407) + \sin(0.235x_2 - 0.031) - 0.007) \\ (0.004x_0 + 0.087x_1 - 0.081x_2 + 3.316) + 6.19$	$0.177\sqrt{(0.58e^{3.132x_0} + 1)} + 0.486\cos(4.706(x_1 + 0.016)^2 - 0.066) - 0.104$	$0.002|31.163x_3 + 5.524(x_0 + 0.023)^4 + 5.498(0.985x_1 - (x_2 - 0.038)^2 + 0.042)^2 - 1.464|$
TPSR	$-0.071x_1(-8.596x_0 - 0.119) - 0.113\cos(x_0 - 0.782) - 0.009$	$(-1.362 + 0.321/(-9.597e^{2.221x_0 - 6.275x_2} - 0.665)) \\ (-0.044x_2 + 1.552 + 1/((0.008x_0 + 0.06) \\ (0.043x_0 - 0.981) - 0.109))$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$0.044x_1 + 0.01(-x_2 - 0.013)^4 + 0.063(-0.355x_0^2 + 0.418x_1 + 0.209x_3 - 1)^2 - 0.126$
SeTGAP	$0.61x_0x_1 + 1.041\sin((2.243x_0 - 1.512)(x_1 - 0.669) + 0.005)$	$-0.497x_0 + 0.063x_0^2 + (0.648\sqrt{0.1x_1 + 1} + 0.002)(4.913\sin(0.205x_2) + 0.003) + 6.497$	$0.151e^{1.499x_0} + 0.499\sin(3x_1 + 1.574)$	$-0.02x_0^2x_1 - 0.004x_0^2 + 0.01x_0^4 + 0.01x_1^2 \\ -0.02x_2^2x_3 + 0.01x_2^4 + 0.01x_3^2 + 0.01$

Table 8: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 2

Method	E5	E6	E7	E8
PySR	$e^{1.2x_3}$	$\frac{\cos(0.2x_2^2)}{\cos(1.947\tanh(\tanh(0.281x_1)))} + \tanh(x_0)$	$-x_1^2\sinh(0.492\cos(6.274x_0 + 1.574) + 0.778) + 1.049$	$2.036\cosh(\tanh(x_0)) + 3.34\cosh(\tanh(\tanh(x_1\tanh(x_1)))) - 5.5$
TaylorGP	$x_3^{0.5}e^{x_3} + \sin^{0.5}(0.056e^{x_3})$	$\tanh(x_0) + 0.327 + \frac{4.831\sin(x_2)}{x_2}$	$-0.974x_1^2 + |x_1|^{0.5} + 0.008$	$\frac{\tanh(x_1)}{\cos(\tanh(x_0))}$
NeSymReS	—	$0.149x_0 + x_1\sin(0.29x_1 + \frac{x_2}{x_1})$	$\frac{0.289x_0 + x_1^2}{\cos(2.667(0.131x_1 - 1)^2) - 2.044}$	$\cos(\frac{\sin(\frac{x_0}{x_1})}{x_0}) + 0.629$
E2E	$0.96e^{1.218x_3} - 0.004 + 0.947\sin(221.795x_1 + 27.668)$	$(5.971|0.169x_1 - 0.005|+0.14) \\ \cos((1.531 - 17.089x_2)(-0.012x_2 - 0.009)) + 0.8\arctan(0.695x_0 + 0.196) - 0.01$	$(0.009\sin(7.746x_0 + 0.025) + 7.454) \\ (0.002(0.055x_0 - 1)^3 - 0.863(x_1 - 0.001)^2 + 0.696) \\ /(6.13\sin(7.746x_0 + 0.025) + 9.52)$	$-1.35\arctan(-0.096|2.433x_1 + (0.328x_0 - 0.01)(-2.094x_1 + (0.008 - 1.796x_0) \\ (0.465 - 13.81x_1)(-0.835x_1 - 0.038) \\ +0.128) + 0.113| - 0.04) - 0.035$
TPSR	$0.003 + 0.987e^{0.001x_2 + 1.206x_3} - 0.036\cos(0.188x_0 - x_1 + 1.229)$	$2795.262 - 2795.746\sin(0.008 \\ (-0.169x_0 + 0.025x_2 + 1)^2 + 1.529)$	$-0.003(-0.124x_0 - 21.52)(x_1 - 0.202) \\ (19.963(-0.322x_1 - 0.484 \\ \cos(6.286x_0 + 10.97) + 1)^2 - 56.092)$	$2.008 - 1.496e^{-0.644|(0.071x_0 - 0.001)(7.973x_1 - 0.124)|}$
SeTGAP	$e^{1.2x_3} - \sin(x_0 + x_1x_2 + 9.43)$	$-(0.997|x_1|+0.008)\sin(0.201x_2^2 - 1.611) + \tanh(0.492x_0) - 0.008$	$\frac{2.785x_1^2 + 0.05\sin(6.283x_0 - 3.149) - 2.793}{2.77\sin(6.283x_0 - 3.149) - 4.167}$	$2 - \frac{12.162}{-0.026x_1 + 0.132x_1^2 + 0.084x_1^4 + 12.008x_1^4 + 12.144} - \frac{14.373}{14.113x_0^4 + 0.232(-x_0)^2 + 14.361}$

Table 9: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 2

Method	E9	E10	E11	E12	E13
PySR	$\tan(1.044\cos(0.627x_0) + 0.242) + \tan(\cos(e^{\cos(0.893x_1^{0.5})})) - 2.117$	$\sin(x_0e^{x_1})$	$2x_0\log(x_1^2)$	$x_0\sin(\frac{1.0}{x_1}) + 1.0$	$(x_0(\tan(\sinh(\sinh(\cosh(\\ \tanh(0.309\cosh(x_1)))))) \\ +7.703))(x_0 + 8.766)$
TaylorGP	$-0.906\log(|x_0|) + \log(|x_1|)\tanh(x_1) \\ -|x_0 + 0.119|^{0.5}$	$\sin(x_0e^{x_1})$	$4x_0\log(|x_1|)$	$0.752x_0\tanh(\frac{0.985}{x_1}) + 0.752$	$2\sqrt{x_0}\log(|x_1|)$
NeSymReS	$\log(|\frac{x_1}{x_0}|) - 1.272$	$\sin(x_0e^{x_1})$	$x_0\log(x_1^4)$	$(x_0 + x_1)\sin(\frac{1}{x_1})$	$0.231x_0 + \log(x_1^2)$
E2E	$((1.9 - 0.6\log(13.214(x_0 + 0.001)^2 \\ +0.6))(0.23x_1 + 0.231) - 0.6)/ \\ (0.23x_1 + 0.231)$	$-0.02x_0 - 0.982\sin(\\ (29.612x_0 + 0.106) \\ (10.592e^{0.571x_1} \\ -0.033)(0.017\cos(\\ 0.07e^{0.902x_1} - 4.53) \\ +0.002)) - 0.003$	$x_0(6.669\log(0.19 \\ (|22.158x_1 + 0.255|+ \\ 0.111)^{0.5} - 0.01) + 0.1)$	$0.997 - 7.62 \\ \sin(\frac{(0.131x_0 - 0.001)(0.004x_1 - 6.7)}{6.661x_1 + 0.137})$	$(0.091\log(0.174x_0 + 0.772) \\ +0.086)(27.6\log(0.711 \\ (x_1 + 0.043)^2 + 0.13) \\ -0.099)$
TPSR	$\left(-3.757 - \frac{2.69}{10.828(x_0 - 0.002)^2 + 13.341}\right) \\ (-1.817\arctan(x_1 + 2.012) - 10.316) \\ -0.045|(39.196x_0 + 0.018) \\ (0.003x_1 - 0.256)|-48.763$	$1.0\sin(0.9976x_0e^{x_1})$	$-1.567x_0 + (0.007 + 1/(-1.1 \\ |12.472x_1 - 0.03|-0.988)) \\ (12.636 - 27.802(x_1 - 0.002)^2) \\ (1.515x_0 + 0.013) - 0.05$	$6.9697135x_1 + \\ 1.654205 - \frac{4.6\cdot10^{-5}}{x_1}$	$(-0.053 + \\ \frac{1}{0.228|6.18x_1 + 0.015|+0.15}) \\ (0.088 - 0.092x_1^2) \\ \left(1.026 - \frac{1.844}{x_0 + 1.941}\right) \\ (-2.037x_0 - 40.447)$
SeTGAP	$-\log(30.042x_0^2 + 7.509) + \\ |\log(13.19x_1 + 6.59)|+0.13$	$\sin(x_0e^{x_1})$	$2x_0\log(x_1^2)$	$x_0\sin(\frac{1}{x_1}) + 1.0$	$2.0\sqrt{x_0}\log(|x_1|)$

In each iteration, a new dataset was generated using a unique initialization seed, ensuring that all methods were trained and evaluated on the same data for a fair comparison. Each table reports the best expression found by each SR method, with shaded cells indicating cases where the learned expression's functional form matches the underlying ground-truth function. These results provide insight into the stability and consistency of each method in discovering accurate symbolic representations across different dataset realizations.

Table 10: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 3

Method	E1	E2	E3	E4
PySR	$0.607x_0x_1 +$ $1.1\cos(-2.25x_0x_1 +$ $1.5x_0 + 1.5x_1 + 0.571)$	$e^{e^{-\sin(0.128x_0)}} +$ $1.214e^{\sqrt{e^{\tanh(x_2)}}} + \tanh(\frac{x_1}{x_2})$	$0.15e^{1.499x_0} +$ $0.497\cos(3.001x_1 + \frac{1.557}{x_0+22.291})$	$-0.02x_2^2(\frac{2.628x_1\sin(x_2)}{x_2} + x_3) -$ $0.007(x_1 - 13.401)(\cosh(x_0) + \cosh(x_2))$
TaylorGP	$0.6x_0x_1$	$-0.503x_0 + 0.001x_1 + 0.211x_2 + 8.607$	$0.346e^{x_0}\log(1.348|(-0.247x_0 + e^{x_0}$ $-0.247e^{0.5|\sin(x_1)|^{0.5}})^{0.5}|) + 0.107$	$x_0 + x_2 - |x_0|^{\frac14}|x_2|^{\frac14} + 0.778$
NeSymReS	$0.586x_0x_1 +$ $\cos(0.585(0.991x_0 - x_1)^2)$	$-x_0 + 0.384e^{0.024x_1} + e^{\sin(0.175x_2)} + 6.813$	$0.582e^{x_0} -$ $0.749\sin(0.57(1 - 0.001x_1)^2)$	—
E2E	$0.255x_0(2.399x_1 + 0.034) +$ $1.11\cos(3089.014x_0 +$ $1035.398) - 0.006$	$(0.776 - 0.111x_0)(0.171x_1 + (0.174x_0 + 0.019)$ $(0.001x_2 - 0.07)(-0.499x_0 +$ $0.042\cos(1.689x_1 + 50.388)$ $+42.271) + 0.017) - 2.94\cos((2.184x_1 + 86.243)$ $(0.025\sin(0.116x_2 - 0.038) + 0.016)) + 6.62$	$0.113e^{1.594x_0} + 0.511\cos(3.08$ $(x_1 + 0.032)^2 + 0.008) + 0.136$	0.0
TPSR	$-0.141x_1(0.053 - 4.272x_0) +$ $0.002\sin(6.904x_1 - 7.997)$ $+0.05$	$(-0.08 + \frac{1}{-468.708e^{3.582x_0 - 9.1x_2} - 11.692})$ $(-0.03x_1 - 7.596)((-0.427x_0 - 0.022x_2 -$ $(-0.006x_0 - 0.303)(1.156x_0 + 2.022) - 0.294)^2$ $+1.84)(-0.008x_1 + 1.435 + \frac{1}{0.002x_0 + 0.394})$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$0.011\left(0.705 - (-x_0 - 0.011)^2\right)^2 +$ $0.049(-0.003x_0 - 0.243x_1$ $+0.41x_2^2 - 0.407x_3 + 1)^2 - 0.065$
SeTGAP	$0.607x_0x_1 +$ $1.123\sin((2.243x_0 - 1.496)$ $(x_1 - 0.654) + 0.014)$	$-0.5x_0 + 0.062x_0^2 +$ $(3.163\sqrt{0.1x_1 + 1} + 0.001)\sin(0.2x_2) + 6.513$	$0.151e^{1.497x_0} -$ $0.5\sin(3.004x_1 - 7.854)$	$-0.004x_0 - 0.02x_0^2x_1 +$ $0.017(-x_0)^2 + 0.009(-x_0)^4 + 0.008(-x_1)^2 +$ $0.01(-x_2)^4 + 0.011(-x_3)^2 - (0.013x_2^2 + 0.003)$ $(1.575x_3 + 0.614) - 0.024$

Table 11: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 3

Method	E5	E6	E7	E8
PySR	$e^{1.2x_3} + \sin(x_0 + x_1x_2)$	$|x_1|\cos(0.2x_2^2) + \tanh(0.502x_0)$	$-0.412x_1^2\cosh(1.055e^{-0.743\sin(6.278x_0)}) + 1$	$2.046\cosh(\tanh(x_0)) +$ $2.046\tanh(\frac{x_1}{\tanh(\sinh(1.875x_1))})$ -3.196
TaylorGP	$2e^{x_3} - \log(3e^{x_3} + 0.856\sqrt{e^{(x_3)}|x_3|} -$ $\log(2.32e^{x_3} + 0.856\sqrt{e^{(x_3)}|x_3|}) - 0.26) - 0.26$	$\frac{x_1\sin(x_2)}{x_2\tanh(x_1)} + \cos(x_2) + \tanh(x_0)$	$-0.974x_1^2 + \sqrt{|x_1|}$	$1.148|x_0|^{\frac14}|x_1|^{\frac14} - 0.105$
NeSymReS	—	$0.144x_0 + x_1\sin(0.065x_1 + \frac{x_2}{x_1})$	$\frac{0.072x_0 + x_1^2}{\cos(3.099(0.012x_1 - 1)^2) - 0.252}$	$\cos\left(\frac{\sin(\frac{x_0}{x_1})}{x_0}\right) + 0.69$
E2E	$0.871e^{1.253x_3} + 0.981\cos(88.561(0.014|3.137x_1 +$ $31752866.296(0.104\arctan(0.087x_2 +$ $5.508) - 1)^2 - 2.465|-1)^{0.5} - 0.006) + 0.13$	$(0.002\sin(0.816x_1 + 0.616) +$ $0.151)(0.036x_1 + \cos(0.711(0.014$ $-x_2)^2 - 0.573) + 3.995)$	$(0.002(1 - 0.256\sin(0.183|-32.468x_0 +$ $0.047x_1 + 14.284|-0.293))^3$ $-0.026)(-0.043x_0 +$ $19.294(x_1 + 0.017)^2 - 0.543)$	0.865
TPSR	$(x_1 + 923.05)(0.001e^{1.2x_3} - 0.01) + 8.903$	$(0.016\cos(0.771(x_1 + 0.145)^2$ $-0.875) - 0.084)(-2.126x_0 -$ $43.836\cos((3.209 - \frac{8.357}{0.028x_2 + 0.476})$ $(0.058x_2 + 0.044)) - 17.68)$	$-0.039x_0 + 294.76(-1 + 0.139/$ $(0.036 - 3.266e^{-7}/(-5.83e^{-9}$ $(-x_1 - 0.092)^2 - 6.046e^{-7})))^2 - 167.907$	$(-0.239x_1 - 4.712)$ $(0.455x_1 - 8.997)$ $(\arctan(4.665(0.02 - x_1)^2$ $(x_0 - 0.039)^2 + 31.821) - 1.521)$
SeTGAP	$1.006e^{1.201x_3} + 1.006\sin(1.008x_0 +$ $(1.005x_1 - 0.006)(x_2 + 0.029) + 12.451) - 0.007$	$1.001\cos(0.2x_2^2 + 0.01)$ $|x_1| + 1.0\tanh(0.5x_0) + 0.002$	$\frac{5.18x_1^2 - 5.18}{5.18\sin(6.283x_0 + 3.142) - 7.77}$	$2.0 - \frac{14.485}{0.142x_0^6 + 14.37(-x_0)^4 + 14.466} +$ $\frac{13.074}{-13.081x_1^4 - 13.074}$

Table 12: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 3

Method	E9	E10	E11	E12	E13
PySR	$1.172\sqrt{|x_1|} - 1.518\sinh(\sinh(\tan(\tanh($ $1.739e^{\cos(\cos(0.217x_0)} + \cos(\tanh(x_0))))))) + 2.232$	$\sin(x_0e^{x_1})$	$2x_0\log(x_1^2)$	$x_0\sin(\frac{1.0}{x_1}) + 1$	$\sqrt{x_0}\log(x_1^2)$
TaylorGP	$-\log(|x_0|) + \tanh(\log(|x_1|)) - \sqrt{|\log(\frac{0.568}{|x_0|})|}$	$\sin(x_0e^{x_1})$	$4.545x_0\log(|x_1|)$	$\frac{2.77\log(\sqrt{|x_0 + x_1|})}{\tanh(\frac{0.5(x_0+x_1)}{x_1})}$	$2\log(|x_1|)\sqrt{|x_0|}$
NeSymReS	$\log\left(\frac{0.286|x_1|}{|x_0|}\right)$	$\sin(x_0e^{x_1})$	$x_0\log(x_1^4)$	$(x_0 + x_1)\sin(\frac{1}{x_1})$	$0.241x_0 + \log(x_1^2)$
E2E	$-0.098 - 0.052/((-0.003(0.029x_1 - 1)^3 + 0.005 +$ $6.94/(-0.001x_0 - 51.126x_1 + (3.21$ $-0.085|32.081x_0 - 1.081|)$ $(2.068x_1 - 69.713) + 126.399)))$	$1.0\sin(0.035$ $x_0(26.26e^{1.16x_1}$ $-0.1)) - 0.001$	$(0.058|15.4\arctan($ $0.258x_1 + 0.008) - 0.009|$ $-0.005)(-0.006x_0 + 59.4$ $\sin(0.168x_0 - 0.001) + 0.011)$	$1.0 + 7.5\sin(0.001x_0$ $(0.1 - \frac{24.6}{-0.159x_1 - 0.005}))$	$1.08\log(0.02(-x_0 - 0.825)^2$ $(-x_1 - 0.057)^2$ $(-x_1 - 0.056)^2 + 0.101) + 0.09$
TPSR	$0.396x_1 + 0.004\left(1.464 + \frac{1}{19.402(0.002-x_0)^2 + 28.549}\right)$ $(0.873x_0 - 46.945)(-10.284x_0 + x_1 - 555.305)$ -149.911	$1.0\sin(1.001x_0e^{x_1})$	$(0.001 - 0.12x_0)(-0.971x_1^2 -$ $42.941 - 103.498(-0.605$ $|3.087x_1 - 0.011|-0.633))$	$(-1.292 - 0.14/(-0.179$ $(0.01x_0 + x_1 + 0.073)^2 -$ $0.106))(-0.005x_1 +$ $\frac{0.195 - 0.883x_0}{x_1 + 0.066} - 0.809)$	$(-0.104 + 1/(0.01|82.844x_1$ $-0.068|+0.111))(49.998 -$ $50.042\cos(0.042x_1))$ $(159.942 + \frac{170.493}{-0.07x_0 - 1.16})$
SeTGAP	$-0.999\log(11.668x_0^2 + 2.91) +$ $1.006|\log(13.4x_1 + 6.834)|-0.861$	$1.0\sin(1.0x_0e^{x_1})$	$2.0x_0\log(x_1^2)$	$1.0x_0\sin(\frac{1}{x_1}) + 1.0$	$\sqrt{1.0x_0}\log(x_1^2)$

Table 13: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 4

Method	E1	E2	E3	E4
PySR	$0.608x_0x_1 +$ $1.1\cos(1.076(1.045x_0$ $-0.697)(2x_1$ $-1.333) - 1.571)$	$(0.198x_1 + 2.985)\sin(0.201x_2) +$ $\cosh(0.836e^{\tanh(2.02e^{0.162x_0})} - 4.76)$	$13.457e^{\frac{1.753x_0 - 5.251}{\cosh(\tanh(\cos(1.525x_1)))}}$	$0.091x_1(\cos(0.643x_0) - 1.803) - 0.044x_3$ $\cosh(0.65x_2) + 0.091\cosh(x_0)$ $+0.091\cosh(x_2) - 0.027$
TaylorGP	$0.602x_0x_1$	$1.946\sqrt{x_0}(\tanh(x_0) - 0.837) +$ $1.946(\log(e^{x_2} + 11.111\sin(\cos(e^{x_0})) +$ $11.111\sqrt{|x_0(x_2 + 3.147\sqrt{|x_0|} + 0.984)|}))$	$\frac{-x_0 + e^{x_0} -}{\sqrt{|x_0 - e^{x_0} - \sqrt{|x_0|} + 0.785\sqrt{|e^{x_0} - 0.083|}}}$	$\frac{x_2}{\tanh(x_2)} + \log(|x_0|)\sqrt{|x_0|} - 1.571$
NeSymReS	$0.586x_0x_1 +$ $\cos(0.586(x_0$ $-0.988x_1)^2)$	$-x_0 + e^{\sin(x_2)} + 7.289$	$0.584e^{x_0} - 0.411$	—
E2E	$0.03x_0(19.977x_1 +$ $0.327) - 1.1\cos(31.768$ $x_1 + 2.118) + 0.016$	$0.075x_0^2 - 0.443x_0 + 0.014x_1x_2 +$ $0.002x_1 - 0.002x_2^2 + 0.083x_2 +$ $3.07\sin(0.227x_2 + 0.025) + 6.23$	$0.107x_0 + 0.156(0.318x_0 + 1)^2(x_0 - 0.002)^2$ $(|0.279x_1 + 0.578| + 0.133)^2 -$ $0.501\sin(0.091x_0 + 3.572x_1 - 1.273) + 0.152$	$(0.095 - 0.001x_1)(0.1(0.559x_1 - (x_0 -$ $0.003)^2 + 0.037)^2 + 0.073) + 0.036$ $(-0.524x_3 + (0.204x_0 - 0.001)(0.142x_2$ $+0.013) + (0.017x_2 + 0.001)$ $(32.565x_2 + 1.169) + 0.024)^2 - 0.001$
TPSR	$-0.069x_0(-8.798x_1 -$ $0.074) + 0.036\sin(5.606$ $x_1 + 2.258) + 0.009$	$0.001(31.887x_1 + x_2 + 12721.866)$ $\frac{e^{(-1.482(0.06x_0+0.651))}}{e^{-41.665e^{0.01x_1}+0.693x_2}-29.007}$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$11.1441\left(1 - 0.944(-0.074x_2 - 1)^2\right)^2$ -0.214569
SeTGAP	$0.607x_0x_1 +$ $1.1\sin((2.25x_0 - 1.489)$ $(x_1 - 0.663))$	$-0.5x_0 + 0.063x_0^2 +$ $(3.148\sqrt{(0.1x_1 + 1)} + 0.012)$ $\sin(0.204x_2) + 6.493$	$0.147e^{1.508x_0} +$ $0.5\sin(2.999x_1 + 1.571) + 0.002$	$-0.02x_0^2x_1 + 0.011x_0^4 + 0.009x_1^2 +$ $0.011x_3^2 - 0.013(-x_0)^2 + 0.009(-x_2)^4 -$ $(0.035x_2^2 - 0.002)(0.584x_3 - 0.46) - 0.016$

Table 14: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 4

Method	E5	E6	E7	E8
PySR	$e^{1.2x_3} + \sin(x_0 + x_1x_2)$	$\sqrt{(x_1^2)}\cos(0.2x_2^2) + \tanh(0.5x_0)$	$\frac{-1.0x_1^2 + \cos(0.011x_1\sin(6.283x_0))}{\sin(6.283x_0) + 1.5}$	$4.404\tanh(\cosh(0.869x_0)$ $\sqrt{\tanh(\cosh(x_1))})$ $\tanh(\cosh(0.869x_1)) - 2.427$
TaylorGP	$e^{x_3}\sqrt{|x_3|} + \tanh(0.515e^{x_3} -$ $0.331\sin(\sqrt{e^{x_3}|x_3 - 0.662|}))$	$\left(-\log(|x_2|) + |x_1|^{0.5}\right)(\cos(x_2) + 0.604) +$ $\cos(x_2) + \tanh(0.142x_0)$	$-0.974x_1^2 + |x_1|^{0.5}$	$1.108|x_0|^{\frac{1}{4}}|x_1|^{\frac{1}{4}}$
NeSymReS	—	$0.173x_0 + x_1\sin\left(0.284x_1 + \frac{x_2}{x_1}\right)$	$\frac{0.059x_0 + x_1^2}{\cos(1.577(-0.668x_1 - 1)^3) - 1.862}$	$\cos\left(\frac{\sin\left(\frac{x_0}{x_1}\right)}{x_0}\right) + 0.644$
E2E	$0.96e^{1.21x_3} - 0.963\sin(-0.001x_0$ $\left(0.03 + \frac{90.6}{0.216x_1 - 0.235}\right) + 0.555x_1 +$ $0.03x_2 + (0.737 - 6.778x_0)$ $(0.007x_1 - 1.38) + 0.898) + 0.055$	$0.011x_1 - (1.57 - 0.001x_2)\arctan(-0.155x_0$ $+ (0.055\sin(0.173x_1 + 0.031) - 0.005)$ $(53.8\cos(0.171x_1 - 0.172(x_2 + 0.075)^2 +$ $0.011) + 0.006) + 0.004) - 0.006$	$(11.393(x_1 - 0.128)^2 + 11.849)$ $(-0.02(-0.021\cos(1254.632$ $(1 - 0.528x_0)^2 + 0.173) - 1)^2$ $-0.022)$	$2.0 - 0.9(0.006(|3.121x_1$ $+0.11)(24.018(x_0 -$ $0.003)^2 + 6.0)| + 0.5)$
TPSR	$0.988e^{1.203x_3} + 0.055\sin(x_0$ $-8.658x_1 + 1.093) + 0.029$	$(0.001 + \frac{1}{x_0 - 411.157})(31.572 - 0.001x_2)$ $(0.214\cos(2.182x_1 + 2.001) - 5.403)$ $((4.295 - 1.096x_0)(-0.044x_1 - 0.586)$ $+14.767\cos(0.533x_2 - 0.05) + 9.177)$	$-0.026x_0 - 1.254 +$ $\frac{0.067 - 0.027(-x_1 - 0.023)^2}{0.006\sin(7.183x_0 + 2.841) + 0.03}$	$2.01 - 1.478$ $e^{-0.00384|x_1(90.031x_0 + 2.401)|}$
SeTGAP	$0.997e^{1.201x_3} + 0.999\sin(x_0 +$ $1.0x_1x_2 + 0.017) + 0.004$	$0.997\cos(0.202x_2^2 + 6.236)|x_1| +$ $1.0\tanh(0.5x_0)$	$\frac{5.94x_1^2 - 5.979}{-5.941\sin(6.283x_0 - 6.283) - 8.91}$ -0.004	$2.0 - \frac{15.123}{15.039x_1^4 + 15.129} -$ $\frac{15.086}{15.06x_0^4 + 15.09}$

Table 15: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 4

Method	E9	E10	E11	E12	E13
PySR	$1.141\sqrt{x_1} + 1.141(-0.895x_0 +$ $(0.059x_0^2 - 1.08)\tanh(3.763x_0))$ $\tanh(x_0)$	$\sin(x_0e^{x_1}\tanh($ $e^{e^{\tanh(\cosh(\tan(e^{x_1} + 0.775)))}}))$	$2x_0\log(x_1^2)$	$x_0\cos\left(1.659 - \frac{0.088x_0x_1 + x_0}{x_0x_1}\right) + 1.0$	$\sqrt{x_0}\log(x_1^2)$
TaylorGP	$0.26x_1 + \log\left(\frac{0.984}{|x_0|}\right) -$ $\sqrt{\left|\log\left(\frac{0.593}{|x_0|}\right) + \log\left(\left|\tanh\left(\frac{x_1}{x_0}\right)\right|\right)\right|}$	$\sin(x_0e^{x_1})$	$4x_0\log(|x_1|)$	$\left(x_0\tanh\left(\frac{0.664}{x_1}\right) + 1\right)\sqrt{|\tanh(x_1)|}$	$2\log(|x_1|)\sqrt{|x_0|}$
NeSymReS	$\log\left(\frac{0.282|x_1|}{|x_0|}\right)$	$\sin(x_0e^{x_1})$	$x_0\log(x_1^4)$	$(x_0 + x_1)\sin\left(\frac{1}{x_1}\right)$	$0.234x_0 + \log(x_1^2)$
E2E	$-0.6\log(10.525(0.003 - x_0)^2$ $+0.6) + 1.9 - \frac{0.6}{0.229x_1 + 0.235}$	$0.001 - 1.0\sin(0.561x_0$ $(-0.002 + 17.5/((-88.065$ $-\frac{0.009}{7.629x_0 + 19.891})(0.001$ $-0.025x_1) + 0.364(0.008$ $x_1 - 1)^2 - 7.514($ $0.732x_1 - 1)^2 - 5.172)))$	$-2.266x_0(0.018 -$ $0.26x_0(13.784(0.002$ $-(x_1 - 0.024)^2)^2$ $+0.008))$	$0.998 - 6.35\sin((0.006$ $+\frac{0.673}{0.173x_1 + 0.007})(-0.04x_0 - 0.002))$	$(5.75 - 8.24/(1.265($ $|x_1| + 0.008)^{0.5} + 0.135))$ $(2.72\log(0.174x_0$ $+0.991) + 0.188)$
TPSR	$-0.001x_0 + 1.386x_1 + 23.694\arctan(($ $0.005x_1 + 0.059)(0.155x_0^2 + 37.746$ $(1 - 0.074x_1)^2 - 1.731|x_0| +$ $25.317) - 4.298) + 12.633$	$0.9998\sin(0.9994x_0e^{x_1})$ -0.000683	$7.994x_0 + 0.052$	$(0.429 - 0.001x_0)(\arctan(224.422$ $((x_1 + 0.121)^3 - 0.052)^2 + 2.305)$ $-1.24)(-0.001x_0(-0.049 - \frac{6021.0}{x_1})$ $+0.081x_0 + 7.034)$	$(0.064x_0 + 0.751 - 119.56/$ $((-41.662 - 117.245e^{-0.175x_0})$ $(-0.108|x_1| - 0.014)))$ $(0.313x_1 + 0.362(0.706$ $-0.561x_1)(x_1 + 2.798) - 0.513)$
SeTGAP	$-1.0\log(14.156x_0^2 + 3.539) +$ $1.0|\log(9.499x_1 + 4.75)| - 0.294$	$1.0\sin(1.0x_0e^{x_1})$	$2.0x_0\log(x_1^2)$	$1.0x_0\sin\left(\frac{1}{x_1}\right) + 1.0$	$\sqrt{1.0x_0}\log(x_1^2)$

Table 16: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 5

Method	E1	E2	E3	E4
PySR	$0.608x_0x_1 + 1.1\sin(2.25x_0(x_1 - 0.667) - 1.5x_1 + 1.0)$	$-0.5x_0 + (0.179x_1 + 2.984)\sin(0.2x_2) + e^{e^{\cos(\cos(0.111x_0))}} + 1.041$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$(\cosh(x_0) + \cosh(x_2))(-0.005x_1 - 0.005x_3 + 0.091)$
TaylorGP	$0.616x_0x_1$	$-0.494x_0 - 0.005x_1 + 0.204x_2 + 8.593$	$\dfrac{-x_0 + e^{x_0} -}{\sqrt{\left|x_0 + 0.838\sqrt{e^{x_0}} - e^{x_0} - \sqrt{\|x_0\|}\right|}}$	$\frac{x_2}{\tanh(x_2)} + \log\left(\left\|\frac{x_0}{\tanh(x_0)}\right\|\right)\sqrt{\|x_0\|} - 1.797$
NeSymReS	$0.59x_0x_1 + \cos(0.584(0.989 x_0 - x_1)^2)$	$-x_0 + e^{\sin(x_2)} + 7.456 - 0.05e^{-0.102x_1}$	$0.585e^{x_0} - 0.437\sin\left(1.612(0.04x_1 + 1)^2\right)$	—
E2E	$0.021x_0(29.332x_1 - 0.194) + 1.1\cos(30.949 x_1 - 0.049) + 0.01$	$-0.016x_0 + 0.049x_2 + (0.086x_0 - 0.704)(0.75x_0 - 0.015) + (0.003x_1 + 0.052)(-0.97x_1 + 53.2\sin(0.221x_2 + 0.035) + 0.064) + 6.444$	$0.224e^{0.188x_0}$ $e^{(0.004x_0 + 0.045)(22.712x_0 + 0.009x_1 - 5.15)}$ $+ 0.514\cos(3.152x_1 + 0.082) + 0.043$	$0.002\|30.73x_3 + 5.334(x_0 + 0.019)^4 + 6.811(x_1 - 0.94(x_2 - 0.021)^2 + 0.028)^2 + 1.185\|$
TPSR	$-0.6\log(12.904(x_0 + 0.008)^2 + 0.6) + 1.9 - \frac{0.6}{0.227x_1 + 0.236}$	$(0.372x_0 - 0.368x_2 + 123.99)$ $((0.001x_0 + 0.127)(x_0 - 0.045x_2 - 254.229) + 0.086$ $\arctan(0.002x_2(2.01x_1 + 71.49)) + 32.38)$	$0.052e^{2.5x_0} + 0.5\cos(4.0x_1 - 0.03) + 0.09$	$-0.13x_1 + 0.032x_3 + 0.081(0.067x_1 + 0.005x_2 + 0.363x_3 - (1 - 0.032 \|7.087x_0 + 28.484\|)^2 - 0.334 (x_2 + 0.005)^2 - 0.028)^2 + 0.319$
SeTGAP	$0.607x_0x_1 - 1.1\sin((2.252x_0 - 1.492)(x_1 - 0.666) + 9.432)$	$-0.5x_0 + 0.063x_0^2 + 3.16\sqrt{0.1x_1 + 1}\sin(0.202x_2) + 6.499$	$0.15e^{1.5x_0} + 0.5\sin(3.0x_1 + 1.571)$	$-0.02x_0^2x_1 + 0.001x_0^2 + 0.01x_0^4 + 0.01x_1^2 - 0.02x_2^2x_3 + 0.01x_4^2 + 0.01(-x_3)^2 - 0.001$

Table 17: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 5

Method	E5	E6	E7	E8
PySR	$e^{1.2x_3} + \sin(x_0 + x_1x_2)$	$\dfrac{0.52\cos(0.2x_2^2) + \tanh(x_0)}{\tan\left(\cos\left(\tanh\left(\frac{1.9}{x_1}\right)\right) + 0.534\right)}$	$4.26x_1^2(\tanh(\sin(6.283 x_0) + 1.8) - 1.1) + 0.967$	$\sinh(0.406\cosh(\tanh(x_0 \sinh(\tanh(\sinh(x_0)))))) + \tanh(\cosh(0.895x_1)) + 0.69) - 3.031$
TaylorGP	$e^{x_3}\|x_3\|^{0.5} + 0.487$	$\frac{x_1\sin(x_2)}{x_2\tanh(x_1)} - 1.185\cos\left(e^{\|x_2\|^{0.5}}\right)$	$-x_1^2 + \sqrt{\|x_1\|}$	$1.07\|x_0x_1\|^{\frac{1}{4}}$
NeSymReS	—	$-0.376x_0 + x_1\sin\left(\frac{x_0}{x_1}\right)$	$\frac{-3967.593x_0 + x_1^2}{\cos(65757.558(-x_1 - 0.157)^3) + 28400.518}$	$\cos(\frac{\sin(\frac{x_0}{x_1})}{x_0}) + 0.653$
E2E	$0.903e^{1.231x_3} + 0.105 + 1.0\cos(0.764/(0.062 \sin((-0.619x_0 - 0.079)(0.344x_1 - 0.212x_2 - 0.857)) - 0.007))$	$(-0.001\|4.547x_1 - 0.068\| - 0.601)$ $\arctan(8.32x_0 - 242.537(0.016x_0 + 1)^3 + 7.105) + 0.005\cos(0.667 (-x_2 - 0.162)^2 - 4.352) - 0.004$	$(0.09 - 0.083(x_1 - 0.002)^2)(6.724(1 - 0.756 \sin(6.11x_0 - 0.012))^2 + 4.0)$	$0.536 - 0.903\arctan((0.046 x_1 - 0.991)(-0.367\sin(2.487 \sqrt{(-\|32.358x_1 + 1.7\| - 0.017)^2 + 0.006} - 49.4) + 0.017\|(0.053(x_0 - 0.007)^2 + 0.016)(0.001x_1 + 425.696 (x_1 + 0.011)^2 - 22.805)\| - 0.286))$
TPSR	$0.97e^{1.211x_3} + 0.087\cos(x_1 + 3.732x_2 + 10.665) + 0.027$	$(0.007x_1 + 0.004)(2.087x_1 + 1.792) + 3.762\sin(0.0044x_0 + 2.3489 \cos(0.523x_2 + 0.03) - 0.73097) + 1.077$	$1.075 - 0.971(-0.022x_0 - x_1 + 0.096\cos(x_0 + 1.524) + 0.078)^2$	$(-0.004x_1 - 0.001) (x_1 + 0.002) + 2.063 - 1.494e^{-0.001}(5.976x_0 - 0.065)(59.997x_1 + 2.475)\|$
SeTGAP	$0.977e^{1.201x_3} + 1.0\sin(x_0 + x_1x_2 + 18.85) + 0.001$	$1.0\cos(0.2x_2^2 + 6.282)\|x_1\| + 1.0\tanh(0.5x_0)$	$\dfrac{(0.947 - 0.948x_1^2)}{((\sin(6.283x_0) + 1.478)^{-1.0} + 0.025)} + 0.02$	$2.0 + \frac{15.997}{(0.026x_1^3 - 15.998x_1^4 - 15.994)} - \frac{16.587}{(-0.001x_0 + 0.077x_0^2 + 0.012x_0^3 + 16.459x_0^4 + 16.583)}$

Table 18: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 5

Method	E9	E10	E11	E12	E13
PySR	$\sqrt{1.148x_1} + \cosh(1.354\cos(x_0) \cos(\tanh(0.652x_0))) - 2.202\cosh(1.613 \tanh(0.481x_0))$	$\sin(x_0e^{x_1})$	$x_0(0.943 - 2.768\log((\log(\cosh(1.303 \tanh(0.652x_1))))^{0.5})) + 3.385x_0 \log(\log(\cosh(x_1)))$	$1.0x_0\sin\left(\frac{1.0}{x_1}\right) + 1.0$	$\sqrt{x_0}\log(x_1^2)$
TaylorGP	$\dfrac{-\log(\|x_0\|) + \tanh(\log(\|x_1\|)) -}{\sqrt{\left\|\log\left(\frac{0.603}{\|x_0\|}\right)\right\|}}$	$\dfrac{\tanh(\sin(e^{x_1}\sin(x_0)))}{\sqrt{\|\cos(\sin(x_1))\|)}}$	$3.947x_0\log(\|x_1\|)$	$\frac{1.217x_0}{1.229x_1 + \frac{0.471}{x_1}} + 0.961$	$2\log(\|x_1\|)\sqrt{\|x_0\|}$
NeSymReS	$\log\left(\frac{0.28\|x_1\|}{\|x_0\|}\right)$	$\sin(x_0e^{x_1})$	$x_0\log(x_1^4)$	$(x_0 + x_1)\sin\left(\frac{1}{x_1}\right)$	$0.252x_0 + \log(x_1^2)$
E2E	$-0.6\log(12.904(x_0 + 0.008)^2 + 0.6) + 1.9 - \frac{0.6}{0.227x_1 + 0.236}$	$0.001 - 1.0\sin(0.257x_0 (0.1 - 3.597e^{1.08x_1}))$	$0.257x_0(30.0\log(0.19 (\|21.905x_1 + 0.102\| + 0.001)^{0.5} - 0.01) + 0.01)$	1	$(0.031 - 0.44\log(23.696 (-x_1 - 0.009)^2(x_1 + 0.201)^2 (0.033x_0 + 0.086\|0.501x_0 - 7.823\| - 1)^2 + 0.02)) (-0.001x_0 - 8.705) (0.032x_0 + 0.057)$
TPSR	$(0.187 - 0.012x_1)(0.179x_1 - 2.902) ((-2.035 + \frac{4.906}{0.282\|x_0 + 0.001\| + 1.048}) (-0.655x_1 - 3.76) - 0.296\|x_0\| + 8.821)$	$1.0\sin(1.0x_0e^{x_1})$	$(-0.318x_0 - 0.004) (-0.392x_1^2 - 15.221 - \frac{30.882}{-0.017\|98.801x_1 + 0.806\| - 0.484})$	$-10.226x_0 + 0.038x_1 + (x_0 + 4639.596 + \frac{371.436}{\arctan(440.318x_1 + 15.611)}) (0.006x_0(-0.002x_1 - 141.195) + 0.796x_0 - 0.001) + 6.082$	$(-0.029 + 19.719/(0.422 \|(0.45x_0 + 23.971)(4.248x_1 + 0.008)\| + 7.133)(3.329 - 3.344(x_1 + 0.002)^2) (-0.207x_0 - 0.669)$
SeTGAP	$-1.001\log(18.044x_0^2 + 4.522) + 0.994\|\log(18.83x_1 + 9.24)\| - 0.705$	$1.0\sin(1.0x_0e^{x_1})$	$2.0x_0\log(x_1^2)$	$1.0x_0\sin\left(\frac{1}{x_1}\right) + 1.0$	$1.0\sqrt{x_0}\log(x_1^2)$

Table 19: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 6

Method	E1	E2	E3	E4
PySR	$0.607x_0x_1$	$-0.499x_0 + e^{\tanh(e^{x_1})}\tanh(x_2) +$ $\log(\cosh(x_0) + 29.144) + 3.287$	$0.15e^{1.5x_0} + 0.5\cos(3x_1)$	$(0.09 - 0.009x_1)\cosh(x_0) +$ $(0.091 - 0.009x_3)\cosh(x_2)$
TaylorGP	$0.62x_0x_1$	$-0.497x_0 - 0.002x_1 + 0.208x_2 + 8.571$	$-x_0 + e^{x_0} - \sqrt{|x_0|} - 0.506 +$ $\tanh(x_0 - e^{x_0} - 0.977|$ $+0.977)$	$\sqrt{\log(|x_0|)}\sqrt{\left\|\frac{x_2\left(-x_2\sqrt{|x_0|}+\log(|x_2|)+0.553\right)}{\tanh(x_2)}\right\|}$
NeSymReS	$0.667x_0x_1 + \cos(0.066$ $(-x_0 - 0.869x_1)^2)$	$93.845e^{0.002x_2} + e^{\sin\left(\frac{x_0}{x_1}\right)} - 86.601$	$0.58e^{x_0} -$ $0.464\sin\left(1.482\left(-0.071x_1 - 1\right)^2\right)$	—
E2E	$0.024x_1(25.373x_0$ $-0.001x_1 + 0.057) -$ $1.1\cos(76.2\tan(2.983$ $x_1 - 4.272) + 0.767)$ $+0.002$	$(35.4 - 0.001x_0)\,(0.022(0.299x_0 - 1)^2$ $+(0.013x_1 + 0.301)(0.319\sin((0.145x_1$ $+3.599)(0.052x_2 - 0.004)) + 0.061)$ $+0.163)$	$0.212e^{1.379x_0} +$ $0.475\cos(3.108x_1 - 0.004) - 0.013$	$0.018(0.573x_1 - (0.033x_0 - 0.002)(21.569x_0$ $-1.005) + 0.036)^2 + 2.72|(0.01\arctan(-$ $(0.079e^{\frac{0.005}{18.338x_2 - 4.735}} - 199.276)e^{-\frac{0.005}{18.338x_2 - 4.735}}$ $+0.001)((1.851x_2 - 2.372)(0.283x_3$ $-0.03) + 0.25((x_2 + 0.005)^2 - 0.003)^2$ $+0.075)|+0.001$
TPSR	$0.074x_1(8.206x_0 +$ $0.136) + 0.043\sin(4.77$ $x_1 - 0.485) + 0.001$	$0.037x_1 - 0.134x_2 + 0.766|9.705x_0 +$ $93.72 + \frac{94.187}{9.425e^{0.022x_0 + 0.035x_1 - 0.524x_2}+9.862}|$ $-1026.309 + \frac{117.7}{0.001x_0 + 0.123}$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$0.003(-0.005x_2 + (0.363 - 0.003x_3))$ $((-0.941x_2 - 1.086)(-1.089x_1 + x_2 + 0.284) -$ $0.001(x_0(0.52x_3 + 69.999) + 0.001x_0 +$ $0.677)^2 - 0.238) - (1 - 0.025x_3)^2(0.148x_1$ $+x_2 - 0.552)^2 - 0.001)^2 + 0.524$
SeTGAP	$0.608x_0x_1 - 1.099$ $\sin((2.24x_0 - 1.544)$ $(x_1 - 0.67) - 9.46)$	$-0.5x_0 + 0.063x_0^2 +$ $3.162\sqrt{0.1x_1 + 1}\sin(0.2x_2) + 6.498$	$0.149e^{1.502x_0} -$ $0.5\sin(3.0x_1 + 4.712)$	$-0.004x_0 - 0.02x_0^2x_1 + 0.017(-x_0)^2 +$ $0.009(-x_0)^4 + 0.008(-x_1)^2 +$ $0.01x_2^4 + 0.011x_3^2 - (0.013x_2^2 + 0.003)$ $(1.575x_3 + 0.614) - 0.023$

Table 20: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 6

Method	E5	E6	E7	E8
PySR	$e^{1.2x_3} + \sin(x_0 + x_1x_2)$	$2.535e^{\tan\left(1.058\cos\left(2.528\tanh\left(\frac{2}{x_1}\right)\right)\right)}$ $\cos(0.2x_2^2) + \tanh(x_0)$	$\frac{1.0 - 1.0x_1^2}{\sin(6.283x_0)+1.5}$	$\sinh(\sinh(\sinh(\tanh(\sinh(\cosh($ $\tanh(x_0\tanh(x_0)))$ $\cosh(\tanh(x_1\tanh(x_1))) - 0.718)))))$
TaylorGP	$e^{x_3}|x_3|^{0.5} + \tanh((x_3 -$ $0.891)e^{x_3} + 1.718e^{x_3})$	$-1.185\cos\left(e^{|x_2|^{0.5}}\right) + \frac{4.854\sin(x_2)}{x_2}$	$-x_1^2 +$ $\log(|x_1^2 - \log(|x_2^2 -$ $\sqrt{|x_1(x_1 - 0.073)|}|)|)$	$1.102|x_0x_1|^{\frac{1}{4}}$
NeSymReS	—	$-0.368x_0 + x_1\sin\left(\frac{x_0}{x_1} - 0.001x_2\right)$	$\frac{0.184x_0 + x_1^2}{\cos(1.965x_0 - x_1) - 1.94}$	$\cos\left(\frac{\sin\left(\frac{x_0}{x_1}\right)}{x_0}\right) + 0.643$
E2E	$0.002x_2 + 1.106e^{1.147x_3} +$ $0.774\cos(0.002x_1) + 0.942$ $\cos(28.703x_2 - 0.956) - 0.977$	0.942	$\left(6.284(x_1 - 0.007)^2 - 7.49\right)$ $(-0.082(0.01x_1 + \sin(6.457x_0 +$ $0.14) - 0.705)^2 - 0.098)$	$2.06 - 30.5/(6.892(0.047 - x_0)^2$ $(x_1 - 0.003)^2 + 18.24((1 - 0.006x_0)^2)^2$ $-9.27\cos(1.329x_1 - 44.203) + 10.058)$
TPSR	$0.98e^{1.207x_3} - 0.039\sin(-x_1 +$ $3.302x_2 + 18.63) + 0.022$	$(-0.068x_0 - 1.576\cos(0.538x_2)$ $-0.674)(0.016803x_0 + 0.001x_1 +$ $4.731\sin(0.015(0.326 - x_1)^2$ $+3.728) + 2.043)$	$\frac{412.541 - 411.968}{\left(0.001(x_1 + 0.002)^2 + 1\right)^2}$	$(0.001 - 0.003x_1)(x_1 - 0.002) + 2.057 -$ $1.494e^{-0.001|(5.993x_0 + 0.026)(59.999x_1 - 1.554)|}$
SeTGAP	$0.995e^{1.201x_3} + 1.0\sin(x_0 +$ $x_1x_2 - 6.283) + 0.002$	$0.999\cos(0.2x_2^2 - 0.011)|x_1| +$ $1.0\tanh(0.5x_0)$	$\left(-0.001 + \frac{0.999}{\sin(6.283x_0 - 9.424) - 1.5}\right)$ $(x_1^2 - 0.977) + 0.018$	$-\frac{16.085}{(16.038x_1^4 + 2.016.09)} + 2.0 +$ $\frac{7.918}{\left(0.158x_0^2 - 8.067x_0^4 - 7.943\right)}$

Table 21: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 6

Method	E9	E10	E11	E12	E13
PySR	$\sinh(\sinh(2.7$ $e^{-\frac{0.259x_0^2}{x_1 + 0.727} - \frac{0.178x_0}{(x_1 + 1.017)\tanh(x_0)}}$ $-1.154)) - 1.787$	$\sin(1.0x_0e^{x_1})$	$2x_0\log(x_1^2)$	$x_0\sin\left(\frac{1}{x_1}\right) + 1.0$	$\sqrt{x_0}\log(x_1^2)$
TaylorGP	$-0.683\log(|x_0|) -$ $1.618\sqrt{|x_0|} + \sqrt{|x_1|}$	$\sin(x_0e^{x_1})$	$2.458x_0\log(|x_1|) +$ $(1.632x_0 - 0.908)\log(|x_1|) +$ $0.875\cos(3.872x_0 + (2.732$ $x_0 - 2.481)\log(|0.754x_1$ $\tanh(x_1)|x_0|^{1.5} + 0.307|) +$ $(4.458x_0 - 2.481)\log(|x_1|))$	$0.806x_0\sin\left(\frac{1.229}{x_1}\right)$ $+0.806$	$(\log(|x_0|)\log(|x_1|) + \tanh(\log(|x_1|)))$ $(\log(|x_0\tanh(\log(|x_1|)) + \log(|x_1|) -$ $0.13|^{0.5}) + |\tanh(x_0\tanh($ $\log(6.162|\log(|x_1|)|)))|^{0.5})$
NeSymReS	$\log\left(\frac{0.281|x_1|}{|x_0|}\right)$	$\sin(x_0e^{x_1})$	$x_0\log(x_1^4)$	$(x_0 + x_1)\sin\left(\frac{1}{x_1}\right)$	$0.245x_0 + \log(x_1^2)$
E2E	$2.964(0.2x_1 - 1)^3 + 4.352 -$ $0.792\log(166.831$ $(x_0 - 0.052)^2 + 17.6)$	$1.0\sin(0.005x_0$ $(167.868e^{1.143x_1} -$ $0.01)) + 0.001$	$(2.024x_0 - 0.006)(0.75$ $\log(1.946(-x_1 - 0.005)^2$ $+0.05) + 0.068)$	$7.12\sin(0.001x_0$ $(0.003 + \frac{31.2}{0.173x_1 + 0.001}))$ $+0.996$	$(0.004 - 5.12\log(0.175x_0 + 1.109))$ $(0.945((|0.004 - \frac{49.1}{0.326x_1 - 0.001}| +$ $0.045)^{0.5} - 0.898)^{0.5} - 3.44)$
TPSR	$\left(-0.01 + \frac{0.001}{x_0 + 11.073}\right)(3.527 -$ $0.23x_1)(x_1 - 15.288)((3.097$ $-6.725/(0.000341|(x_0 - 0.003)$ $(0.02x_1 + 0.399)(0.954x_1 +$ $19.312)|+0.016)(-0.005x_1 -$ $0.063) + 0.665|x_0|-25.78)$	$1.0\sin(0.999x_0e^{x_1})$	$(26.232 - 0.008x_1)$ $(-0.523x_0 - 0.01)$	$2.867x_0 + 1.066$	$(0.023 - \frac{0.25}{0.829|x_1 - 0.001| + 0.087})$ $(0.948 - 0.97x_1)(43.582 - 0.84x_0)$ $(0.055x_0 + 0.137)(0.494x_1 + 0.482)$
SeTGAP	$-1.0\log(11.241x_0^2 + 2.809) +$ $1.0|\log(9.824x_1 + 4.914)|$ -0.559	$1.0\sin(1.0x_0e^{x_1})$	$2.0x_0\log(x_1^2)$	$1.0x_0\sin\left(\frac{1}{x_1}\right) + 1.0$	$2.0\sqrt{x_0}\log(|x_1|)$

Table 22: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 7

Method	E1	E2	E3	E4
PySR	$0.608x_0x_1 +$ $1.1\sin((x_0 - 0.667)$ $(x_1 - 0.667))$	$3.017\sin(0.2x_2) +$ $3.017\cosh(0.162x_0 - 0.723) + 2.818$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$-0.02x_0^2x_1 - 0.02x_2^2x_3 +$ $0.091\cosh(x_0) +$ $0.091\cosh(x_2)$
TaylorGP	$0.607x_0x_1$	$-0.499x_0 - 0.008x_1 + 0.207x_2 + 8.586$	$-2x_0 + e^{x_0} - 0.363$	$0.756x_2\tanh(x_2) +$ $0.756\log(|x_0|)\sqrt{|x_0|}$
NeSymReS	$0.61x_0x_1 - \sin(0.001$ $(0.965x_0 - 1)^2)$	$96.368e^{0.002x_2} + e^{\sin(x_0x_1)} - 89.13$	$0.579e^{x_0} -$ $0.437\sin\left(1.633\left(0.043x_1 + 1\right)^2\right)$	—
E2E	$0.598x_0x_1 - 0.004x_0$ $+0.001x_1 + 0.143$	$(0.011x_2 + 0.445)(0.174x_1 - 0.051x_2$ $+5.52)(0.387(0.235x_0 - 1)^2 + \sin($ $0.225x_2 + 0.023) + 0.144) + 4.88$	$((7.16(0.001x_0 - 1)^2 + 0.01)$ $(0.198e^{1.42x_0} + 0.519\cos(3.037x_1$ $-0.034 - 0.072) + 0.057)/$ $(7.16(0.001x_0 - 1)^2 + 0.01)$	$0.002|5.04((x_0 + 0.033)^2 -$ $0.001)^2 + 5.21(0.818x_1 +$ $0.67x_3 - (x_2 + 0.025)^2 +$ $0.01)^2 - 0.11| + 0.001$
TPSR	$0.011x_1(56.515x_0$ $+1.528) - 0.031\sin($ $4.588x_1 + 0.776)$ -0.005	$(2.621(0.060401x_0 - 0.019x_2 - 1.0 -$ $\frac{0.1813}{-0.544e^{-2.096x_1 + 6.406x_2 - 1.507}})^2 + 0.508)$ $(0.073x_0 + 0.011x_1 - 0.028x_2 + 2.346)$	$0.15e^{1.5x_0} +$ $0.5\cos(3.0x_1)$	$0.198(-x_2 - 0.027)^2 +$ $0.127(0.236x_1 - 0.234(0.006$ $-x_0)^2 - 1)^2 - 0.662$
SeTGAP	$0.607x_0x_1 +$ $1.098\sin((2.24$ $x_0 - 1.544)(x_1 -$ $0.67)) - 0.002$	$-0.501x_0 + 0.064x_0^2 + (3.166$ $(0.1x_1 + 1)^{0.5} + 0.001)$ $\sin(0.201x_2) + 6.452$	$0.151e^{1.498x_0} -$ $0.5\sin(3.001x_1 + 4.712)$	$-0.02x_0^2x_1 + 0.01x_4^4 - 0.021$ $x_3(x_2)^2 + 0.005x_3 + 0.01(x_0)^4$ $+0.01(x_1)^2 - (0.004x_0 - 0.708)$ $(0.014x_3^2 - 0.019) + 0.028$

Table 23: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 7

Method	E5	E6	E7	E8
PySR	$e^{1.2x_3} + \sin(x_0 + x_1x_2)$	$x_1\cos(0.2x_2^2)\tanh(236.582x_1) +$ $\tanh(0.496x_0)$	$(1.863x_1^2 - 1.843)(\tanh(\sinh($ $\sin(6.281x_0)) + 1.345) - 1.215$	$1.644\log(0.976/(\cos(\tanh(1.094x_0$ $\tanh(\sinh(\sinh(x_0)))))$ $\cos(\tanh(0.812\sinh(x_1)))))$
TaylorGP	$2e^{x_3} - \log(|-3e^{x_3} + \log(|$ $2.287e^{x_3} - 1.447\log(|\sin(x_3$ $)|)|) + 1.447\log(|\sin(x_3)|)|)$	$\frac{x_1\sin(x_2)}{x_2\tanh(x_1)} + \cos(x_2) + \tanh(x_0)$	$-x_1^2 + \log\left(\left|x_1^2 - \log\left(\left|x_1^2 - 0.828\sqrt{|x_1^2|}\right|\right)\right|\right)$	$\tanh((x_0 + 0.504)\tanh(x_0)) +$ $\tanh(x_1^2 - 0.017)$
NeSymReS	—	$-0.356x_0 + x_1\sin\left(\frac{x_0}{x_1} - 0.009x_2\right)$	$\frac{0.08x_0 + x_1^2}{\cos(2.445(0.924x_1 - 1)^2) - 1.916}$	$\cos\left(\frac{\sin\left(\frac{x_0}{x_1}\right)}{x_0}\right) + 0.695$
E2E	$1.138e^{1.163x_3} - 0.123 + 0.953$ $\cos(22.045x_2 - 14.032 + 1.11/$ $(-0.742x_0 + 0.002x_1 + 19.256) +$ $2.11/((31.854x_2 + 0.793)(0.001x_3$ $+2.26)) - 0.033/(3.382x_2 + 0.147))$	0.806	$(0.368 - \frac{4.78}{8.105 - 0.291x_1})(0.338x_1 + 23.594)$ $(0.348x_1 - 4.637)(0.136(0.019 - x_1)^2$ $-0.117)(-0.219(1 - 0.329$ $\cos(6.186x_0 - 7.784))^3 - 0.046)$	$2.1 - 0.9/(0.03(0.309x_0 + 0.074)$ $(0.687x_0 - 0.059)(2.086x_1 - 0.139)$ $(17.421x_1 - 0.429)| + 0.6)$
TPSR	63.518 $e^{(0.5 - 0.15x_3)(-0.068\cos(x_1 + 1.23x_2 + 1) - 8.195)}$ -0.024	$(-0.396\cos(0.876x_2 - 0.03)$ $-0.167)((-0.016x_2 - 0.304)(x_0 -$ $13.652x_2 + 269.341) + \cos(2.997$ $x_1 + 0.599) + 70.235)$	$-0.09x_0 - 0.01x_1$ $(90.02x_1 + 6.533) + 0.94$	$2.011 -$ $1.427e^{-0.0039|x_1(98.999x_0 + 0.009)|}$
SeTGAP	$0.999e^{1.201x_3} + 1.0\sin(x_0 +$ $x_1x_2 - 12.566) + 0.003$	$1.0\cos(0.2x_2^2 - 6.275)|x_1| +$ $1.0\tanh(0.501x_0)$	$\frac{2.907x_1^2 - 2.919}{2.904\sin(6.283x_0 + 3.148) - 4.358}$	$2.0 - \frac{14.033}{13.947x_1^4 - 0.122x_0^3 + 14.032} - \frac{14.438}{14.532x_1^4 + 14.434}$

Table 24: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 7

Method	E9	E10	E11	E12	E13
PySR	$0.226\cos(\cos(x_0)) + \sinh(\sinh($ $\sinh(0.974\cos(0.643x_0) + 0.167)))$ $-\frac{8.714}{x_1 + 2.537}$	$\sin(1.0x_0e^{x_1})$	$2.0x_0\log(x_1^2)$	$x_0\sin\left(\frac{1}{x_1}\right) + 1$	$\sqrt{x_0}\log(x_1^2)$
TaylorGP	$0.26x_1 + \log\left(\frac{0.952}{|x_0|}\right) -$ $\sqrt{\left|\log\left(\frac{0.593}{|x_0|}\right) + \log\left(\left|\tanh\left(\frac{x_1}{x_0}\right)\right|\right)\right|}$	$\sin(x_0e^{x_1})$	$4x_0\log(|x_1|)$	$\frac{x_0\tanh(x_1^2)}{x_1} + 1$	$2\log(|x_1|)\sqrt{|x_0|}$
NeSymReS	$\log\left(\frac{0.274|x_1|}{|x_0|}\right)$	$\sin(x_0e^{x_1})$	$x_0\log(x_1^4)$	$(x_0 + x_1)\sin\left(\frac{1}{x_1}\right)$	$0.23x_0 + \log(x_1^2)$
E2E	$2.0 - 0.6\log(6.862(1 -$ $\frac{0.129}{-0.056x_1 - 0.02})^2(x_0 - 0.009)^2 + 0.8)$	$-0.998\sin((0.214 -$ $33.248x_0)(0.026$ $e^{1.125x_1} + 0.001))$	$0.241x_0(31.0$ $\log(0.19(|25.419x_1 +$ $0.12| + 0.011)^{0.5}$ $-0.01) + 0.5)$	$(0.629 - 4.58\sin((-0.007 +$ $\frac{9.78}{0.014 - 3.043x_1})(0.042x_0 +$ $0.001)))(1.581 - 0.006x_0)$	$(-0.052 + \frac{0.004}{-0.019 - \frac{4.486}{5.98x_0 - 68.5}})$ $(48.802\log(0.012(0.001 - 6.156/$ $((0.243x_1 - 0.002)(0.806\sin($ $0.085x_0 - 0.168) + 0.054))|$ $+0.025) + 0.04)$
TPSR	$0.612x_1 + (0.995 - 1.013\arctan(x_1$ $+0.926))(0.808x_1 - 1.554) - 1.0$ $\log(51.139x_0^2 + 12.793) + 2.955$	$0.9998\sin(1.0x_0e^{x_1})$	$-0.03x_0(0.196x_1 -$ $261.804 + 40.943/(0.07$ $|1.488x_1 + 0.003| + 0.058))$	$-0.772x_0 + (10.323 +$ $\frac{1}{x_0 - 0.027})(0.029\arctan($ $15.95x_1 - 5.238) + 0.059)$ $(x_0 + (0.128 - 0.034x_1)$ $(1.088 - 0.098x_1)(4.661x_0 -$ $0.086) - 0.027) + 0.946$	$(-1.365 +$ $\frac{1.705}{0.0007x_0 - 0.55113(x_1 + 0.006)^2 + 0.376})$ $(-0.457x_0 - 2.791)$
SeTGAP	$-1.0\log(2.785x_0^2 + 0.696) +$ $1.0|\log(13.845x_1 + 6.92)| - 2.296$	$1.0\sin(1.0x_0e^{x_1})$	$4.0x_0\log(|x_1|)$	$1.0x_0\sin\left(\frac{1}{x_1}\right) + 1.0$	$2.0\sqrt{x_0}\log(|x_1|)$

Table 25: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 8

Method	E1	E2	E3	E4
PySR	$0.609x_0x_1 +$ $0.931\cos(1.488x_0 - 1.126x_1$ $(2x_0 - 3.019) - 1.9x_1 + 0.556)$	$\dfrac{2.278\tanh(0.356x_2) + 6.388}{\sqrt{(0.016x_0^2 + e^{-0.402x_0})^{0.5} + 0.127\tanh(x_1x_2)}}$	$0.15e^{1.499x_0} -$ $0.502\cos(2.999x_1 + 3.143)$	$(\cosh(x_0) + \cosh(x_2))$ $(-0.005x_1 - 0.005x_3 + 0.091)$
TaylorGP	$0.613x_0x_1$	$-0.506x_0 - 0.001x_1 + 0.206x_2 + 8.575$	$-0.612\left(e^{x_0}e^{2\tanh(x_0)}\right)^{0.5} + e^{x_0}$	$\left(-\sin(x_0) + \sqrt{|x_2|}\right)e^{\tanh\left(\log(\sqrt{|x_0||x_2|})\right)}$
NeSymReS	$0.666x_0x_1 +$ $\cos(0.068(-x_0 - 0.83x_1)^2)$	$e^{\sin(x_0x_1)} + 124.329 - 116.894e^{-0.002x_2}$	$0.579e^{x_0} - 0.444\cos(0.228x_1)$	—
E2E	$(0.001x_0 - 1.463)\sin(0.482$ $\cos(\frac{0.121x_0 + 78.526}{0.007x_0 + 0.29}) - 0.004)$ $+(3.222x_0 + 0.007)$ $(0.186x_1 + 0.003)$	$(0.01x_0 - 0.099)(5.434x_0 + 0.438) + 6.3 +$ $3.0\sin((0.001x_1 + 0.02)(10.39x_2 + 0.291))$	$0.143e^{1.497x_0} + 0.043 +$ $0.506\cos(3.559x_1 + 0.061)$	$0.079|0.162(-x_1 + 0.885(x_0 + 0.05)^2$ $+0.031)^2 + 0.123(0.135x_0x_1 + 0.004x_0$ $+0.011x_1 - x_2^2 - 0.022x_2 + 0.884x_3$ $+0.024)^2 + 0.523| - 0.048$
TPSR	$-2.937x_1(0.001 - 0.208x_0) +$ $0.04\sin(4.355x_1 - 0.296)$ -0.002	$(-20.407 - \frac{0.002}{x_1 - 0.182})(-0.01x_2 - 1.038 + 1/$ $(0.014(0.201x_0 + 0.007x_2 - 1)^2(0.442x_0 +$ $0.001x_2 - 1)^2 + 1.342))$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$-0.00034(0.347x_1 - 7.955)(x_3 - (90.12$ $-0.35x_3)(0.111x_1 - 2.6) - 1.987) + 0.01$ $(-0.001x_1 + 0.995x_2^2 - x_3 + 0.062)^2 - 0.621$
SeTGAP	$0.608x_0x_1 +$ $1.099\sin((2.251$ $x_0 - 1.52)(x_1 - 0.655))$	$-0.5x_0 + 0.062x_0^2 + 6.541 +$ $\sqrt{0.1x_1 + 1}(3.166\sin(0.201x_2) - 0.033)$	$0.15e^{1.5x_0} + 0.5\cos(3.001x_1)$	$-0.005x_0^2 + 0.01x_0^4 - 0.02x_1(-x_0)^2 +$ $0.01x_1^2 + 0.01x_2^4 - 0.02x_3(-x_2)^2$ $+0.01x_3^2 + 0.015$

Table 26: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 8

Method	E5	E6	E7	E8
PySR	$e^{1.2x_3} +$ $\sin(x_0 + x_1x_2)$	$0.279e^{3.915\cos\left(\frac{\tanh(1.761\tanh(\sin(x_1)))}{x_1}\right)}$ $\cdots\cos\left(\tanh\left(\frac{4.667}{x_1}\right)\right)$ $\cos(0.2x_2^2) + \tanh(x_0)$	$\cos(x_1) + \sinh(\sin(6.285x_0) +$ $\cos(0.35x_1) + \cos(0.668x_1) - 2.288)$	$\log(0.705\sinh(1.975$ $\cosh(\tanh(x_0)) - \frac{1.184}{\sinh(\cosh(x_1))}))$
TaylorGP	$0.855e^{x_3}\log(|2x_3 +$ $\sqrt{|\tanh(e^{2x_3})|} + 0.748|)$	$\cos(x_2) + 0.422 +$ $\frac{5.65\log(|x_1|^{0.5})\sin(x_2)}{x_2}$	$-x_1^2 + \log(|x_1^2 -$ $\log(|x_1^2 - \log(|x_1^2 + 0.943|)|)|)$	$\frac{0.829|x_0|^{\frac{1}{4}}}{\cos(\tanh(x_1))}$
NeSymReS	—	$0.126x_0 + x_1\sin\left(0.294x_1 + \frac{x_2}{x_1}\right)$	$\frac{0.042x_0 + x_1^2}{\cos(5.419(x_1 - 0.517)^2) - 1.896}$	$\cos\left(\frac{\sin\left(\frac{x_0}{x_1}\right)}{x_0}\right) + 0.644$
E2E	$1.392e^{1.085x_3} + 0.949\sin($ $-25.875x_2 + 100.707 +$ $\frac{0.007}{0.055 - 0.1x_1}) - 0.484$	$(6.048|0.168x_1 + 0.001| + 0.14)$ $\cos((0.21 - 17.355x_2)(-0.012x_2$ $-0.01)) + 0.8\arctan(0.699x_0 +$ $0.269) - 0.01$	$(-10.369(1 - 0.002x_0)^2(0.346x_1 +$ $(0.01 - 0.003x_1)(0.029x_0 - 0.003) +$ $0.018)^2 + 1.7)(0.373(\sin($ $6.398x_0 + 0.342) - 0.814)^2 + 0.42)$	$2.065(-1 - 0.556/(-0.361(x_0 + 0.028)^2$ $(0.001x_1 - 1)^2(x_1 + 0.046)^2 + 0.409\cos($ $2.271x_1 + 0.076) - 1.369))^2 + 0.001$
TPSR	$0.983e^{-0.001x_2 + 1.206x_3}$ $e^{0.001\arctan((-59.047x_0}$ $\cdots -117.779)(-70.903x_1 -$ $\cdots 32.111x_3 + 8.033)) + 0.01$	$(0.003\cos(0.044x_0 + 0.023x_1^2 - 2.266)$ $+0.003)(5.010^{-6}x_0 - 0.005x_1 - 465.626$ $(0.025 - \arctan(-0.473 -$ $\frac{9.002}{-0.668(x_2 - 0.082)^2 - 0.109}))^3 - 0.002)$	$1.045 -$ $0.926(0.012x_0 - x_1 + 0.031)^2$	$2.007 - \frac{1.09}{0.179(0.01 - x_1)^2(-x_0 - 0.03)^2 + 0.823}$
SeTGAP	$0.999e^{1.2x_3} + 1.0\sin($ $x_0 + x_1x_2 +$ $12.567) + 0.003$	$1.0\cos(0.2x_2^2 + 12.564)|x_1| +$ $1.0\tanh(0.501x_0)$	$\frac{3.767 - 3.787x_1^2}{(3.786\sin(6.283x_0) + 5.68)} + 0.003$	$\frac{14.946}{0.121x_0^3 - 15.228x_0^4 + 0.176(-x_0)^2 - 14.95} -$ $\frac{18.513}{0.135x_1 - 0.322x_1^3 + 18.515(-x_1)^4 + 18.515} + 2.0$

Table 27: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 8

Method	E9	E10	E11	E12	E13
PySR	$0.843e^{1.412\cos(x_0)} + \log(x_1 + 0.504)$ $-3.313 + 0.843\cos(\cos(0.866x_0))$ $-0.036\cosh(x_0)$	$\sin(x_0e^{x_1})$	$2x_0\log(x_1^2)$	$x_0\sin(\cos(0.001/$ $((1581.494x_1)/(\cosh(x_1))$ $+\cosh(x_1)))/x_1) + 1$	$\sqrt{x_0}\log(x_1^2)$
TaylorGP	$\log\left(\frac{0.764}{|x_0|}\right) + \tanh(x_1) - (|\log\left(\frac{0.792}{|x_0|}\right)$ $+\log(|\tanh(\frac{x_1}{x_0})|)|)^{0.5}$	$\sin(x_0e^{x_1})$	$4x_0\log(|x_1|)$	$0.938 + \frac{-x_0x_1 - 0.344}{-x_1^2 - 0.307}$	$0.255x_0\log(|x_1|) + 2.807\log(0.943$ $|x_1|)|\log(0.826\sqrt{|x_1|}|\log(4.902$ $|\log(4.902|\log(|x_1|)|)|)|) + 0.13|^{0.5}$
NeSymReS	$\log\left(\frac{0.282|x_1|}{|x_0|}\right)$	$\sin(x_0e^{x_1})$	$x_0\log(x_1^4)$	$(x_0 + x_1)\sin\left(\frac{1}{x_1}\right)$	$0.239x_0 + \log(x_1^2)$
E2E	$-6.44|0.81\arctan(0.295x_0 + 0.004)$ $+0.001| + 3.33 + 96.1/((0.021x_0 -$ $47.199)(0.062x_1 + 0.262)(0.023x_0$ $+0.692x_1 + 2.901))$	$0.994\sin((-5.26x_0 -$ $0.008)(0.273e^{0.573x_1} +$ $0.006)(-0.31x_1 +$ $0.118\log(0.055x_1 +$ $0.756) - 0.782))$	$(0.331x_0 + 0.001)$ $(20.8\log(0.327(|20.18$ $x_1 + 0.198| + 0.009)^{0.5}$ $-0.003) - 4.11)$	$7.72\sin(0.003x_0(0.008 +$ $\frac{6.35}{0.173x_1 + 0.002})) + 0.078$ $|0.006x_1 - 0.247| + 0.987$	$(0.034 - 0.052\sqrt{|0.603x_0 + 1|})$ $(0.08 - 57.9\log(|7.09|0.162x_1$ $-0.004| - 0.001|))$
TPSR	$0.13x_1 + 2.261\arctan(x_1 + 0.751)$ $-1.0\log(13.419x_0^2 + 3.359) - 0.1994$	$1.0\sin(1.0x_0e^{x_1})$	$-17.697x_0 + (0.185x_0 +$ $0.002)(0.673(0.001 - x_1)^2$ $+122.071 +$ $\frac{112.993}{-0.116|29.974x_1 + 0.112| - 1.036})$ -0.157	$(7.035\cdot10^{-9}x_0 + 8.736\cdot10^{-5})$ $(5.096x_1 + 11119.99 +$ $\frac{(0.002 - 0.068x_0)(288275.2x_1 + 22166.6)}{-1.46(x_1 + 0.084)^2 - 0.985})$	$(0.089 - \frac{0.153}{0.125|x_1| + 0.01})(-15.537x_0$ $-72.042)(0.013(0.004 - x_1)^2$ $(1 - 0.005x_0)^2 - 0.01)$
SeTGAP	$-0.999\log(3.882x_0^2 + 0.963) +$ $0.999|\log(10.173x_1 + 5.052)| - 1.656$	$1.0\sin(1.0x_0e^{x_1})$	$2.0x_0\log(x_1^2)$	$1.0x_0\sin\left(\frac{1}{x_1}\right) + 1.0$	$1.0\sqrt{x_0}\log(x_1^2)$

Table 28: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 9

Method	E1	E2	E3	E4
PySR	$0.607x_0x_1 - 1.1\cos(x_0(2.25x_1 - 1.5) - 1.5x_1 + 2.571)$	$0.178x_1\sin(0.199x_2) + 2.983\sin(0.2x_2) + 2.983\log(\cosh(0.464x_0 - 1.809) + 5.42)$	$0.15e^{1.5x_0} + 0.5\cos(3x_1)$	$(0.092 - 0.008x_1)(\cosh(x_0) + 6.478) + (0.092 - 0.009x_3)\cosh(x_2) - 0.643$
TaylorGP	$0.613x_0x_1$	$\tanh(x_2) + |(x_0(-x_0 + (e^{x_0})^{\frac{1}{4}} + \log(55.556|x_0|) + 1.704 + \cos(e^{1.424\log(55.556|x_0|)^{\frac{1}{4}}})))(\tanh(x_0))|^{0.5} + 1.305|-0.587x_0 + 0.587(e^{x_0})^{\frac{1}{4}} + 0.587\log(55.556|x_0|) + 0.587\cos(x_0) + 1|^{0.5}$	$-x_0 + e^{x_0} + \cos(x_0) - 0.815$	$x_0 + 0.118x_2^2 - 0.412$
NeSymReS	$0.61x_0x_1 + \sin(0.002(-x_0 - 0.698)^2)$	$93.728e^{0.002x_2} + e^{\sin(x_0x_1)} - 86.434$	$0.583e^{x_0} - 0.465\cos(0.204x_1)$	—
E2E	$0.021x_1(29.356x_0 + 0.027) + 1.1\cos(30.874x_1 - 1.378) - 0.001$	$0.019x_1 + (0.007x_1 - 0.056)(8.915x_0 + 0.563) + (0.437x_1 + 8.091)(0.371\sin(0.194x_2 + 0.034) - 0.007) + 6.279$	$0.042x_0 - 0.004x_1 + (0.001x_0 + 0.216)(2.33\cos(3.43x_1 + 0.075) - 0.07) + 0.166\sqrt{0.779e^{2.954x_0} + 1} - 0.008$	$0.004|2.358(x_3 - 0.048)^2 + 2.231((x_0 + 0.028)^2 + 0.039)^2 + 2.177(-0.921x_1 + (x_2 + 0.01)^2 + 0.024)^2 - 0.255| + 0.003$
TPSR	$0.072x_1(8.488x_0 + 0.065) + 0.056\sin(6.353x_1 + 1.19) - 0.032$	$(-0.009 + \frac{1}{-89.564(0.13x_2 - 1)^2 - 79.916})(-1126.481((-0.001 + \frac{0.002}{0.283|(0.259x_2 + 0.033)(9.015x_2 - 0.006)| + 0.093})(0.441x_2 + 0.076)(-0.947x_0 - 0.688x_1 - 0.205x_2 + 0.213) + 1)^3 - 7.999)(-0.002x_1 + 0.122(0.134x_0 - 1)^2 + 0.303)$	$0.007x_1(82.733x_0 + 27.9) + 0.351(0.166x_1 - 1)^3(-0.361\cos((0.014 - 0.032x_1)(33.244 - 77.329x_0)) - 1)^3 - 0.553$	$-0.1167x_1 - 0.149x_3 + 0.0016(x_2 - 0.123)^2 + 0.00512(-x_2 - (-5.082x_0 - 0.035)(-0.255x_0 - 0.072) - 0.002)^2 + 1.551$
SeTGAP	$0.608x_0x_1 + 1.1\sin((2.256x_0 - 1.491)(x_1 - 0.655) + 0.047) - 0.001$	$-0.499x_0 + 0.064x_0^2 + \sqrt{0.1x_1 + 1}(3.161\sin(0.2x_2) + 0.011) + 6.46$	$0.15e^{1.5x_0} + 0.5\cos(3.0x_1)$	$-0.02x_0^2x_1 + 0.004x_0^2 + 0.01x_0^4 + 0.01x_1^2 - 0.02x_2^2x_3 + 0.01x_2^4 + 0.01x_3^2 - 0.009$

Table 29: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 9

Method	E5	E6	E7	E8
PySR	$(e^{1.2x_3}) + \sin(x_0 + x_1x_2)$	$\tanh(x_0) + 9.715\cos(0.2x_2^2)\cos(\frac{5.902}{\tanh(\cosh(0.095x_1))})$	$(2.482x_1^2 - 2.467)(\cos(0.435e^{\cos(6.278x_0 + 1.57)}) - 1.16)$	$0.8\tan(\sinh(\tanh(\cosh(0.827\cosh(\tan(\tanh(x_1))))\tanh(\cosh(x_0)))))\cosh(\tanh(x_0)) - 0.957$
TaylorGP	$(\cos(\sin(\tanh(1.515e^{x_3}))) + \tanh(1.119|(0.799\cos(\sin(\tanh(1.515e^{x_3}))) + 0.799\tanh(0.534e^{0.5(x_3)}))e^{x_3} - e^{x_3}|^{0.5}))e^{x_3}$	$\cos(x_2) + \tanh(x_0) + \frac{7.407\log(|x_1|^{0.5})\sin(x_2)}{x_2}$	$-x_1^2 + \log(|x_1^2 + \frac{-x_1^2 + \sqrt{|x_1|} + \sqrt{|x_1 - 0.479\tanh(e^{-0.538x_2^2})|}}{x_1^2 + 0.392}|)$	$e^{\tanh(\tanh(\log(0.864\sqrt{|x_0||x_1|})))}$
NeSymReS	—	$0.145x_0 + x_1\sin(0.302x_1 + \frac{x_2}{x_1})$	$-0.05632x_0 - 0.8403x_1^2$	$\cos(\frac{\sin(\frac{x_0}{x_1})}{x_0}) + 0.634$
E2E	$-0.001x_1 + 0.002x_3 + 0.023(0.024 - x_3)^2 + 0.947e^{1.206x_3} + 0.933\cos(12297.635(0.404x_1 + 1)^2 + 0.225(0.001x_0 - 0.001x_2 + 1)^{0.5} - 15.41) + 0.032$	$(0.035x_1 - 0.002)(4.984\arctan(-0.03 - \frac{4.462}{-0.539x_0 - 0.118})(0.004x_0 + 0.001)) - 0.16) + 0.929$	$(0.09 - 0.082(x_1 + 0.007)^2)(6.131(0.766 - \sin(6.516x_0 + 0.216))^2 + 5.0)$	$2.1 - \frac{0.9}{0.002|(3.087x_1 - 0.168)(37.54(x_0 + 0.029)^3 + 6.0)| + 0.6}$
TPSR	$2.627214 - 0.004x_2$	$(0.397\cos(0.017x_1 + 0.524x_2 + 0.076) + 0.24)(0.234x_0 + \cos(1.716x_1 - 2.174) + 6.18)$	$-0.934(0.086 - x_1)^2(0.001x_0 + 1 + \frac{0.116}{x_1})^2 + 1.098$	$0.001x_1 + 0.61(0.002 - \arctan(-0.01x_0(89.997x_1 - 2.242)))^2 + 0.654$
SeTGAP	$1.001e^{1.2x_3} + 1.0\sin(x_0 + x_1x_2 + 12.566) - 0.008$	$-1.0\cos(0.2x_2^2 - 9.425)|x_1| + 1.0\tanh(0.5x_0)$	$\frac{8.18x_1^2 - 8.247}{(-8.185\sin(6.283x_0 - 6.283) - 12.273)} - 0.006$	$\frac{16.205}{(-0.015x_0 - 16.207x_0^4 - 0.033(-x_0)^3 - 16.206)} + 2.0 + \frac{11.914}{(0.013x_1 - 0.118x_1^2 - 0.016x_1^3 - 11.817x_1^4 - 11.901)}$

Table 30: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 9

Method	E9	E10	E11	E12	E13
PySR	$\log(\frac{x_1 + 0.5}{2x_0^2 + 0.5})$	$\sin(x_0e^{x_1})$	$2.0x_0\log(x_1^2)$	$x_0\sin(\frac{1}{x_1}) + 1.0$	$\sqrt{x_0}\log(x_1^2)$
TaylorGP	$-\log(|x_0|) + \tanh(x_1) - 0.298 - \sqrt{|\log(\frac{0.873}{|x_0|}) + \tanh(x_1) - \sqrt{|x_0|}|}$	$\sin(\tanh(x_0))\cos(\sqrt{|x_1|})$	$4x_0\log(|x_1|)$	$\frac{0.998x_0\sin(\frac{x_0}{x_1(x_0 + 0.048)})}{\sqrt{|\log(\sqrt{|x_0|})|} + 0.998}$	$(\sqrt{x_0}\log(\sqrt{|x_1|})^2 + 0.689)(\log(\sqrt{|x_1|}) + 0.446) + \log(|x_1|)(\log(|x_0|) + 1) + \log(|x_1\tanh(x_1)|))$
NeSymReS	$\log(\frac{0.281|x_1|}{|x_0|})$	$\sin(x_0e^{x_1})$	$x_0\log(x_1^4)$	$(x_0 + x_1)\sin(\frac{1}{x_1})$	$0.243x_0 + \log(x_1^2)$
E2E	$-0.6\log(10.053(0.009 - x_0)^2 + 0.6) + 1.9 - \frac{0.7}{0.274x_1 + 0.301}$	$-0.994\sin((3.608x_0 + 0.013)(-0.009 - 0.234e^{(0.595 - 0.053x_1)} \ldots(1.976x_1 + 0.061))) - 0.001$	$(2.922x_0 + 0.006)(4.14\arctan(0.336 - 0.008|0.26x_1 + 0.695 - \frac{84.4}{3.242x_1 - 0.071}|) - 0.047)$	$0.588 + 7.1\sin(0.058 - 9.51/((0.01 - \frac{103.409}{13.96x_0 - 0.071})(-0.001x_0 + 10.994x_1 - 0.059)))$	$(0.741 - 4.65\arctan(0.175x_0 + 0.35))(0.174e^{0.175x_0} - 20.8)(0.008\log(0.001x_1 + 12.9) + 0.072)$
TPSR	$59.31(-0.003x_1 - 1)^2 - 66.389 - 82.366/(-4.5|(31.371 + \frac{17.165}{0.218x_1 + 0.124})(0.301x_0 - 0.001)(0.007x_1 + 0.021)| - 10.235)$	$1.0\sin(0.9992x_0e^{x_1})$	$7.953x_0 - 0.188$	$0.827 + 5.558\sin((0.056x_0 - 4.862)(0.006x_0(0.1 + \frac{8.807}{0.002 - 1.613x_1}) - 0.009))$	$\frac{-0.684x_0 + 97.57 + \frac{50.936}{-0.009x_0 - 0.535}}{} + 15.513/((66.734 + 207.296e^{-0.219x_0})(-0.012(x_1 - 0.002)^2 - 0.007))$
SeTGAP	$-1.001\log(13.041x_0^2 + 3.264) + 0.998|\log(7.365x_1 + 3.647)| - 0.108$	$1.0\sin(1.0x_0e^{x_1})$	$4.0x_0\log(|x_1|)$	$1.0x_0\sin(\frac{1}{x_1}) + 1.0$	$2.0\sqrt{x_0}\log(|x_1|)$

Table 31: Comparison of predicted equations (E1—E4) with rounded numerical coefficients — Iteration 10

Method	E1	E2	E3	E4
PySR	$0.608 x_0 x_1$	$x_0 + 2.969 \sin (0.2 x_2) + \cosh (0.083 x_0 - 3.519) - 10.302$	$0.15 e^{1.5 x_0} + 0.5 \sin (3.0 x_1 + 1.571)$	$-0.165 x_1 - 0.165 x_3 + 1.564 \log((\cosh(\sinh(0.083 x_0^2)) - 0.553) \cosh(x_2)) + 1.564 \cos(x_2)$
TaylorGP	$\frac{0.604 x_0 x_1 (x_0^2 x_1^2)^{\frac{1}{4}}}{\sqrt{|x_0 x_1|}}$	$-0.513 x_0 - 0.001 x_1 + 0.204 x_2 + 8.572$	$-x_0 \tanh (0.632 e^{x_0} - 1) + e^{x_0} - \log (e^{x_0}) - 0.682$	$x_0 + x_2 - 2$
NeSymReS	$0.589 x_0 x_1 + \cos(0.598(0.966 x_0 - x_1)^2)$	$-x_0 + e^{\sin (x_1)} + 63.876 - 56.599 e^{-0.004 x_2}$	$0.575 e^{x_0} - 0.418 \sin (1.451(0.052 x_1 + 1)^2)$	—
E2E	$(23.064 x_0 - 0.034) (0.026 x_1 + 0.001) - 0.001 + 1.09 \sin(0.158 x_0 + (1.872 x_1 + 0.125)(0.003|0.021 x_1 - 2.239| + 81.7) + 0.021)$	$-0.008 x_0 (0.172 x_2 - 1.607)(0.007 x_1 - 40.299) + 6.26 + (0.008 x_1 + 0.149) (19.2 \sin((0.001 x_2 + 0.073) (3.077 x_2 + 0.001)) + 0.003)$	$0.003 x_0 + 0.06 e^{1.804 x_0} - 0.499 \cos(0.003 x_0 - 2.985 x_1 + 59.557) + 0.159$	$0.843|0.07 x_3 - 0.013(0.836 x_1 - (x_0 + 0.059)^2(0.004 x_1 - 1)^2 + 0.039)^2 - 0.01(0.96 x_3 - (x_2 - 0.046)^2 + 0.046)^2 + 0.017|$
TPSR	$-x_0 + (1.009 - 73.098 x_0) (-0.008 x_1 - 0.013) + 0.007$	$(0.098 + \frac{1}{x_0 + 44.83})(0.217 - 0.991 x_2)(- 0.006 x_0 - (-0.303 + \frac{23.261}{x_2 - 0.219})(0.033 x_1 + 38.009)(-0.003 x_2 - 0.213)(-0.613 x_2 + (0.499 - 0.007 x_0)(27.627(-0.01 x_0 - 1)^2 + 7.021) - 17.446) + 111.696 + \frac{0.001}{x_1})$	$-0.008 x_0 (5.0 - 75.398 x_1) + 0.04 x_0 + 1.1 \cos(8.906(0.168 - 0.253 x_1) (x_0 - 0.667) + 1.567)$	$-0.182 x_1 + 0.005(1 - 0.015 (0.007 - |9.838 x_0 + 0.02|)^2)^2 + 0.04(0.011 x_0^2 + 0.45 x_2^2 - 0.425 x_3 + 1)^2 + 0.045$
SeTGAP	$0.607 x_0 x_1 + 1.099 \sin((2.242 x_0 - 1.492) (x_1 - 0.685))$	$-0.5 x_0 + 0.062 x_0^2 + 3.162\sqrt{0.1 x_1 + 1} \sin (0.201 x_2) + 6.501$	$0.147 e^{1.507 x_0} - 0.5 \sin (3.001 x_1 + 4.712)$	$-0.02 x_0^2 x_1 + 0.01 x_0^4 + 0.01 x_1^2 + 0.01 x_1^4 + 0.01 x_3^2 - (0.02 x_2^2 + 0.002)(x_3 + 0.212) + 0.009$

Table 32: Comparison of predicted equations (E5—E8) with rounded numerical coefficients — Iteration 10

Method	E5	E6	E7	E8
PySR	$1.0 e^{1.2 x_3} + \sin (x_0 + x_1 x_2)$	$|x_1| \cos(0.2 x_2^2) + \tanh (0.5 x_0)$	$\frac{2.119 - 2.118 x_1^2}{2.118 \sin (6.283 x_0) + 3.176}$	$\log(\tan(1.442(\tanh(\cosh(x_1) \tanh(\sqrt{\cosh(x_0)})) \tanh(\cosh(x_0)))^{0.5}) - 0.635)$
TaylorGP	$2 e^{x_3} - 0.24 - \tanh(e^{x_3} + \sin((1.598 - 0.126\sqrt{|x_3|}) (e^{x_3} - 0.24)))\sqrt{|x_3 + e^{x_3}|}$	$\cos (x_2) + \tanh (x_0) + \frac{4.95 \sin (x_2)}{x_2}$	$-x_1^2 + 0.802\sqrt{|x_1^2 + \sin (x_0)|}$	$2.037 \tanh (0.521\sqrt{|x_0||x_1|})$
NeSymReS	—	$-0.368 x_0 + x_1 \sin (\frac{x_0}{x_1} + 0.005 x_2)$	$\frac{0.166 x_0 + x_1^2}{\cos (2.498(0.137 x_1 - 1)^2) - 2.073}$	$\cos (\frac{\sin (\frac{x_0}{x_1})}{x_0}) + 0.64$
E2E	$0.002 x_2 + 0.273 + 0.845 e^{1.25 x_3} e^{0.008 \arctan (-0.065 x_0 + 0.002 x_1 - 14.76)} + 0.993 \cos(-16.985 x_2 + (-0.008 x_0 - 37.102)(67.0|0.925 + \frac{0.008}{-0.573 x_3 - 0.015}| - 59.1) + 0.088 + \frac{46.3}{0.012 - 0.001 x_0})$	$(-0.007 x_1 - 0.015)(9.191 \cos((0.021 - 0.906 x_2)(21.206|3.697 x_0 + 11.042 (-x_2 - 0.077)^2 + 0.689| + 0.008)) - 0.121) - 0.219 \arctan(-0.489 x_0 - 0.094) + 0.566$	$0.003 x_0 + 0.144 + (0.002 - 0.005(-x_1 - 0.014)^2) /(0.006 - 0.001(- \sin(7.343 x_0 + 3.499) - 0.628)^2)$	$2.0 - 0.9/(0.006|(2.068 x_0 - 0.4) (3.413 x_0 + 0.229)(2.073 x_1 + 0.038) (3.109 x_1 + 0.206)| + 0.5)$
TPSR	$14.264(0.045 e^{x_3} + 0.002 \cos(7.628 x_2 - 12.078) + 1)^2 - 14.456$	$1.56(0.05 x_0 + 1)^2 - 0.88$	$1.0 - 0.89(-x_1 - 0.05)^2$	$2.015 - \frac{0.793}{0.124(0.001 - x_0)^2(-x_1 - 0.005)^2 + 0.595}$
SeTGAP	$1.002 e^{1.2 x_3} + 0.999 \sin (x_0 + x_1 x_2 - 6.281)$	$0.998 \cos (0.201 x_2^2 - 18.884)|x_1| + 1.0 \tanh (0.501 x_0)$	$\frac{1.005 x_1^2 - 0.873}{(- \sin (6.283 x_0 - 6.283) - 1.507)} + 0.084$	$-\frac{11.534}{(11.542 x_1^4 + 11.535)} + 2.0 - \frac{11.534}{(0.045 x_0 + 0.062 x_0^2 + 11.474 x_0^4 + 11.528)}$

Table 33: Comparison of predicted equations (E9—E13) with rounded numerical coefficients — Iteration 10

Method	E9	E10	E11	E12	E13
PySR	$\sqrt{1.148 x_1} - 4.723 + \frac{4.49}{\cosh (x_0 \tanh (\frac{2.296}{1.849 - 0.771 \cos (x_0)}))}$	$\sin (x_0 e^{x_1})$	$2 x_0 \log (x_1^2)$	$1.0 x_0 \sin (\frac{1.0}{x_1}) + 1.0$	$\log(3.711 \sinh(\sqrt{|\log(|\cosh (0.381 x_1)|)|})) \tanh(\sqrt{e^{-2.032 \cosh (0.209 x_1)}} + 2.45)2.015\sqrt{|x_0|}$
TaylorGP	$\log (\sqrt{|\frac{x_1}{x_0}|}) - 1.486\sqrt{|x_0|} + 0.889$	$\cos (\sqrt{|x_1|}) \tanh (\tanh (x_0))$	$4 x_0 \log (|x_1|)$	$\log (e^{\tanh (\frac{x_0}{x_1})}) + \tanh (\frac{x_0}{x_1}) + \sqrt{|\tanh (x_1)|}$	$2 \log (|x_1|)\sqrt{|x_0|}$
NeSymReS	$\log (\frac{0.284|x_1|}{|x_0|})$	$\sin (x_0 e^{x_1})$	$x_0 \log (x_1^4)$	$(x_0 + x_1) \sin (\frac{1}{x_1})$	$0.23 x_0 + \log (x_1^2)$
E2E	$-0.986 \log(26.495(-0.179 - \frac{1}{0.691 x_1 + 7.596})^2(-0.063 - \frac{1}{0.691 x_1 + 0.626})^2(0.031 - x_0)^2 + 0.224) - 0.087$	$-1.0 \sin(0.026 x_0 (1.0 - 36.638 e^{1.091 x_1})) - 0.001$	$(-0.214 x_0 - 0.001) (6.0 \log(1742.4 (0.02 - \frac{1}{1.244 x_1 + 0.023})^2 + 0.01) - 50.0)$	$0.996 - 7.12 \sin((0.01 + \frac{19.5}{0.006 x_0 - 9.741 x_1 - 0.305}) (0.067 x_0 + 0.001))$	$(2.27 - 0.465(0.086(|(92.8 \arctan(0.172 x_0 + 20.71) + 29.3) (-0.015 x_1 + (0.913 + \frac{61.081}{x_1})(0.056 x_0 - 2.916) + 0.736)| + 0.001)^{0.5} - 1)^{0.5})(3.87 \log(|21.928|0.027 x_0 - 1|^{0.5} - 26.9|) + 0.05)$
TPSR	$-97.127 x_1 - 0.029(-x_0 - 0.024)^2 + (-0.013 x_0 + x_1 - 1169.375)(-0.084 x_1 - 0.002 \arctan(0.29 - 5.115 |0.129 x_0 + 0.001|) + 0.118) + 137.562$	$1.003 \sin (0.99667 x_0 e^{x_1})$	$(-0.524 x_0 - 0.02) (-0.277 x_1^2 - 7.971 - \frac{32.033}{-5.252|0.701 x_1 + 0.001| - 0.726})$	$0.026 x_0(-0.068 + \frac{1}{-0.634(-x_1 - 0.964)^2 - 0.887}) (-36.911 \arctan(44.002 x_1 - 16.916) - 34.003) - 0.075 x_0 + 0.978$	$(-0.015 + \frac{1}{0.539|9.48 x_1 + 0.014| + 0.511}) (0.417 - 0.44(0.001 - x_1)^2) (1.325 - \frac{2.356}{x_0 + 1.916}) (-1.316 x_0 - 22.185)$
SeTGAP	$-1.0 \log (10.831 x_0^2 + 2.707) + 1.0|\log (21.911 x_1 + 10.961)| - 1.399$	$1.0 \sin (1.0 x_0 e^{x_1})$	$4.0 x_0 \log (|x_1|)$	$1.0 x_0 \sin (\frac{1}{x_1}) + 1.0$	$1.0\sqrt{x_0} \log (x_1^2)$

Table 34: Comparison of expressions learned by SeTGAP Under Noisy Conditions

Problem	$\sigma_a = 0.01$	$\sigma_a = 0.03$	$\sigma_a = 0.05$
E1	$0.608\,x_0\,x_1 + 1.089\sin((2.263$ $x_0 - 1.525)(x_1 - 0.665) - 0.011)$	$(0.611\,x_0 + 0.001)(x_1 - 0.043) - 0.885$ $\sin((2.253\,x_0 - 1.544)(x_1 - 0.721) - 3.214) + 0.006$	$0.61\,x_1(x_0 - 0.006) -$ $0.47\sin((2.381\,x_0 - 1.171)(x_1 - 0.798)$ $+8.501) + 0.006$
E2	$0.063\,x_0^2 - 0.499\,x_0 + (3.078$ $\sqrt{0.098\,x_1 + 1} + 0.062)$ $(\sin(0.199\,x_2) - 0.002) + 6.481$	$0.063\,x_0^2 - 0.499\,x_0 + (3.222\sqrt{0.099\,x_1 + 1} -$ $0.058)(\sin(0.201\,x_2) - 0.007) + 6.511$	$(-(0.507\,x_0 - 6.557)(0.181\,x_1 - 18.566) -$ $(0.374\,x_0^2 + 18.893\sin(0.198\,x_2))$ $\log(x_1 + 19.824))/(0.181\,x_1 - 18.566)$
E3	$0.143e^{1.516\,x_0} + 0.506$ $\sin(2.999\,x_1 + 1.572) + 0.009$	$0.148e^{1.505\,x_0} - 0.498\sin(3.0\,x_1 - 1.572) + 0.003$	$0.149e^{1.502\,x_0} -$ $0.496\sin(3.0\,x_1 - 1.581) + 0.003$
E4	$0.01x_0^4 - 0.02x_0^2x_1 - 0.007x_0^2$ $+0.009x_1^2 + 0.011x_2^2 - 0.02x_2^2x_3$ $-0.02x_2^2 + 0.008x_3^2 + 0.113$	$0.01x_0^4 + 0.01x_2^4 - 0.02x_2^2x_3 + 0.01x_3^2 +$ $0.242\cosh(10.69\sqrt{1 - 0.016x_1} - 8.186) - 1.473$	$0.01x_1^4 - 0.021x_3^2x_4 + 0.01x_3^4 +$ $0.011x_4 + 0.01x_4^4 + (0.093x_1^2 + 0.007$ $|x_4 + 1|)\sin(0.248x_2 - 2.981) + 0.124$
E5	$1.004e^{1.199\,x_3} + 0.997$ $\sin(1.0\,x_0 + 1.0\,x_1\,x_2) - 0.005$	$(-(0.015\sin(0.508\,x_2 + 1.574) - 0.001)(|4.779$ $\sin(1.0\,x_0 + 3.141) - 6.283| - 6.283) +$ $(6.226\sin(0.508\,x_2 + 1.574) - 6.283)(0.002$ $e^{1.202|x_3 + 5.282|} + 0.145\sin(1.0|x_1 - 6.283| -$ $1.847) + 0.067\sin(1.0|x_1 + 6.202| - 4.425)|2.649$ $\sin(1.0\,x_0 + 3.24) - 2.097| + 0.005))/$ $(6.226\sin(0.508\,x_2 + 1.574) - 6.283)$	$0.996e^{1.202\,x_3} +$ $1.0\sin(1.0\,x_0 + 1.0\,x_1\,x_2) + 0.004$
E6	$1.0\cos(0.2\,x_2^2 + 0.008)|x_1|$ $+1.0\tanh(0.5\,x_0) + 0.005$	$\cos(0.199\,x_2^2 + 0.028)(1.001|x_1| - 0.001) +$ $1.002\tanh(0.486\,x_0) + 0.02$	$1.0\cos(0.2\,x_2^2)|x_1| + 0.997\tanh(0.502\,x_0)$
E7	$\frac{1.001\,x_1^2 - 0.9914}{\sin(6.283\,x_0 + 3.15) - 1.501} + 0.011$	$\frac{-1.001\,x_1^2 + 1.006}{\sin(6.283\,x_0) + 1.501} + 0.007$	$-0.982\frac{x_1^2 - 0.993}{\sin(6.283\,x_0 + 3.139) - 1.492} + 0.019$
E8	$2.0 - 11.356/(11.313x_1^4 - 0.116x_1^3$ $+0.047x_1^2 + 0.078x_1 + 11.352) -$ $\frac{10.636}{10.804x_0^4 - 0.113x_0^2 + 10.646}$	$2.0 - \frac{15.531}{15.414x_1^4 - 0.051x_1^3 + 0.165x_1^2 + 0.018x_1 + 15.517} -$ $\frac{12.561}{12.523x_0^4 - 0.085x_0^3 + 0.062x_0^2 + 0.061x_0 + 12.553}$	$2.0 - \frac{12.837}{12.956x_0^4 + 0.007x_0^3 - 0.125x_0^2 + 12.861} -$ $\frac{16.19}{16.217x_1^4 - 0.007x_1^3 + 16.202}$
E9	$-1.0\log(4.884\,x_0^2 + 1.22) +$ $1.0|\log(9.415\,x_1 + 4.709)| - 1.35$	$-1.0\log(17.004\,x_0^2 + 4.251) +$ $0.999|\log(10.158\,x_1 + 5.072)| - 0.176$	$5.965\sqrt{0.625\log(9.36\,x_1 + 6.35) + 1} -$ $0.999\log(11.259\,x_0^2 + 2.817) - 7.713$
E10	$1.0\sin(1.0\,x_0e^{x_1})$	$1.0\sin(1.0\,x_0e^{x_1})$	$1.0\sin(1.0\,x_0e^{1.0\,x_1})$
E11	$2.0\,x_0\log(x_1^2)$	$(4.001\,x_0 - 0.004)\log(|x_1|) + 0.003$	$(4.001\,x_0 - 0.008)\log(|x_1|) + 0.006$
E12	$1.0\,x_0\sin(1/x_1) + 1.0$	$1.0\,x_0\sin(1/x_1) + 1.0$	$1.0\,x_0\sin(1/x_1) + 1.0$
E13	$(1.001\sqrt{x_0} - 0.001)\log(x_1^2)$	$(0.993\sqrt{x_0} - 0.002)\log(x_1^2) + 0.029$	$(1.001\sqrt{x_0} - 0.003)\log(x_1^2)$

C   SeTGAP Under Noisy Conditions

This appendix presents the expressions learned by SeTGAP under noisy conditions, shown in Table 34, where shaded cells indicate an incorrectly identified functional form. While the expressions obtained in the noiseless setting are provided in Tables 2–4, here we focus on analyzing the impact of noise on the recovered functional forms. These expressions correspond to the results reported in Table 6, where different noise levels were introduced to assess the robustness of SeTGAP.