Interpreting diffusion score matching using normalizing flow

Abstract

Scoring matching (SM), and its related counterpart, Stein discrepancy (SD) have achieved great success in model training and evaluations. However, recent research shows their limitations when dealing with certain types of distributions. One possible fix is incorporating the original score matching (or Stein discrepancy) with a diffusion matrix, which is called *diffusion score matching* (DSM) (or *diffusion Stein discrepancy* (DSD)) . However, the lack of the interpretation of the diffusion limits its usage within simple distributions and manually chosen matrix. In this work, we plan to fill this gap by interpreting the diffusion matrix using normalizing flows. Specifically, we theoretically prove that DSM (or DSD) is equivalent to the original score matching (or score matching) evaluated in the transformed space defined by the normalizing flow, where the diffusion matrix is the inverse of the flow's Jacobian matrix. In addition, we also build its connection to Riemannian manifolds, and further extend it to continuous flows, where the change of DSM is characterized by an ODE.

1. Introduction

Recently, score matching (Hyvärinen & Dayan, 2005) and its closely related counterpart, Stein discrepancy (Gorham, 2017) have made great progress in both understanding their theoretical properties and practical usage. Particularly, unlike Kullback–Leibler (KL) divergence which can only be used for distributions with known normalizing constant, SM (or SD) can be evaluated for unnormalized densities, and requires fewer assumptions for the probability distributions (Fisher et al., 2021). Such useful properties enable them to be widely applied in training energy-based model (EBM) (Song et al., 2020a; Grathwohl et al., 2020; Wenliang et al.,

2019), state-of-the-art score-based generative model (Song & Ermon, 2019; Song et al., 2020b), statistical tests (Liu et al., 2016; Chwialkowski et al., 2016) and variational inference (Hu et al., 2018; Liu & Wang, 2016).

Despite their elegant statistical properties, recent work (Barp et al., 2019) demonstrated their failure when dealing with certain type of distributions (e.g. heavy-tailed distributions). For instance, when the data and the model are heavy tailed distributions, the model can fail to recover the true mode even in one dimensional case. The root of this problem is that the SM (or SD) objective is highly non-convex and does not correlate well with likelihood. To fix it, Barp et al. (2019) proposed a variant called *diffusion score matching* (and *diffusion Stein discrepancy*), where a diffusion matrix is introduced. However, the author did not provide us an interpretation of this diffusion matrix. In fact, the diffusion used by the author (Barp et al., 2019) is manually chosen for toy densities. Such lack of interpretation hinders further development of a proper training method of the diffusion matrix.

In this paper, we aim to give an interpretation based on normalizing flows, which sheds light on developing training method for the diffusion. We summarize our contributions as follows:

- We theoretically prove that DSM (or DSD) is equivalent to the original SM (or SD) performed in the transformed space defined by the normalizing flow. The diffusion matrix is exactly the same as the inverse of the flow's Jacobian matrix.

- We further show that its connection to Riemannian manifold. Specifically, we show the diffusion matrix is closely related to the Riemannian metric tensor.

- We further extend DSM to their continuous version. Namely, we derive an ODE to characterize its instantaneous change.

We hope that by building these connections, a broad range of techniques from normalizing flow communities can be leveraged to develop training methods for the diffusion matrix.

[1]Anonymous Institution, Anonymous City, Anonymous Region, Anonymous Country. Correspondence to: Anonymous Author <anon.email@domain.com>.

Preliminary work. Under review by INNF+ 2021. Do not distribute.

2. Background: Diffusion Stein discrepancy

2.1. Score matching and Stein discrepancy

Let $\mathcal{P}$ be the space of Borel probability measures on $\mathbb{R}^D$, $\mathbb{Q} \in \mathcal{P}$ to be a probability measure, the objective for model learning is to find a sequence of probability measures $\{\mathbb{P}_\theta : \theta \in \Theta\} \subset \mathcal{P}$ that approximates $\mathbb{Q}$ in an appropriate sense. One common way to achieve this is by defining a discrepancy measure $\mathcal{D} : \mathcal{P} \times \mathcal{P} \to \mathbb{R}$, which quantifies the differences between two probability measures. Thus, the optimal parameters θ^* can be obtained by $\theta^* = \mathrm{argmin} \mathcal{D}(\mathbb{Q}||\mathbb{P}_\theta)$. The choice of discrepancy depends on the properties of the probability measures, the efficiency and its robustness. The one we are focused on is called *Fisher divergence*. Assuming for probability measures $\mathbb{Q}$ and $\mathbb{P}_\theta$, we have corresponding twice differentiable densities $q(\boldsymbol{x})$, $p_\theta(\boldsymbol{x})$. The *Fisher divergence* (Johnson, 2004) is defined as

$$\mathcal{F}(q,p) = \frac{1}{2}\mathbb{E}_q[||\boldsymbol{s}_p(\boldsymbol{x}) - \boldsymbol{s}_q(\boldsymbol{x})||^2] \tag{1}$$

where $\boldsymbol{s}_p(\boldsymbol{x}) = \nabla_{\boldsymbol{x}} \log p_\theta(\boldsymbol{x})$ is called the score of p_θ, and $\boldsymbol{s}_q$ is defined accordingly. Despite that q is often used for underlying data densities with the intractable $\boldsymbol{s}_q$, $\boldsymbol{s}_q$ in fact acts as a constant for parameter θ. Thus, one can use integration-by-part to derive the following:

$$\mathcal{F}(q,p_\theta) = \underbrace{\mathbb{E}_q[\frac{1}{2}||\boldsymbol{s}_p(\boldsymbol{x})||^2 + Tr(\nabla_{\boldsymbol{x}}\boldsymbol{s}_p(\boldsymbol{x}))]}_{\mathcal{L}(\theta)} + C_q \tag{2}$$

where C_q is a constant w.r.t parameter θ. This alternative objective $\mathcal{L}(\theta)$ is referred as *score matching* (Hyvärinen & Dayan, 2005).

Another discrepancy measure we are interested is called *Stein discrepancy*, which is defined as

$$\mathcal{S}(q,p_\theta) = \sup_{\boldsymbol{f} \in \mathcal{H}} \mathbb{E}_q[\boldsymbol{s}_p(\boldsymbol{x})^T \boldsymbol{f}(\boldsymbol{x}) + \nabla_{\boldsymbol{x}}^T \boldsymbol{f}(\boldsymbol{x})] \tag{3}$$

where $\boldsymbol{f} : \mathbb{R}^D \to \mathbb{R}^D$ is a test function, and $\mathcal{H}$ is an appropriate test function family, e.g. reproducing kernel Hilbert space (Liu et al., 2016; Chwialkowski et al., 2016) or Stein class (Gorham, 2017; Liu et al., 2016). Recent work (Hu et al., 2018) proved a strong connection between Stein discrepancy and Fisher divergence by showing the optimal test function:

$$\boldsymbol{f}^*(\boldsymbol{x}) \propto \boldsymbol{s}_p(\boldsymbol{x}) - \boldsymbol{s}_q(\boldsymbol{x}) \tag{4}$$

Thus, by substitution back into eq. 3, we can show Stein discrepancy is equivalent to Fisher divergence up to a multiplicative constant.

Recent works (Barp et al., 2019; Gorham et al., 2019) further extend the score matching and Stein discrepancy by incorporating a diffusion matrix $\boldsymbol{m}(\boldsymbol{x}) : \mathbb{R}^D \to \mathbb{R}^{D \times D}$, which are

Figure 1. The SM loss computed for different θ. Orange line: Original SM loss with orange dashed line indicating valid initialization region. Blue line: DSM loss with dashed blue line indicating the optimum solution θ^*. The **black** dashed line indicates true θ.

called *diffusion score matching* (DSM) and *diffusion Stein discrepancy* (DSD). DSM is defined as

$$DSM_m(q,p_\theta) = \frac{1}{2}\mathbb{E}_q[||\boldsymbol{m}(\boldsymbol{x})^T(\boldsymbol{s}_p(\boldsymbol{x}) - \boldsymbol{s}_q(\boldsymbol{x}))||^2] \tag{5}$$

where $\boldsymbol{m}(\boldsymbol{x})$ is a matrix-valued function. Similarly, DSD is defined as

$$DSD_m(q,p_\theta)$$
$$= \sup_{\boldsymbol{f} \in \mathcal{H}} \mathbb{E}_q[(\boldsymbol{m}(\boldsymbol{x})^T\boldsymbol{s}_p(\boldsymbol{x}))^T\boldsymbol{f}(\boldsymbol{x}) + \nabla_{\boldsymbol{x}}^T(\boldsymbol{m}(\boldsymbol{x})\boldsymbol{f}(\boldsymbol{x}))] \tag{6}$$

It can be shown that as long as $\boldsymbol{m}(\boldsymbol{x})$ is invertible, $DSM_m(q,p_\theta)$ and $DSD_m(q,p_\theta)$ are valid divergences.

These two extensions have demonstrated superior performances when dealing with certain type of distributions. In the following, we give a motivating example similar to Barp et al. (2019).

2.2. Motivating example: Student-t distribution

Let assume q, p_θ to be 1 dimensional student-t distribution. The target is to approximate q by p_θ. The training set is 300 i.i.d data sampled from q with mean 0 and scale 0.3. We assume the scale parameter for p_θ is the same as q, and the only trainable parameter θ is the mean. The degree of freedom is 5 for both q, p_θ. Figure 1 shows the score matching loss computed for different θ. We can observe that for original SM loss, it is highly non-convex, and the loss value does not correlate well with likelihood. Indeed, we can see the true location $\theta = 0$ is protected by two high 'walls'. In other words, unlike maximum likelihood estimator, a parameter θ that is closer to the ground truth does not necessarily produce low SM loss. One important consequence is that unless the initialized θ is within the narrow valid region, the gradient-based optimization will never recover the ground truth.

On the other hand, if we chose $\boldsymbol{m}(\boldsymbol{x}) = (1 + \frac{(\boldsymbol{x}-\theta)^2}{0.6})$ as the diffusion matrix, the corresponding $DSM_m(q, p_\theta)$ loss is convex. The ground truth can be recovered by minimizing DSM with a proper gradient-based optimizer.

The selection of the diffusion matrix is crucial to the success of the estimator. Unfortunately, the interpretation of this matrix is unclear, not mentioning a selection algorithm. In the following, we aim to shed lights on this problem by connecting this diffusion with normalizing flows.

3. Diffusion matrix as normalizing flow

3.1. Interpreting DSM/DSD using normalizing flow

Let assume we have two densities $q_X(\boldsymbol{x})$, $p_X(\boldsymbol{x})$ defined on $\mathbb{R}^D$ and are twice differentiable. We further define a differentiable invertible transformation $\boldsymbol{T}(\boldsymbol{x}) : \mathbb{R}^D \to \mathbb{R}^D$. Let's define

$$\boldsymbol{y} = \boldsymbol{T}(\boldsymbol{x}) \tag{7}$$

with corresponding densities $q_Y(\boldsymbol{y})$ and $p_Y(\boldsymbol{y})$. We can prove the following theorem:

Theorem 3.1. *For twice differentiable densities $q_X(\boldsymbol{x})$, $p_X(\boldsymbol{x})$ and an invertible differentiable transformation $T :$ $\mathbb{R}^D \to \mathbb{R}^D$, the* diffusion score matching *objective (Eq.5) is equivalent to the original* score matching

$$\mathcal{F}(q_Y, p_Y) = \frac{1}{2}\mathbb{E}_{q_Y}[||\boldsymbol{s}_{p_Y}(\boldsymbol{y}) - \boldsymbol{s}_{q_Y}(\boldsymbol{y})||^2] \tag{8}$$

where $\boldsymbol{y} = T(\boldsymbol{x})$, and p_Y, q_Y are corresponding densities after the transformation. The diffusion matrix $\boldsymbol{m}(\boldsymbol{x})$ is the inverse of the Jacobian matrix $(\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1}$

Proof. From the change of variable formula, the corresponding densities $p_Y(\mathbf{y})$, $q_Y(\mathbf{y})$ can be defined as:

$$p_Y(\mathbf{y}) = p_X(T^{-1}(\mathbf{y}))|\frac{\partial T^{-1}(\mathbf{y})}{\partial \mathbf{y}}|,$$

$$q_Y(\mathbf{y}) = q_X(T^{-1}(\mathbf{y}))|\frac{\partial T^{-1}(\mathbf{y})}{\partial \mathbf{y}}|.$$

Then the Fisher divergence (Eq.1) for p_Y, q_Y can be formulated as follows (we use the notation $\mathcal{F}_T$ to emphasise that p_Y, q_Y are obtained using transformation T):

$$F(p_Y, q_Y) := \frac{1}{2}\mathbb{E}_{q_Y}[||\nabla_\mathbf{y}\log p_Y(\mathbf{y}) - \nabla_\mathbf{y}\log q_Y(\mathbf{y})||_2^2]$$

$$= \frac{1}{2}\mathbb{E}_{q_Y}[||\nabla_\mathbf{y}\log p_X(T^{-1}(\mathbf{y})) - \nabla_\mathbf{y}\log q_X(T^{-1}(\mathbf{y}))||_2^2]$$

$$= \frac{1}{2}\mathbb{E}_{q_Y}[||\nabla_\mathbf{y}T^{-1}(\mathbf{y})^\top$$
$$(\nabla_{T^{-1}(\mathbf{y})}\log p_X(T^{-1}(\mathbf{y})) - \nabla_{T^{-1}(\mathbf{y})}\log q_X(T^{-1}(\mathbf{y})))||_2^2]$$

$$= \frac{1}{2}\mathbb{E}_{q_X}[||(\nabla_{\boldsymbol{x}}T(\boldsymbol{x}))^{-\top}(\nabla_{\boldsymbol{x}}\log p_X(\boldsymbol{x}) - \nabla_{\boldsymbol{x}}\log q_X(\boldsymbol{x}))||_2^2], \tag{9}$$

where the last step comes from changing the variable to $\boldsymbol{x} = T^{-1}(\mathbf{y})$ and noticing that $\nabla_\mathbf{y}T^{-1}(\mathbf{y}) = (\nabla_{\boldsymbol{x}}T(\boldsymbol{x}))^{-1}$ from the inverse function theorem. This objective coincides with the *diffusion score matching* (Eq.5). Importantly, $DSM_m(q_X, p_X)$ is a valid divergence (i.e. $DSM_m(p_X, q_X) = 0$ iff. $p_X = q_X$) when $m(\boldsymbol{x})$ is an invertible matrix for every $\boldsymbol{x}$. As normalising flow transformations naturally give invertible Jacobian matrices, we can easily extablish the connection $\mathcal{F}(p_Y, q_Y) = DSM_m(p_X, q_X)$ with $m(\boldsymbol{x}) = (\nabla_{\boldsymbol{x}}T(\boldsymbol{x}))^{-1}$. $\square$

We also include additional plots for the student-t example. Specifically, we plot the log likelihood function after the transformation defined by $\boldsymbol{m}(\boldsymbol{x}) = (1 + \frac{(\boldsymbol{x}-\theta)}{0.6})$ in figure 2 (Appendix D).

Similarly, we can prove the connections between DSD (Eq.6) and normalizing flow. The proof is in appendix A.

Theorem 3.2. *For twice differentiable densities $q_X(\boldsymbol{x})$, $p_X(\boldsymbol{x})$, an invertible differentiable transformation $\boldsymbol{T}(\boldsymbol{x}) :$ $\mathbb{R}^D \to \mathbb{R}^D$ and differentiable test function in suitable test function family $\mathcal{H}$: $\boldsymbol{f} : \mathbb{R}^D \to \mathbb{R}^D \in \mathcal{H}$, the diffusion Stein discrepancy (Eq.6) is equivalent to the original Stein discrepancy*

$$\mathcal{S}(q_Y, p_Y) = \sup_{\boldsymbol{g}\in\mathcal{H}'} \mathbb{E}_{q_Y}[\boldsymbol{s}_{p_Y}(\boldsymbol{y})^T\boldsymbol{g}(\boldsymbol{y}) + \nabla_{\boldsymbol{y}}^T\boldsymbol{g}(\boldsymbol{y})] \tag{10}$$

where $\boldsymbol{g}(\boldsymbol{y}) = \boldsymbol{f}(\boldsymbol{T}^{-1}(\boldsymbol{y}))$, $\mathcal{H}'$ is the corresponding function space for $\boldsymbol{g}$, p_Y and q_Y are transformed densities by $\boldsymbol{T}(\cdot)$. The diffusion matrix $\boldsymbol{m}(\boldsymbol{x}) = (\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1}$.

Based on the above two theorems, we formally establish the connections between the *DSM/DSD* with normalizing flows. This gives us an interpretation of the diffusion matrix as the inverse of the Jacobian matrix defined by the flow.

We can further interpret this diffusion using the distributions defined on a Riemannian manifold. In fact, we can show that the *DSM* is equivalent to the original SM performed on two densities defined on Riemannian manifolds with the metric tensor $\boldsymbol{G}(\boldsymbol{x}) = \boldsymbol{m}^{-T}(\boldsymbol{x})\boldsymbol{m}(\boldsymbol{x})^{-1}$

3.2. Interpreting DSM using Riemannian manifold

Assume we have a Riemannian manifold $(\mathcal{M}, \boldsymbol{g})$ with Riemannian metric tensor $\boldsymbol{g}$. For each point $\boldsymbol{a} \in \mathcal{M}$, we assume it has a local coordinates $\boldsymbol{x}_a = [x_a^1, \ldots, x_a^D]$. We can prove the following proposition:

Proposition 3.1. *Assume the Riemannian manifold $(\mathcal{M}, \boldsymbol{g})$ as defined above, we define two probability measures $\mathbb{Q}$, $\mathbb{P}$. We denote the corresponding densities (in terms of local coordinates $\boldsymbol{x}$) w.r.t. Riemannian manifold as $\tilde{p}(\boldsymbol{x}) = \frac{d\mathbb{P}}{d\mathcal{M}(\boldsymbol{x})}$ and $\tilde{q}(\boldsymbol{x}) = \frac{d\mathbb{Q}}{d\mathcal{M}(\boldsymbol{x})}$. Then, the score matching loss for $\tilde{p}$*

and $\tilde{q}$ is

$$\mathcal{F}_{\mathcal{M}}(\tilde{q}, \tilde{p}) = \frac{1}{2}\mathbb{E}_q[\boldsymbol{\Delta}(\boldsymbol{x})^T \boldsymbol{G}(\boldsymbol{x})^{-1}\boldsymbol{\Delta}(\boldsymbol{x})] \qquad (11)$$

where $p(\boldsymbol{x}) = \frac{d\mathbb{P}}{d\mathcal{M}(\boldsymbol{x})}\frac{d\mathcal{M}(\boldsymbol{x})}{d\boldsymbol{x}}$, $q(\boldsymbol{x})$ is defined similarly, and $\boldsymbol{\Delta}(\boldsymbol{x}) = \boldsymbol{s}_p(\boldsymbol{x}) - \boldsymbol{s}_q(\boldsymbol{x})$. $\boldsymbol{G}(\boldsymbol{x})$ is an symmetric positive definite matrix representing the Riemannian metric tensor. Particularly, if $\boldsymbol{G}(\boldsymbol{x}) = \boldsymbol{m}(\boldsymbol{x})^{-T}\boldsymbol{m}(\boldsymbol{x})^{-1}$, then $\mathcal{F}_{\mathcal{M}}(\tilde{q}, \tilde{p})$ is equivalent to DSM (Eq.5) with diffusion matrix $\boldsymbol{m}(\boldsymbol{x})$.

The proof is in appendix B.

This result is more general than theorem 3.1. Specifically, theorem 3.1 only proves a sufficient condition for the DSM to be a valid discrepancy. Namely, if we have an invertible flow, the corresponding diffusion matrix $\boldsymbol{m}(x)$ must be invertible. However, the converse is not true. On the other hand, proposition 3.1 only requires $\boldsymbol{m}(\boldsymbol{x})$ to be invertible, which is more general. Indeed, from the topological point of view, if we have an invertible and differentiable flow $\boldsymbol{T}$, then the transformed space (Riemannian manifold) is actually diffeomorphic to the original space (e.g. $\mathbb{R}^D$). Thus, this flow can be viewed as a special case of Gemici et al. (2016). But in general, Riemannian manifold may not be diffeomorphic to $\mathbb{R}^D$, which explains why theorem 3.1 is only a sufficient condition.

3.3. Continuous DSM with ODE flow

Previous sections assume a deterministic transformation $\boldsymbol{T}(\boldsymbol{x})$. Recent work has shown promising results for continuous flows characterised by an ODE (Chen et al., 2018; Grathwohl et al., 2018). Namely, we can consider a transformation

$$d\boldsymbol{x} = \boldsymbol{g}(\boldsymbol{x}(t))dt \qquad (12)$$

where $\boldsymbol{g}(\boldsymbol{x}(t))$ is a deterministic drift that is uniformly Lipschitz continuous w.r.t. $\boldsymbol{x}$. We define p_t and q_t to be the corresponding densities for $\boldsymbol{x}(t)$. Inspired by Chen et al. (2018), we can characterise the instantaneous change of the score matching loss $\frac{d\mathcal{F}(q_t, p_t)}{dt}$ by the following proposition:

Proposition 3.2. *Let $p_t(\boldsymbol{x}(t))$, $q_t(\boldsymbol{x}(t))$ be two probability density functions, where $\boldsymbol{x}(t)$ is characterized by an ODE defined in eq.12. Assume $\boldsymbol{g}(\boldsymbol{x}(t))$ is uniformly Lipschitz continuous w.r.t. $\boldsymbol{x}(t)$. Then, the instantaneous change of score matching loss follows:*

$$\frac{d\mathcal{F}(q_t, p_t)}{dt} = -\frac{1}{2}\mathbb{E}_{q_t}[\Delta(\boldsymbol{x})^T(\nabla_{\boldsymbol{x}}\boldsymbol{g}(\boldsymbol{x})+\nabla_{\boldsymbol{x}}\boldsymbol{g}(\boldsymbol{x})^T)\Delta(\boldsymbol{x})]$$
$$(13)$$

where $\Delta(\boldsymbol{x}) = \boldsymbol{s}_{p_t}(\boldsymbol{x}) - \boldsymbol{s}_{q_t}(\boldsymbol{x})$.

The proof is in appendix C.

4. Conclusion

In this paper, we discuss the connections of the *diffusion score matching* and *diffusion Stein discrepancy* to normalizing flows. Specifically, we theoretically prove that the DSM (or DSD) is equivalent to performing the original *score matching* (or *Stein discrepancy*) on the transformed densities. The diffusion matrix $\boldsymbol{m}(\boldsymbol{x})$ is defined by the inverse of the flow's Jacobian matrix. We also establish the connection of DSM with densities defined on Riemannian manifolds by showing that the diffusion matrix is closely related to the Riemannian metric tensor. In the end, we extend the DSM by continuous flow, and derive an ODE characterizing its instantaneous changes. By building the connections, we hope to shed lights on developing training method for the diffusion matrix to enable the practical usage of DSM (or DSD) for large models.

References

Barp, A., Briol, F.-X., Duncan, A. B., Girolami, M., and Mackey, L. Minimum stein discrepancy estimators. *arXiv preprint arXiv:1906.08283*, 2019.

Chen, R. T., Rubanova, Y., Bettencourt, J., and Duvenaud, D. Neural ordinary differential equations. *arXiv preprint arXiv:1806.07366*, 2018.

Chwialkowski, K., Strathmann, H., and Gretton, A. A kernel test of goodness of fit. In *International conference on machine learning*, pp. 2606–2615. PMLR, 2016.

Fisher, M., Nolan, T., Graham, M., Prangle, D., and Oates, C. Measure transport with kernel stein discrepancy. In *International Conference on Artificial Intelligence and Statistics*, pp. 1054–1062. PMLR, 2021.

Gemici, M. C., Rezende, D., and Mohamed, S. Normalizing flows on riemannian manifolds. *arXiv preprint arXiv:1611.02304*, 2016.

Gorham, J. *Measuring sample quality with Stein's method*. Stanford University, 2017.

Gorham, J., Duncan, A. B., Vollmer, S. J., Mackey, L., et al. Measuring sample quality with diffusions. *Annals of Applied Probability*, 29(5):2884–2928, 2019.

Grathwohl, W., Chen, R. T., Bettencourt, J., Sutskever, I., and Duvenaud, D. Ffjord: Free-form continuous dynamics for scalable reversible generative models. *arXiv preprint arXiv:1810.01367*, 2018.

Grathwohl, W., Wang, K.-C., Jacobsen, J.-H., Duvenaud, D., and Zemel, R. Learning the stein discrepancy for training and evaluating energy-based models without sampling. In *International Conference on Machine Learning*, pp. 3732–3747. PMLR, 2020.

Hu, T., Chen, Z., Sun, H., Bai, J., Ye, M., and Cheng, G. Stein neural sampler. *arXiv preprint arXiv:1810.03545*, 2018.

Hyvärinen, A. and Dayan, P. Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6(4), 2005.

Johnson, O. *Information theory and the central limit theorem*. World Scientific, 2004.

Liu, Q. and Wang, D. Stein variational gradient descent: A general purpose bayesian inference algorithm. *arXiv preprint arXiv:1608.04471*, 2016.

Liu, Q., Lee, J., and Jordan, M. A kernelized stein discrepancy for goodness-of-fit tests. In *International conference on machine learning*, pp. 276–284. PMLR, 2016.

Song, Y. and Ermon, S. Generative modeling by estimating gradients of the data distribution. *arXiv preprint arXiv:1907.05600*, 2019.

Song, Y., Garg, S., Shi, J., and Ermon, S. Sliced score matching: A scalable approach to density and score estimation. In *Uncertainty in Artificial Intelligence*, pp. 574–584. PMLR, 2020a.

Song, Y., Sohl-Dickstein, J., Kingma, D. P., Kumar, A., Ermon, S., and Poole, B. Score-based generative modeling through stochastic differential equations. *arXiv preprint arXiv:2011.13456*, 2020b.

Wenliang, L., Sutherland, D., Strathmann, H., and Gretton, A. Learning deep kernels for exponential family densities. In *International Conference on Machine Learning*, pp. 6737–6746. PMLR, 2019.

A. Proof of theorem 3.2

Proof. Let's first define the Stein operator as

$$\mathcal{S}_{p_Y}[\boldsymbol{g}] = \boldsymbol{s}_{p_Y}(\boldsymbol{y})^T\boldsymbol{g}(\boldsymbol{x}) + \nabla_{\boldsymbol{y}}^T\boldsymbol{g}(\boldsymbol{y}) \qquad (14)$$

for the test function $\boldsymbol{g}(\boldsymbol{y})$ and density $p_Y(\mathbf{y})$. Thus, the Stein discrepancy can be rewritten as

$$\mathcal{S}(q_Y, p_Y) = \sup_{\boldsymbol{g}\in\mathcal{H}} \mathbb{E}_{q_Y}[\mathcal{S}_{p_Y}[\boldsymbol{g}]] \qquad (15)$$

In the following, we will focus on the Stein operator. From the change of variable formula $\boldsymbol{y} = \boldsymbol{T}(\boldsymbol{x})$, we have

$$p_Y(\mathbf{y}) = p_X(\boldsymbol{T}^{-1}(\mathbf{y}))\left|\frac{\partial\boldsymbol{T}^{-1}(\mathbf{y})}{\partial\mathbf{y}}\right|, \quad \boldsymbol{g}(\mathbf{y}) = \boldsymbol{f}(\boldsymbol{T}^{-1}(\mathbf{y})) \qquad (16)$$

Now we can rewrite the Stein operator:

$$\mathcal{S}_{p_Y}[\boldsymbol{g}] = \nabla_{\mathbf{y}}\log p_Y(\mathbf{y})^T\boldsymbol{g}(\mathbf{y}) + \nabla_{\mathbf{y}}^T\boldsymbol{g}(\mathbf{y})$$
$$=\nabla_{\mathbf{y}}\log p_X(\boldsymbol{T}^{-1}(\mathbf{y}))^T\boldsymbol{g}(\mathbf{y}) + (\nabla_{\mathbf{y}}\log\left|\frac{\partial\boldsymbol{T}^{-1}(\mathbf{y})}{\partial\mathbf{y}}\right|)^T\boldsymbol{g}(\mathbf{y})$$
$$+\nabla_{\mathbf{y}}\boldsymbol{g}(\mathbf{y})$$
$$= \left[(\nabla_{\mathbf{y}}\boldsymbol{T}^{-1}(\mathbf{y}))^T(\nabla_{\boldsymbol{T}^{-1}(\mathbf{y})}\log p_X(\boldsymbol{T}^{-1}(\mathbf{y})))\right]^T\boldsymbol{g}(\mathbf{y})$$
$$+ \underbrace{(\nabla_{\mathbf{y}}\log\left|\frac{\partial\boldsymbol{T}^{-1}(\mathbf{y})}{\partial\mathbf{y}}\right|)^T\boldsymbol{g}(\mathbf{y})}_{\textcircled{1}}$$
$$+ Tr[(\nabla_{\mathbf{y}}\boldsymbol{T}^{-1}(\mathbf{y}))\nabla_{\boldsymbol{T}^{-1}(\mathbf{y})}\boldsymbol{f}(\boldsymbol{T}^{-1}(\mathbf{y}))] \qquad (17)$$

The second equality is from the chain rule and definition of divergence operator ∇^T. For the layout of the matrix calculus, we follow the column vector layout as the following: for a function $\boldsymbol{h}:\mathbb{R}^D\to\mathbb{R}$, and $\boldsymbol{f}:\mathbb{R}^D\to\mathbb{R}^N$, we have

$$\frac{\partial h(\boldsymbol{x})}{\partial\boldsymbol{x}} = \begin{bmatrix}\frac{\partial h(\boldsymbol{x})}{\partial x_1}\\ \vdots\\ \frac{\partial h(\boldsymbol{x})}{\partial x_D}\end{bmatrix}$$
$$\frac{\partial\boldsymbol{f}(\boldsymbol{x})}{\partial\boldsymbol{x}} = \begin{bmatrix}\frac{\partial f_1(\boldsymbol{x})}{\partial x_1} & \cdots & \frac{\partial f_1(\boldsymbol{x})}{\partial x_D}\\ \vdots & \vdots & \vdots\\ \frac{\partial f_N(\boldsymbol{x})}{\partial x_1} & \cdots & \frac{\partial f_N(\boldsymbol{x})}{\partial x_D}\end{bmatrix} \qquad (18)$$

Now, we focus on $\textcircled{1}$ term:

$$\nabla_{\mathbf{y}}\log\left|\frac{\partial\boldsymbol{T}^{-1}(\mathbf{y})}{\partial\mathbf{y}}\right| = Tr[(\nabla_{\mathbf{y}}\boldsymbol{T}^{-1}(\mathbf{y}))^{-1}\nabla_{\mathbf{y}}\nabla_{\mathbf{y}}\boldsymbol{T}^{-1}(\mathbf{y})]$$
$$= Tr[\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x})\nabla_{\mathbf{y}}(\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1}]$$
$$= Tr[\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x})\nabla_{\mathbf{y}}\boldsymbol{T}^{-1}(\mathbf{y})\nabla_{\boldsymbol{x}}(\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1}]$$
$$= Tr[\nabla_{\boldsymbol{x}}(\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1}] \qquad (19)$$

where we use the inverse function theorem $\nabla_{\mathbf{y}}\boldsymbol{T}^{-1}(\mathbf{y}) = (\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1}$. In addition, we define $\nabla_{\boldsymbol{x}}^T(\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1} = Tr[\nabla_{\boldsymbol{x}}(\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1}]$.

So we can set $\boldsymbol{m}(\boldsymbol{x}) = (\nabla_{\boldsymbol{x}}\boldsymbol{T}(\boldsymbol{x}))^{-1}$, we can obtain:

$$\mathcal{S}_{p_Y}[\boldsymbol{g}] = (\boldsymbol{m}(\boldsymbol{x})^T\boldsymbol{s}_p(\boldsymbol{x}))^T\boldsymbol{f}(\boldsymbol{x}) + (\nabla_{\boldsymbol{x}}^T\boldsymbol{m}(\boldsymbol{x}))^T\boldsymbol{f}(\boldsymbol{x})$$
$$+ Tr[\boldsymbol{m}(\boldsymbol{x})\nabla_x\boldsymbol{f}(\boldsymbol{x})]$$
$$= (\boldsymbol{m}(\boldsymbol{x})^T\boldsymbol{s}_p(\boldsymbol{x}))^T\boldsymbol{f}(\boldsymbol{x}) + \nabla_{\boldsymbol{x}}^T[\boldsymbol{m}(\boldsymbol{x})\boldsymbol{f}(\boldsymbol{x})] \qquad (20)$$

which is exactly the same as the inner part of DSD (Eq.6). So with change of variable formula, we can easily show

$$\mathcal{S}(q_Y, p_Y) = DSD_m(q_X, p_X) \qquad (21)$$

$\square$

B. Proof of proposition 3.1

With the definition of the Riemannian manifold $(\mathcal{M}, \boldsymbol{g})$, for any point $\boldsymbol{a}\in\mathcal{M}$ with local coordinates $\boldsymbol{x}\in\mathbb{R}^D$, and two vectors $\boldsymbol{u}, \boldsymbol{v}$ from its tangent plane $T_a\mathcal{M}$, we can represents $\boldsymbol{u}, \boldsymbol{v}$ using the basis $(\frac{\partial}{\partial x_i})_{\boldsymbol{a}}$ as

$$\boldsymbol{u} = \sum_{i=1}^D u_i(\frac{\partial}{\partial x_i})_{\boldsymbol{a}}, \quad \boldsymbol{v} = \sum_{i=1}^D v_i(\frac{\partial}{\partial x_i})_{\boldsymbol{a}} \qquad (22)$$

The inner product defined by the metric $\boldsymbol{g}$ can be expressed as

$$\boldsymbol{g}(\boldsymbol{u}, \boldsymbol{v}) = \sum_{i,j}^D u_iv_j\langle(\frac{\partial}{\partial x_i})_{\boldsymbol{a}}, (\frac{\partial}{\partial x_j})_{\boldsymbol{a}}\rangle_g = \sum_{i,j}^D u_ig_{ij}(\boldsymbol{x})v_j \qquad (23)$$

where $g_{ij}(\boldsymbol{x})$ is the $ij-$th element of matrix $\boldsymbol{G}(\boldsymbol{x})$ and $\langle\cdot,\cdot\rangle_g$ is the inner product defined by Riemannian metric $\boldsymbol{g}$.

We assume the measure $\mathcal{M}(\boldsymbol{x})$ is absolutely continuous w.r.t. Lebesgue measure, then we have the following change of variable formula

$$d\mathcal{M}(\boldsymbol{x}) = \sqrt{|\boldsymbol{G}(\boldsymbol{x})|}d\boldsymbol{x} \qquad (24)$$

Then we can represents the densities $\tilde{p}, \tilde{q}$ under Lebessgue measure

$$p(\boldsymbol{x}) = \frac{d\mathbb{P}}{d\mathcal{M}(\boldsymbol{x})}\frac{d\mathcal{M}(\boldsymbol{x})}{d\boldsymbol{x}} = \tilde{p}(\boldsymbol{x})\sqrt{|\boldsymbol{G}(\boldsymbol{x})|} \qquad (25)$$

and $q(\boldsymbol{x})$ is defined accordingly. The score matching loss for $\tilde{p}$ and $\tilde{q}$ is

$$\mathcal{F}_{\mathcal{M}}(\tilde{q}, \tilde{p}) = \frac{1}{2}\int\tilde{q}(\boldsymbol{x})||\nabla\log\tilde{p}(\boldsymbol{x}) - \nabla\log\tilde{q}(\boldsymbol{x})||_g^2d\mathcal{M}(\boldsymbol{x})$$
$$= \frac{1}{2}\int q(\boldsymbol{x})||\nabla\log\tilde{p}(\boldsymbol{x}) - \nabla\log\tilde{q}(\boldsymbol{x})||_g^2d\boldsymbol{x} \qquad (26)$$

Now let's define $\nabla \log \tilde{p}(\boldsymbol{x})$. From the basics of Riemannian manifold, for a point $\boldsymbol{a} \in \mathcal{M}$ with local coordinate $\boldsymbol{x}$, and $\mathcal{X}$ is a vector field on $\mathcal{M}$, we have the following definition

$$\langle \sum_{i=1}^{D} (\nabla \log \tilde{p}(\boldsymbol{x}))_i (\frac{\partial}{\partial x_i})_a, \sum_{j=1}^{D} \mathcal{X}_j (\frac{\partial}{\partial x_j}) \rangle_g = \sum_{i=1}^{D} \mathcal{X}_i \frac{\partial \log \tilde{p}}{\partial x_i} \tag{27}$$

Written in terms of matrix form, assume $\boldsymbol{X} = [\mathcal{X}_1, \ldots, \mathcal{X}_D]^T$, and $g_{ij}(\boldsymbol{x})$ is the element of symmetric positive definite matrix $\boldsymbol{G}(\boldsymbol{x})$, we have

$$(\nabla \log \tilde{p})^T \boldsymbol{G}(\boldsymbol{x}) \boldsymbol{X} = (\frac{\partial \log \tilde{p}}{\partial \boldsymbol{x}})^T \boldsymbol{X}$$
$$\Longrightarrow \nabla \log \tilde{p} = \boldsymbol{G}^{-1}(\boldsymbol{x})(\frac{\partial \log \tilde{p}}{\partial \boldsymbol{x}}) \tag{28}$$

Therefore, we have

$$||\nabla \log \tilde{p}(\boldsymbol{x}) - \nabla \log \tilde{q}(\boldsymbol{x})||_g^2$$
$$= \langle \nabla \log \tilde{p}(\boldsymbol{x}) - \nabla \log \tilde{q}(\boldsymbol{x}), \nabla \log \tilde{p}(\boldsymbol{x}) - \nabla \log \tilde{q}(\boldsymbol{x}) \rangle_g$$
$$= \langle \boldsymbol{G}^{-1}(\boldsymbol{x}) \underbrace{(\frac{\partial \log \tilde{p}}{\partial \boldsymbol{x}} - \frac{\partial \log \tilde{q}}{\partial \boldsymbol{x}})}_{\tilde{\Delta}(\boldsymbol{x})}, \boldsymbol{G}^{-1}(\boldsymbol{x})(\frac{\partial \log \tilde{p}}{\partial \boldsymbol{x}} - \frac{\partial \log \tilde{q}}{\partial \boldsymbol{x}}) \rangle_g$$
$$= \tilde{\Delta}(\boldsymbol{x})^T \boldsymbol{G}^{-1}(\boldsymbol{x}) \boldsymbol{G}(\boldsymbol{x}) \boldsymbol{G}^{-1}(\boldsymbol{x}) \tilde{\Delta}(\boldsymbol{x})$$
$$= \tilde{\Delta}(\boldsymbol{x})^T \boldsymbol{G}^{-1}(\boldsymbol{x}) \tilde{\Delta}(\boldsymbol{x}) \tag{29}$$

By change of variable formula, it is also easy to show that

$$\tilde{\Delta}(\boldsymbol{x}) = \underbrace{(\frac{\partial \log p}{\partial \boldsymbol{x}} - \frac{\partial \log q}{\partial \boldsymbol{x}})}_{\Delta(\boldsymbol{x})} \tag{30}$$

Therefore, we have

$$||\nabla \log \tilde{p}(\boldsymbol{x}) - \nabla \log \tilde{q}(\boldsymbol{x})||_g^2 = \Delta^T(\boldsymbol{x}) \boldsymbol{G}^{-1}(\boldsymbol{x}) \Delta(\boldsymbol{x}) \tag{31}$$

Substitute back to $\mathcal{F}_{\mathcal{M}}(\tilde{q}, \tilde{p})$ (Eq.26), we can obtain the result. Particularly, compare to *DSM* (Eq.5), we can observe that if $\boldsymbol{G}(\boldsymbol{x}) = \boldsymbol{m}(\boldsymbol{x})^{-T} \boldsymbol{m}(\boldsymbol{x})^{-1}$, the $\mathcal{F}_{\mathcal{M}}(\tilde{q}, \tilde{p})$ is equivalent to *DSM*. Indeed, as $\boldsymbol{m}(\boldsymbol{x}) \in \mathbb{R}^{D \times D}$ is an invertible matrix, then $\boldsymbol{m}(\boldsymbol{x})^{-T} \boldsymbol{m}(\boldsymbol{x})^{-1}$ must be symmetric positive definite, which satisfies the requirements for $\boldsymbol{G}(\boldsymbol{x})$.

C. Proof of proposition 3.2

An ODE flow is defined by the solution of the following ODE:

$$d\boldsymbol{x} = \boldsymbol{g}(\boldsymbol{x})dt \tag{32}$$

with $\boldsymbol{g}(\boldsymbol{x})$ the deterministic drift term. Let us consider the forward Euler discretisation of the ODE, which gives

$$\boldsymbol{x}(t + \delta) = \boldsymbol{x}(t) + \delta \boldsymbol{g}(\boldsymbol{x}(t)) := \boldsymbol{T}_\delta(\boldsymbol{x}(t)). \tag{33}$$

With $\delta \approx 0$ we see that T_δ is an invertible transformation. Now consider $\mathbf{y} = \boldsymbol{x}(t + \delta)$ and $\boldsymbol{x}(t) = \boldsymbol{x}$. This again pushes $p_X(\boldsymbol{x})$ and $q_X(\boldsymbol{x})$ to $p_Y(\mathbf{y})$ and $q_Y(\mathbf{y})$, respectively. Therefore we can reuse results from theorem 3.1 and derive

$$F(p_Y, q_Y) = \frac{1}{2} \mathbb{E}_{q_X}[||\boldsymbol{m}(\boldsymbol{x})^T (\boldsymbol{s}_{p_X}(\boldsymbol{x}) - \boldsymbol{s}_{q_X}(\boldsymbol{x}))||^2], \tag{34}$$

where $\boldsymbol{m}(\boldsymbol{x}) = (\nabla_{\boldsymbol{x}} \boldsymbol{T}_\delta(\boldsymbol{x}))^{-1}$ Notice that $T_\delta(\boldsymbol{x}) = \boldsymbol{x}$ when $\delta = 0$. This means we can compute the change of score matching at time t as:

$$\frac{\partial}{\partial t} F(p_Y, q_Y) = \lim_{\delta \to 0^+} \frac{F(p_Y, q_Y) - F(p_X, q_X)}{\delta}$$
$$= \frac{1}{2} \lim_{\delta \to 0^+} \mathbb{E}_{q_X(\boldsymbol{x})}[\Delta(\boldsymbol{x})^\top \delta^{-1}(m(\boldsymbol{x})m(\boldsymbol{x})^\top - \mathbf{I})\Delta(\boldsymbol{x})], \tag{35}$$

with $\Delta(\boldsymbol{x}) = \nabla_{\boldsymbol{x}} \log p_X(\boldsymbol{x}) - \nabla_{\boldsymbol{x}} \log q_X(\boldsymbol{x})$. As $\nabla_{\boldsymbol{x}} T_\delta(\boldsymbol{x}) = \mathbf{I} + \delta \nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x})$, simple calculation shows that

$$\delta^{-1}(m(\boldsymbol{x})m(\boldsymbol{x})^\top - \mathbf{I})$$
$$= \delta^{-1}[[(\mathbf{I} + \delta \nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x}))^\top (\mathbf{I} + \delta \nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x}))]^{-1} - \mathbf{I}]$$
$$= \delta^{-1}[[\mathbf{I} + \delta(\nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x}) + \nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x})^\top) + \mathcal{O}(\delta^2)]^{-1} - \mathbf{I}]$$
$$= -[\nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x}) + \nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x})^\top + \mathcal{O}(\delta)]$$
$$[\mathbf{I} + \delta(\nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x}) + \nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x})^\top) + \mathcal{O}(\delta^2)]^{-1}, \tag{36}$$

which leads to

$$\frac{\partial}{\partial t} F(p_Y, q_Y)$$
$$= \lim_{\delta \to 0^+} \frac{1}{2} \mathbb{E}_{q_X(\boldsymbol{x})}[\Delta(\boldsymbol{x})^\top \delta^{-1}(m(\boldsymbol{x})m(\boldsymbol{x})^\top - \mathbf{I})\Delta(\boldsymbol{x})]$$
$$= -\frac{1}{2} \mathbb{E}_{q_X(\boldsymbol{x})}[\Delta(\boldsymbol{x})^\top (\nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x}) + \nabla_{\boldsymbol{x}} \boldsymbol{g}(\boldsymbol{x})^\top)\Delta(\boldsymbol{x})] \tag{37}$$

As this quantifies the instantaneous changes, replacing p_t and q_t for p_Y, p_X, q_Y and q_X gives the instantaneous change of score matching loss.

D. Additional plots

From the motivating example and theorem 3.1, we know $\boldsymbol{m}(\boldsymbol{x}) = (1 + \frac{(\boldsymbol{x}-\theta)^2}{b})$. Therefore, by simple calculus, the corresponding transformation $y = \boldsymbol{T}(\boldsymbol{x})$ can be defined as

$$\mathbf{y} = \boldsymbol{T}(\boldsymbol{x}) = \frac{1}{b\sqrt{b}} \tan^{-1}(\frac{\boldsymbol{x} - \theta}{\sqrt{b}})$$
$$\boldsymbol{x} = \boldsymbol{T}^{-1}(\mathbf{y}) = \sqrt{b} \tan(b\sqrt{b}\mathbf{y}) + \theta \tag{38}$$

Let's define the transformed densities $p_Y(\mathbf{y})$ and $q_Y(\mathbf{y})$ as

$$p_Y(\mathbf{y}) = p(\boldsymbol{T}^{-1}(\mathbf{y})) |\nabla_{\mathbf{y}} \boldsymbol{T}^{-1}(\mathbf{y})|$$
$$q_Y(\mathbf{y}) = q(\boldsymbol{T}^{-1}(\mathbf{y})) |\nabla_{\mathbf{y}} \boldsymbol{T}^{-1}(\mathbf{y})| \tag{39}$$

Therefore, we can plot the log likelihood for the original densities p,q and transformed densities p_Y, q_Y as figure 2.

Figure 2. **Left**:The log likelihood plot for original densities q, p. **Middle**: The log likelihood function for transformed density p_Y **Right**: The log likelihood function for q_Y. We choose $\theta = -2.5$ and $b = 0.6$. Notice that the transformed densities p_Y, q_Y are periodic as we consider $y \in \mathbb{R}$. This won't happen if we consider $\boldsymbol{y} = \boldsymbol{T}(\boldsymbol{x})$. Because all $\boldsymbol{x}$ value will be squeezed inside the period containing 0, i.e. $\boldsymbol{y}$ will inside $[-3.37, 3.37]$ in this case.

In this case, we set $p(\boldsymbol{x})$ has the mean -2.5 with the same scale 0.3 as q, whereas q has mean 0. For the transformation $\boldsymbol{T}$, we set $\theta = -2.5$ with $b = 0.6$.