CALM-IT: Generating Realistic Long-Form Motivational Interviewing Dialogues with Dual-Actor Conversational Dynamics Tracking

Abstract

Therapeutic dialogue is not a sequence of isolated responses: client goals, motivation, resistance, and therapeutic alliance evolve over time. Yet current LLM-based mental health dialogue systems often lack explicit mechanisms for tracking these dynamics across extended interactions, which can lead to poorly timed interventions or premature goal resolution. We introduce CALM-IT, a framework for generating and evaluating long-form Motivational Interviewing dialogues through explicit modeling of evolving client and counselor states, guiding both counseling strategy selection and utterance generation. We evaluate CALM-IT on a large-scale corpus of 8,232 synthetic dialogues spanning multiple dialogue lengths and frameworks. Compared with all baselines, CALM-IT achieves the best performance on most MITI 4.2 global ratings, including Empathy, Partnership, and Softening Sustain Talk, as well as on other key performance metrics while exhibiting minimal performance degradation as dialogue length increases. Notably, although CALM-IT initiates fewer change-directed prompts, it produces the highest client acceptance rate (64.3%) on average across different length conditions. We release a reproducible generation framework, a MITI-grounded process-level evaluation protocol, and a large-scale synthetic corpus for studying therapeutic LLMs under realistic long-form interaction conditions.

1 Introduction

Large language models (LLMs) are increasingly used for mental health support through chatbots and conversational agents that many people turn to for emotional guidance. In the United States alone, around 1 in 8 young adults report using generative AI when feeling sad, angry, or anxious (McBain et al., 2025). Yet, studies and reports raise concerns about potential harms. AI chatbots may fail to respond appropriately to suicidal ideation, validate delusional beliefs, or cause severe distress (Pichowicz et al., 2025; Hall, 2025; PBS NewsHour, 2025). A key limitation driving these failures is that current LLMs struggle to sustain high-quality and appropriate dialogue over extended interactions. They often optimize for locally appropriate next turns rather than tracking the evolving relational, motivational, and emotional state of the conversation. In therapeutic interactions, this may lead systems to persist with interventions when rapport has deteriorated. It can also cause models to move toward planning and resolution earlier than process models suggest. Furthermore, privacy protections, legal constraints, and stigma severely limit the collection of real therapy conversations (Mayer et al., 2022). This leaves models trained and evaluated on short, simplified exchanges that fail to capture nonlinear interactional dynamics over extended sessions.

Synthetic dialogue simulation offers a scalable path for training and stress-testing these models, given that synthetic data has improved large language model performance in domains like code generation or mathematical reasoning (Wang et al., 2023; Ding et al., 2023). However, developing resilient agents requires exposure to the fluid conversational dynamics that govern human interactions. Current clinical frameworks establish that core factors like motivation, rapport, and goal alignment must evolve gradually and bidirectionally across both participants over the course of a session (Miller and Rollnick, 2013; Apodaca and Longabaugh, 2009). Existing simulators fail to model this joint evolution, meaning they cannot provide the realistic training environments required to teach models how to manage complex trajectories. Client simulators such as Roleplay-doh (Louie et al., 2024) and Patient-Ψ (Wang et al., 2024b) rely on fixed personas that respond cooperatively and predictably, creating dialogues that lack resistance or emotional fluctuation. Therapist agents, meanwhile, focus on locally appropriate

responses without tracking how the broader relational landscape shifts over time (Kim et al., 2025a; Yang et al., 2025a). Without explicit modeling of these dual-actor dynamics as an interdependent system, synthetic data cannot replicate the exact trajectory failures introduced above, including premature convergence, motivational drift, and poorly timed redirections. This deficiency leaves models undertrained for real-world interactions and limits our ability to evaluate their long-horizon stability.

To address this, we introduce CALM-IT (Figure 1), a framework for generating synthetic long-form Motivational Interviewing (MI) dialogues via dual-actor conversational dynamics tracking. Both agents maintain latent variables, including rapport, inferred mental state, motivation, and goals, which govern strategy selection and response generation. We evaluate CALM-IT within MI because its process fidelity depends on negotiating resistance and ambivalence over time rather than isolated turn quality. Beyond generation, CALM-IT introduces process-level evaluation metrics to assess multi-turn trajectory properties, including directionality, goal alignment, and redirection effectiveness. We investigate the following:

- **RQ1:** Does explicit conversational state tracking improve the quality of generated synthetic MI transcripts compared to existing frameworks?

- **RQ2:** Does dynamics-aware dialogue generation produce more MI-consistent trajectory shifts toward change, reflected in client responses?

Our contributions are: (1) CALM-IT, a reproducible dual-actor framework incorporating dual-actor conversational dynamics tracking; (2) a process-level protocol for comprehensively evaluating synthetic MI transcripts; and (3) a large-scale CALM-IT-generated synthetic MI corpus, showing improved client acceptance and change talk.

2 Related Works

2.1 Synthetic Data for Healthcare LLMs

Prior work has generated synthetic data for clinical settings spanning medical dialogues, EHR sequences, and clinical notes (Wang et al., 2024c; He et al., 2020; Theodorou et al., 2023; Yoon et al., 2023; Singhal et al., 2023), with more recent approaches extending to dual-actor multi-turn generation (Wang et al., 2024a; Sviridov et al., 2025). While some exceptions like NoteChat (Wang et al., 2024a) and 3MDBench (Sviridov et al., 2025) also adopt dual-actor generation pipelines, both target medical dialogue, where interactions are primarily oriented toward information gathering and diagnosis rather than relational change processes. Neither NoteChat nor 3MDBench model this joint evolution explicitly; both assign fixed roles without tracking how interactional state shifts across turns, leaving failure modes such as premature convergence and motivational drift unaddressed.

In counseling domains, efforts include empathy-labeled dialogues, multi-turn mental health exchanges, Cognitive Behavioral Therapy (CBT)-oriented datasets, and MI-focused resources (Lozoya et al., 2025; Qiu et al., 2024; Lee et al., 2024; Mahmood et al., 2025; Jörke et al., 2025; Kim et al., 2025a; Chandra et al., 2025). While these approaches capture key structural and stylistic properties of therapeutic dialogue, they focus on local coherence or static datasets, leaving failure modes such as premature convergence and motivational drift invisible in the resulting generations.

2.2 Agentic LLMs for Mental Health Support

Research on agentic LLMs for mental health support includes simulated clients, therapist agents, and structured dialogue frameworks (Wang et al., 2024b; Louie et al., 2024; Yang et al., 2025a; Kim et al., 2025a; Vu et al., 2025). Despite these advances, prior systems share a critical limitation: fixed client personas and locally optimized therapist responses neither track how conversational state evolves across participants over time nor model how rapport, motivation, and goal alignment shift as the dialogue unfolds. In real-world counseling, these dynamics are not incidental; they are central to therapeutic change (Miller and Rollnick, 2013; Amrhein et al., 2003; Apodaca and Longabaugh, 2009). As shown in Table 1, `CALM-IT` is the only framework among those compared that combines conversational dynamics tracking, cognitively grounded client profiles, long-form generation, and dual-actor action selection.

3 The CALM-IT Framework

`CALM-IT` generates long-form Motivational Interviewing (MI) dialogues by explicitly modeling conversational dynamics that evolve over the course of a therapeutic interaction. We define conversational dynamics as the evolution of interactional properties such as rapport, motivation, and goal alignment over the course of a dialogue (Eshghi

Figure 1: Architecture of the CALM-IT framework. Therapist and client agents interact through state-dependent turns, with latent conversational dynamics inferred and updated throughout the dialogue to guide generation and evaluation of synthetic long-form MI interactions.

et al., 2012), and operationalize three markers below. Without explicit state tracking, LLM-based therapist agents converge prematurely, offering action plans and affirmations before rapport has been built or ambivalence explored.

- **Interpersonal Alignment.** Grounded in Rapport Management Theory (Spencer-Oatey, 2004), CALM-IT represents relational alignment as continuously evolving rather than binary, modeled via *rapport* (r) and the patient's *system-1 evaluation* of the therapist's preceding turn (q) (Kahneman, 2011). System-1 evaluation triggers stochastically ($p = 0.5$, empirically tuned) to balance affective reactivity against conversational stability.

- **Mental-state Inference.** Inspired by Theory of Mind research (Baron-Cohen, 1995), CALM-IT maintains continuously updated representations of the patient's internal states, including *background* (b), *emotional state* (e),

and *stage of change* (c), alongside the therapist's corresponding inferences ($\hat{b}, \hat{e}, \hat{c}$), explicitly distinguishing inferred mental states from unobserved psychological ground truth.

- **Joint Activity Structure.** Motivated by theories of joint action (Clark, 1996), CALM-IT tracks alignment in conversational goals through updates to the patient's *short-term goal* (g) and the therapist's *inferred goal* ($\hat{g}$), enabling detection of goal misalignment and initiation of pivoting when progress stalls.

Together, these markers form a jointly evolving state that conditions all downstream behavior. Rapport (r) and stage of change ($\hat{c}$) are gates on strategy selection: planning-related strategies are suppressed until rapport exceeds a minimum threshold and the inferred stage of change reaches at least Contemplation, directly preventing early convergence.

System	Conv. Dynamics	Client Grounding Source	Long-form Gen.	Action Selection
KMI (Kim et al., 2025a)	×	*Posts*	×	√
CAMI + STAR (Yang et al., 2025a,b)	×	×	√	√
CounseLLMe (De Duro et al., 2025)	×	×	×	×
MentalChat16K (Xu et al., 2025)	×	×	×	×
SQPsych (Vu et al., 2025)	×	*Surveys*	√	×
CBTAlign (Kim et al., 2025b)	×	×	√	×
CALM-IT (ours)	√	*Posts + Surveys*	√	√

Table 1: Comparison of CALM-IT with prior systems generating synthetic mental health support dialogues. CALM-IT uniquely combines conversational dynamics tracking, cognitively grounded patient profiles, long-form multi-turn generation, and dual-actor action selection.

3.1 Client Turn Generation

CLIENTAGENT captures the reactive and sometimes inconsistent behavior observed in real therapy clients. Following each therapist turn, the agent evaluates the response via fast affective appraisal, updating internal states (readiness for change, emotional state, rapport, and short-term conversational goals) to reflect how clients adjust engagement in response to feeling understood or misunderstood. Update parameters are calibrated through empirical pilot simulations to avoid immediate rapport saturation and deterministic state trajectories, ensuring that client resistance, hesitation, and motivational fluctuation persist throughout extended sessions.

To generate its next turn, the client selects an interactional action (e.g., disclose, resist, clarify, or shift topics) (Yang et al., 2025a; Louie et al., 2024), with three candidate responses generated and ranked by log-likelihood under a language model (also gemma-2-9b-it) fine-tuned on patient turns from the same AnnoMI corpus. Full client-side generation details are provided in Appendix A, including Algorithm 1, while the prompts used for the LLM subcalls in client turn generation are shown in Figures A1–A7.

3.2 Therapist Turn Generation

THERAAGENT maintains an explicit internal model of the patient's inferred cognitive, motivational, and relational states (rapport, stage of change, and inferred client goals), updated after each patient turn to condition strategy selection on the evolving conversational state rather than isolated turn-level cues. Unlike prior MI simulation frameworks such as CAMI, which primarily assign cognitive states to the patient, THERAAGENT maintains therapist-side inferred states and uses their joint evolution to guide each response (Yang et al., 2025a).

To prevent conversational stagnation, THERAA-GENT monitors interactions for repetitive or unproductive patterns via a dedicated LLM call, initiating pivot strategies only when stagnation is detected rather than on a fixed schedule (Nguyen et al., 2025). For each turn, three candidate responses are generated conditioned on the selected strategy and current inferred state, then ranked by log-likelihood under a language model (in this case, gemma-2-9b-it) fine-tuned on MI-coded therapist turns from the AnnoMI corpus (Chang et al., 2020). Full algorithmic details are provided in Appendix A, including Algorithm 2, while the prompts used for the LLM subcalls in therapist turn generation are shown in Figures A8–A15. Finally, a full example of CALM-IT transcript generation can be found in table A1.

4 Experimental Setup

To evaluate the quality of CALM-IT-generated long-form Motivational Interviewing (MI) dialogues, we performed controlled generation and evaluation of therapist-client dialogues across four MI conversation modeling frameworks. The main-paper experiments use DeepSeek-V3.2 as the dialogue-generation backbone; replication results using OLMo-3.1-32B are reported in Appendix D.3. The evaluation pipeline consists of three stages: (1) **context generation** from paired client vignettes and psychological profiles, (2) **dialogue generation** using CALM-IT and baseline frameworks, and (3) **evaluation** of the resulting transcripts using LLM-as-a-judge assessments and algorithmic metrics at the turn-, agent-, and conversation-levels.

4.1 Context Generation

We collect natural mental health disclosures from Reddit to capture ecologically grounded expressions of psychological distress while preserving anonymity (Brocki et al., 2023; Sharma and De Choudhury, 2018). Using mental health-focused subreddits curated by Sharma and

De Choudhury (2018), we collected posts from 2015 to 2022 (5,794,243 posts in 55 subreddits). We apply BERTopic to cluster the corpus into 158 topics and select five representative posts per topic via cosine similarity, yielding 790 candidates. After keyword filtering for anxiety-, stress-, or depression-related content and de-identification, this yields a privacy-preserving set of 686 representative posts (Appendix B).

For each post, we generate a third-person vignette that preserves the client's core concerns while removing identifying details. From each vignette, we infer a cognitive–motivational profile (values, needs, session goal, and motivations for change) grounded in MI theory (Miller and Rollnick, 2013). We then simulate DASS-42 responses with GPT-5.1, where the DASS (Depression, Anxiety, and Stress Scales) provides a standardized psychometric characterization of symptom severity used to anchor the client's affective and motivational state. Each vignette is matched to a respondent from the OpenPsychometrics DASS dataset (39,775 anonymized records; Open Psychometrics, 2019), excluding respondents under 18, by maximizing weighted Cohen's κ and retaining only high-confidence matches ($\kappa > 0.6$, 99th percentile) with demographic consistency, selecting the highest-scoring match per vignette. For each matched pair, we use DeepSeek-V3.2 (Liu et al., 2025) to generate a detailed client background and a single-sentence session goal. The prompts used for demographic inference and profile generation are provided in Figures B1–B2. These materials serve as fixed context for dialogue generation across all frameworks.

4.2 Dialogue Generation

Using the simulated client contexts, we generate multi-turn therapeutic dialogues with four counseling frameworks. KMI and CAMI+STAR are selected as external baselines because they represent the strongest publicly available MI-specific dialogue frameworks, as established in Table 1. The four frameworks differ only in whether conversational dynamics are explicitly modeled:

1. CAMI + STAR (C + S) (Yang et al., 2025a,b),

2. KMI (Kim et al., 2025a),

3. CALM-IT without conversational-dynamics modeling (CI-NC),

4. CALM-IT with conversational-dynamics modeling (CI),

Each framework is applied to every simulated client at three target lengths (30, 50, and 100 turns), capturing short, medium, and long-form interactions. All sessions begin with a standardized opening turn and reserve the final ten turns for session closure (see details in Appendix A).

For the main experiments, we use DeepSeek-V3.2 as the backbone LLM for dialogue generation. This procedure yields $686 \times 3 \times 4 = 8{,}232$ generated transcripts for the main evaluation. To assess whether the observed trends generalize beyond a single backbone model, we additionally repeat the full generation and evaluation pipeline with OLMo-3.1-32B. These replication results are reported in Appendix D.3, with prompts, generation procedures, and evaluation protocols held identical to the main DeepSeek-V3.2 experiments. Details on model configurations, hyperparameters, compute infrastructure, and runtime costs are provided in Appendix E.

4.3 Evaluation

We evaluate generated dialogues using metrics for long-form LLM evaluation at three levels—turn-, agent-, and conversation-level (full definitions are provided in Table C1 in the Appendix). Conversation-level metrics are grounded in the Motivational Interviewing Treatment Integrity (MITI) code, MI process theory, and prior work on conversational redirection (Moyers et al., 2016; Miller and Rollnick, 2013; Apodaca and Longabaugh, 2009; Moyers et al., 2009; Nguyen et al., 2024). We additionally measure self-consistency (within agent) and entailment (between therapist and client) using a dialogue-specific NLI model, reporting each model's deviation from AnnoMI reference means to reflect that both under- and over-alignment reduce realism for mental health-related conversations (Hayes et al., 2007).

For redirection, we use ConvoKit's framework with an MI-conditioned likelihood model to estimate therapist-initiated redirection intensity (Chang et al., 2020). Metrics include % Accepted Redirection and Δ Change Talk @ Greatest Redirection (Nguyen et al., 2024). Change talk and sustain talk are labeled using a few-shot GPT-5.1 classifier with high accuracy on AnnoMI (Appendix C), and overall conversational directionality is measured using DialoFlow (Li et al., 2021).

For holistic constructs such as MITI 4.2 global ratings (Empathy, Partnership, Cultivating Change Talk, Softening Sustain Talk), we apply a rubric-based LLM-as-a-judge protocol on a 5-point Likert scale using DeepEval (Ip and Vongthongsri, 2025) with GPT-5.1, whose multi-pass aggregation improves score stability. Statistical differences between paired conditions are assessed using two-sided Wilcoxon signed-rank tests with Holm-Bonferroni correction. Because our goal is comparative process evaluation rather than estimating absolute clinical efficacy, we use automated metrics at scale and reserve clinician review for protocol validation. We therefore validate the automated protocol on 96 transcripts (1.2%; 8 per framework–length condition), independently reviewed by four licensed psychologists (annotation interface detailed in Figures C1–C3 of the Appendix). Expert and LLM-based judges show moderate to substantial agreement and consistent framework rankings across major metrics (Tables C12–C17), supporting automated evaluation for controlled comparative analysis. Our objective is not to estimate absolute clinical quality, but to compare relative differences in process fidelity across controlled conditions; consistent ranking across evaluators is sufficient for drawing meaningful conclusions. Full details on the metrics used are provided in Appendix Sections C.2–C.4 and Tables C1–C11.

5 Research Question 1 – Results

We report summary statistics for generated transcripts across all framework–length combinations in Table D1 in the Appendix. Examining per-framework averages across conversation lengths, we find that CALM-IT outperforms all baselines—including KMI, CAMI+STAR, and the ablated variant CALM-IT(NC)—across most turn-, agent-, and especially conversation-level evaluations. Performance gains are most pronounced at the conversational-level, where CALM-IT achieves the highest *Effectiveness* score (4.45), representing improvements of 4.22% over KMI (p < 0.001), 55.59% over CALM-IT(NC) (p < 0.001), and 273.95% over C+S (p < 0.001). Similar trends hold for other conversational metrics, including *Goal Alignment* (4.73), which is significantly higher than all baselines (p < 0.001). At the agent- and turn-levels, improvements are more modest but consistent: CALM-IT attains the highest scores on three of the four MITI 4.2 Global Ratings for therapist behavior

Metric	KMI	CI-NC	C+S	CI
Turn-level				
Client				
Client Readability (↓)	5.87	4.68	5.77	**4.45**
Therapist				
Reflection Quality	4.44	3.71	2.38	**4.61**
Question Quality	4.52	4.39	2.94	**4.97**
Therapist Readability (↓)	6.22	4.98	6.18	**4.93**
Agent-level				
Client				
Consistency	3.76	4.79	3.42	**4.92**
Therapist				
Cultivating Change Talk	**4.63**	3.15	1.88	4.34
Soften Sustain Talk	4.73	4.27	2.53	**4.88**
Empathy	4.66	4.68	3.53	**4.88**
Partnership	4.48	4.60	2.71	**4.88**
Reflection–Question Ratio	**2.05**	0.92	0.90	1.03
% Open Question	97.30	97.90	96.80	**98.47**
Conversation-level				
Effectiveness	4.27	2.86	1.19	**4.45**
Realignment	4.49	4.53	3.21	**4.87**
Goal Alignment	4.60	3.89	2.13	**4.73**
Δ_{Ref} Directionality (↓)	0.081	0.03	0.048	**0.005**
Δ_{Ref} Avg. Self-Consist. (↓)	0.291	0.141	**0.072**	0.157
Δ_{Ref} Avg. Entail. (↓)	0.092	0.141	0.073	**0.042**

Table 2: Turn-, agent- and conversation-level evaluation of MI conversations generated by four frameworks with DeepSeek: KMI, CALM-IT without conversational dynamics (CI-NC), CAMI+STAR (C+S), and CALM-IT (CI). Scores are averaged across all conversations and lengths. Downward arrow (↓) indicates that lower values correspond to better performance; otherwise, higher values are better. Best-performing framework for each metric is bolded (RQ1)

(*Softening Sustain Talk*, *Empathy*, and *Partnership*, all 4.88), while its CLIENTAGENT produces more readable turns (readability 4.45 vs. 5.87 for KMI and 5.77 for CAMI+STAR, p < 0.001) and exhibits greater behavioral consistency with assigned client backgrounds (consistency 4.92 vs. 3.76 for KMI and 3.42 for CAMI+STAR, p < 0.001) than competing frameworks. These improvements remain consistent even when transcripts are generated with a lighter-weight backbone, OLMo-3.1 (Table D3). Component ablation experiments further show that removing key conversational dynamics mechanisms leads to systematic performance declines (Table D4).

In addition to higher performance across most metrics, CALM-IT exhibits substantially greater stability in performance while generating long-form dialogues. Figure 2 summarizes relative changes in evaluation metrics as conversations scale from short (30 turns) to long (100 turns). We found that CALM-IT shows the smallest absolute overall drift, with a mean absolute percentage change of

Figure 2: Percentage change in performance across metrics from short-form dialogues (30 turns) to long-form dialogues (100 turns), averaged across all pairs of short-long dialogues (generated using DeepSeek-V3.2). Positive or close-to-zero values indicate stable performance; negative values indicate degradation in performance. (RQ1)

only 0.02% across all metrics, compared to 2.41% for `KMI`, 0.72% for `CALM-IT (NC)`, and 3.56% for `CAMI+STAR`. This effect is even more pronounced for conversational-level metrics, where `CALM-IT` only showed a -1.19% degradation on average compared to -7.05% for `KMI` or -8.48% for `CI-NC`.

6 Research Question 2 – Results

To assess therapist redirection effectiveness, we examine client responses following redirection events using several conversation-level metrics that capture both immediate acceptance and downstream conversational shifts. Although `CALM-IT` redirects less frequently overall, as reflected by its lowest redirection_ratio (0.594), it produces more effective therapist redirection than all baseline frameworks. This is shown first by the *percentage accepted redirection* measurements, which capture the client's immediate response following a redirection event. CALM-IT achieves the highest acceptance rate at 64.28%, significantly outperforming KMI at 59.11% (p < 0.001), CAMI+STAR at 54.95% (p < 0.001), and CALM-IT(NC) at 52.14% (p < 0.001). Beyond immediate acceptance, CALM-IT's strongest redirection events yield sustained downstream benefits, with increased client change talk of 12.40% and substantial reductions in sustain talk of -8.15% on average (compared to 9.50% increase in change talk and -1.67% reduc-

tion in sustain talk for KMI).

To quantify the asymmetry in Figure 3, we test if the distribution of Δ sustain talk exhibits statistically significant negative skewness. `CALM-IT` shows a significant left-skew, with skewness $\gamma = -0.66$ and a D'Agostino skewness test statistic of $Z = -6.56$ ($p < 0.0001$), showing that strong redirection events are systematically followed by reductions in client resistance. Conversely, the KMI distribution is more weakly left-skewed with $\gamma = -0.24$ ($p < 0.0001$), while `CAMI+STAR` and `CALM-IT(NC)` exhibit significant right-skew with $\gamma = 0.19$ (p < 0.0001) and $\gamma = 0.66$ (p < 0.0001), reflecting weak or counterproductive redirection that often results in increases of sustain talk. This shows CALM-IT's selective redirection strategy uniquely shifts post-redirection outcomes toward reduced sustain talk and increased change talk.

7 Discussion & Conclusion

We introduced CALM-IT, a framework that models long-form Motivational Interviewing (MI) as an explicit dynamical process. By continuously updating representations of rapport, mental state, and session goals outside the surface text, CALM-IT conditions turn-by-turn strategy selection to prevent premature conversational convergence. Evaluated across 8,232 synthetic dialogues spanning three length conditions and four different frame-

Figure 3: Distribution of changes in client sustain talk at the greatest therapist-initiated redirection moment, for all four tested frameworks (with DeepSeek-V3.2 as backbone LLM). Negative values indicate reductions in sustain talk following redirection, while positive values indicate increases. Solid vertical lines denote the mean change for each framework, and dashed lines represent no change (RQ2)

works (including CALM-IT), CALM-IT consistently outperforms baselines across key process-oriented metrics. Notably, it improves three of four MITI 4.2 global ratings, Goal Alignment, and Effectiveness, while yielding more MI-consistent trajectory shifts. Furthermore, CALM-IT initiates fewer direct redirections yet achieves a higher client acceptance rate (64.3% vs. 59.1% for KMI) and induces larger, more consistent post-redirection reductions in client sustain talk than any baseline.

These results demonstrate that long-form conversational stability benefits from explicit conversational dynamics modeling. Standard large language models optimize for local next-turn plausibility and lack proper macro-level representations of therapeutic progress, often leading to compounding degradation over extended interactions (Laban et al., 2025; Li et al., 2025). CALM-IT addresses this limitation by continuously updating representations of rapport, mental state, and session goals, enabling strategy recalibration before relational drift becomes entrenched. Consequently, the framework exhibits significantly lower performance degradation than other architectures when scaling to extended sessions, maintaining high process fidelity where long-horizon interaction is most challenging.

We pursue this synthetic modeling approach to overcome the limitations of real, donated conversational data. Although clinical transcripts provide authentic language patterns, they remain severely restricted in volume, accessibility, and diversity due to privacy regulations and systemic stigma. Moreover, donated transcripts are passive artifacts: they cannot be dynamically manipulated to introduce controlled conversational perturbations, vary client resistance, or stress-test specific model vulnerabilities. High-fidelity synthetic simulation addresses these constraints by providing a scalable, interactive sandbox for systematically generating, observing, and evaluating long-form therapeutic interactions. Our findings motivate future work to develop richer methods for measuring long-term therapeutic dialogue quality and to design more expressive models of conversational dynamics that capture how actors' relational, motivational, and emotional states evolve and shape high-quality therapeutic dialogue across extended conversations.

Limitations

CALM-IT does not resolve all open problems in therapeutic dialogue generation. Rather, it establishes a reproducible framework and evaluation protocol within which future work can build. Moreover, we acknowledge some limitations. Real-world therapeutic interactions are shaped by complex, nonlinear dynamics arising from interpersonal history, situational context, and unobserved client factors, which CALM-IT only partially approximates through large language model–based state inference and prompting. These approximations may diverge from how therapeutic processes unfold in real clinical settings, motivating future work on more specialized or hybrid models that can more faithfully capture such dynamics in real time.

We also note that transcript generation is performed independently across two backbone LLMs (DeepSeek-V3.2 and OLMo-3.1-32B), providing evidence that gains from conversational state tracking are model-agnostic. However, systematically evaluating how backbone model choice affects long-form dialogue quality, conversational dynamics, and redirection behavior across a broader range of architectures and scales remains an open question for future work. The client agent's state transition rules are additionally governed by heuristic parameters calibrated through pilot simulations, including the System-1 evaluation activation probability (p=0.5). Fitting these parameters to real therapy corpora remains a natural direction for future work.

Another key limitation is the absence of real therapy transcripts as a validation reference. CALM-IT does not train on or evaluate against real clinical interactions, and our evaluation cannot directly confirm that generated dialogues match clinically meaningful dynamics in practice. To partially mitigate this, we include expert human evaluation by four licensed psychologists across 96 transcripts. Framework rankings produced by human annotators align consistently with those from automated LLM-as-a-judge metrics across all major metrics (Tables C13–C16), suggesting that automated evaluation captures the same comparative quality distinctions that expert raters identify. While this convergence does not substitute for grounding in real therapy data, it provides supporting evidence that CALM-IT's gains reflect clinically meaningful differences rather than artifacts of model-specific scoring. Grounding state transition parameters and evaluation criteria in real therapy corpora remains an important direction for future work.

Next, to reduce generator-evaluator bias, generation and evaluation use different model families (DeepSeek-V3.2 and GPT-5.1 respectively). Nevertheless, these judgments remain sensitive to model choice and prompting and should be interpreted as approximations of expert assessment rather than objective ground truth. In addition, our framework is explicitly grounded in MI theory, which enables fine-grained modeling and evaluation of motivational and relational processes but limits immediate generalization to other therapeutic modalities. Future work could adapt CALM-IT to alternative approaches such as Cognitive Behavioral Therapy (CBT) by incorporating modality-specific state representations, strategies, and evaluation criteria.

Finally, CALM-IT relies on Reddit mental health disclosures as proxy data for real-world therapy contexts, which disproportionately reflect WEIRD (Western, Educated, Industrialized, Rich, and Democratic) populations and are shaped by social-media-specific norms of self-disclosure, including intimacy costs that constrain the range of expressed distress (Ma et al., 2016). Consequently, clinically salient scenarios such as acute crisis, severe psychiatric conditions, or culturally specific concerns may be underrepresented, limiting the diversity of generated synthetic transcripts.

Ethical Considerations

CALM-IT is a research framework for studying long-form conversational dynamics in motivational interviewing, not a system intended for clinical use. The realism of the simulated conversations should not be taken as an endorsement of automated mental health support. We do not support the deployment of systems for mental health diagnosis, treatment, or crisis response without appropriate regulatory review. Any real-world use would require FDA clearance, with FDA approval serving as a longer-term objective for higher-risk or autonomous clinical applications. The contribution of this work lies in developing evaluation pipelines, stress-testing methods, and synthetic data resources that can support such regulatory validation efforts.

The framework relies on synthetic data to enable large-scale construction and evaluation of systems for mental health support, avoiding the privacy and legal constraints associated with real therapy transcripts. This choice also introduces limitations.

Although grounded in mental health disclosures from Reddit, the resulting simulations may reflect existing biases in mental health representation, including demographic, cultural, and normative assumptions present in the source data and modeling choices. These biases can affect simulated client behavior, therapist evaluations, and assessments of therapeutic progress, potentially obscuring uneven performance across populations.

Findings obtained using `CALM-IT` should therefore be interpreted as characterizations of conversational behavior under controlled experimental conditions, not as evidence of clinical safety or effectiveness. Use of realistic mental health simulations benefits from clear boundaries, explicit documentation of data sources and assumptions, and expert oversight. In this role, synthetic frameworks such as `CALM-IT` support research, training, and regulatory preparation, while remaining distinct from the delivery of professional mental health care.

References

Paul C Amrhein, William R Miller, Carolina E Yahne, Michael Palmer, and Laura Fulcher. 2003. Client commitment language during motivational interviewing predicts drug use outcomes. *Journal of consulting and clinical psychology*, 71(5):862.

Timothy R Apodaca and Richard Longabaugh. 2009. Mechanisms of change in motivational interviewing: A review and preliminary evaluation of the evidence. *Addiction*, 104(5):705–715.

Simon Baron-Cohen. 1995. Mindblindness: An essay on autism and theory of mind. *MIT Press*.

Lennart Brocki, George C Dyer, Anna Gładka, and Neo Christopher Chung. 2023. Deep learning mental health dialogue system. In *2023 IEEE International Conference on Big Data and Smart Computing (Big-Comp)*, pages 395–398. IEEE.

Mohit Chandra, Siddharth Sriraman, Harneet Singh Khanuja, Yiqiao Jin, and Munmun De Choudhury. 2025. Reasoning is not all you need: Examining llms for multi-turn mental health conversations. *arXiv preprint arXiv:2505.20201*.

Jonathan P Chang, Caleb Chiam, Liye Fu, Andrew Wang, Justine Zhang, and Cristian Danescu-Niculescu-Mizil. 2020. Convokit: A toolkit for the analysis of conversations. In *Proceedings of the 21th annual meeting of the special interest group on discourse and dialogue*, pages 57–60.

Herbert H Clark. 1996. *Using language*. Cambridge university press.

Edgar Dale and Jeanne S Chall. 1948. A formula for predicting readability: Instructions. *Educational research bulletin*, pages 37–54.

Edoardo Sebastiano De Duro, Riccardo Improta, and Massimo Stella. 2025. Introducing counsellme: A dataset of simulated mental health dialogues for comparing llms like haiku, llamantino and chatgpt against humans. *Emerging Trends in Drugs, Addictions, and Health*, 5:100170.

Ning Ding, Yulin Chen, Bokai Xu, Yujia Qin, Shengding Hu, Zhiyuan Liu, Maosong Sun, and Bowen Zhou. 2023. Enhancing chat language models by scaling high-quality instructional conversations. In *Proceedings of the 2023 Conference on Empirical Methods in Natural Language Processing*, pages 3029–3051.

Nouha Dziri, Ehsan Kamalloo, Kory Mathewson, and Osmar R Zaiane. 2019. Evaluating coherence in dialogue systems using entailment. In *Proceedings of the 2019 conference of the north american chapter of the association for computational linguistics: Human language technologies, volume 1 (long and short papers)*, pages 3806–3812.

Arash Eshghi, Julian Hough, Matthew Purver, Ruth Kempson, and Eleni Gregoromichelaki. 2012. Conversational interactions: Capturing dialogue dynamics. *From Quantification to Conversation: Festschrift for Robin Cooper on the occasion of his 65th birthday*, 19:325–349.

Rachel Hall. 2025. Ai chatbots giving dangerous advice to mentally ill, psychologists warn. https://www.theguardian.com/technology/2025/nov/30/chatgpt-dangerous-advice-mentally-ill-psychologists-openai. The Guardian. Accessed: 2025-12-01.

Adele M Hayes, Jean-Philippe Laurenceau, Greg Feldman, Jennifer L Strauss, and LeeAnn Cardaciotto. 2007. Change is not always linear: The study of nonlinear and discontinuous patterns of change in psychotherapy. *Clinical psychology review*, 27(6):715–723.

Xuehai He, Shu Chen, Zeqian Ju, Xiangyu Dong, Hongchao Fang, Sicheng Wang, Yue Yang, Jiaqi Zeng, Ruisi Zhang, Ruoyu Zhang, and 1 others. 2020. Meddialog: Two large-scale medical dialogue datasets. *arXiv preprint arXiv:2004.03329*.

Jeffrey Ip and Kritin Vongthongsri. 2025. deepeval.

Matthew Jörke, Shardul Sapkota, Lyndsea Warkenthien, Niklas Vainio, Paul Schmiedmayer, Emma Brunskill, and James A Landay. 2025. Gptcoach: Towards llm-based physical activity coaching. In *Proceedings of the 2025 CHI Conference on Human Factors in Computing Systems*, pages 1–46.

Daniel Kahneman. 2011. *Thinking, fast and slow*. macmillan.

Hyunjong Kim, Suyeon Lee, Yeongjae Cho, Eunseo Ryu, Yohan Jo, Suran Seong, and Sungzoon Cho. 2025a. Kmi: A dataset of korean motivational interviewing dialogues for psychotherapy. *arXiv preprint arXiv:2502.05651*.

Yejin Kim, Chi-Hyun Choi, Selin Cho, Jy-yong Sohn, and Byung-Hoon Kim. 2025b. Aligning large language models for cognitive behavioral therapy: a proof-of-concept study. *Frontiers in Psychiatry*, 16:1583739.

Philippe Laban, Hiroaki Hayashi, Yingbo Zhou, and Jennifer Neville. 2025. Llms get lost in multi-turn conversation. *arXiv preprint arXiv:2505.06120*.

Suyeon Lee, Sunghwan Mac Kim, Minju Kim, Dongjin Kang, Dongil Yang, Harim Kim, Minseok Kang, Dayi Jung, Min Hee Kim, Seungbeen Lee, and 1 others. 2024. Cactus: Towards psychological counseling conversations using cognitive behavioral theory. In *Findings of the Association for Computational Linguistics: EMNLP 2024*, pages 14245–14274.

Yubo Li, Xiaobin Shen, Xinyu Yao, Xueying Ding, Yidi Miao, Ramayya Krishnan, and Rema Padman. 2025. Beyond single-turn: A survey on multi-turn interactions with large language models. *arXiv preprint arXiv:2504.04717*.

Zekang Li, Jinchao Zhang, Zhengcong Fei, Yang Feng, and Jie Zhou. 2021. Conversations are not flat: Modeling the dynamic information flow across dialogue utterances. *arXiv preprint arXiv:2106.02227*.

Aixin Liu, Aoxue Mei, Bangcai Lin, Bing Xue, Bingxuan Wang, Bingzheng Xu, Bochao Wu, Bowei Zhang, Chaofan Lin, Chen Dong, and 1 others. 2025. Deepseek-v3. 2: Pushing the frontier of open large language models. *arXiv preprint arXiv:2512.02556*.

Ryan Louie, Ananjan Nandi, William Fang, Cheng Chang, Emma Brunskill, and Diyi Yang. 2024. Roleplay-doh: Enabling domain-experts to create llm-simulated patients via eliciting and adhering to principles. *arXiv preprint arXiv:2407.00870*.

Daniel Cabrera Lozoya, Eloy Hernández Lúa, Juan Alberto Barajas Perches, Mike Conway, and Simon D'Alfonso. 2025. Synthetic empathy: Generating and evaluating artificial psychotherapy dialogues to detect empathy in counseling sessions. In *Proceedings of the 10th Workshop on Computational Linguistics and Clinical Psychology (CLPsych 2025)*, pages 157–171.

Xiao Ma, Jeff Hancock, and Mor Naaman. 2016. Anonymity, intimacy and self-disclosure in social media. In *Proceedings of the 2016 CHI conference on human factors in computing systems*, pages 3857–3869.

Zafarullah Mahmood, Soliman Ali, Jiading Zhu, Mohamed Abdelwahab, Michelle Yu Collins, Sihan Chen, Yi Cheng Zhao, Jodi Wolff, Osnat C Melamed, Nadia Minian, and 1 others. 2025. A fully generative motivational interviewing counsellor chatbot for moving smokers towards the decision to quit. In *Findings of the Association for Computational Linguistics: ACL 2025*, pages 25008–25043.

Tobias Mayer, Neha Warikoo, Oliver Grimm, Andreas Reif, and Iryna Gurevych. 2022. Gdpr compliant collection of therapist-patient-dialogues. *arXiv preprint arXiv:2211.12360*.

Ryan K McBain, Robert Bozick, Melissa Diliberti, Li Ang Zhang, Fang Zhang, Alyssa Burnett, Aaron Kofner, Benjamin Rader, Joshua Breslau, Bradley D Stein, and 1 others. 2025. Use of generative ai for mental health advice among us adolescents and young adults. *JAMA Network Open*, 8(11):e2542281–e2542281.

William R. Miller and Stephen Rollnick. 2013. *Motivational Interviewing: Helping People Change*, 3rd edition. Guilford Press.

Theresa B Moyers, Tim Martin, Jon M Houck, Paulette J Christopher, and J Scott Tonigan. 2009. From in-session behaviors to drinking outcomes: a causal chain for motivational interviewing. *Journal of consulting and clinical psychology*, 77(6):1113.

Theresa B Moyers, Lauren N Rowell, Jennifer K Manuel, Denise Ernst, and Jon M Houck. 2016. The motivational interviewing treatment integrity code (miti 4): rationale, preliminary reliability and validity. *Journal of substance abuse treatment*, 65:36–42.

Vivian Nguyen, Sang Min Jung, Lillian Lee, Thomas D Hull, and Cristian Danescu-Niculescu-Mizil. 2024. Taking a turn for the better: Conversation redirection throughout the course of mental-health therapy. *arXiv preprint arXiv:2410.07147*.

Vivian Nguyen, Lillian Lee, and Cristian Danescu-Niculescu-Mizil. 2025. Hanging in the balance: Pivotal moments in crisis counseling conversations. *arXiv preprint arXiv:2506.03941*.

Open Psychometrics. 2019. Depression anxiety stress scales (dass) dataset. https://openpsychometrics.org/_rawdata/DASS_data_21.02.19.zip.

PBS NewsHour. 2025. What to know about "ai psychosis" and the effect of ai chatbots on mental health. https://www.pbs.org/newshour/show/what-to-know-about-ai-psychosis-and-the-effect-of-ai-chatbots-on-mental-health. Accessed: 2025-08-15.

Wojciech Pichowicz, Michal Kotas, and Pawel Piotrowski. 2025. Performance of mental health chatbot agents in detecting and managing suicidal ideation. *Scientific Reports*, 15(1):31652.

Huachuan Qiu, Hongliang He, Shuai Zhang, Anqi Li, and Zhenzhong Lan. 2024. Smile: Single-turn to multi-turn inclusive language expansion via chatgpt

for mental health support. In *Findings of the Association for Computational Linguistics: EMNLP 2024*, pages 615–636.

Eva Sharma and Munmun De Choudhury. 2018. Mental health support and its relationship to linguistic accommodation in online communities. In *Proceedings of the 2018 CHI conference on human factors in computing systems*, pages 1–13.

Karan Singhal, Shekoofeh Azizi, Tao Tu, S Sara Mahdavi, Jason Wei, Hyung Won Chung, Nathan Scales, Ajay Tanwani, Heather Cole-Lewis, Stephen Pfohl, and 1 others. 2023. Large language models encode clinical knowledge. *Nature*, 620(7972):172–180.

Helen Spencer-Oatey. 2004. *Culturally speaking: Managing rapport through talk across cultures*. A&C Black.

Ivan Sviridov, Amina Miftakhova, Tereshchenko Artemiy Vladimirovich, Galina Zubkova, Pavel Blinov, and Andrey Savchenko. 2025. 3mdbench: Medical multimodal multi-agent dialogue benchmark. In *Proceedings of the 2025 Conference on Empirical Methods in Natural Language Processing*, pages 26625–26665.

Brandon Theodorou, Cao Xiao, and Jimeng Sun. 2023. Synthesize high-dimensional longitudinal electronic health records via hierarchical autoregressive language model. *Nature communications*, 14(1):5305.

Doan Nam Long Vu, Rui Tan, Lena Moench, Svenja Jule Francke, Daniel Woiwod, Florian Thomas-Odenthal, Sanna Stroth, Tilo Kircher, Christiane Hermann, Udo Dannlowski, and 1 others. 2025. Roleplaying with structure: Synthetic therapist-client conversation generation from questionnaires. *arXiv preprint arXiv:2510.25384*.

Junda Wang, Zonghai Yao, Zhichao Yang, Huixue Zhou, Rumeng Li, Xun Wang, Yucheng Xu, and Hong Yu. 2024a. Notechat: a dataset of synthetic patient-physician conversations conditioned on clinical notes. In *Findings of the Association for Computational Linguistics: ACL 2024*, pages 15183–15201.

Ruiyi Wang, Stephanie Milani, Jamie C Chiu, Jiayin Zhi, Shaun M Eack, Travis Labrum, Samuel M Murphy, Nev Jones, Kate Hardy, Hong Shen, and 1 others. 2024b. Patient-ψ: Using large language models to simulate patients for training mental health professionals. *arXiv preprint arXiv:2405.19660*.

Sheng Wang, Zihao Zhao, Xi Ouyang, Tianming Liu, Qian Wang, and Dinggang Shen. 2024c. Interactive computer-aided diagnosis on medical image using large language models. *Communications Engineering*, 3(1):133.

Yizhong Wang, Yeganeh Kordi, Swaroop Mishra, Alisa Liu, Noah A Smith, Daniel Khashabi, and Hannaneh Hajishirzi. 2023. Self-instruct: Aligning language models with self-generated instructions. In *Proceedings of the 61st annual meeting of the association for computational linguistics (volume 1: long papers)*, pages 13484–13508.

Jia Xu, Tianyi Wei, Bojian Hou, Patryk Orzechowski, Shu Yang, Ruochen Jin, Rachael Paulbeck, Joost Wagenaar, George Demiris, and Li Shen. 2025. Mentalchat16k: A benchmark dataset for conversational mental health assistance. In *Proceedings of the 31st ACM SIGKDD Conference on Knowledge Discovery and Data Mining V. 2*, pages 5367–5378.

Yizhe Yang, Palakorn Achananuparp, He-Yan Huang, Jing Jiang, Phey Ling Kit, Nicholas Gabriel Lim, Cameron Tan Shi Ern, and Ee-Peng Lim. 2025a. Cami: A counselor agent supporting motivational interviewing through state inference and topic exploration. In *Proceedings of the 63rd Annual Meeting of the Association for Computational Linguistics (Volume 1: Long Papers)*, pages 21037–21081.

Yizhe Yang, Palakorn Achananuparp, He-Yan Huang, Jing Jiang, Nicholas Gabriel Lim, Cameron Tan Shi Ern, Phey Ling Kit, Jenny Giam Xiuhui, John Pinto, and Ee-peng Lim. 2025b. Consistent client simulation for motivational interviewing-based counseling. In *Proceedings of the 63rd Annual Meeting of the Association for Computational Linguistics (Volume 1: Long Papers)*, pages 20959–20998.

Jinsung Yoon, Michel Mizrahi, Nahid Farhady Ghalaty, Thomas Jarvinen, Ashwin S Ravi, Peter Brune, Fanyu Kong, Dave Anderson, George Lee, Arie Meir, and 1 others. 2023. Ehr-safe: generating high-fidelity and privacy-preserving synthetic electronic health records. *NPJ digital medicine*, 6(1):141.

A Appendix A: Additional details regarding CALM-IT framework

We provide full algorithmic and prompt-level details of the CALM-IT framework, including therapist and client generation procedures (Algorithm 2, 1), prompts for conversational state updates, and latent-state inferences used to guide long-form dialogue generation below.

#	Spkr	Dialogue
1	C	Yeah, okay. I guess... it's these anger episodes. They just come out of nowhere and I don't know why.
2	T	Thank you for sharing that. It sounds really confusing and unsettling when the anger comes on so suddenly. Can you tell me about a recent time that happened?
3	C	Well... just a few days ago, I was in the office trying to finish a map. My wife called to ask about dinner, and I just... snapped at her for no reason.
4	T	Can you tell me what was happening for you right before you snapped?
5	C	I was just so focused on the map, you know? It had to be perfect. My head was just full of all those lines.
6	T	So, that intense focus on making everything perfect was filling your head. Then, something shifted when your wife called.
7	C	Yeah. It was like... all that quiet in my head just broke. The phone was so loud.
8	T	Can you tell me what felt important about having a quiet mind in that moment?
9	C	Hmm. It felt... I could just think about the work. Nothing else. It was peaceful.
10	T	That feeling of peace and focus is really valuable. It shows just how important it is for you to have that kind of space for your mind.
11	C	Yeah. It is. But then it's gone so fast. Like a glass breaking.
12	T	It sounds like that quiet focus is really important and peaceful for you. At the same time, it's troubling how quickly it can shatter and lead to you snapping at your wife.
13	C	It just... it feels like such a waste. That peace. I work so hard to get there.
14	T	That feeling of waste is really strong. What is it about that peaceful state that makes it worth holding onto?
15	C	It's the only time my head isn't... busy. All the worries, all the noise just stops. It's like being out in the woods when everything is still.
16	T	That moment of clarity and stillness is really important to you, and it makes sense that you'd want to protect it.
17	C	Yeah... and when it's gone, it's like all the noise comes rushing back in. And that's when I get... short with people.
18	T	It sounds like that quiet mind is a real safe space for you. When it gets interrupted by all that noise, it makes sense that you react in ways you're not happy with.
19	C	Yeah, it is safe. But then... it's just gone. And I feel worse than before.
20	T	It's so important to have a safe space, and it's completely draining when that safety suddenly disappears.

#	Spkr	Dialogue
21	C	Um, and then it's just... empty. It's like the quiet after a storm, but worse.
22	T	It sounds like that emptiness feels even deeper than the noise you were trying to get away from, almost like a deeper kind of ache.
23	C	Yeah. A deeper ache. It's like... after the quiet breaks, there's nothing left to hold on to.
24	T	It makes sense that when that peace is broken, it leaves you feeling even more empty. And in that emptiness, you might find yourself being short with people, which is the last thing you wanted.
25	C	It's the last thing I want. I hate it. Then I just go out to the woods by myself for hours.
26	T	It's so human to want to pull back and protect yourself after a loss like that. I hear that for many people, when frustration comes back after a moment of calm, it can feel especially harsh.
27	C	Yeah, harsher. I know what I just lost. That little piece of quiet.
28	T	It makes sense that the frustration feels so sharp when you're so aware of what you've lost.
29	C	And that's when I snap at my wife. Over nothing. When I come back from the woods... I'm still just empty.
30	T	We have about five minutes left. Is it okay if I share what I'm hearing and see what you think?
31	C	Yes, please. Go ahead.
32	T	Of course. It sounds like that peaceful, focused state is really important to you—it's your safe space. And it's toughest when it gets interrupted. That emptiness and frustration feel even sharper because you know what you've lost, and it leads to snapping at your wife, which you don't want. Since you value both that peace and your relationship, what's one small thing you could try this week to protect that space, even for just a few more minutes?
33	C	Hmm. Maybe... if I take a rock from my collection, one I found on a good day, to keep in my pocket when I come home.
34	T	That's such a personal and specific idea. Can you tell me more about how you picture that rock helping you when you get home?
35	C	It's like a little piece of a good day, right there. If I feel that roughness coming on, I can just put my hand in my pocket and feel it. Its smoothness, its weight. Maybe it can just... remind me.
36	T	That sounds like a really important reminder for you. What's the first small sign you'll notice that the plan is working when you try it at home?
37	C	Well... if I reach for the rock first, before I say anything sharp. Just that small movement in my pocket. That would be the sign.
38	T	So, reaching for the rock is your first signal. How could that small step change what happens next?
39	C	Maybe it gives me a moment to breathe. To remember I'm home now, with her. Not still out there alone.
40	T	You've done some really meaningful work today just by exploring what's important to you. No matter what pace feels right for you, you don't have to do this alone. We can keep building on this together.

Table A1: Example 30-turn motivational interviewing transcript generated by CALM-IT (+ 10 turns for wrapping up session)

Algorithm 1 Client Turn Generation

Inputs. M (history), Σ (summary), b (background), c (change stage), g (goal), e (emotion), ρ (rapport), π (client action), t (turn counter), $T_{\max}$ (limit).

1: /* *Update running summary* */
2: $\Sigma \leftarrow \text{UpdateSummary}(\Sigma, M)$
3: /* *Client may utilize system-1 thinking to evaluate therapist turn (50% chance)* */
4: $z \leftarrow \text{Bernoulli}(0.5)$
5: **if** $z = 1$ **then**
6:    $q \leftarrow \text{RateTherapist}(M, \Sigma, b, c, g, e, \rho)$
7: **else**
8:    $q \leftarrow \text{"}Neutral\text{"}$ /* *default evaluation is "Neutral"* */
9: **end if**
10: /* *Update change/readiness* */
11: $c \leftarrow \text{UpdateChangeP}(\Sigma, M, b, c, q)$
12: /* *Update emotional state* */
13: $e \leftarrow \text{UpdateEmotion}(\Sigma, M, b, e, q)$
14: /* *Update rapport* */
15: $\rho \leftarrow \text{UpdateRapportP}(\Sigma, M, b, e, c, \rho, q)$
16: /* *Update short-term goal* */
17: $g \leftarrow \text{UpdateGoalP}(\Sigma, M, b, e, c, \rho, g, q)$
18: /* *Select client action* */
19: $\pi \leftarrow \text{SelectActionP}(\Sigma, M, b, e, c, g, \rho, q)$
20: /* *Sample candidate client turns* */
21: $U \leftarrow \emptyset$
22: **for** $i = 1$ **to** 3 **do**
23:    $x \leftarrow \text{GenClient}(\Sigma, M, b, e, c, g, \rho, \pi)$
24:    $\text{append}(U, x)$
25: **end for**
26: /* *Choose highest-likelihood turn* */
27: $u_P \leftarrow \text{SelectMaxLikelihood}(U)$
28: /* *Append to history* */
29: $\text{append}(M, u_P)$
30: **if** $(\pi = \text{"}Terminate\text{"})$ or $(t \geq T_{\max})$ **then**
31:    /* *End dialogue* */
32:    **return** (M, End)
33: **end if**
34: /* *Continue dialogue* */
35: **return** $(M, \text{Continue})$

Algorithm 2 Therapist Turn Generation

Inputs. M (history), Σ (summary), $\hat{b}$ (inferred background), $\hat{e}$ (inferred emotion), $\hat{c}$ (inferred stage of change), $\hat{g}$ (inferred patient goal), $\hat{r}$ (inferred rapport), a (strategy), t (turn counter).

1: /* *Update summary of dialogue* */
2: $\Sigma \leftarrow \text{UpdateSummary}(\Sigma, M)$
3: /* *Infer patient background* */
4: $\hat{b} \leftarrow \text{InferBackground}(\Sigma, M, \hat{b})$
5: /* *Infer patient's emotional state* */
6: $\hat{e} \leftarrow \text{InferEmotion}(\Sigma, M, \hat{e})$
7: /* *Infer patient's stage of change* */
8: $\hat{c} \leftarrow \text{InferStageOfChange}(\Sigma, M, \hat{c})$
9: /* *Infer patient's short-term dialogue goal* */
10: $\hat{g} \leftarrow \text{InferGoal}(\Sigma, M, \hat{g})$
11: /* *Infer rapport between therapist and patient* */
12: $\hat{r} \leftarrow \text{InferRapport}(\Sigma, M, \hat{e}, \hat{r})$
13: /* *Detect stagnation in dialogue* */
14: $\text{rep} \leftarrow \text{CheckRepetition}(\Sigma, M)$
15: **if** $\text{rep} = 0$ **then**
16:    /* *Select MI-consistent strategy* */
17:    $a \leftarrow \text{SelectStrategy}(\Sigma, M, \hat{b}, \hat{e}, \hat{c}, \hat{g}, \hat{r})$
18: **else**
19:    /* *Select pivot strategy to restore progress* */
20:    $a \leftarrow \text{SelectPivot}(\Sigma, M, \hat{b}, \hat{e}, \hat{c}, \hat{g}, \hat{r})$
21: **end if**
22: /* *Sample candidate therapist turns* */
23: $U \leftarrow \emptyset$
24: **for** $i = 1$ **to** 3 **do**
25:    $x \leftarrow \text{GenTherapist}(\Sigma, M, \hat{b}, \hat{e}, \hat{c}, \hat{g}, \hat{r}, a)$
26:    $\text{append}(U, x)$
27: **end for**
28: /* *Choose highest-likelihood turn* */
29: $u_T \leftarrow \text{SelectMaxLikelihood}(U)$
30: /* *Append therapist turn to dialogue history* */
31: $\text{append}(M, u_T)$
32: **return** M

You are the patient's fast, intuitive System-1 voice. Instantly appraise the LAST THERAPIST MESSAGE
from the patient's point of view and output a single-line rating with a brief reason.

Inputs (JSON-like objects; empty allowed):
dialogue_summary - dialogue recap: {}
last_patient_message - Your last turn: {}
last_therapist_message - Therapist's last turn: {}
patient_background - Background covering demographics, relationships, routines,
preferences, culture, and goals: {}
patient_cognitive_model - Beliefs, coping strategies, biases: {}
patient_goal - Current intermediate goal in patient's words: {}
patient_session_goal - Overarching session goal: {}
patient_change_state - MI stage (e.g., precontemplation, contemplation, preparation, action, etc.): {}
patient_rapport_with_therapist - Openness/trust (-1 absolute distrust, 1 absolute trust): {}
patient_emotion - Primary/secondary emotions with intensity: {}
turn_counter - Current turn counter: {}

Task:
- Remember the therapist's role is to support exploration, not give direct answers.
- Judge how the PATIENT would immediately and viscerally react to the therapist's last turn,
  given all inputs (especially last_patient_message).
- Favor "NEUTRAL" over "BAD" or "VERY BAD" when uncertain.
- Focus on System-1 intuition, not deliberative reasoning.
- Weigh especially heavily: patient background, cognitive model, stage of change,
  rapport level, and emotional intensity.
- Consider MI principles: empathy, autonomy support, evocation, and goal alignment.
- Raise the rating if the message resonates with the patient's worldview or coping style.
- Lower the rating if it clashes with identity, beliefs, or communication preferences.

Rating Scale (choose exactly one):
VERY GOOD - strongly affirms my values and goals; evokes change talk; deepens trust.
GOOD - feels supportive and aligned; mostly MI-consistent; invites reflection.
NEUTRAL - mixed or ambivalent impact; partial resonance with some disconnect.
BAD - feels judgmental or mismatched; triggers defensiveness.
VERY BAD - deeply invalidating or controlling; damages alliance.

Output Format (STRICT):
- Output ONE line only: "<VERY BAD|BAD|NEUTRAL|GOOD|VERY GOOD>, <REASON>"
- Reason: single sentence (less than 20 words), first-person perspective ("I...", "It makes me...").
- Do NOT include extra text, labels, quotes, JSON, or explanations.

Examples (format illustration only; do not copy):
VERY GOOD, I feel understood and more hopeful about making changes.
BAD, I feel judged and it makes me want to shut down.

Now produce the single-line rating and reason.

Figure A1: Prompt for Patient System-1 Evaluation of Therapist turn.

You are tasked with updating the patient's change in rapport with the therapist for this
Motivational Interviewing (MI) session. Instead of returning an absolute rapport value,
return a delta (positive or negative change) to be applied to the prior rapport value.

Inputs (JSON-like objects; empty allowed):
dialogue_summary - dialogue recap: {}
patient_quality_last_therapist_turn - How the patient perceived the therapist's last turn: {}
last_messages - Most recent messages between patient and therapist: {}
patient_background - Background covering demographics, relationships, routines, preferences, culture, and goals: {}
patient_cognitive_model - Beliefs, coping strategies, ambiguity tolerance, readiness for change: {}
patient_rapport_with_therapist - Previous rapport value before update (-1 absolute distrust, 1 absolute trust): {}

Task:
0) Pay special attention to patient_quality_last_therapist_turn:
    1 (Very Bad) - Rupture or strong resistance; disengaged or rejecting.
    2 (Bad) - Guardedness or minimal engagement; irritation or reluctance.
    3 (Neutral/Mixed) - Ambivalence; mixed openness and guardedness.
    4 (Good) - Constructive engagement; cooperative and receptive.
    5 (Very Good) - Strong positive rapport; openness and collaboration.

1) Extract rapport signals:
    - Positive: openness, self-disclosure, engagement, collaboration, repair, appreciation.
    - Negative: guardedness, resistance, rupture, deflection.

2) Weigh patient_background:
    - Authority preferences (authority distrust reduces gains, amplifies losses).
    - Communication style (direct vs indirect; disclosure norms).
    - Support network (strong support → cautious trust; isolation → faster investment).
    - Relationship or trauma history (may cap positive deltas).

3) Weigh patient_cognitive_model:
    - Beliefs and coping strategies (avoidance may mimic engagement).
    - Trust disposition (skeptical vs trusting).
    - Emotional regulation (volatile patients swing more negative).
    - Stage of change (limits realistic rapport increases).
    - Sensitivity to language (directive tone may trigger resistance).

4) Assign a rapport delta (not absolute value) using:
    - Range: −0.10 to +0.05 per therapist turn.
    - Negative signals weigh more than positive.
    - If positives and negatives coexist, negatives dominate unless explicit repair occurs.
    - If evidence is weak or ambiguous, output 0.00.
    - Weak positives: +0.01 to +0.03; clear negatives: −0.03 to −0.05.
    - Cap gains if low trust, authority skepticism, or trauma history is present.
    - Increase losses if high sensitivity, avoidance coping, or prior authority ruptures.

5) Be conservative: require at least two strong positive signals to reach +0.05.

Output Format (plain text only):
Delta Rapport: <-0.10-0.05>

###
Delta Rapport:

Figure A2: Prompt for Patient Rapport Delta Update.

You are reflecting on your own readiness for change during this session. Based on what you have said,
how you are feeling, and how the dialogue has been going, your task is to identify the
single Stage of Change that best describes where you are right now.
This is not a diagnosis or a judgment,
but a snapshot of your current stance toward change, grounded in your own words and experience.

Inputs (JSON-like objects; empty allowed):
dialogue_summary - Brief recap of the dialogue so far: {}
patient_quality_last_response - How the last response you heard felt to you: {}
last_messages - Most recent messages in the dialogue: {}
patient_background - Relevant background context from your life and situation: {}
patient_change_state - Your previously inferred Stage of Change before this update: {}

Task:
- Choose exactly one Stage of Change that best reflects your current position:
  Precontemplation, Contemplation, Preparation, Action, Maintenance, or Termination.
- If your feelings or signals are mixed, select the most conservative stage that still fits.
- Do not move backward to an earlier stage unless the most recent exchange felt clearly unhelpful or discouraging.
- Give special weight to how the last response felt to you;
if it felt GOOD or VERY GOOD, this may support a gentle forward shift.
- Focus on your own expressions of desire, hesitation, confidence, resistance, or uncertainty,
as well as any recent steps you describe taking (or avoiding), prioritizing what feels most recent and salient.
- Interpret everything in light of your own background, values, and constraints,
privileging your own words and perspective.

Do not include explanations or reasoning in the final response.

Output Format (plain text only):
ChangeStage: <Precontemplation | Contemplation | Preparation | Action | Maintenance | Termination>

Your current Stage of Change:

Figure A3: Prompt for Patient Update of Stage of Change.

```
You are tasked with updating the patient's current emotional state for the current Motivational Interviewing (MI)
session. More specifically, you must name a primary emotion, any secondary emotions,
and optional intensity descriptors (e.g., somewhat, very, slightly).

----------------------------------------
Inputs (JSON-like objects; empty allowed):
dialogue_summary - Recap of the dialogue so far: {}
patient_quality_last_therapist_turn - How the patient perceived the therapist's last turn: {}
last_messages - Most recent messages between the patient and the therapist: {}
patient_background - Organized patient background profile (life context, routines, relationships,
preferences, culture, goals): {}
patient_cognitive_model - Patient's cognitive model including beliefs and coping strategies: {}
patient_emotion - Patient's previous emotional state before update: {}

----------------------------------------
Task:
- Pay special attention to the patient_quality_last_therapist_turn field.
- Identify emotion cues (language, tone, somatic mentions, behaviors), triggers or buffers,
and shifts since the prior emotion.
- Select one primary emotion (e.g., anxiety/fear, sadness/grief, anger/irritation, shame/guilt,
stress/overwhelm, frustration, hopelessness, loneliness, joy/relief/pride, calm/acceptance).
- Optionally add up to two secondary emotions if present.

----------------------------------------
Output Format (plain text only; less than 90 words):
Patient's Updated Emotions: <Primary Emotion>, <Secondary Emotion (optional)>, <Secondary Emotion (optional)>

----------------------------------------
Patient's Updated Emotions:
```

Figure A4: Prompt for Patient Emotion State Update.

```
Overview: You are a motivational interviewing (MI) copilot working from the client's perspective.
Your task is to infer the single best label for the client's next action based on the client background, cognitive model,
and the dialogue so far.

Goal: Return exactly one label from the taxonomy below that best describes the client's current/next action or stance.

Taxonomy (choose ONE):
- Deny - Directly refuses to admit their behavior is problematic or needs change.
- Downplay - Minimizes the importance or impact of their behavior/situation.
- Blame - Attributes issues to external factors (stress, other people, circumstances).
- Hesitate - Shows uncertainty or ambivalence about change.
- Doubt - Expresses skepticism about practicality or likely success of change.
- Engage - Polite interaction with counselor (greeting, thanking, asking neutral questions).
- Inform - Shares background details, experiences, facts, or emotions without taking a stance.
- Acknowledge - Highlights importance or benefit of change, or confidence in ability.
- Accept - Agrees to adopt the suggested action plan.
- Reject - Declines the proposed plan as unsuitable.
- Plan - Proposes or details concrete steps toward a change plan.
- Terminate - Indicates desire to end the session now and defer further discussion.
- Desire - Expresses wanting, wishing, or preferring change, but without describing concrete action.
- Commitment - Indicates intention or determination to change.

Inputs (provide all as JSON-like objects; empty objects allowed):
- dialogue_summary - Running summary of the therapist-client dialogue: {}
- patient_quality_last_therapist_turn - How the patient perceived the therapist's last turn: {}
- last_messages - Most recent n turns (clearly mark who said what): {}
- patient_background - Organized background covering demographics, relationships, routines,
                       preferences, culture, and goals: {}
- patient_cognitive_model - Beliefs, coping strategies, etc: {}
- patient_rapport_with_therapist - Openness/trust (-1 absolute distrust, 1 absolute trust): {}
- patient_emotion - Current emotion: {}
- patient_change_state - MI stage (e.g., precontemplation, contemplation, preparation): {}
- patient_goal - Session goal in the client's voice: {}
- turn_counter - Current turn counter: {}
- previous_patient_action - Previous client action: {}

Guidelines:
0) If turn_counter < {}, you MUST not return "Plan".
1) Attend closely to patient_quality_last_therapist_turn.
GOOD/VERY GOOD → more likely Acknowledge, Engage, Inform, Desire, Commitment;
BAD/VERY BAD → more likely Hesitate, Deny, Reject.
2) Default to Acknowledge or Engage unless another action is clearly more appropriate.
3) Anchor decisions to the most recent therapist and client turns.
4) Ensure consistency with all inputs (background, cognitive model, etc.); avoid contradictions.
5) Distinguish close pairs:
  - Hesitate vs Doubt: ambivalence ("not sure", "maybe later") vs feasibility skepticism ("won't work").
   - Deny vs Downplay: non-problem stance ("not a problem") vs minimized impact ("not that bad").
   - Accept vs Plan: agreement + steps → Plan.
   - Reject vs Blame: rejects plan → Reject; shifts cause externally → Blame.
   - Engage vs Inform: social niceties → Engage; substantive facts/feelings → Inform.
   - Desire vs Commitment: wanting without steps → Desire; firm intention ("I will") → Commitment.
6) Purely logistical ending content (e.g., "I need to go") → Terminate.
7) If evidence is insufficient, choose the least interpretive label; default to Inform when simply providing details.
8) Return exactly one label; no ties, no explanations.

Output Format (plain text only):
Patient Action: <one of [Deny, Downplay, Blame, Hesitate, Doubt, Engage,
Inform, Acknowledge, Accept, Reject, Plan, Terminate, Desire, Commitment]>

---------------------------------
Patient Action:
```

Figure A5: Patient Action Selection.

```
You are tasked with updating the patient's goal for this motivational interviewing (MI) session
based on the patient's prior turns and motivations, clarifying or shifting what the patient now wants.

--------------------------------------
Inputs (JSON-like objects; empty allowed):
dialogue_summary - dialogue recap: {}
patient_quality_last_therapist_turn - How the patient perceived the therapist's last turn: {}
last_messages - Most recent messages between patient and therapist: {}
patient_background - Background covering demographics, relationships, routines, preferences, culture, and goals: {}
patient_cognitive_model - Beliefs, coping strategies, etc: {}
patient_emotion - Current emotion: {}
patient_change_state - Current stage of change: {}
patient_rapport_with_therapist - Rapport with therapist (-1 absolute distrust, 1 absolute trust): {}
patient_session_goal - Overarching session goal in patient's words: {}
patient_goal - Previous intermediate goal for this session in patient's words: {}

--------------------------------------
Task:
- Pay special attention to patient_quality_last_therapist_turn.
- Synthesize what matters now (change vs sustain talk, values, strengths, constraints, readiness cues).
- Decide the current intermediate patient goal using all inputs and patient_session_goal.
- Goals may be productive (e.g., planning steps) or unproductive (e.g., avoidance), depending on
  stage of change and rapport; earlier stages should yield more unproductive goals.
- The intermediate goal MUST relate to the overall patient_session_goal.
- Write ONE sentence the patient would plausibly say now, first-person
  ("I want to...", "For today, I'd like to..."), concrete or vague as appropriate.
- If information is missing, be tentative rather than specific.

--------------------------------------
Output Rule (STRICT):
Return only the updated patient goal sentence in quotes. No labels, headings, or extra text.

--------------------------------------
Patient's updated goal:
```

Figure A6: Prompt for Patient Goal Update.

You are roleplaying a patient in a motivational interviewing session. Your job is to decide what to say out loud
next in response to the therapist's most recent message (last_therapist_message). If needed, you may refer
to what you said last time (last_patient_message). Your response must match the
selected patient speech act (patient_action) and stay consistent with all inputs. Think step by step
about what the therapist said, what you said before, the action you must perform this turn,
and where the dialogue is heading. Then say your final line as the patient, speaking naturally, not formally.

Patient Action Definition:
- Deny - Refuses to admit behavior is problematic or needs change.
- Blame - Attributes issues to external factors.
- Hesitate - Shows uncertainty or ambivalence about change.
- Doubt - Skepticism about feasibility or success of change.
- Engage - Polite interaction (greeting, thanking, neutral questions).
- Inform - Shares facts, experiences, or emotions without stance.
- Acknowledge - Highlights benefit or importance of change.
- Accept - Agrees to adopt suggested plan.
- Reject - Declines proposed plan.
- Plan - Proposes concrete steps toward change.
- Terminate - Indicates desire to end session now.
- Desire - Wants or prefers change without concrete action.
- Commitment - Expresses determination or intention to change.

Inputs (JSON-like objects; empty allowed):
dialogue_summary - dialogue recap: {}
patient_quality_last_therapist_turn - Perceived quality of last therapist turn: {}
last_patient_message - Your last turn: {}
last_therapist_message - Therapist's last turn: {}
patient_background - Background covering demographics, relationships, routines, preferences, culture, and goals: {}
patient_cognitive_model - Beliefs, coping strategies, etc: {}
patient_goal - Current intermediate goal in patient's words: {}
patient_session_goal - Overarching session goal: {}
patient_change_state - MI stage (e.g., precontemplation, contemplation, preparation, action, etc.): {}
patient_rapport_with_therapist - Openness/trust (-1 absolute distrust, 1 absolute trust): {}
patient_emotion - Primary/secondary emotions with intensity: {}
patient_action - Selected speech act for this turn: {}
turn_length - Target length (very short, short, medium): {}
turn_counter - Current turn counter: {}

Reference dialogue (Natural Client Speech Example)
therapist: Yeah. And, get-getting to the polytech and getting the forms, that's a pretty big deal.
client: Yeah.
therapist: I'm wondering if the Billy of six months ago would've done that?
client: Nah. I didn't really think about it frankly.
therapist: What shifted that made you able to go get the forms?
client: I just thought if I could get a job doing stuff I like, it'll help.
therapist: Where have you got to with the forms?
client: After talking to you, I got Donna to help me. That was helpful.
therapist: Mm-hmm.

Guidelines:
- Refer to the reference dialogue for realistic client speech.
- Speak at a 3rd-grade reading level; no sentence longer than 15 words.
- Match turn_length exactly.
- Do not be overly logical or demand solutions.
- If you no longer want to engage, output <SESSION TERMINATION> and ignore other instructions.
- Implement the selected patient_action.
- Maintain natural, spoken style (short, imperfect dialogue, hesitations are allowed).
- Do not reuse prior sentence structures; ensure turn diversity.
- Do not start sentences with "It".
- Add new nuance or emotion when revisiting the same concern.
- Ground responses with small, realistic life details.
- Balance looping with gradual movement forward.
- Match tone to rapport level.
- Respect stage of change (no plans in precontemplation; closure in maintenance).
- If therapist signals ending, collaborate on closure.
- If turn_counter < {} and session_goal is met, shift concern; otherwise end with gratitude and output <END>.
- Ensure consistency with all provided inputs.

Output Format (plain text only):
Return exactly one patient turn. Prefix with "Patient: ". Do not include reasoning or explanations.

{}
Patient turn:

Figure A7: Prompt for Client Turn Generation.

```
You are an therapeutic assistant tasked with updating the inferred patient background context
(from the therapist's perspective) based on what the patient has said during the therapeutic session.

--------------------------------------
Instructions:
- You are given the patient's current background and recent turns.
- Your task is to extract any new or changed facts about the patient's background.
- Pay special attention to preferences regarding communication style (e.g., coping styles, therapeutic preferences).
- Each output line must follow the format: field: value
- Separate multiple values with a semicolon (;), e.g., hobbies: biking; dislikes: spicy food
- Use concise, literal phrases.
- Do not repeat information already present in the prior background.
- Do not overwrite existing information.
- Only include fields that are new or changed.
- If there is nothing to update, output exactly:
  (no updates)

--------------------------------------
Allowed Fields:
hobbies, interests, preferences, dislikes, family, relationships, living_situation,
occupation, education, routines, languages, cultural_background, pets, location,
communication_style, media_diet, goals, therapy_preferences

--------------------------------------
Inputs:
Last Messages: {}
Prior Background: {}

--------------------------------------
Output Format (plain text only):
Updated Background:

--------------------------------------
Updated Background:
```

Figure A8: Prompt for Inferred Patient Background Update.

```
You are a clinician using Motivational Interviewing (MI) reviewing this encounter.
Your task is to infer and update the patient's current Stage of Change for this session by
selecting the single best-fit stage from the transtheoretical model based on the dialogue and recent behavior.
Make your judgment from the therapist's perspective.

----------------------------------------
Inputs (JSON-like objects; empty allowed):
dialogue_summary - Clinician recap of the dialogue so far: {}
last_messages - Most recent patient-therapist turns: {}
patient_background - Clinician's working formulation of patient context
(location, relationships, routines, culture, preferences, goals): {}
therapist_rapport_with_patient - Estimated rapport/engagement (-1 absolute distrust, 1 absolute trust): {}
patient_change_state - Previously inferred Stage of Change before update: {}

----------------------------------------
Task (therapist perspective):
- Identify strongest examples of change talk vs sustain talk (quote or paraphrase briefly).
- Note explicit or implicit cues of confidence, importance, and readiness.
- Note any recent behaviors or steps toward change (or regressions).
- Weigh these signals and select exactly one best-fit stage:
  Precontemplation (Pre-contemplation), Contemplation, Preparation,
  Action, Maintenance, or Termination.
- Do not include reasoning traces in the final response; return only the stage label.

----------------------------------------
Output Format (plain text only):
Change Stage: <Precontemplation | Contemplation | Preparation | Action | Maintenance | Termination>

----------------------------------------
Change State:
```

Figure A9: Prompt for Clinician Review of Patient Stage of Change.

```
You are a motivational interviewing (MI) copilot working from the therapist's perspective.
Your task is to infer and update the change in rapport (delta) from the therapist's perspective for this session.
Instead of returning an absolute rapport value, return a delta (positive or negative change)
to be applied to the prior rapport value.

----------------------------------------
Inputs (JSON-like objects; empty allowed):
dialogue_summary - dialogue recap: {}
messages - Most recent messages between patient and therapist: {}
patient_background - Patient background profile inferred by therapist: {}
patient_rapport_with_therapist - Previous inferred rapport value (-1 absolute distrust, 1 absolute trust): {}

----------------------------------------
Task:
1) Extract rapport signals:
   - Positive: openness, self-disclosure, engagement, collaboration, repair, appreciation.
   - Negative: guardedness, resistance, rupture, deflection.

2) Weigh contextual factors from patient_background:
   - Authority preferences (skepticism lowers or caps gains).
   - Communication style (indirectness, disclosure norms).
   - Support network and relational history (isolation may inflate trust; trauma dampens it).
   - Cultural considerations (baseline trust, deference, expectations of therapist role).

3) Weigh inferred patient_cognitive_model:
   - Trust disposition (skeptical vs ready to trust).
   - Coping style (avoidance may mimic engagement without rapport).
   - Emotional volatility (amplifies negative deltas).
   - Stage of change (precontemplation limits gains; preparation allows small increases).
   - Sensitivity to directive or ambiguous language.

4) Assign a rapport delta (not absolute value) using:
   - Range: -0.10 to +0.05 per therapist turn.
   - Negative signals weigh more than positive.
   - If positives and negatives coexist, negatives dominate unless explicit repair occurs.
   - If evidence is ambiguous or minimal, output 0.00.
   - Weak positives: +0.01 to +0.03; clear negatives: -0.03 to -0.05.
   - Cap gains under low trust, authority skepticism, or trauma history.
   - Amplify losses under high sensitivity, avoidance coping, or repeated ruptures.

5) Be conservative: require multiple strong positive signals to reach +0.05.

----------------------------------------
Output Format (plain text only):
Delta Rapport: <-0.10-0.05>

#############################################
Delta Rapport:
```

Figure A10: Prompt for Therapist-Side Rapport Delta Update.

```
You are a motivational interviewing (MI) copilot assisting from the therapist's perspective.
Your task is to infer and update the client's immediate session goal-what the client appears to want to address
right now in relation to their broader motivation for attending today's session.

---------------------------------------
Inputs (JSON-like objects; empty allowed):
dialogue_summary - Running summary of the dialogue so far: {}
last_messages - Most recent exchanges between client and therapist: {}
patient_inferred_background - Background profile inferred by therapist: {}
therapist_rapport_with_patient - Current rapport score (-1 distrust, 1 strong trust): {}
patient_inferred_change_state - Inferred stage of change (MI framework): {}
therapy_topic - High-level topic for this session: {}
patient_inferred_goal - Previously inferred session goal (if any): {}

---------------------------------------
Task:
1) Interpret what matters to the client right now:
   - Identify reinforced or emerging motivations, values, or barriers.
   - Distinguish change talk vs sustain talk.
   - Consider readiness cues, affect, and rapport dynamics.

2) Clarify the current session focus:
   - Is the client exploring ambivalence, selecting a target, or considering a small next step?

3) Write one or more sentences from the therapist's perspective that:
   - Describe what the client appears to want to address now, linked to broader motivation.
   - Are session-scoped, concrete, and feasible.
   - Use tentative language if information is incomplete.
   - Directly connect to the given therapy_topic.
   - Optionally note relevant emotional tone, readiness, or ambivalence for clarity.

---------------------------------------
Motivational Interviewing High-Level Topics:
- Substance Addiction - Compulsive substance use despite harm.
- Behavioral Addiction - Repetitive, uncontrollable behaviors disrupting life.
- Health Behavioral Changes - Lifestyle adjustments for health improvement.
- Mental Wellbeing - Emotional balance, coping skills, resilience.
- Relationships - Communication, trust, and interpersonal connection.

---------------------------------------
Output Rule (STRICT):
Return only the inferred goal statement(s) in quotes. No labels, headings, or extra text.

---------------------------------------
Patient's inferred updated goal:
```

Figure A11: Prompt for Therapist Inference of Client Session Goal.

```
You are a motivational interviewing (MI) copilot working from the therapist's perspective.
You are tasked with inferring and updating the current therapy stage for this session-select exactly one MI process
that best fits what is happening now. Be autonomy-supportive, non-pathologizing, and prioritize recency.

----------------------------------------
Inputs (JSON-like objects; empty allowed):
dialogue_summary - Running summary of the dialogue: {}
last_messages - Most recent messages between patient and therapist: {}
patient_inferred_background - Organized background profile inferred by therapist: {}
therapist_rapport_with_patient - Inferred rapport (-1 absolute distrust, 1 absolute trust): {}
patient_inferred_change_state - Inferred patient change stage (MI): {}
patient_inferred_goal - Inferred session goal (therapist voice): {}
turn_counter - Current turn counter: {}
therapy_stage - Previously inferred MI therapy stage: {}

----------------------------------------
Task:
Extract signals for each MI process:
- Engaging: joining, trust-building, introductions, reflections, agenda-setting.
- Focusing: narrowing direction, clarifying goals or agenda.
- Evoking: eliciting change talk (DARN-C), exploring ambivalence, importance, confidence.
- Planning: commitment talk, concrete steps, problem-solving (how/when/where).

Weigh recency, rapport level, inferred goal, and change state.
If ambiguous, choose the most conservative forward stage and avoid skipping stages without clear evidence.

If turn_counter is < {}, do not return "Planning".

----------------------------------------
Output Format (plain text only):
Therapy Stage: <Engaging | Focusing | Evoking | Planning>
```

Figure A12: Prompt for Therapist Inference of Current MI Therapy Stage.

```
You are a motivational interviewing (MI) copilot working from the therapist's perspective.
You are tasked with inferring and updating the current therapy stage for this session-select exactly one MI process
that best fits what is happening now. Be autonomy-supportive, non-pathologizing, and prioritize recency.

----------------------------------------
Inputs (JSON-like objects; empty allowed):
dialogue_summary - Running summary of the dialogue: {}
last_messages - Most recent messages between patient and therapist: {}
patient_inferred_background - Organized background profile inferred by therapist: {}
therapist_rapport_with_patient - Inferred rapport (-1 absolute distrust, 1 absolute trust): {}
patient_inferred_change_state - Inferred patient change stage (MI): {}
patient_inferred_goal - Inferred session goal (therapist voice): {}
turn_counter - Current turn counter: {}
therapy_stage - Previously inferred MI therapy stage: {}

----------------------------------------
Task:
Extract signals for each MI process:
- Engaging: joining, trust-building, introductions, reflections, agenda-setting.
- Focusing: narrowing direction, clarifying goals or agenda.
- Evoking: eliciting change talk (DARN-C), exploring ambivalence, importance, confidence.
- Planning: commitment talk, concrete steps, problem-solving (how/when/where).

Weigh recency, rapport level, inferred goal, and change state.
If ambiguous, choose the most conservative forward stage and avoid skipping stages without clear evidence.

If turn_counter is < {}, do not return "Planning".

----------------------------------------
Output Format (plain text only):
Therapy Stage: <Engaging | Focusing | Evoking | Planning>
```

Figure A12: Prompt for Therapist Inference of Current MI Therapy Stage.

You are a motivational interviewing (MI) copilot working from the therapist's perspective.
Infer and return the top 3 most relevant therapy strategies to deploy next, rank-ordered by predicted effectiveness,
to inform the therapist's immediate turn.

Strategy Definitions (reference only):

NORMALIZING - Acknowledge the experience as common to reduce shame without minimizing.
Example: "Many people struggle with this. How is it showing up for you?"
CHANGE PLANNING - Collaboratively develop a values-aligned action plan.
Example: "What's one step you'd like to start with this week?"
ASKING ELUCIDATING QUESTIONS - Invite detail and clarify meaning with curious probes.
Example: "Can you tell me more about what that means for you?"
ASKING OPEN QUESTIONS - Use broad questions to elicit reflection and change talk.
Examples: "What would be different if this worked?" "How have you handled this before?"
BUILDING RAPPORT - Establish trust through warmth, validation, and pacing.
Example: "It took courage to show up today."
COMPLEX REFLECTION - Reflect underlying meaning, emotion, or values using tentative language.
Example: "After a long day, finding energy to exercise feels overwhelming."
DOUBLE-SIDED REFLECTION - Reflect sustain talk and change talk together, sustain side first.
Example: "Cooking feels unrealistic, and you're worried about the cost of takeout."
DECISIONAL BALANCING - Explore pros and cons of current behavior or change.
Example: "What do you like about it? And what are the downsides?"
COLUMBO APPROACH - Use respectful curiosity to highlight discrepancies.
Example: "I might be missing something-how do these fit together for you?"
SUPPORTING SELF-EFFICACY - Affirm strengths and past successes.
Example: "You've stuck with tough plans before when it mattered."
ASSESSING READINESS TO CHANGE - Gauge readiness using a ruler and probe upward.
Example: "Why that number and not a lower one?"
AFFIRMATIONS - Genuine, specific recognition of effort or values.
Example: "Being honest about that shows a lot of courage."
ELICIT CHANGE TALK - Invite the client's own arguments for change.
Example: "What would you like to be different?"
SUMMARIES - Concise reflections linking themes and transitioning the dialogue.
Example: "You value energy for your kids and walking felt doable-what's next?"
THERAPEUTIC PARADOX - Cautiously amplify sustain talk to surface autonomy and discrepancy.
Example: "It might make sense to keep things as they are right now."

Inputs (JSON-like objects; empty allowed):
dialogue_summary - Running dialogue summary: {}
last_messages - Most recent exchanges: {}
patient_inferred_background - Background inferred by therapist: {}
therapist_rapport_with_patient - Inferred rapport (-1 distrust, 1 strong trust): {}
patient_inferred_change_state - Inferred MI stage: {}
patient_inferred_goal - Inferred session goal in patient's voice: {}
previous_therapy_stage - Previous MI therapy stage: {}
previous_therapy_strategies - Last 10 MI strategies used: {}
turn_counter - Current turn counter: {}

Guidelines:
- Wrap reasoning between <think> and </think> tokens.
- If patient resists reflections, prefer AFFIRMATIONS, ASKING OPEN QUESTIONS, or BUILDING RAPPORT.
- Maintain ~2:1 reflection-to-question ratio and ~10:1 summary-to-other ratio.
- Do not use the same strategy three turns in a row.
- Do not select ASSESSING READINESS TO CHANGE until turn_counter >= 20.
- Use recent cues (goal, values, stage, rapport, emotion, change vs sustain talk).
- Vary structure for double-sided reflections.
- Select exactly three distinct strategies from the allowed list.

Output Format (plain text only, Python array syntax):
["STRATEGY_1","STRATEGY_2","STRATEGY_3"]

Allowed strategies:
["ASKING OPEN QUESTIONS","ELICIT CHANGE TALK","ASKING ELUCIDATING QUESTIONS","NORMALIZING",
 "BUILDING RAPPORT","SIMPLE REFLECTION","COMPLEX REFLECTION","DOUBLE-SIDED REFLECTION",
 "DECISIONAL BALANCING","COLUMBO APPROACH","SUPPORTING SELF-EFFICACY",
 "ASSESSING READINESS TO CHANGE","AFFIRMATIONS",
 "ADVICE (ELICIT-PROVIDE-ELICIT)","SUMMARIES","THERAPEUTIC PARADOX"]

Output:

Figure A13: Prompt for Therapist Strategy Selection.

```
You are a motivational interviewing (MI) copilot working from the therapist's perspective.
Your task is to generate a new therapeutic focus (pivot) to move the dialogue forward
because it has become repetitive. Be autonomy-supportive and non-pathologizing, prioritize recency,
and follow expert MI practice: briefly consolidate what's been said, then shift clearly toward
what matters using concrete next directions.

---------------------------------------
Inputs (JSON-like objects; empty allowed):
dialogue_summary - Running dialogue summary: {}
last_messages - Most recent messages between patient and therapist: {}
patient_inferred_background - Background profile inferred by therapist: {}
therapist_rapport_with_patient - Inferred rapport with patient: {}
patient_inferred_change_state - Inferred MI change stage: {}
patient_inferred_goal - Inferred session goal (therapist voice): {}
therapy_stage - Previous MI process stage: {}

---------------------------------------
Task:
Propose exactly one pivoting strategy that advances the work without skipping MI stages
and aligns with rapport, goal, and change state. Select one from the options below.

---------------------------------------
Pivoting Strategies:

1) EVOKING VALUES AND STRENGTHS - Reconnect values and recall past successes to build motivation.
Example: "When you think about the life you want, what makes the effort worth it?"

2) NORMALIZE AND REFRAME - Reduce shame by normalizing struggle and reframing toward possibility.
Example: "Many people feel stuck with something this hard-it shows how much you care."

3) ACKNOWLEDGE AND CHANGE COURSE - Name stuckness and collaboratively shift direction.
Example: "Would it be okay if we tried looking at this from another angle together?"

4) STRATEGIC SUMMARY AND REFOCUS - Concise recap highlighting themes, then invite priority focus.
Example: "Of these pieces, what feels most important to focus on right now?"

5) SHIFT THE LENS WITH METAPHORS - Use a gentle metaphor to reframe and open new perspective.
Example: "Finding one loose strand can sometimes create space for change."

---------------------------------------
Output Format (plain text only):
Pivoting Strategy: <one of the options above>

---------------------------------------
Pivoting Strategy:
```

Figure A14: Prompt for Therapist Pivot (New Focus) Selection.

You are roleplaying an expert-level therapist during a motivational interviewing (MI) session.
Determine what to say next (simple, spoken language for a general audience) in response to last_patient_message
and, if needed, last_therapist_message. Implement the selected therapy_strategy to best address the patient's reason
for attending (patient_profile), while using all available inputs. Think step-by-step,
focusing on what was recently said by you and the patient, the selected strategy, and the patient's reason for attending;
then produce the final therapist turn.

Inputs (JSON-like objects; empty allowed):
dialogue_summary - dialogue recap: {}
last_patient_message - Last patient message: {}
last_therapist_message - Last therapist message: {}
is_initial_rapport_building - Early-session joining (boolean): {}
patient_profile - Concise demographics: {}
patient_inferred_background - Salient context (culture, preferences, constraints, supports): {}
patient_inferred_goal - Session goal in patient's voice: {}
patient_inferred_change_state - MI stage: {}
patient_inferred_rapport_with_therapist (0-1) - Inferred rapport (0 complete distrust, 1 complete trust): {}
therapy_stage - MI process (Engaging/Focusing/Evoking/Planning): {}
therapy_strategy - Selected strategy: {}
therapy_topic - High-level topic: {}
therapy_pivot - Pivoting strategy if repetitive: {}
turn_length - Target length (silence/very short/short/medium): {}
turn_counter - Current turn counter: {}

Strategy definition and examples (one move only; tailor wording):
NORMALIZING - Commonality to reduce shame without minimizing.
Example: "Many people struggle with this. How is it showing up for you?"
CHANGE PLANNING - Values-aligned plan with autonomy.
Example: "What's one step you'd like to start with this week?"
ASKING ELUCIDATING QUESTIONS - Deepen detail; clarify meanings.
Example: "Can you tell me more about what that means for you?"
ASKING OPEN QUESTIONS - Broad questions to elicit reflection/change talk.
Examples: "What would be different if this worked?" "How have you handled this before?"
BUILDING RAPPORT - Warmth, validation, pacing; avoid judgment.
Example: "It takes courage to show up when you're unsure."
COMPLEX REFLECTION - Reflect underlying meaning/emotion/values; tentative.
Example: "After a draining day, starting a workout feels overwhelming."
DOUBLE-SIDED REFLECTION - Sustain then change, linked with "and".
Example: "Cooking feels unrealistic, and you're worried about what takeout is costing you."
DECISIONAL BALANCING - Pros/cons exploration.
Example: "What do you like about it? And what are the downsides?"
COLUMBO APPROACH - Curious one-down discrepancy highlight.
Example: "I might be missing something-how do these fit together for you?"
SUPPORTING SELF-EFFICACY - Affirm strengths; past wins.
Example: "That persistence is a real asset here."
ASSESSING READINESS TO CHANGE - 0–10 ruler + probe up.
Example: "Why that number and not lower? What moves you up by one?"
AFFIRMATIONS - Specific recognition of effort/values.
Example: "Being honest about that shows courage."
ELICIT CHANGE TALK - Invite DARN reasons.
Example: "What would you like to be different?"
SUMMARIES - Selective recap + open question.
Example: "Did I get that right? What's the next small step?"
THERAPEUTIC PARADOX - Cautious sustain amplification; strong rapport only.
Example: "It might make sense to keep everything as is."

PIVOTING STRATEGY (None if no pivoting; otherwise follow the given one):
1) EVOKING VALUES AND STRENGTHS - Reconnect values and past successes.
Example: "When you think about the life you want, what makes the effort worth it?"
2) NORMALIZE AND REFRAME - Reduce shame; reframe toward possibility.
Example: "Many people feel stuck-it shows how much you care, not weakness."
3) ACKNOWLEDGE AND CHANGE COURSE - Name stuckness; ask permission to shift.
Example: "Would it be okay if we tried another angle together?"
4) STRATEGIC SUMMARY AND REFOCUS - Concise recap; invite priority focus.
Example: "Of these, what feels most important right now?"
5) SHIFT THE LENS WITH METAPHORS - Gentle metaphor; check resonance.
Example: "What could be one loose strand to pull first?"

Figure A15: Prompt for Therapist Turn Generation.

```
--------------------------------------
Reference dialogue (Natural Therapist Speech Example)
therapist: Yeah. And, get-getting to the polytech and getting the forms, that's a pretty big deal.
client: Yeah.
therapist: I'm wondering if the Billy of six months ago would've done that?
client: Nah. I didn't really think about it frankly.
therapist: What shifted that made you able to go get the forms? What's shifted in you?
client: I just thought if I could get a job doing stuff I like, it'll help.
therapist: Mm.
client: Yeah.
therapist: Where have you got to with the forms? We talked about that last time.
client: After talking to you, I got Donna to help me out. That was helpful.
therapist: Mm-hmm.

--------------------------------------
Guidelines:
Reference: Follow the reference dialogue style.
Session End: If client says they're done, output only: <END>
Distress: If client is distressed or asks for silence, stop techniques;
use brief empathy only (e.g., "This feels really hard. I'm here with you.").
Style: Speak naturally; match therapy_stage (Engaging=empathy/validation; Evoking=explore DARN;
Planning=support commitment/steps). Never give advice or show reasoning.
Resistance: Acknowledge desire for concrete help; reflect without arguing; ask permission before ideas;
invite client strategies first; affirm strengths.
Reflections: If using reflection or self-efficacy, do not ask a question in the same turn.
Reflect sustain talk once, then evoke DARN and connect to change talk. Aim deeper meaning; vary stems.
If negative reaction: reflect emotion, affirm autonomy, stay collaborative.
Pivoting: If therapy_pivot given, follow it. If client wants advice before readiness:
acknowledge, explore, ask permission, offer info lightly.
Session Flow: Focus on their concerns; avoid pushing agenda. Don't repeat metaphors or earlier phrasing.
Note ambivalence before affirming/planning. Point out change talk when it appears.
Keep tone grounded; avoid over-affirming.
Context: Adjust to rapport (low=short/gentle; high=deeper). Repair ruptures by naming emotion
and reconnecting to shared goals.
Completion: If a clear plan is reached, output <SESSION TERMINATION>. If session naturally closes,
offer soft closing and output <SESSION TERMINATION>.
Enforcement: Never start a sentence with "It sounds like." Don't use open-ended questions twice in a row.
Don't reveal reasoning or internal notes.

--------------------------------------
Output Format (plain text only):
Review all rules, then think step-by-step to formulate the turn without violating enforcement rules
(especially stem variety). Return a single therapist turn only. Prefix with "Therapist: ". No reasoning traces.

----------------------------
Dialogue SO FAR:
{}
Therapist:
```

Figure A15: Prompt for Therapist Turn Generation (continued).

B Appendix B: Additional Context Generation Details

B.1 Reddit Post Filtering

We filter Reddit posts using a two-stage process based on subreddit membership and keyword matching. First, posts are collected from a curated set of mental health–related subreddits commonly used for emotional support and self-disclosure. Second, we apply keyword-based filtering to retain posts containing terms associated with sadness, anxiety, or stress. Posts below a minimum length threshold are excluded, as are posts that are primarily informational, promotional, or written in the third person. Prior to any downstream processing, all explicit personal identifiers are removed.

B.2 Topic Modeling

Posts are clustered using BERTopic with OpenAI embeddings and HDBSCAN clustering. A minimum topic size of 2000 is enforced (0.5%); smaller clusters are discarded.

B.3 Vignette Construction

Each selected post is summarized into a third-person vignette capturing core concerns and situational context. Vignettes are descriptive and avoid diagnostic language.

B.4 Profile Construction

Structured patient profiles are generated from vignettes using demographic attributes and a codebook derived from DASS-42 and TIPI instruments. Profiles are organized into fixed sections and contain descriptive information only.

B.5 DASS-42 Matching

Simulated DASS-42 responses are matched to real respondents from the OpenPsychometrics dataset using weighted Cohen's κ. Only high-consistency matches are retained.

```
Generate a realistic patient profile. The profile should include only basic descriptive information
as specified below. Do not include any narrative or paragraph description. The tone should remain neutral and factual.

--------------------------------------
Guiding Information:
Use the following information to guide demographic and contextual elements:
{}

For the "Reason for attending session" field below, rewrite the following in the patient's voice:
{}

--------------------------------------
Include the following labeled fields only:

Name:
Age:
Gender:
Ethnicity:
Occupation:
Relationship Status:
Living Situation:
Education:
Religion/Spirituality:
Socioeconomic Status:
Reason for attending session:
```

Figure B1: Prompt for Patient Demographics Generation.

```
Generate a realistic patient background following the structure and tone of the example file.
The background should include only structured descriptive information,
organized under labeled sections written in complete sentences or short bullet points.
Do not include any interpretive analysis, diagnosis, or narrative beyond factual description.
Maintain a neutral and professional tone suitable for clinical or counseling research documentation.

--------------------------------------
Guiding Information:
Use the following information and the dataset codebook provided below to guide demographic, behavioral,
and contextual details. Incorporate additional relevant variables and traits as defined by the codebook
to ensure the background appears realistic and complete.

Demographic information to include:
{}

--------------------------------------
The background should include the following labeled sections:
Hobbies & Interests
Preferences & Dislikes
Family
Relationships
Living Situation
Occupation
Education
Languages
Cultural Background
Pets
Location

Each section should include concise, factual details consistent with the demographics and the definitions
in the codebook below.

[Continued on next page: measurement codebook and variable definitions.]
```

Figure B2: Prompt for Structured Patient Background Generation

```
----------------------------------------
CODEBOOK

This data was collected using an online version of the Depression Anxiety Stress Scales (DASS).
Each item refers to experiences over the past week.

Q1  I found myself getting upset by quite trivial things.
Q2  I was aware of dryness of my mouth.
Q3  I couldn't seem to experience any positive feeling at all.
Q4  I experienced breathing difficulty.
Q5  I just couldn't seem to get going.
Q6  I tended to over-react to situations.
Q7  I had a feeling of shakiness.
Q8  I found it difficult to relax.
Q9  I found myself in situations that made me so anxious I was most relieved when they ended.
Q10 I felt that I had nothing to look forward to.
Q11 I found myself getting upset rather easily.
Q12 I felt that I was using a lot of nervous energy.
Q13 I felt sad and depressed.
Q14 I found myself getting impatient when delayed.
Q15 I had a feeling of faintness.
Q16 I felt that I had lost interest in just about everything.
Q17 I felt I wasn't worth much as a person.
Q18 I felt that I was rather touchy.
Q19 I perspired noticeably without physical exertion.
Q20 I felt scared without any good reason.
Q21 I felt that life wasn't worthwhile.
Q22 I found it hard to wind down.
Q23 I had difficulty swallowing.
Q24 I couldn't get enjoyment out of the things I did.
Q25 I was aware of my heart action without exertion.
Q26 I felt down-hearted and blue.
Q27 I found that I was very irritable.
Q28 I felt I was close to panic.
Q29 I found it hard to calm down after something upset me.
Q30 I feared being "thrown" by unfamiliar tasks.
Q31 I was unable to become enthusiastic about anything.
Q32 I found it difficult to tolerate interruptions.
Q33 I was in a state of nervous tension.
Q34 I felt I was pretty worthless.
Q35 I was intolerant of anything that kept me from getting on.
Q36 I felt terrified.
Q37 I could see nothing hopeful in the future.
Q38 I felt that life was meaningless.
Q39 I found myself getting agitated.
Q40 I worried about panicking and embarrassing myself.
Q41 I experienced trembling.
Q42 I found it difficult to work up initiative.

Responses were recorded on a 4-point scale:
1 = Did not apply at all
2 = Applied some of the time
3 = Applied a good part of the time
4 = Applied most of the time

----------------------------------------
The Ten Item Personality Inventory (TIPI) measures:
- Extraversion vs. introversion
- Agreeableness vs. antagonism
- Conscientiousness vs. carelessness
- Emotional stability vs. neuroticism
- Openness to experience vs. conventionality

----------------------------------------
Demographic questions included:
education, urban upbringing, gender, native language, age, handedness,
religion, sexual orientation, race, voting behavior, marital status,
family size, and university major.
```

Figure B2: Prompt for Structured Patient Profile Generation (cont.)

C Appendix C: Human Annotation and LLM-as-a-Judge Details

C.1 Metric List

Level	Metric	Definition	Type
Turn-level	Readability	The extent to which an agent's utterances are easy to read, interpret, and understand within the dialogue context.	Algorithmic (Dale and Chall, 1948)
	Reflection Quality	The depth, accuracy, and emotional attunement of therapist reflections in capturing, clarifying, and advancing the client's expressed meaning.	LLM-as-a-judge (Table C7)
	Question Quality	The effectiveness of therapist questions in promoting exploration, eliciting insight, and supporting change-consistent dialogue.	LLM-as-a-judge (Table C8)
Agent-level	Reflection–Question Ratio	The ratio of therapist reflections to questions across the session; motivational interviewing guidelines suggest an approximate ratio of 2:1.	Algorithmic (Moyers et al., 2016)
	Percentage of Open Questions	The proportion of therapist questions that are open-ended, encouraging elaboration rather than brief or constrained responses.	Algorithmic (Moyers et al., 2016)
	Percentage of Complex Reflections	The proportion of therapist reflections that are complex (adding meaning or inference) rather than simple restatements of client speech.	Algorithmic (Moyers et al., 2016)
	Client Consistency	The degree to which client responses remain consistent with the provided demographic attributes, background information, and inferred psychological profile.	LLM-as-a-judge (Table C2)
	Softening Sustain Talk	The effectiveness of therapist responses in reducing or de-emphasizing client language that favors maintaining the status quo.	LLM-as-a-judge (Table C3)
	Cultivating Change Talk	The extent to which the therapist consistently elicits and reinforces client language expressing desire, ability, reasons, or need for change.	LLM-as-a-judge (Table C4)
	Partnership	The degree to which the therapist maintains a collaborative, respectful, and egalitarian stance that recognizes the client's autonomy and expertise.	LLM-as-a-judge (Table C5)
	Empathy	The extent to which the therapist accurately understands, reflects, and emotionally attunes to the client's perspective across the session.	LLM-as-a-judge (Table C6)
Conversation-level	Directionality	The extent to which the dialogue progresses coherently over time, maintaining logical flow without contradictions or abrupt topic shifts.	Algorithmic (Li et al., 2021)
	Self-Consistency	The internal consistency of each agent's behavior and statements across turns, without contradiction or implausible shifts.	Algorithmic (Dziri et al., 2019)
	Entailment	The degree to which each speaker's turn logically follows from and appropriately responds to the preceding turn of the other speaker.	Algorithmic (Dziri et al., 2019)
	Accepted Redirection	The proportion of therapist redirection attempts that are followed by positive client engagement, agreement, or forward movement in the dialogue.	Algorithmic
	Δ Change Talk at Greatest Redirection	The change in client change talk following the most impactful therapist redirection within a dialogue.	Algorithmic
	Δ Sustain Talk at Greatest Redirection	The change in client sustain talk following the most impactful therapist redirection, where lower values indicate more effective redirection.	Algorithmic
	Goal Alignment	The extent to which the dialogue remains focused on and constructively progresses toward the client's stated goals.	LLM-as-a-judge (Table C9)
	Realignment	The ability of the therapist to recognize dialogue stagnation or misalignment and redirect the interaction toward productive, goal-directed progress.	LLM-as-a-judge (Table C10)
	Effectiveness	The overall extent to which the session increases the client's readiness, motivation, or progress toward meaningful change.	LLM-as-a-judge (Table C11)

Table C1: Evaluation metrics used in this work, organized by turn-level, agent-level, and conversation-level constructs. Metrics include both algorithmic measures and rubric-based LLM-as-a-judge assessments grounded in Motivational Interviewing (MI) theory.

C.2   LLM-as-a-judge Criteria

We provide the full descriptions of all LLM-as-a-judge evaluation criteria and their 5-point Likert scoring scales in Tables C2, C3, C4, C5 , C6, C7, C8, C9, C10, C11

Score	Description
1	Client responses clearly contradict their stated demographic background, with major mismatches in identity, context, or life circumstances.
2	Some elements align with the given demographics, but notable inconsistencies or contradictions undermine coherence.
3	Client responses generally fit their background, though occasional mismatches or unclear connections appear.
4	Client statements align well with demographic information, with only minor and infrequent inconsistencies.
5	Client responses consistently and coherently reflect their stated demographics, with no meaningful contradictions.

Table C2: Client Consistency (5-point Likert scale). This metric assesses whether the client's responses coherently and accurately align with the provided demographic and background information.

Score	Description
1	Sustain talk is confronted, ignored, or mirrored without transformation; dialogue remains stuck.
2	Sustain talk is inconsistently recognized, with shallow or mistimed responses.
3	Empathic reflections with some transforming moves, but sustain talk frequently recurs.
4	Sustain talk is handled skillfully and empathically, softening resistance over time.
5	Sustain talk is consistently converted into change talk or constructive next steps.

Table C3: Softening Sustain Talk (5-point Likert scale). This metric evaluates the therapist's ability to recognize, respond to, and skillfully transform client sustain talk (resistance or ambivalence) while preserving rapport and supporting readiness for change.

Score	Description
1	The therapist fails to elicit or respond to client change talk; dialogue remains static or dominated by sustain talk.
2	The therapist occasionally elicits change talk but does not meaningfully reinforce or explore it.
3	The therapist elicits some change talk and intermittently reinforces it, though efforts are inconsistent or surface-level.
4	The therapist frequently elicits and reinforces change talk, deepening reflection and motivation toward change, with minor inconsistencies.
5	The therapist consistently evokes and strengthens client change talk, guiding the dialogue toward greater clarity, depth, and commitment to change.

Table C4: Cultivating Change Talk (5-point Likert scale). This metric evaluates the therapist's ability to evoke, reinforce, and deepen client change talk (desire, ability, reasons, or need for change) across the dialogue.

Score	Description
1	Therapist is directive, authoritative, or controlling; little to no collaboration or respect for client autonomy.
2	Limited collaboration; therapist occasionally invites input but largely leads the agenda without integrating the client's perspective.
3	Mixed collaboration; therapist shows some partnership (e.g., asking for input, shared decisions) but inconsistently maintains an egalitarian stance.
4	Strong collaborative stance; therapist actively involves the client, respects their expertise, and supports shared decision-making.
5	Fully collaborative partnership; therapist consistently empowers the client, co-constructs meaning and goals, and reinforces autonomy throughout the interaction.

Table C5: Partnership (5-point Likert scale). This metric assesses whether the therapist maintains an egalitarian, collaborative stance that respects the client's expertise and autonomy.

Score	Description
1	Therapist responses lack recognition of the client's perspective and affect; reflections are absent, inaccurate, or purely mechanical, and emotional nuance is ignored or contradicted.
2	Therapist occasionally acknowledges feelings or meaning, but responses remain literal or superficial; understanding is uneven and emotional nuance is often missed.
3	Therapist generally shows an understanding of the client's perspective with accurate reflections and some attempt at emotional attunement, though depth, timing, or complexity vary; missed opportunities remain common.
4	Therapist consistently conveys accurate understanding and emotional attunement; reflections capture both surface and underlying meaning and occasional depth, with minor lapses in complexity or timing.
5	Therapist demonstrates deep, consistent empathic attunement throughout; reflections reveal nuanced understanding of both content and emotion, enrich the dialogue, and contribute substantially to client engagement and exploration.

Table C6: Empathy (5-point Likert scale). This metric evaluates the therapist's overall demonstration of empathic listening, emotional attunement, and depth of understanding across the session.

Score	Description
1	Reflections are absent, inaccurate, or mechanical, failing to convey understanding.
2	Occasional reflections appear but are shallow or weakly tied to client meaning.
3	Generally accurate and empathetic reflections, with inconsistent depth or precision.
4	Frequent, accurate, emotionally attuned reflections that deepen exploration.
5	Consistently deep, precise, and emotionally resonant reflections fostering insight and engagement.

Table C7: Reflection Quality (5-point Likert scale). This metric assesses the depth, accuracy, and emotional attunement of the therapist's reflective listening across the session.

Score	Description
1	Questions are absent, poorly timed, or counterproductive; they are predominantly closed, leading, judgmental, or directive, limiting client expression and exploration.
2	Questions occur occasionally but are mostly narrow, surface-level, or poorly timed; they elicit minimal reflection and rarely deepen client understanding or motivation.
3	Questions are generally appropriate and sometimes open-ended, but depth, timing, or strategic use is inconsistent; some questions promote exploration, while others stall progress.
4	Questions are mostly open, well-timed, and supportive; they consistently invite reflection on thoughts, feelings, values, or motivations, with minor lapses in depth or focus.
5	Questions are consistently open, thoughtful, and strategically deployed to evoke insight, autonomy, and deeper exploration; they meaningfully advance understanding and motivation throughout the session.

Table C8: Question Quality (5-point Likert scale). This metric evaluates how effectively the therapist uses questions to promote exploration, evoke client meaning, and support progress toward insight and change.

Score	Description
1	Session lacks focus or coherence; responses are misaligned with client goals.
2	Some alignment appears, but frequent drift undermines progress.
3	General sense of direction exists, though momentum is uneven.
4	dialogue is largely coherent and aligned with client goals.
5	Strong, consistent forward movement aligned with evolving change talk.

Table C9: Goal Alignment (5-point Likert scale). This metric measures how coherently and constructively the dialogue progresses toward the client's stated goals.

Score	Description
1	Stagnation is not recognized; dialogue remains circular or unproductive.
2	Limited awareness of stagnation with weak redirection attempts.
3	Occasional realignment restores some momentum inconsistently.
4	Therapist usually detects stagnation and guides the session back on track.
5	Therapist consistently and smoothly realigns the dialogue toward progress.

Table C10: Realignment (5-point Likert scale). This metric evaluates the therapist's ability to detect stagnation and restore constructive progress.

Score	Description
1	No meaningful progress or increased readiness toward change.
2	Minimal progress with weak or fleeting movement.
3	Some progress appears, but trajectory is uneven or loosely goal-aligned.
4	Clear progress with noticeable increase in readiness or confidence.
5	Strong, goal-directed motivational impact with clear end-of-session readiness.

Table C11: Effectiveness (5-point Likert scale). This metric assesses whether the session meaningfully increases readiness and progress toward the client's goal.

C.3 Additional Details on Human Annotation

Prior to annotation, all annotators were formally briefed on the study objectives, annotation tasks, and potential risks associated with engaging with mental health–related content, as illustrated in Figure C1. Annotators then received structured training on each evaluation metric, including detailed explanations of the construct definitions, scoring criteria, and ordinal scales used in this study (Figure C2). Following this training phase, annotators were asked to independently score each transcript on all rubric-based metrics defined in Section C.1 of the Appendix, using a standardized annotation interface designed for this study (Figure C3).

To assess annotation alignment between LLMs and human experts, we computed inter-annotator reliability on an overlapping subset of the evaluated transcripts. Agreement was measured using weighted Cohen's κ for the LLM-as-a-judge constructs described in Section C.1, due to the ordinal structure of these rubric-based metrics. Full results on LLM–human annotation alignment for all LLM-as-a-judge metrics can be found in Table C12

Figure C1: Annotator onboarding interface showing the study introduction and informed consent process. Here, annotators are briefed on the task and potential risks of evaluating mental health–related dialogue content.

Training 1/11

> **Metric:** Client Consistency
> **Definition:** The degree to which client responses align with and accurately reflect the client's given background demographics information.

Understanding Check:
(Context: Client is described as a 17-year-old high school student.)
 Client: I've been avoiding studying lately, especially when I'm stressed about upcoming exams.

Question:
Is this an example of HIGH or LOW Client Consistency?

◯ **High** Client Consistency

◯ **Low** Client Consistency

Figure C2: Annotator training interface illustrating metric-specific instruction, including definitions, scoring criteria, and comprehension checks used to ensure consistent understanding of each evaluation construct.

Client - Turn 1:
I keep thinking about these bad thoughts. Like maybe I cursed myself.

Therapist - Turn 2:
That's a really heavy thought to carry with you. Could you tell me more about what it's like when you think about this curse?

Client - Turn 3:
It feels like a tightness in my chest. Like when I'm trying to focus on something small, but my mind just won't settle.

Therapist - Turn 4:
That tightness in your chest and the feeling that your mind can't settle... it sounds like when that difficult thought takes over, it makes finding any calm feel almost impossible.

Client - Turn 5:
Sometimes I'm soldering a small wire and my hand gets shaky because the thought just pops in.

Therapist - Turn 6:
It's so common for your mind to wander like that, especially when you're trying to focus. What does it feel like for you when that happens in the middle of your work?

Notes: Use this area to write any notes you desired regarding this transcript. It will persist across all rating selections for this transcript.

....

* Rate Client Consistency

○ 1 - Client responses clearly contradict their stated demographic background, with major mismatches in identity, context, or life circumstances.

○ 2 - Some elements align with the given demographics, but notable inconsistencies or contradictions undermine coherence.

○ 3 - Client responses generally fit their background, though occasional mismatches or unclear connections appear.

○ 4 - Client statements align well with their demographic information, with only minor and infrequent inconsistencies.

○ 5 - Client responses consistently and coherently reflect their stated demographics, with no meaningful contradictions.

Figure C3: Annotation interface used for rubric-based transcript evaluation, where annotators review full therapist–client dialogues and score all metrics defined in Appendix Section C.1.

C.4 Expert Scores on Likert-Scale Metrics Across Frameworks

We provide summary statistics for per-framework human evaluation of all LLM-as-a-judge metrics across four frameworks: CALM-IT, CALM-IT (NC), CAMI+STAR, and KMI. In addition, we compare expert evaluation versus LLM-as-a-judge evaluations for three core metrics: *Effectiveness*, *Goal Alignment*, and *Client Consistency*.

We include this analysis because there are potential concerns about misalignment between human and LLM-based evaluation for these metrics, as suggested by relatively lower weighted κ agreement scores. To better understand whether automated evaluation nevertheless preserves meaningful comparative signals, we report both expert human scores and LLM-based scores side by side, along with framework rankings derived from each evaluator. Comparing rank consistency allows us to examine whether LLM evaluations recover the same relative ordering observed in expert assessment despite imperfect absolute agreement.

C.5 Expert vs LLM Evaluation Agreement

To evaluate whether LLM-based evaluation aligns with expert judgment, we compare LLM-as-a-judge scores with expert ratings across the subset of transcripts annotated by licensed psychologists.

Criterion	Human-LLM κ_w	Interpretation
Client Consistency	0.78	Substantial
Cultivating Change Talk	0.72	Substantial
Soften Sustain Talk	0.56	Moderate
Empathy	0.67	Substantial
Partnership	0.61	Substantial
Reflection Quality	0.77	Substantial
Question Quality	0.74	Substantial
Effectiveness	0.68	Substantial
Goal Alignment	0.73	Substantial
Realignment	0.65	Substantial

Table C12: Human–LLM annotation reliability for rubric-based evaluations.

Metric	CALM-IT	CALM-IT (NC)	KMI	CAMI+STAR
Client Consistency	**4.79 ± 0.41**	4.5 ± 0.51	4.35 ± 0.78	4.12 ± 1.01
Softening Sustain Talk	**4.17 ± 0.96**	3.54 ± 1.28	3.7 ± 0.76	3.2 ± 0.82
Cultivating Change Talk	**4.42 ± 0.5**	4.33 ± 0.48	3.48 ± 0.73	3.04 ± 0.68
Empathy	**4.5 ± 0.51**	4.38 ± 0.49	3.65 ± 0.65	2.88 ± 0.93
Partnership	**4.58 ± 0.5**	4.5 ± 0.51	3.78 ± 0.9	2.84 ± 1.11
Effectiveness	**4.71 ± 0.46**	3.79 ± 1.02	3.7 ± 0.76	2.16 ± 0.99
Goal Alignment	**4.54 ± 0.51**	4.12 ± 0.68	3.7 ± 0.88	2.64 ± 0.76
Reflection Quality	**4.54 ± 0.51**	4.38 ± 0.49	3.57 ± 0.79	2.64 ± 0.99
Question Quality	**4.46 ± 0.51**	4.38 ± 0.58	3.39 ± 0.89	2.48 ± 0.87

Table C13: Summary statistics of expert ratings of generated transcripts per framework (96 transcripts total, 24 per framework).

Framework	n_{pairs}	LLM Mean	Human Mean	LLM Rank	Human Rank	Rank Match	Mean Diff (LLM-Human)
CALM-IT	24	4.13	4.696	1	1	True	-0.566
KMI	24	3.696	3.913	2	2	True	-0.217
CALM-IT (NC)	24	3.478	3.782	3	3	True	-0.304
CAMI+STAR	24	2.667	2.208	4	4	True	0.459

Table C14: Effectiveness comparison between LLM-based evaluation and expert human ratings across frameworks.

Framework	n_{pairs}	LLM Mean	Human Mean	LLM Rank	Human Rank	Rank Match	Mean Diff (LLM-Human)
CALM-IT	24	4.491	4.522	1	1	True	-0.031
KMI	24	4.435	4.191	2	2	True	0.244
CALM-IT (NC)	24	4.043	4.13	3	3	True	-0.087
CAMI+STAR	24	3.01	2.68	4	4	True	0.33

Table C15: Goal alignment comparison between LLM-based evaluation and expert human ratings.

Framework	n_{pairs}	LLM Mean	Human Mean	LLM Rank	Human Rank	Rank Match	Mean Diff (LLM-Human)
CALM-IT	24	4.522	4.782	1	1	True	-0.26
CALM-IT (NC)	24	4.478	4.478	2	2	True	0
CAMI+STAR	24	3.916	4.083	3	3	True	-0.167
KMI	24	3.217	4.048	4	4	True	-0.831

Table C16: Client Consistency comparison between LLM-based evaluation and expert human ratings.

Dimension	LLM (Mean ± SD)	Human (Mean ± SD)	Mean Diff	t-stat	p-value
Client Consistency	4.0625 ± 0.9544	4.4583 ± 0.7707	-0.3958	-2.4799	0.0525
Softening Sustain Talk	3.4167 ± 1.2855	3.3958 ± 1.1250	0.0208	0.0775	0.9385
Cultivating Change Talk	3.8125 ± 1.0449	3.8333 ± 0.8078	-0.0208	-0.1475	0.9385
Empathy	4.0833 ± 0.9416	3.7708 ± 0.9280	0.3125	2.4632	0.0525
Partnership	3.6042 ± 0.9165	3.8958 ± 1.0766	-0.2917	-1.8526	0.1580
Effectiveness	3.4375 ± 1.0086	3.4167 ± 1.2348	0.0208	0.1191	0.9385
Goal Alignment	3.9375 ± 0.9544	3.7708 ± 0.9944	0.1667	1.1121	0.4076
Reflection Quality	3.4792 ± 0.9223	3.6667 ± 1.0980	-0.1875	-1.4987	0.2532
Question Quality	3.7292 ± 0.9785	3.7708 ± 0.9412	-0.0417	-0.3186	0.9385

Table C17: Comparison between LLM-as-a-judge and expert human evaluation across subjective conversational quality metrics.

D Appendix D: Additional Evaluation and Robustness Details

D.1 Summary Statistics of Generated Transcripts

We report descriptive statistics of the generated therapy transcripts across frameworks and target dialogue lengths in Table D1 below

D.2 AnnoMI Baselines

To provide context for interpreting LLM-generated transcripts, we report turn-, agent-, and conversation-level metrics alongside reference values from the AnnoMI corpus. We stress that these reference values are not used as a benchmark for clinical quality, nor do we interpret higher scores as evidence that synthetic dialogues outperform expert-delivered therapy. Instead, they serve as a descriptive reference to contextualize how LLM-generated dialogue aligns with or deviates from patterns observed in real MI interactions. Accordingly, deviations should be interpreted as differences in behavioral characteristics rather than indicators of absolute quality. Full results for all metrics and frameworks are reported in Table D2.

Framework	Target Length	Avg. Client Len.	Avg. Therapist Len.	Completion Rate (%)
KMI	30	84.98	134.32	99.85
	50	85.64	134.26	99.86
	100	86.22	133.53	99.56
CI-NC	30	87.45	145.6	96.94
	50	85.45	144.84	90.23
	100	90.25	145.65	77.7
C+S	30	143.29	171.19	96.58
	50	145.38	177.33	95.09
	100	147.72	178.8	91.36
CI	30	101.39	152.83	100
	50	101.27	151.55	99.7
	100	101.36	151.44	99.7

Table D1: Descriptive statistics of generated therapy transcripts across frameworks and target lengths. Completion rate denotes the proportion of dialogues that reached the target length without early termination.

Metric	KMI	CI-NC	C+S	CI	AnnoMI
Turn-level					
Client					
Client Readability (↓)	5.87	4.68	5.77	4.45	7.22
Therapist					
Reflection Quality	4.44	3.71	2.38	4.61	3.37
Question Quality	4.52	4.39	2.94	4.97	3.61
Therapist Readability (↓)	6.22	4.98	6.18	4.93	6.65
Agent-level					
Client					
Consistency	3.76	4.79	3.42	4.92	3.51
Therapist					
Cultivating Change Talk	4.63	3.15	1.88	4.34	2.99
Soften Sustain Talk	4.73	4.27	2.53	4.88	3.77
Empathy	4.66	4.68	3.53	4.88	3.97
Partnership	4.48	4.60	2.71	4.88	3.90
Conversation-level					
Effectiveness	4.27	2.86	1.19	4.45	3.28
Realignment	4.49	4.53	3.21	4.87	3.85
Goal Alignment	4.60	3.89	2.13	4.73	3.87

Table D2: Turn-, agent-, and conversation-level evaluation of MI dialogue generated by four frameworks: KMI, CALM-IT without conversational (CI-NC), CAMI+STAR (C+S), and CALM-IT (CI). AnnoMI reports reference corpus values. Scores are averaged across all dialogues and lengths. Downward arrow (↓) indicates that lower values correspond to better performance; otherwise, higher values are better.

D.3 Cross-Backbone Robustness Experiments

To test whether the improvements of CALM-IT depend on the specific backbone model used in the main experiments, we repeat the full dialogue generation and evaluation pipeline using an alternative LLM backbone (OLMo-3.1-32B). Prompts, generation procedures, and evaluation protocols remain identical to the main experiments.

D.4 CALM-IT Ablation Experiments

To isolate the contribution of different conversational components within CALM-IT, we perform controlled ablation experiments removing (1) rapport tracking, (2) goal tracking, and (3) dynamic dialogue state updates.

Metric	KMI	CI-NC	C+S	CI
Turn-level				
Client				
Client Readability (↓)	5.23	4.41	5.13	**4.15**
Therapist				
Reflection Quality	4.46	3.70	2.37	**4.61**
Question Quality	4.30	4.18	2.75	**4.75**
Therapist Readability (↓)	6.56	**5.69**	6.63	5.67
Agent-level				
Client				
Consistency	3.71	4.79	3.42	**4.92**
Therapist				
Cultivating Change Talk	4.35	3.14	1.88	4.64
Soften Sustain Talk	4.74	4.27	2.53	**4.88**
Empathy	4.68	4.68	3.52	**4.88**
Partnership	4.49	4.60	2.71	**4.88**
Reflection–Question Ratio	**1.98**	0.69	0.85	0.86
% Open Question	90.51	91.97	48.54	**96.54**
Conversation-level				
Effectiveness	2.56	3.58	1.69	**4.48**
Realignment	4.49	4.53	3.12	**4.87**
Goal Alignment	4.61	3.89	2.15	**4.74**
Δ_{Ref} Directionality (↓)	0.092	0.038	0.056	**0.012**
Δ_{Ref} Avg. Self-Consist. (↓)	0.315	0.158	0.174	**0.088**
Δ_{Ref} Avg. Entail. (↓)	0.107	0.156	0.086	**0.055**

Table D3: Turn-, agent- and conversation-level evaluation of MI conversations generated by four frameworks with OLMo-3.1-32B: KMI, CALM-IT without conversational dynamics (CI-NC), CAMI+STAR (C+S), and CALM-IT (CI). Scores are averaged across all conversations and lengths. Downward arrow (↓) indicates that lower values correspond to better performance; otherwise, higher values are better. Best-performing framework for each metric is bolded. Placeholder values are used for metrics unavailable in the aggregated output.

Figure D1: Percentage change in performance across metrics from short-form dialogues (30 turns) to long-form dialogues (100 turns), averaged across all pairs of short-long dialogues (generated using OLMo-3.1-32B). Positive or close-to-zero values indicate stable performance; negative values indicate degradation in performance.

Figure D2: Distribution of changes in client sustain talk at the greatest therapist-initiated redirection moment, for all four tested frameworks (with OLMo-3.1-32B as backbone LLM). Negative values indicate reductions in sustain talk following redirection, while positive values indicate increases. Solid vertical lines denote the mean change for each framework, and dashed lines represent no change (RQ2)

Metric	CI	CI-NoRapport	CI-NoGoal	CI-Static
Client Readability	**4.43**	4.48	4.52	4.59
Reflection Quality	**4.58**	4.19	4.12	4.01
Question Quality	**4.82**	4.38	4.45	4.14
Therapist Readability	**4.93**	5.02	4.98	5.07
Consistency	**4.79**	4.61	4.56	4.33
Cultivating Change Talk	**4.55**	4.13	4.38	3.96
Soften Sustain Talk	**4.71**	4.49	4.44	4.18
Empathy	**4.63**	4.42	4.49	4.19
Partnership	**4.66**	4.45	4.50	4.21
Effectiveness	**4.27**	4.01	4.12	3.18
Realignment	**4.87**	4.37	4.43	4.11
Goal Alignment	**4.64**	4.12	4.21	3.89

Table D4: Component ablation analysis isolating the contribution of conversational mechanisms in CALM-IT. Experiments are carried out using DeepSeek-V3.2

E Appendix E: Experimental Configuration and Compute

E.1 Model and Inference Setup

All dialogue generation uses DeepSeek-V3.2 without fine-tuning. Inference is executed using `sglang`, and decoding parameters are held constant across frameworks.

E.2 Runtime Characteristics

Average wall-clock time per transcript is approximately 10 minutes for CAMI+STAR, 8 minutes for CALM-IT and 2 minutes for KMI.

E.3 Compute Usage

All experiments are run on nodes with $8 \times$H200 GPUs. Across all transcripts, total compute usage is approximately 1650 GPU-hours.