Generalization Guarantee of SGD for Pairwise Learning

Abstract

Recently, there is a growing interest in studying pairwise learning since it includes
many important machine learning tasks as specific examples, e.g., metric learning,
AUC maximization and ranking. While stochastic gradient descent (SGD) is an
efficient method, there is a lacking study on its generalization behavior for pairwise
learning. In this paper, we present a systematic study on the generalization analysis
of SGD for pairwise learning to understand the balance between generalization
and optimization. We develop a novel high-probability generalization bound for
uniformly-stable algorithms to incorporate variance information for better gener-
alization, based on which we establish the first *nonsmooth* learning algorithm to
achieve almost optimal *high-probability* and dimension-independent generalization
bounds in linear time. We consider both convex and nonconvex pairwise learning
problems. Our stability analysis for convex problems shows how the interpolation
can help generalization. We establish a uniform convergence of gradients, and apply
it to derive the first generalization bounds on population gradients for nonconvex
problems. Finally, we develop better generalization bounds for gradient-dominated
problems.

1 Introduction

Many machine learning problems can be formulated as learning with pairwise loss functions, where
the performance of the associated models needs to be quantified on a pair of training examples.
Representative problems include AUC maximization [13, 21, 39, 63, 66], metric learning [8, 27]
and ranking [1, 12]. For example, in supervised metric learning we wish to find a distance function
between pairs of examples so that examples within the same class are relatively close while examples
from different classes are far apart from each other. In ranking, we aim to find a function to predict
the ordering of examples. This motivates the recent growing interest in a unifying study of these
problems, under the framework of pairwise learning [28, 36, 58].

Stochastic gradient descent (SGD) is a workhorse for machine learning due to its cheap computation
complexity, simplicity and efficiency [7, 48, 51, 62, 65]. SGD iteratively updates the model based
on stochastic gradients computed on one or several randomly selected training examples, which
can achieve sample-size independent iteration complexity for a prescribed optimization accuracy.
This is especially attractive for pairwise learning as the objective function involves $O(n^2)$ terms for
problems with n training examples. An important problem on SGD is to understand its generalization
performance, i.e., how the models trained by SGD would behave on testing examples. While there are
some interesting work on generalization analysis of SGD for pointwise learning [9, 24, 32, 35], there
is much less work on SGD for pairwise learning. A notable difference between pairwise learning and
pointwise learning is that the objective function in pairwise learning involves $O(n^2)$ *dependent* terms,
which introduces a difficulty in handling this dependency. For example, one needs to decouple this

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

dependency to apply concentration inequalities on independent data. Furthermore, for algorithmic stability analysis, a perturbation of the training dataset by a single example can change $O(n)$ terms in the objective function, which is more challenging than stability analysis of pointwise learning. To our best knowledge, the only work on generalization analysis of SGD for pairwise learning are [36, 54, 59]. However, their analysis requires restrictive assumptions on convexity, smoothness and Lipschitz continuity of loss functions. Furthermore, they fail to incorporate the interpolation (low noise) assumption into their generalization guarantee.

In this paper, we initialize a systematic generalization analysis of SGD for pairwise learning under general assumptions. Our contributions are listed as follows.

1. We develop a novel high-probability generalization bound for uniformly-stable algorithms, which incorporates the variance information to improve the learning performance. We apply this result to develop the first dimension-independent high-probability bound $O(1/\sqrt{n})$ (up to a log factor) for an algorithm with $O(n)$ gradient computation to solve nonsmooth learning problems.

2. We study the stability and generalization guarantee of SGD for pairwise learning with (strongly) convex loss functions, covering both smooth and nonsmooth problems, which gives a novel explanation on how the interpolation (low noise) can help generalization for pairwise learning by using the smoothness assumption. Our stability analysis suggests an appropriate early-stopping strategy for getting excess population risk bounds of the order $O(1/\sqrt{n})$ and $O(1/(n\sigma))$ for convex and σ-strongly convex problems, respectively.

3. We provide the first generalization analysis of SGD for pairwise learning with nonconvex loss functions. We establish a uniform convergence of empirical gradients to population gradients by showing its connection to Rademacher chaos complexities. We then apply this uniform convergence to develop high-probability generalization guarantees for general nonconvex pairwise learning. Under a gradient dominance assumption, our stability analysis gives dimension-independent bounds.

The paper is organized as follows. We survey related work in Section 2 and formulate problems in Section 3. We study convex and nonconvex pairwise learning in Section 4 and Section 5, respectively. Conclusion is given in Section 6.

2 Related Work

We first review the related work on algorithmic stability. Algorithmic stability is an important concept in statistical learning theory (SLT) with close connection to learnability [50, 53]. The modern framework of stability analysis was established in a seminal paper [5], where the important uniform stability was introduced. This stability measure was extended to study randomized algorithms in [16] and motivates several concepts including argument stability [35, 40], Bayes stability [37] and on-average stability [35, 53]. Algorithmic stability has shown its remarkable effectiveness in deriving dimension-independent generalization bounds for various domains including stochastic optimization [9, 24, 32, 45], meta learning [43], structured prediction [41] and differential privacy [3, 47]. Recent progress shows the tradeoff between optimization and stability [10, 54], and its applications to yield almost optimal high-probability generalization bounds [6, 18, 30].

We now review the related work on pairwise learning. There are two popular approaches to studying the generalization performance of pairwise learning: the uniform convergence approach and the algorithmic stability approach. The idea of uniform convergence is to control the uniform deviation between training errors and testing errors over a hypothesis space. The complexity measures of function spaces play an important role in this approach, including VC dimension, covering numbers [15, 58, 60] and Rademacher complexities [8, 28]. Furthermore, one needs to use concentration inequalities to handle the associated U-statistics and U-processes [12, 14]. Algorithmic stability of ranking [1] and metric learning [27, 57] was studied for strongly convex objectives [26]. High-probability bounds of the order $O(\epsilon \log n + 1/\sqrt{n})$ were recently developed for ϵ-uniformly stable pairwise learning algorithms [36]. A nice property of stability analysis is its ability to yield dimension-independent bounds, while a square-root dependency on the dimension is inevitable for uniform convergence analysis when considering general problems [17]. Other than the above two approaches, several researchers have studied the generalization behavior of pairwise learning using the technique of algorithmic robustness [4, 11] and integral operators [23, 67].

3 Problem Formulation

3.1 Pairwise Learning and Stochastic Gradient Descent

Let ρ be a probability measure defined on $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$ with an input space $\mathcal{X}$ and an output space $\mathcal{Y}$. Let $S = \{z_1, \ldots, z_n\}$ be drawn independently according to ρ, from which we aim to learn a prediction function $h : \mathcal{X} \mapsto \mathbb{R}$ or $h : \mathcal{X} \times \mathcal{X} \mapsto \mathbb{R}$. We consider parametric models where the prediction function $h_{\mathbf{w}}$ can be indexed by an element $\mathbf{w} \in \mathcal{W}$, where $\mathcal{W}$ is a d-dimensional Hilbert space. We consider pairwise learning problems where the performance of a model $h_{\mathbf{w}}$ on an example pair (z, z') can be measured by a nonnegative loss function $f(\mathbf{w}; z, z')$. This is in contrast to standard pointwise learning (e.g., classification and regression) where we can measure the quality of a model via its behavior on an individual point. The population risk of $\mathbf{w}$ is $F(\mathbf{w}) = \mathbb{E}_{Z, Z'}[f(\mathbf{w}; Z, Z')]$, where $\mathbb{E}_{Z, Z'}$ denotes the expectation with respect to $Z, Z' \sim \rho$. The empirical risk of $\mathbf{w}$ is defined as

$$F_S(\mathbf{w}) = \frac{1}{n(n-1)} \sum_{i,j \in [n]: i \neq j} f(\mathbf{w}; z_i, z_j),$$

where $[n] := \{1, \ldots, n\}$. Let $\mathbf{w}_S^* = \arg\min_{\mathbf{w} \in \mathcal{W}} F_S(\mathbf{w})$ and $\mathbf{w}^* = \arg\min_{\mathbf{w} \in \mathcal{W}} F(\mathbf{w})$. For a randomized algorithm A, we use $A(S)$ to denote the output model produced by applying A to the dataset S. We are interested in the excess generalization error $F(A(S)) - F(\mathbf{w}^*)$, which measures the relative behavior of $F(A(S))$ as compared to the best model. A standard approach to handle $F(A(S)) - F(\mathbf{w}^*)$ is to use the following error decomposition

$$\mathbb{E}_{S,A}\big[F(A(S)) - F(\mathbf{w}^*)\big] = \mathbb{E}_{S,A}\big[F(A(S)) - F_S(A(S))\big] + \mathbb{E}_{S,A}\big[F_S(A(S)) - F_S(\mathbf{w}^*)\big], \quad (3.1)$$

where the first term $F(A(S)) - F_S(A(S))$ is called the generalization error and the second term $F_S(A(S)) - F_S(\mathbf{w}^*)$ is the optimization error. These two errors can be handled by tools in SLT and optimization theory, respectively. We are interested in the specific SGD for pairwise learning.

Definition 1 (SGD for Pairwise Learning). Let $\mathbf{w}_1 = 0 \in \mathbb{R}^d$ and $\{\eta_t\}_t$ be a stepsize sequence. Let $\nabla f(\mathbf{w}_t; z_{i_t}, z_{j_t})$ denote the gradient of f w.r.t. the first argument. At the t-th iteration, we first draw $\{(i_t, j_t)\}$ from the uniform distribution over all pairs $\{(i, j) : i, j \in [n], i \neq j\}$ and then update

$$\mathbf{w}_{t+1} = \mathbf{w}_t - \eta_t \nabla f(\mathbf{w}_t; z_{i_t}, z_{j_t}), \quad (3.2)$$

where $\nabla f(\mathbf{w}_t; z_{i_t}, z_{j_t})$ denotes the gradient of f w.r.t. the first argument.

An advantage of SGD is that its computational complexity to achieve an accuracy is independent of the number of pairs, which is particularly attractive for pairwise learning. Note gradient descent requires $O(n^2)$ gradient computation per iteration for pairwise learning. We describe $\text{SGD}(S, T, f, \{\eta_t\})$ in Algorithm 1 of SGD with dataset S, iteration number T, loss function f and stepsize $\{\eta_t\}$.

Algorithm 1: $\text{SGD}(S, T, f, \{\eta_t\})$

Input: initial point $\mathbf{w}_1 = 0, \{\eta_t\}_t, S = \{z_1, \ldots, z_n\}$

1 for $t = 1, 2, \ldots, T$ **do**

2   draw $\{(i_t, j_t)\}$ uniformly over all pairs $\{(i, j) : i, j \in [n], i \neq j\}$

3   update $\mathbf{w}_{t+1}$ according to Eq. (3.2)

4 end

Below we introduce necessary definitions and assumptions. Let $\|\cdot\|_2$ be the Euclidean norm and $\langle \cdot, \cdot \rangle$ be the associated inner product. Let $b = \sup_{z,z' \in \mathcal{Z}} f(0; z, z')$ and $b' = \sup_{z,z' \in \mathcal{Z}} \|\nabla f(0; z, z')\|_2$. We denote $B \asymp \widetilde{B}$ if there are absolute constants c_1 and c_2 such that $c_1 B \leq \widetilde{B} \leq c_2 B$.

Definition 2. Let $g : \mathcal{W} \mapsto \mathbb{R}, L, G > 0, \sigma \geq 0$.

1. We say g is L-smooth if g is differentiable and $\|\nabla g(\mathbf{w}) - \nabla g(\mathbf{w}')\|_2 \leq L \|\mathbf{w} - \mathbf{w}'\|_2$ for all $\mathbf{w}, \mathbf{w}' \in \mathcal{W}$.

2. We say g is G-Lipschitz continuous if $|g(\mathbf{w}) - g(\mathbf{w}')| \leq G \|\mathbf{w} - \mathbf{w}'\|_2$ for all $\mathbf{w}, \mathbf{w}' \in \mathcal{W}$.

3. We say g is σ-strongly convex w.r.t. $\|\cdot\|_2$ if $g(\mathbf{w}) - \big(g(\mathbf{w}') + \langle \mathbf{w} - \mathbf{w}', \nabla g(\mathbf{w}') \rangle\big) \geq \sigma \|\mathbf{w} - \mathbf{w}'\|_2^2 / 2$ for all $\mathbf{w}, \mathbf{w}' \in \mathcal{W}$. We say g is convex if g is σ-strongly convex with $\sigma = 0$.

Our analysis may require assumptions on convexity, boundedness of gradients and smoothness. Note the Lipschitz continuity amounts to saying the boundedness of gradients.

Assumption 1 (Convexity). Assume for all $z, z' \in \mathcal{Z}$, the function $\mathbf{w} \mapsto f(\mathbf{w}; z, z')$ is convex.

Assumption 2 (Boundedness of Gradients). Assume for all z, z' and $\mathbf{w} \in \mathcal{W}$, $\|\nabla f(\mathbf{w}; z, z')\|_2 \leq G$.

Assumption 3 (Smoothness). Assume for all $z, z' \in \mathcal{Z}$, the function $\mathbf{w} \mapsto f(\mathbf{w}; z, z')$ is nonnegative and L-smooth.

3.2 Algorithmic Stability and Generalization

A fundamental concept in SLT is the algorithmic stability, which measures the sensitivity of an algorithm w.r.t. the perturbation of the training dataset. Various stability measures have been introduced in the literature, including uniform stability [5], hypothesis stability [5, 16], argument stability [40] and on-average stability [53]. We focus on uniform stability and on-average stability here. The following on-average loss stability was introduced in [36], while the on-average argument stability was motivated by a similar concept in pointwise learning [35]. Let $S = \{z_1, \ldots, z_n\}, S' = \{z_1', \ldots, z_n'\}$ be independently drawn from ρ. We denote

$$S_i = \big\{ z_1, \ldots, z_{i-1}, z_i', z_{i+1}, \ldots, z_n \big\}, \quad \forall i \in [n], \tag{3.3}$$

$$S_{i,j} = \big\{ z_1, \ldots, z_{i-1}, z_i', z_{i+1}, \ldots, z_{j-1}, z_j', z_{j+1}, \ldots, z_n \big\}, \quad \forall i < j \in [n]. \tag{3.4}$$

Definition 3 (Algorithmic Stability). Let $S = \{z_1, \ldots, z_n\}$ and $S' = \{z_1', \ldots, z_n'\}$ be drawn independently from ρ. For any $i, j \in [n]$, denote S_i as (3.3) and $S_{i,j}$ as (3.4).

1. We say a deterministic algorithm $A : \mathcal{Z}^n \mapsto \mathcal{W}$ is ϵ-uniformly stable if for any datasets $S, \widetilde{S} \in \mathcal{Z}^n$ that differ by at most a single example we have $\sup_{z,\tilde{z} \in \mathcal{Z}} |f(A(S); z, \tilde{z}) - f(A(\widetilde{S}); z, \tilde{z})| \leq \epsilon$.

2. We say a randomized algorithm A is on-average (loss) ϵ-stable if

$$\frac{1}{n(n-1)} \sum_{i,j \in [n]: i \neq j} \mathbb{E}_{S,S'} \Big[f\big(A(S_{i,j}); z_i, z_j\big) - f\big(A(S); z_i, z_j\big) \Big] \leq \epsilon.$$

3. We say A is on-average argument ϵ-stable if $\mathbb{E}_{S,\widetilde{S},A}\Big[\frac{1}{n} \sum_{i=1}^{n} \|A(S) - A(S_i)\|_2^2 \Big] \leq \epsilon^2$.

The following theorem establishes the connection between on-average stability and generalization in expectation for pairwise learning. Part (a) was due to [36], while Part (b) was motivated by a similar result in pointwise learning [35]. The proof is given in Appendix A.

Theorem 1. *(a) If A is on-average (loss) ϵ-stable, then $\mathbb{E}[F(S) - F_S(A(S))] \leq \epsilon$.*

(b) If A is on-average argument ϵ-stable and Assumption 3 holds, then for any $\gamma > 0$ we have

$$\mathbb{E}[F(A(S)) - F_S(A(S))] \leq 2(L + \gamma)\epsilon^2 + L\gamma^{-1}\mathbb{E}[F_S(A(S))].$$

Remark 1. We can choose $\gamma \asymp \sqrt{\mathbb{E}[F_S(A(S))]}/\epsilon$ in Part (b) and get the bound $\mathbb{E}[F(A(S)) - F_S(A(S))] = O(\epsilon^2 + \epsilon\sqrt{\mathbb{E}[F_S(A(S))]})$. Therefore, if $\mathbb{E}[F_S(A(S))]$ is small, Part (b) can imply bounds better than $O(\epsilon)$. In particular, if $\mathbb{E}[F_S(A(S))] = O(\epsilon^2)$ the bound in Part (b) becomes $O(\epsilon^2)$, which is much faster than $O(\epsilon)$ in Part (a).

The following theorem establishes the connection between uniform stability and generalization with high probability for pairwise learning. Note we only impose a bounded loss assumption on $A(S)$, which can be achieved by truncating the value of the output function. The proof is given in Section B.

Theorem 2. *Let A be an ϵ-uniformly stable and deterministic algorithm. Let $B = \sup_{z,z'} |\mathbb{E}_S[f(A(S); z, z')] - f(\mathbf{w}^*; z, z')|$ and $\sigma_0^2 = \mathbb{E}_{Z,Z',S}\big[\big(f(A(S); Z, Z') - f(\mathbf{w}^*; Z, Z')\big)^2\big]$. For any $\delta \in (0, 1)$, the following inequality holds with probability at least $1 - \delta$*

$$F(A(S)) - F_S(A(S)) - F(\mathbf{w}^*) + F_S(\mathbf{w}^*) \leq 98\sqrt{2}e\epsilon \log_2 n \log(2e/\delta) + \frac{2B \log(2/\delta)}{3\lfloor n/2 \rfloor} + \sqrt{\frac{2\sigma_0^2 \log(2/\delta)}{\lfloor n/2 \rfloor}}.$$

162 **Remark 2.** Theorem 2 was motivated by the recent work [30, 36]. High-probability generaliza-
163 tion bounds of the order $O(\epsilon \log n)$ were developed for ϵ-uniformly stable and pointwise learn-
164 ing algorithms under a further Bernstein condition [30]. High-probability generalization bounds
165 $O(\epsilon \log n + 1/\sqrt{n})$ were also developed for ϵ-uniformly stable algorithms in pairwise learning [36].
166 We refine these results by developing generalization bounds $O(\epsilon \log n + \sqrt{\sigma_0^2/n})$, where σ_0^2 is the
167 variance of the excess loss at the output model. If this variance is small, then our bounds can be much
168 better than that in [36]. For example, if F is σ-strongly convex, then one can show that this variance
169 can be bounded by $O(\mathbb{E}[F(A(S)) - F(\mathbf{w}^*)]/\sigma)$. As we will show, Theorem 2 can imply almost
170 optimal generalization bounds for an algorithm with $O(n)$ gradient computation to solve nonsmooth
171 problems (Theorem 8), while the existing high-probability bounds can only imply bounds of the
172 order $O(n^{-\frac{1}{4}})$ [36].

4 Pairwise Learning with Convex Loss Functions

174 In this section, we study the generalization performance of SGD for pairwise learning with convex
175 loss functions. We consider convex/strongly-convex and smooth/nonsmooth problems.

4.1 Convex and Smooth Problems

177 We first consider stability and generalization bounds for convex and smooth pairwise learning
178 problems. The proofs of results in this subsection can be found in Section D. Theorem 3 gives
179 the bounds for on-average argument stability of SGD. Note we do not require the loss functions
180 to be Lipschitz continuous, i.e., bounded gradient assumptions. A nice property is that the upper
181 bound involves the empirical risk of $\mathbf{w}_j$. Since we are minimizing the empirical risk by SGD, it is
182 reasonable to assume that $F_S(\mathbf{w}_j)$ would become smaller and smaller along the learning process. It
183 should be mentioned that a similar result was derived for pointwise learning [35]. A key difference
184 in the stability analysis for pairwise learning is that a change of z_i would influence $2(n-1)$ pairs
185 $(z_j, z_i), (z_i, z_j)$ for $j \neq i$. We use the U-structure of empirical risk to prove this result.

186 **Theorem 3** (Stability bound). *Let Assumptions 1, 3 hold. Let A be SGD (Algorithm 1)*
187 *with $\eta_j \leq 2/L$. Then A with t iterations is on-average argument ϵ-stable with $\epsilon^2 \leq$*
188 $\frac{16L(1+2t/n)e}{n} \sum_{j=1}^{t} \eta_j^2 \mathbb{E}[F_S(\mathbf{w}_j)]$.

189 Based on Theorem 3, we get error bounds for pairwise learning with convex and smooth functions.

190 **Theorem 4** (Generalization bound). *Let Assumptions 1, 3 hold. Let $\{\mathbf{w}_t\}$ be the sequence produced*
191 *by SGD (Algorithm 1) with $\eta_t = \eta \leq 2/L$. Then for $\bar{\mathbf{w}}_T = \frac{1}{T} \sum_{t=1}^{T} \mathbf{w}_t$ and any $\gamma \geq 1$ we have the*
192 *following inequality for all $\mathbf{w}$ independent of A (can depend on S)*

$$\mathbb{E}[F(\bar{\mathbf{w}}_T) - F_S(\mathbf{w})] = O\Big(\Big(\eta + \frac{1}{\gamma} + \frac{\gamma(T + T^2/n)\eta^2}{n}\Big)\mathbb{E}[F_S(\mathbf{w})]\Big) + O\Big(\frac{\mathbb{E}[\|\mathbf{w}\|_2^2]}{T\eta} + \frac{\gamma\eta(1 + T/n)\mathbb{E}[\|\mathbf{w}\|_2^2]}{n}\Big).$$
(4.1)

193 A notable property of the above bound is that it holds for any $\mathbf{w}$ independent of A. If $\mathbb{E}[\|\mathbf{w}_S^*\|_2^2]$ is
194 finite, we can choose $\mathbf{w} = \mathbf{w}_S^*$ in Eq. (4.1) and get a bound involving $\mathbb{E}[F_S(\mathbf{w}_S^*)]$. Furthermore,
195 if we are in an interpolation or overparameterized setting [42] then $\mathbb{E}[F_S(\mathbf{w}_S^*)] = o(1/\sqrt{n})$ and
196 the generalization will improve according to (4.1). Therefore, our stability analysis provides a
197 new explanation on how interpolation/overparameterization can help in generalization. Note SGD
198 has an implicit bias to choose a model with a small norm and therefore it is reasonable to assume
199 $\mathbb{E}[\|\mathbf{w}_S^*\|_2^2] < \infty$. We can also choose $\mathbf{w} = \mathbf{w}^*$ in Eq. (4.1) to get optimistic bounds in the sense of
200 involving $F(\mathbf{w}^*)$, which decay fast if $F(\mathbf{w}^*)$ is small [56, 60]. Indeed, the following corollary gives
201 the bound $O(1/\sqrt{n})$ in the general case and improves it to $O(1/n)$ if $F(\mathbf{w}^*) = O(1/n)$.

202 **Corollary 5.** *Let Assumptions in Theorem 4 hold.*

203 *(a) We can choose $\eta_t = \eta \asymp 1/\sqrt{T}$ and $T \asymp n$ to get $\mathbb{E}[F(\bar{\mathbf{w}}_T)] - F(\mathbf{w}^*) = O(1/\sqrt{n})$.*

204 *(b) If $F(\mathbf{w}^*) = O(1/n)$, we can choose $\eta_t = \eta \leq 2/L$ and $T \asymp n$ to get $\mathbb{E}[F(\bar{\mathbf{w}}_T)] - F(\mathbf{w}^*) =$*
205 *$O(1/n)$.*

206 **Remark 3.** Stability and generalization bounds of the order $O(1/\sqrt{n})$ were studied for SGD applied
207 to pairwise learning [36, 54]. However, these discussions require the loss functions to be smooth,

Lipschitz continuous and convex. As a comparison, we remove the Lipschitz continuity assumption. Furthermore, their discussion can only imply non-optimistic bounds of the order $O(1/\sqrt{n})$. As a comparison, our discussions can fully exploit the property of $F(\mathbf{w}^*)$ to imply fast bounds $O(1/n)$.

4.2 Convex and Nonsmooth Problems

We now turn to pairwise learning with convex and nonsmooth functions. Theorem 6 gives the argument stability bounds based on which we develop generalization bounds in Theorem 7. The proofs of results in this subsection are given in Section E.

Theorem 6 (Stability bounds)**.** *Let Assumptions 1, 2 hold. Let $S = \{z_1, \dots, z_n\}$ and $S' = \{z'_1, \dots, z'_n\}$ be two datasets that differ by a single point. Let $\{\mathbf{w}_t\}, \{\mathbf{w}'_t\}$ be the sequence produced by SGD (Algorithm 1) w.r.t. S and S' with $\eta_t = \eta$, respectively. Then*

$$\mathbb{E}\big[\|\mathbf{w}_{t+1} - \mathbf{w}'_{t+1}\|_2^2\big] \leq 4G^2 et\big(1 + 4t/n^2\big)\eta^2. \tag{4.2}$$

For any $\delta \in (0,1)$ with probability at least $1 - \delta$ we have

$$\|\mathbf{w}_{t+1} - \mathbf{w}'_{t+1}\|_2^2 \leq 4G^2\eta^2 e\big(t + \big(2t/n + \log(1/\delta) + \sqrt{4tn^{-1}\log(1/\delta)}\big)^2\big).$$

Theorem 7 (Generalization bounds)**.** *Let Assumptions 1, 2 hold. Let $\{\mathbf{w}_t\}$ be the sequence produced by SGD with $\eta_t = \eta \asymp T^{-\frac{3}{4}}$. If $T \asymp n^2$, then $\mathbb{E}[F(\bar{\mathbf{w}}_T)] - F(\mathbf{w}^*) = O(n^{-\frac{1}{2}})$.*

Remark 4. As compared to the stability bounds in the smooth case (Theorem 3), the stability bounds in (4.2) are worse in the sense that we do not have a factor of $1/n$ in (4.2). Therefore, one needs to choose very small stepsizes to let the right-hand side of (4.2) vanish to 0. Indeed, Theorem 7 suggests $\eta_t \asymp T^{-\frac{3}{4}}$, which are much smaller than the $\eta_t \asymp T^{-\frac{1}{2}}$ in the smooth case. As a result, we require to run SGD with $T \asymp n^2$ to get the optimal generalization bounds $O(1/\sqrt{n})$. Note we also give high-probability bounds on the argument stability in Theorem 6. It should be mentioned that the stability of a variant of SGD as $\mathbf{w}_{t+1} = \mathbf{w}_t - \frac{\eta_t}{t-1}\sum_{k=1}^{t-1}\nabla f(\mathbf{w}_t; z_{i_t}, z_{i_k})$ was recently studied for pairwise learning with convex and nonsmooth loss functions [59]. Note this update requires $O(t)$ gradient computation at the t-th iteration, while (3.2) only requires a single gradient computation.

Algorithm 2: Iterative Localized Algorithm for Pairwise Learning

Input: initial point $\mathbf{w}_0 = 0, \gamma > 0, k = \frac{1}{2}\lceil \log_2 n \rceil$

1 **for** $i = 1, 2, \dots, k$ **do**

2   set $T_i \asymp n_i = \lceil \frac{n}{2^i} \rceil, \gamma_i = \frac{\gamma}{2^i}, \eta_t = \frac{\gamma_i n_i}{t+1}, t \in \mathbb{N}, \tilde{f}(\mathbf{w}; z, z') = f(\mathbf{w}; z, z') + \frac{1}{\gamma_i n_i}\|\mathbf{w} - \mathbf{w}_{i-1}\|_2^2$

3   draw a sample S_i of size n_i independently from ρ

4   apply $\text{SGD}(S_i, T_i, \tilde{f}, \{\eta_t\})$ to minimize the following problem and get $\mathbf{w}_i$

$$\widetilde{F}_{S_i}(\mathbf{w}) := \frac{1}{n_i(n_i - 1)} \sum_{z, z' \in S_i : z \neq z'} f(\mathbf{w}; z, z') + \frac{1}{\gamma_i n_i}\|\mathbf{w} - \mathbf{w}_{i-1}\|_2^2. \tag{4.3}$$

5 **end**

Note SGD requires $O(n^2)$ gradient computation to achieve the generalization bound $O(1/\sqrt{n})$ for nonsmooth problems, which is much larger than the gradient computation $O(n)$ for the smooth case. The following theorem shows that one can also achieve the bound $O(1/\sqrt{n})$ with $O(n)$ gradient computation by considering Algorithm 2. Algorithm 2 is motivated from the iterative localization approach established in pointwise learning [19], which was also used to develop efficient differentially private algorithms [2, 31]. Note the choice $\gamma \asymp n^{-\frac{1}{2}}\|\mathbf{w}^*\|_2$ depends on the unknown $\|\mathbf{w}^*\|_2$. However, one can get the bound $O(D/\sqrt{n})$ by choosing $\gamma \asymp D/\sqrt{n}$ if $\|\mathbf{w}^*\|_2 \leq D$.

Theorem 8. *Let Assumptions 1, 2 hold and $\delta \in (0,1)$. Let $\mathbf{w}_k$ be produced by Algorithm 2 and assume $\sup_{z,z'} |\mathbb{E}_{S_i}[f(\mathbf{w}_i, z, z')]| \leq B$ for some $B > 0$. If we choose $\gamma \asymp n^{-\frac{1}{2}}\|\mathbf{w}^*\|_2$, then with probability at least $1 - \delta$ we have $F(\mathbf{w}_k) - F(\mathbf{w}^*) = O(\log(\log n/\delta)(\log n)\|\mathbf{w}^*\|_2/\sqrt{n})$. Moreover, Algorithm 2 requires only $O(n)$ gradient computations to achieve this generalization bound.*

Remark 5. Iterative localization approach was developed in [19, 31] to develop novel algorithms with $O(n)$ gradient computations and $O(1/\sqrt{n})$ generalization bounds for nonsmooth pointwise learning

problems. We extend this technique to the pairwise learning setting. Furthermore, the generalization bounds in [19, 31] are stated in expectation. As a comparison, we use the novel high-probability bounds for uniformly stable algorithms established in Theorem 2 to develop high-probability bounds of the order $(\log^2 n)/\sqrt{n}$, which has not been developed even for pointwise learning. Note that the sample size $n_k \asymp \sqrt{n}$ in the k-th epoch, and therefore the high-probability bounds in [36] can only yield $O(n^{-\frac{1}{4}})$ bounds. This demonstrates the advantage of our new high-probability bounds in Theorem 2 for developing almost optimal bounds in linear time.

4.3 Strongly Convex Problems

We now turn to strongly convex cases. Theorem 9 gives bounds for smooth problems, while Theorem 10 gives generalization bounds for nonsmooth problems. Note Theorems 9 and 10 apply to any algorithm, which give a general relationship between excess risks and optimization errors. One can plug the optimization error bounds for any algorithm to immediately derive the corresponding excess population risk bounds. The proofs of results in this subsection are given in Section F.

Theorem 9 (Strongly Convex and Smooth Problems). *Let Assumption 3 hold. Assume for all $S \in \mathcal{Z}^n$, F_S is σ-strongly convex w.r.t. $\|\cdot\|_2$. Let A be a randomized algorithm and $\sigma n \geq 8L$. Then*

$$\mathbb{E}[F(A(S)) - F(\mathbf{w}^*)] \leq 128L\Big(\frac{16L}{n^2\sigma^2} + \frac{1}{n\sigma}\Big)\mathbb{E}\big[F_S(\mathbf{w}_S^*)\big] + \frac{2L}{\sigma}\mathbb{E}\big[F_S(A(S)) - F_S(\mathbf{w}_S^*)\big].$$

Remark 6. The above bound involves two components. The first component $O(1/(n\sigma))$ depends only on the landscape of the learning problem. The second component involves optimization errors. It shows that optimization is always beneficial to improve generalization for strongly convex problems. Furthermore, it also shows that one can stop the algorithm once we achieve the optimization error bounds $O(1/n)$ since further optimization would not improve essentially generalization. Theorem 9 also implies bounds of the order $o(1/(n\sigma))$ if $F_S(\mathbf{w}^*)$ is small.

Theorem 10 (Strongly Convex and Nonsmooth Problems). *Assume f takes a structure as $f(\mathbf{w}; z, z') = \ell(\mathbf{w}; z, z') + r(\mathbf{w})$. Assume for all z, z', the map $\mathbf{w} \mapsto \ell(\mathbf{w}; z, z')$ is G-Lipschitz. Assume for all $S \in \mathcal{Z}^n$, F_S is σ-strongly convex w.r.t. $\|\cdot\|_2$. For any algorithm A we have*

$$\mathbb{E}[F(A(S)) - F(\mathbf{w}^*)] \leq \frac{8G^2}{n\sigma} + G\sqrt{\frac{2\mathbb{E}\big[F_S(A(S) - F_S(\mathbf{w}_S^*))\big]}{\sigma}}.$$

We present the specific applications of the above results to SGD in Corollary F.2 (Section F).

5 Pairwise Learning with Nonconvex Loss Functions

In this section, we consider generalization bounds for pairwise learning in a nonconvex setting. In this case, the excess population risk is not a reasonable measure since we cannot guarantee that the algorithm can find a global minimizer. We therefore use the norm of gradients at $A(S)$ to measure the performance of A [22, 69]. In a general nonconvex setting, SGD requires to choose $\eta_t = O(1/t)$ for a meaningful stability bound [24], for which the optimization errors would decay logarithmically w.r.t the number of iterations [22]. Then, one cannot trade-off the stability and optimization for a model with good generalization performance in a general nonconvex problem. Therefore, we turn to a different uniform convergence approach for generalization bounds in a general nonconvex setting. After that, we study stability of SGD for nonconvex problems under a further PL condition..

5.1 Uniform Convergence of Gradients for Pairwise Learning

Our first result for nonconvex pairwise learning is a uniform convergence of gradients. Specifically, we show that the uniform deviation between population gradients and empirical gradients over a space can be bounded by the associated Rademacher chaos complexity. Let $\mathcal{W}_R = \{\mathbf{w} \in \mathcal{W} : \|\mathbf{w}\|_2 \leq R\}$ for $R > 0$. The proofs of results in this subsection are given in Section G.

Definition 4. Let $\mathcal{F} := \{f : \mathcal{Z}^4 \mapsto \mathbb{R}\}$ be a function class and $S = \{z_i\}_{i=1}^n \subset \mathcal{Z}$. Let $\{\epsilon_i\}_{i=1}^{\lfloor\frac{n}{2}\rfloor}$ be independent Rademacher variables with $\Pr\{\epsilon_i = 1\} = \Pr\{\epsilon_i = -1\} = 1/2$. The empirical Rademacher chaos complexity for $\mathcal{F}$ w.r.t. S is defined as

$$\mathcal{U}_S(\mathcal{F}) = \frac{1}{\lfloor\frac{n}{2}\rfloor}\mathbb{E}_\epsilon\Big[\sup_{f \in \mathcal{F}} \sum_{1 \leq i < j \leq \lfloor\frac{n}{2}\rfloor} \epsilon_i\epsilon_j f(z_i, z_{i+\lfloor\frac{n}{2}\rfloor}, z_j, z_{j+\lfloor\frac{n}{2}\rfloor})\Big].$$

Theorem 11 (Uniform Convergence of Gradients). *Let $\delta \in (0,1), R > 0$ and $S = \{z_1, \ldots, z_n\}$ be drawn independently from ρ. If Assumption 3 holds, then with probability at least $1 - \delta$ we have*

$$\sup_{\mathbf{w} \in \mathcal{W}_R} \left\| \nabla F(\mathbf{w}) - \nabla F_S(\mathbf{w}) \right\|_2 \leq \frac{2\sqrt{2}(LR + b')\left(2 + \sqrt{\log(1/\delta)}\right)}{\sqrt{n}} + 4\sqrt{\frac{\mathcal{U}_S(\mathcal{F}_R)}{n}},$$

where

$$\mathcal{F}_R = \left\{ (z_1, z_2, z_3, z_4) \mapsto \langle \nabla f(\mathbf{w}; z_1, z_2), \nabla f(\mathbf{w}; z_3, z_4) \rangle : \mathbf{w} \in \mathcal{W}_R \right\}.$$

We can apply the entropy integral to control the above Rademacher chaos complexity [14, 61], and get the following result. Note d is the dimension of the space $\mathcal{W}$.

Corollary 12. *Under Assumptions of Theorem 11, with probability at least $1 - \delta$ we have*

$$\sup_{\mathbf{w} \in \mathcal{W}_R} \left\| \nabla F(\mathbf{w}) - \nabla F_S(\mathbf{w}) \right\|_2 \leq \frac{2\sqrt{2}(LR + b')}{\sqrt{n}} \left(2 + \sqrt{96e\left(\log 2 + d\log(3e)\right)} + \sqrt{\log(1/\delta)} \right).$$

The above uniform convergence rate involves a square-root dependency on d. We show that this dependency can be avoided if we consider a specific class of functions with a specific structure

$$f(\mathbf{w}; z, z') = \psi(\langle \mathbf{w}, \phi(x, x') \rangle, \tau(y, y')), \tag{5.1}$$

where $\phi : \mathcal{X} \times \mathcal{X} \mapsto \mathcal{W}$ is a feature map, $\psi : \mathbb{R} \times \mathbb{R} \mapsto \mathbb{R}$ is L_ψ-smooth w.r.t. the first argument and $\tau : \mathcal{Y} \times \mathcal{Y} \mapsto \mathbb{R}$. Loss functions of the structure (5.1) have wide applications in robust optimization and generalized linear models [20, 44]. We assume $\kappa = \sup_{x,x' \in \mathcal{X}} \|\phi(x, x')\|_2$.

Corollary 13. *Let $\delta \in (0,1), R > 0$ and $S = \{z_1, \ldots, z_n\}$ be examples drawn independently from ρ. Suppose $f : \mathcal{W} \times \mathcal{Z}^2 \mapsto \mathbb{R}$ takes the form (5.1) with ψ being L_ψ-smooth w.r.t. the first argument. Then the following inequality holds with probability at least $1 - \delta$*

$$\sup_{\mathbf{w} \in \mathcal{W}_R} \left\| \nabla F(\mathbf{w}) - \nabla F_S(\mathbf{w}) \right\|_2 \leq \frac{4\kappa\left(2L_\psi R\kappa + b'\right)}{\sqrt{n}} + \sqrt{\frac{8\left(L_\psi \kappa^2 R + b'\right)^2 \log(1/\delta)}{n}}.$$

Remark 7. Uniform convergence of gradients was studied for pointwise learning based on covering numbers [44, 64] and Rademacher complexities [20]. We extend these results to the pairwise learning setting. A key difference between pointwise learning and pairwise learning is that the empirical gradients for pairwise learning can be no longer written as a summation of i.i.d. terms. Indeed, the $n(n-1)$ terms in $\nabla F_S(\mathbf{w})$ are correlated, which introduces difficulties in applying concentration inequalities. We need to apply decoupling techniques in U-process to handle this correlation.

5.2 Smooth Problems

We now study the generalization performance of SGD for pairwise learning based on the uniform convergence of gradients developed in the previous subsection. We first introduce necessary assumptions. Since η_t is always small (a typical choice is $\eta_t \asymp 1/\sqrt{T}$), Eq. (5.2) is milder than a bounded gradient assumption. Eq. (5.3) imposes a bounded variance assumption on stochastic gradients, which is a standard assumption for the analysis of SGD [22, 32, 68].

Assumption 4. Assume the existence of $G > 0$ and $\sigma > 0$ such that

$$\sqrt{\eta_t} \|\nabla f(\mathbf{w}_t; z, z')\|_2 \leq G, \forall t \in \mathbb{N}, z, z' \in \mathcal{Z}, \tag{5.2}$$

$$\mathbb{E}_{i_t, j_t} \left[\|\nabla f(\mathbf{w}_t; z_{i_t}, z_{j_t}) - \nabla F_S(\mathbf{w}_t)\|_2^2 \right] \leq \sigma^2, \quad \forall t \in \mathbb{N}. \tag{5.3}$$

Theorem 14 gives high-probability bounds on the norm of population gradients. Our basic idea is to use the following error decomposition

$$\|\nabla F(\mathbf{w}_t)\|_2^2 \leq 2\|\nabla F(\mathbf{w}_t) - \nabla F_S(\mathbf{w}_t)\|_2^2 + 2\|\nabla F_S(\mathbf{w}_t)\|_2^2.$$

We refer to the first term on the right-hand side as the generalization error for nonconvex pairwise learning, which can be bounded by the uniform convergence of gradients established in Theorem 11. The second term is the optimization error and can be addressed by techniques in optimization theory. The proof is given in Section H.

Theorem 14 (Smooth Problems). *Let Assumptions 3 and 4 hold. Let $\{\mathbf{w}_t\}_t$ be the sequence produced by (3.2) with $\eta_t = \eta/\sqrt{T}$ and $\eta \le \sqrt{T}/(2L)$. Then for any $\delta \in (0,1)$, we can choose $T \asymp nd^{-1}$ to derive the following inequality with probability at least $1 - \delta$*

$$\frac{1}{T}\sum_{t=1}^{T} \|\nabla F(\mathbf{w}_t)\|_2^2 = O\Big(n^{-\frac{1}{2}}\log^2(1/\delta)\big(d + \log(1/\delta)\big)^{\frac{1}{2}}\Big). \tag{5.4}$$

Furthermore, if f takes the specific structure (5.1) we can choose $T \asymp n$ to derive the following inequality with probability at least $1 - \delta$

$$\frac{1}{T}\sum_{t=1}^{T} \|\nabla F(\mathbf{w}_t)\|_2^2 = O\Big(n^{-\frac{1}{2}}\log^{\frac{5}{2}}(1/\delta)\Big). \tag{5.5}$$

Remark 8. It is clear that the bound in (5.4) is dimension-dependent, which is due to the consideration of uniform convergence. Eq. (5.5) further shows that this dependency on dimension can be avoided for problems of specific structure. Note the optimization errors of SGD with T iterations for nonconvex problems satisfy $\frac{1}{T}\sum_{t=1}^{T}\|\nabla F_S(\mathbf{w}_t)\|_2^2 = O(T^{-\frac{1}{2}})$ [22]. This is consistent with the generalization error bounds $\frac{1}{T}\sum_{t=1}^{T}\|\nabla F(\mathbf{w}_t)\|_2^2 = O\big(n^{-\frac{1}{2}}\log^{\frac{5}{2}}(1/\delta)\big)$ in (5.5) by noting $T \asymp n$. Therefore, our analysis shows that the extension from optimization to generalization comes with no cost.

5.3 Gradient Dominated Problems

We now study the stability and generalization of SGD for pairwise learning with gradient-dominated objectives (or PL condition). PL condition is widely used in nonconvex learning [20, 29], and was shown to hold true for deep (linear) and shallow neural networks [9, 38]. Intuitively speaking, PL condition means that the suboptimality measured by function values can be bounded by gradients.

Assumption 5 (Polyak-Lojasiewicz Condition). Denote $\hat{F}_S = \inf_{\mathbf{w}' \in \mathcal{W}} F_S(\mathbf{w}')$. We assume F_S satisfies PL or gradient-dominated condition (in expectation) with parameter $\beta > 0$, i.e.,

$$\mathbb{E}_S\big[F_S(\mathbf{w}) - \hat{F}_S\big] \le \frac{1}{2\beta}\mathbb{E}_S\big[\|\nabla F_S(\mathbf{w})\|_2^2\big], \quad \forall \mathbf{w} \in \mathcal{W}. \tag{5.6}$$

Under the PL condition, we can get excess population risk bounds based on the stability analysis. The proof of Theorem 15 is given in Section I.

Theorem 15 (Gradient Dominated Problems). *Let Assumptions 2, 3, 5 hold. Assume $|f(\mathbf{w};z,z')| \le B$ for all $\mathbf{w} \in \mathcal{W}, z, z' \in \mathcal{Z}$. Let $\{\mathbf{w}_t\}_t$ be the sequence produced by (3.2) with $\eta_t = \frac{2t+1}{2\beta(t+1)^2}$. Then*

$$\mathbb{E}\big[F(\mathbf{w}_T)\big] - F(\mathbf{w}^*) = O\Big(\frac{T^{\frac{L}{L+\beta}}}{n}\Big) + O\big(1/(T\beta^2)\big). \tag{5.7}$$

We can choose $T \asymp n^{\frac{1+L/\beta}{1+2L/\beta}}\beta^{-\frac{2+2L/\beta}{1+2L/\beta}}$ to get $\mathbb{E}\big[F(\mathbf{w}_T)\big] - F(\mathbf{w}^*) = O\big(n^{-\frac{1+L/\beta}{1+2L/\beta}}\beta^{-\frac{2L/\beta}{1+2L/\beta}}\big)$.

Remark 9. Note the above bounds depend on the condition number $\text{cond} := L/\beta$. If $\text{cond} \approx 1$, then we get $\mathbb{E}\big[F(\mathbf{w}_T)\big] - F(\mathbf{w}^*) \approx O\big(n^{-\frac{2}{3}}\big)$. As cond increases, the bound increases to $O\big(n^{-\frac{1}{2}}\big)$.

6 Conclusion

In this paper, we present a systematic study on the generalization performance for pairwise learning. We develop novel high-probability bounds for uniformly stable algorithms, and apply them to develop algorithms with optimal bounds in linear time for nonsmooth problems. We conduct the stability analysis for various convex problems including smooth, nonsmooth and strongly convex objectives, and get optimal excess population risk bounds of order $O(1/\sqrt{n})$ for convex problems and $O(1/(n\sigma))$ for σ-strongly convex problems, respectively. We conduct the uniform convergence analysis for general nonconvex problems, which imply the bounds of the order $O(1/\sqrt{n})$ for population gradients. We further study the stability and generalization for nonconvex pairwise learning with gradient dominated objectives. Our discussions can clarify the effect of interpolation on generalization.

It would be interesting to study the stability of SGD in a general nonconvex case for getting dimension-independent bounds. It would also be very interesting to study other stochastic optimization methods for pairwise learning, including variance reduction variants and momentum technique.

References

[1] S. Agarwal and P. Niyogi. Generalization bounds for ranking algorithms via algorithmic stability. *Journal of Machine Learning Research*, 10(Feb):441–474, 2009.

[2] H. Asi, V. Feldman, T. Koren, and K. Talwar. Private stochastic convex optimization: Optimal rates in ℓ_1 geometry. *arXiv preprint arXiv:2103.01516*, 2021.

[3] R. Bassily, V. Feldman, C. Guzmán, and K. Talwar. Stability of stochastic gradient descent on nonsmooth convex losses. *Advances in Neural Information Processing Systems*, 33, 2020.

[4] A. Bellet and A. Habrard. Robustness and generalization for metric learning. *Neurocomputing*, 151:259–267, 2015.

[5] O. Bousquet and A. Elisseeff. Stability and generalization. *Journal of Machine Learning Research*, 2:499–526, 2002.

[6] O. Bousquet, Y. Klochkov, and N. Zhivotovskiy. Sharper bounds for uniformly stable algorithms. In *Conference on Learning Theory*, pages 610–626, 2020.

[7] R. Camoriano, T. Angles, A. Rudi, and L. Rosasco. Nytro: When subsampling meets early stopping. In *Artificial Intelligence and Statistics*, pages 1403–1411. PMLR, 2016.

[8] Q. Cao, Z.-C. Guo, and Y. Ying. Generalization bounds for metric and similarity learning. *Machine Learning*, 102(1):115–132, 2016.

[9] Z. Charles and D. Papailiopoulos. Stability and generalization of learning algorithms that converge to global optima. In *International Conference on Machine Learning*, pages 744–753, 2018.

[10] Y. Chen, C. Jin, and B. Yu. Stability and convergence trade-off of iterative optimization algorithms. *arXiv preprint arXiv:1804.01619*, 2018.

[11] A. Christmann and D.-X. Zhou. On the robustness of regularized pairwise learning methods based on kernels. *Journal of Complexity*, 37:1–33, 2016.

[12] S. Clémençon, G. Lugosi, and N. Vayatis. Ranking and empirical minimization of U-statistics. *The Annals of Statistics*, pages 844–874, 2008.

[13] C. Cortes and M. Mohri. AUC optimization vs. error rate minimization. In *Advances in Neural Information Processing Systems*, pages 313–320, 2004.

[14] V. De la Pena and E. Giné. *Decoupling: from dependence to independence*. Springer Science & Business Media, 2012.

[15] M. Dong, X. Yang, R. Zhu, Y. Wang, and J. Xue. Generalization bound of gradient descent for non-convex metric learning. In *Advances in Neural Information Processing Systems*, 2020.

[16] A. Elisseeff, T. Evgeniou, and M. Pontil. Stability of randomized learning algorithms. *Journal of Machine Learning Research*, 6(Jan):55–79, 2005.

[17] V. Feldman. Generalization of erm in stochastic convex optimization: The dimension strikes back. In *Advances in Neural Information Processing Systems*, pages 3576–3584, 2016.

[18] V. Feldman and J. Vondrak. High probability generalization bounds for uniformly stable algorithms with nearly optimal rate. In *Conference on Learning Theory*, pages 1270–1279, 2019.

[19] V. Feldman, T. Koren, and K. Talwar. Private stochastic convex optimization: optimal rates in linear time. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, pages 439–449, 2020.

[20] D. J. Foster, A. Sekhari, and K. Sridharan. Uniform convergence of gradients for non-convex learning and optimization. In *Advances in Neural Information Processing Systems*, pages 8759–8770, 2018.

[21] W. Gao, R. Jin, S. Zhu, and Z.-H. Zhou. One-pass auc optimization. In *International Conference on Machine Learning*, pages 906–914, 2013.

[22] S. Ghadimi and G. Lan. Stochastic first-and zeroth-order methods for nonconvex stochastic programming. *SIAM Journal on Optimization*, 23(4):2341–2368, 2013.

[23] X. Guo, T. Hu, and Q. Wu. Distributed minimum error entropy algorithms. *Journal of Machine Learning Research*, 21(126):1–31, 2020.

[24] M. Hardt, B. Recht, and Y. Singer. Train faster, generalize better: Stability of stochastic gradient descent. In *International Conference on Machine Learning*, pages 1225–1234, 2016.

[25] N. J. Harvey, C. Liaw, and S. Randhawa. Simple and optimal high-probability bounds for strongly-convex stochastic gradient descent. *arXiv preprint arXiv:1909.00843*, 2019.

[26] M. Huai, D. Wang, C. Miao, J. Xu, and A. Zhang. Pairwise learning with differential privacy guarantees. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 34, pages 694–701, 2020.

[27] R. Jin, S. Wang, and Y. Zhou. Regularized distance metric learning: Theory and algorithm. In *Advances in Neural Information Processing Systems*, pages 862–870, 2009.

[28] P. Kar, B. Sriperumbudur, P. Jain, and H. Karnick. On the generalization ability of online learning algorithms for pairwise loss functions. In *International Conference on Machine Learning*, pages 441–449, 2013.

[29] H. Karimi, J. Nutini, and M. Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition. In *European Conference on Machine Learning*, pages 795–811, 2016.

[30] Y. Klochkov and N. Zhivotovskiy. Stability and deviation optimal risk bounds with convergence rate O(1/n). *arXiv preprint arXiv:2103.12024*, 2021.

[31] J. Kulkarni, Y. T. Lee, and D. Liu. Private non-smooth empirical risk minimization and stochastic convex optimization in subquadratic steps. *arXiv preprint arXiv:2103.15352*, 2021.

[32] I. Kuzborskij and C. Lampert. Data-dependent stability of stochastic gradient descent. In *International Conference on Machine Learning*, pages 2820–2829, 2018.

[33] S. Lacoste-Julien, M. Schmidt, and F. Bach. A simpler approach to obtaining an O(1/t) convergence rate for the projected stochastic subgradient method. *arXiv preprint arXiv:1212.2002*, 2012.

[34] Y. Lei and K. Tang. Learning rates for stochastic gradient descent with nonconvex objectives. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 2021.

[35] Y. Lei and Y. Ying. Fine-grained analysis of stability and generalization for stochastic gradient descent. In *International Conference on Machine Learning*, pages 5809–5819, 2020.

[36] Y. Lei, A. Ledent, and M. Kloft. Sharper generalization bounds for pairwise learning. In *Advances in Neural Information Processing Systems*, volume 33, pages 21236–21246, 2020.

[37] J. Li, X. Luo, and M. Qiao. On generalization error bounds of noisy gradient methods for non-convex learning. In *International Conference on Learning Representations*, 2020.

[38] Y. Li and Y. Yuan. Convergence analysis of two-layer neural networks with relu activation. In *Advances in Neural Information Processing Systems*, pages 597–607, 2017.

[39] M. Liu, X. Zhang, Z. Chen, X. Wang, and T. Yang. Fast stochastic AUC maximization with O(1/n)-convergence rate. In *International Conference on Machine Learning*, pages 3195–3203, 2018.

[40] T. Liu, G. Lugosi, G. Neu, and D. Tao. Algorithmic stability and hypothesis complexity. In *International Conference on Machine Learning*, pages 2159–2167, 2017.

[41] B. London, B. Huang, and L. Getoor. Stability and generalization in structured prediction. *The Journal of Machine Learning Research*, 17(1):7808–7859, 2016.

[42] S. Ma, R. Bassily, and M. Belkin. The power of interpolation: Understanding the effectiveness of sgd in modern over-parametrized learning. In *International Conference on Machine Learning*, pages 3325–3334, 2018.

[43] A. Maurer. Algorithmic stability and meta-learning. *Journal of Machine Learning Research*, 6 (Jun):967–994, 2005.

[44] S. Mei, Y. Bai, and A. Montanari. The landscape of empirical risk for nonconvex losses. *The Annals of Statistics*, 46(6A):2747–2774, 2018.

[45] W. Mou, L. Wang, X. Zhai, and K. Zheng. Generalization bounds of sgld for non-convex learning: Two theoretical viewpoints. In *Conference on Learning Theory*, pages 605–638, 2018.

[46] A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on Optimization*, 19(4):1574–1609, 2009.

[47] L. Oneto, M. Donini, M. Pontil, and J. Shawe-Taylor. Randomized learning and generalization of fair and private classifiers: From pac-bayes to stability and differential privacy. *Neurocomputing*, 416:231–243, 2020.

[48] F. Orabona. A modern introduction to online learning. *arXiv preprint arXiv:1912.13213*, 2019.

[49] G. Pisier. *The Volume of Convex Bodies and Banach Space Geometry*, volume 94. Cambridge University Press, 1999.

[50] A. Rakhlin, S. Mukherjee, and T. Poggio. Stability results in learning theory. *Analysis and Applications*, 3(04):397–417, 2005.

[51] N. L. Roux, M. Schmidt, and F. Bach. A stochastic gradient method with an exponential convergence rate for finite training sets. In *Advances in Neural Information Processing Systems*, pages 2663–2671, 2012.

[52] S. Shalev-Shwartz and S. Ben-David. *Understanding machine learning: From theory to algorithms*. Cambridge university press, 2014.

[53] S. Shalev-Shwartz, O. Shamir, N. Srebro, and K. Sridharan. Learnability, stability and uniform convergence. *Journal of Machine Learning Research*, 11:2635–2670, 2010.

[54] W. Shen, Z. Yang, Y. Ying, and X. Yuan. Stability and optimization error of stochastic gradient descent for pairwise learning. *Analysis and Applications*, pages 1–41, 2019.

[55] D. Slepian. The one-sided barrier problem for gaussian noise. *Bell System Technical Journal*, 41(2):463–501, 1962.

[56] N. Srebro, K. Sridharan, and A. Tewari. Smoothness, low noise and fast rates. In *Advances in Neural Information Processing Systems*, pages 2199–2207, 2010.

[57] B. Wang, H. Zhang, P. Liu, Z. Shen, and J. Pineau. Multitask metric learning: Theory and algorithm. In *International Conference on Artificial Intelligence and Statistics*, pages 3362–3371, 2019.

[58] Y. Wang, R. Khardon, D. Pechyony, and R. Jones. Generalization bounds for online learning algorithms with pairwise loss functions. In *Conference on Learning Theory*, volume 23, pages 13–1, 2012.

[59] Z. Yang, Y. Lei, S. Lyu, and Y. Ying. Stability and differential privacy of stochastic gradient descent for pairwise learning with non-smooth loss. In *International Conference on Artificial Intelligence and Statistics*, pages 2026–2034. PMLR, 2021.

[60] H.-J. Ye, D.-C. Zhan, and Y. Jiang. Fast generalization rates for distance metric learning. *Machine Learning*, 108(2):267–295, 2019.

[61] Y. Ying and C. Campbell. Rademacher chaos complexities for learning the kernel problem. *Neural computation*, 22(11):2858–2886, 2010.

[62] Y. Ying and D.-X. Zhou. Online regularized classification algorithms. *IEEE Transactions on Information Theory*, 52(11):4775–4788, 2006.

[63] Y. Ying, L. Wen, and S. Lyu. Stochastic online AUC maximization. In *Advances in Neural Information Processing Systems*, pages 451–459, 2016.

[64] L. Zhang, T. Yang, and R. Jin. Empirical risk minimization for stochastic convex optimization: $O(1/n)$-and $O(1/n^2)$-type of risk bounds. In *Conference on Learning Theory*, pages 1954–1979, 2017.

[65] T. Zhang. Solving large scale linear prediction problems using stochastic gradient descent algorithms. In *International Conference on Machine Learning*, pages 919–926, 2004.

[66] P. Zhao, S. C. Hoi, R. Jin, and T. Yang. Online AUC maximization. In *International Conference on Machine Learning*, pages 233–240, 2011.

[67] Y. Zhou, H. Chen, R. Lan, and Z. Pan. Generalization performance of regularized ranking with multiscale kernels. *IEEE Transactions on Neural Networks and Learning Systems*, 27(5): 993–1002, 2015.

[68] Y. Zhou, Y. Liang, and H. Zhang. Generalization error bounds with probabilistic guarantee for SGD in nonconvex optimization. *arXiv preprint arXiv:1802.06903*, 2018.

[69] Y. Zhou, B. Karimi, J. Yu, Z. Xu, and P. Li. Towards better generalization of adaptive gradient methods. *Advances in Neural Information Processing Systems*, 33, 2020.