Dynamics of SGD with Stochastic Polyak Stepsizes: Truly Adaptive Variants and Convergence to Exact Solution

Abstract

Recently Loizou et al. (2021), proposed and analyzed stochastic gradient descent (SGD) with stochastic Polyak stepsize (SPS). The proposed SPS comes with strong convergence guarantees and competitive performance; however, it has two main drawbacks when it is used in non-over-parameterized regimes: (i) It requires a priori knowledge of the optimal mini-batch losses, which are not available when the interpolation condition is not satisfied (e.g., regularized objectives), and (ii) it guarantees convergence only to a neighborhood of the solution. In this work, we study the dynamics and the convergence properties of SGD equipped with new variants of the stochastic Polyak stepsize and provide solutions to both drawbacks of the original SPS. We first show that a simple modification of the original SPS that uses lower bounds instead of the optimal function values can directly solve issue (i). On the other hand, solving issue (ii) turns out to be more challenging and leads us to valuable insights into the method's behavior. We show that if interpolation is not satisfied, the correlation between SPS and stochastic gradients introduces a bias, which effectively distorts the expectation of the gradient signal near minimizers, leading to non-convergence - even if the stepsize is scaled down during training. To fix this issue, we propose DecSPS, a novel modification of SPS, which guarantees convergence to the exact minimizer - without a priori knowledge of the problem parameters. For strongly-convex optimization problems, DecSPS is the first stochastic adaptive optimization method that converges to the exact solution without restrictive assumptions like bounded iterates/gradients.

1   Introduction

We consider the stochastic optimization problem:

$$\min_{x \in \mathbb{R}^d} \left[f(x) = \frac{1}{n} \sum_{i=1}^{n} f_i(x) \right], \tag{1}$$

where each f_i is convex and lower bounded. We denote by $\mathcal{X}^*$ the non-empty set of optimal points x^* of equation (1). We set $f^* := \min_{x \in \mathbb{R}^d} f(x)$, and $f_i^* := \inf_{x \in \mathbb{R}^d} f_i(x)$.

In this setting, the algorithm of choice is often Stochastic Gradient Descent (SGD), i.e. $x^{k+1} = x^k - \gamma_k \nabla f_{\mathcal{S}_k}(x^k)$, where $\gamma_k > 0$ is the stepsize at iteration k, $\mathcal{S}_k \subseteq [n]$ a random subset of datapoints (minibatch) with cardinality B sampled independently at each iteration k, and $\nabla f_{\mathcal{S}_k}(x) := \frac{1}{B} \sum_{i \in \mathcal{S}_k} \nabla f_i(x)$ is the minibatch gradient.

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

A careful choice of γ_k is crucial for most applications [4, 14]. The simplest option is to pick γ_k to be constant over training, with its value inversely proportional to the Lipschitz constant of the gradient. While this choice yields fast convergence to the neighborhood of a minimizer, two main problems arise: (a) the optimal γ depends on (often unknown) problem parameters — hence often requires heavy tuning ; and (b) it cannot be guaranteed that $\mathcal{X}^*$ is reached in the limit [13, 16, 15]. A simple fix for the last problem is to allow polynomially decreasing stepsizes (second option) [23]: this choice for γ_k often leads to convergence to $\mathcal{X}^*$, but hurts the overall algorithm speed. The third option, which became very popular with the rise of deep learning, is to implement an *adaptive* stepsize. These methods do not commit to a fixed schedule, but instead use the optimization statistics (e.g. gradient history, cost history) to tune the value of γ_k at each iteration. These stepsizes are known to work very well in deep learning [35], and include Adam [19], Adagrad [11], and RMSprop [29].

Ideally, a theoretically grounded adaptive method should yield fast convergence to $\mathcal{X}^*$ without knowledge of problem dependent parameters, such as the gradient Lipshitz constant or the strong convexity constant. As a result, an ideal adaptive method should require very little tuning by the user, while matching the performance of a fine-tuned γ_k. However, while in practice this is the case for common adaptive methods such as Adam and AdaGrad, the associated convergence rates often rely on strong assumptions — e.g. that the iterates live on a bounded domain, or that gradients are uniformly bounded in norm [11, 32, 31]. While the above assumptions are valid in the constrained setting, they are problematic for problems defined in the whole $\mathbb{R}^d$.

A promising new direction in the adaptive stepsizes literature is based on the idea of Polyak stepsizes, introduced by [25] in the context of deterministic convex optimization. Recently [22] successfully adapted Polyak stepsizes to the stochastic setting, and provided convergence rates matching fine-tuned SGD — while the algorithm does not require knowledge of the unknown quantities such as the gradient Lipschitz constant. The results especially shines in the overparameterized strongly convex setting, where linear convergence to x^* is shown. This result is especially important since, under the same assumption, no such rate exists for AdaGrad (see e.g. [31] for the latest results) or other adaptive stepsizes. Moreover, the method was shown to work surprisingly well on deep learning problems, without requiring heavy tuning [22].

Even if the stochastic Polyak stepsize (SPS) [22] comes with strong convergence guarantees, it has two main drawbacks when it is used in non-over-parameterized regimes: (i) It requires *a priori* knowledge of the optimal mini-batch losses, which are not often available for big batch sizes or regularized objectives (see discussion in §1.1) and (ii) it guarantees convergence only to a neighborhood of the solution. In this work, we study the dynamics and the convergence properties of SGD equipped with new variants of SPS for solving general convex optimization problems. Our new proposed variants provide solutions to both drawbacks of the original SPS.

1.1 Background and Technical Preliminaries

The stepsize proposed by [22] is

$$\gamma_k = \min\left\{\frac{f_{\mathcal{S}_k}(x^k) - f_{\mathcal{S}_k}^*}{c\|\nabla f_{\mathcal{S}_k}(x^k)\|^2}, \gamma_b\right\}, \tag{SPS$_{\max}$}$$

where $\gamma_b, c > 0$ are problem-independent constants, $f_{\mathcal{S}_k} := \frac{1}{|\mathcal{S}_k|}\sum_{i \in \mathcal{S}_k} f_i$, $f_{\mathcal{S}_k}^* = \inf_{x \in \mathbb{R}^d} f_{\mathcal{S}_k}(x)$.
Dependency on $f_{\mathcal{S}_k}$. Crucially the algorithm requires knowledge of $f_{\mathcal{S}_k}^*$ for every realization of the mini-batch $\mathcal{S}_k$. In the non-regularized overparametrized setting (e.g. neural networks), $f_{\mathcal{S}_k}$ is often zero for every subset $\mathcal{S}$ [34]. However, this is not the only setting where $f_{\mathcal{S}}^*$ is computable: e.g., in the regularized logistic loss with batch size 1, it is possible to recover a cheap closed form expression for each f_i^* [22]. Unfortunately, if the batch-size is bigger than 1 or the loss becomes more demanding (e.g. cross-entropy), then *no such closed-form computation is possible*.

Rates and comparison with AdaGrad. In the convex overparametrized setting (more precisely, under the interpolation condition, i.e. $\exists\, x^* \in \mathcal{X}^*: \inf_{x \in \mathbb{R}^d} f_{\mathcal{S}}(x) = f_{\mathcal{S}}(x^*)$ for all $\mathcal{S}$, see also §2), SPS$_{\max}$ enjoys a convergence speed of $\mathcal{O}(1/k)$, without requiring knowledge of the gradient Lipschitz constant or other problem parameters. Recently, [31] showed that the same rate can be achieved for

AdaGrad in the same setting. However, there is an important difference: the rate of [31] is technically $\mathcal{O}(dD^2/k)$, where d is the problem dimension and D^2 is a global bound on the squared distance to the minimizer, which is assumed to be finite. Not only does $\text{SPS}_{\max}$ not have this dimension dependency, which dates back to crucial arguments in the AdaGrad literature [11, 21], but also does not require bounded iterates. While this assumption is satisfied in the constrained setting, it has no reason to hold in the unconstrained scenario. Unfortunately, this is a common problem of all AdaGrad variants: with the exception of [33] (which works in a slightly different scenario), no rate can be provided in the stochastic setting without the bounded iterates/gradients [12] assumption — even after assuming strong convexity. However, in the non-interpolated setting, AdaGrad enjoys a convergence guarantee of $\mathcal{O}(1/\sqrt{k})$ (with the bounded iterates assumption). A similar rate does not yet exist for SPS, and our work aims at filling this gap.

1.2 Main Contributions

As we already mentioned, in the non-interpolated setting $\text{SPS}_{\max}$ has the following issues:

Issue (1): For $B > 1$ (minibatch setting), $\text{SPS}_{\max}$ requires the exact knowledge of $f_{\mathcal{S}}^*$. This is not practical.

Issue (2): $\text{SPS}_{\max}$ guarantees convergence to a neighborhood of the solution. It is not clear how to modify it to yield convergence to the exact minimizer.

Having the above two issues in mind, the main contributions of our work (see also Table 1 in the appendix, for a summary of the main complexity results obtained in this paper) are summarized as follows:

- In §3, we provide a direct solution for Issue (1). We explain how only a lower bound on $f_{\mathcal{S}}^*$ (trivial if all f_is are non-negative) is required for convergence to a neighborhood of the solution. While this neighborhood is bigger that the one for $\text{SPS}_{\max}$, our modified version provides a practical baseline for the solution to the second issue.

- We explain why Issue (2) is highly non-trivial and requires an in-depth study of the bias induced by the interaction between gradients and Polyak stepsizes. Namely, we show that simply multiplying the stepsize of $\text{SPS}_{\max}$ by $1/\sqrt{k}$ — which would work for vanilla SGD [23] — yields a bias in the solution found by SPS (§4), regardless of the estimation of $f_{\mathcal{S}}^*$.

- In §5, we provide a solution to the problem (Issue (2)) by introducing additional structure — as well as the fix to Issue (1) — into the stepsize. We call the new algorithm *Decreasing SPS* (DecSPS), and provide a convergence guarantee under the bounded domain assumption — matching the standard AdaGrad results.

- In §5.2 we go one step further and show that, if strong convexity is assumed, iterates are bounded with probability 1 and hence we can remove the bounded iterates assumption. To the best of our knowledge, DecSPS, is the first stochastic adaptive optimization method that converges to the exact solution without assuming strong assumptions like bounded iterates/gradients.

- In §5.3 we provide extensions of our approach to the non-smooth setting.

- In §6, we corroborate our theoretical results with experimental testing.

2 Background on Stochastic Polyak Stepsize

In this section, we provide a concise overview of the results in [22], and highlight the main assumptions and open questions.

To start, we remind the reader that problem (1) is said to be interpolated if there exists a problem solution $x^* \in \mathcal{X}^*$ such that $\inf_{x \in \mathbb{R}^d} f_i(x) = f_i(x^*)$ for all $i \in [n]$. The degree of interpolation at batch size B can be quantified by the following quantity, introduced by [22] and studied also in [31, 9]: fix a batch size B, and let $\mathcal{S} \subseteq [n]$ with $|\mathcal{S}| = B$.

$$\sigma_B^2 := \mathbb{E}_{\mathcal{S}}[f_{\mathcal{S}}(x^*) - f_{\mathcal{S}}^*] = f(x^*) - \mathbb{E}_{\mathcal{S}}[f_{\mathcal{S}}^*] \tag{2}$$

It is easy to realize that as soon as problem (1) is interpolated, then $\sigma_B^2 = 0$ for each $B \leq n$. In addition, note that σ_B^2 is non-increasing as a function of B.

125 We now comment on the main result from [22].

126 **Theorem 2.1** (Main result of [22]). *Let each f_i be L_i-smooth convex functions. Then SGD with*

127 *SPS_{max}, mini-batch size B, and $c = 1$, converges as:* $\mathbb{E}\left[f(\bar{x}^K) - f(x^*)\right] \leq \frac{\|x^0 - x^*\|^2}{\alpha K} + \frac{2\gamma_b \sigma_B^2}{\alpha}$,

128 *where $\alpha = \min\left\{\frac{1}{2cL_{max}}, \gamma_b\right\}$ and $\bar{x}^K = \frac{1}{K}\sum_{k=0}^{K-1} x^k$. If in addition f is μ-strongly convex, then,*

129 *for any $c \geq 1/2$, SGD with SPS_{max} converges as:* $\mathbb{E}\|x^k - x^*\|^2 \leq (1 - \mu\alpha)^k \|x^0 - x^*\|^2 + \frac{2\gamma_b \sigma_B^2}{\mu\alpha}$,

130 *where again $\alpha = \min\{\frac{1}{2cL_{max}}, \gamma_b\}$ and $L_{max} = \max\{L_i\}_{i=1}^n$ is the maximum smoothness constant.*

131 In the overparametrized setting, the result guarantees convergence to the exact minimizer, without

132 knowledge of the gradient Lipschitz constant (as vanilla SGD would instead require) and without

133 assuming bounded iterates (in contrast to [31]).

134 As soon as (1) a *regularizer* is applied to the loss (e.g. $L2$ penalty), or (2) the number of datapoints

135 gets comparable to the dimension, then the problem is not interpolated and SPS_{max} only converges

136 to a neighborhood and it gets impractical to compute f_S^* — *this is the setting we study in this paper.*

137 *Remark* 1 (What if $\|\nabla f_{S_k}\| = 0$?). In the rare case that $\|\nabla f_{S_k}(x^k)\|^2 = 0$, there is no need to

138 evaluate the stepsize. In this scenario, the update direction $\nabla f_{S_k}(x^k) = 0$ and thus the iterate is

139 not updated irrespective of the choice of step-size. If this happens, the user should simply sample a

140 different minibatch. We note that in our experiments (see §6), such event never occurred.

141 **Related work on Polyak stepsize:** The classical Polyak stepsize [25] has been successfully used

142 in the analysis of deterministic subgradient methods in different settings [5, 7, 18]. First attempts

143 on providing an efficient variant of the stepsize that works well in the stochastic setting were made

144 in [3, 24]. However, as explained in [22], none of these approaches provide a natural stochastic

145 extension with strong theoretical convergence guarantees, and thus Loizou et al. [22] proposed the

146 stochastic Polyak stepsize SPS_{max} as a better alternative.[1] Despite its recent appearance, SPS_{max} has

147 already been used and analyzed as a stepsize for SGD for solving structured non-convex problems

148 [15], in combination with other adaptive methods [31], with a moving target [17] and in the update

149 rule of stochastic mirror descent [9]. These extensions are orthogonal to our approach, and we

150 speculate that our proposed variants can also be used in the above settings. We leave such extensions

151 for future work.

3 Removing f_S^* from SPS

153 As motivated in the last sections, computing f_S^* in the non-interpolated setting is not practical. In this

154 section, we explore the effect of using a lower bound $\ell_S^* \leq f_S^*$ instead in the SPS_{max} definition.

$$\gamma_k = \min\left\{\frac{f_{S_k}(x^k) - \ell_{S_k}^*}{c\|\nabla f_{S_k}(x^k)\|^2}, \gamma_b\right\}, \tag{SPS_{max}^ℓ}$$

156 Such a lower bound is easy to get for many problems of interest: indeed, for standard regularized

157 regression and classification tasks, the loss is non-negative hence one can pick $\ell_S^* = 0$, for any

158 $S \subseteq [n]$.

159 The obvious question is: what is the effect of estimating ℓ_S^* on the convergence rates in Thm. 2.1?

160 We found that the proof of [22] is easy to adapt to this case, by using the following fundamental

161 bound (see also Lemma B.2): $\frac{1}{2cL_{S_k}} \leq \frac{f_{S_k}(x^k) - f_{S_k}^*}{c\|\nabla f_{S_k}(x^k)\|^2} \leq \frac{f_{S_k}(x^k) - \ell_{S_k}^*}{c\|\nabla f_{S_k}(x^k)\|^2}$.

162 The following results can be seen as an easy extension of the main result of [22], under a newly

163 defined suboptimality measure:

$$\hat{\sigma}_B^2 := \mathbb{E}_{S_k}[f_{S_k}(x^*) - \ell_{S_k}^*] = f(x^*) - \mathbb{E}_{S_k}[\ell_{S_k}^*]. \tag{3}$$

---

[1] Unknown to these authors, a variant of SGD with SPS_{max} was already proposed by Asi and Duchi [2] as a special case of a model-based method called the lower-truncated model. Asi and Duchi [2] also proposed a decreasing step-size variant of SPS_{max} which is closely related but different than the DecSPS that we propose in §5. Among some differences, they assume interpolation for their convergence results whereas we do not in §5. We describe the differences between our work and Asi and Duchi [2] in more detail in Appendix A.

Figure 1: We consider a 100 dim problem with $n = 100$ datapoints where each $f_i = \frac{1}{2}(x - x_i^*)^\top H_i(x - x_i^*) + f_i^*$, with $f_i^* = 1$ for all $i \in [n]$ and H_i a random SPD matrix generated using the standard Gaussian matrix $A_i \in \mathbb{R}^{d \times 3d}$ as $H_i = A_i A_i^\top /(3d)$. If $x_i^* \neq x_j^*$ for $i \neq j$, then the problem does **not satisfy interpolation (left plot)**. Instead, if all x_i^*s are equal, **then the problem is interpolated (central plot)**. The plot shows the behaviour of $\text{SPS}_{\max}^\ell$ ($\gamma_b = 2$) for different choices of the approximated suboptimality ℓ_i^*. We plot (mean and std deviation over 10 runs) the function suboptimality level $f(x) - f(x^*)$ for different values of ℓ_i^*. Note that, if instead $f_i^* = 0$ for all i then all the shown **algorithms coincide (right plot)** and converge to the solution.

Theorem 3.1. *Under $\text{SPS}_{\max}^\ell$, the same exact rates in Thm. 2.1 hold (under the corresponding assumptions), after replacing σ_B^2 with $\hat{\sigma}_B^2$.*

And we also have an easy practical corollary.

Corollary 3.2. *In the context of Thm. 3.1, assume all f_is are non-negative and estimate $\ell_S^* = 0$ for all $S \subseteq [n]$. Then the same exact rates in Thm. 2.1 hold for $\text{SPS}_{\max}^\ell$, after replacing σ_B^2 with $f^* = f(x^*)$.*

A numerical illustration of this result can be found in Fig. 1. In essence, both theory and experiments confirm that, if interpolation is not satisfied, then we have a rate until a convergence ball, where the size is optimal under exact knowledge of f_S^*. Instead, under interpolation, if all the f_is are non-negative and $f^* = 0$, then $\text{SPS}_{\max} = \text{SPS}_{\max}^\ell$. Finally, in the less common case in practice where $f^* > 0$ but we still have interpolation, then $\text{SPS}_{\max}$ converges to the exact solution while $\text{SPS}_{\max}^\ell$ does not. To conclude $\text{SPS}_{\max}^\ell$ does not (of course) work better than $\text{SPS}_{\max}$, but it is a practical variant which we can use as a baseline in §5 for an adaptive stochastic Polyak stepsize with convergence to the true x^* in the non-interpolated setting.

4 Bias in the SPS dynamics.

In this section, we study convergence of the standard $\text{SPS}_{\max}$ in the non-interpolated regime, under an additional (decreasing) multiplicative factor, in the most ideal setting: batch size 1, and we have knowledge of each f_i^*. That is, we consider $\gamma_k = \min\{\frac{f_{i_k}(x^k) - f_{i_k}^*}{c_k \|\nabla f_{i_k}(x^k)\|^2}, \gamma_b\}$ with $c_k \to \infty$, e.g. $c_k = \mathcal{O}(\sqrt{k})$ or $c_k = \mathcal{O}(k)$. We note that, in the SGD case, simply picking e.g. $\gamma_k = \gamma_0/\sqrt{k+1}$ would guarantee convergence of $f(x^k)$ to $f(x^*)$, in expectation and with high probability [20, 23]. Therefore, it is natural to expect a similar behavior for SPS, if $1/c_k$ safisfies the usual Robbins-Monro conditions [27]: $\sum_{k=0}^\infty 1/c_k = \infty, \sum_{k=0}^\infty 1/c_k^2 < \infty$.

We show that this is not the case: quite interestingly, $f(x^k)$ converges to a biased solution due to the *correlation between ∇f_{i_k} and γ_k.* we show this formally, in the case of non-interpolation (otherwise both SGD and SPS do not require a decreasing learning rate).

Counterexample. Consider the following finite-sum setting: $f(x) = \frac{1}{2}f_1(x) + \frac{1}{2}f_2(x)$ with $f_1(x) = \frac{a_1}{2}(x - 1)^2, f_2(x) = \frac{a_2}{2}(x + 1)^2$. To make the problem interesting, we choose $a_1 = 2$ and $a_2 = 1$: this introduces asymmetry in the average landscape with respect to the origin. During optimization, we sample f_1 and f_2 independently and seek convergence to the unique minimizer $x^* = \frac{a_1 - a_2}{a_1 + a_2} = 1/3$. The first thing we notice is that x^* is not a stationary point for the dynamics under SPS. Indeed note that since $f_i^* = 0$ for $i = 1, 2$ we have (assuming γ_b large enough): $\gamma_k \nabla f_{i_k}(x) = \frac{x-1}{2c_k}$, if $i_k = 1$, and $\gamma_k \nabla f_{i_k}(x) = \frac{x+1}{2c_k}$ if $i_k = 2$.

Crucially, note that this update is *curvature-independent*. The expected update is $\mathbb{E}_{i_k}[\gamma_k \nabla f_{i_k}(x)] = \frac{x-1}{4c_k} + \frac{x+1}{4c_k} = \frac{1}{2c_k}x$. Hence, the iterates can only converge to $x = 0$ — because this is the only fixed point for the update rule. The proof naturally extends to the multidimensional setting, an illustration can be found in Fig. 2.

In the same picture, we show how our modified variant of the vanilla stepsize — we call this new algorithm DecSPS, see §5 — instead converges to the correct solution.

Remark 2. SGD with (non-adaptive) stepsize γ_k instead keeps the curvature, and therefore is able to correctly estimate the average $\mathbb{E}_{i_k}[\gamma_k \nabla f_{i_k}(x)] = \frac{\gamma_k}{2}(a_1 + a_2)\left[x - \frac{a_1 - a_2}{a_1 + a_2}\right]$ — precisely because γ_k is independent from ∇f_{i_k}. From this we can see that SGD can only converge to the correct stationary point $x^* = \frac{a_1 - a_2}{a_1 + a_2}$ — because again this is the only fixed point for the update rule.

In the appendix, we go one step further and provide an analysis of the bias of SPS in the one-dimensional quadratic case (Prop. D.2). Yet, we expect the precise characterization of the bias phenomenon in the non-quadratic setting to be particularly challenging. We provide additional insights in §D.2. Instead, in the next section, we show how to effectively modify γ_k to yield convergence to x^* without further assumptions.

No interpolation   **Interpolation**

Figure 2: Dynamics of $\text{SPS}_{\max}$ with decreasing multiplicative constant (SGD style) compared with DecSPS. We compared both in the **interpolated setting (right)** and in the **non-interpolated setting (left)**. In the non-interpolated setting, a simple multiplicative factor introduces a bias in the final solution, as discussed in this section. We consider two dimensional $f_i = \frac{1}{2}(x - x_i^*)^\top H_i(x - x_i^*)$, for $i = 1, 2$ and plot the contour lines of the corresponding landscapes, as well as the average landscape $(f_1 + f_2)/2$ we seek to minimize. Solution is denoted with a gold star.

5   DecSPS: Convergence to the exact solution

We propose the following modification of the vanilla SPS proposed in [22], designed to yield convergence to the exact minimizer while keeping the main adaptiveness properties[2]. We call it Decreasing SPS (DecSPS), since it combines a steady stepsize decrease with the adaptiveness of SPS.

$$\gamma_k := \frac{1}{c_k} \min\left\{ \frac{f_{\mathcal{S}_k}(x^k) - \ell_{\mathcal{S}_k}^*}{\|\nabla f_{\mathcal{S}_k}(x^k)\|^2},\ c_{k-1}\gamma_{k-1} \right\}, \tag{DecSPS}$$

for $k \in \mathbb{N}$, where $c_k \neq 0$ for every $k \in \mathbb{N}$. We set $c_{-1} = c_0$ and $\gamma_{-1} = \gamma_b > 0$ (stepsize bound, similar to [22]), to get $\gamma_0 := \frac{1}{c_0} \cdot \min\left\{ \frac{f_{\mathcal{S}_k}(x^k) - \ell_{\mathcal{S}_k}^*}{\|\nabla f_{\mathcal{S}_k}(x^k)\|^2},\ c_0\gamma_b \right\}$.

Lemma 5.1. *Let each f_i be L_i smooth and let $(c_k)_{k=0}^\infty$ be any non-decreasing positive sequence of real numbers. Under DecSPS, we have* $\min\left\{ \frac{1}{2c_k L_{\max}}, \frac{c_0\gamma_b}{c_k} \right\} \leq \gamma_k \leq \frac{c_0\gamma_b}{c_k}$, *and* $\gamma_{k-1} \leq \gamma_k$

Remark 3. As stated in the last lemma, under the assumption of c_k non-decreasing, γ_k is trivially *non-increasing* since $\gamma_k \leq c_{k-1}\gamma_{k-1}/c_k$.

The proof can be found in the appendix, and is based on a simple induction argument.

5.1   Convergence under bounded iterates

The following result provides a proof of convergence of SGD for the γ_k sequence defined above.

Theorem 5.2. *Consider SGD with DecSPS and let $(c_k)_{k=0}^\infty$ be any non-decreasing sequence such that $c_k \geq 1, \forall k \in \mathbb{N}$. Assume that each f_i is convex and L_i smooth. We have:*

$$\mathbb{E}[f(\bar{x}^K) - f(x^*)] \leq \frac{2c_{K-1}\tilde{L}D^2}{K} + \frac{1}{K}\sum_{k=0}^{K-1} \frac{\hat{\sigma}_B^2}{c_k}, \tag{4}$$

where $D^2 := \max_{k \in [K-1]} \|x^k - x^*\|^2$, $\tilde{L} := \max\left\{ \max_i\{L_i\}, \frac{1}{2c_0\gamma_b} \right\}$ and $\bar{x}^K = \frac{1}{K}\sum_{k=0}^{K-1} x^k$.

[2] Similar choices are possible. We found that this leads to the biggest stepsize magnitude, allowing for faster convergence in practice.

If $\hat{\sigma}_B^2 = 0$, then $c_k = 1$ for all $k \in \mathbb{N}$ leads to a rate $\mathcal{O}(\frac{1}{K})$, well known from [22]. If $\hat{\sigma}_B^2 > 0$, as for the standard SGD analysis under decreasing stepsizes, the choice $c_k = \mathcal{O}(\sqrt{k})$ leads to an optimal asymptotic trade-off between the deterministic and the stochastic terms, hence to the asymptotic rate $\mathcal{O}(1/\sqrt{k})$ since $\sum_{k=0}^{K-1} \frac{1}{\sqrt{k+1}} \leq 2\sqrt{K}$. Moreover, picking $c_0 = 1$ minimizes the speed of convergence for the deterministic factor. Under the assumption that $\hat{\sigma}_B^2 \ll \tilde{L}D^2$ (e.g. reasonable distance initialization-solution and $L_{\max} > 1/\gamma_b$), this factor is dominant compared to the factor involving $\hat{\sigma}_B^2$. For this setting, the rate simplifies as follows.

Corollary 5.3. *Under the setting of Thm. 5.2, for $c_k = \sqrt{k+1}$ ($c_{-1} = c_0$) we have*

$$\mathbb{E}[f(\bar{x}^K) - f(x^*)] \leq \frac{2\tilde{L}D^2 + 2\hat{\sigma}_B^2}{\sqrt{K}}. \tag{5}$$

Remark 4 (Beyond bounded iterates). The result above crucially relies on the bounded iterates assumption: $D^2 < \infty$. To the best of our knowledge, if no further regularity is assumed, modern convergence results for adaptive methods (e.g. variants of AdaGrad) in convex stochastic programming require[3] this assumption, or else require gradients to be globally bounded. To mention a few: [11, 26, 32, 8, 31]. A simple algorithmic fix to this problem is adding a cheap projection step onto a large bounded domain [21]. We can of course include this projection step in DecSPS, and the theorem above will hold with no further modification. Yet we found this to be not necessary: the strong guarantees of SPS in the strongly convex setting [22] let us go one step beyond: in §5.2 we show that, if each f_i is strongly convex (e.g. regularizer is added), then one can bound the iterates globally with probability one, without knowledge of the gradient Lipschitz constant. To the best of our knowledge, no such result exist for AdaGrad — except [30], for the deterministic case.

Remark 5 (Dependency on the problem dimension). In standard results for AdaGrad, a dependency on the problem dimension often appears (e.g. Thm. 1 in [31]). This dependency follows from a bound on the AdaGrad preconditioner that can be found e.g. in Thm. 4 in [21]. In the SPS case no such dependency appears — specifically because the stepsize is lower bounded by $1/(2c_k L_{\max})$.

5.2 Removing the bounded iterates assumption

We prove that under DecSPS the iterates live in a set of diameter $D_{\max}$ almost surely. This can be done by assuming strong convexity of each f_i.

The result uses this alternative definition of *neighborhood*: $\hat{\sigma}_{B,\max}^2 := \max_{\mathcal{S} \subseteq [n], |\mathcal{S}|=B}[f_{\mathcal{S}}(x^*) - \ell_{\mathcal{S}}^*]$.

Note that trivially $\hat{\sigma}_{B,\max}^2 < \infty$ under the assumption that all f_i are lower bounded and $n < \infty$.

Proposition 5.4. *Let each f_i be μ_i-strongly convex and L_i-smooth. The iterates of SGD with DecSPS with $c_k = \sqrt{k+1}$ (and $c_{-1} = c_0$) are such that $\|x^k - x^*\|^2 \leq D_{\max}^2$ almost surely $\forall k \in \mathbb{N}$, where $D_{\max}^2 := \max\left\{\|x^0 - x^*\|^2, \frac{2c_0\gamma_b\hat{\sigma}_{B,\max}^2}{\min\left\{\frac{\mu_{\min}}{2L_{\max}}, \mu_{\min}\gamma_b\right\}}\right\}$, with $\mu_{\min} = \min_{i \in [n]} \mu_i$ and $L_{\max} = \max_{i \in [n]} L_i$.*

The proof relies on the variations of constants formula and an induction argument — it is provided in the appendix. We are now ready to state the main theorem for the unconstrained setting, which follows from Prop. 5.4 and Thm. 5.2.

Theorem 5.5. *Consider SGD with the DecSPS stepsize $\gamma_k := \frac{1}{\sqrt{k+1}} \cdot \min\left\{\frac{f_{\mathcal{S}_k}(x^k) - f_{\mathcal{S}_k}^*}{\|\nabla f_{\mathcal{S}_k}(x^k)\|^2}, \gamma_{k-1}\sqrt{k}\right\}$, for $k \geq 1$ and γ_0 defined as at the beginning of this section. Let each f_i be μ_i-strongly convex and L_i-smooth:*

$$\mathbb{E}[f(\bar{x}^K) - f(x^*)] \leq \frac{2\tilde{L}D_{\max}^2 + 2\hat{\sigma}_B^2}{\sqrt{K}}. \tag{6}$$

Remark 6. The careful reader might notice that, while we assumed strong convexity, our rate is slower than the optimal $\mathcal{O}(1/K)$. This issue is due to the adaptive nature of DecSPS. It is indeed notoriously

---

[3] Perhaps the only exception is the result of [32], where the authors work on a different setting: i.e. they introduce the RUIG inequality.

hard to achieve a convergence rate of $\mathcal{O}(1/K)$ for adaptive methods in the strongly convex regime. While further investigations will for sure shed light on this interesting problem, we note that *the result we provide is somewhat unique in the literature*: we are not aware of any adaptive method that enjoys such a convergence rate without either (a) assuming bounded iterates/gradients or (b) assuming knowledge of the gradient Lipschitz constant and or the strong convexity constant.

5.3 Extension to the non-smooth setting

For any $\mathcal{S} \subseteq [n]$, we denote in this section by $g_{\mathcal{S}}(x)$ the subgradient of $f_{\mathcal{S}}$ evaluated at x. We discuss the extension of DecSPS to the non-smooth setting.

A straightforward application of DecSPS leads to a stepsize γ_k which is no longer lower bounded (see Lemma 5.1) by the positive quantity $\min\left\{\frac{1}{2c_k L_{\max}}, \frac{c_0 \gamma_b}{c_k}\right\}$. Indeed, the gradient Lipschitz constant in the non-smooth case is formally $L_{\max} = \infty$. Hence, γ_k prescribed by DecSPS can get arbitrarily small[4] for finite k. One easy solution to the problem is to enforce a lower bound, and adopt a new proof technique. Specifically we propose the following:

$$\gamma_k := \frac{1}{c_k} \cdot \min\left\{\max\left\{c_0 \gamma_\ell, \frac{f_{\mathcal{S}_k}(x^k) - \ell_{\mathcal{S}_k}^*}{\|g_{\mathcal{S}_k}(x^k)\|^2}\right\}, c_{k-1}\gamma_{k-1}\right\}, \qquad \text{(DecSPS-NS)}$$

where $c_k \neq 0$ for every $k \geq 0$, $\gamma_\ell \geq \gamma_b$ is a small positive number and all the other quantities are defined as in DecSPS. In particular, as for DecSPS, we set $c_{-1} = c_0$ and $\gamma_{-1} = \gamma_b$. Intuitively, γ_k is selected to live in the interval $[c_0\gamma_\ell/c_k, c_0\gamma_b/c_k]$ (see proof in §F, appendix), but has subgradient-dependent adaptive value. In addition, this stepsize is enforced to be monotonically decreasing.

Theorem 5.6. *For any non-decreasing positive sequence* $(c_k)_{k=0}^{\infty}$*, consider SGD with DecSPS-NS. Assume that each f_i is convex and lower bounded. We have*

$$\mathbb{E}[f(\bar{x}^K) - f(x^*)] \leq \frac{c_{K-1}D^2}{\gamma_\ell c_0 K} + \frac{1}{K}\sum_{k=0}^{K-1}\frac{c_0\gamma_b G^2}{c_k}, \qquad (7)$$

where $D^2 := \max_{k\in[K-1]}\|x^k - x^*\|^2$ *and* $G^2 := \max_{k\in[K-1]}\|g_{\mathcal{S}_k}(x^k)\|^2$.

One can then easily derive an $\mathcal{O}(1/\sqrt{k})$ convergence rate. This is presented in §F (appendix).

6 Numerical Evaluation

We evaluate the performance of DecSPS with $c_k = c_0\sqrt{k+1}$ on binary classification tasks, with regularized logistic loss $f(x) = \frac{1}{n}\sum_{i=1}^{n}\log(1 + \exp(y_i \cdot a_i^\top x)) + \frac{\lambda}{2}\|x\|^2$, where $a_i \in \mathbb{R}^d$ is the feature vector for the i-th datapoint and $y_i \in \{-1, 1\}$ is the corresponding binary target.

We study performance on three datasets: (1) a Synthetic Dataset, (2) The A1A dataset [6] and (3) the Breast Cancer dataset [10]. We choose different regularization levels and batch sizes bigger than 1. Details are reported in §G, and the code is available in the supplementary material.

At the batch sizes and regularizer levels we choose, the problems do not satisfy interpolation. Indeed, running full batch gradient descent yields $f^* > 0$. While running $\text{SPS}_{\max}$ on these problems (1) does not guarantee convergence to f^* and (2) requires full knowledge of the set of optimal function values $\{f_{\mathcal{S}}^*\}_{|\mathcal{S}|=B}$, in DecSPS we can simply pick the lower bound $0 = \ell_{\mathcal{S}}^* \leq f_{\mathcal{S}}^*$ for every $\mathcal{S}$. Supported by Theorems 5.2 & 5.5 & 5.6, we expect SGD with DecSPS to converge to the minimum f^*.

Stability of DecSPS. DecSPS has two hyperparameters: the upper bound γ_b on the first stepsize and the scaling constant c_0. While Thm. 5.6 guarantees convergence for any positive value of these hyperparameters, the result of Thm. 5.2 suggests that using $c_0 = 1$ yields the best performance under the assumption that $\hat{\sigma}_B^2 \ll \tilde{L}D^2$ (e.g. reasonable distance of initialization from the solution, and $L_{\max} > 1/\gamma_b$). In Fig. 3, we show on the synthetic dataset that (1) $c_0 = 1$ is indeed the best choice in this setting and (2) the performance of SGD with DecSPS is almost independent of γ_b. Similar

---

[4] Take for instance the deterministic setting one-dimensional setting $f(x) = |x|$. As $x \to 0$, the stepsize prescribed by DecSPS converges to zero. This is not the case e.g. in the quadratic setting.

findings are reported and commented in Figure 9 (Appendix) for the other datasets. Hence, *for all further experiments*, we choose the hyperparameters $\gamma_b = 10, c_0 = 1$.

Comparison with vanilla SGD with decreasing step-size. We compare the performance of DecSPS against the classical decreasing SGD stepsize $\eta/\sqrt{k+1}$, which guarantees convergence to the exact solution at the same asymptotic rate as DecSPS. We show that, while the asymptotics are the same, DecSPS with the theoretically inspired hyperparameters $c_0 = 1, \gamma_b = 10$ performs competitively to a fine-tuned η — where crucially the optimal value of η depends on the problem. This behavior is shown on all the considered datasets, and is reported in Figures 4 (*Breast* and *Synthetic* reported in the appendix for space constraints).

Figure 3: DecSPS ($c_k = c_0\sqrt{k+1}$) sensitivity to hyperparameters on the *Synthetic Dataset*, with $\lambda = 0$. Experiments are repeated 3 times and plotted is mean and std.

DecSPS yields a truly adaptive stepsize. We inspect the value of γ_k returned by DecSPS, shown in Figures 4 & 10 (in the appendix). Compared to the vanilla SGD stepsize $\eta/\sqrt{k+1}$, a crucial difference appears: γ_k decreases faster than $\mathcal{O}(1/\sqrt{k})$. This showcases that, while the factor $\sqrt{k+1}$ can be found in the formula of DecSPS[5], the algorithm structure provides additional adaptation to curvature. Indeed, in (regularized) logistic regression, the local gradient Lipschitz constant increases as we approach the solution. Since the optimal stepsize for steadily-decreasing SGD is $1/(L\sqrt{k+1})$, where L is the global Lipschitz constant [13], it is pretty clear that η should be decreased over training for optimal converge (as L effectively increases). This is precisely what DecSPS is doing.

Comparison with AdaGrad stepsizes. Last, we compare DecSPS with another adaptive coordinate-independent stepsize with strong theoretical guarantees: the norm version of AdaGrad (a.k.a. AdaGrad-Norm, AdaNorm), which guarantees the exact solution at the same asymptotic rate as DecSPS [32]. AdaGrad-norm at each iteration updates the scalar $b_{k+1}^2 = b_k^2 + \|\nabla f_{\mathcal{S}_k}(x_k)\|^2$ and then selects the next step as $x_{k+1} = x_k - \frac{\eta}{b_{k+1}}\nabla f_i(x_k)$. Hence, it has tuning parameters b_0 and η. In Fig. 4 we show that, on the Breast Cancer dataset, after fixing $b_0 = 0.1$ as recommended in [32] (see their Figure 3), tuning η cannot quite match the performance of DecSPS. This behavior is also observed on the other two datasets we consider (see Fig. 12 in the Appendix). Last, in Fig. 11 in the Appendix, we show that further tuning of b_0 likely does not yield a substantial improvement.

Figure 4: **Left:** performance of DecSPS, on the A1A Dataset ($\lambda = 0.01$). **Right:** performance of DecSPS on the Breast Cancer Dataset ($\lambda = 1e - 1$). Further experiments can be found in §G (appendix).

Comparison with Adam and AMSgrad. In Figures 13&14 we compare DecSPS with Adam [19] and AMSgrad [26] on the A1A and Breast Cancer datasets. The results show that DecSPS with the usual hyperparameters is comparable to the fine-tuned version of both these algorithms — which however do not enjoy comparable convergence guarantees in the unbounded domain setting.

7   Conclusions and Future Work

We provided a practical variant of SPS [22] which converges to the true problem solution without the interpolation assumption in convex stochastic problems — matching the rate of AdaGrad. If in

---

[5] We pick $c_k = c_0\sqrt{k+1}$, as suggested by Cor. 5.3 & F.2

addition strong convexity as assumed, then we show how, in contrast to current results for AdaGrad, the bounded iterates assumption can be dropped. The main open direction is a proof of a faster rate $\mathcal{O}(1/K)$ under strong convexity. There are also other possible extensions of our work: using the proposed new variants of SPS with accelerated methods, studying further the effect of mini-batching and non-uniform sampling of DecSPS, and extensions to the distributed and decentralized settings.

References

[1] H. Asi and J. C. Duchi. The importance of better models in stochastic optimization. *Proceedings of the National Academy of Sciences*, 116(46):22924–22930, 2019.

[2] H. Asi and J. C. Duchi. Stochastic (approximate) proximal point methods: Convergence, optimality, and adaptivity. *SIAM Journal on Optimization*, 29(3):2257–2290, 2019.

[3] L. Berrada, A. Zisserman, and M. P. Kumar. Training neural networks for and by interpolation. In *International Conference on Machine Learning*, 2020.

[4] L. Bottou, F. E. Curtis, and J. Nocedal. Optimization methods for large-scale machine learning. *Siam Review*, 60(2):223–311, 2018.

[5] S. Boyd, L. Xiao, and A. Mutapcic. Subgradient methods. *Lecture Notes of EE392o, Stanford University, Autumn Quarter*, 2004:2004–2005, 2003.

[6] C.-C. Chang. LIBSVM: a library for support vector machines. *ACM Transactions on Intelligent Systems and Technology*, 2011.

[7] D. Davis, D. Drusvyatskiy, K. J. MacPhee, and C. Paquette. Subgradient methods for sharp weakly convex functions. *Journal of Optimization Theory and Applications*, 179(3):962–982, 2018.

[8] A. Défossez, L. Bottou, F. Bach, and N. Usunier. A simple convergence proof of Adam and Adagrad. *arXiv preprint arXiv:2003.02395*, 2020.

[9] R. D'Orazio, N. Loizou, I. Laradji, and I. Mitliagkas. Stochastic mirror descent: Convergence analysis and adaptive variants via the mirror stochastic Polyak stepsize. *arXiv preprint arXiv:2110.15412*, 2021.

[10] D. Dua and C. Graff. UCI machine learning repository, 2017. URL http://archive.ics.uci.edu/ml.

[11] J. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of Machine Learning Research*, 12(7), 2011.

[12] A. Ene and H. L. Nguyen. Adaptive and universal algorithms for variational inequalities with optimal convergence. *arXiv preprint arXiv:2010.07799*, 2020.

[13] S. Ghadimi and G. Lan. Stochastic first-and zeroth-order methods for nonconvex stochastic programming. *SIAM Journal on Optimization*, 23(4), 2013.

[14] I. Goodfellow, Y. Bengio, and A. Courville. *Deep Learning*. MIT press, 2016.

[15] R. Gower, O. Sebbouh, and N. Loizou. SGD for structured nonconvex functions: Learning rates, minibatching and interpolation. In *International Conference on Artificial Intelligence and Statistics*, 2021.

[16] R. M. Gower, N. Loizou, X. Qian, A. Sailanbayev, E. Shulgin, and P. Richtárik. SGD: General analysis and improved rates. In *International Conference on Machine Learning*, 2019.

[17] R. M. Gower, A. Defazio, and M. Rabbat. Stochastic Polyak stepsize with a moving target. *arXiv preprint arXiv:2106.11851*, 2021.

[18] E. Hazan and S. Kakade. Revisiting the Polyak step size. *arXiv preprint arXiv:1905.00313*, 2019.

[19] D. P. Kingma and J. Ba. Adam: A method for stochastic optimization. *arXiv:1412.6980*, 2014.

[20] H. Kushner and G. G. Yin. *Stochastic Approximation and Recursive Algorithms and Applications*, volume 35. Springer Science & Business Media, 2003.

[21] K. Y. Levy, A. Yurtsever, and V. Cevher. Online adaptive methods, universality and acceleration. *Advances in Neural Information Processing Systems*, 31, 2018.

[22] N. Loizou, S. Vaswani, I. H. Laradji, and S. Lacoste-Julien. Stochastic Polyak step-size for SGD: An adaptive learning rate for fast convergence. In *International Conference on Artificial Intelligence and Statistics*, 2021.

[23] A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on Optimization*, 19(4):1574–1609, 2009.

[24] A. M. Oberman and M. Prazeres. Stochastic gradient descent with Polyak's learning rate. *arXiv preprint arXiv:1903.08688*, 2019.

[25] B. Polyak. *Introduction to Optimization*. Inc., Publications Division, New York, 1987.

[26] S. J. Reddi, S. Kale, and S. Kumar. On the convergence of Adam and beyond. In *International Conference on Learning Representations*, 2018.

[27] H. Robbins and S. Monro. A stochastic approximation method. *The Annals of Mathematical Statistics*, 1951.

[28] M. Rolinek and G. Martius. L4: Practical loss-based stepsize adaptation for deep learning. *Advances in neural information processing systems*, 31, 2018.

[29] T. Tieleman and G. Hinton. Lecture 6.5 - RMSprop, Coursera: Neural networks for machine learning. *University of Toronto, Technical Report*, 2012.

[30] C. Traoré and E. Pauwels. Sequential convergence of AdaGrad algorithm for smooth convex optimization. *Operations Research Letters*, 49(4):452–458, 2021.

[31] S. Vaswani, I. Laradji, F. Kunstner, S. Y. Meng, M. Schmidt, and S. Lacoste-Julien. Adaptive gradient methods converge faster with over-parameterization (but you should do a line-search). *arXiv preprint arXiv:2006.06835*, 2020.

[32] R. Ward, X. Wu, and L. Bottou. Adagrad stepsizes: Sharp convergence over nonconvex landscapes. In *International Conference on Machine Learning*, 2019.

[33] Y. Xie, X. Wu, and R. Ward. Linear convergence of adaptive stochastic gradient descent. In *International Conference on Artificial Intelligence and Statistics*, 2020.

[34] C. Zhang, S. Bengio, M. Hardt, B. Recht, and O. Vinyals. Understanding deep learning (still) requires rethinking generalization. *Communications of the ACM*, 64(3):107–115, 2021.

[35] J. Zhang, S. P. Karimireddy, A. Veit, S. Kim, S. Reddi, S. Kumar, and S. Sra. Why are adaptive methods good for attention models? *Advances in Neural Information Processing Systems*, 33, 2020.

Figure 5: **Requested by Reviewer 98CJ:** Toy dataset, sensitivity of DecSPS to c_0 and γ_b with 10 experiments. Mean and 3 standard deviations are plotted.

Figure 6: **Requested by Reviewer CpM5** : Tuning b_0 in A1A (left plots) and Breast cancer (right plot) datasets. $b_0 = 0.1$ already shown in the appendix.

Figure 7: **Requested by Reviewer CpM5** : Results on A1A for $\lambda = 1e - 4$ (left) and $\lambda = 1e - 6$ (right). Additional tuning of SPS is required but performance of the tuned method is solid.