RANDOM COORDINATE LANGEVIN MONTE CARLO

ABSTRACT

Langevin Monte Carlo (LMC) is a popular Markov chain Monte Carlo sampling method. One drawback is that it requires the computation of the full gradient at each iteration, an expensive operation if the dimension of the problem is high. We propose a new sampling method: Random Coordinate LMC (RC-LMC). At each iteration, a single coordinate is randomly selected to be updated by a multiple of the partial derivative along this direction plus noise, and all other coordinates remain the same. We investigate the total complexity of RC-LMC and compare it with the classical LMC for log-concave probability distributions. When the gradient of the log-density is Lipschitz, RC-LMC is less expensive than classical LMC when f is highly skewed for high dimensional problems, and when both gradient and Hessian of the log-density are Lipschitz, RC-LMC is always cheaper than LMC, by a factor proportional to the square root of problem dimension. In the latter case, our estimate of complexity is sharp with respect to the dimension.

1 INTRODUCTION

Monte Carlo sampling plays an important role in machine learning (Andrieu et al., 2003) and Bayesian statistics. In applications, the need for sampling is found in atmospheric science (Fabian, 1981), epidemiology (Li et al., 2020), petroleum engineering (Nagarajan et al., 2007), in the form of data assimilation (Reich, 2011), volume computation (Vempala, 2010) and bandit optimization (Russo et al., 2018).

In many of these applications, the dimension of the problem is extremely high. For example, for weather prediction, one measures the current state temperature and moisture level, to infer the flow in the air, before running the Navier–Stokes equations into the near future (Evensen, 2009). The quantities to be inferred is the air flow, and in a global numerical weather prediction model, the degrees of freedom is as high as 10^9. Another example is from epidemiology: when a disease is spreading, one measures the everyday new infection cases to infer the transmission rate in different regions. On a county-level modeling, one treats $3,141$ different counties in the US separately, and the parameter to be inferred has dimension at least $3,141$ (Li et al., 2020).

In this work, we focus on Monte Carlo sampling of log-concave probability distributions on $\mathbb{R}^d$, meaning the probability density can be written as $p(x) \propto e^{-f(x)}$ where a $f(x)$ is a convex function. The goal is to generate (approximately) i.i.d. samples according to the target probability distribution with density $p(x)$. Several sampling frameworks have been proposed in the literature, including importance sampling and sequential Monte Carlo (Geweke, 1989; Neal, 2001; Del Moral et al., 2006), ensemble methods (Reich, 2011; Iglesias et al., 2013), Markov chain Monte Carlo (MCMC) (Roberts and Rosenthal, 2004), including Metropolis-Hasting based MCMC (MH-MCMC) (Metropolis et al., 1953; Hastings, 1970; Roberts and Tweedie, 1996), Gibbs samplers (Geman and Geman, 1984; Casella and George, 1992), and Hamiltonian Monte Carlo (Neal, 1993; Duane et al., 1987). Langevin Monte Carlo (LMC) (Rossky et al., 1978; Parisi, 1981; Roberts and Tweedie, 1996) is a popular MCMC method that has received quite intensive attention in recent years for the new progress in non-asymptotic analysis of its convergence property (Durmus and Moulines, 2017; Dalalyan, 2017; Dalalyan and Karagulyan, 2019; Durmus et al., 2019).

Denoting by x^m the location of the sample at m-th iteration, LMC obtains the next location as follows:
$$x^{m+1} = x^m - \nabla f(x^m)h + \sqrt{2h}\xi_d^m, \tag{1}$$

where h is the time stepsize, and ξ_d^m is drawn i.i.d. from $\mathcal{N}(0, I_d)$, where I_d denotes identity matrix of size $d \times d$. LMC can be viewed as the Euler-Maruyama discretization of the following stochastic differential equation (SDE):

$$\mathrm{d}X_t = -\nabla f(X_t)\,\mathrm{d}t + \sqrt{2}\,\mathrm{d}B_t\,, \tag{2}$$

where B_t is a d-dimensional Brownian motion with independent components. It is well known that under mild conditions, the SDE converges exponentially fast to the target distribution $p(x)$ (see e.g., (Markowich and Villani, 1999)). Since (1) approximates the SDE (2) with an $\mathcal{O}(h)$ discretization error, the probability distribution of x^m produced by LMC (1) converges exponentially to the target distribution up to a discretization error (Dalalyan and Karagulyan, 2019).

One main drawback of LMC is that its dependence on the problem dimension d is rather bad. In each iteration, the full gradient needs to be evaluated. However, in most practical problems, since the analytical expression of the gradient is not available, each partial derivative component in the gradient needs to be computed separately, either through finite differencing or automatic differentiation, making the cost roughly d times the number of required iterations. In the weather prediction and epidemiology problems discussed above, f stands for the map from the parameter space to the measured quantities via the underlying partial differential equations (PDEs), and each dimensional partial derivative calls for one forward and one adjoint PDE solve. That leads to $2d$ PDE solves in total per one iteration. Another example comes from the study of directed graphs with multiple nodes. Denote the nodes $\mathcal{N} = \{1, 2, \ldots, d\}$ and directed edges $\mathcal{E}$. Suppose there is a scalar variable x_i associated with each node, and the function f has the form $f(x) = \sum_{(i,j)\in\mathcal{E}} f_{ij}(x_i, x_j)$, then the partial derivative of f with respect to x_i is given by

$$\frac{\partial f}{\partial x_i} = \sum_{j:(i,j)\in\mathcal{E}} \frac{\partial f_{ij}}{\partial x_i}(x_i, x_j) + \sum_{l:(l,i)\in\mathcal{E}} \frac{\partial f_{li}}{\partial x_i}(x_l, x_i)\,.$$

Note that the number of terms in the summations equals the number of edges that touch node i, the expected value of which is about $2/d$ times the total number of edges in the graph. Meanwhile, evaluation of the full gradient would require evaluation of both partial derivatives of each f_{ij} for *all* edges in the graph. Hence, the cost difference between these two operations is a factor of order d.

In this paper, we study how to modify the updating strategies of LMC to reduce the numerical cost, with the focus on reducing dependence on d. In particular, we will develop and analyze a method called Random Coordinate Langevin Monte Carlo (RC-LMC). This idea is inspired by the random coordinate descent (RCD) algorithm from optimization (Nesterov, 2012; Wright, 2015). RCD is a version of Gradient Descent (GD). In each step, one coordinate (or a block of coordinates) is selected at random for updating at each iteration using the associated partial derivative. In optimization, RCD can be significantly cheaper than GD, especially when the objective function is skewed and the dimensionality of the problem is high. In RC-LMC, we use the same basic strategy: At iteration m, a single coordinate of x^m is randomly selected for updating, while all others are left untouched.

Generally speaking, the cost of updating a single coordinate is less than the cost of updating all coordinates, but more iterations are required to achieve the target accuracy. The total cost requires delicate analysis. As for RCD versus GD in optimization, the savings of RC-LMC by comparison with LMC depends strongly on the structure of the dimensional Lipschitz constants. Under an assumption that there is a factor-of-d difference in per-iteration costs, we conclude the following:

1. (Theorem 4.2) When gradient of f is Lipschitz but the Hessian is not, RC-LMC costs $O(d^2/\epsilon^2)$ for an ϵ-accurate solution, and it is cheaper than the classical LMC if f is skewed and the dimension of the problem is high. The optimal numerical cost in this setting is achieved when the probability of choosing the i-th direction is proportional to the i-th directional Lipschitz constant;.

2. (Theorem 4.3) When both gradient and Hessian of f are Lipschitz, RC-LMC requires $O(d^{3/2}/\epsilon)$ iterations to achieve ϵ accuracy. On the other hand, the currently available result indicates that the classical LMC costs $O(d^2/\epsilon)$. Thus, RC-LMC saves a factor of at least $\sqrt{d}$.

3. (Proposition 4.1) The $O(d^{3/2}/\epsilon)$ complexity bound for RC-LMC is sharp when both gradient and Hessian of f are Lipschitz.

We make two additional remarks. (a) Throughout the paper we assume that one element of the gradient is available at an expected cost of approximately $1/d$ of the cost of the full gradient evaluation.

Although this property is intuitive, and often holds in many situations (such as the graph-based example presented above), it might not hold for certain problems (Wright, 2015). (b) Besides replacing gradient evaluation by coordinate algorithms, one might also improve the dimension dependence of LMC algorithm by utilizing a faster converging underlying SDEs than (2). One such example is to use underdamped Langevin dynamics instead, see e.g., (Rossky et al., 1978; Dalalyan and Riou-Durand, 2018; Cheng et al., 2018; Eberle et al., 2019; Shen and Lee, 2019; Cao et al., 2019), which can be also combined with coordinate sampling. For the clarity of presentation, we will focus only on LMC in this work and leave the extension to underdampled samplers for future works.

The paper is organized as follows. We present the new coordinate algorithm in Section 2. Notations and assumptions on f are listed in Section 3, where we also recall theoretical results for the classical LMC method. We present our main results regarding the numerical cost in Section 4 and numerical experiments in Section 5. Proofs of the main results are deferred to the Appendix.

2 RANDOM COORDINATE LANGEVIN MONTE CARLO

We introduce the Random Coordinate Langevin Monte Carlo (RC-LMC) method in this section. At each iteration, one coordinate is chosen at random and updated, while the other components of x are unchanged. Specifically, denoting by r^m the index of the random coordinate chosen at m-th iteration, we obtain $x_{r^m}^{m+1}$ according to a single-coordinate version of (1) and set $x_j^{m+1} = x_j^m$ for $j \neq r^m$.

For generality, we do not necessarily require r^m to be chosen uniformly from $\{1, 2, \ldots, d\}$, but rather denote ϕ_i the probability of component i being chosen. We note the distribution from which r_m is drawn by Φ, where

$$\Phi := \{\phi_1, \phi_2, \ldots, \phi_d\}, \quad \text{where } \phi_i > 0 \text{ for all } i \text{ and } \sum_{i=1}^d \phi_i = 1. \tag{3}$$

The stepsize may depend on the choice of coordinate; we denote the tepsizes by $\{h_1, h_2, \ldots, h_d\}$ and assume that they do not change across iterations. In this paper, we choose h_i to be inversely dependent on probabilities ϕ_i, as follows:

$$h_i = \frac{h}{\phi_i}, \quad i = 1, 2, \ldots, d, \tag{4}$$

where $h > 0$ is a parameter that can be viewed as the expected stepsize. In Section 4.2-4.3, we will find the optimal form of Φ under different scenarios. The initial iterate x^0 is drawn from a distribution q_0, which can be any distribution that is easy to draw from (the normal distribution, for example). We present the complete method in Algorithm 1.

Algorithm 1 Random Coordinate Langevin Monte Carlo (RC-LMC)

Input: Coordinate distribution $\Phi := \{\phi_1, \phi_2, \ldots, \phi_d\}$; parameter $h > 0$ and stepsize set $\{h_1, h_2, \ldots, h_d\}$ defined in (3)–(4); M (stop index).

Sample x^0 from a initial distribution q_0
for $m = 0, 1, 2, \ldots M - 1$ **do**
    1. Draw $r^m \in \{1, \ldots, d\}$ according to probability distribution Φ;
    2. Draw ξ^m from $\mathcal{N}(0, 1)$;
    3. Update x^{m+1} by

$$x_i^{m+1} = \begin{cases} x_i^m - h_i \partial_i f(x^m) + \sqrt{2h_i}\, \xi^m, & i = r^m \\ x_i^m, & i \neq r^m. \end{cases} \tag{5}$$

end for
return x^M

When we compare (5) with the classical LMC (1), we see that in the updating formula, the gradient is replaced by a partial derivative in a random direction r^m:

$$\nabla f(x^m) \to \partial_{r^m} f(x^m) e_{r^m},$$

where e_i is the unit vector for i-th direction.

Define the elapsed time at m-th iteration as

$$T^m := \sum_{n=0}^{m-1} h_{r^n}, \quad \text{and} \quad T^0 := 0, \tag{6}$$

then for $t \in (T^m, T^{m+1}]$, the updating formula (5) can be viewed as the Euler approximation to the following SDE:

$$\begin{cases} X_{r^m}(t) = X_{r^m}(T^m) - \int_{T^m}^t \partial_{r^m} f(X(s)) \, ds + \sqrt{2} \int_{T^m}^t dB_s, \\ X_i(t) = X_i(T^m), \quad \forall i \neq r^m. \end{cases} \tag{7}$$

We note that the SDE preserves the invariant measure, that is, $X(t) \sim p$ for any $t \geq 0$. We discuss further the convergence property of the SDE (7) in Section 4.1.

3 NOTATIONS, ASSUMPTIONS AND CLASSICAL RESULTS

We unify notations and assumptions in this section, and summarize and discuss the classical results on LMC. Throughout the paper, to quantify the distance between two probability distributions, we use the Wasserstein distance defined by

$$W(\mu, \nu) = \left(\inf_{(X,Y) \in \Gamma(\mu,\nu)} \mathbb{E} |X - Y|^2 \right)^{1/2},$$

where $\Gamma(\mu, \nu)$ is the set of distribution of $(X, Y) \in \mathbb{R}^{2d}$ whose marginal distributions, for X and Y respectively, are μ and ν. The distributions in $\Gamma(\mu, \nu)$ are called the *couplings* of μ and ν. Due to the use of power 2 in the definition, this is sometimes called (more specifically) the Wasserstein-2 distance.

We assume that f is strongly convex, so that p is strongly log-concave. We obtain results under two different assumptions: First, Lipschitz continuity of the gradient of f (Assumption 3.1) and second, Lipschitz continuity of the Hessian of f (Assumption 3.2 together with Assumption 3.1).

Assumption 3.1. *The function f is twice differentiable, f is μ-strongly convex for some $\mu > 0$ and its gradient ∇f is L-Lipschitz. That is, for all $x, x' \in \mathbb{R}^d$, we have*

$$f(x) - f(x') - \nabla f(x')^\top (x - x') \geq \frac{\mu}{2} |x - x'|^2, \tag{8}$$

and

$$|\nabla f(x) - \nabla f(x')| \leq L|x - x'|. \tag{9}$$

It is an elementary consequence of (8) that

$$(\nabla f(x') - \nabla f(x))^\top (x' - x) \geq \mu \|x' - x\|^2, \quad \text{for all } x, x' \in \mathbb{R}^d. \tag{10}$$

Since each coordinate direction plays a distinct role in RC-LMC, we distinguish the Lipschitz constants in each such direction. When Assumption 3.1 holds, partial derivatives in all coordinate directions are also Lipschitz. Denoting them as L_i for each $i = 1, 2, \ldots, d$, we have

$$|\partial_i f(x + t\mathbf{e}_i) - \partial_i f(x)| \leq L_i |t| \tag{11}$$

for any $x \in \mathbb{R}^d$ and any $t \in \mathbb{R}$. We further denote $L_{\max} := \max_i L_i$ and define condition numbers as follows:

$$\kappa = L/\mu \geq 1, \quad \kappa_i = L_i/\mu \geq 1, \quad \kappa_{\max} = \max_i \kappa_i. \tag{12}$$

As shown in (Wright, 2015), we have

$$L_i \leq L_{\max} \leq L \leq dL_{\max}, \quad \kappa_i \leq \kappa_{\max} \leq \kappa \leq d\kappa_{\max}. \tag{13}$$

These assumptions together imply that the spectrum of the Hessian is bounded above and below for all x, specifically, $\mu I_d \preceq \nabla^2 f(x) \preceq L I_d$ and $[\nabla^2 f(x)]_{ii} \leq L_i \leq L_{\max}$ for all $x \in \mathbb{R}^d$.

Both upper and lower bounds of L in term of $L_{\max}$ in (13) are tight. If $\nabla^2 f$ is a diagonal matrix, then $L_{\max} = L$, both being the biggest eigenvalue of $\nabla^2 f$. Thus, $\kappa_{\max} = \kappa$ in this case. This is the

case in which all coordinates are independent of each other, for example $f = \sum_i \lambda_i x_i^2$. On the other hand, if $\nabla^2 f = \mathbf{e} \cdot \mathbf{e}^\top$ where $\mathbf{e} \in \mathbb{R}^d$ satisfies $\mathbf{e}_i = 1$ for all i, then $L = dL_{\max}$ and $\kappa = d\kappa_{\max}$. This is a situation in which f is highly skewed, that is, $f = (\sum_i x_i)^2/2$.

The next assumption concerns higher regularity for f.

Assumption 3.2. *The function f is three times differentiable and $\nabla^2 f$ is H-Lipschitz, that is*

$$\|\nabla^2 f(x) - \nabla^2 f(x')\|_2 \le H|x - x'|, \quad \text{for all } x, x' \in \mathbb{R}^d. \tag{14}$$

When this assumption holds, we further define H_i to satisfy

$$|\partial_{ii} f(x + t\boldsymbol{e}_i) - \partial_{ii} f(x)| \le H_i |t|, \tag{15}$$

for any $i = 1, 2, \ldots, d$, all $x \in \mathbb{R}^d$, and all $t \in \mathbb{R}$, where $\partial_{ii} f$ is the (i, i) diagonal entry of the Hessian matrix $\nabla^2 f$.

We summarize existing results for LMC in the following theorem.

Theorem 3.1 ((Durmus et al., 2019, Theorem 9), (Dalalyan and Karagulyan, 2019, Theorem 5)). *Let q_m be the probability distribution of the m-th iteration of LMC (1), and p be the target distribution. Using the notation $W_m := W(q_m, p)$, we have the following.*

- *Suppose that Assumption 3.1 holds and let $h \le 1/L$. Then*

$$W_m \le \exp\left(-\mu h m/2\right) W_0 + 2(\kappa h d)^{1/2}; \tag{16}$$

- *Suppose that Assumptions 3.1 and 3.2 hold and let $h < 2/(\mu + L)$. Then*

$$W_m \le \exp\left(-\mu h m\right) W_0 + \frac{Hhd}{2\mu} + 3\kappa^{3/2}\mu^{1/2}hd^{1/2}. \tag{17}$$

This theorem yields stopping criteria for the number of iterations M to achieve a user-defined accuracy of ϵ. When the gradient of f is Lipschitz, to achieve ϵ-accuracy, we can require both terms on the right hand side of (16) to be smaller than $\epsilon/2$, which occurs when

$$h = \Theta(\epsilon^2/d\kappa), \quad M = \Theta\left(\frac{1}{\mu h} \log\left(\frac{W_0}{\epsilon}\right)\right) = \Theta\left(\frac{d\kappa}{\mu\epsilon^2} \log\left(\frac{W_0}{\epsilon}\right)\right), \tag{18}$$

leading to a cost of $\tilde{O}(d^2\kappa/(\mu\epsilon^2))$ evaluations of gradient components (when we assume that each full gradient can be obtained at the cost of d individual components of the gradient). When both the gradient and the Hessian are Lipschitz, to achieve ϵ-accuracy, we require all three terms on the right hand side of (17) to be smaller than $\epsilon/3$. Assuming $d \gg 1$ and all other constants are $O(1)$, we thus obtain

$$h = \Theta(\epsilon\mu/dH), \quad M = \Theta\left(\frac{dH}{\mu^2\epsilon} \log\left(\frac{W_0}{\epsilon}\right)\right), \tag{19}$$

which yields a cost of $\tilde{O}(d^2 H/(\mu^2\epsilon))$ evaluations of gradient components.

4 MAIN RESULTS

We discuss the main results from two perspectives. In Section 4.1 we examine the convergence of the underlying SDE (7), laying the foundation for the convergence in the discrete setting. We then build upon this result and show the convergence of the RC-LMC algorithm in Section 4.2 and 4.3 under two different assumptions. We show in Section 4.4 that when both Assumption 3.1 and 3.2 are satisfied, our bound is tight with respect to d and ϵ.

4.1 CONVERGENCE OF THE SDE (7)

To study the convergence of (7), we first let $X^m = X(T^m)$ and denote the probability filtration by $\mathcal{F}^m = \left\{x^0, r^{n \le m}, B_{s \le T^m}\right\}$. Then $\{X^m\}_{m=0}^\infty$ is a Markov chain and the following theorem shows its geometric ergodicity.

Theorem 4.1. *Denote by $q_m(x)$ the probability density function of X^m. If f satisfies Assumption 3.1 and $h \leq \frac{\mu \min\{\phi_i\}}{4 + 8L^2 + 32L^4}$, then $p(x)$ is the density of the stationary distribution of the Markov chain $\{X^m\}_{m=0}^{\infty}$. Furthermore, if the second moment of q_0 is finite and X^0 is drawn from q_0, then there are constants $R > 0$ and $r > 1$, independent of m, such that for any $m \geq 0$ we have*

$$\int_{\mathbb{R}^d} |q_m(x) - p(x)| \, \mathrm{d}x \leq R r^{-m} \,. \tag{20}$$

See proof in Appendix A. This theorem states that the solution to the SDE converges to the target distribution. Since the discrepancy between q_m and p decays exponentially in time on the continuous level, the discrete version (as computed in the algorithm) can be expected to converge as well. We will establish this fact in subsequent subsections.

4.2 CONVERGENCE OF RC-LMC. CASE 1: LIPSCHITZ GRADIENT

Under Assumption 3.1, we have the following result. The proof can be found in Appendix B.

Theorem 4.2. *Assume f satisfies Assumption 3.1, and $h_i = h/\phi_i$ with $h \leq \frac{\mu \min\{\phi_i\}}{8L^2}$. Let q_m be the probability distribution of x^m computed in (5), let p be the target distribution, and denote $W_m := W(q_m, p)$. Then we have*

$$W_m \leq \exp\left(-\frac{\mu h m}{4}\right) W_0 + \frac{5h^{1/2}}{\mu} \sqrt{\sum_{i=1}^{d} \frac{L_i^2}{\phi_i}} \,. \tag{21}$$

We make a few comments here: (1) the requirement on h is rather weak. When both μ and L are moderate (both $O(1)$ constants), the requirement is essentially $h \lesssim 1/d$. (1) The estimate (21) consists of two terms. The first is exponentially decaying term and the second comes from the variance of random coordinate selection. If we assume all Lipschitz constants L_i are of $O(1)$, this remainder term is roughly $O(h^{1/2}d)$. (3) The theorem suggests a stopping criterion: to have $W_m \leq \epsilon$, we roughly need $h < \epsilon^2/d^2$, and $m = \widetilde{O}(d^2/\epsilon^2)$, assuming $L_i = O(1)$. In terms of ϵ and d dependence, this puts m at the same order as (18), as required by the classical LMC.

Theorem 4.2 holds for all choices of $\{\phi_i\}$ satisfying (3). From the explicit formula (21) we can choose $\{\phi_i\}$ to minimize the right-hand side of the bound. Nesterov (2012) proposed distributions Φ that depend on the dimensional Lipschitz constants $L_i, i = 1, 2, \ldots, d$ from (11). For $\alpha \in \mathbb{R}$, we can let $\phi_i(\alpha) \propto L_i^\alpha$, specifically,

$$\phi_i(\alpha) := \frac{L_i^\alpha}{\sum_j L_j^\alpha}, \quad \text{and} \quad \Phi(\alpha) := \{\phi_1(\alpha), \phi_2(\alpha), \ldots, \phi_d(\alpha)\} \,. \tag{22}$$

Note that when $\alpha = 0$, $\phi_i(0) = 1/d$ for all i: the uniform distribution among all coordinates. When $\alpha > 0$, the directions that with larger Lipschitz constants have higher probability to be chosen. Since $h_i = h/\phi_i$, one uses smaller stepsizes for stiffer directions. (On the other hand, when $\alpha < 0$, the directions with larger Lipschitz constants are less likely to be chosen, and the stepsizes are larger in stiffer directions, a situation that is not favorable and should be avoided.) Given α, we can define the quantities

$$K_\alpha := \sum_{i=1}^{d} \kappa_i^\alpha \quad \text{so that} \quad K_0 = d \quad \text{and} \quad K_1 = \sum_{i=1}^{d} \kappa_i \,. \tag{23}$$

Note that when the condition number κ_i is $O(1)$ for each $i = 1, 2, \ldots, d$, then K_1 is a quantity of size $O(d)$. The following corollary discusses various choices of α and the corresponding computational cost.

Corollary 4.1. *Under the same conditions as in Theorem 4.2, with $\phi_i = \phi_i(\alpha)$ defined in (22), the number of iterations M required to attain $W_M \leq \epsilon$ is $M = \Theta\left(\frac{K_{2-\alpha} K_\alpha}{\mu \epsilon^2} \log\left(\frac{W_0}{\epsilon}\right)\right)$. The expression in parentheses is optimized when $\alpha = 1$, for which we have*

$$M = \Theta\left(\frac{(\sum_i \kappa_i)^2}{\mu \epsilon^2} \log\left(\frac{W_0}{\epsilon}\right)\right) \,. \tag{24}$$

See proof in Appendix B. We note that the initial error W_0 enters through a $\log$ term and is essentially negligible. To compare RC-LMC with the classical LMC, we compare (24) with (18), adjusting (18) by a factor of d to account for the higher cost per iteration. RC-LMC has more favorable computational cost if $d^2\kappa \geq \left(\sum_i \kappa_i\right)^2$. Since $\kappa_i \leq \kappa_{\max}$, this is guaranteed if $\kappa \geq \kappa_{\max}^2$, which in turn is true when $\kappa \sim d\kappa_{\max}$ and $d > \kappa_{\max}$, that is, for highly skewed f in high dimensional space.

Our proof of Theorem 4.2 follows from a coupling approach similar to that used by Dalalyan and Karagulyan (2019) for LMC. We emphasize that for the coordinate algorithm, we need to overcome the additional difficulty that the process of each coordinate is not contracting (see proof in Appendix C). This is a different situation from the classical LMC (Dalalyan and Karagulyan, 2019), where the SDE (2) has contraction property and we need only to quantify the error caused by discretization. In RC-LMC, we use the contraction property of the algorithm, which ensures that the distance between two different trajectories following the algorithm contracts. However, this contraction property is not component-wise, so we need to choose Young's constant wisely and take summation of every coordinate. The summation will also produce some extra terms, which we need to bound. Dalalyan and Karagulyan (2019) obtain an estimate for the cost of classical LMC of $\widetilde{O}(d^2\kappa^2/(\mu\epsilon^2))$. Compared with this estimate, our estimate for the cost of RC-LMC is always cheaper (since $\kappa^2 \geq \kappa_{\max}^2$). The improved estimate of the cost of LMC (18) was obtained by Durmus et al. (2019) using a quite different approach based on optimal transportation. It is not clear whether their technique can be adapted to the coordinate setting to obtain an improved estimate.

4.3 CONVERGENCE OF RC-LMC. CASE 2: LIPSCHITZ HESSIAN

We now assume that Assumption 3.1 and 3.2 hold, that is, both the gradient and the Hessian of f are Lipschitz continuous. In this setting, we obtain the following improved convergence estimate. The proof can be found in Appendix C.

Theorem 4.3. *Assume f satisfies Assumptions 3.1 and 3.2 and let $h_i = h/\phi_i$, with $h \leq \frac{\mu\min\{\phi_i\}}{8L^2}$ Denoting by $q_m(x)$ the probability density function of x^m computed from (5) and by p the target distribution, and letting $W_m := W(q_m, p)$, we have:*

$$W_m \leq \exp\left(-\frac{\mu hm}{4}\right) W_0 + \frac{3h}{\mu}\sqrt{\sum_{i=1}^{d} \frac{(L_i^3 + H_i^2)}{\phi_i^2}}\,. \tag{25}$$

We see again two terms in the bound, an exponentially decaying term and a variance term. Assuming all Lipschitz constants are $O(1)$, the variance term is of $O(hd^{3/2})$. By comparing with Theorem 4.2, we see that ϵ error can be achieved with the looser stepsize requirement $h \lesssim \frac{\epsilon}{d^{3/2}}$.

By choosing $\{\phi_i\}$ to optimize the bound in Theorem 4.3, we obtain the following corollary.

Corollary 4.2. *Under the same conditions as in Theorem 4.3, the optimal choice of $\{\phi_i\}$ is to set:*

$$\phi_i = \frac{\left(L_i^3 + H_i^2\right)^{1/3}}{\sum_{i=1}^{d}\left(L_i^3 + H_i^2\right)^{1/3}}\,.$$

For this choice, the number of iterations M required to guarantee $W_M \leq \epsilon$ satisfies

$$M = \Theta\left(\frac{\left(\sum_{i=1}^{d}\left(L_i^3 + H_i^2\right)^{1/3}\right)\left(\sum_{i=1}^{d}\left(L_i^3 + H_i^2\right)^{2/3}\right)^{1/2}}{\mu^2\epsilon}\log\left(\frac{W_0}{\epsilon}\right)\right)\,. \tag{26}$$

If μ, κ_i and H_i are all constants of $O(1)$, then the total cost is $\tilde{O}(d^{3/2}/\epsilon)$.

See proof in Appendix C. This is a significant improvement compared to the cost of classical LMC (which requires $\tilde{O}(d^2/\epsilon)$ (Dalalyan and Riou-Durand, 2018)), regardless of the structure of f. Indeed, the cost is reduced by a factor of $d^{1/2}$, which can be significant for high dimensional problems.

4.4 TIGHTNESS OF THE COMPLEXITY BOUND

When both the gradient and Hessian are Lipschitz, we claim that estimate $\tilde{O}(d^{3/2}/\epsilon)$ obtained in Corollary 4.2 is tight. An example is presented in the following proposition.

Proposition 4.1. *Let $\phi_i(\alpha) = 1/d$ for all i, and set the initial distribution and the target distribution to be:*

$$q_0(x) = \frac{1}{(4\pi)^{d/2}} \exp(-|x - \mathsf{e}|^2/4), \quad p(x) = \frac{1}{(2\pi)^{d/2}} \exp(-|x|^2/2), \qquad (27)$$

where $\mathsf{e} \in \mathbb{R}^d$ satisfies $\mathsf{e}_i = 1$ for all i. Let q_m be the probability distribution of x^m generated by Algorithm 1, and denote $W_m := W(q_m, p)$. Then we have

$$W_m \geq \exp\left(-2mh\right)\frac{\sqrt{d}}{3} + \frac{d^{3/2}h}{6}, \quad m \geq 1. \qquad (28)$$

In particular, to have $W_m \leq \epsilon$, one needs at least $m = \widetilde{O}(d^{3/2}/\epsilon)$.

See proof in Appendix D. For the given initial data, we have $W_0 = \sqrt{4 - 2\sqrt{2}} \, d^{1/2}$. To have $W_m \leq \epsilon$, both terms in (28) must be smaller than ϵ. Thus we have $h \lesssim \epsilon/d^{3/2}$, $m \sim \frac{1}{h}\log(\epsilon/\sqrt{d})$, which implies a cost of $\widetilde{O}(d^{3/2}/\epsilon)$, as claimed.

5 NUMERICAL RESULTS

We provide some numerical results in this section. Since it is extremely challenging to estimate the Wasserstein distance between two distributions in high dimensions, we demonstrate instead the convergence of estimated expectation for a given observable. Denoting by $\{x^{(i),M}\}_{i=1}^N$ the list of N samples, with each of them computed through Algorithm 1 independently with M iterations, we define the error as follows:

$$\text{Error}_M = \left| \frac{1}{N} \sum_{i=1}^{N} \psi(x^{(i),M}) - \mathbb{E}_p(\psi) \right|, \qquad (29)$$

where ψ is a test function and $\mathbb{E}_p(\psi)$ is the expectation of ψ under the target distribution p. As $h \to 0$ and $Mh \to \infty$, we have $W_M \to 0$, and $x^{(i),M}$ can be regarded as approximately sampled from p. Thus, according to the central limit theorem, we have $\text{Error}_M = O(1/\sqrt{N})$.

In this example, we set the target and initial distributions to be Gaussian $p(x) \propto p_1(\mathsf{x})p_2(x)$ and $q_0(x) \propto p_1(\mathsf{x} - \mathsf{e})p_2(x)$ with

$$p_1(\mathsf{x}) = \exp\left(-\frac{1}{2}\mathsf{x}\left(\mathsf{T} + (d/10)I\right)^\top \left(\mathsf{T} + (d/10)I\right)\mathsf{x}^\top\right), \quad p_2 = \exp\left(-\frac{1}{2}\sum_{i=11}^{100}|x_i|^2\right),$$

where $\mathsf{x} = (x_1, x_2, \ldots, x_{10})^\top$, $\mathsf{e} = (1, 1, \ldots, 1)^\top \in \mathbb{R}^{10}$, I is the identity matrix and T is a random matrix with each entry i.i.d. drawn from $\mathcal{N}(0, 1)$. We run the simulation with $N = 10^6$, and we compute Error_M with $\psi(x) = \|\mathsf{x}\mathsf{x}^\top\|_2$. This measures the spectral norm of the covariance matrix of the first 10 entries. As shown in Figure 1, RC-LMC with $\alpha = 1$ converges faster than RC-LMC with $\alpha = 0$, and both converge faster than the classical LMC.

Figure 1: The decay of error with respect to the cost (number of ∂f calculations).

REFERENCES

Andrieu, C., Freitas, N., Doucet, A., and Jordan, M. (2003). An introduction to MCMC for Machine Learning. *Machine Learning*, 50:5–43.

Cao, Y., Lu, J., and Wang, L. (2019). On explicit L^2-convergence rate estimate for underdamped langevin dynamics. *arXiv preprint arXiv:1908.04746*.

Casella, G. and George, E. I. (1992). Explaining the gibbs sampler. *The American Statistician*, 46(3):167–174.

Cheng, X., Chatterji, N., Bartlett, P., and Jordan, M. (2018). Underdamped Langevin MCMC: A non-asymptotic analysis. In *Proceedings of the 31st Conference On Learning Theory*, volume 75, pages 300–323.

Dalalyan, A. (2017). Theoretical guarantees for approximate sampling from smooth and log-concave densities. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(3):651–676.

Dalalyan, A. and Karagulyan, A. (2019). User-friendly guarantees for the Langevin Monte Carlo with inaccurate gradient. *Stochastic Processes and their Applications*, 129(12):5278 – 5311.

Dalalyan, A. S. and Riou-Durand, L. (2018). On sampling from a log-concave density using kinetic langevin diffusions. *arXiv*, abs/1807.09382.

Del Moral, P., Doucet, A., and Jasra, A. (2006). Sequential monte carlo samplers. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 68(3):411–436.

Duane, S., Kennedy, A., Pendleton, B. J., and Roweth, D. (1987). Hybrid monte carlo. *Physics Letters B*, 195(2):216 – 222.

Durmus, A., Majewski, S., and Miasojedow, B. (2019). Analysis of langevin monte carlo via convex optimization. *Journal of Machine Learning Research*, 20:73:1–73:46.

Durmus, A. and Moulines, É. (2017). Non-asymptotic convergence analysis for the Unadjusted Langevin Algorithm. *Ann. Appl. Probab.*, 27(3):1551–1587.

Eberle, A., Guillin, A., and Zimmer, R. (2019). Couplings and quantitative contraction rates for Langevin dynamics. *Annals of Probability*, 47(4):1982–2010.

Evensen, G. (2009). *Data Assimilation: The Ensemble Kalman Filter*. Springer-Verlag Berlin Heidelberg.

Fabian, P. (1981). Atmospheric sampling. *Advances in Space Research*, 1(11):17 – 27.

Geman, S. and Geman, D. (1984). Stochastic relaxation, gibbs distributions, and the bayesian restoration of images. *IEEE Trans. Pattern Anal. Mach. Intell.*, 6:721–741.

Geweke, J. (1989). Bayesian inference in econometric models using Monte Carlo integration. *Econometrica*, 57(6):1317–1339.

Hastings, W. (1970). Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, 57(1):97–109.

Iglesias, M., Law, K., and Stuart, A. (2013). Ensemble Kalman methods for inverse problems. *Inverse Problems*, 29(4):045001.

Li, R., Pei, S., Chen, B., Song, Y., Zhang, T., Yang, W., and Shaman, J. (2020). Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (sars-cov-2). *Science*, 368(6490):489–493.

Markowich, P. and Villani, C. (1999). On the trend to equilibrium for the Fokker-Planck equation: An interplay between physics and functional analysis. In *Physics and Functional Analysis, Matematica Contemporanea (SBM) 19*, pages 1–29.

Mattingly, J., Stuart, A., and Higham, D. (2002). Ergodicity for sdes and approximations: locally lipschitz vector fields and degenerate noise. *Stochastic Processes and their Applications*, 101(2):185 – 232.

Metropolis, N., Rosenbluth, A., Rosenbluth, M., Teller, A., and Teller, E. (1953). Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21(6):1087–1092.

Nagarajan, N., Honarpour, M., and Sampath, K. (2007). Reservoir-fluid sampling and characterization — key to efficient reservoir management. *Journal of Petroleum Technology*, 59.

Neal, R. M. (1993). Probabilistic inference using Markov Chain Monte Carlo methods. *Technical Report CRG-TR-93-1. Dept. of Computer Science, University of Toronto.*

Neal, R. M. (2001). Annealed importance sampling. *Statistics and Computing*, 11:125–139.

Nesterov, Y. (2012). Efficiency of coordinate descent methods on huge-scale optimization problems. *SIAM Journal on Optimization*, 22(2):341–362.

Parisi, G. (1981). Correlation functions and computer simulations. *Nuclear Physics B*, 180(3):378–384.

Reich, S. (2011). A dynamical systems framework for intermittent data assimilation. *BIT Numerical Mathematics*, 51(1):235–249.

Roberts, G. and Rosenthal, J. (2004). General state space Markov chains and MCMC algorithms. *Probability Surveys*, 1.

Roberts, G. and Tweedie, R. (1996). Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2(4):341–363.

Rossky, P. J., Doll, J. D., and Friedman, H. L. (1978). Brownian dynamics as smart monte carlo simulation. *The Journal of Chemical Physics*, 69(10):4628–4633.

Russo, D., Roy, B., Kazerouni, A., Osband, I., and Wen, Z. (2018). A tutorial on Thompson sampling. *Foundations and Trends in Machine Learning*, 11(1):1–96.

Shen, R. and Lee, Y. T. (2019). The randomized midpoint method for log-concave sampling. In *Advances in Neural Information Processing Systems*, pages 2100–2111.

Vempala, S. (2010). Recent progress and open problems in algorithmic convex geometry. In *IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science*, volume 8, pages 42–64.

Wright, S. J. (2015). Coordinate descent algorithms. *Mathematical Programming, Series B*, 151(1):3–34.

A PROOF OF THEOREM 4.1

We recall the SDE (7):

$$\begin{cases} X_{r^m}(t) = X_{r^m}(T^m) - \int_{T^m}^t \partial_{r^m} f(X(s)) \, \mathrm{d}s + \sqrt{2} \int_{T^m}^t \mathrm{d}B_s \,, \\ X_i(t) = X_i(T^m) \,, \quad \forall i \neq r^m \,, \end{cases} \tag{30}$$

where r^m is randomly selected from $1, \ldots, d$. Moreover, recall that $X^{m+1} = X\left(T^{m+1}\right)$ is a Markovian process. We denote its transition kernel by Ξ, meaning that

$$X^{m+1} \overset{d}{=} \Xi(X^m, \cdot) \,.$$

Moreover, we denote Ξ^n the n-step transition kernel. The following proposition establishes the exponential convergence of the Markov chain.

Proposition A.1. *Under conditions of Theorem 4.1, there are constants $R_1 > 0, r_1 > 1$, such that for any $x^0 \in \mathbb{R}^d$*

$$\sup_{A \in \mathcal{B}(\mathbb{R}^d)} \left| \Xi^{md}(x^0, A) - \int_A p(x) \, \mathrm{d}x \right| \leq \left(|x^0 - x^*|^2 + 1 \right) R_1 r_1^{-m} \,, \tag{31}$$

where x^* is the minimal point of $f(x)$ and Ξ is the transition kernel for $\{X^m\}_{m=0}^\infty$.

We postpone the proof of Proposition A.1 to Section A.1. Now, we are ready to prove the theorem.

Proof of Theorem 4.1. First, suppose the distribution of X^m is induced by p. Then for $i \neq r^m$, the distribution of $X_i(t)$ between $[T^m, T^{m+1}]$ is preserved. Meanwhile, we have

$$\mathrm{d}X_{r^m} = -\partial_{r^m} f(X(s)) \, \mathrm{d}t + \sqrt{2} \, \mathrm{d}B_s \,,$$

and the marginal distribution of $X_{r^m}(t)$ is also preserved. Therefore $X^{m+1} \sim p$, proving that $p(x)$ is the density of the stationary distribution.

Second, to prove (20), let $x^0 \sim q_0$ that has finite second moment, we multiply q_0 on both sides of (31) and integrate, to obtain

$$\int_{\mathbb{R}^d} |q_{md}(x) - p(x)| \, \mathrm{d}x \leq C_0 r_1^{-m} \,,$$

where C_0 is a constant.

By using (30) with Itô's formula, we have

$$\frac{\mathrm{d}\mathbb{E}|X_{r^m}(t)|^2}{\mathrm{d}t} = -2\mathbb{E}\left(\partial_{r^m} f(X_{r^m}(t)) X_{r^m}(t) \right) + 2 \leq 2 + \mathbb{E}|\partial_{r^m} f(X_{r^m}(t))|^2 + \mathbb{E}|X_{r^m}(t)|^2$$

$$\leq 2 + L_{r^m}^2 \mathbb{E}|X_{r^m}(t) - x_{r^m}^*|^2 + \mathbb{E}|X_{r^m}(t)|^2 \leq C_{1,r^m} \mathbb{E}|X_{r^m}(t)|^2 + C_{2,r^m} \,,$$

where C_{1,r^m} and C_{2,r^m} are constants that depend only on x^* and L_{r^m}. From Grönwall's inequality, we obtain

$$\mathbb{E}\left(|X_i^{m+1}|^2 \big| r^m = i \right) \leq \exp(C_{1,i} h_i) \left[\mathbb{E}(|X_i^m|^2) + C_{2,i} h_i \right] \,, \quad \text{for all } i = 1, 2, \ldots, d.$$

Then, if $\mathbb{E}|X^m|^2 < \infty$, we have for any $i = 1, 2, \ldots, d$ that

$$\mathbb{E}\left(|X_i^{m+1}|^2 \right) = \frac{1}{d} \mathbb{E}\left(|X_i^{m+1}|^2 \big| r^m = i \right) + \left(1 - \frac{1}{d} \right) \mathbb{E}\left(|X_i^{m+1}|^2 \big| r^m \neq i \right)$$

$$\leq \frac{1}{d} \exp(C_{1,i} h_i) \left[\mathbb{E}(|X_i^m|^2) + C_{2,i} h_i \right] + \left(1 - \frac{1}{d} \right) \mathbb{E}(|X_i^m|^2) < \infty \,,$$

which implies $\mathbb{E}|X^{m+1}|^2 < \infty$. Therefore, if q_0 has finite second moment, then q_i all have finite second moments for $i = 1, \ldots, d-1$.

Letting $x^0 \sim q_i$, multiplying q_i on both sides of (31) and integrating, we obtain

$$\int_{\mathbb{R}^d} |q_{md+i}(x) - p(x)| \, \mathrm{d}x \leq C_i r_1^{-m} \,,$$

where C_i is a constant. Since this bound holds for all $0 \leq i \leq d-1$, we set $R = (\max_i C_i) r_1$ and $r = r_1^{1/d}$ to obtain (20). $\qquad \square$

A.1 PROOF OF PROPOSITION A.1

Before we prove the Proposition, we first recall a result from (Mattingly et al., 2002) for the convergence of Markov chain using Lyapunov condition together with minorization condition.

Theorem A.1. *[(Mattingly et al., 2002, Theorem 2.5)] Let $\{X^n\}_{n=0}^\infty$ denote the Markov chain on $\mathbb{R}^d$ with transition kernel Ξ and filtration $\mathcal{F}^n$. Let $\{X^n\}_{n=0}^\infty$ satisfy the following two conditions:*

Lyapunov condition: There is a function $L : \mathbb{R}^d \to [1, \infty)$, with $\lim_{x \to \infty} L(x) = \infty$, and real numbers $\alpha \in (0, 1)$, and $\beta \in [0, \infty)$ such that

$$\mathbb{E}\left(L(X^{n+1}) \big| \mathcal{F}^n\right) \leq \alpha L(X^n) + \beta \,.$$

Minorization condition: For L from the Lyqpunov condition, define the set $C \subset \mathbb{R}^d$ as follows:

$$C = \left\{ x \in \mathbb{R}^d \mid L(x) \leq \frac{2\beta}{\gamma - \alpha} \right\} \,, \tag{32}$$

for some $\gamma \in (\alpha^{1/2}, 1)$. Then there exists an $\eta > 0$ and a probability measure $\mathcal{M}$ supported on C (that is, $\mathcal{M}(C) = 1$), such that

$$\Xi(x, A) \geq \eta \mathcal{M}(A), \quad \forall A \in \mathcal{B}(\mathbb{R}^d),\ x \in C \,.$$

Under these conditions, the Markov chain $\{X^n\}_{n=0}^\infty$ has a unique invariant measure π. Furthermore, there are constants $r \in (0, 1)$ and $R \in (0, \infty)$ such that, for any $x_0 \in \mathbb{R}^d$, we have

$$\sup_{A \in \mathcal{B}(\mathbb{R}^d)} \left| \Xi^n(x^0, A) - \pi(A) \right| \leq L(x_0) R r^{-n} \,. \tag{33}$$

To use this result to prove Proposition A.1, we will consider the d-step chain of $\{X^n\}$ and verify the two conditions, as in the following two lemmas for the Lyapunov function and the minorization over a small set, respectively.

Lemma A.1. *Assume f satisfies Assumption 3.1 and*

$$h \leq \frac{\mu \min\{\phi_i\}}{4 + 8L^2 + 32L^4} \,, \tag{34}$$

where L is the Lipschitz constant defined in (9). Let the Lyapunov function be $L(x) = |x - x^*|^2 + 1$, then we have:

$$\mathbb{E}\left(L(X^{m+1}) \big| \mathcal{F}^m\right) \leq \alpha_1 L(X^m) + \beta_1 \tag{35}$$

with

$$\alpha_1 = 1 - \mu h \,, \quad \beta_1 = (24 + 120L^2 + \mu)h \,.$$

Lemma A.2. *Under conditions of Lemma A.1, with $L(x) = |x - x^*|^2 + 1$, let Ξ denote the transition kernel. Define the set $C \subset \mathbb{R}^d$ as in (32), for some $\gamma \in (\alpha^{1/2}, 1)$. Then there exists an $\eta > 0$ and a probability measure $\mathcal{M}$ with $\mathcal{M}(C) = 1$, such that*

$$\Xi^d(x, A) \geq \eta \mathcal{M}(A), \quad \forall A \in \mathcal{B}(\mathbb{R}^d),\ x \in C \,. \tag{36}$$

Proposition A.1 follows easily from these results.

Proof of Proposition A.1. It suffices to show d-step chain $\left\{X^{md}\right\}_{m=0}^\infty$ satisfies the conditions in Theorem A.1 with $L(x) = |x - x^*|^2 + 1$, $\alpha = \alpha_1^d$ and $\beta = d\beta_1$, and π is induced by p. We apply (35) from Lemma A.1 iteratively, d times, to obtain

$$\mathbb{E}\left(L\left(X^{(m+1)d}\right) \big| \mathcal{F}^{md}\right) \leq \alpha_1^d L\left(X^{md}\right) + d\beta_1 \,,$$

which implies that $\left\{X^{md}\right\}_{m=0}^\infty$ satisfies Lyapunov condition in Theorem A.1 with $\alpha = \alpha_1^d$. Moreover, Lemma A.2 directly implies that the d-step transition kernel satisfies the minorization condition. Therefore, by Theorem A.1, we have

$$\sup_{A \in \mathcal{B}(\mathbb{R}^d)} \left| \Xi^{md}(x^0, A) - \pi(A) \right| \leq L(x_0) R r^{-m} \,,$$

which concludes the proof of the proposition when we substitute $\pi(A) = \int_A p(x)\,\mathrm{d}x$. $\qquad\square$

Proof of Lemma A.1. We assume without loss of generality that $x^* = 0 \in \mathbb{R}^d$ (so that $L(x) = |x|^2 + 1$) and drop the filtration $\mathcal{F}^m$ in the formula for simplicity of notation. Then

$$\mathbb{E}\left(L\left(X^{m+1}\right)\right) = \sum_{i=1}^{d} \phi_i \mathbb{E}\left(L\left(X^{m+1}\right)\big|r^m = i\right). \tag{37}$$

Since

$$L\left(X^{m+1}\right) = |X^{m+1}|^2 + 1 = |X^m + (X^{m+1} - X^m)|^2 + 1$$
$$= L\left(X^m\right) + 2X^m(X^{m+1} - X^m) + |X^{m+1} - X^m|^2,$$

we have

$$\mathbb{E}\left(L\left(X^{m+1}\right)\big|r^m = i\right) = L\left(X^m\right) + 2\mathbb{E}\left[X_i^m\left(X_i^{m+1} - X_i^m\right)\big|r^m = i\right]$$
$$+ \mathbb{E}\left[\left(X_i^{m+1} - X_i^m\right)^2\big|r^m = i\right]. \tag{38}$$

To deal with second term and third term in (38), we first note that, under condition $r^m = i$:

$$X_i^{m+1} - X_i^m = -\int_{T^m}^{T^m + h_i} \partial_i f(X(s))\,\mathrm{d}s + \sqrt{2}\int_{T^m}^{T^m + h_i} \mathrm{d}B_s. \tag{39}$$

This means

$$2\mathbb{E}\left[X_i^m\left(X_i^{m+1} - X_i^m\right)\big|r^m = i\right]$$
$$= -2\mathbb{E}\left[X_i^m\int_{T^m}^{T^m + h_i} \partial_i f(X(s))\,\mathrm{d}s\bigg|r^m = i\right] \tag{40}$$
$$= -2h_i X_i^m \partial_i f(X^m) - 2\mathbb{E}\left[X_i^m\int_{T^m}^{T^m + h_i} (\partial_i f(X(s)) - \partial_i f(X^m))\,\mathrm{d}s\bigg|r^m = i\right].$$

We further bound the second term of (40):

$$\left|\mathbb{E}\left[X_i^m\int_{T^m}^{T^m + h_i} (\partial_i f(X(s)) - \partial_i f(X^m))\,\mathrm{d}s\bigg|r^m = i\right]\right|$$
$$\leq h_i \mathbb{E}\left[X_i^m\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t)) - \partial_i f(X^m)|\right)\bigg|r^m = i\right]$$
$$\overset{(\mathrm{I})}{\leq} 2h_i^2 |X_i^m|^2 + 2\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t)) - \partial_i f(X^m)|^2\bigg|r^m = i\right) \tag{41}$$
$$\overset{(\mathrm{II})}{\leq} 2h_i^2 |X_i^m|^2 + 2L_i^2 \mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |X_i(t) - X_i^m|^2\bigg|r^m = i\right)$$
$$\overset{(\mathrm{III})}{\leq} 2h_i^2 |X_i^m|^2 + 16h_i^2 L_i^2 |\partial_i f(X^m)|^2 + 60h_i L_i^2$$
$$\overset{(\mathrm{IV})}{\leq} (2 + 16L_i^4)h_i^2 |X_i^m|^2 + 60h_i L_i^2,$$

where we used Young's inequality in (I), the Lipschitz condition in (II), Lemma A.3 below (specifically, inequality (44)) in (III), and the Lipschitz condition again in (IV). This, when substituted into (40), gives

$$2\mathbb{E}\left[X_i^m\left(X_i^{m+1} - X_i^m\right)\big|r^m = i\right] \leq -2h_i X_i^m \partial_i f(X^m) + (4 + 32L_i^4)h_i^2 |X_i^m|^2 + 120h_i L_i^2.$$

To bound the third term in (38), again for the case $r^m = i$, we use (39) again for:

$$
\mathbb{E}\left[\left(X_i^{m+1} - X_i^m\right)^2 \middle| r^m = i\right]
$$

$$
= \mathbb{E}\left[\left(\int_{T^m}^{T^m+h_i} \partial_i f(X(s))\,\mathrm{d}s - \sqrt{2}\int_{T^m}^{T^m+h_i}\,\mathrm{d}B_s\right)^2 \middle| r^m = i\right]
$$

$$
\overset{(\mathrm{I})}{\leq} 2h_i^2\,\mathbb{E}\left(\sup_{T^m \leq t \leq T^m+h_i}|\partial_i f(X(t))|^2 \middle| r^m = i\right) + 4\mathbb{E}\left(\left|\int_{T^m}^{T^m+h_i}\,\mathrm{d}B_s\right|^2 \middle| r^m = i\right) \tag{42}
$$

$$
= 2h_i^2\,\mathbb{E}\left(\sup_{T^m \leq t \leq T^m+h_i}|\partial_i f(X(t))|^2 \middle| r^m = i\right) + 4h_i
$$

$$
\overset{(\mathrm{II})}{\leq} 8h_i^2|\partial_i f(X^m)|^2 + 88h_i^3 L_i^2 + 4h_i
$$

$$
\overset{(\mathrm{III})}{\leq} 8L_i^2 h_i^2|X_i^m|^2 + 24h_i\,,
$$

where we used Young's inequality in (I), Lemma A.3 below (specifically, inequality (43)) in (II), and Lipschitz continuity in (III), together with $88h_i^2 L_i^2 \leq 20$ by (34).

Finally, we have

$$
\mathbb{E}\left(L\left(X^{m+1}\right)\middle| r^m = i\right) \leq L\left(X^m\right) - 2h_i X_i^m \partial_i f(X^m) + (4 + 8L_i^2 + 32L_i^4)h_i^2|X_i^m|^2 + (24 + 120L_i^2)h_i\,.
$$

By summing according to (37), and using (4) and $L_i \leq L$ for all $i = 1, 2, \ldots, d$, we obtain

$$
\mathbb{E}\left(L\left(X^{m+1}\right)\right) = \sum_{i=1}^{d}\phi_i\,\mathbb{E}\left(L\left(X^{m+1}\right)\middle| r^m = i\right)
$$

$$
\leq L\left(X^m\right) - 2h\left\langle X^m, \nabla f(X^m)\right\rangle + \frac{\left(4 + 8L^2 + 32L^4\right)h^2}{\min\{\phi_i\}}(L\left(X^m\right) - 1) + (24 + 120L^2)h\,.
$$

Finally, using $\left\langle X^m, \nabla f(X^m)\right\rangle \geq \mu(L\left(X^m\right) - 1)$ (from (10) with $x' = X^m$ and $x = x^* = 0$) and (34), we obtain (35). $\qquad\square$

Proof of Lemma A.2. To prove (36), we construct a new Markov process $\widetilde{X}^m$. Defining $\widetilde{X}^0 = x^0$, we obtain $\widetilde{X}^{m+1}$ from $\widetilde{X}^m$ by running the following process:

$$
\widetilde{T}^n = \sum_{i=1}^{n} h_i,\quad \widetilde{T}^0 = 0,\quad Z(0) = \widetilde{X}^m\,.
$$

Then for $\widetilde{T}^{n-1} \leq t \leq \widetilde{T}^n$ and $n \leq d$, let

$$
\begin{cases}
Z_n(t) = Z_n\left(\widetilde{T}^{n-1}\right) - \int_{\widetilde{T}^{n-1}}^{t}\partial_n f\left(Z(s)\right)\,\mathrm{d}s + \sqrt{2}\int_{\widetilde{T}^{n-1}}^{t}\,\mathrm{d}B_s\,, \\
Z_i(t) = Z_i\left(\widetilde{T}^{n-1}\right),\quad i \neq n\,,
\end{cases}
$$

and set $\widetilde{X}^{m+1} = Z\left(\widetilde{T}^d\right)$. Denote the transition kernel by Ξ_{cyc} (corresponding to one round of a cyclic version of the coordinate algorithm). We then have the following properties:

- For any $x \in C$ and $A \in \mathcal{B}(\mathbb{R}^d)$, we have

$$
\Xi^d(x, A) \geq \Pi_{i=1}^{d}\phi_i \Xi_{\mathrm{cyc}}(x, A) > 0\,.
$$

- Ξ_{cyc} possesses a positive jointly continuous density.

According to (Mattingly et al., 2002, Lemma 2.3), since Ξ_{cyc} has a positive jointly continuous density, there exists an $\eta' > 0$ and a probability measure $\mathcal{M}$ with $\mathcal{M}(C) = 1$, such that

$$
\Xi_{\mathrm{cyc}}(x, A) > \eta'\mathcal{M}(A),\quad \forall A \in \mathcal{B}\left(\mathbb{R}^d\right), x \in C\,,
$$

which implies

$$\Xi^d(x, A) \geq \Pi_{i=1}^d \phi_i \Xi_{\mathrm{cyc}}(x, A) > \Pi_{i=1}^d \phi_i \eta' \mathcal{M}(A), \quad \forall A \in \mathcal{B}\left(\mathbb{R}^d\right), x \in C \,.$$

This proves (36) by setting $\eta = \Pi_{i=1}^d \phi_i \eta'$. $\qquad\square$

In the proof of Lemma A.1, we used several estimates in inequalities (41) and (42). We prove these estimates in the following lemma.

Lemma A.3. *Suppose that the assumptions of Lemma A.1 hold, and let X_i evolve according to* (39). *Then we have the following bounds:*

$$\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t))|^2\right) \leq 4|\partial_i f(X^m)|^2 + 44 h_i L_i^2 \,, \tag{43}$$

$$\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |X_i(t) - X_i^m|^2\right) \leq 8 h_i^2 |\partial_i f(X^m)|^2 + 30 h_i \,. \tag{44}$$

Proof. To obtain (43), we have

$$
\begin{aligned}
&\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t))|^2\right)\\
&\leq \mathbb{E}\left[\sup_{T^m \leq t \leq T^m + h_i} \left(|\partial_i f(X^m)| + L_i |X_i(t) - X_i^m|\right)^2\right]\\
&\leq 2|\partial_i f(X^m)|^2 + 2 L_i^2 \mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |X_i(t) - X_i^m|^2\right) \,.
\end{aligned}
\tag{45}
$$

To bound the second term, we use (39) again:

$$
\begin{aligned}
&\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |X_i(t) - X_i^m|^2\right)\\
&= \mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} \left|\int_{T^m}^t \partial_i f\left(X(s)\right) \, \mathrm{d}s - \sqrt{2}\int_{T^m}^t \mathrm{d}B_s\right|^2\right)\\
&\leq 2 h_i^2 \mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t))|^2\right) + 4\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} \left|\int_{T^m}^t \mathrm{d}B_s\right|^2\right)\\
&\leq 2 h_i^2 \mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t))|^2\right) + 16 h_i \,,
\end{aligned}
\tag{46}
$$

where we use Young's inequality and

$$\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} \left|\int_{T^m}^t \mathrm{d}B_s\right|^2\right) \leq 4\mathbb{E}\left(\left|\int_{T^m}^{T^m + h_i} \mathrm{d}B_s\right|^2\right) = 4 h_i$$

by Doob's maximal inequality. By substituting (46) into (45), we obtain

$$
\begin{aligned}
&\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t))|^2\right)\\
&\leq 4 h_i^2 L_i^2 \mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t))|^2\right) + 2|\partial_i f(X^m)|^2 + 32 h_i L_i^2 \,.
\end{aligned}
$$

Using $h_i L_i \leq \frac{1}{4}$, we move the first term on the right to the left to obtain

$$\frac{3}{4}\mathbb{E}\left(\sup_{T^m \leq t \leq T^m + h_i} |\partial_i f(X(t))|^2\right) \leq 2|\partial_i f(X^m)|^2 + 32 h_i L_i^2 \,,$$

leading to (43). Then we obtain (44) by plugging this in (46) and using the fact that $88 h_i^3 L_i^2 < 14 h_i$ by (34). $\qquad\square$

B   PROOF OF THEOREM 4.2

The proof of this theorem requires us to design a reference solution to explicitly bound $W(q_m, p)$. Let $\tilde{x}^0$ be a random vector drawn from target distribution induced by p, so that $W_2^2(q_0, p) = \mathbb{E}|x^0 - \tilde{x}^0|^2$. We then require $\tilde{x}$ to solve the following SDE: for $t \in (T^m, T^{m+1}]$, with T^m defined in (6):

$$\begin{cases} \tilde{x}_{r^m}(t) = \tilde{x}_{r^m}(T^m) - \int_{T^m}^t \partial_{r^m} f(\tilde{x}(s)) \, \mathrm{d}s + \sqrt{2} \int_{T^m}^t \mathrm{d}B_s \,, \\ \tilde{x}_i(t) = \tilde{x}_i(T^m), \quad i \neq r^m \,. \end{cases} \tag{47}$$

If we use the same Brownian motion as in (5), we have

$$\tilde{x}^{m+1} = \tilde{x}^m + \left[-\int_{T^m}^{T^{m+1}} \partial_{r^m} f(\tilde{x}(s)) \, \mathrm{d}s + \sqrt{2h_{r^m}} \xi^m \right] \boldsymbol{e}_{r^m} \,, \tag{48}$$

where $\boldsymbol{e}_{r^m}$ is the unit vector in r^m direction. Since the r^m-th marginal distribution of $\tilde{x}(t)$ is preserved in each time step according to (47), the whole distribution of $\tilde{x}(t)$ is preserved to be p for all t. Therefore, by the definition $W_m = W(q_m, p)$, we have

$$W_m^2 \leq \mathbb{E}|\Delta^m|^2 = \mathbb{E}|x^m - \tilde{x}^m|^2 \,,$$

where

$$\Delta^m := \tilde{x}^m - x^m \,. \tag{49}$$

This means bounding W_m amounts to evaluating $\mathbb{E}|\Delta^m|^2$. Under Assumption 3.1, we have the following result.

Proposition B.1. *Suppose the assumptions of Theorem 4.2 are satisfied and let $\{x^m\}$, $\{\tilde{x}^m\}$, and $\{\Delta^m\}$ be defined in* (5), (47), *and* (49), *respectively. Then, we have*

$$\mathbb{E}|\Delta^{m+1}|^2 \leq \left(1 - \frac{h\mu}{2}\right) \mathbb{E}|\Delta^m|^2 + \frac{10h^2}{\mu} \sum_{i=1}^d \frac{L_i^2}{\phi_i} \,. \tag{50}$$

The proof of this result appears in Appendix B.1. The proof for Theorem 4.2 is now immediate.

Proof of Theorem 4.2. By iterating (50), we obtain

$$\mathbb{E}|\Delta^m|^2 \leq \left(1 - \frac{h\mu}{2}\right)^m \mathbb{E}|\Delta^0|^2 + \frac{20h}{\mu^2} \sum_{i=1}^d \frac{L_i^2}{\phi_i} \,,$$

and since $h\mu/2 \in (0, 1)$, we have

$$\mathbb{E}|\Delta^m|^2 \leq \exp\left(-\frac{\mu h m}{2}\right) \mathbb{E}|\Delta^0|^2 + \frac{20h}{\mu^2} \sum_{i=1}^d \frac{L_i^2}{\phi_i} \,. \tag{51}$$

By construction, we have $W^2(q_0, p) = \mathbb{E}|\Delta^0|^2$ and $W^2(q_m, p) \leq \mathbb{E}|\Delta^m|^2$. By taking the square root of both sides and using $a^2 \leq b^2 + c^2 \Rightarrow a \leq b + c$ for any nonnegative a, b, and c, we arrive at (21). □

The proof for Corollary 4.1 is also obvious.

Proof of Corollary 4.1. To ensure that $W_m \leq \epsilon$, we set the two terms on the right hand side of (21) to be smaller than $\epsilon/2$, which implies that

$$h = O\left(\frac{\mu^2 \epsilon^2}{100 \sum_{i=1}^d \frac{L_i^2}{\phi_i(\alpha)}}\right) \quad \text{and} \quad m \geq \frac{4}{\mu h} \log\left(\frac{2W_0}{\epsilon}\right) \,. \tag{52}$$

By using the definition of $\phi_i(\alpha)$ according to (22), we obtain

$$\sum_{i=1}^d \frac{L_i^2}{\phi_i(\alpha)} = \left(\sum_{i=1}^d \frac{L_i^2}{L_i^\alpha}\right) \left(\sum_{j=1}^d L_j^\alpha\right) = \mu^2 K_{2-\alpha} K_\alpha \,,$$

which implies that $m = \tilde{O}\left((K_{2-\alpha}K_\alpha)/(\mu\epsilon^2)\right)$. Furthermore, $\alpha = 1$ gives the optimal cost, because:

$$K_{2-\alpha}K_\alpha = \left(\sum \kappa_i^\alpha\right)\left(\sum \kappa_i^{2-\alpha}\right) \geq \left(\sum_i \kappa_i\right)^2 = K_1^2\,,$$

due to Hölder's inequality. $\qquad\square$

B.1 PROOF OF PROPOSITION B.1

We prove the Proposition by means of the following lemma.

Lemma B.1. *Under the conditions of Proposition B.1, for $m \geq 0$ and $i = 1, 2, \ldots, d$, we have*

$$\mathbb{E}|\Delta_i^{m+1}|^2 \leq \left(1 + h\mu + \frac{h^2\mu^2}{\phi_i}\right)\mathbb{E}|\Delta_i^m|^2 - 2h\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right]$$
$$+ \frac{3h^2}{\phi_i}\mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2 + \left(\frac{2h^3 L_i^3}{\mu\phi_i^2} + \frac{8h^2 L_i^2}{\mu\phi_i}\right)\,. \tag{53}$$

Proof. In the m-th time step, we have

$$\mathbb{P}(r^m = i) = \phi_i, \quad \mathbb{P}(r^m \neq i) = 1 - \phi_i\,,$$

so that

$$\mathbb{E}|\Delta_i^{m+1}|^2 = \phi_i\mathbb{E}\left(|\Delta_i^{m+1}|^2 \mid r^m = i\right) + (1 - \phi_i)\mathbb{E}\left(|\Delta_i^{m+1}|^2 \mid r^m \neq i\right)$$
$$= \phi_i\mathbb{E}\left(|\Delta_i^{m+1}|^2 \mid r^m = i\right) + (1 - \phi_i)\mathbb{E}|\Delta_i^m|^2\,. \tag{54}$$

We now analyze the first term on the right hand side under condition $r^m = i$. By definition of Δ_i^{m+1}, we have

$$\Delta_i^{m+1} = \Delta_i^m + (\tilde{x}_i^{m+1} - \tilde{x}_i^m) - (x_i^{m+1} - x_i^m)$$
$$= \Delta_i^m + \left(-\int_{T^m}^{T^m+h_i}\partial_i f(\tilde{x}(s))\,\mathrm{d}s + \sqrt{2h_i}\xi_m\right) - \left(-\int_{T^m}^{T^m+h_i}\partial_i f(x^m)\,\mathrm{d}s + \sqrt{2h_i}\xi_m\right)$$
$$= \Delta_i^m - \int_{T^m}^{T^m+h_i}\left(\partial_i f(\tilde{x}(s)) - \partial_i f(x^m)\right)\,\mathrm{d}s$$
$$= \Delta_i^m - \int_{T^m}^{T^m+h_i}\left(\partial_i f(\tilde{x}(s)) - \partial_i f(\tilde{x}^m) + \partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\,\mathrm{d}s$$
$$= \Delta_i^m - h_i\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right) - \int_{T^m}^{T^m+h_i}\left(\partial_i f(\tilde{x}(s)) - \partial_i f(\tilde{x}^m)\right)\,\mathrm{d}s$$
$$= \Delta_i^m - h_i\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right) - V^m\,, \tag{55}$$

where we have defined

$$V^m := \int_{T^m}^{T^m+h_i}\left(\partial_i f(\tilde{x}(s)) - \partial_i f(\tilde{x}^m)\right)\,\mathrm{d}s\,. \tag{56}$$

By Young's inequality, we have

$$\mathbb{E}\left(|\Delta_i^{m+1}|^2 \mid r^m = i\right)$$
$$= \mathbb{E}\left(|\Delta_i^{m+1} + V^m - V^m|^2 \mid r^m = i\right)$$
$$\leq (1 + a)\mathbb{E}\left(|\Delta_i^{m+1} + V^m|^2 \mid r^m = i\right) + \left(1 + \frac{1}{a}\right)\mathbb{E}\left(|V^m|^2 \mid r^m = i\right)\,, \tag{57}$$

where $a > 0$ is a parameter to be specified later.

For the first term on the right hand side of (57), we have

$$\mathbb{E}\left(|\Delta_i^{m+1} + V^m|^2 \mid r^m = i\right)$$
$$= \mathbb{E}|\Delta_i^m - h_i\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)|^2$$
$$= \mathbb{E}|\Delta_i^m|^2 - 2h_i\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right] + h_i^2\mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2\,. \tag{58}$$

Note that the second term will essentially become the second line in (53), and the third term will become the third line in (53) (upon the proper choice of a). For very small h, this term is negligible.

For the second term on the right-hand side of (57), we recall the definition (56) and obtain

$$
\begin{aligned}
\mathbb{E}\left(|V^m|^2\big|r^m = i\right) &\overset{(I)}{\leq} h_i \int_{T^m}^{T^m + h_i} \mathbb{E}\left(|\partial_i f(\tilde{x}(s)) - \partial_i f(\tilde{x}^m)|^2 \big| r^m = i\right) \mathrm{d}s \\
&\overset{(II)}{\leq} h_i L_i^2 \int_{T^m}^{T^m + h_i} \mathbb{E}\left(|\tilde{x}(s) - \tilde{x}^m|^2 \big| r^m = i\right) \mathrm{d}s \\
&= h_i L_i^2 \int_{T^m}^{T^m + h_i} \mathbb{E}\left(\left|\int_{T^m}^{s} \partial_i f(\tilde{x}(t))\,\mathrm{d}t + \sqrt{2}(B_s - B_{T^m})\right|^2 \bigg| r^m = i\right) \mathrm{d}s \\
&\overset{(III)}{\leq} 2h_i^2 L_i^2 \int_{T^m}^{T^m + h_i} \int_{T^m}^{s} \mathbb{E}\left(|\partial_i f(\tilde{x}(t))|^2 \big| r^m = i\right) \mathrm{d}t\,\mathrm{d}s \\
&\quad + 4h_i^2 L_i^2 \int_{T^m}^{T^m + h_i} \mathbb{E}|\xi^m|^2\,\mathrm{d}s \\
&\overset{(IV)}{=} h_i^4 L_i^2 \mathbb{E}\left(|\partial_i f(\tilde{x}^m)|^2\right) + 4h_i^3 L_i^2 \\
&\overset{(V)}{=} h_i^4 L_i^2 \mathbb{E}_p|\partial_i f|^2 + 4h_i^3 L_i^2 \overset{(VI)}{\leq} h_i^4 L_i^3 + 4h_i^3 L_i^2\,,
\end{aligned}
\tag{59}
$$

where (II) comes from L-Lipschitz condition (11), (I) and (III) come from the use of Young's inequality and Jensen's inequality when we move the $|\cdot|^2$ from outside to inside of the integral, and (IV) and (V) hold true because $\tilde{x}(t) \sim p$ for all t. In (VI) we use $\mathbb{E}_p|\partial_i f|^2 \leq L_i$ using (Dalalyan and Karagulyan, 2019, Lemma 3).

By substituting (58) and (59) into the right hand side of (57), we obtain

$$
\begin{aligned}
\mathbb{E}&\left(|\Delta_i^{m+1}|^2 \mid r^m = i\right) \\
&\leq (1+a)\mathbb{E}|\Delta_i^m|^2 - 2h_i(1+a)\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right] \\
&\quad + h_i^2(1+a)\mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2 + \left(1 + \frac{1}{a}\right)\left(h_i^4 L_i^3 + 4h_i^3 L_i^2\right).
\end{aligned}
\tag{60}
$$

By substituting (60) into (54), we have

$$
\begin{aligned}
\mathbb{E}|\Delta_i^{m+1}|^2 &\leq (1 + a\phi_i)\,\mathbb{E}|\Delta_i^m|^2 - 2(1+a)h\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right] \\
&\quad + \frac{(1+a)h^2}{\phi_i}\mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2 + \left(1 + \frac{1}{a}\right)\left(\frac{h^4 L_i^3}{\phi_i^3} + \frac{4h^3 L_i^2}{\phi_i^2}\right),
\end{aligned}
\tag{61}
$$

where we have used $h_i \phi_i = h$.

Now, we need to choose a value of $a > 0$ appropriate to establish (53). By comparing the two formulas, we see the need to set

$$
a\phi_i = h\mu \quad \Rightarrow \quad a = h_i \mu = \frac{h\mu}{\phi_i} \leq 1\,.
$$

since $h \leq \min\{\phi_i\}/\mu$. It follows that $1 + \frac{1}{a} \leq \frac{2\phi_i}{h\mu}$. By substituting into (61), we obtain

$$
\begin{aligned}
\mathbb{E}|\Delta_i^{m+1}|^2 &\leq (1 + h\mu)\,\mathbb{E}|\Delta_i^m|^2 - 2h\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right] \\
&\quad - \frac{2h^2\mu}{\phi_i}\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right] + \frac{2h^2}{\phi_i}\mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2 \\
&\quad + \left(\frac{2h^3 L_i^3}{\mu\phi_i^2} + \frac{8h^2 L_i^2}{\mu\phi_i}\right).
\end{aligned}
\tag{62}
$$

We conclude the lemma by using the following Cauchy-Schwartz inequality to control the third term on the right hand side of this expression:

$$
-\frac{2h^2\mu}{\phi_i}\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right] \leq \frac{h^2\mu^2}{\phi_i}\mathbb{E}|\Delta_i^m|^2 + \frac{h^2}{\phi_i}\mathbb{E}|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)|^2\,. \quad \square
$$

Proposition B.1 is obtained by simply summing all components in the lemma.

Proof of Proposion B.1. Noting

$$\mathbb{E}|\Delta^{m+1}|^2 = \sum_{i=1}^{d} \mathbb{E}|\Delta_i^{m+1}|^2,$$

we bound the right hand side by (53) and get

$$\mathbb{E}|\Delta^{m+1}|^2 \leq \left(1 + h\mu + \frac{h^2\mu^2}{\min\{\phi_i\}}\right)\mathbb{E}|\Delta^m|^2 - 2h\mathbb{E}\left\langle\Delta^m, \nabla f(\tilde{x}^m) - \nabla f(x^m)\right\rangle$$

$$+ \frac{3h^2}{\min\{\phi_i\}}\mathbb{E}\left|\nabla f(\tilde{x}^m) - \nabla f(x^m)\right|^2 + \left(\frac{2h^3}{\mu}\sum_{i=1}^{d}\frac{L_i^3}{\phi_i^2} + \frac{8h^2}{\mu}\sum_{i=1}^{d}\frac{L_i^2}{\phi_i}\right). \tag{63}$$

The second and third terms on the right-hand side can be bounded in terms of $\mathbb{E}|\Delta^m|^2$:

- By convexity, we have

$$\mathbb{E}\left\langle\Delta^m, \nabla f(\tilde{x}^m) - \nabla f(x^m)\right\rangle \geq \mu\mathbb{E}|\Delta^m|^2. \tag{64}$$

- As the gradient is L-Lipschitz, we have

$$\mathbb{E}\left|\nabla f(\tilde{x}^m) - \nabla f(x^m)\right|^2 \leq L^2\mathbb{E}|\Delta^m|^2. \tag{65}$$

By substituting (64) and (65) into (63) and using $\mu \leq L$, we obtain

$$\mathbb{E}|\Delta^{m+1}|^2 \leq \left(1 - h\mu + \frac{4h^2L^2}{\min\{\phi_i\}}\right)\mathbb{E}|\Delta^m|^2 + \left(\frac{2h^3}{\mu}\sum_{i=1}^{d}\frac{L_i^3}{\phi_i^2} + \frac{8h^2}{\mu}\sum_{i=1}^{d}\frac{L_i^2}{\phi_i}\right). \tag{66}$$

If we take h sufficiently small, the coefficient in front of $\mathbb{E}|\Delta^m|^2$ is strictly smaller than 1, ensuring the decay of the error. Indeed, by setting $h \leq \frac{\mu\min\{\phi_i\}}{8L^2}$, we have

$$\frac{4h^2L^2}{\min\{\phi_i\}} \leq \frac{h\mu}{2}, \quad \text{and} \quad \frac{hL_i}{\phi_i} \leq \frac{\mu}{8L} \leq 1,$$

which leads to the iteration formula (50). $\qquad\square$

C PROOF OF THEOREM 4.3

Theorem 4.3 is based on the following proposition.

Proposition C.1. *Suppose the assumptions of Theorem 4.3 and let $\{x^m\}$, $\{\tilde{x}^m\}$, and $\{\Delta_m\}$ be defined as in* (5), (47), *and* (49), *respectively. Then we have*

$$\mathbb{E}|\Delta^{m+1}|^2 \leq \left(1 - \frac{h\mu}{2}\right)\mathbb{E}|\Delta^m|^2 + \frac{4h^3}{\mu}\sum_{i=1}^{d}\frac{(L_i^3 + H_i^2)}{\phi_i^2}. \tag{67}$$

We prove this result in Appendix C.1. The proof of the theorem is now immediate.

Proof of Theorem 4.3. Use (67) iteratively, we have

$$\mathbb{E}|\Delta^{m+1}|^2 \leq \left(1 - \frac{h\mu}{2}\right)^m \mathbb{E}|\Delta^0|^2 + \frac{8h^2}{\mu^2}\sum_{i=1}^{d}\frac{(L_i^3 + H_i^2)}{\phi_i^2}$$

$$\leq \exp\left(-\frac{\mu h m}{2}\right)\mathbb{E}|\Delta^0|^2 + \frac{8h^2}{\mu^2}\sum_{i=1}^{d}\frac{(L_i^3 + H_i^2)}{\phi_i^2}.$$

Using $W^2(q_0, p) = \mathbb{E}|\Delta^0|^2$ and $W^2(q_m, p) \leq \mathbb{E}|\Delta^m|^2$, we take the square root on both sides, we obtain (25). $\qquad\square$

The proof of Corollary 4.2 is also immediate.

Proof of Corollary 4.2. Use (25), to ensure $W_m \leq \epsilon$, we set two terms on the right hand side of (25) to be smaller than $\epsilon/2$, which implies that

$$
h = O\left(\frac{\epsilon\mu}{\sqrt{\sum_{i=1}^{d} \frac{(L_i^3 + H_i^2)}{\phi_i^2}}}\right), \quad m \geq \frac{4}{\mu h} \log\left(\frac{2W_0}{\epsilon}\right). \tag{68}
$$

To find optimal choice of ϕ_i, we need to minimize

$$
\sum_{i=1}^{d} \frac{(L_i^3 + H_i^2)}{\phi_i^2}
$$

under constraint $\sum_i^d \phi_i = 1$ and $\phi_i > 0$. Introducing a Lagrange multiplier $\lambda \in \mathbb{R}$, define the Lagrangian function as follows:

$$
F(\phi_1, \phi_2, \ldots, \phi_d, \lambda) = \sum_{i=1}^{d} \frac{(L_i^3 + H_i^2)}{\phi_i^2} + \lambda\left(\sum_{i=1}^{d} \phi_i - 1\right).
$$

By setting $\partial F/\partial \phi_i = 0$ for all i, and substituting into the constraint $\sum_i^d \phi_i = 1$ to find the appropriate value of λ, we find that the optimal $(\phi_1, \phi_2, \ldots, \phi_d)$ satisfies

$$
\phi_i = \frac{(L_i^3 + H_i^2)^{1/3}}{\sum_{i=1}^{d} (L_i^3 + H_i^2)^{1/3}}, \quad i = 1, 2, \ldots, d.
$$

By substituting into (68), we obtain (26). $\square$

C.1 PROOF OF PROPOSITION C.1

The strategy of the proof for this proposition is almost identical to that of the previous section. The reference solution $\tilde{x}$ is defined as in (47). We will use the following lemma:

Lemma C.1. *Under the conditions of Proposition C.1, for $m \geq 0$ and $i = 1, 2, \ldots, d$, we have*

$$
\mathbb{E}|\Delta_i^{m+1}|^2 \leq \left(1 + h\mu + \frac{h^2\mu^2}{\phi_i}\right)\mathbb{E}|\Delta_i^m|^2 - 2h\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right]
$$
$$
+ \frac{3h^2}{\phi_i}\mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2 + \frac{4h^3\left(L_i^3 + H_i^2\right)}{\phi_i^2 \mu}. \tag{69}
$$

Proof. In the m-th time step, we have

$$
\mathbb{P}(r^m = i) = \phi_i, \quad \mathbb{P}(r^m \neq i) = 1 - \phi_i,
$$

meaning that

$$
\mathbb{E}|\Delta_i^{m+1}|^2 = \phi_i \mathbb{E}\left(|\Delta_i^{m+1}|^2 \mid r^m = i\right) + (1 - \phi_i)\mathbb{E}\left(|\Delta_i^{m+1}|^2 \mid r^m \neq i\right)
$$
$$
= \phi_i \mathbb{E}\left(|\Delta_i^{m+1}|^2 \mid r^m = i\right) + (1 - \phi_i)\mathbb{E}|\Delta_i^m|^2. \tag{70}
$$

To bound the first term in (54) we use the definition of Δ_i^{m+1}. Under the condition $r^m = i$, we have, with the same derivation as in (55):

$$
\Delta_i^{m+1} = \Delta_i^m - h_i\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right) - \int_{T^m}^{T^m + h_i}\left(\partial_i f(\tilde{x}(s)) - \partial_i f(\tilde{x}^m)\right)\,\mathrm{d}s \tag{71}
$$
$$
= \Delta_i^m - h_i\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right) - V^m,
$$

where we denoted $V^m = \int_{T^m}^{T^m + h_i}\left(\partial_i f(\tilde{x}(s)) - \partial_i f(\tilde{x}^m)\right)\,\mathrm{d}s$.

However, different from (59), since f has higher regularity, we can find a tighter bound for the integral. Denote

$$U^m = \int_{T^m}^{T^m+h_i} \left(\partial_i f(\tilde{x}(s)) - \partial_i f(\tilde{x}^m) - \sqrt{2} \int_{T^m}^s \partial_{ii} f(\tilde{x}(z)) \, \mathrm{d}B_z \right) \mathrm{d}s \tag{72}$$

and

$$\Phi^m = \sqrt{2} \int_{T^m}^{T^m+h_i} \int_{T^m}^s \partial_{ii} f(\tilde{x}(z)) \, \mathrm{d}B_z \, \mathrm{d}s \,. \tag{73}$$

Then (71) can be written as

$$\Delta_i^{m+1} = \Delta_i^m - h_i \left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m) \right) - \Phi^m - U^m \,, \tag{74}$$

which implies, according to Young's inequality, that, for any a:

$$\mathbb{E}\left(|\Delta_i^{m+1}|^2 \big| r^m = i \right) = \mathbb{E}\left(|\Delta_i^{m+1} + U^m - U^m|^2 \big| r^m = i \right)$$
$$\leq (1+a)\mathbb{E}\left(|\Delta_i^{m+1} + U^m|^2 \big| r^m = i \right) + \left(1 + \frac{1}{a} \right) \mathbb{E}\left(|U^m|^2 \big| r^m = i \right) \,. \tag{75}$$

Both terms on the right-hand side of (75) are small. We now control the first term. Plug in the definition (74), we have:

$$\mathbb{E}\left(|\Delta_i^{m+1} + U^m|^2 \mid r^m = i \right) = \mathbb{E}\left(|\Delta_i^m - h_i \left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m) \right) - \Phi^m|^2 \big| r^m = i \right) \,. \tag{76}$$

Noting that

$$\mathbb{E}\left((\Delta_i^m - h_i \left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m) \right)) \cdot \Phi^m \right)$$
$$= \sqrt{2} \int_{T^m}^{T^m+h_i} \mathbb{E}\left[\int_{T^m}^s \left(\Delta_i^m - h_i \left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m) \right) \right) \cdot \partial_{ii} f(\tilde{x}(z)) \, \mathrm{d}B_z \right] \mathrm{d}s = 0$$

because

$$\mathbb{E}\left[\int_{T^m}^s \left(\Delta_i^m - h_i \left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m) \right) \right) \cdot \partial_{ii} f(\tilde{x}(z)) \, \mathrm{d}B_z \right] = 0 \,,$$

according to the property of Itô's integral, we can discard the cross terms with Φ^m in (76) to obtain

$$\mathbb{E}\left(|\Delta_i^{m+1} + U^m|^2 \mid r^m = i \right) = \mathbb{E}|\Delta_i^m|^2 - 2h_i \mathbb{E}\left[\Delta_i^m \left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m) \right) \right]$$
$$+ h_i^2 \mathbb{E}\left| \partial_i f(\tilde{x}^m) - \partial_i f(x^m) \right|^2 + \mathbb{E}\left(|\Phi^m|^2 \big| r^m = i \right) \,. \tag{77}$$

For the last term of (77), we have the following control:

$$\mathbb{E}\left(|\Phi^m|^2 \big| r^m = i \right) = \mathbb{E}\left(2 \left| \int_{T^m}^{T^m+h_i} \int_{T^m}^s \partial_{ii} f(\tilde{x}(z)) \, \mathrm{d}B_z \, \mathrm{d}s \right|^2 \bigg| r^m = i \right)$$

$$\overset{(\mathrm{I})}{\leq} 2\mathbb{E}\left[\left(\int_{T^m}^{T^m+h_i} \mathrm{d}s \right) \left(\int_{T^m}^{T^m+h_i} \left| \int_{T^m}^s \partial_{ii} f(\tilde{x}(z)) \, \mathrm{d}B_z \right|^2 \mathrm{d}s \right) \bigg| r^m = i \right]$$

$$\leq 2h_i \int_{T^m}^{T^m+h_i} \mathbb{E}\left(\left| \int_{T^m}^s \partial_{ii} f(\tilde{x}(z)) \, \mathrm{d}B_z \right|^2 \bigg| r^m = i \right) \mathrm{d}s$$

$$\overset{(\mathrm{II})}{=} 2h_i \int_{T^m}^{T^m+h_i} \int_{T^m}^s \mathbb{E}\left(|\partial_{ii} f(\tilde{x}(z))|^2 \big| r^m = i \right) \mathrm{d}z \, \mathrm{d}s$$

$$\overset{(\mathrm{III})}{=} h_i^3 \mathbb{E}_p |\partial_{ii} f|^2 = h_i^3 L_i^2 \,,$$

where we use Hölder's inequality in I and $\tilde{x}(t) \sim p$ for all t in III. In II, we use the following property of Itô's integral:

$$\mathbb{E}\left(\left| \int_{T^m}^s \partial_{ii} f(\tilde{x}(z)) \, \mathrm{d}B_z \right|^2 \bigg| r^m = i \right) = \int_{T^m}^s \mathbb{E}\left(|\partial_{ii} f(\tilde{x}(z))|^2 \big| r^m = i \right) \mathrm{d}z \,.$$

By substituting into (77), we obtain

$$
\begin{aligned}
\mathbb{E}\left(|\Delta_i^{m+1} + U^m|^2 \mid r^m = i\right) \leq & \mathbb{E}|\Delta_i^m|^2 - 2h_i \mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right] \\
& + h_i^2 \mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2 + h_i^3 L_i^2
\end{aligned}
\tag{78}
$$

To bound the second term on the right-hand side of (75), we first note that f is three times continuously differentiable, and (15) implies $\|\partial_{iii} f\|_\infty \leq H_i$. Take $\mathrm{d}t$ on both sides of (47), under condition $r^m = i$, we first have

$$
\mathrm{d}\tilde{x}_i(t) = -\partial_i f(\tilde{x}(s))\,\mathrm{d}s + \sqrt{2}\,\mathrm{d}B_s\,.
\tag{79}
$$

According to Itô's formula, we obtain

$$
\partial_i f(\tilde{x}(t)) - \partial_i f(\tilde{x}^m) = \int_{T^m}^t \partial_{ii} f(\tilde{x}(s))\,\mathrm{d}\tilde{x}_i(s) + \int_{T^m}^t \partial_{iii} f(\tilde{x}(s))\,\mathrm{d}s\,.
\tag{80}
$$

Substituting (79) into (80), we have

$$
\begin{aligned}
& \partial_i f(\tilde{x}(t)) - \partial_i f(\tilde{x}^m) - \sqrt{2}\int_{T_m}^t \partial_{ii} f(\tilde{x}(s))\,\mathrm{d}B_s \\
& = \int_{T^m}^t -\partial_{ii} f(\tilde{x}(s))\partial_i f(\tilde{x}(s)) + \partial_{iii} f(\tilde{x}(s))\,\mathrm{d}s\,.
\end{aligned}
\tag{81}
$$

By substituting into (72), we obtain

$$
\begin{aligned}
& \mathbb{E}\left(|U^m|^2 \mid r^m = i\right) \\
& \overset{(\mathrm{I})}{\leq} h_i \int_{T^m}^{T^m + h_i} \mathbb{E}\left(\left|\partial_i f(\tilde{x}(s)) - \partial_i f(\tilde{x}^m) - \sqrt{2}\int_{T^m}^s \partial_{ii} f(\tilde{x}(z))\,\mathrm{d}B_r\right|^2 \middle| r^m = i\right)\,\mathrm{d}s \\
& \overset{(\mathrm{II})}{=} q h_i \int_{T^m}^{T^m + h_i} \mathbb{E}\left(\left|\int_{T^m}^s \left(-\partial_{ii} f(\tilde{x}(z))\partial_i f(\tilde{x}(z)) + \partial_{iii} f(\tilde{x}(z))\right)\,\mathrm{d}z\right|^2 \middle| r^m = i\right)\,\mathrm{d}s \\
& \overset{(\mathrm{III})}{\leq} h_i^2 \int_{T^m}^{T^m + h_i} \int_{T^m}^s \mathbb{E}\left(\left|\partial_{ii} f(\tilde{x}(z))\partial_i f(\tilde{x}(z)) + \partial_{iii} f(\tilde{x}(z))\right|^2 \middle| r^m = i\right)\,\mathrm{d}z\,\mathrm{d}s \\
& \overset{(\mathrm{IV})}{\leq} 2h_i^2 \int_{T^m}^{T^m + h_i} \int_{T^m}^s \mathbb{E}\left(\left|\partial_{ii} f(\tilde{x}(z))\partial_i f(\tilde{x}(z))\right|^2 \middle| r^m = i\right)\,\mathrm{d}z\,\mathrm{d}s \\
& \quad + 2h_i^2 \int_{T^m}^{T^m + h_i} \int_{T^m}^s \mathbb{E}\left(\left|\partial_{iii} f(\tilde{x}(z))\right|^2 \middle| r^m = i\right)\,\mathrm{d}z\,\mathrm{d}s \\
& \overset{(\mathrm{V})}{\leq} h_i^4\left(L_i^3 + H_i^2\right)\,.
\end{aligned}
\tag{82}
$$

In the derivation, (II) comes from plugging in (81), and (I) and (III) come from the use of Jensen's inequality, (V) comes from the use of Lipschitz continuity in the first and the second derivative ((11) and (15) in particular), and the fact that $\tilde{x}(t) \sim p$ for all t. Note also $\mathbb{E}_p|\partial_i f|^2 \leq L_i$ by (Dalalyan and Karagulyan, 2019, Lemma 3).

By plugging (78) and (82) into (70) and (75), we obtain

$$
\begin{aligned}
& \mathbb{E}|\Delta_i^{m+1}|^2 \\
& \leq (1 + a\phi_i)\,\mathbb{E}|\Delta_i^m|^2 - 2(1 + a)h\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right] \\
& \quad + \frac{(1 + a)h^2}{\phi_i}\mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2 + \frac{(1 + a)h^3 L_i^2}{\phi_i^2} + \left(1 + \frac{1}{a}\right)\frac{h^4\left(L_i^3 + H_i^2\right)}{\phi_i^3}\,,
\end{aligned}
\tag{83}
$$

where we use $h_i \phi_i = h$. Comparing with (69), we need to set

$$
a = h_i \mu = \frac{h\mu}{\phi_i} < 1\,,
$$

where we use $h < \frac{\mu \min\{\phi_i\}}{8L^2}$. This leads to $1 + \frac{1}{a} \leq \frac{2\phi_i}{h\mu}$. By substituting into (61), we obtain

$$\mathbb{E}|\Delta_i^{m+1}|^2 \leq \left(1 + h\mu + \frac{h^2\mu^2}{\phi_i}\right) \mathbb{E}|\Delta_i^m|^2 - 2h\mathbb{E}\left[\Delta_i^m\left(\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right)\right]$$
$$+ \frac{3h^2}{\phi_i}\mathbb{E}\left|\partial_i f(\tilde{x}^m) - \partial_i f(x^m)\right|^2 + \frac{2h^3 L_i^2}{\phi_i^2} + \frac{2h^3\left(L_i^3 + H_i^2\right)}{\phi_i^2\mu}.$$

Noting $L_i/\mu > 1$, we conclude the lemma. $\qquad\square$

The proof of Proposition C.1 is obtained by summing up all components and applying Lemma C.1.

Proof of Proposition C.1. Noting that

$$\mathbb{E}|\Delta^{m+1}|^2 = \sum_{i=1}^{d} \mathbb{E}|\Delta_i^{m+1}|^2,$$

we substitute using (69) to obtain

$$\mathbb{E}|\Delta^{m+1}|^2 \leq \left(1 + h\mu + \frac{h^2\mu^2}{\min\{\phi_i\}}\right) \mathbb{E}|\Delta^m|^2 - 2h\mathbb{E}\left\langle \Delta^m, \nabla f(\tilde{x}^m) - \nabla f(x^m)\right\rangle$$
$$+ \frac{3h^2}{\min\{\phi_i\}}\mathbb{E}\left|\nabla f(\tilde{x}^m) - \nabla f(x^m)\right|^2 + \frac{4h^3}{\mu}\sum_{i=1}^{d}\frac{\left(L_i^3 + H_i^2\right)}{\phi_i^2}. \tag{84}$$

The second and third terms in the right-hand side of this bound can be controlled by $\mathbb{E}|\Delta^m|^2$, as follows. By convexity, we have

$$\mathbb{E}\left\langle \Delta^m, \nabla f(\tilde{x}^m) - \nabla f(x^m)\right\rangle \geq \mu\mathbb{E}|\Delta^m|^2. \tag{85}$$

By the L-Lipschitz property, we have

$$\mathbb{E}\left|\nabla f(\tilde{x}^m) - \nabla f(x^m)\right|^2 \leq L^2\mathbb{E}|\Delta^m|^2. \tag{86}$$

By substituting (85) and (86) into (63), and using $\mu < L$, we have

$$\mathbb{E}|\Delta^{m+1}|^2 \leq \left(1 - h\mu + \frac{4h^2 L^2}{\min\{\phi_i\}}\right) \mathbb{E}|\Delta^m|^2 + \frac{4h^3}{\mu}\sum_{i=1}^{d}\frac{\left(L_i^3 + H_i^2\right)}{\phi_i^2}. \tag{87}$$

Since $h < \frac{\mu \min\{\phi_i\}}{8L^2}$, we obtain (67). $\qquad\square$

D PROOF OF PROPOSITION 4.1

Proof of Proposition 4.1. For this special target distribution p, the objective function is $f(x) = \sum_{i=1}^{d}\frac{|x_i|^2}{2}$. With $\alpha = 0$ and $\phi_i = 1/d$, we have: $x_i^{m+1} = x_i^m$ for all $i \neq r^m$ and

$$x_{r^m}^{m+1} = (1 - dh)x_{r^m}^m + \sqrt{2dh}\xi^m.$$

Therefore for all $i = 1, 2, \ldots, d$, we have

$$\mathbb{E}|x_i^{m+1}|^2 = \frac{1}{d}\mathbb{E}\left(|x_i^{m+1}|^2 \big| r^m = i\right) + \left(1 - \frac{1}{d}\right)\mathbb{E}\left(|x_i^{m+1}|^2 \mid r^m \neq i\right)$$
$$= \frac{1}{d}\mathbb{E}\left(|(1 - dh)x_i^m + \sqrt{2dh}\xi^m|^2 \big| r^m = i\right) + \left(1 - \frac{1}{d}\right)\mathbb{E}\left(|x_i^m|^2\right)$$
$$= \left(1 - 2h + dh^2\right)\mathbb{E}|x_i^m|^2 + 2h \tag{88}$$

where we use $\mathbb{E}_\xi\left|x_i^m - dhx_i^m + \sqrt{2dh}\xi^m\right|^2 = (1 - dh)^2|x_i^m|^2 + 2dh$ in the last equation. By summing (88) over i, we obtain

$$\mathbb{E}|x^{m+1}|^2 = \left(1 - 2h + dh^2\right)\mathbb{E}|x^m|^2 + 2dh.$$

Using it iteratively, and considering $\mathbb{E}|x^0|^2 = 3d$, we have:

$$
\begin{aligned}
\mathbb{E}|x^m|^2 &\geq 3d\left(1 - 2h + dh^2\right)^m + \left(1 - \left(1 - 2h + dh^2\right)^m\right)\frac{2dh}{2h - dh^2} \\
&= d\left(1 - 2h + dh^2\right)^m + \frac{2d}{2 - dh} + 2d\left(1 - \frac{1}{2 - dh}\right)\left(1 - 2h + dh^2\right)^m \\
&\geq d\left(1 - 2h\right)^m + \frac{2d}{2 - dh}\,,
\end{aligned}
$$

where we use $dh \leq 1$ in the last inequality.

Since

$$
W(q_m, p) \geq \left(\int |x|^2 q_m(x)\,\mathrm{d}x\right)^{1/2} - \left(\int |x|^2 p(x)\,\mathrm{d}x\right)^{1/2} = \left(\int |x|^2 q_m(x)\,\mathrm{d}x\right)^{1/2} - \sqrt{d}\,,
$$

we have

$$
\begin{aligned}
W(q_m, p) \geq \left(\int |x|^2 q_m(x)\,\mathrm{d}x\right)^{1/2} - \sqrt{d} &\geq \frac{d\left(1 - 2h\right)^m + \frac{2d}{2 - dh} - d}{\sqrt{d\left(1 - 2h\right)^m + \frac{2d}{2 - dh}} + \sqrt{d}} \\
&\geq \frac{\sqrt{d}}{3}\left(1 - 2h\right)^m + \frac{d^{3/2}h}{6} \\
&\geq \exp\left(-2mh\right)\frac{\sqrt{d}}{3} + \frac{d^{3/2}h}{6}\,,
\end{aligned}
$$

where in the last inequality we use

$$
\sqrt{d\left(1 - 2h\right)^m + \frac{2d}{2 - dh}} + \sqrt{d} \leq 3\sqrt{d}.
$$

Therefore, we finally prove (28). $\qquad\square$