Polynomial time guarantees
for the Burer-Monteiro method

Abstract

The Burer-Monteiro method is one of the most widely used techniques for solving large-scale semidefinite programs (SDP). The basic idea is to solve a nonconvex program in Y, where Y is an $n \times p$ matrix such that $X = YY^T$. We show that this method can solve SDPs in polynomial time in a smoothed analysis setting. More precisely, we consider an SDP whose domain satisfies some compactness and smoothness assumptions, and slightly perturb the cost matrix and the constraints. We show that if $p \gtrsim \sqrt{2(1+\eta)m}$, where m is the number of constraints and $\eta > 0$ is any fixed constant, then the Burer-Monteiro method can solve SDPs to any desired accuracy in polynomial time, in the setting of smooth analysis. The bound on p approaches the celebrated Barvinok-Pataki bound in the limit as η goes to zero, beneath which it the nonconvex program can be suboptimal. Our main technical contribution, which is key for our tight bound on p, is to connect spurious approximately critical points of the nonconvex program to tubular neighborhoods of certain algebraic varieties, and then estimate the volume of such tubes.

1 Introduction

Let $\mathbb{S}^n$ be the set of $n \times n$ symmetric matrices. Given matrices $A_1, \ldots, A_m, C \in \mathbb{S}^n$ and a vector $b \in \mathbb{R}^m$, consider the *semidefinite program* (SDP):

$$\min_{X \in \mathbb{S}^n} C \bullet X \quad \text{s.t.} \quad \mathcal{A}(X) = b, \quad X \succeq 0, \tag{SDP}$$

where $\mathcal{A} : \mathbb{S}^n \to \mathbb{R}^m$, $X \mapsto (A_1 \bullet X, \ldots, A_m \bullet X)$ is a linear map and the notation $X \succeq 0$ in indicates that X is a positive semidefinite (PSD) matrix. Also consider the *least squares* SDP

$$\min_{X \in \mathbb{S}^n} \|\mathcal{A}(X) - b\|^2 \quad \text{s.t.} \quad X \succeq 0, \tag{SDP_{ls}}$$

Though interior point methods solve (SDP) and (SDP_{ls}) in polynomial time, they often have memory problems for large values of n. The Burer-Monteiro method [11, 12] is one of the most widely used procedures for large scale problems. Several papers have worked in understanding the practical success of this method, e.g., [9, 10, 12]. Although various results have been shown, they all fall short of showing that one can reach an approximately optimal solution of (SDP)/(SDP_{ls}) in polynomial time. This paper provides the first polynomial time guarantees for the Burer-Monteiro method.

The Burer-Monteiro method consists in writing $X = YY^T$ for some $Y \in \mathbb{R}^{n \times p}$, and solving a nonconvex optimization problem in terms of Y. For (SDP) we get the problem

$$\min_{Y \in \mathfrak{M}} C \bullet YY^T, \qquad \mathfrak{M} := \{Y \in \mathbb{R}^{n \times p} : \mathcal{A}(YY^T) = b\}, \tag{BM}$$

and for (SDP_{ls}) we get the unconstrained problem

$$\min_{Y \in \mathbb{R}^{n \times p}} \|\mathcal{A}(YY^T) - b\|^2. \tag{BM_{ls}}$$

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

We focus now on problem (*SDP*), though our results apply to (*SDP$_{ls}$*) as well. Let $\tau(k) := \binom{k+1}{2}$ be the k-th triangular number. It is known that problems (*SDP*) and (*BM*) have the same optimal value for any p such that $\tau(p) \geq m$; this is known as the *Barvinok-Pataki bound* [3, 29]. But due to nonconvexity, local optimization methods may converge to a critical point of (*BM*) which is not globally optimal [34]. In this paper we are mainly interested in 2nd-order critical points (abbreviated as 2-critical points).

Problem (*BM*) has been well studied, most remarkably in the sequel of papers [5, 9, 10, 30]. Boumal et al. [9, 10] showed that (*BM*) has no spurious 2-critical points when $\tau(p) > m$, assuming that the feasible set $\mathfrak{M}$ is a smooth manifold and that the cost matrix C is *generic*. Hence, the Burer-Monteiro method converges to the global optimal solution as long as the cost C is outside a certain "bad" set, which has measure zero. However, this analysis only applies to the limit points. The follow up works [5] and [30] provide finite-iteration guarantees, though for values of p larger than the Barvinok-Pataki bound. They introduce a smoothed analysis [32] setting in which the cost matrix C is subject to a small random perturbation of magnitude σ. Such a perturbation ensures that C stays outside the "bad" region, so the limit points are globally optimal. They then argue that certain notions of approximate criticality for (*BM*) guarantee approximate optimality for (*SDP*) when $\tau(p) \gtrsim \frac{9}{2} m \log(\Omega(\sqrt{n}/\sigma))$.

Despite the earlier work, the question of whether the Burer-Monteiro method can be used to completely solve problem (*SDP*) in polynomial time has not been yet resolved. The works [5, 30] do not provide end-to-end polynomial time guarantees and make structural assumptions on the problem, as will be further discussed in the related work section. In particular, finding a feasible solution to (*SDP*) that could serve as initial point is a complicated and practically relevant problem, but it has not been addressed before. In addition, the lower bound on p is qualitatively wrong, as it is not only larger than the Barvinok-Pataki bound, but also gets worse as the magnitude σ of the perturbation decreases.

In order to fully solve (*SDP*), we propose a 2-stage procedure. First, solve (*BM$_{ls}$*) to either get a point Y_0 that is approximately feasible, i.e,. $\mathcal{A}(Y_0 Y_0^T) \approx b$, or to prove that no such point exists. Second, solve (*BM*) to get an approximately optimal point, starting with point Y_0 from the first stage. We prove that this scheme is successful when $\tau(p) > (1+\eta)m$, for an arbitrary constant $\eta > 0$.

The next theorem provides polynomial time guarantees for the second stage. The complexity is with respect to the setting of a smoothed analysis, in which the cost matrix C is perturbed.

Theorem 1 ((*SDP*) in polytime)**.** *Let p such that $\tau(p) > (1+\eta)m$, for a fixed $\eta > 0$. Apply a random perturbation of magnitude σ to the cost matrix C. Assume that $\mathfrak{M}$ is compact and smooth (LICQ holds). Solve (BM) using a constrained optimization method with 2nd-order guarantees (e.g., Theorem 4) initialized at an approximately feasible point Y_0. Then, after $\mathrm{poly}(n, \sigma^{-1})$ iterations, we get a point Y such that YY^T is approximately optimal for (SDP) with high probability.*

Note that our bound on p does not depend on the magnitude σ of the perturbation, as opposed to earlier works. This improved bound relies on a technical novelty of this paper, namely, using sophisticated tools from real algebraic geometry for concentration bounds. Indeed, we characterize the presence of spurious approximately critical points through tubular neighborhoods of algebraic varieties, and make use of effective bounds for the volume of such tubes [4, 23, 27]. On the other hand, [9, 10] uses simple dimension arguments, while [5, 30] rely on random matrix theory.

The following theorem provides polynomial time guarantees for the first stage. We again consider a smoothed analysis setting, but this time perturbing the matrices A_i that define the linear map $\mathcal{A}$.

Theorem 2 ((*SDP$_{ls}$*) in polytime)**.** *Let p such that $\tau(p) > (1+\eta)m$, for a fixed $\eta > 0$. Apply a random perturbation of magnitude σ to the constraint map $\mathcal{A}$. Assume that the sublevel sets $\{Y : \|\mathcal{A}(YY^T) - b\| \leq \alpha\}$ are compact. Solve (BM$_{ls}$) using an unconstrained optimization method with 2nd-order guarantees (e.g., [15, 16]). Then, after $\mathrm{poly}(n, \sigma^{-1})$ iterations, we get a point Y such that YY^T is approximately optimal for (SDP$_{ls}$) with high probability.*

Theorems 1 and 2 together provide end-to-end guarantees for solving (*SDP*), with polynomial smoothed analysis complexity. The main assumptions are that p is slightly above the Barvinok-Pataki bound, while the feasible set is compact and smooth. In the general case, we require that both $C, \mathcal{A}$ are slightly perturbed. If a feasible point of the (*SDP*) is known, then Theorem 1 alone is enough.

We point out that problem (*SDP$_{ls}$*) is a matrix sensing problem which is of interest in its own right, independent of its connection to SDP feasibility. The guarantees from Theorem 2 apply even if the

optimal value of (SDP_{ls}) is strictly positive (i.e., there is no $X \succeq 0$ with $\mathcal{A}(X) = b$). There are earlier works [6, 22, 26] proving global optimality guarantees for the nonconvex problem (BM_{ls}), but they all rely on the restricted isometry property (RIP). To the best of our knowledge, Theorem 2 provides the first global guarantees for (BM_{ls}) that do not rely on RIP.

Related work

We first discuss (SDP). Early work by Burer and Monteiro [12] and by Journée et al. [25] gave strong indications that (BM) has no spurious critical points above the Barvinok-Pataki bound. A formal proof was given by Boumal et al. [5, 9] under a genericity assumption. Subsequent work by Bhojanapalli et al. [5] and Pumir et al. [30] investigated the case of approximately critical points under smoothed analysis. The Barvinok-Pataki bound was recently shown to be optimal up to lower order terms for general SDPs [34], though for structured families of SDPs a smaller rank might suffice [2, 28]. The paper [19] extended the results from [9] to SDPs with multiple PSD constraints.

The works [5, 30] are the closest to this paper, but they do not provide end-to-end polynomial guarantees for solving general SDPs (nor it is claimed). In particular, neither of them discuss initialization or feasibility. The paper [30] relies on Riemannaian optimization, which is only practical for very special SDPs (like the maxcut SDP). In the general case, the retraction operator is computationally intractable. The paper [5] relies on a penalized version of (SDP). It has the crucial catch that they rely on solving an auxiliary problem that in general is as hard as solving another SDP. In addition, their global optimality guarantees are with respect to the solution of the penalized problem, as opposed to the solution of the original SDP. Furthermore, complexity estimates are not provided, and are nontrivial due to the interdependence between the penalty parameter, the perturbation magnitude, and the target precision. In contrast, our paper does provide end-to-end polynomial time guarantees, and even better, our results hold down to the Barvinok-Pataki bound.

Problem (SDP_{ls}) has been well studied in the matrix sensing community. Bhojanapalli et al. [6] showed that (BM_{ls}) has no spurious local minima under the RIP condition, and they also provided polynomial time guarantees. Similar results have been derived later, e.g., [22, 26]. Note that RIP is a very strong assumption about the condition number of the linear map $\mathcal{A}$, particularly since the RIP constant needs to be small [35]. In contrast, our result in Theorem 2 simply assumes a small perturbation around a worst-case instance $\mathcal{A}$. The comparison of smoothed analysis with RIP is akin to assuming that a matrix has a nonzero smallest singular value (and getting bounds that depend on inverse polynomials on this quantity), rather than assuming its condition number is close to one.

2 Critical points in nonlinear programming

Unconstrained case. Consider the optimization problem

$$\min_{y \in \mathbb{R}^n} f(y), \tag{P_{un}}$$

with $f : \mathbb{R}^n \to \mathbb{R}$ twice continuously differentiable. A vector $y \in \mathbb{R}^n$ is a *2nd-order critical* point for (P_{un}), abbreviated 2-critical, if $\nabla f(y) = 0$, $\nabla^2 f(y) \succeq 0$. Practical optimization algorithms cannot obtain a solution satisfying these equations exactly. Hence, we relax these conditions.

Definition 1. Given $\varepsilon_1, \varepsilon_2 \in \mathbb{R}_+$, a vector y is $(\varepsilon_1, \varepsilon_2)$-*approximately 2-critical* (AC) for (P_{un}) if:

$$\|\nabla f(y)\| \leq \varepsilon_1, \qquad \nabla^2 f(y) \succeq -\varepsilon_2 I_n. \tag{1}$$

Several algorithms for unconstrained optimization with provable convergence guarantees are known. Recent work has focused on deriving algorithms with finite-time guarantees. In particular, the trust region method computes an $(\varepsilon_1, \varepsilon_2)$-AC point in $O(\max\{\varepsilon_1^{-2}\varepsilon_2^{-1}, \varepsilon_2^{-3}\})$ iterations [16], and the adaptive regularization with cubics (ARC) method takes $O(\max\{\varepsilon_1^{-2}, \varepsilon_2^{-3}\})$ iterations [15]. We formally state the result for the ARC method.

Theorem 3 ([15]). *Assume that there exists $\alpha > 0$ such that a point y_0 with $f(y_0) \leq \alpha$ is known, and the functions $f, \nabla f, \nabla^2 f$ are uniformly bounded and Lipschitz continuous on the set $\{y : f(y) \leq \alpha\}$. The ARC method initialized at y_0 requires $O(\max\{\varepsilon_1^{-2}, \varepsilon_2^{-3}\})$ iterations to produce an $(\varepsilon_1, \varepsilon_2)$-AC point y. Furthermore, each iteration requires $O(1)$ evaluations of f and its derivatives.*

129 **Constrained case.** Consider the nonlinear program

$$\min_{y \in \mathfrak{M}} f(y), \qquad \mathfrak{M} := \{y \in \mathbb{R}^m : h(y) = 0\}, \tag{P_{con}}$$

130 with $f : \mathbb{R}^n \to \mathbb{R}$, $h : \mathbb{R}^n \to \mathbb{R}^m$ twice continuously differentiable. The Lagrangian function is
131 $L(y, \lambda) = f(y) + \lambda \cdot h(y)$. A vector $y \in \mathbb{R}^n$ is a *2-critical point* if there are multipliers $\lambda \in \mathbb{R}^m$ satisfying
132 some first order and second order optimality conditions. As before, for practical optimization we
133 need to consider a notion of approximate criticality.

134 **Definition 2.** Given $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$, $\gamma \in \mathbb{R}_+$, a pair (y, λ) is (ε, γ)-*approximately feasible*
135 *approximately 2-critical* (AFAC) for (P_{con}) if:

$$\|h(y)\| \leq \varepsilon_0, \qquad \|\nabla_y L(y, \lambda)\| \leq \varepsilon_1, \tag{2a}$$

$$u^T \nabla_{yy}^2 L(y, \lambda) u \geq -\varepsilon_2 \quad (\forall\, u : \|u\| = 1, \|\nabla h(y) u\| \leq \gamma). \tag{2b}$$

136 In the constrained case, for a local minima to be a critical point we need some regularity conditions.
137 One such condition is the *linear independence constraint qualification* (LICQ), that states that $\nabla h(y)$
138 is full rank. This is equivalent to $\mathfrak{M}$ being smooth at y for the case of complete intersections (i.e.,
139 $\text{codim}\, \mathfrak{M} = m$). We next introduce a quantitative version of LICQ.

140 **Definition 3.** For $\varrho > 0$, say that ϱ-*LICQ* holds at y if the singular values of $\nabla h(y)$ are at least ϱ.

141 Several local optimization methods for (P_{con}) with provable convergence guarantees are known. In
142 particular, augmented Lagrangians [1] and trust-region methods [20, §15.4] converge to 2-critical
143 points. Recent work has focused on finding algorithms with finite-time guarantees. The complexity
144 of computing approximately 1-critical points was studied in, e.g., [7, 17, 18, 21].

145 As for approximately 2-critical points, we are only aware of [14, 31]. But both papers use a different
146 2nd-order condition, which is not easy to translate into our setting. Nonetheless, in Theorem 4 below
147 we show that AFAC points can be computed in polynomial time. The proof of this theorem is in
148 Appendix A, and relies on a variant of the method from [14]. To the best of our knowledge, this is the
149 first polynomial time bound for computing 2-critical points using the standard notions of criticality.

150 **Theorem 4.** *Assume that there exist* $\beta, \varrho \in \mathbb{R}_+$ *such that a point* y_0 *in the set* $\mathfrak{M}_\beta := \{y : \|h(y)\| \leq \beta\}$
151 *is known, the functions* $f, \nabla f, \nabla^2 f, \nabla h_i, \nabla^2 h_i$ $(i \in [m])$ *are uniformly bounded and Lipschitz*
152 *continuous on* $\mathfrak{M}_\beta$, *and* ϱ-*LICQ holds at all* $y \in \mathfrak{M}_\beta$. *Let* $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$, $\gamma \in \mathbb{R}_+$ *such that*

$$\varepsilon_0 \leq \beta, \quad \varepsilon_1 \leq 1, \quad \varepsilon_1^2 \leq \tfrac{1}{16} R_\lambda^{-1} \varepsilon_0\, \varepsilon_2, \quad \gamma^2 \leq \tfrac{1}{16} R_\lambda^{-3} \varepsilon_0\, \varepsilon_2, \tag{3}$$

153 *where* $R_\lambda := 2 + 2\, \varrho^{-1} L_f$, $L_f := \max_y \|\nabla f(y)\|$. *There is an algorithm that, when initialized*
154 *at* y_0, *requires* $O(\max\{\varepsilon_0^{-2}\, \varepsilon_1^{-2},\, \varepsilon_0^{-3}\, \varepsilon_2^{-3}\})$ *evaluations of* f, h *and their derivatives to produce an*
155 (ε, γ)-*AFAC pair* (y, λ), *with* $\|\lambda\| \leq R_\lambda$.

156 *Remark.* Theorem 4 assumes that ϱ-LICQ holds everywhere in $\mathfrak{M}_\beta$, an assumption which can only
157 be guaranteed when β is very small. Hence the initial point y_0 must be approximately feasible.

3 Optimality of critical points of (*BM*)

159 In this section we will show that problem (*BM*) has no *spurious* approximately feasible approxi-
160 mately critical (AFAC) points with high probability. This means that any AFAC point of (*BM*) is
161 approximately optimal for (*SDP*). We consider a smoothed analysis setting in which the constraint
162 variables $\mathcal{A}, b$ are fixed, and the cost matrix C is subject to a small random perturbation. We will
163 restrict our attention to AFAC pairs (Y, λ) of bounded norm. Hence, we assume that $\|Y\| \leq R_Y$ and
164 $\|\lambda\| \leq R_\lambda$ for some fixed constants R_Y, R_λ.

165 A crucial step toward our theorem is a geometric characterization of the spurious AFAC points in
166 terms of tubes around algebraic varieties. We then take advantage of known effective bounds for the
167 volume of such tubes [4, 27].

168 From now on, we use the Frobenius norm for all matrices.

3.1 Spurious approximately critical points

170 The optimality conditions for (*SDP*) are well known: $\mathcal{A}(X) = b, S(\lambda) X = 0, X \succeq 0, S(\lambda) \succeq 0$, where
171 $S(\lambda)$ is the following *slack* matrix

$$S(\lambda) := C - \mathcal{A}^*(\lambda) \in \mathbb{S}^n,$$

172   and $\mathcal{A}^* : \mathbb{R}^m \to \mathbb{S}^n$, $\lambda \mapsto \sum_i \lambda_i A_i$ is the adjoint of $\mathcal{A}$. We now relax the optimality conditions.

173   **Definition 4.** Let $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$. A pair (X, λ) is ε-*approximately optimal* for (*SDP*) if:

$$\|\mathcal{A}(X) - b\| \leq \varepsilon_0, \quad \|S(\lambda)X\| \leq \varepsilon_1, \quad X \succeq 0, \quad S(\lambda) \succeq -\varepsilon_2 \, I_n. \tag{4}$$

174   It is known that an ε-approximately optimal solution is at distance $O(\|\varepsilon\|)$ from an optimal solution
175   under nondegeneracy assumptions [33]. We can also give a simple bound on the optimality gap.

176   **Lemma 1.** *If* $(\bar{X}, \bar{\lambda})$ *is* ε-*approximately optimal for* (*SDP*) *then*

$$C \bullet \bar{X} \; \leq \; C \bullet X + \varepsilon_0 \|\bar{\lambda}\| + \varepsilon_1 \sqrt{n} + \varepsilon_2 \|X\| \sqrt{n} \qquad \forall \, \textit{feasible } X.$$

177   *Proof.* The lemma follows from the following equations:

$$
\begin{aligned}
C \bullet X \;&=\; \bar{\lambda} \cdot \mathcal{A}(X) + S(\bar{\lambda}) \bullet X \;\geq\; \bar{\lambda} \cdot b - (\varepsilon_2 I_n) \bullet X \;=\; \bar{\lambda} \cdot b - \varepsilon_2 \|X\| \sqrt{n}, \\
\bar{\lambda} \cdot b \;&\geq\; \bar{\lambda} \cdot \mathcal{A}(\bar{X}) - \|\bar{\lambda}\| \, \|b - \mathcal{A}(\bar{X})\| \;\geq\; \bar{\lambda} \cdot \mathcal{A}(\bar{X}) - \varepsilon_0 \|\bar{\lambda}\|, \\
\bar{\lambda} \cdot \mathcal{A}(\bar{X}) \;&\geq\; \bar{\lambda} \cdot \mathcal{A}(\bar{X}) + S(\bar{\lambda}) \bullet \bar{X} - \|S(\bar{\lambda})\bar{X}\|_* \;\geq\; C \bullet \bar{X} - \varepsilon_1 \sqrt{n}. \qquad \square
\end{aligned}
$$

178   We proceed to problem (*BM*). This is a special instance of the nonlinear program (P_{con}) with $f(Y) =$
179   $C \bullet YY^T$ and $h(Y) = \mathcal{A}(YY^T) - b$. The Lagrangian function is $L(y, \lambda) = S(\lambda) \bullet YY^T + b^T \lambda$.
180   The criticality conditions for (*BM*) are obtained by specializing (2).

181   **Definition 5.** Let $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$, $\gamma \in \mathbb{R}_+$. A pair (Y, λ) is (ε, γ)-AFAC for (*BM*) if:

$$\|\mathcal{A}(YY^T) - b\| \leq \varepsilon_0, \qquad \|S(\lambda)Y\| \leq \varepsilon_1, \tag{5a}$$

$$S(\lambda) \bullet UU^T \geq -\varepsilon_2 \qquad (\forall U \in \mathbb{R}^{n \times p} : \|U\| = 1, \, \|\mathcal{A}(UY^T)\| \leq \gamma). \tag{5b}$$

182   We are ready to formalize the concept of spurious critical points.

183   **Definition 6.** Let $R_Y, R_\lambda \in \mathbb{R}_+$ be fixed and let (Y, λ) such that $\|Y\| \leq R_Y$ and $\|\lambda\| \leq R_\lambda$. Given
184   $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$, $\gamma \in \mathbb{R}_+$, (Y, λ) is *spurious* (ε, γ)-AFAC if (Y, λ) is (ε, γ)-AFAC for (*BM*)
185   but (YY^T, λ) is not ε'-approx. optimal for (*SDP*) for $\varepsilon' := (\varepsilon_0, R_Y \varepsilon_1, \varepsilon_2)$. A pair (Y, λ) is spurious
186   *exactly* critical if the above holds for $\varepsilon = 0, \gamma = 0$.

3.2   Statement of the theorem

188   We present the main result of this section. Let $\mathcal{A}, b$ be fixed, and let C be obtained from a random
189   perturbation of magnitude σ around some fixed $\bar{C}$. Concretely, C is *uniformly* distributed on the
190   Frobenius ball $\mathbf{B}_\sigma(\bar{C}) \subset \mathbb{S}^n$ of radius σ centered at $\bar{C}$. Consider the set $\mathscr{C}_{\varepsilon, \gamma} \subset \mathbb{S}^n$, consisting of all
191   cost matrices for which there is a spurious AFAC point:

$$\mathscr{C}_{\varepsilon, \gamma} := \{C \in \mathbb{S}^n : \exists (Y, \lambda) \text{ an spurious } (\varepsilon, \gamma)\text{-AFAC pair}\}.$$

192   We show that if $\tau(p) > m$, then the probability $\Pr[C \in \mathscr{C}_{\varepsilon, \gamma}] \to 0$ as the ratio $\varepsilon_1 / \gamma \to 0$.

193   **Theorem 5** (critical $\Rightarrow$ optimal)**.** *Let* p *such that* $\tau(p) > m$. *Let* $\varepsilon \in \mathbb{R}_+^3$, $\gamma \in \mathbb{R}_+$. *Let* C *be uniformly*
194   *distributed on the Frobenius ball* $\mathbf{B}_\sigma(\bar{C})$. *Then*

$$\Pr[C \in \mathscr{C}_{\varepsilon, \gamma}] \; \leq \; 4e \, \delta^{\tau(p) - m} \, (3\kappa)^m \, (4n^3/\sigma)^{\tau(p)},$$

195   *where* $\delta := \varepsilon_1 \|\mathcal{A}\| / \gamma$ *and* $\kappa := R_\lambda \|\mathcal{A}\|$, *provided that* $\delta < \sigma / 4n^3$.

196   The following corollary shows that when the stronger condition $\tau(p) > (1 + \eta)m$ holds, where η
197   is a fixed constant, then we can derive a high probability bound while maintaining δ polynomially
198   bounded. Its proof is a straightforward manipulation.

199   **Corollary 1.** *Consider the setup from Theorem 5. Assume that* $\tau(p) \geq (1 + \eta)m + \eta t$ *and* $\delta \leq$
200   $(1/3\kappa)^{1/\eta} (\sigma/4n^3)^{1 + 1/\eta}$ *for some* $\eta, t > 0$. *Then* $\Pr[C \in \mathscr{C}_{\varepsilon, \gamma}] \; \leq \; 4e \, (\sigma/12\kappa n^3)^t$.

3.3 Tubes around varieties

Our proof of Theorem 5 relies on a geometric characterization of the set $\mathscr{C}_{\varepsilon,\gamma}$. Such characterization is known for the case $\varepsilon = 0, \gamma = 0$, corresponding to *exactly* critical points. It was shown in [10], see also [19], that the existence of a spurious exactly critical point implies that C lies in a certain algebraic variety of $\mathbb{S}^n$, as follows:

$$\exists \text{ (spurious exactly critical point)} \implies C \in \mathbb{S}^n_{n-p} + \operatorname{Im}\mathcal{A}^*,$$

where $\mathbb{S}^n_{n-p} := \{X : \operatorname{rank} X \leq n-p\}$ is a variety of bounded rank matrices, and $\operatorname{Im}\mathcal{A}^*$ is the linear space spanned by $A_1, \ldots, A_m$. Hence, we have that $\mathscr{C}_{0,0} \subset \mathbb{S}^n_{n-p} + \operatorname{Im}\mathcal{A}^*$. When $\tau(p) > m$, the variety $\mathbb{S}^n_{n-p} + \operatorname{Im}\mathcal{A}^*$ is properly contained in $\mathbb{S}^n$. It follows that $\mathscr{C}_{0,0}$ has measure zero and hence, for generic C, there are no spurious exactly critical points.

We show below that for approximately critical points the situation is analogous, except that we need to consider a *tubular* neighborhood around the variety $\mathbb{S}^n_{n-p} + \operatorname{Im}\mathcal{A}^*$.

Definition 7. Given $\mathcal{W} \subset \mathbb{S}^n, \delta \in \mathbb{R}_+$, let $\operatorname{tube}_\delta \mathcal{W} := \{X \in \mathbb{S}^n : \exists W \in \mathcal{W} \text{ s.t. } \|X - W\| \leq \delta\}$.

Proposition 1. *Let* $\delta := \varepsilon_1 \|\mathcal{A}\|/\gamma$ *and* $B_\lambda := \{\lambda \in \mathbb{R}^m : \|\lambda\| \leq R_\lambda\}$. *Then*

$$\mathscr{C}_{\varepsilon,\gamma} \subset \operatorname{tube}_\delta(\mathbb{S}^n_{n-p}) + \mathcal{A}^*(B_\lambda) = \operatorname{tube}_\delta(\mathbb{S}^n_{n-p} + \mathcal{A}^*(B_\lambda)).$$

The next lemma is the key ingredient for Proposition 1. It shows that if Y is a spurious AFAC point, then its smallest singular value $\sigma_p(Y)$ is bounded from below. This is a generalization of a previous result by Burer and Monteiro [12] about exactly critical points. An analogue of this lemma is found in [30, Lem.3.2] for the case of EFAC points.

Lemma 2. *Let* Y *be an* (ε, γ)-*AFAC point of (BM). If* $\sigma_p(Y) \leq \gamma/\|\mathcal{A}\|$, *then* YY^T *is* ε'-*approximately optimal for (SDP), with* $\varepsilon' := (\varepsilon_0, R_Y \varepsilon_1, \varepsilon_2)$.

Proof. Let (Y, λ) satisfy (5), and let us show that (YY^T, λ) satisfies (4). The first three conditions in (4) are easy to check. We proceed to show the last one: $S(\lambda) \succeq -\varepsilon_2 I_n$. Given a unit vector $u \in \mathbb{R}^n$, we need to show that $u^T S(\lambda) u \geq -\varepsilon_2$. There is a unit vector $z \in \mathbb{R}^p$ such that $\|Yz\| = \sigma_p(Y)$. The matrix $U := uz^T$ satisfies $\|U\| = 1$ and

$$\|UY^T\| \leq \|u\|\|Yz\| = \sigma_p(Y) \leq \gamma/\|\mathcal{A}\|.$$

Then $\|\mathcal{A}(UY^T)\| \leq \gamma$, so by (5b) we have

$$-\varepsilon_2 \leq S(\lambda) \bullet UU^T = \|z\|^2 (u^T S(\lambda) u) = u^T S(\lambda) u. \qquad \square$$

Proof of Proposition 1. Let $C \in \mathscr{C}_{\varepsilon,\gamma}$, so there is a spurious (ε, γ)-AFAC pair (Y, λ). By Lemma 2, we must have $\sigma_p(Y) > \gamma/\|\mathcal{A}\|$. Let $S := S(\lambda)$, and note that $\|SY\| \leq \varepsilon_1 = \gamma\delta/\|\mathcal{A}\|$ by (5a). Then

$$\operatorname{dist}(S, \mathbb{S}^n_{n-p}) = \sqrt{\textstyle\sum_{i=1}^p \sigma_{n-i+1}^2(S)} \leq \|SY\|/\sigma_p(Y) < \delta, \tag{6}$$

so that $C = S(\lambda) + \mathcal{A}^*(\lambda) \in \operatorname{tube}_\delta(\mathbb{S}^n_{n-p}) + \mathcal{A}^*(B_\lambda). \qquad \square$

By Proposition 1, the set of "bad" cost matrices $\mathscr{C}_{\varepsilon,\gamma}$ is contained in a tube around a variety. To prove Theorem 5, we need an upper bound on the probability mass of this tube. The computation of integrals over tubes has a long history in differential geometry [23]. Effective bounds for these integrals were shown in [4, 13, 27]; they were used for smooth analysis in the first reference. We will use the following recent bound.

Theorem 6 ([4, Thm.1.1]). *Let* $V \subset \mathbb{R}^k$ *be a real variety of codimension c defined by polynomials of degree at most D. Let x be uniformly distributed on the Euclidean ball* $\mathbf{B}_\sigma(\bar{x}) \subset \mathbb{R}^k$. *If the ratio* $\sigma/\delta \geq (4D+1)(k-c)$, *then we have that* $\Pr[x \in \operatorname{tube}_\delta(V)] \leq 4e(4kD\delta/\sigma)^c$,

Proof of Theorem 5. Let $B_\lambda \subset \mathbb{R}^m$ be the ball of radius R_λ centered at zero. Consider an ϵ-net $\mathcal{N}$ of B_λ, where $\epsilon := \delta/\|\mathcal{A}\|$. It is known that $(3R_\lambda/\epsilon)^m = (3\kappa/\delta)^m$ points suffices. Observe that

$$\mathcal{A}^*(B_\lambda) \subset \mathcal{A}^*(\operatorname{tube}_\epsilon(\mathcal{N})) \subset \operatorname{tube}_\delta(\mathcal{A}^*(\mathcal{N})), \tag{7a}$$

$$\mathscr{C}_{\varepsilon,\gamma} \subset \operatorname{tube}_\delta(\mathbb{S}^n_{n-p}) + \mathcal{A}^*(B_\lambda) \subset \operatorname{tube}_{2\delta}(\mathbb{S}^n_{n-p}) + \mathcal{A}^*(\mathcal{N}). \tag{7b}$$

238 Recall that $\mathbb{S}^n_{n-p}$ is a variety of codimension $\tau(p)$ defined by equations of degree $n-p+1$. For any
239 $\ell \in \mathcal{N}$, Theorem 6 gives

$$\Pr[C-\mathcal{A}^*(\ell) \in \text{tube}_{2\delta}(\mathbb{S}^n_{n-p})] \; < \; 4e\big(8\tau(n)(n-p+1)\delta/\sigma\big)^{\tau(p)} \; < \; 4e\big(4n^3\delta/\sigma\big)^{\tau(p)}.$$

240 Finally, the union bound gives

$$\Pr[C \in \mathscr{C}_{\varepsilon,\gamma}] \; \leq \; \#\mathcal{N} \cdot \Pr[C \in \text{tube}_{2\delta}(\mathbb{S}^n_{n-p})+\mathcal{A}^*(\ell)] \; < \; (3\kappa/\delta)^m \cdot 4e\left(4n^3\delta/\sigma\right)^{\tau(p)}. \qquad \square$$

4 Optimality of critical points of (BM_{ls})

242 In this section we will show that (BM_{ls}) has no *spurious* approximately critical (AC) points with
243 high probability. This means that any AC point of (BM_{ls}) is approximately optimal for (SDP_{ls}).
244 Hence, we can solve the least squares problem (SDP_{ls}) to global optimality using the Burer-Monteiro
245 method. Note that if (SDP) is feasible, then the optimal value of (SDP_{ls}) is zero. Nevertheless, the
246 results of this section apply to an arbitrary instance of (SDP_{ls}), even if the optimal value is nonzero.
247 Appendix B contains the proofs of the lemmas, propositions, and theorems from this section.

248 We proceed to formalize the notion of spurious critical points. The optimality conditions for the
249 convex problem (SDP_{ls}) are: $S(X)X = 0, X \succeq 0, S(X) \succeq 0$, where $S(X)$ is the gradient of the
250 least squares objective function $f(X) = \|\mathcal{A}(X)-b\|^2$:

$$S(X) \; := \; \nabla f(X) \; = \; 2\mathcal{A}^*(\mathcal{A}(X)-b) \; \in \; \mathbb{S}^n.$$

251 We call X approximately optimal if either $\mathcal{A}(X) \approx b$ or the above conditions are almost satisfied.

252 **Definition 8.** Let $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}^3_+$. A matrix $X \succeq 0$ is ε-*approximately optimal* for (SDP_{ls}) if

$$\|\mathcal{A}(X)-b\| \leq \varepsilon_0 \qquad \text{or} \qquad (\;\|S(X)X\| \leq \varepsilon_1 \text{ and } S(X) \succeq -\varepsilon_2\, I_n\;). \qquad (8)$$

253 The following lemma bounds the optimality gap for the second case in (8).

254 **Lemma 3.** *Let $\bar{X} \in \mathbb{S}^n_+$ such that $\|S(\bar{X})\bar{X}\| \leq \varepsilon_1$, $S(\bar{X}) \succeq -\varepsilon_2 I_n$. Then*

$$f(\bar{X}) \; \leq \; f(X) + \varepsilon_1\sqrt{n} + \varepsilon_2\|X\|\sqrt{n} \qquad \forall\, X \in \mathbb{S}^n_+.$$

255 The Burer-Monteiro problem (BM_{ls}) is a special instance of the unconstrained optimization prob-
256 lem (P_{un}). The criticality conditions for (BM_{ls}) are obtained by specializing (1).

257 **Definition 9.** Let $(\varepsilon_1, \varepsilon_2) \in \mathbb{R}^2_+$. $Y \in \mathbb{R}^{n \times p}$ is $(\varepsilon_1, \varepsilon_2)$-*approximately 2-critical* (AC) for (BM_{ls}) if

$$\|S(YY^T)Y\| \leq \varepsilon_1, \qquad (9a)$$

$$S(YY^T) \bullet UU^T + 4\|\mathcal{A}(UY^T)\|^2 \geq -\varepsilon_2 \quad (\forall U \in \mathbb{R}^{n \times p} : \|U\|=1) \qquad (9b)$$

258 We are ready to formalize the concept of spurious critical points.

259 **Definition 10.** Let $R_Y \in \mathbb{R}_+$ be fixed and let Y with $\|Y\| \leq R_Y$. Given $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}^3_+$, Y
260 is *spurious ε-AC* if Y is $(\varepsilon_1, \varepsilon_2)$-AC for (BM_{ls}) but YY^T is not ε'-approx. optimal for (SDP_{ls}) for
261 $\varepsilon' := (\varepsilon_0, R_Y\varepsilon_1, 5\varepsilon_2)$. Y is spurious *exactly* critical if the above holds for $\varepsilon = 0$.

262 We assume here that $b \in \mathbb{R}^m$ is fixed, and we vary the linear map $\mathcal{A}$. We identify $\mathcal{A}$ with the tuple of
263 matrices $(A_1, \ldots, A_m)$, and hence view it as an element of the Euclidean space $(\mathbb{S}^n)^m$. We assume
264 that $\mathcal{A}$ is *uniformly* distributed on the Euclidean ball $\mathbf{B}_\sigma(\bar{A}) \subset (\mathbb{S}^n)^m$ of radius σ centered at $\bar{A}$.
265 Consider the set $\mathscr{A}_\varepsilon \subset (\mathbb{S}^n)^m$, consisting of all $\mathcal{A}$ for which there is a spurious AC point:

$$\mathscr{A}_\varepsilon \; := \; \{\mathcal{A} \in (\mathbb{S}^n)^m : \exists Y \text{ a spurious } \varepsilon\text{-AC point}\}.$$

266 We show that if $\tau(p) > m$, then the probability $\Pr[\mathcal{A} \in \mathscr{A}_\varepsilon] \to 0$ as the ratio $\varepsilon_1/\sqrt{\varepsilon_2} \to 0$.

267 **Theorem 7** (critical $\Rightarrow$ feasible). *Let p such that $\tau(p)>m$, and $\varepsilon \in \mathbb{R}^3_+$. Let $\mathcal{A}$ be uniformly distributed*
268 *on the Euclidean ball $\mathbf{B}_\sigma(\bar{A})$. Then*

$$\Pr[\mathcal{A} \in \mathscr{A}_\varepsilon] \; \leq \; 4e\, \delta^{\tau(p)-m}\, (3\kappa)^m\, (2n^3m/\sigma\varepsilon_0)^{\tau(p)},$$

269 *where $\delta := \varepsilon_1 R_A/\sqrt{\varepsilon_2}$, $\kappa := 2(R_A R_Y^2+\|b\|)R_A$, and $R_A := \|\bar{A}\|+\sigma$, provided that $\delta < \sigma\varepsilon_0/4n^3m$.*

270 As before, we can derive a high probability bound when $\tau(p) > (1+\eta)m$.

271 **Corollary 2.** *Consider the setup from Theorem 7. Assume that $\tau(p) \geq (1+\eta)m + \eta t$, $\varepsilon_0 \geq \rho\sigma$ and*
272 $\delta \leq (1/3\kappa)^{1/\eta}(\rho\sigma^2/2n^3m)^{1+1/\eta}$ *for some $\eta, t, \rho > 0$. Then $\Pr[\mathcal{A} \in \mathscr{A}_\varepsilon] \leq 4e\,(\rho\sigma^2/6\kappa n^3m)^t$.*

273 As in Section 3, our proof of Theorem 5 relies on a geometric characterization of the spurious AC
274 points. First consider the simpler case of spurious *exactly* critical points ($\varepsilon = 0$). The following
275 equation is a consequence of our analysis:

$$\exists \text{ (spurious exactly critical point)} \implies \mathbb{S}^n_{n-p} \cap \mathrm{Im}\mathcal{A}^* \text{ is nontrivial.}$$

276 This implies that the set $\mathscr{A}_0$ has measure zero when $\tau(p) > m$. In order to handle the case $\varepsilon > 0$ we
277 rely again on tubular neighborhoods, as stated next.

278 **Proposition 2.** *Let $\delta := \varepsilon_1 R_A/\sqrt{\varepsilon_2}$, $D_\lambda := \{\lambda \in \mathbb{R}^m : 2\varepsilon_0 \leq \|\lambda\| \leq R_\lambda\}$, $R_\lambda := 2(R_A R_Y^2 + \|b\|)$.*
279 *Then for $\mathcal{A} \in \mathscr{A}_\varepsilon$ we have that $\mathrm{tube}_\delta(\mathbb{S}^n_{n-p}) \cap \mathcal{A}^*(D_\lambda) \neq \emptyset$.*

5 Polynomial time guarantees

281 We are ready to derive polynomial time guarantees for the Burer-Monteiro method. The proof of
282 Theorem 1 relies on two facts: (i) the method from Theorem 4 finds an AFAC point of (*BM*) in
283 polynomial time; (ii) Corollary 1 shows that an AFAC point of (*BM*) is approximately optimal for
284 (*SDP*) with high probability. Hence, the Burer-Monteiro method solves (*SDP*) in polynomial time
285 in the setting of smooth analysis. The proof of Theorem 2 is analogous, except that it relies on
286 Theorem 3 and corollary 2.

287 Appendix C provides explicit complexity estimates for solving (*SDP*) and (*SDP$_{ls}$*) using the Burer-
288 Monteiro method, indicating the precise optimality guarantees of the output.

6 Experiments

290 We present some experimental results to complement our theorems. We rely on the library `NLopt` [24]
291 (MIT license) to solve the nonlinear problem (*BM*). Concretely, we use the *augmented Lagrangian*
292 *method* (ALM) implemented in `NLopt` (which is based on [8]), and we use the preconditioned
293 truncated Newton method as the subroutine. We also rely on the commercial solver `Mosek` for SDPs.

294 For our first experiment we consider a random SDP with a planted solution. More precisely, we take
295 a matrix $X_0 \in \mathbb{S}^{50}$, $X_0 \succeq 0$ of rank $r \in \{4, 7, 12\}$, and generate a random SDP for which X_0 is
296 an optimal solution. To do so, we generate $m := \tau(r)$ random constraints that are satisfied at X_0,
297 and then find a cost matrix C in the normal cone of X_0. We generate 100 random SDPs for each r
298 We solve these SDPs with the Burer-Monteiro method, using different values of p (the rank of Y)
299 and random initializations. The initial points are matrices with i.i.d. normalized Gaussian entries.
300 Figure 1 shows the percentage of experiments solved correctly for each value of r and p. We regard
301 an experiment as "correct" if the criticality conditions from (4) are satisfied.

Figure 1: Performance of the BM method for random SDPs with a planted solution of rank r.

302 Figure 1 illustrates that there is a sharp phase transition at the Barvinok-Pataki bound $p = r$. Above
303 the Barvinok-Pataki bound, the Burer-Monteiro method solves most instances. Beneath the Barvinok-
304 Pataki bound, it is not just that our techniques stop working, but that the method itself usually

fails. Note that previous work [5, 30] required p to be larger than $3r/\sqrt{2}$. However, we see from experiments that the phase transition is much sharper than this. Observe that, even for $p \geq r$, the number of experiments solved is always below 100%. Nonetheless, the number of bad instances seems to get smaller for larger values of p. This agrees with our result from Theorem 5.

For the second experiment we fix the parameters $n := 50$, $m := 28$, and $p := r := 7$. Among the 100 random SDPs considered in Figure 1, we take an instance for which the Burer-Monteiro problem performed badly. We then perturb this seemingly bad SDP by adding varying amounts of noise σ. For each noise level we solve 70 random experiments, in which both the perturbations and the initializations are random. The perturbations consist in adding to each matrix a random matrix with i.i.d. Gaussian entries scaled by the noise level. Figure 2 summarizes the results obtained.

(a) Fraction of SDPs solved

(b) Iteration progress

Figure 2: Performance of the BM method after perturbing a "bad" SDP with different levels of noise.

Figure 2a shows the percentage of instances for which the Burer-Monteiro method succeeded for each noise level. The percentage is with respect to the random perturbation and the random initialization. For the unperturbed problem ($\sigma = 0$) the method succeeds only for 52% of the random initializations. This percentage increases as we add noise. For $\sigma = 0.2$ the method succeeds 87% of the time. Figure 2b shows the progress of the algorithm for the cases $\sigma = 0$ and $\sigma = 0.2$. The progress is measured in terms of the residual of the criticality conditions. The figure shows the mean and standard deviation of the residual for each iteration of ALM.

Figures 2a and 2b illustrate the advantages of smoothing a badly behaved SDP. Theorem 9 predicts that the complexity of the algorithm is proportional to σ^{-d} for some exponent d. So for a fixed number of iterations N, we should set the noise level proportional to $N^{-1/d}$. However, our bounds were shown for the algorithm from Theorem 4. We do not know if they also apply to ALM.

7   Discussion

We study the Burer-Monteiro method for solving the problems (*SDP*) and (*SDP*$_{ls}$). We provide the first polynomial time guarantees, using a smoothed analysis setting. Previous analyses for (*SDP*) did not provide end-to-end guarantees and made structural assumptions on the problem instance, while previous work for (*SDP*$_{ls}$) relied on the RIP condition, a strong assumption.

The main technical novelty of this paper is the use of volumes of tubular neighborhoods of varieties for deriving concentration bounds. This allows our results to hold even as p approaches the Barvinok-Pataki bound arbitrarily close. Although the Barvinok-Pataki bound is optimal for general SDPs, there are better bounds for some special classes of SDPs. An important open question is whether our results could be extended to allow for smaller values of p in those special classes.

The Burer-Monteiro method is not a single algorithm, but a family of algorithms, one for each local optimization method for solving (*BM*). Similarly, our guarantees in Theorems 1 and 2 apply for any local method with 2nd order guarantees. The only method we are aware that provably computes AFAC points of (*BM*) efficiently is the one in Theorem 4. It would be interesting to extend our results to classical optimization methods, such as ALM, which have robust implementations.

References

[1] Roberto Andreani, Ernesto G Birgin, José Mario Martínez, and María Laura Schuverdt. Second-order negative-curvature methods for box-constrained and general constrained optimization. *Comput. Optim. Appl.*, 45(2):209–236, 2010.

[2] Afonso S Bandeira, Nicolas Boumal, and Vladislav Voroninski. On the low-rank approach for semidefinite programs arising in synchronization and community detection. In *Proc. Conf. Learn. Theory*, pages 361–382, 2016.

[3] Alexander I. Barvinok. Problems of distance geometry and convex properties of quadratic maps. *Discrete Comput. Geom.*, 13(2):189–202, 1995.

[4] Saugata Basu and Antonio Lerario. Hausdorff approximations and volume of tubes of singular algebraic sets. *arXiv:2104.05053*, 2021.

[5] Srinadh Bhojanapalli, Nicolas Boumal, Prateek Jain, and Praneeth Netrapalli. Smoothed analysis for low-rank solutions to semidefinite programs in quadratic penalty form. In *Conf. Learn. Theory*, pages 3243–3270, 2018.

[6] Srinadh Bhojanapalli, Behnam Neyshabur, and Nati Srebro. Global optimality of local search for low rank matrix recovery. In *Adv. Neural Inf. Process. Syst.*, pages 3873–3881, 2016.

[7] Ernesto G Birgin, JL Gardenghi, José Mario Martinez, Sandra A Santos, and Ph L Toint. Evaluation complexity for nonlinear constrained optimization using unscaled KKT conditions and high-order models. *SIAM J. Optim.*, 26(2):951–967, 2016.

[8] Ernesto G Birgin and José Mario Martínez. Improving ultimate convergence of an augmented Lagrangian method. *Optim. Methods Software*, 23(2):177–195, 2008.

[9] Nicolas Boumal, Vlad Voroninski, and Afonso Bandeira. The non-convex Burer-Monteiro approach works on smooth semidefinite programs. In *Adv. Neural Inf. Process. Syst.*, pages 2757–2765, 2016.

[10] Nicolas Boumal, Vladislav Voroninski, and Afonso Bandeira. Deterministic guarantees for Burer-Monteiro factorizations of smooth semidefinite programs. *Commun. Pure Appl. Math.*, 2019.

[11] Samuel Burer and Renato DC Monteiro. A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization. *Math. Program.*, 95(2):329–357, 2003.

[12] Samuel Burer and Renato DC Monteiro. Local minima and convergence in low-rank semidefinite programming. *Math. Program.*, 103(3):427–444, 2005.

[13] Peter Bürgisser, Felipe Cucker, and Martin Lotz. The probability that a slightly perturbed numerical analysis problem is difficult. *Math. Comput.*, 77(263):1559–1583, 2008.

[14] Coralia Cartis, Nicholas IM Gould, and Ph L Toint. Optimality of orders one to three and beyond: characterization and evaluation complexity in constrained nonconvex optimization. *J. Complexity*, 2018.

[15] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Adaptive cubic regularisation methods for unconstrained optimization. Part II: worst-case function-and derivative-evaluation complexity. *Math. Program.*, 130(2):295–319, 2011.

[16] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Complexity bounds for second-order optimality in unconstrained optimization. *J. Complexity*, 28(1):93–108, 2012.

[17] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. On the evaluation complexity of cubic regularization methods for potentially rank-deficient nonlinear least-squares problems and its relevance to constrained nonlinear optimization. *SIAM J. Optim.*, 23(3):1553–1574, 2013.

[18] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. On the complexity of finding first-order critical points in constrained nonlinear optimization. *Math. Program.*, 144(1-2):93–106, 2014.

[19] Diego Cifuentes. On the Burer–Monteiro method for general semidefinite programs. *Optimization Letters*, pages 1–11, 2021.

[20] Andrew R Conn, Nicholas IM Gould, and Ph L Toint. *Trust region methods*, volume 1 of *MOS-SIAM Series Optim.* SIAM, 2000.

[21] Frank E Curtis, Daniel P Robinson, and Mohammadreza Samadi. Complexity analysis of a trust funnel algorithm for equality constrained optimization. *SIAM J. Optim.*, 28(2):1533–1563, 2018.

[22] Rong Ge, Chi Jin, and Yi Zheng. No spurious local minima in nonconvex low rank problems: A unified geometric analysis. In *Proc. Int. Conf. Mach. Learn.*, pages 1233–1242, 2017.

[23] Alfred Gray. *Tubes*, volume 221. Birkhäuser, 2012.

[24] Steven G. Johnson. The NLopt nonlinear-optimization package. Available at `http://github.com/stevengj/nlopt`, 2019.

[25] Michel Journée, Francis Bach, P-A Absil, and Rodolphe Sepulchre. Low-rank optimization on the cone of positive semidefinite matrices. *SIAM J. Optim.*, 20(5):2327–2351, 2010.

[26] Qiuwei Li and Gongguo Tang. The nonconvex geometry of low-rank matrix optimizations with general objective functions. In *IEEE Global Conf. Signal Inf. Process.*, pages 1235–1239, 2017.

[27] Martin Lotz. On the volume of tubular neighborhoods of real algebraic varieties. *Proc. Am. Math. Soc.*, 143(5):1875–1889, 2015.

[28] Song Mei, Theodor Misiakiewicz, Andrea Montanari, and Roberto Imbuzeiro Oliveira. Solving SDPs for synchronization and maxcut problems via the Grothendieck inequality. In *Proc. Conf. Learn. Theory*, pages 1476–1515, 2017.

[29] Gábor Pataki. On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues. *Math. Oper. Res.*, 23(2):339–358, 1998.

[30] Thomas Pumir, Samy Jelassi, and Nicolas Boumal. Smoothed analysis of the low-rank approach for smooth semidefinite programs. In *Adv. Neural Inf. Process. Syst.*, pages 2287–2296, 2018.

[31] Mehmet Fatih Sahin, Armin Eftekhari, Ahmet Alacaoglu, Fabian Latorre, and Volkan Cevher. An inexact augmented Lagrangian framework for nonconvex optimization with nonlinear constraints. *Proc. Neural Inf. Process. Syst.*, 32:13965–13977, 2019.

[32] Daniel A Spielman and Shang-Hua Teng. Smoothed analysis of algorithms: Why the simplex algorithm usually takes polynomial time. *JACM*, 51(3):385–463, 2004.

[33] Jos F Sturm. Error bounds for linear matrix inequalities. *SIAM J. Optim.*, 10(4):1228–1248, 2000.

[34] Irene Waldspurger and Alden Waters. Rank optimality for the Burer–Monteiro factorization. *SIAM Journal on Optimization*, 30(3):2577–2602, 2020.

[35] Richard Zhang, Cédric Josz, Somayeh Sojoudi, and Javad Lavaei. How much restricted isometry is needed in nonconvex matrix recovery? In *Adv. Neural Inf. Process. Syst.*, pages 5586–5597, 2018.