Polynomial time guarantees
for the Burer-Monteiro method

Diego Cifuentes
School of Industrial and Systems Engineering
Georgia Institute of Technology
Atlanta, GA 30332
diego.cifuentes@isye.gatech.edu

Ankur Moitra
Department of Mathematics
Massachusetts Institute of Technology
Cambridge, MA 02139
moitra@mit.edu

Abstract

The Burer-Monteiro method is one of the most widely used techniques for solving large-scale semidefinite programs (SDP). The basic idea is to solve a nonconvex program in Y, where Y is an $n \times p$ matrix such that $X = YY^T$. We show that this method can solve SDPs in polynomial time in a smoothed analysis setting. More precisely, we consider an SDP whose domain satisfies some compactness and smoothness assumptions, and slightly perturb the cost matrix and the constraints. We show that if $p \gtrsim \sqrt{2(1+\eta)m}$, where m is the number of constraints and $\eta > 0$ is any fixed constant, then the Burer-Monteiro method can solve SDPs to any desired accuracy in polynomial time, in the setting of smooth analysis. The bound on p approaches the celebrated Barvinok-Pataki bound in the limit as η goes to zero, beneath which it the nonconvex program can be suboptimal. Our main technical contribution, which is key for our tight bound on p, is to connect spurious approximately critical points of the nonconvex program to tubular neighborhoods of certain algebraic varieties, and then estimate the volume of such tubes.

1   Introduction

Let $\mathbb{S}^n$ be the set of $n \times n$ symmetric matrices. Given matrices $A_1, \ldots, A_m, C \in \mathbb{S}^n$ and a vector $b \in \mathbb{R}^m$, consider the *semidefinite program* (SDP):

$$\min_{X \in \mathbb{S}^n} \ C \bullet X \quad \text{s.t.} \quad \mathcal{A}(X) = b, \quad X \succeq 0, \tag{SDP}$$

where $\mathcal{A} : \mathbb{S}^n \to \mathbb{R}^m$, $X \mapsto (A_1 \bullet X, \ldots, A_m \bullet X)$ is a linear map and the notation $X \succeq 0$ in indicates that X is a positive semidefinite (PSD) matrix. Also consider the *least squares* SDP

$$\min_{X \in \mathbb{S}^n} \ \|\mathcal{A}(X) - b\|^2 \quad \text{s.t.} \quad X \succeq 0, \tag{SDP_{ls}}$$

Though interior point methods solve (SDP) and (SDP_{ls}) in polynomial time, they often have memory problems for large values of n. The Burer-Monteiro method [11, 12] is one of the most widely used procedures for large scale problems. Several papers have worked in understanding the practical success of this method, e.g., [9, 10, 12]. Although various results have been shown, they all fall short of showing that one can reach an approximately optimal solution of (SDP)/(SDP_{ls}) in polynomial time. This paper provides the first polynomial time guarantees for the Burer-Monteiro method.

The Burer-Monteiro method consists in writing $X = YY^T$ for some $Y \in \mathbb{R}^{n \times p}$, and solving a nonconvex optimization problem in terms of Y. For (SDP) we get the problem

$$\min_{Y \in \mathfrak{M}} \ C \bullet YY^T, \qquad \mathfrak{M} := \{Y \in \mathbb{R}^{n \times p} : \mathcal{A}(YY^T) = b\}, \tag{BM}$$

and for (SDP_{ls}) we get the unconstrained problem

$$\min_{Y \in \mathbb{R}^{n \times p}} \ \|\mathcal{A}(YY^T) - b\|^2. \tag{BM_{ls}}$$

We focus now on problem (*SDP*), though our results apply to (SDP_{ls}) as well. Let $\tau(k) := \binom{k+1}{2}$ be the k-th triangular number. It is known that problems (*SDP*) and (*BM*) have the same optimal value for any p such that $\tau(p) \geq m$; this is known as the *Barvinok-Pataki bound* [3, 29]. But due to nonconvexity, local optimization methods may converge to a critical point of (*BM*) which is not globally optimal [34]. In this paper we are mainly interested in 2nd-order critical points (abbreviated as 2-critical points).

Problem (*BM*) has been well studied, most remarkably in the sequel of papers [5, 9, 10, 30]. Boumal et al. [9, 10] showed that (*BM*) has no spurious 2-critical points when $\tau(p) > m$, assuming that the feasible set $\mathfrak{M}$ is a smooth manifold and that the cost matrix C is *generic*. Hence, the Burer-Monteiro method converges to the global optimal solution as long as the cost C is outside a certain "bad" set, which has measure zero. However, this analysis only applies to the limit points. The follow up works [5] and [30] provide finite-iteration guarantees, though for values of p larger than the Barvinok-Pataki bound. They introduce a smoothed analysis [32] setting in which the cost matrix C is subject to a small random perturbation of magnitude σ. Such a perturbation ensures that C stays outside the "bad" region, so the limit points are globally optimal. They then argue that certain notions of approximate criticality for (*BM*) guarantee approximate optimality for (*SDP*) when $\tau(p) \gtrsim \frac{9}{2} m \log(\Omega(\sqrt{n}/\sigma))$.

Despite the earlier work, the question of whether the Burer-Monteiro method can be used to completely solve problem (*SDP*) in polynomial time has not been yet resolved. The works [5, 30] do not provide end-to-end polynomial time guarantees and make structural assumptions on the problem, as will be further discussed in the related work section. In particular, finding a feasible solution to (*SDP*) that could serve as initial point is a complicated and practically relevant problem, but it has not been addressed before. In addition, the lower bound on p is qualitatively wrong, as it is not only larger than the Barvinok-Pataki bound, but also gets worse as the magnitude σ of the perturbation decreases.

In order to fully solve (*SDP*), we propose a 2-stage procedure. First, solve (BM_{ls}) to either get a point Y_0 that is approximately feasible, i.e., $\mathcal{A}(Y_0 Y_0^T) \approx b$, or to prove that no such point exists. Second, solve (*BM*) to get an approximately optimal point, starting with point Y_0 from the first stage. We prove that this scheme is successful when $\tau(p) > (1+\eta)m$, for an arbitrary constant $\eta > 0$.

The next theorem provides polynomial time guarantees for the second stage. The complexity is with respect to the setting of a smoothed analysis, in which the cost matrix C is perturbed.

Theorem 1 ((*SDP*) in polytime). *Let p such that $\tau(p) > (1+\eta)m$, for a fixed $\eta > 0$. Apply a random perturbation of magnitude σ to the cost matrix C. Assume that $\mathfrak{M}$ is compact and smooth (LICQ holds). Solve (BM) using a constrained optimization method with 2nd-order guarantees (e.g., Theorem 4) initialized at an approximately feasible point Y_0. Then, after $\mathrm{poly}(n, \sigma^{-1})$ iterations, we get a point Y such that YY^T is approximately optimal for (SDP) with high probability.* A more detailed version of this result, with explicit optimality and complexity constants, appears in Theorem 9.

Note that our bound on p does not depend on the magnitude σ of the perturbation, as opposed to earlier works. This improved bound relies on a technical novelty of this paper, namely, using sophisticated tools from real algebraic geometry for concentration bounds. Indeed, we characterize the presence of spurious approximately critical points through tubular neighborhoods of algebraic varieties, and make use of effective bounds for the volume of such tubes [4, 23, 27]. On the other hand, [9, 10] uses simple dimension arguments, while [5, 30] rely on random matrix theory.

The following theorem provides polynomial time guarantees for the first stage. We again consider a smoothed analysis setting, but this time perturbing the matrices A_i that define the linear map $\mathcal{A}$.

Theorem 2 ((SDP_{ls}) in polytime). *Let p such that $\tau(p) > (1+\eta)m$, for a fixed $\eta > 0$. Apply a random perturbation of magnitude σ to the constraint map $\mathcal{A}$. Assume that the sublevel sets $\{Y : \|\mathcal{A}(YY^T) - b\| \leq \alpha\}$ are compact. Solve (BM_{ls}) using an unconstrained optimization method with 2nd-order guarantees (e.g., [15, 16]). Then, after $\mathrm{poly}(n, \sigma^{-1})$ iterations, we get a point Y such that YY^T is approximately optimal for (SDP_{ls}) with high probability.* A more detailed version of this result, with explicit optimality and complexity constants, appears in Theorem 10.

Theorems 1 and 2 together provide end-to-end guarantees for solving (*SDP*), with polynomial smoothed analysis complexity. The main assumptions are that p is slightly above the Barvinok-Pataki bound, while the feasible set is compact and smooth. In the general case, we require that both $C, \mathcal{A}$ are slightly perturbed. If a feasible point of the (*SDP*) is known, then Theorem 1 alone is enough.

We point out that problem (SDP_{ls}) is a matrix sensing problem which is of interest in its own right, independent of its connection to SDP feasibility. The guarantees from Theorem 2 apply even if the optimal value of (SDP_{ls}) is strictly positive (i.e., there is no $X \succeq 0$ with $\mathcal{A}(X)=b$). There are earlier works [6, 22, 26] proving global optimality guarantees for the nonconvex problem (BM_{ls}), but they all rely on the restricted isometry property (RIP). To the best of our knowledge, Theorem 2 provides the first global guarantees for (BM_{ls}) that do not rely on RIP.

Related work

We first discuss (*SDP*). Early work by Burer and Monteiro [12] and by Journée et al. [25] gave strong indications that (*BM*) has no spurious critical points above the Barvinok-Pataki bound. A formal proof was given by Boumal et al. [5, 9] under a genericity assumption. Subsequent work by Bhojanapalli et al. [5] and Pumir et al. [30] investigated the case of approximately critical points under smoothed analysis. The Barvinok-Pataki bound was recently shown to be optimal up to lower order terms for general SDPs [34], though for structured families of SDPs a smaller rank might suffice [2, 28]. The paper [19] extended the results from [9] to SDPs with multiple PSD constraints.

The works [5, 30] are the closest to this paper, but they do not provide end-to-end polynomial guarantees for solving general SDPs (nor it is claimed). In particular, neither of them discuss initialization or feasibility. The paper [30] relies on Riemannaian optimization, which is only practical for very special SDPs (like the maxcut SDP). In the general case, the retraction operator is computationally intractable. The paper [5] relies on a penalized version of (*SDP*). It has the crucial catch that they require a preprocessing step that in general is as hard as solving another SDP. Since solving an auxiliary SDP is required, it fails to explain why the BM method improves over general SDP solvers. In addition, the optimality guarantees in [5] are with respect to the solution of the penalized problem. Translating their results into polynomial time guarantees with respect to the original SDP is nontrivial due to interdependence between the penalty parameter, the perturbation magnitude, and the target precision[1]. In contrast, our paper does provide end-to-end polynomial time guarantees, and even better, our results hold down to the Barvinok-Pataki bound.

Problem (SDP_{ls}) has been well studied in the matrix sensing community. Bhojanapalli et al. [6] showed that (BM_{ls}) has no spurious local minima under the RIP condition, and they also provided polynomial time guarantees. Similar results have been derived later, e.g., [22, 26]. Note that RIP is a very strong assumption about the condition number of the linear map $\mathcal{A}$, particularly since the RIP constant needs to be small [35]. In contrast, our result in Theorem 2 simply assumes a small perturbation around a worst-case instance $\mathcal{A}$. The comparison of smoothed analysis with RIP is akin to assuming that a matrix has a nonzero smallest singular value (and getting bounds that depend on inverse polynomials on this quantity), rather than assuming its condition number is close to one.

2 Critical points in nonlinear programming

Unconstrained case. Consider the optimization problem

$$\min_{y \in \mathbb{R}^n} f(y), \qquad (P_{\text{un}})$$

with $f : \mathbb{R}^n \to \mathbb{R}$ twice continuously differentiable. A vector $y \in \mathbb{R}^n$ is a *2nd-order critical* point for (P_{un}), abbreviated 2-critical, if $\nabla f(y) = 0$, $\nabla^2 f(y) \succeq 0$. Practical optimization algorithms cannot obtain a solution satisfying these equations exactly. Hence, we relax these conditions.

Definition 1. Given $\varepsilon_1, \varepsilon_2 \in \mathbb{R}_+$, a vector y is $(\varepsilon_1, \varepsilon_2)$-*approximately 2-critical* (AC) for (P_{un}) if:

$$\|\nabla f(y)\| \leq \varepsilon_1, \qquad \nabla^2 f(y) \succeq -\varepsilon_2 I_n. \qquad (1)$$

---

[1]Translating optimality guarantees of the penalized problem to the original SDP requires tuning the penalty parameter μ. The straightforward choice is $\mu = \Omega(\epsilon^{-2})$, where ϵ is the target precision. This leads to high Lipschitz numbers, complicating the complexity analysis (e.g., Theorem 3 does not apply). In addition, [5, Thm 13] constraints the perturbation magnitude by $\sigma^2 \geq \Omega(\mu \epsilon^{3/2} \gamma^{-1}) \geq \Omega(\epsilon^{-1/2} \gamma^{-1})$. Hence, the results seem to require large perturbations, while in smoothed analysis we are interested in small perturbations.

Several algorithms for unconstrained optimization with provable convergence guarantees are known. Recent work has focused on deriving algorithms with finite-time guarantees. In particular, the trust region method computes an $(\varepsilon_1, \varepsilon_2)$-AC point in $O(\max\{\varepsilon_1^{-2}\varepsilon_2^{-1}, \varepsilon_2^{-3}\})$ iterations [16], and the adaptive regularization with cubics (ARC) method takes $O(\max\{\varepsilon_1^{-2}, \varepsilon_2^{-3}\})$ iterations [15]. We formally state the result for the ARC method.

Theorem 3 ([15]). *Assume that there exists $\alpha > 0$ such that a point y_0 with $f(y_0) \leq \alpha$ is known, and the functions $f, \nabla f, \nabla^2 f$ are uniformly bounded and Lipschitz continuous on the set $\{y : f(y) \leq \alpha\}$. The ARC method initialized at y_0 requires $O(\max\{\varepsilon_1^{-2}, \varepsilon_2^{-3}\})$ iterations to produce an $(\varepsilon_1, \varepsilon_2)$-AC point y. Furthermore, each iteration requires $O(1)$ evaluations of f and its derivatives.*

Constrained case.   Consider the nonlinear program

$$\min_{y \in \mathfrak{M}} \; f(y), \qquad \mathfrak{M} := \{y \in \mathbb{R}^m : h(y) = 0\}, \tag{P_{con}}$$

with $f : \mathbb{R}^n \to \mathbb{R}$, $h : \mathbb{R}^n \to \mathbb{R}^m$ twice continuously differentiable. The Lagrangian function is $L(y, \lambda) = f(y) + \lambda \cdot h(y)$. A vector $y \in \mathbb{R}^n$ is a *2-critical point* if there are multipliers $\lambda \in \mathbb{R}^m$ satisfying some first order and second order optimality conditions. As before, for practical optimization we need to consider a notion of approximate criticality.

Definition 2. Given $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$, $\gamma \in \mathbb{R}_+$, a pair (y, λ) is (ε, γ)-*approximately feasible approximately 2-critical* (AFAC) for (P_{con}) if:

$$\|h(y)\| \leq \varepsilon_0, \qquad\quad \|\nabla_y L(y, \lambda)\| \leq \varepsilon_1, \tag{2a}$$

$$u^T \nabla_{yy}^2 L(y, \lambda) u \geq -\varepsilon_2 \quad (\forall\, u : \|u\| = 1, \|\nabla h(y) u\| \leq \gamma). \tag{2b}$$

In the constrained case, for a local minima to be a critical point we need some regularity conditions. One such condition is the *linear independence constraint qualification* (LICQ), that states that $\nabla h(y)$ is full rank. This is equivalent to $\mathfrak{M}$ being smooth at y for the case of complete intersections (i.e., $\operatorname{codim} \mathfrak{M} = m$). We next introduce a quantitative version of LICQ.

Definition 3. For $\varrho > 0$, say that ϱ-*LICQ* holds at y if the singular values of $\nabla h(y)$ are at least ϱ.

Several local optimization methods for (P_{con}) with provable convergence guarantees are known. In particular, augmented Lagrangians [1] and trust-region methods [20, §15.4] converge to 2-critical points. Recent work has focused on finding algorithms with finite-time guarantees. The complexity of computing approximately 1-critical points was studied in, e.g., [7, 17, 18, 21].

As for approximately 2-critical points, we are only aware of [14, 31]. But both papers use a different 2nd-order condition, which is not easy to translate into our setting. Nonetheless, in Theorem 4 below we show that AFAC points can be computed in polynomial time. The proof of this theorem is in Appendix A, and relies on a variant of the method from [14]. To the best of our knowledge, this is the first polynomial time bound for computing 2-critical points using the standard notions of criticality.

Theorem 4. *Assume that there exist $\beta, \varrho \in \mathbb{R}_+$ such that a point y_0 in the set $\mathfrak{M}_\beta := \{y : \|h(y)\| \leq \beta\}$ is known, the functions $f, \nabla f, \nabla^2 f, \nabla h_i, \nabla^2 h_i$ $(i \in [m])$ are uniformly bounded and Lipschitz continuous on $\mathfrak{M}_\beta$, and ϱ-LICQ holds at all $y \in \mathfrak{M}_\beta$. Let $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$, $\gamma \in \mathbb{R}_+$ such that*

$$\varepsilon_0 \leq \beta, \quad \varepsilon_1 \leq 1, \quad \varepsilon_1^2 \leq \tfrac{1}{16} R_\lambda^{-1} \varepsilon_0\, \varepsilon_2, \quad \gamma^2 \leq \tfrac{1}{16} R_\lambda^{-3} \varepsilon_0\, \varepsilon_2, \tag{3}$$

where $R_\lambda := 2 + 2\varrho^{-1} L_f$, $L_f := \max_y \|\nabla f(y)\|$. There is an algorithm that, when initialized at y_0, requires $O(\max\{\varepsilon_0^{-2}\varepsilon_1^{-2}, \varepsilon_0^{-3}\varepsilon_2^{-3}\})$ evaluations of f, h and their derivatives to produce an (ε, γ)-AFAC pair (y, λ), with $\|\lambda\| \leq R_\lambda$.

Remark. Theorem 4 assumes that ϱ-LICQ holds everywhere in $\mathfrak{M}_\beta$, an assumption which can only be guaranteed when β is very small. Hence the initial point y_0 must be approximately feasible.

3   Optimality of critical points of (*BM*)

In this section we will show that problem (*BM*) has no *spurious* approximately feasible approximately critical (AFAC) points with high probability. This means that any AFAC point of (*BM*) is approximately optimal for (*SDP*). We consider a smoothed analysis setting in which the constraint

variables $\mathcal{A}, b$ are fixed, and the cost matrix C is subject to a small random perturbation. We will restrict our attention to AFAC pairs (Y, λ) of bounded norm. Hence, we assume that $\|Y\| \le R_Y$ and $\|\lambda\| \le R_\lambda$ for some fixed constants R_Y, R_λ.

A crucial step toward our theorem is a geometric characterization of the spurious AFAC points in terms of tubes around algebraic varieties. We then take advantage of known effective bounds for the volume of such tubes [4, 27].

From now on, we use the Frobenius norm for all matrices.

3.1 Spurious approximately critical points

The optimality conditions for (*SDP*) are well known: $\mathcal{A}(X) = b, S(\lambda)X = 0, X \succeq 0, S(\lambda) \succeq 0$, where $S(\lambda)$ is the following *slack* matrix

$$S(\lambda) := C - \mathcal{A}^*(\lambda) \in \mathbb{S}^n,$$

and $\mathcal{A}^* : \mathbb{R}^m \to \mathbb{S}^n, \lambda \mapsto \sum_i \lambda_i A_i$ is the adjoint of $\mathcal{A}$. We now relax the optimality conditions.

Definition 4. Let $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}^3_+$. A pair (X, λ) is ε-*approximately optimal* for (*SDP*) if:

$$\|\mathcal{A}(X) - b\| \le \varepsilon_0, \quad \|S(\lambda)X\| \le \varepsilon_1, \quad X \succeq 0, \quad S(\lambda) \succeq -\varepsilon_2 I_n. \tag{4}$$

It is known that an ε-approximately optimal solution is at distance $O(\|\varepsilon\|)$ from an optimal solution under nondegeneracy assumptions [33]. We can also give a simple bound on the optimality gap.

Lemma 1. If $(\bar{X}, \bar{\lambda})$ is ε-approximately optimal for (*SDP*) then

$$C \bullet \bar{X} \le C \bullet X + \varepsilon_0 \|\bar{\lambda}\| + \varepsilon_1 \sqrt{n} + \varepsilon_2 \|X\| \sqrt{n} \qquad \forall \text{ feasible } X.$$

Proof. The lemma follows from the following equations:

$$C \bullet X = \bar{\lambda} \cdot \mathcal{A}(X) + S(\bar{\lambda}) \bullet X \ge \bar{\lambda} \cdot b - (\varepsilon_2 I_n) \bullet X = \bar{\lambda} \cdot b - \varepsilon_2 \|X\| \sqrt{n},$$
$$\bar{\lambda} \cdot b \ge \bar{\lambda} \cdot \mathcal{A}(\bar{X}) - \|\bar{\lambda}\| \|b - \mathcal{A}(\bar{X})\| \ge \bar{\lambda} \cdot \mathcal{A}(\bar{X}) - \varepsilon_0 \|\bar{\lambda}\|,$$
$$\bar{\lambda} \cdot \mathcal{A}(\bar{X}) \ge \bar{\lambda} \cdot \mathcal{A}(\bar{X}) + S(\bar{\lambda}) \bullet \bar{X} - \|S(\bar{\lambda})\bar{X}\|_* \ge C \bullet \bar{X} - \varepsilon_1 \sqrt{n}. \qquad \square$$

We proceed to problem (*BM*). This is a special instance of the nonlinear program (P_{con}) with $f(Y) = C \bullet YY^T$ and $h(Y) = \mathcal{A}(YY^T) - b$. The Lagrangian function is $L(y, \lambda) = S(\lambda) \bullet YY^T + b^T \lambda$. The criticality conditions for (*BM*) are obtained by specializing (2).

Definition 5. Let $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}^3_+, \gamma \in \mathbb{R}_+$. A pair (Y, λ) is (ε, γ)-AFAC for (*BM*) if:

$$\|\mathcal{A}(YY^T) - b\| \le \varepsilon_0, \qquad \|S(\lambda)Y\| \le \varepsilon_1, \tag{5a}$$
$$S(\lambda) \bullet UU^T \ge -\varepsilon_2 \qquad (\forall U \in \mathbb{R}^{n \times p} : \|U\| = 1, \|\mathcal{A}(UY^T)\| \le \gamma). \tag{5b}$$

We are ready to formalize the concept of spurious critical points.

Definition 6. Let $R_Y, R_\lambda \in \mathbb{R}_+$ be fixed and let (Y, λ) such that $\|Y\| \le R_Y$ and $\|\lambda\| \le R_\lambda$. Given $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}^3_+, \gamma \in \mathbb{R}_+, (Y, \lambda)$ is *spurious* (ε, γ)-AFAC if (Y, λ) is (ε, γ)-AFAC for (*BM*) but (YY^T, λ) is not ε'-approx. optimal for (*SDP*) for $\varepsilon' := (\varepsilon_0, R_Y \varepsilon_1, \varepsilon_2)$. A pair (Y, λ) is spurious *exactly* critical if the above holds for $\varepsilon = 0, \gamma = 0$.

3.2 Statement of the theorem

We present the main result of this section. Let $\mathcal{A}, b$ be fixed, and let C be obtained from a random perturbation of magnitude σ around some fixed $\bar{C}$. Concretely, C is *uniformly* distributed on the Frobenius ball $\mathbf{B}_\sigma(\bar{C}) \subset \mathbb{S}^n$ of radius σ centered at $\bar{C}$. Consider the set $\mathscr{C}_{\varepsilon, \gamma} \subset \mathbb{S}^n$, consisting of all cost matrices for which there is a spurious AFAC point:

$$\mathscr{C}_{\varepsilon, \gamma} := \{C \in \mathbb{S}^n : \exists (Y, \lambda) \text{ an spurious } (\varepsilon, \gamma)\text{-AFAC pair}\}.$$

We show that if $\tau(p) > m$, then the probability $\Pr[C \in \mathscr{C}_{\varepsilon, \gamma}] \to 0$ as the ratio $\varepsilon_1/\gamma \to 0$.

Theorem 5 (critical ⇒ optimal)**.** *Let p such that $\tau(p) > m$. Let $\boldsymbol{\varepsilon} \in \mathbb{R}_+^3$, $\gamma \in \mathbb{R}_+$. Let C be uniformly distributed on the Frobenius ball $\mathbf{B}_\sigma(\bar{C})$. Then*

$$\Pr[C \in \mathscr{C}_{\boldsymbol{\varepsilon}, \gamma}] \ \le \ 4e \, \delta^{\tau(p)-m} \, (3\kappa)^m \, (4n^3/\sigma)^{\tau(p)},$$

where $\delta := \varepsilon_1 \|\mathcal{A}\|/\gamma$ and $\kappa := R_\lambda \|\mathcal{A}\|$, provided that $\delta < \sigma/4n^3$.

The following corollary shows that when the stronger condition $\tau(p) > (1+\eta)m$ holds, where η is a fixed constant, then we can derive a high probability bound while maintaining δ polynomially bounded. Its proof is a straightforward manipulation.

Corollary 1. *Consider the setup from Theorem 5. Assume that $\tau(p) \ge (1+\eta)m + \eta t$ and $\delta \le (1/3\kappa)^{1/\eta}(\sigma/4n^3)^{1+1/\eta}$ for some $\eta, t > 0$. Then $\Pr[C \in \mathscr{C}_{\boldsymbol{\varepsilon}, \gamma}] \ \le \ 4e \, (\sigma/12\kappa n^3)^t$.*

3.3 Tubes around varieties

Our proof of Theorem 5 relies on a geometric characterization of the set $\mathscr{C}_{\boldsymbol{\varepsilon}, \gamma}$. Such characterization is known for the case $\varepsilon = 0, \gamma = 0$, corresponding to *exactly* critical points. It was shown in [10], see also [19], that the existence of a spurious exactly critical point implies that C lies in a certain algebraic variety of $\mathbb{S}^n$, as follows:

$$\exists \, (\text{spurious exactly critical point}) \quad \implies \quad C \in \mathbb{S}_{n-p}^n + \operatorname{Im} \mathcal{A}^*,$$

where $\mathbb{S}_{n-p}^n := \{X : \operatorname{rank} X \le n-p\}$ is a variety of bounded rank matrices, and $\operatorname{Im} \mathcal{A}^*$ is the linear space spanned by $A_1, \dots, A_m$. Hence, we have that $\mathscr{C}_{0,0} \subset \mathbb{S}_{n-p}^n + \operatorname{Im} \mathcal{A}^*$. When $\tau(p) > m$, the variety $\mathbb{S}_{n-p}^n + \operatorname{Im} \mathcal{A}^*$ is properly contained in $\mathbb{S}^n$. It follows that $\mathscr{C}_{0,0}$ has measure zero and hence, for generic C, there are no spurious exactly critical points.

We show below that for approximately critical points the situation is analogous, except that we need to consider a *tubular* neighborhood around the variety $\mathbb{S}_{n-p}^n + \operatorname{Im} \mathcal{A}^*$.

Definition 7. Given $\mathcal{W} \subset \mathbb{S}^n, \delta \in \mathbb{R}_+$, let $\operatorname{tube}_\delta \mathcal{W} := \{X \in \mathbb{S}^n : \exists W \in \mathcal{W} \text{ s.t. } \|X - W\| \le \delta\}$.

Proposition 1. *Let $\delta := \varepsilon_1 \|\mathcal{A}\|/\gamma$ and $B_\lambda := \{\lambda \in \mathbb{R}^m : \|\lambda\| \le R_\lambda\}$. Then*

$$\mathscr{C}_{\boldsymbol{\varepsilon}, \gamma} \subset \operatorname{tube}_\delta(\mathbb{S}_{n-p}^n) + \mathcal{A}^*(B_\lambda).$$

The next lemma is the key ingredient for Proposition 1. It shows that if Y is a spurious AFAC point, then its smallest singular value $\sigma_p(Y)$ is bounded from below. This is a generalization of a previous result by Burer and Monteiro [12] about exactly critical points. An analogue of this lemma is found in [30, Lem.3.2] for the case where $\varepsilon_0 = 0$.

Lemma 2. *Let Y be an $(\boldsymbol{\varepsilon}, \gamma)$-AFAC point of (BM). If $\sigma_p(Y) \le \gamma/\|\mathcal{A}\|$, then YY^T is $\boldsymbol{\varepsilon}'$-approximately optimal for (SDP), with $\boldsymbol{\varepsilon}' := (\varepsilon_0, R_Y \varepsilon_1, \varepsilon_2)$.*

Proof. Let (Y, λ) satisfy (5), and let us show that (YY^T, λ) satisfies (4). The first three conditions in (4) are easy to check. We proceed to show the last one: $S(\lambda) \succeq -\varepsilon_2 I_n$. Given a unit vector $u \in \mathbb{R}^n$, we need to show that $u^T S(\lambda) u \ge -\varepsilon_2$. There is a unit vector $z \in \mathbb{R}^p$ such that $\|Yz\| = \sigma_p(Y)$. The matrix $U := uz^T$ satisfies $\|U\| = 1$ and

$$\|UY^T\| \le \|u\| \|Yz\| = \sigma_p(Y) \le \gamma/\|\mathcal{A}\|.$$

Then $\|\mathcal{A}(UY^T)\| \le \gamma$, so by (5b) we have

$$-\varepsilon_2 \ \le \ S(\lambda) \bullet UU^T \ = \ \|z\|^2 (u^T S(\lambda) u) \ = \ u^T S(\lambda) u. \qquad \square$$

Proof of Proposition 1. Let $C \in \mathscr{C}_{\boldsymbol{\varepsilon}, \gamma}$, so there is a spurious $(\boldsymbol{\varepsilon}, \gamma)$-AFAC pair (Y, λ). By Lemma 2, we must have $\sigma_p(Y) > \gamma/\|\mathcal{A}\|$. Let $S := S(\lambda)$, and note that $\|SY\| \le \varepsilon_1 = \gamma \delta/\|\mathcal{A}\|$ by (5a). Then

$$\operatorname{dist}(S, \mathbb{S}_{n-p}^n) \ = \ \sqrt{\textstyle\sum_{i=1}^p \sigma_{n-i+1}^2(S)} \ \le \ \|SY\|/\sigma_p(Y) \ < \ \delta, \qquad (6)$$

so that $C = S(\lambda) + \mathcal{A}^*(\lambda) \in \operatorname{tube}_\delta(\mathbb{S}_{n-p}^n) + \mathcal{A}^*(B_\lambda). \qquad \square$

By Proposition 1, the set of "bad" cost matrices $\mathscr{C}_{\varepsilon,\gamma}$ is contained in a tube around a variety. To prove Theorem 5, we need an upper bound on the probability mass of this tube. The computation of integrals over tubes has a long history in differential geometry [23]. Effective bounds for these integrals were shown in [4, 13, 27]; they were used for smooth analysis in the first reference. We will use the following recent bound.

Theorem 6 ([4, Thm.1.1]). *Let $V \subset \mathbb{R}^k$ be a real variety of codimension c defined by polynomials of degree at most D. Let x be uniformly distributed on the Euclidean ball $\mathbf{B}_\sigma(\bar{x}) \subset \mathbb{R}^k$. If the ratio $\sigma/\delta \geq (4D+1)(k-c)$, then we have that $\Pr[x \in \text{tube}_\delta(V)] \leq 4e(4kD\delta/\sigma)^c$,*

Proof of Theorem 5. Let $B_\lambda \subset \mathbb{R}^m$ be the ball of radius R_λ centered at zero. Consider an ϵ-net $\mathcal{N}$ of B_λ, where $\epsilon := \delta/\|\mathcal{A}\|$. It is known that $(3R_\lambda/\epsilon)^m = (3\kappa/\delta)^m$ points suffices. Observe that

$$\mathcal{A}^*(B_\lambda) \subset \mathcal{A}^*(\text{tube}_\epsilon(\mathcal{N})) \subset \text{tube}_\delta(\mathcal{A}^*(\mathcal{N})), \tag{7a}$$

$$\mathscr{C}_{\varepsilon,\gamma} \subset \text{tube}_\delta(\mathbb{S}_{n-p}^n) + \mathcal{A}^*(B_\lambda) \subset \text{tube}_{2\delta}(\mathbb{S}_{n-p}^n) + \mathcal{A}^*(\mathcal{N}). \tag{7b}$$

Recall that $\mathbb{S}_{n-p}^n$ is a variety of codimension $\tau(p)$ defined by equations of degree $n-p+1$. For any $\ell \in \mathcal{N}$, Theorem 6 gives

$$\Pr[C - \mathcal{A}^*(\ell) \in \text{tube}_{2\delta}(\mathbb{S}_{n-p}^n)] \; < \; 4e\big(8\tau(n)(n-p+1)\delta/\sigma\big)^{\tau(p)} \; < \; 4e\big(4n^3\delta/\sigma\big)^{\tau(p)}.$$

Finally, the union bound gives

$$\Pr[C \in \mathscr{C}_{\varepsilon,\gamma}] \; \leq \; \#\mathcal{N} \cdot \Pr[C \in \text{tube}_{2\delta}(\mathbb{S}_{n-p}^n) + \mathcal{A}^*(\ell)] \; < \; (3\kappa/\delta)^m \cdot 4e\left(4n^3\delta/\sigma\right)^{\tau(p)}. \qquad \square$$

4 Optimality of critical points of (BM_{ls})

In this section we will show that (BM_{ls}) has no *spurious* approximately critical (AC) points with high probability. This means that any AC point of (BM_{ls}) is approximately optimal for (SDP_{ls}). Hence, we can solve the least squares problem (SDP_{ls}) to global optimality using the Burer-Monteiro method. Note that if (SDP) is feasible, then the optimal value of (SDP_{ls}) is zero. Nevertheless, the results of this section apply to an arbitrary instance of (SDP_{ls}), even if the optimal value is nonzero. Appendix B contains the proofs of the lemmas, propositions, and theorems from this section.

We proceed to formalize the notion of spurious critical points. The optimality conditions for the convex problem (SDP_{ls}) are: $S(X)X = 0, X \succeq 0, S(X) \succeq 0$, where $S(X)$ is the gradient of the least squares objective function $f(X) = \|\mathcal{A}(X) - b\|^2$:

$$S(X) := \nabla f(X) = 2\mathcal{A}^*(\mathcal{A}(X) - b) \in \mathbb{S}^n.$$

We call X approximately optimal if either $\mathcal{A}(X) \approx b$ or the above conditions are almost satisfied.

Definition 8. Let $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$. A matrix $X \succeq 0$ is ε-*approximately optimal* for (SDP_{ls}) if

$$\|\mathcal{A}(X) - b\| \leq \varepsilon_0 \qquad \text{or} \qquad (\|S(X)X\| \leq \varepsilon_1 \text{ and } S(X) \succeq -\varepsilon_2 I_n). \tag{8}$$

The following lemma bounds the optimality gap for the second case in (8).

Lemma 3. *Let $\bar{X} \in \mathbb{S}_+^n$ such that $\|S(\bar{X})\bar{X}\| \leq \varepsilon_1, S(\bar{X}) \succeq -\varepsilon_2 I_n$. Then*

$$f(\bar{X}) \; \leq \; f(X) + \varepsilon_1\sqrt{n} + \varepsilon_2\|X\|\sqrt{n} \qquad \forall X \in \mathbb{S}_+^n.$$

The Burer-Monteiro problem (BM_{ls}) is a special instance of the unconstrained optimization problem (P_{un}). The criticality conditions for (BM_{ls}) are obtained by specializing (1).

Definition 9. Let $(\varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^2$. $Y \in \mathbb{R}^{n \times p}$ is $(\varepsilon_1, \varepsilon_2)$-*approximately 2-critical* (AC) for (BM_{ls}) if

$$\|S(YY^T)Y\| \leq \varepsilon_1, \tag{9a}$$

$$S(YY^T) \bullet UU^T + 4\|\mathcal{A}(UY^T)\|^2 \geq -\varepsilon_2 \quad (\forall U \in \mathbb{R}^{n \times p} : \|U\| = 1) \tag{9b}$$

We are ready to formalize the concept of spurious critical points.

Definition 10. Let $R_Y \in \mathbb{R}_+$ be fixed and let Y with $\|Y\| \leq R_Y$. Given $\varepsilon = (\varepsilon_0, \varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^3$, Y is *spurious* ε-AC if Y is $(\varepsilon_1, \varepsilon_2)$-AC for (BM_{ls}) but YY^T is not ε'-approx. optimal for (SDP_{ls}) for $\varepsilon' := (\varepsilon_0, R_Y\varepsilon_1, 5\varepsilon_2)$. Y is spurious *exactly* critical if the above holds for $\varepsilon = 0$.

We assume here that $b \in \mathbb{R}^m$ is fixed, and we vary the linear map $\mathcal{A}$. We identify $\mathcal{A}$ with the tuple of matrices $(A_1, \ldots, A_m)$, and hence view it as an element of the Euclidean space $(\mathbb{S}^n)^m$. We assume that $\mathcal{A}$ is *uniformly* distributed on the Euclidean ball $\mathbf{B}_\sigma(\bar{\mathcal{A}}) \subset (\mathbb{S}^n)^m$ of radius σ centered at $\bar{\mathcal{A}}$. Consider the set $\mathscr{A}_\varepsilon \subset (\mathbb{S}^n)^m$, consisting of all $\mathcal{A}$ for which there is a spurious AC point:

$$\mathscr{A}_\varepsilon \ := \ \{\mathcal{A} \in (\mathbb{S}^n)^m : \exists\, Y \text{ a spurious } \varepsilon\text{-AC point}\}.$$

We show that if $\tau(p) > m$, then the probability $\Pr[\mathcal{A} \in \mathscr{A}_\varepsilon] \to 0$ as the ratio $\varepsilon_1/\sqrt{\varepsilon_2} \to 0$.

Theorem 7 (critical $\Rightarrow$ feasible). *Let p such that $\tau(p) > m$, and $\varepsilon \in \mathbb{R}_+^3$. Let $\mathcal{A}$ be uniformly distributed on the Euclidean ball $\mathbf{B}_\sigma(\bar{\mathcal{A}})$. Then*

$$\Pr[\mathcal{A} \in \mathscr{A}_\varepsilon] \ \le \ 4e\, \delta^{\tau(p)-m}\, (3\kappa)^m \, (2n^3 m/\sigma\varepsilon_0)^{\tau(p)},$$

where $\delta := \varepsilon_1 R_A/\sqrt{\varepsilon_2}$, $\kappa := 2(R_A R_Y^2 + \|b\|)R_A$, and $R_A := \|\bar{\mathcal{A}}\| + \sigma$, provided that $\delta < \sigma\varepsilon_0/4n^3 m$.

As before, we can derive a high probability bound when $\tau(p) > (1+\eta)m$.

Corollary 2. *Consider the setup from Theorem 7. Assume that $\tau(p) \ge (1+\eta)m + \eta t$, $\varepsilon_0 \ge \rho\sigma$ and $\delta \le (1/3\kappa)^{1/\eta}(\rho\sigma^2/2n^3 m)^{1+1/\eta}$ for some $\eta, t, \rho > 0$. Then $\Pr[\mathcal{A} \in \mathscr{A}_\varepsilon] \ \le \ 4e\,(\rho\sigma^2/6\kappa n^3 m)^t$.*

As in Section 3, our proof of Theorem 5 relies on a geometric characterization of the spurious AC points. First consider the simpler case of spurious *exactly* critical points ($\varepsilon = 0$). The following equation is a consequence of our analysis:

$$\exists \text{ (spurious exactly critical point)} \quad \Longrightarrow \quad \mathbb{S}_{n-p}^n \cap \operatorname{Im} \mathcal{A}^* \text{ is nontrivial.}$$

This implies that the set $\mathscr{A}_0$ has measure zero when $\tau(p) > m$. In order to handle the case $\varepsilon > 0$ we rely again on tubular neighborhoods, as stated next.

Proposition 2. *Let $\delta := \varepsilon_1 R_A/\sqrt{\varepsilon_2}$, $D_\lambda := \{\lambda \in \mathbb{R}^m : 2\varepsilon_0 \le \|\lambda\| \le R_\lambda\}$, $R_\lambda := 2(R_A R_Y^2 + \|b\|)$. Then for $\mathcal{A} \in \mathscr{A}_\varepsilon$ we have that $\operatorname{tube}_\delta(\mathbb{S}_{n-p}^n) \cap \mathcal{A}^*(D_\lambda) \ne \emptyset$.*

5 Polynomial time guarantees

We are ready to derive polynomial time guarantees for the Burer-Monteiro method. The proof of Theorem 1 relies on two facts: (i) the method from Theorem 4 finds an AFAC point of (*BM*) in polynomial time; (ii) Corollary 1 shows that an AFAC point of (*BM*) is approximately optimal for (*SDP*) with high probability. Hence, the Burer-Monteiro method solves (*SDP*) in polynomial time in the setting of smooth analysis. The proof of Theorem 2 is analogous, except that it relies on Theorem 3 and corollary 2.

Appendix C provides explicit complexity estimates for solving (*SDP*) and (*SDP$_{ls}$*) using the Burer-Monteiro method, indicating the precise optimality guarantees of the output.

6 Experiments

We present some experimental results to complement our theorems. We rely on the library `NLopt` [24] (MIT license) to solve the nonlinear problem (*BM*). Concretely, we use the *augmented Lagrangian method* (ALM) implemented in `NLopt` (which is based on [8]), and we use the preconditioned truncated Newton method as the subroutine. We also rely on the commercial solver `Mosek` for SDPs.

For our first experiment we consider a random SDP with a planted solution. More precisely, we take a matrix $X_0 \in \mathbb{S}^{50}$, $X_0 \succeq 0$ of rank $r \in \{4, 7, 12\}$, and generate a random SDP for which X_0 is an optimal solution. To do so, we generate $m := \tau(r)$ random constraints that are satisfied at X_0, and then find a cost matrix C in the normal cone of X_0. We generate 100 random SDPs for each r. We solve these SDPs with the Burer-Monteiro method, using different values of p (the rank of Y) and random initializations. The initial points are matrices with i.i.d. normalized Gaussian entries. Figure 1 shows the percentage of experiments solved correctly for each value of r and p. We regard an experiment as "correct" if the criticality conditions from (4) are satisfied.

Figure 1 illustrates that there is a sharp phase transition at the Barvinok-Pataki bound $p = r$. Above the Barvinok-Pataki bound, the Burer-Monteiro method solves most instances. Beneath the Barvinok-Pataki bound, it is not just that our techniques stop working, but that the method itself usually

Figure 1: Performance of the BM method for random SDPs with a planted solution of rank r.

fails. Note that previous work [5, 30] required p to be larger than $3r/\sqrt{2}$. However, we see from experiments that the phase transition is much sharper than this. Observe that, even for $p \geq r$, the number of experiments solved is always below 100%. Nonetheless, the number of bad instances seems to get smaller for larger values of p. This agrees with our result from Theorem 5.

For the second experiment we fix the parameters $n := 50$, $m := 28$, and $p := r := 7$. Among the 100 random SDPs considered in Figure 1, we take an instance for which the Burer-Monteiro problem performed badly. We then perturb this seemingly bad SDP by adding varying amounts of noise σ. For each noise level we solve 70 random experiments, in which both the perturbations and the initializations are random. The perturbations consist in adding to each matrix a random matrix with i.i.d. Gaussian entries scaled by the noise level. Figure 2 summarizes the results obtained.

(a) Fraction of SDPs solved   (b) Iteration progress

Figure 2: Performance of the BM method after perturbing a "bad" SDP with different levels of noise.

Figure 2a shows the percentage of instances for which the Burer-Monteiro method succeeded for each noise level. The percentage is with respect to the random perturbation and the random initialization. For the unperturbed problem ($\sigma = 0$) the method succeeds only for 52% of the random initializations. This percentage increases as we add noise. For $\sigma = 0.2$ the method succeeds 87% of the time. Figure 2b shows the progress of the algorithm for the cases $\sigma = 0$ and $\sigma = 0.2$. The progress is measured in terms of the residual of the criticality conditions. The figure shows the mean and standard deviation of the residual for each iteration of ALM.

Figures 2a and 2b illustrate the advantages of smoothing a badly behaved SDP. Theorem 9 predicts that the complexity of the algorithm is proportional to σ^{-d} for some exponent d. So for a fixed number of iterations N, we should set the noise level proportional to $N^{-1/d}$. However, our bounds were shown for the algorithm from Theorem 4. We do not know if they also apply to ALM.

7 Discussion

We study the Burer-Monteiro method for solving the problems (*SDP*) and (*SDP$_{ls}$*). We provide the first polynomial time guarantees, using a smoothed analysis setting. Previous analyses for (*SDP*) did not provide end-to-end guarantees and made structural assumptions on the problem instance, while previous work for (*SDP$_{ls}$*) relied on the RIP condition, a strong assumption.

The main technical novelty of this paper is the use of volumes of tubular neighborhoods of varieties for deriving concentration bounds. This allows our results to hold even as p approaches the Barvinok-Pataki bound arbitrarily close. Although the Barvinok-Pataki bound is optimal for general SDPs, there are better bounds for some special classes of SDPs. An important open question is whether our results could be extended to allow for smaller values of p in those special classes.

The Burer-Monteiro method is not a single algorithm, but a family of algorithms, one for each local optimization method for solving (*BM*). Similarly, our guarantees in Theorems 1 and 2 apply for any local method with 2nd order guarantees. The only method we are aware that provably computes AFAC points of (*BM*) efficiently is the one in Theorem 4. It would be interesting to extend our results to classical optimization methods, such as ALM, which have robust implementations.

Acknowledgments and Funding Disclosure

Ankur Moitra was supported in part by a Microsoft Trustworthy AI Grant, NSF CAREER Award CCF-1453261, NSF Large CCF-1565235, a David and Lucile Packard Fellowship, an Alfred P. Sloan Fellowship and an ONR Young Investigator Award.

References

[1] Roberto Andreani, Ernesto G Birgin, José Mario Martínez, and María Laura Schuverdt. Second-order negative-curvature methods for box-constrained and general constrained optimization. *Comput. Optim. Appl.*, 45(2):209–236, 2010.

[2] Afonso S Bandeira, Nicolas Boumal, and Vladislav Voroninski. On the low-rank approach for semidefinite programs arising in synchronization and community detection. In *Proc. Conf. Learn. Theory*, pages 361–382, 2016.

[3] Alexander I. Barvinok. Problems of distance geometry and convex properties of quadratic maps. *Discrete Comput. Geom.*, 13(2):189–202, 1995.

[4] Saugata Basu and Antonio Lerario. Hausdorff approximations and volume of tubes of singular algebraic sets. *Mathematische Annalen*, 2022.

[5] Srinadh Bhojanapalli, Nicolas Boumal, Prateek Jain, and Praneeth Netrapalli. Smoothed analysis for low-rank solutions to semidefinite programs in quadratic penalty form. In *Conf. Learn. Theory*, pages 3243–3270, 2018.

[6] Srinadh Bhojanapalli, Behnam Neyshabur, and Nati Srebro. Global optimality of local search for low rank matrix recovery. In *Adv. Neural Inf. Process. Syst.*, pages 3873–3881, 2016.

[7] Ernesto G Birgin, JL Gardenghi, José Mario Martinez, Sandra A Santos, and Ph L Toint. Evaluation complexity for nonlinear constrained optimization using unscaled KKT conditions and high-order models. *SIAM J. Optim.*, 26(2):951–967, 2016.

[8] Ernesto G Birgin and José Mario Martínez. Improving ultimate convergence of an augmented Lagrangian method. *Optim. Methods Software*, 23(2):177–195, 2008.

[9] Nicolas Boumal, Vlad Voroninski, and Afonso Bandeira. The non-convex Burer-Monteiro approach works on smooth semidefinite programs. In *Adv. Neural Inf. Process. Syst.*, pages 2757–2765, 2016.

[10] Nicolas Boumal, Vladislav Voroninski, and Afonso Bandeira. Deterministic guarantees for Burer-Monteiro factorizations of smooth semidefinite programs. *Commun. Pure Appl. Math.*, 2019.

[11] Samuel Burer and Renato DC Monteiro. A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization. *Math. Program.*, 95(2):329–357, 2003.

[12] Samuel Burer and Renato DC Monteiro. Local minima and convergence in low-rank semidefinite programming. *Math. Program.*, 103(3):427–444, 2005.

[13] Peter Bürgisser, Felipe Cucker, and Martin Lotz. The probability that a slightly perturbed numerical analysis problem is difficult. *Math. Comput.*, 77(263):1559–1583, 2008.

[14] Coralia Cartis, Nicholas IM Gould, and Ph L Toint. Optimality of orders one to three and beyond: characterization and evaluation complexity in constrained nonconvex optimization. *J. Complexity*, 2018.

[15] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Adaptive cubic regularisation methods for unconstrained optimization. Part II: worst-case function-and derivative-evaluation complexity. *Math. Program.*, 130(2):295–319, 2011.

[16] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Complexity bounds for second-order optimality in unconstrained optimization. *J. Complexity*, 28(1):93–108, 2012.

[17] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. On the evaluation complexity of cubic regularization methods for potentially rank-deficient nonlinear least-squares problems and its relevance to constrained nonlinear optimization. *SIAM J. Optim.*, 23(3):1553–1574, 2013.

[18] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. On the complexity of finding first-order critical points in constrained nonlinear optimization. *Math. Program.*, 144(1-2):93–106, 2014.

[19] Diego Cifuentes. On the Burer–Monteiro method for general semidefinite programs. *Optimization Letters*, pages 1–11, 2021.

[20] Andrew R Conn, Nicholas IM Gould, and Ph L Toint. *Trust region methods*, volume 1 of *MOS-SIAM Series Optim.* SIAM, 2000.

[21] Frank E Curtis, Daniel P Robinson, and Mohammadreza Samadi. Complexity analysis of a trust funnel algorithm for equality constrained optimization. *SIAM J. Optim.*, 28(2):1533–1563, 2018.

[22] Rong Ge, Chi Jin, and Yi Zheng. No spurious local minima in nonconvex low rank problems: A unified geometric analysis. In *Proc. Int. Conf. Mach. Learn.*, pages 1233–1242, 2017.

[23] Alfred Gray. *Tubes*, volume 221. Birkhäuser, 2012.

[24] Steven G. Johnson. The NLopt nonlinear-optimization package. Available at `http://github.com/stevengj/nlopt`, 2019.

[25] Michel Journée, Francis Bach, P-A Absil, and Rodolphe Sepulchre. Low-rank optimization on the cone of positive semidefinite matrices. *SIAM J. Optim.*, 20(5):2327–2351, 2010.

[26] Qiuwei Li and Gongguo Tang. The nonconvex geometry of low-rank matrix optimizations with general objective functions. In *IEEE Global Conf. Signal Inf. Process.*, pages 1235–1239, 2017.

[27] Martin Lotz. On the volume of tubular neighborhoods of real algebraic varieties. *Proc. Am. Math. Soc.*, 143(5):1875–1889, 2015.

[28] Song Mei, Theodor Misiakiewicz, Andrea Montanari, and Roberto Imbuzeiro Oliveira. Solving SDPs for synchronization and maxcut problems via the Grothendieck inequality. In *Proc. Conf. Learn. Theory*, pages 1476–1515, 2017.

[29] Gábor Pataki. On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues. *Math. Oper. Res.*, 23(2):339–358, 1998.

[30] Thomas Pumir, Samy Jelassi, and Nicolas Boumal. Smoothed analysis of the low-rank approach for smooth semidefinite programs. In *Adv. Neural Inf. Process. Syst.*, pages 2287–2296, 2018.

[31] Mehmet Fatih Sahin, Armin Eftekhari, Ahmet Alacaoglu, Fabian Latorre, and Volkan Cevher. An inexact augmented Lagrangian framework for nonconvex optimization with nonlinear constraints. *Proc. Neural Inf. Process. Syst.*, 32:13965–13977, 2019.

[32] Daniel A Spielman and Shang-Hua Teng. Smoothed analysis of algorithms: Why the simplex algorithm usually takes polynomial time. *JACM*, 51(3):385–463, 2004.

[33] Jos F Sturm. Error bounds for linear matrix inequalities. *SIAM J. Optim.*, 10(4):1228–1248, 2000.

[34] Irene Waldspurger and Alden Waters. Rank optimality for the Burer–Monteiro factorization. *SIAM Journal on Optimization*, 30(3):2577–2602, 2020.

[35] Richard Zhang, Cédric Josz, Somayeh Sojoudi, and Javad Lavaei. How much restricted isometry is needed in nonconvex matrix recovery? In *Adv. Neural Inf. Process. Syst.*, pages 5586–5597, 2018.

A Computing AFAC points

In this appendix we prove Theorem 4, which states that AFAC points can be computed in polynomial time.

We focus on the constrained optimization problem (P_{con}) throughout this appendix. We will later need the following lemma, which gives a simple bound on the magnitude of the Lagrange multipliers of (P_{con}).

Lemma 4. *Let (y, λ) be such that $\|\nabla_y L(y, \lambda)\| \leq \varepsilon_1$. If ϱ-LICQ holds at y, then $\|\lambda\| \leq \varrho^{-1}(\varepsilon_1 + \|\nabla f(y)\|)$.*

Proof. Let $J := \nabla h(y)$. Since $\nabla L(y, \lambda) = \nabla f(y) + J^T \lambda$, and J is full rank, then $\lambda = (J^\dagger)^T (\nabla L(y, \lambda) - \nabla f(y))$, where $J^\dagger$ is the pseudo-inverse of J. Hence $\|\lambda\| \leq \varrho^{-1}(\varepsilon_1 + \|\nabla f(y)\|)$. $\square$

A.1 The algorithm

Cartis et al. [14] proposed a method for computing q-th order critical points for $q \in \{1, 2, 3\}$. However, they use a nonstandard notion of criticality which is not easy to translate into our setting. We present here a slight modification of this algorithm that accommodates more general criticality conditions.

Consider the least squares functions

$$\nu(y) := \|h(y)\|^2, \qquad \mu(t, y) := (f(y) - t)^2 + \|h(y)\|^2.$$

We denote $\mu_t = \mu(t, \cdot)$ the function obtained by fixing the value of t. Algorithm 1 below is a variant of the method from [14]. It consists of two phases. The first phase attempts to find an approximately feasible solution through the unconstrained problem $\min_y \nu(y)$. If successful, the second phase minimizes f while preserving feasibility. To do so, it solves a sequence of problems $\min_y \mu(t_k, y)$, where the values $\{t_k\}_{k \geq 0}$ are decreasing.

Algorithm 1 Constrained optimization algorithm based on [14]

Input: Initial point $y_0 \in \mathbb{R}^n$, tolerances $\epsilon_0 \in \mathbb{R}_+$, $\boldsymbol{\epsilon} \in \mathbb{R}_+^q$, constant $\delta \in (0, 1)$.
Output: A point $y \in \mathbb{R}^n$ and a number $t \leq f(y)$.

$\qquad\qquad\qquad\qquad\qquad$ PHASE I
$\quad y_1 := \text{local} \min_y \nu(y)$ starting with y_0
$\quad t_0 := f(y_1)$
$\quad$ **if** $\nu(y_1) > (\delta \epsilon_0)^2$ **then return** (y_1, t_0)

$\qquad\qquad\qquad\qquad\qquad$ PHASE II
$\quad t_1 := f(y_1) - (\epsilon_0^2 - \nu(y_1))^{1/2}$
$\quad$ **for** $k = 2, 3, 4, \ldots$ **do**
$\qquad y_k := \text{local} \min_y \mu(t_{k-1}, y)$ starting with y_{k-1}
$\qquad$ **if** $\mu(t_{k-1}, y_k) < (\delta \epsilon_0)^2$ **then** $\qquad\qquad\qquad\qquad\qquad\qquad$ ▷ case (a)
$\qquad\qquad t_k := f(y_k) - (\epsilon_0^2 - \nu(y_k))^{1/2}$
$\qquad\qquad$ **if** $\chi(\mu_{t_k}, y_k) \leq \boldsymbol{\epsilon}$ **then return** (y_k, t_k)
$\qquad$ **if** $\mu(t_{k-1}, y_k) \geq (\delta \epsilon_0)^2$ & $f(y_k) < t_{k-1}$ **then** $\qquad\qquad$ ▷ case (b)
$\qquad\qquad t_k := 2f(y_k) - t_{k-1}$
$\qquad\qquad$ **if** $\chi(\mu_{t_k}, y_k) \leq \boldsymbol{\epsilon}$ **then return** (y_k, t_k)
$\qquad$ **if** $\mu(t_{k-1}, y_k) \geq (\delta \epsilon_0)^2$ & $f(y_k) \geq t_{k-1}$ **then** $\qquad\qquad$ ▷ case (c)
$\qquad\qquad$ **return** (y_k, t_k), with $t_k := t_{k-1}$

Algorithm 1 relies on an *inner method* for solving the unconstrained problem $\min_y \psi(y)$, where ψ is either ν or $\mu_t = \mu(t, \cdot)$. Given $\boldsymbol{\epsilon} = (\epsilon_1, \ldots, \epsilon_q) \in \mathbb{R}_+^q$, the inner method looks for a point y such that $\chi(\psi, y) \leq \boldsymbol{\epsilon}$, for some *criticality measure* $\chi = (\chi_1, \ldots, \chi_q)$. We assume that the j-th component $\chi_j(\psi, y)$ only involves derivatives $\{\nabla^d \psi(y)\}_{d \leq j}$ up to order j. For instance, the AC-criticality condition from (1) corresponds to the case

$$\chi^{\text{AC}}(\psi, y) := \left(\|\nabla \psi(y)\|, -\min \text{eig}(\nabla^2 \psi(y)) \right). \tag{10}$$

Given an initial point y^0 and tolerances $\epsilon \in \mathbb{R}_+^q$, the inner method produces iterates $\{y^i\}_{i=1}^N$. We assume that the final point y^N achieves these tolerances and that the objective function decreases proportionately to N:

$$\chi(\psi, y^N) \leq \epsilon \qquad \text{and} \qquad \psi(y^0) - \psi(y^N) \geq N \, \kappa_\psi \, p(\epsilon), \tag{11}$$

for some $\kappa_\psi > 0$ and some function p. Hence, the number of iterations N is proportional to $p(\epsilon)^{-1}$.

The next theorem provides guarantees for Algorithm 1. Our proof closely follows that of [14, Thm.4.5] but has the advantage that it applies to a general class of criticality measures, as opposed to [14], which relies on a particular nonstandard measure of criticality. However our complexity is larger than in [14] by a factor of ϵ_0^{-1}.

Theorem 8. *Assume that:*

- *The inner method satisfies* (11) *for the function ν with constant κ_ν.*

- *The inner method satisfies* (11) *for the function μ_t, and the constant κ_μ is independent of t.*

- *There exists $\beta > \epsilon_0$ and $f_{\text{low}} \in \mathbb{R}$ such that $f(y) \geq f_{\text{low}}$ for all $y \in \mathfrak{M}_\beta$, where $\mathfrak{M}_\beta := \{y : \|h(y)\| \leq \beta\}$.*

Then the total number of inner iterations made in Algorithm 1 is at most

$$p(\epsilon)^{-1} \left(\kappa_\nu^{-1} \, \nu(y_0) \, + \, \epsilon_0 \, \kappa_\mu^{-1} \, (1-\delta)^{-1} \, (f(y_1) - f_{\text{low}} + \beta) \right), \tag{12}$$

and the algorithm returns a pair (y, t) such that:

either	$t < f(y),$	$\|h(y)\| \leq \epsilon_0,$	$\chi(\mu_t, y) \leq \epsilon,$	(13a)
or	$t = f(y),$	$\|h(y)\| > \delta\epsilon_0,$	$\chi_1(\nu, y) \leq \epsilon_1.$	(13b)

A.2 Proof of Theorem 8

Let K be the number of outer iterations of Algorithm 1. Consider the sets of indices:

$$A := \{1\} \cup \{k : 2 \leq k \leq K \text{ and case (a) is applied}\},$$
$$B := \{k : 2 \leq k \leq K \text{ and case (b) is applied}\}.$$

The following lemma gives a few properties of Algorithm 1. Its proof is identical to [14, Lem.3.1].

Lemma 5. *If the algorithm reaches Phase II, then:*

$$\nu(y_k) \leq \mu(t_k, y_k) \leq \epsilon_0^2, \qquad 0 \leq f(y_k) - t_k \leq \epsilon_0, \qquad \text{for } k \geq 1, \tag{14}$$
$$\mu(t_k, y_k) = \epsilon_0^2, \qquad t_{k-1} - t_k \geq (1-\delta)\epsilon_0, \qquad \text{for } k \in A, \tag{15}$$
$$\mu(t_k, y_k) = \mu(t_{k-1}, y_k), \qquad t_{k-1} > t_k, \qquad \text{for } k \in B, \tag{16}$$
$$\mu(t_k, y_k) \geq (\delta\epsilon_0)^2, \qquad \chi(\mu_{t_k}, y_k) \leq \epsilon, \qquad \text{for } k = K. \tag{17}$$

Let (y, t) be the output of Algorithm 1, and let us show (13). Assume first that the algorithm terminates in Phase I. Then y is a local minimum of ν, $\nu(y) > (\delta\epsilon_0)^2$, and $t = f(y)$. Hence (13b) holds. Assume now that the algorithm terminates in Phase II. By (14) and (17), we have

$$t \leq f(y), \qquad (\delta\epsilon_0)^2 \leq \mu_t(y) \leq \epsilon_0^2, \qquad \chi(\mu_t, y) \leq \epsilon.$$

If $f(y) < t$ then $\|h(y)\| \leq \sqrt{\mu_t(y)} \leq \epsilon_0$, so (13a) holds. Consider now the case that $f(y) = t$. Note that $\mu_t(y) = \nu(y)$, $\nabla \mu_t(y) = \nabla \nu(y)$. Then $\chi_1(\mu_t, y) = \chi_1(\nu, y)$, as they only involve derivatives up to order 1. Since $\|h(y)\| = \sqrt{\mu_t(y)} \geq \delta\epsilon_0$, then (13b) holds.

We proceed to show that the number of inner iterations is bounded by (12). Each outer iteration k of Algorithm 1 calls the inner method once. Let N_k be the number of inner iterations made in this call. The total number of inner iterations is $\sum_{k=1}^K N_k$.

We first analyze Phase I. The inner method is applied to the problem $\min_y \nu(y)$, starting with y_0 and terminating with y_1. By (11), we have

$$\nu(y_0) \geq \nu(y_0) - \nu(y_1) \geq N_1 \, \kappa_\nu \, p(\epsilon).$$

It follows that $N_1 \leq \nu(y_0)/\kappa_\nu p(\epsilon)$.

We proceed to Phase II. For each $a \in A$, let $n(a)$ be the next integer that lies in A. For the largest $a \in A$ we define $n(a) := K$, where K is the final iteration. We can group the indices $k \geq 2$ as follows:

$$\{2, 3, \ldots, K\} = \bigcup_{a \in A} K_a, \qquad K_a := \{a+1, a+2, \ldots, n(a)\}.$$

We will show that for any $a \in A$ we have that

$$N(K_a) := \sum_{k \in K_a} N_k \leq \epsilon_0^2/\kappa_\mu p(\epsilon). \tag{18}$$

Consider an iteration $k \in K_a$. The inner method is applied to $\min_y \mu(t_{k-1}, y)$, starting with y_{k-1} and terminating with y_k. By (11), we have

$$\mu(t_{k-1}, y_{k-1}) - \mu(t_{k-1}, y_k) \geq N_k \kappa_\mu p(\epsilon).$$

Observe that $K_a \setminus \{n(a)\} \subset B$. By (16), we have

$$\mu(t_{k-1}, y_k) = \mu(t_k, y_k) \quad \text{for} \quad k \in K_a \setminus \{n(a)\}.$$

Also note that $\mu(t_a, y_a) = \epsilon_0^2$ by (15). Therefore,

$$\begin{aligned}
\epsilon_0^2 &\geq \mu(t_a, y_a) - \mu(t_{n(a)}-1, y_{n(a)}) \\
&= \sum_{k \in K_a} \mu(t_{k-1}, y_{k-1}) - \mu(t_{k-1}, y_k) \geq \sum_{k \in K_a} N_k \kappa_\mu p(\epsilon).
\end{aligned}$$

By rearranging the above inequality we get (18).

Let us now upper bound the cardinality of A. By (15) and (16) we have that $t_{k-1}-t_k$ is at least $(1-\delta)\epsilon_0$ for $k \in A$, and is positive for $k \in B$. Also note that $t_0 = f(y_1)$ and $t_K \geq f(y_K)-\epsilon_0 \geq f_{\text{low}}-\beta$ by (14). Then

$$f(y_1)-f_{\text{low}}+\beta \geq t_0-t_K = \sum_{k=1}^{K}(t_{k-1}-t_k) \geq \sum_{k \in A}(t_{k-1}-t_k) \geq |A|\,(1-\delta)\epsilon_0,$$

and hence $|A| \leq (f(y_1)-f_{\text{low}}+\beta)/(1-\delta)\epsilon_0$.

Combining everything, we derive

$$\sum_{k=1}^{K} N_k \leq N_1 + |A| \cdot \max_{a \in A} N(K_a) \leq \frac{\nu(y_0)}{\kappa_\nu p(\epsilon)} + \frac{f(y_1)-f_{\text{low}}+\beta}{(1-\delta)\epsilon_0} \cdot \frac{\epsilon_0^2}{\kappa_\mu\, p(\epsilon)},$$

which is equal to (12).

A.3 Proof of Theorem 4

We finally show that AFAC points can be computed in polynomial time. Let $\varepsilon_0, \varepsilon_1, \varepsilon_2, \gamma, R_\lambda$ be as in the statement of Theorem 4. We consider Algorithm 1 with parameters

$$\delta := 1/2, \qquad q := 2, \qquad \boldsymbol{\epsilon} := (\epsilon_1, \epsilon_2),$$
$$\epsilon_0 := \varepsilon_0, \quad \epsilon_1 := R_\lambda^{-1}\varepsilon_0\varepsilon_1, \quad \epsilon_2 := \tfrac{1}{2}R_\lambda^{-1}\varepsilon_0\varepsilon_2.$$

For the inner method we use the ARC algorithm from Theorem 3, using the criticality measure (10). Algorithm 1 returns a pair (y, t). The associated multiplier is $\lambda := (f(y)-t)^{-1}\,h(y) \in \mathbb{R}^m$, which is defined only if $f(y) \neq t$.

In order to apply Theorem 8, we have to check that the functions ν and $\mu_t = \mu(t, \cdot)$ are smooth enough so that the inner algorithm satisfies (11).

Lemma 6 ([14, Lem.4.1]). *Assume that $\{\nabla^j f\}_{j=0}^q$, $\{\nabla^j h\}_{j=0}^q$ are uniformly bounded and Lipschitz continuous on a set $D \subset \mathbb{R}^n$. Then*

(i) $\{\nabla^j \nu\}_{j=0}^q$ are uniformly bounded and Lipschitz continuous on D.

(ii) $\{\nabla^j \mu_t\}_{j=0}^q$ are uniformly bounded and Lipschitz continuous on $D \cap B_t$, with $B_t := \{y : |f(y)-t|\leq 1\}$, and the constants are independent of t.

The above lemma shows that ν is smooth on $\mathfrak{M}_\beta$ and μ_t is smooth on $\mathfrak{M}_\beta \cap B_t$, with $B_t := \{y : |f(y)-t|\leq 1\}$. Note that all points y_k produced by Algorithm 1 lie in $\mathfrak{M}_\beta \cap B_t$ because of (14). Since ν, μ_t are sufficiently smooth, we can apply Theorem 3 (see also [15]). We conclude that the inner method satisfies (11) with

$$p(\boldsymbol{\epsilon}) = \min\{\epsilon_1^2, \epsilon_2^3\} = \Omega(\min\{\varepsilon_0^2 \varepsilon_1^2, \varepsilon_0^3 \varepsilon_2^3\}).$$

Hence, by Theorem 8, the total number of inner iterations is $O(p(\boldsymbol{\epsilon})^{-1}) = O(\max\{\varepsilon_0^{-2} \varepsilon_1^{-2}, \varepsilon_0^{-3} \varepsilon_2^{-3}\})$. Since each inner iteration requires $O(1)$ function evaluations (see Theorem 3), then the total number of function evaluations has the same order of magnitude.

Let us see that the conditions (2) hold. Let (y, t) be the output of Algorithm 1. By Theorem 8, this pair satisfies either (13a) or (13b). Let us see that (13b) cannot occur. Assume that

$$\|h(y)\| > \epsilon_0/2, \qquad \|\nabla\nu(y)\| \leq \epsilon_1.$$

Observe that $\|h(y)\|\leq \epsilon_0 \leq \beta$ by (14), and hence ϱ-LICQ holds at y. Then

$$\varrho\,\epsilon_0 < 2\,\varrho\|h(y)\| \leq 2\,\|h(y)^T\nabla h(y)\| = \|\nabla\nu(y)\| \leq \epsilon_1.$$

Also note that

$$R_\lambda^{-1} \leq \tfrac{1}{2}\varrho(1+L_f)^{-1} \leq \tfrac{1}{2}\varrho,$$
$$\epsilon_1/\epsilon_0 = R_\lambda^{-1}\varepsilon_1 \leq \tfrac{1}{2}\varrho\,\varepsilon_1 \leq \tfrac{1}{2}\varrho.$$

The last two equations give a contradiction.

Then the output (y, t) satisfies (13a). Hence, $t < f(y)$ and

$$\|h(y)\| \leq \epsilon_0, \quad \|\nabla\mu_t(y)\| \leq \epsilon_1, \quad \nabla^2\mu_t(y) \succeq -\epsilon_2 I_n.$$

Let $\alpha := (f(y)-t)^{-1}$, so that $\lambda = \alpha\,h(y)$. It can be checked that $\alpha^2 \mu_t(y) = \|(1, \lambda)\|^2$. Note that $\mu_t(y) \geq (\epsilon_0/2)^2$ by (17), and hence

$$\alpha = \mu_t(y)^{-1/2}\,\|(1, \lambda)\| \leq 2\varepsilon_0^{-1}\|(1, \lambda)\|.$$

The Lagrangian function $L(y, \lambda) = f(y)+\lambda \cdot h(x)$ is closely related to $\mu_t(y)$. A simple calculation gives that

$$\begin{aligned} \nabla L(y, \lambda) &= \alpha \cdot \tfrac{1}{2}\nabla\mu_t(y), \\ \nabla^2 L(y, \lambda) &= \alpha\left(\tfrac{1}{2}\nabla^2\mu_t(y) - \tilde{J}^T\tilde{J}\right), \end{aligned} \tag{19}$$

where $\tilde{J} := \begin{pmatrix} \nabla f(y) \\ \nabla h(y) \end{pmatrix}$ is the augmented Jacobian.

We proceed to verify (2a). We already have that $\|h(y)\| \leq \varepsilon_0$. Note that

$$\|\nabla L(y, \lambda)\| = \tfrac{1}{2}\alpha\,\|\nabla\mu_t(y)\| \leq \varepsilon_0^{-1}\|(1, \lambda)\|\epsilon_1 = R_\lambda^{-1}\|(1, \lambda)\|\varepsilon_1. \tag{20}$$

We claim that $\|(1, \lambda)\|\leq R_\lambda$. By Lemma 4 and $R_\lambda^{-1}\leq \varrho/2$, $\varepsilon_1 \leq 1$, we have

$$\|\lambda\| \leq \varrho^{-1}\,(R_\lambda^{-1}\varepsilon_1\|(1, \lambda)\| + \|\nabla f(y)\|) \leq \tfrac{1}{2}(1+\|\lambda\|) + \varrho^{-1}L_f.$$

It follows that $\|\lambda\| \leq 1+2\varrho^{-1}L_f = R_\lambda - 1$ and hence $\|(1, \lambda)\| \leq R_\lambda$, as we claimed. Then $\|\nabla L(y, \lambda)\| \leq \varepsilon_1$ by (20).

We now verify (2b). Let $u \in \mathbb{R}^n$ of unit norm such that $\|Ju\| \leq \gamma$, where $J := \nabla h(y)$. We need to show that $u^T\nabla^2 L(y, \lambda)u \geq -\varepsilon_2$. By (19), we have

$$u^T\nabla^2 L(y, \lambda)u = \alpha\left(\tfrac{1}{2}u^T\nabla^2\mu_t(y)u - \|\tilde{J}u\|^2\right). \tag{21}$$

Note that $u^T\nabla^2\mu_t(y)u \geq -\epsilon_2 = -\tfrac{1}{2}R_\lambda^{-1}\varepsilon_0\varepsilon_2$. We bound $\|\tilde{J}u\|$ next:

$$\tilde{J} = \begin{pmatrix} \nabla f(y) \\ \nabla h(y) \end{pmatrix} = \begin{pmatrix} \nabla L(y, \lambda) \\ 0 \end{pmatrix} + \begin{pmatrix} -\lambda^T J \\ J \end{pmatrix},$$

$$\|\tilde{J}u\| \leq \|\nabla L(y, \lambda)\| + \|(1, \lambda)\|\,\|Ju\| \leq \varepsilon_1 + \gamma\,\|(1, \lambda)\| \leq \tfrac{1}{2}(R_\lambda^{-1}\varepsilon_0\varepsilon_2)^{1/2},$$

where we used that ε_1 and γR_λ are at most $\tfrac{1}{4}(R_\lambda^{-1}\varepsilon_0\varepsilon_2)^{1/2}$ by (3). Hence

$$\alpha\left(\tfrac{1}{2}u^T\nabla^2\mu_t(y)u - \|\tilde{J}u\|^2\right) \geq -(2\varepsilon_0^{-1}\|(1, \lambda)\|) \cdot (\tfrac{1}{2}R_\lambda^{-1}\varepsilon_0\varepsilon_2) \geq -\varepsilon_2.$$

Together with (21), we get that $u^T\nabla^2 L(y, \lambda)u \geq -\varepsilon_2$.

B Proofs from Section 4

Proof of Lemma 3. Let $L(X) := f(X) - \bar{S} \bullet X$, with $\bar{S} := S(\bar{X})$. This is a convex function with $\nabla L(\bar{X}) = 0$, so $\bar{X}$ is its global minimum. Note that

$$
\begin{aligned}
f(X) &= L(X) + \bar{S} \bullet X \geq L(X) - (\varepsilon_2 I_n) \bullet X \geq L(X) - \varepsilon_2 \|X\| \sqrt{n}, \\
L(\bar{X}) &= f(\bar{X}) - \bar{S} \bullet \bar{X} \geq f(\bar{X}) - \|\bar{S}\bar{X}\|_* \geq f(\bar{X}) - \varepsilon_1 \sqrt{n}.
\end{aligned}
$$

Since $L(X) \geq L(\bar{X})$, the result follows from the above equations. □

The next lemma is an analogue of Lemma 2.

Lemma 7. *Let Y be an $(\varepsilon_1, \varepsilon_2)$-AC point of (BM_{ls}). If $\sigma_p(Y) \leq \sqrt{\varepsilon_2}/R_A$, then YY^T is ε'-approximately optimal for (SDP_{ls}), with $\varepsilon' := (0, R_Y \varepsilon_1, 5\varepsilon_2)$.*

Proof. Let Y satisfy (9), and let us show that YY^T satisfies (8). The first-order condition is easy to check. We proceed to show that $u^T S(X) u \geq -\varepsilon_2'$ for any unit vector $u \in \mathbb{R}^n$. Let $z \in \mathbb{R}^p$ be a unit vector such that $\|Yz\| = \sigma_p(Y)$. The matrix $U := uz^T$ satisfies $\|U\| = 1$ and $\|UY^T\| \leq \|u\| \|Yz\| = \sigma_p(Y)$. Then $\|\mathcal{A}(UY^T)\| \leq \sqrt{\varepsilon_2}$ and by (9b) we have

$$
u^T S(X) u = S(YY^T) \bullet UU^T \geq -\varepsilon_2 - 4\|\mathcal{A}(UY^T)\|^2 \geq -5\varepsilon_2. \qquad \square
$$

Proof of Proposition 2. As $\mathcal{A} \in \mathscr{A}_{\boldsymbol{\varepsilon}}$, there is a spurious $\boldsymbol{\varepsilon}$-AC point Y. By Lemma 7, we must have $\sigma_p(Y) > \sqrt{\varepsilon_2}/R_A$. Note that $\|S(YY^T)Y\| \leq \varepsilon_1$. Together with (6), we conclude that $S(YY^T) \in \text{tube}_\delta(\mathbb{S}_{n-p}^n)$. Let $\lambda := 2(\mathcal{A}(YY^T) - b)$. Note that $\|\lambda\| > 2\varepsilon_0$ and

$$
\|\lambda\| = 2\|\mathcal{A}(YY^T) - b\| \leq 2(\|\mathcal{A}\| \|Y\|^2 + \|b\|) \leq R_\lambda.
$$

Then $\lambda \in D_\lambda$ and $\mathcal{A}^*(\lambda) = S(YY^T) \in \text{tube}_\delta(\mathbb{S}_{n-p}^n)$. □

Proof of Theorem 7. The result in Proposition 2 can be expressed as:

$$
\mathscr{A}_{\boldsymbol{\varepsilon}} \subset \{\mathcal{A} \in (\mathbb{S}^n)^m : 0 \in \text{tube}_\delta(\mathbb{S}_{n-p}^n) + \mathcal{A}^*(D_\lambda)\},
$$

which is closer to the formula in Proposition 1. Consider an ϵ-net $\mathcal{N}$ of D_λ, where $\epsilon := \delta/R_A$. It suffices to take $(3R_\lambda/\epsilon)^m = (3\kappa/\delta)^m$ points for the ϵ-net. A reasoning similar to (7) gives

$$
\begin{aligned}
\mathscr{A}_{\boldsymbol{\varepsilon}} &\subset \{\mathcal{A} \in (\mathbb{S}^n)^m : 0 \in \text{tube}_{2\delta}(\mathbb{S}_{n-p}^n) + \mathcal{A}^*(\mathcal{N}))\} \\
&= \bigcup_{\ell \in \mathcal{N}} \{\mathcal{A} \in (\mathbb{S}^n)^m : \mathcal{A}^*(\ell) \in \text{tube}_{2\delta}(\mathbb{S}_{n-p}^n))\}.
\end{aligned}
$$

Let $\ell \in \mathcal{N}$, and consider the linear map

$$
\phi_\ell : (\mathbb{S}^n)^m \to \mathbb{S}^n, \quad \mathcal{A} \mapsto \mathcal{A}^*(\ell).
$$

This is a surjective map. Moreover, the scaled map $\frac{1}{\|\ell\|} \phi_\ell$ gives an isometry $(\ker \phi_\ell)^\perp \cong \mathbb{S}^n$. It follows that

$$
\phi_\ell(\mathcal{A}) \in \text{tube}_{2\delta}(\mathbb{S}_{n-p}^n) \iff \mathcal{A} \in \text{tube}_{2\delta/\|\ell\|}\left(\phi_\ell^{-1}(\mathbb{S}_{n-p}^n)\right).
$$

Since $\|\ell\| \geq 2\varepsilon_0$, we conclude that

$$
\mathscr{A}_{\boldsymbol{\varepsilon}} \subset \bigcup_{\ell \in \mathcal{N}} \text{tube}_{\delta/\varepsilon_0}(V_\ell), \quad \text{with } V_\ell := \phi_\ell^{-1}(\mathbb{S}_{n-p}^n).
$$

The final part of the proof is similar to the one in Theorem 5. The variety V_ℓ is a cylinder over $\mathbb{S}_{n-p}^n$, so it has the same codimension $\tau(p)$ and degree $n-p+1$ as $\mathbb{S}_{n-p}^n$, The ambient space is $(\mathbb{S}^n)^m$, of dimension $\tau(n)m$. Using the union bound and Theorem 6, we get

$$
\Pr[\mathcal{A} \in \mathscr{A}_{\boldsymbol{\varepsilon}}] < \#\mathcal{N} \cdot \Pr[\mathcal{A} \in \text{tube}_{\delta/\varepsilon_0}(V_\ell)] < (3\kappa/\delta)^m \cdot 4e \left(2n^3 m\delta/\sigma\varepsilon_0\right)^{\tau(p)}. \qquad \square
$$

C Explicit complexity estimates

In this section we provide explicit complexity estimates for Theorems 1 and 2. We first introduce some notation. Consider constants $\alpha \geq \beta > 0$ and sets $\mathfrak{M}_\alpha \supset \mathfrak{M}_\beta$, where $\mathfrak{M}_t := \{Y : \|\mathcal{A}(YY^T) - b\| \leq t\}$. We assume that β is small enough so that ϱ-LICQ holds globally on $\mathfrak{M}_\beta$. On the other hand, $\alpha > 0$ is sufficiently large so that a point $Y_0 \in \mathfrak{M}_\alpha$ is always known. We further assume that $\mathfrak{M}_\alpha$ is compact. This is satisfied, for instance, when the feasible set of (*SDP*) is compact and satisfies Slater's condition, as shown next.

Lemma 8. *Assume that the set $\{X : \mathcal{A}(X) = b, X \succeq 0\}$ is compact and satisfies Slater's condition (i.e., $\exists X : \mathcal{A}(X) = b, X \succ 0$). Then $\mathcal{M}_t$ is compact for any $t \geq 0$.*

Proof. Consider the SDP $\max\{I \bullet X : \mathcal{A}(X) = b, X \succeq 0\}$ and its dual $\min\{b^T \lambda : \mathcal{A}^*(\lambda) \succeq I\}$. Let R_0^2 be the primal optimal value, which is finite by compactness. Strong duality holds by Slater's condition. So the dual optimum is attained at some $\bar{\lambda}$, and $\mathcal{A}^*(\bar{\lambda}) \succeq I$, $b^T\bar{\lambda} = R_0^2$. Given $Y \in \mathcal{M}_t$,

$$\|Y\|^2 = I \bullet YY^T \leq \mathcal{A}^*(\bar{\lambda}) \bullet YY^T = \bar{\lambda} \cdot \mathcal{A}(YY^T) = \bar{\lambda} \cdot b + \bar{\lambda} \cdot (\mathcal{A}(YY^T) - b) \leq R_0^2 + t\|\bar{\lambda}\|.$$

We conclude that $\mathcal{M}_t$ is contained in a ball of radius $(R_0^2 + t\|\bar{\lambda}\|)^{1/2}$. $\qquad\square$

Notice that a suitable value α can be obtained from an arbitrary point $Y_0 \in \mathbb{R}^{n \times p}$. On the other hand, β should be $\Omega(\varrho_{\min})$, where $\varrho_{\min}$ is the smallest LICQ constant among all feasible points $Y \in \mathfrak{M}_0$.

C.1 Solving (*SDP*)

Assume that an approximately feasible solution Y_0 is known. Consider the following setting:

- p satisfies $\tau(p) \geq (1+\eta)m + \eta t$ for some given constants $\eta, t \in \mathbb{R}_+$.
- $\mathcal{A}, b$ are fixed and C is uniformly distributed on a ball $\mathbf{B}_\sigma(\bar{C})$.
- $\exists \beta \in \mathbb{R}_+$ such that: $\mathfrak{M}_\beta$ is compact, a point $Y_0 \in \mathfrak{M}_\beta$ is known, and ϱ-LICQ holds on $\mathfrak{M}_\beta$.
- $R_Y, L_f \in \mathbb{R}_+$ are constants that bound $\|Y\|$ and $\|CY\|$, for $Y \in \mathfrak{M}_\beta$.
- (*BM*) is solved with the method from Theorem 4 initialized at Y_0.

The next theorem shows that the Burer-Monteiro method solves (*SDP*) in polynomial time with high probability.

Theorem 9. *Let $\rho \in (0, 1]$ arbitrary, and let*

$$\varepsilon_0 := \gamma := \epsilon, \quad \varepsilon_1 := \epsilon^2, \quad \varepsilon_2 := 16\,R_\lambda^3\,\epsilon,$$
$$\text{with} \quad \epsilon := K^{-1}\,\rho\,(\sigma/4n^3)^{1+1/\eta},$$

where R_λ and K are the problem dependent constants

$$R_\lambda := 2 + 2\varrho^{-1}L_f, \quad K := \|\mathcal{A}\|\,(3\kappa)^{1/\eta}, \quad \kappa := R_\lambda\|\mathcal{A}\|.$$

The algorithm from Theorem 4 returns a pair (Y, λ) after $O(\epsilon^{-6})$ function evaluations. With probability at least $1 - O(\sigma/n^3)^t$, the pair (YY^T, λ) is $(\epsilon, \epsilon^2 R_Y, 16R_\lambda^3 \epsilon)$-approximately optimal for (SDP).

Proof. The smoothness assumptions in Theorem 4 are satisfied since $\mathfrak{M}_\beta$ is compact. Then (Y, λ) is an (ε, γ)-AFAC pair with $\|\lambda\| \leq R_\lambda$. Note that

$$\delta := \varepsilon_1\|\mathcal{A}\|/\gamma = \epsilon\,\|\mathcal{A}\| \leq (1/3\kappa)^{1/\eta}\,(\sigma/2en^3)^{1+1/\eta}$$

is as in Corollary 1. Hence (YY^T, λ) is $(\varepsilon_0, \varepsilon_1 R_Y, \varepsilon_2)$-approximately optimal for (*SDP*) with probability $1 - O(\sigma/n^3)^t$. $\qquad\square$

The above theorem shows that YY^T obtained is approximately optimal for the perturbed problem (*SDP*) with high probability. Let $(\overline{SDP})$ denote the SDP problem in which we use the unperturbed cost matrix $\bar{C}$. We can also show that YY^T is also approximately optimal for $(\overline{SDP})$.

Corollary 3. *Consider the setup of Theorem 9. With probability at least $1 - O(\sigma/n^3)^t$, the pair (YY^T, λ) is $(\varepsilon_0'', \varepsilon_1'', \varepsilon_2'')$-approximately optimal for $(\overline{SDP})$, where $\varepsilon_0'', \varepsilon_1'', \varepsilon_2'' = O(\sigma)$.*

Proof. Let $X := YY^T$. We know that (X, λ) is $(\varepsilon'_0, \varepsilon'_1, \varepsilon'_2)$-approximately optimal for *(SDP)* with high probability. Let $S := C - \mathcal{A}^*(\lambda)$, $\bar{S} := \bar{C} - \mathcal{A}^*(\lambda)$ be the slack matrices for *(SDP)* and $(\overline{SDP})$. Observe that

$$\|\mathcal{A}(X) - b\| \leq \varepsilon'_0 \leq O(\sigma), \tag{22}$$

$$\|\bar{S}X\| \leq \|SX\| + \|(\bar{S}-S)X\| \leq \varepsilon'_1 + \sigma\|X\| \leq O(\sigma), \tag{23}$$

$$\bar{S} \succeq S - \|\bar{S}-S\|I_n \succeq -(\varepsilon'_2+\sigma)I_n \succeq -O(\sigma)I_n. \tag{24}$$

So the optimality conditions of $(\overline{SDP})$ hold with $\varepsilon''_0, \varepsilon''_1, \varepsilon''_2 = O(\sigma)$. $\qquad\square$

C.2 Solving (SDP_{ls})

Consider the following setting:

- p satisfies $\tau(p) \geq (1+\eta)m + \eta\, t$ for some given constants $\eta, t \in \mathbb{R}_+$.
- b is fixed and $\mathcal{A}$ is uniformly distributed on a ball $\mathbf{B}_\sigma(\bar{\mathcal{A}})$.
- $\exists \alpha \in \mathbb{R}_+$ and a matrix Y_0 such that $\mathfrak{M}_\alpha$ is compact and $Y_0 \in \mathfrak{M}_\alpha$
- $R_Y \in \mathbb{R}_+$ is a constant that bounds $\|Y\|$, for $Y \in \mathfrak{M}_\alpha$.
- (BM_{ls}) is solved with the method from Theorem 3 initialized at Y_0.

Theorem 10. *Let $\rho \in (0, 1]$ arbitrary, and let*

$$\varepsilon_1 := \epsilon^{3/2}, \quad \varepsilon_2 := \epsilon, \quad \epsilon := K^{-1}\left(\rho\,\sigma^2/2n^3m\right)^{1+1/\eta},$$

where $K := R_A\,(3\kappa)^{1/\eta}$, expressed in terms of

$$\kappa := 2(R_A R_Y^2 + \|b\|)R_A, \quad R_A := \|\bar{\mathcal{A}}\| + \sigma.$$

The algorithm from Theorem 3 returns a point Y after $O(\epsilon^{-3})$ function evaluations. With probability at least $1 - O(\sigma^2/n^3m)^t$, we have that YY^T is $(\rho\,\sigma, \epsilon^{3/2}R_Y, 5\epsilon)$-approximately optimal for (SDP_{ls}).

Proof. The smoothness assumptions in Theorem 3 are satisfied since $\mathfrak{M}_\alpha$ is compact. Therefore Y is an $(\varepsilon_1, \varepsilon_2)$-AC point. Note that

$$\delta := \varepsilon_1 R_A/\sqrt{\varepsilon_2} = \epsilon R_A = (1/3\kappa)^{1/\eta}(\rho\,\sigma^2/2n^3m)^{1+1/\eta}$$

is as in Corollary 2. Hence YY^T is $(\rho\,\sigma, \varepsilon_1 R_Y, 5\varepsilon_2)$-approximately optimal for (SDP_{ls}) with probability $1 - O(\sigma^2/n^3m)^t$. $\qquad\square$

Remark. The above theorem holds even if the optimal value of (SDP_{ls}) is nonzero. In the special case that the optimal value is zero, then by Lemma 3 we have that

$$\|\mathcal{A}(YY^T) - b\| \leq \max\{\varepsilon'_0,\ n^{1/4}(\varepsilon'_1 + \varepsilon'_2 R_Y)^{1/2}\},$$

where $\varepsilon'_0 = \rho\,\sigma, \varepsilon'_1 = \epsilon^{3/2}R_Y, \varepsilon'_2 = 5\epsilon$ are the optimality constants from Theorem 10.

Let $(\overline{SDP}_{ls})$ denote the instance of problem (SDP_{ls}) in which we use the unperturbed constraints $\bar{\mathcal{A}}$. We next show that YY^T is also approximately optimal for $(\overline{SDP}_{ls})$.

Corollary 4. *Consider the setup of Theorem 10. With probability at least $1 - O(\sigma^2/n^3m)^t$, the matrix YY^T is $(\varepsilon''_0, \varepsilon''_1, \varepsilon''_2)$-approximately optimal for $(\overline{SDP}_{ls})$, where $\varepsilon''_0, \varepsilon''_1, \varepsilon''_2 = O(\sigma)$.*

Proof. We know that the matrix $X := YY^T$ is $(\varepsilon'_0, \varepsilon'_1, \varepsilon'_2)$-approximately optimal for *(SDP)* with high probability. There are two cases. The first case is that $\|\mathcal{A}(X) - b\| \leq \varepsilon'_0$, which implies that

$$\|\bar{\mathcal{A}}(X) - b\| \leq \|\mathcal{A}(X) - b\| + \|(\bar{\mathcal{A}}-\mathcal{A})X\| \leq \varepsilon'_0 + \sigma\|X\| \leq O(\sigma).$$

This means that $\varepsilon''_0 = O(\sigma)$. Consider the variables:

$$\lambda := 2(\mathcal{A}(X) - b), \quad S := \mathcal{A}^*(\lambda),$$

$$\bar{\lambda} := 2(\bar{\mathcal{A}}(X) - b), \quad \bar{S} := \bar{\mathcal{A}}^*(\bar{\lambda}).$$

The second case is that $\|SX\| \leq \varepsilon_1, S \succeq -\varepsilon_2 I_n$. Note that

$$\|\bar{\lambda} - \lambda\| \leq 2\|\bar{\mathcal{A}} - \mathcal{A}\|\,\|X\| \leq O(\sigma),$$

$$\|\bar{S} - S\| \leq \|\bar{\mathcal{A}}^*(\bar{\lambda} - \lambda)\| + \|(\bar{\mathcal{A}}^* - \mathcal{A}^*)\lambda\| \leq O(\sigma).$$

From (23) and (24) we get that $\|\bar{S}X\| \leq O(\sigma)$ and $\bar{S} \succeq -O(\sigma)I_n$. So the optimality conditions of $(\overline{SDP}_{ls})$ hold with $\varepsilon''_0, \varepsilon''_1, \varepsilon''_2 = O(\sigma)$. $\qquad\square$