REVISITING BISIMULATION: A SAMPLING-BASED STATE SIMILARITY PSEUDO-METRIC

1 INTRODUCTION

In reinforcement learning (RL), we typically deal with systems with large or continuous states encoded in an unstructured way. Because it is not possible to represent the value of each state, it is necessary to learn a structured representation from which we can express the value function in a more meaningful way. One approach to do so is to endow the set of states with a behavioral metric, such that two states that are close in the metric space are also close in the space of value functions. While there exists some notions of state similarity, they are either not amenable to sample-based algorithms (Ferns et al., 2004; 2005), need additional assumptions (Castro, 2020; Zhang et al., 2020; Agarwal et al., 2021) or yield limited theoretical guarantees (Castro et al., 2021). In this paper, we present a new behavioural pseudo-metric, PMiCo, to overcome these shortcomings. PMiCo is based on a recent sampling-based behavioural distance, MICo (Matching under Independent Couplings; Castro et al., 2021), but enjoys more interesting theoretical properties, which we also illustrate empirically.

2 BACKGROUND

Reinforcement learning. We consider a Markov decision process (MDP) $\mathcal{M} = \langle \mathcal{X}, \mathcal{A}, r, \mathcal{P}, \gamma \rangle$ (Puterman, 1994) with finite state space $\mathcal{X}$, finite action space $\mathcal{A}$, transition dynamics $\mathcal{P} : \mathcal{X} \times \mathcal{A} \to \mathscr{P}(\mathcal{X})$, reward function $r : \mathcal{X} \times \mathcal{A} \to \mathbb{R}$, and discount factor $\gamma \in [0, 1)$. A stationary policy $\pi : \mathcal{X} \to \mathscr{P}(\mathcal{A})$ is a mapping from states to distributions over actions. We denote the set of all policies by Π. For any policy $\pi \in \Pi$, the value function $V^\pi(x)$ measures the expected discounted sum of rewards received when starting from state $x \in \mathcal{X}$ and acting according to π and satisfies Bellman's equation $V^\pi(x) := \mathbb{E}_{a \sim \pi(x)} \left[r_x^a + \gamma \mathbb{E}_{x' \sim P_x^a} \left[V^\pi(x') \right] \right]$. In RL, the goal is to find the optimal policy $\pi^* = \operatorname{argmax}_{\pi' \in \Pi} V^{\pi'}(x)$ for all state $x \in \mathcal{X}$ from interactions with *sample trajectories* with the environment.

Bisimulation. The *bisimulation metric* (Ferns et al., 2004) describes two states as similar if their immediate rewards are close and they transition to next states which are also considered similar. More formally, denoting $\mathcal{M}(\mathcal{X}) = \{d \in [0, \infty)^{\mathcal{X} \times \mathcal{X}} \mid \forall x, y, z \in \mathcal{X}, d(x, x) = 0; d(x, y) = d(y, x); d(x, y) \le d(x, z) + d(z, y)\}$ the set of *pseudo-metrics* on $\mathcal{X}$, the bisimulation metric $d^\sim$ is defined as the unique fixed-point of the operator $T_W : \mathcal{M}(X) \to \mathcal{M}(X)$ where $T_W(d)(x, y) = \max_{a \in \mathcal{A}} [|r(x, a) - r(y, a)| + \gamma W_d (P(x, a), P(y, a))]$. Here, W_d is the 1-Wasserstein metric (Villani, 2008) and describes the minimal cost of transporting probability mass from $\mu \in \mathscr{P}(\mathcal{X})$ to $\nu \in \mathscr{P}(\mathcal{X})$. Castro (2020) extended the canonical bisimulation to an on-policy counterpart. The π-*bisimulation* $d_\sim^\pi$ is the fixed point of the operator $T_W^\pi(d)(x, y) = \left| r_x^\pi - r_y^\pi \right| + \gamma \mathcal{W}_d \left(P_x^\pi, P_y^\pi \right)$ with $r_x^\pi = \sum_{a \in \mathcal{A}} \pi(a \mid x) r_x^a$, $P_x^\pi = \sum_{a \in \mathcal{A}} \pi(a \mid x) P(x, a)(\cdot)$ for all $x \in \mathcal{X}$. V^π (resp. V^*) is Lipschitz continuous with respect to $d_\sim^\pi$ (resp. $d^\sim$), that is $|V^\pi(x) - V^\pi(y)| \le d_\sim^\pi(x, y)$ for any $x, y \in \mathcal{X}$ and any $\pi \in \Pi$.

More recently, Castro et al. (2021) introduced the concept of *MICo distance* U^π as the fixed point of the operator $T_M^\pi : \mathbb{R}^{\mathcal{X} \times \mathcal{X}} \to \mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ where $(T_M^\pi U)(x, y) = \left| r_x^\pi - r_y^\pi \right| + \gamma \mathbb{E}_{x' \sim P_x^\pi, y' \sim P_y^\pi} [U(x', y')]$ for all $U : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$. The MICo operator **does not** define a pseudo-metric as there exists states $x \in \mathcal{X}$ such that $d(x, x) \ne 0$. Since most algorithmic methods used to measure distances need the property of zero self-distance, Castro et al. (2021) rely in practice on the projection ΠU^π of the MICo distance.

DISTANCE	SAMPLE-BASED	PSEUDO-METRIC	V^π-LIPSCHITZ CONTINUITY
π-BISIMULATION	✗	✓	✓
MICO	✓	✗	✓
PROJECTED MICO	✓	✗	✗
PMICO	✓	✓	✓

Table 1: Categorization of behavioural distances according to their sample-based definition, pseudo-metric properties and continuity implications.

3   THE PMICO PSEUDO-METRIC

As an alternative to the above distances, we introduce a new (pseudo) metric, inspired by the MICo distance, which we call PMiCo (**P**seudo-metric **M**atching under **i**ndependent **C**ouplings). This metric is desirable because it can be learnt from transition samples and yields strong theoretical guarantees. Given $\pi \in \Pi$, the PMiCo update operator $T_P^\pi : \mathbb{R}^{\mathcal{X} \times \mathcal{X}} \to \mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ is

$$(T_P^\pi U)(x,y) = |r^\pi(x) - r^\pi(y)| + \mathbf{1}_{[x \neq y]} \gamma \mathbb{E}_{x' \sim P_x^\pi, y' \sim P_y^\pi}[U(x',y')] \tag{1}$$

where $\mathbf{1}$ is the indicator function. The PMiCo distance between two distinct points is always positive, making it an appealing candidate for function approximation, without the need of a projection.

Theorem 1. *The PMiCo operator T_P^π is a contraction mapping on $\mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ w.r.t. L^∞ norm and its fixed point U_P^π is a pseudo-metric.*

One question that naturally arises is what are the guarantees of a representation induced by the PMiCo metric. Similarly to the MICo distance, we have the following on-policy guarantee for PMiCo.

Theorem 2. *For any policy $\pi \in \Pi$ and states $x, y \in \mathcal{X}$, $|V^\pi(x) - V^\pi(y)| \leq U_P^\pi(x,y) \leq U^\pi(x,y)$.*

All the proofs can be found in Appendix A. We summarize in Table 1 the properties of the distances mentioned in section 2 and section 3.

We now conduct an empirical evaluation to illustrate the correctness of Theorem 2. Following Castro et al. (2021), we use *Garnet MDPs* (Archibald et al., 1995; Piot et al., 2014). For a fixed number of states and actions, we sample 100 stochastic policies $\{\pi_i\}$ and measure the average signed gap $\frac{1}{100|\mathcal{X}|^2} \sum_i \sum_{x,y} d(x,y) - |V^{\pi_i}(x) - V^{\pi_i}(y)|$. We consider four distances: the PMiCo metric, the MICo distance, the projected MICo distance and the π-bisimulation. Figure 1 clearly shows that the PMiCo metric yields a tighter bound than the MICo distance on average, although it is larger than the π-bisimulation and projected MICo distances. However, unlike PMiCo, π-bisimulation is biased when approximated using only sampled transitions while projected MICo can be negative and may not upper bound the value differences.

Figure 1: Gap between the difference in values and various distances for Garnet MDPs with 3 actions. Shaded areas represent 95 % confidence intervals.

4   DISCUSSION

In this paper, we presented PMiCo, a novel sample-based pseudo-metric. We saw it supports the Lipschitz continuity of the value function and induces a more coarse representation on the state space than the previously introduced MICo distance (Castro et al., 2021), suggesting it can lead to better generalization (Le Lan et al., 2021). Our experiments on Garnet MDPs show the tighter bound yielded by the PMiCo metric. In deep RL, behavioural metrics are often incorporated into the learning process as auxiliary tasks, to help shape the representation learnt by the agent (Gelada et al., 2019; Agarwal et al., 2021; Zhang et al., 2020; Castro et al., 2021). As such, training a network on auxiliary predictions such that the distance between two latent states corresponds to PMiCo could be a simple and effective auxiliary task for deep RL.

URM STATEMENT

The authors acknowledge that at least one key author of this work meets the URM criteria of ICLR 2023 Tiny Papers Track.

REFERENCES

Rishabh Agarwal, Marlos C. Machado, Pablo Samuel Castro, and Marc G. Bellemare. Contrastive behavioral similarity embeddings for generalization in reinforcement learning. In *International Conference on Learning Representations*, 2021.

T. W. Archibald, K. I. M. McKinnon, and L. C. Thomas. On the generation of Markov decision processes. *The Journal of the Operational Research Society*, 46(3):354–361, 1995.

Pablo Samuel Castro. Scalable methods for computing state similarity in deterministic markov decision processes. In *Proceedings of the AAAI Conference on Artificial Intelligence*, 2020.

Pablo Samuel Castro, Tyler Kastner, Prakash Panangaden, and Mark Rowland. Mico: Improved representations via sampling-based state similarity for markov decision processes. *Advances in Neural Information Processing Systems*, 34:30113–30126, 2021.

Norm Ferns, Prakash Panangaden, and Doina Precup. Metrics for finite markov decision processes. In *Proceedings of the 20th conference on Uncertainty in artificial intelligence*, pp. 162–169. AUAI Press, 2004.

Norman Ferns, Prakash Panangaden, and Doina Precup. Metrics for Markov Decision Processes with infinite state spaces. In *Proceedings of the 21st Conference on Uncertainty in Artificial Intelligence, UAI 2005*, 2005.

Carles Gelada, Saurabh Kumar, Jacob Buckman, Ofir Nachum, and Marc G Bellemare. Deepmdp: Learning continuous latent space models for representation learning. In *International Conference on Machine Learning*, pp. 2170–2179. PMLR, 2019.

Dexter Kozen. Coinductive proof principles for stochastic processes. In *21st Annual IEEE Symposium on Logic in Computer Science (LICS'06)*, pp. 359–366. IEEE, 2006.

Charline Le Lan, Marc G Bellemare, and Pablo Samuel Castro. Metrics and continuity in reinforcement learning. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 35, pp. 8261–8269, 2021.

Bilal Piot, Matthieu Geist, and Olivier Pietquin. Difference of convex functions programming for reinforcement learning. In Z. Ghahramani, M. Welling, C. Cortes, N. D. Lawrence, and K. Q. Weinberger (eds.), *Advances in Neural Information Processing Systems 27*, pp. 2519–2527. Curran Associates, Inc., 2014.

Martin L Puterman. Markov decision processes: Discrete stochastic dynamic programming. 1994.

Cédric Villani. *Optimal transport: old and new*, volume 338. Springer Science & Business Media, 2008.

Amy Zhang, Rowan McAllister, Roberto Calandra, Yarin Gal, and Sergey Levine. Learning invariant representations for reinforcement learning without reconstruction. *arXiv preprint arXiv:2006.10742*, 2020.

A   PROOFS FOR SECTION 3

Lemma 1. *The PMiCo operator T_P^π is a contraction mapping on $\mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ with respect to the L^∞ norm.*

Proof. Let $U, U' \in \mathbb{R}^{\mathcal{X} \times \mathcal{X}}$. Then, by definition of the PMiCo pseudo-metric, it follows that for any $x, y \in \mathcal{X}$,

$$|(T_P^\pi U)(x,y) - (T_P^\pi U')(x,y)| = \left| \gamma \mathbf{1}_{[x \neq y]} \mathbb{E}_{x' \sim P_x^\pi, y' \sim P_y^\pi}(U - U')(x', y') \right| \leq \gamma \|U - U'\|_\infty .$$

□

Theorem 1. *The PMiCo operator T_P^π is a contraction mapping on $\mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ w.r.t. L^∞ norm and its fixed point U_P^π is a pseudo-metric.*

Proof. The first part of Theorem 1 follows from Lemma 1.

By application of Banach's fixed-point theorem and by the completeness of $\mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ under the L^∞ norm, it follows that the PMiCo operator has a unique fixed point $U_P^\pi \in \mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ and applying T_P^π repeatedly to an initial function $U \in \mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ converges to U_P^π.

It it easy to see that U_P^π is symmetric, non-negative and has zero self-distance ($\forall x \in \mathcal{X}, U_P^\pi(x,x) = 0$).

To prove the triangle inequality, we adapt the proof of proposition 4.10 from Castro et al. (2021) and rely on a proof by induction.

We define a sequence of iterates $(U_k)_{k \geq 0}$ in $\mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ by $U_0(x,y) = 0$ for all $x, y \in \mathcal{X}$, and $U_{k+1} = T_M^\pi U_k$ for each $k \geq 0$.

The base case of the inductive argument is clear from the choice of U_0.
For the inductive step, assume that for some $k \geq 0$, $U_k(x,y) \leq U_k(x,z) + U_k(z,y)$ for all $x, y, z \in \mathcal{X}$. For $x, y, z \in \mathcal{X}$,
Case I. If $z = x$ or $z = y$, we have $U_{k+1}(x,y) = U_{k+1}(x,z) + U_{k+1}(z,y)$ by zero self-distance argument.
Case II. If $x = y$, we have $U_{k+1}(x,y) = 0 \leq U_{k+1}(x,z) + U_{k+1}(z,y)$ by non-negativity of U_{k+1}.
Case III. If $x \neq y \neq z$, we have

$$
\begin{aligned}
U_{k+1}(x,y) &= |r^\pi(x) - r^\pi(y)| + \gamma \mathbb{E}_{x' \sim P_x^\pi, y' \sim P_y^\pi}[U_k(x', y')] \qquad \text{because } x \neq y \\
&\leq |r^\pi(x) - r^\pi(z)| + |r^\pi(z) - r^\pi(y)| + \gamma \mathbb{E}_{x' \sim P_x^\pi, y' \sim P_y^\pi, z' \sim P_z^\pi}[U_k(x', z') + U_k(z', y')] \\
&= \left(|r^\pi(x) - r^\pi(z)| + \gamma \mathbb{E}_{x' \sim P_x^\pi, z' \sim P_z^\pi}[U_k(x', z')] \right) + \Big(|r^\pi(z) - r^\pi(y)| \\
&\quad + \gamma \mathbb{E}_{z' \sim P_z^\pi, y' \sim P_y^\pi}[U_k(z', y')] \Big) \\
&= U_{k+1}(x,z) + U_{k+1}(z,y) \qquad \text{because } x \neq z, z \neq y
\end{aligned}
$$

as required. Hence, for any $k \geq 0$, U_{k+1} satisfy the triangle inequality. $U_k \to U_P^\pi$ (using lemma 1), so by taking limits on either side of the inequality, we recover that U_P^π itself satisfies the triangle inequality. □

Lemma 2. *For any policy $\pi \in \mathscr{P}(\mathcal{A})^{\mathcal{X}}$ and states $x, y \in \mathcal{X}$, we have $|V^\pi(x) - V^\pi(y)| \leq U_P^\pi(x,y)$.*

Proof. We adapt the proof of proposition 4.8 from Castro et al. (2021) and apply a coinductive argument (Kozen, 2006) to show that if

$$|V^\pi(x) - V^\pi(y)| \leq U(x,y) \text{ for all } x, y \in \mathcal{X} , \tag{2}$$

for some $U \in \mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ symmetric in its two arguments, then we also have

$$|V^\pi(x) - V^\pi(y)| \leq (T_M^\pi U)(x,y) \text{ for all } x, y \in \mathcal{X} .$$

Since the hypothesis holds for the constant function

$$U(x,y) = \begin{cases} 2R_{\max}/(1-\gamma), & \text{for } x \neq y \\ 0, & \text{for } x = y \end{cases} \quad (3)$$

and T_P^π contracts around U_P^π, the conclusion then follows.

Case I. If $x = y$, $V^\pi(x) - V^\pi(y) = 0 = (T_P^\pi U)(x,y)$.

Case II. If $x \neq y$, suppose Equation 2 holds. Then we have

$$V^\pi(x) - V^\pi(y) = r_x^\pi - r_y^\pi \;\; + \gamma \sum_{x' \in \mathcal{X}} P_x^\pi(x')V(x') - \gamma \sum_{y' \in \mathcal{X}} P_y^\pi(y')V(y')$$

$$\leq |r_x^\pi - r_y^\pi| \;\; + \gamma \sum_{x',y' \in \mathcal{X}} P_x^\pi(x')P_y^\pi(y')(V^\pi(x') - V^\pi(y'))$$

$$\leq |r_x^\pi - r_y^\pi| \;\; + \gamma \sum_{x',y' \in \mathcal{X}} P_x^\pi(x')P_y^\pi(y')U(x',y') \text{ by the inductive hypothesis}$$

$$= |r_x^\pi - r_y^\pi| \;\; + \mathbf{1}_{[x \neq y]}\mathbb{E}_{x' \sim P_x^\pi, y' \sim P_y^\pi} U(x',y') \qquad \text{because } x \neq y$$

$$= (T_P^\pi U)(x,y).$$

By symmetry, $V^\pi(y) - V^\pi(x) \leq (T_P^\pi U)(x,y)$, as required. $\qquad \square$

Theorem 2. *For any policy $\pi \in \Pi$ and states $x,y \in \mathcal{X}$, $|V^\pi(x) - V^\pi(y)| \leq U_P^\pi(x,y) \leq U^\pi(x,y)$.*

Proof. The left hand side of the inequality follows by application of lemma 2.

To prove the right hand side, we use a proof by induction. To do so, we define a sequence of iterates $(U_k)_{k \geq 0}$ and $(O_k)_{k \geq 0}$ in $\mathbb{R}^{\mathcal{X} \times \mathcal{X}}$ by $U_0(x,y) = O_0(x,y) = 0$ for all $x,y \in \mathcal{X}$, and $U_{k+1} = T_P^\pi U_k$ and $O_{k+1} = T_M^\pi O_k$ for each $k \geq 0$.

For $k = 0$, $U_0(x,y) - O_0(x,y) \leq 0$.

Assume that there exists $k > 0$ such that $U_k(x,y) \leq O_k(x,y)$. Then for any $x,y \in \mathcal{X}$

$$U_{k+1}(x,y) - O_{k+1}(x,y) = \gamma \mathbb{E}_{x' \sim P_x^\pi, y' \sim P_y^\pi}[\mathbf{1}_{[x \neq y]}[U_k(x',y') - O_k(x',y')]$$

$$\leq \mathbb{E}_{x' \sim P_x^\pi, y' \sim P_y^\pi}[U_k(x',y') - O_k(x',y')] \qquad \text{because } \mathbf{1}_{[x \neq y]} \leq 1$$

$$\leq 0 \qquad \text{because } U_k(x',y') - O_k(x,y) \leq 0$$

T_P^π (resp. T_M^π) are contraction mappings with respect to the L^∞ norm by Lemma 1 (resp. by corollary 4.3 in Castro et al. (2021)), so as $k \to \infty$, $U_k \to U_P^\pi$ and $O_k \to U^\pi$. By taking limit on the left hand side of the above inequality, we recover that $U_P^\pi(x,y) \leq U^\pi(x,y)$. $\qquad \square$