SHARP CONVERGENCE ANALYSIS OF GRADIENT DESCENT FOR OVERPARAMETERIZED DEEP LINEAR NEURAL NETWORKS

ABSTRACT

This paper provides sharp rates of convergence of the gradient descent (GD) method for overparameterized deep linear neural networks with different random initializations. This study touches upon one major open theoretical problem in machine learning: why deep neural networks trained with GD methods are efficient in many practical applications. While the solution of this problem is still beyond reach for general nonlinear deep neural networks, there have been extensive efforts in the literature in studying relevant questions for deep linear neural networks and there are many interesting results in this research direction. For example, recent results on the loss landscape show that even though the loss function of deep linear neural networks is non-convex, every local minimizer is also a global minimizer. When the GD method is applied to train the deep linear networks, it has been shown in the literature that the convergence behavior of the GD method depends on the initialization. In this paper, we obtain the sharp rate of convergence of GD for deep linear networks, and we show that this rate does not depend on the types of random initialization. Furthermore, we show that the depth of the network does not affect the optimal rate of convergence, provided that the width of each hidden layer is appropriately large. Last but not least, we explain that why GD for the overparameterized deep linear network automatically avoid bad saddles.

1 INTRODUCTION

Deep linear neural networks, as a class of toy models, are frequently used to understand loss surfaces and the gradient-based optimization methods related to non-convex problems. The work by Dauphin et al. (2014), and Choromanska et al. (2015a) explored the loss function of deep nonlinear networks based on random matrix theory (such as spherical spin-glass model). This essentially converted the loss surface of deep nonlinear neural networks into that of deep linear neural networks under certain assumptions, some of which are unrealistic. Then, Choromanska et al. (2015b) suggested an open problem to establish a connection between the loss function of the neural networks and the Hamiltonian of the spherical spin-glass models under milder assumptions. Later, Kawaguchi (2016) successfully discarded most of these assumptions via analyzing the loss surface of the deep linear neural networks.

The landscape for deep linear neural network (Kawaguchi, 2016; Kawaguchi & Lu, 2017; Laurent & Brecht, 2018) focused on several properties of critical points: (i) every local minimum is a global minimum; (ii) every critical point that is not a local minimum is a saddle point; and (iii) there exists a saddle that all eigenvalues of its Hessian are zeros if the network is deeper than three layers. Thus, the convergence to the global minimum for deep linear neural networks suffers from the existence of poor saddles.

Lee et al. (2016) shows that the gradient method almost surely never converges to a strict saddle point, but the time cost can depend exponentially in dimension (Du et al., 2017). The GD with perturbations (Ge et al., 2015; Jin et al., 2017) can find a local minimizer in polynomial time. Thus, the trajectory approach combined with random initialization or random algorithm circumvents the obstacle of existence of poor saddles.

Studies on continuous time dynamics of gradient flow (Du et al., 2018; Arora et al., 2018b) for the deep linear network preserved balance property if the initialization is balanced. Several papers by Arora et al. (2018a;b), Du & Hu (2019), and Hu et al. (2020) successfully proved that the GD with corresponding initialization schemes converges to a global minimizer of the deep linear neural networks with high probability. Furthermore, the rate of convergence is linear, and it behaves like GD for a convex problem.

Hu et al. (2020) established that the convergence for Gaussian initialization can be very slow for large depth deep linear neural networks, unless the width is almost linear in depth. They also show that the orthogonal initialization in deep linear neural networks will speed up convergence. Thus, the convergence behavior of the GD method to train deep linear neural networks depends crucially on the initialization.

Recent discoveries found the connection between deep learning and kernel methods (Daniely, 2017; Arora et al., 2019a;b; Chizat et al., 2019; Lee et al., 2019; Du et al., 2019; Cao & Gu, 2019; Woodworth et al., 2020), especially the neural tangent kernel (NTK), introduced by Jacot et al. (2018). For most common neural networks, the NTK becomes constant (Jacot et al., 2018; Liu et al., 2020), and the NTK remains constant throughout training in the limit of large layer width. The neural networks are well described throughout training by their first order Taylor expansion around their parameters at initialization (Lee et al., 2019).

In this paper, we first study the convergence region, the set of initialization which lead to the linear convergence of the GD for the deep linear neural networks (see Lemma 4.1 or Lemma D.1). Next, we demonstrate that if the minimum width among all hidden layers is sufficiently large, then the random initialization will fall into the convergence region with high probability (see Theorem 3.1, Theorem B.1, Theorem B.2, and Theorem B.3). Furthermore, the worst-case convergence rate of the GD for the deep linear neural networks is almost the same as the original convex problem with a corresponding learning rate. We also demonstrate that the trajectories of the GD for the deep linear neural networks are arbitrarily close to those for the convex problem. The precise statement is related to remark 3, Theorem 3.2, Lemma 4.4 (also see Lemma D.5) and Corollary 1.

This work is inspired by recent work (Du & Hu, 2019; Hu et al., 2020), which carefully constructed the upper and lower bounds of eigenvalues of this Gram matrix along the GD and established linear convergence. In this paper, we generalize their results to strongly convex loss functions with layer-varying width and obtain sharper results. We also show that, our rate of convergence for GD in deep linear neural networks is sharp in the sense that it matches the worst-case convergence rate for the original convex problem. The trajectories between the GD for deep linear neural networks and the original convex problem (1) can be arbitrary close. Furthermore we show that the optimal rate neither depends on the types of random initialization nor the depth of the network, provided that the width of each hidden layer is appropriately large. Last but not least, we explain that why GD for the overparameterized deep linear network automatically avoid bad saddles.

2 PRELIMINARIES

2.1 PROBLEM SETUP

Let $x \in \mathbb{R}^{n_x}$ and $y \in \mathbb{R}^{n_y}$ be an input vector and a target vector, respectively. Define $\{(x_i, y_i)\}_{i=1}^m$ as a training dataset of size m, and let $X = [x_1, x_2, \cdots, x_m] \neq 0$ and $Y = [y_1, y_2, \cdots, y_m]$. Denote the weight parameters by $W \in \mathbb{R}^{n_y \times n_x}$.

Consider the well studied convex optimization problem:

$$\underset{W}{\text{minimize}} \quad L(W) := \frac{1}{m} \sum_{i=1}^m l(Wx_i, y_i). \tag{1}$$

The GD for convex problem (1) with learning rate η_* is given by:

$$W(t+1) = W(t) - \eta_* \nabla L(W(t)), t = 0, 1, 2, \cdots. \tag{2}$$

For any matrix A, let $\sigma_{max}(A)$ and $\sigma_{min}(A)$ be the largest and smallest singular values of A respectively. In this paper, we consider two types of matrix norms and one type of semi-norm for A,

$\|A\| := \sigma_{max}(A), \|A\|_F^2 := tr(AA^T)$, and $\|A\|_X := \|AP_X\|_F$, where $P_X = X(X^TX)^\dagger X^T$ is the orthogonal projection matrix onto the column space of X, and $(X^TX)^\dagger$ is the Moore–Penrose inverse.

For two real matrices A, B with the same size, we consider their Frobenius inner product as well as their semi-inner product, $\langle A, B \rangle = \langle A, B \rangle_F := tr(A^TB), \langle A, B \rangle_X := \langle AP_X, BP_X \rangle$.

We list some basic properties for the semi-norm and semi-inner product.

Lemma 2.1. *The loss function $L(W)$ defined in (1) satisfies the following properties: for any $W, V \in \mathbb{R}^{n_y \times n_x}$,*

$1. L(W) = L(WP_X), 2. \nabla L(W) = \nabla L(WP_X)P_X, 3. \langle \nabla L(W), V \rangle_F = \langle \nabla L(W), V \rangle_X,$

$4. \|\nabla L(W)\|_F = \|\nabla L(W)\|_X, 5. \|W\|_X \equiv \|W\|_F$ *if and only if X is full row rank.*

The next lemma illustrates why semi-norm $\|\cdot\|_X$ is crucial in our analysis.

Lemma 2.2. *Assume that $l(\cdot, y)$ is $\alpha(l)-$strongly convex. The following statements hold.*

1. *If X is not full row rank, then $L(W)$ is neither strictly convex nor strongly convex with respect to $\|\cdot\|_F$.*

2. *$L(W)$ is $\frac{\alpha(l)\lambda_{min}(XX^T)}{m}-$strongly convex with respect to $\|\cdot\|_X$, where $\lambda_{min}(XX^T)$ is the smallest non-zero eigenvalue of XX^T.*

The proofs of the two lemmas above can be found in appendix A. From now on, if we do not specify the inner product, we will always consider the semi-inner product.

Assume that L is $\alpha-$strongly convex ($\alpha > 0$), and ∇L is $\beta-$Lipschitz (with respect to semi-norm $\|\cdot\|_X$), that means, for any $W, V \in \mathbb{R}^{n_y \times n_x}$,

$$L(W) \geq L(V) + \langle \nabla L(V), W - V \rangle_X + \frac{\alpha}{2} \|W - V\|_X^2,$$

$$\|\nabla L(W) - \nabla L(V)\|_X = \|\nabla L(W) - \nabla L(V)\|_F \leq \beta \|W - V\|_X.$$

Without loss of generality, we assume that α and β are the best constants. Then, Lemma 2.2 implies $\alpha \geq \frac{\alpha(l)\lambda_{min}(XX^T)}{m}$. Similarly, we can also show that $\beta \leq \frac{\beta(l)\lambda_{max}(XX^T)}{m}$, where $\nabla l(\cdot, y)$ is $\beta(l)-$Lipschitz and $\lambda_{max}(XX^T)$ is the largest eigenvalue of XX^T.

Define the effective condition number of the convex function L by $\kappa = \kappa(L) = \frac{\beta}{\alpha} < \infty$. κ appears naturally in the rate of convergence of GD. Let W_* be a global minimizer of $L(W)$, that is $L(W_*) = \min_W L(W)$. Notice that W_* might not be unique, but W_*P_X is unique.

The well known results on rate of convergence of GD (2) state

$$\eta_* = \frac{1}{\beta} \implies \mathcal{E}(t) \leq \left(1 - \frac{1}{\kappa}\right)^t \mathcal{E}(0), t = 1, 2, \cdots, \text{ as well as,} \tag{3}$$

$$\eta_* = \frac{2}{\alpha + \beta} \implies \mathcal{E}(t) \leq \frac{\beta}{2} \left(1 - \frac{4\kappa}{(1 + \kappa)^2}\right)^t \|W(0) - W_*\|_X^2, t = 1, 2, \cdots, \tag{4}$$

where $\mathcal{E}(t) = L(W(t)) - L(W_*)$.

2.2 DEEP LINEAR NETWORK SETUP

Let $N - 1$ be the number of hidden layers. Assume $rank(X) = r$. Denote the weight parameters by $W_k \in \mathbb{R}^{n_k \times n_{k-1}}, k = 1, 2 \cdots, N$, with $n_N = n_y, n_0 = n_x$, where the n_k is the width of the k-th layer. Set $n_{min} = \min\{n_1, n_2, \cdots, n_{N-1}\}$, and $n_{max} = \max\{n_1, n_2, \cdots, n_{N-1}\}$. For notational convenience, we denote $n_{j:i} = \prod_{i \leq k \leq j} n_k$ and denote $W_{j:i} = W_jW_{j-1} \cdots W_i$ for each $1 \leq i \leq j \leq N$. Define $n_{i-1:i} = 1$ and $W_{i-1:i} = I$ (of appropriate dimension) for completeness.

Consider the implicit regularization $W = W_{N:1}$ for the convex problem (1). Thus, we obtain the following non-convex optimization problem of deep linear neural networks:

$$\underset{W_1, \cdots, W_N}{\text{minimize}} \quad L(W_{N:1}) = \frac{1}{m} \sum_{i=1}^{m} l(W_{N:1}x_i, y_i). \tag{5}$$

Example 2.1. Specifically, if we set the loss to be $l(Wx_i, y_i) = \|Wx_i - y_i\|_2^2$, then $L(W) = \frac{1}{m}\|WX - Y\|_F^2$ is $\frac{2\lambda_{min}(XX^T)}{m}$−strongly convex, and ∇L is $\frac{2\lambda_{max}(XX^T)}{m}$−Lipschitz.

Example 2.2. The deep linear neural networks with regularization $\lambda\|W_N \cdots W_1 P_X\|_F^2$ can be converted into a new optimization problem

$$\underset{W_1, \cdots, W_N}{\text{minimize}} \quad L(W_{N:1}) + \lambda\|W_{N:1}\|_X^2.$$

Let $L_\lambda(W) = L(W) + \lambda\|W\|_X^2$. Then $L_\lambda(\cdot)$ is $\alpha + 2\lambda$-strongly convex, and $\nabla L_\lambda(\cdot)$ is $\beta + 2\lambda$-Lipschitz.

More general, if we consider regularization with form $R(W) = \lambda \cdot g(WP_X)$, and $g(\cdot)$ is α'-strongly convex, and β'-Lipschitz, then for the optimization problem

$$\underset{W_1, \cdots, W_N}{\text{minimize}} \quad L(W_{N:1}) + R(W_N \cdots W_1) =: L_R(W_N \cdots W_1),$$

we know that $L_R(\cdot)$ is $\alpha + \lambda\alpha'$-strongly convex, and $\nabla L_R(\cdot)$ is $\beta + \lambda\beta'$-Lipschitz.

2.3 INITIALIZATION SCHEMES

In literature, researchers consider the following form of the deep linear network instead of (5):

$$\underset{W_1, \cdots, W_N}{\text{minimize}} \quad L(a_N W_{N:1}) = \frac{1}{m} \sum_{i=1}^m l(a_N W_{N:1} x_i, y_i), \tag{6}$$

where $a_N = 1/\sqrt{n_1 n_2 \cdots n_N}$ is a normalization constant.

Applying GD on (6), where we update W_j simultaneously for j,

$$W_j(t+1) = W_j(t) - \eta \cdot a_N \left(W_{N:j+1}(t)\right)^T \nabla L\left(a_N W_{N:1}(t)\right)\left(W_{j-1:1}(t)\right)^T, j = 1, \cdots, N. \tag{7}$$

Recent work consider GD (7), initializing $W_j(0)$ with Gaussian initialization (Du & Hu, 2019) or scaled orthogonal initialization (Hu et al., 2020).

In this paper, we consider the following three kinds of random initialization which generalizes their idea.

Gaussian initialization: Let $W_1(0), \cdots, W_N(0)$ be the weight matrices at initialization. We assume that all entries of $W_j, 1 \leq j \leq N$ are independent Gaussian random variables with zero mean and unit variance.

Then a_N is a normalization constant in the sense that for any $x \in \mathbb{R}^{n_0}$, we have

$$\mathbb{E}\left[\|a_N W_{N:1}(0)x\|_2^2\right] = \|x\|_2^2. \tag{8}$$

In fact, all initialization we discussed in this paper satisfy (8).

Remark 1. Let $V_i(t) = \frac{1}{\sqrt{n_i}} W_i(t)$, for $1 \leq i \leq N$. Then, the GD (7) with unit variance Gaussian initialization is equivalent to

$$V_j(t+1) = V_j(t) - \frac{\eta}{n_j}\left(V_{N:j+1}(t)\right)^T \nabla L\left(V_{N:1}(t)\right)\left(V_{j-1:1}(t)\right)^T, \tag{9}$$

with mean zero and variance $\frac{1}{n_i}$ Gaussian initialization for $V_i, i = 1, \cdots, N$.

The GD (9) for the loss (5) is equivalent to the GD (7) for the loss (6). From now on, we will only consider GD (7) for deep linear neural networks (6).

Orthogonal initialization: We consider a so-called one peak random orthogonal projections and embeddings initialization, which generalizes the idea of orthogonal initialization (Hu et al., 2020).

Definition 2.1. A initialization $W_{N:1}(0) = W_N(0)W_{N-1}(0) \cdots W_1(0)$ is said to be one peak random orthogonal projections and embeddings initialization if there exists $1 \leq p < N$, such that $n_0 \leq n_1 \leq n_2 \leq \cdots \leq n_p, n_p \geq n_{p+1} \geq n_{p+2} \geq \cdots n_{N-1} \geq n_N$, and

$W_1(0), W_2(0), \cdots, W_p(0), W_{p+1}(0), W_{p+2}(0), \cdots, W_{n_N}(0)$ are independent and uniformly distributed over rectangular matrices which satisfy

$$\begin{cases} W_i^T(0)W_i(0) = n_i I_{n_{i-1}}, 1 \leq i \leq p, \\ W_j(0)W_j^T(0) = n_{j-1} I_{n_j}, p+1 \leq j \leq N. \end{cases}$$

Remark 2. In this definition, $\sqrt{\frac{1}{n_i}} W_i(0), 1 \leq i \leq p$ are random embeddings and $\sqrt{\frac{1}{n_{j-1}}} W_j(0), p+1 \leq j \leq N$ are random orthogonal projections. Notice that, A is a random orthogonal projection if and only if A^T is a random embedding.

Arora et al. (2018a) studied the rate of convergence to global optimum for GD to train deep linear neural network for balanced initialization. Here we will consider a special case of balanced initialization which is described as follows.

Special balanced initialization: Assume $n_1 = \cdots = n_{N-1} = n$. Consider initialization $W_N(0) = \sqrt{n} U_N [I_{n_y}, 0_{n_y \times (n-n_y)}] V_N^T$, $W_1(0) = \sqrt{n} U_1 [I_{n_x}, 0_{n_x \times (n-n_x)}]^T V_1^T$ and $W_i(0) = \sqrt{n} U_i I_n V_i^T, 2 \leq i \leq N-1$, where $U_{N-1}, U_N, V_1, V_i = U_{i-1}, 2 \leq i \leq N-1$ are orthogonal matrices (random or deterministic), and V_N has uniform distribution over orthogonal matrices. Notice that only V_N is required to be random.

A simple estimation of the loss at the initialization is given by the following lemma.

Lemma 2.3. *If the initialization satisfies (8) for all x, then with probability at least $1 - \frac{\delta}{2}$, we have*

$$L(a_N W_{N:1}(0)) - L(W_*) \leq \beta B_\delta, \text{ where } B_\delta = \left(\frac{2 \cdot rank(X)}{\delta} + \|W_*\|_X^2 \right).$$

Note that using sharp concentration inequality, the bound B_δ can be improved.

3 MAIN THEOREMS

Assume the thinnest layer is either the input layer or the output layer, that is $n_{min} \geq \max\{n_0, n_N\}$ and the ratio between the width of any hidden layer is bounded from above, precisely we have $\frac{n_{max}}{n_{min}} \leq C_0 < \infty$. The quantities C_2, C_5 and C_6 are defined in appendix D, which depend on hyperparameters $n_N, \kappa, \delta, rank(X), C_0$ and N.

For notational convenience, we denote

$$\mathcal{E}(t) = L(W(t)) - L(W_*), \text{ and } \mathcal{E}_{DLN}(t) = L(a_N W_{N:1}(t)) - L(W_*).$$

Our assumptions and notation are now in place. We next state our main theorems in this section.

3.1 LINEAR CONVERGENCE OF DEEP LINEAR NEURAL NETWORKS

In appendix B, we present a sharp estimate on the linear convergence of the GD for deep linear neural networks, in Theorem B.1 for Gaussian initialization, Theorem B.2 for orthogonal initialization, and Theorem B.3 for special balanced initialization, respectively. In particular, with the special choice of learning rate $\eta = \frac{n_N}{\beta N}$, Theorem B.1 and Theorem B.2 lead us to obtain the following optimum rate of convergence:

Theorem 3.1. *Given any $\delta, \varepsilon \in (0, \frac{1}{2})$, there exists a constant $C := C(\varepsilon)$, such that if one of the following two overparameterization condition holds:*

1. $n_{min} \geq C \cdot C_2 \cdot N$ *with the Gaussian initialization,*

2. $n_{min} \geq C \cdot C_5$ *with the one peak random orthogonal projections and embeddings initialization,*

with probability at least $1 - \delta$ we have

$$\mathcal{E}_{DLN}(t) \leq \left(1 - \frac{1-\varepsilon}{\kappa} \right)^t \mathcal{E}_{DLN}(0), t = 1, 2, \cdots.$$

Remark 3. Consider the GD (2) with learning rate $\eta_* = \frac{1}{\beta}$, and initialization $W(0) = a_N W_{N:1}(0)$. The well known result on the rate of convergence (3) for the GD (2) of convex problem (1) matches the rates in Theorem B.1 and Theorem B.2.

Remark 4. Du & Hu (2019), and Hu et al. (2020) already showed that the number of iterations needed to reach precision ε is $O\left(\kappa \log \frac{1}{\epsilon}\right)$ for l_2 loss. We only improved the rate of convergence and generalized their results to any strongly convex loss.

3.2 RESULTS ON TRAJECTORIES

Theorem 3.1 and remark 3 establish that the rate of convergence to a global optimum for GD to train a deep linear neural networks is almost the same as the trajectories for the GD to train the corresponding convex problem with high probability, if the width is large enough. Moreover, the GD for the fully-connected deep linear neural networks (7) and that for the GD (2) have almost the same trajectories.

Let $\eta_1 = \frac{2n_N}{\beta N}$ be an upper bound of learning rate η. We can show that the trajectories of GD (7) for deep linear neural networks (6) with learning rate $\eta < \eta_1$ are close to those of GD (2) with learning rate $\eta_* = \frac{N}{n_N}\eta$ for the corresponding convex problem (1) with high probability, if the width of each hidden layer is sufficiently large. The precise statement is as follows.

Theorem 3.2. *Consider the GD for the deep linear neural networks (7) with learning rate $\eta < \eta_1$ for $a_N W_{N:1}(t)$, $t = 0, 1, \cdots$, and the GD (2) with learning rate $\eta_* = \frac{N}{n_N}\eta$ for $W(t)$, $t = 0, 1, \cdots$. Given $\tau, \delta \in (0, 1)$, there exists a constant $C := C(\tau, \eta/\eta_1)$ such that if one of the following three overparameterization conditions holds:*

1. $n_{min} \geq C \cdot C_2 \cdot N$ *with the Gaussian initialization,*

2. $n_{min} \geq C \cdot C_5$ *with the one peak random orthogonal projections and embeddings initialization,*

3. $n_{min} \geq C \cdot C_6$ *with the special balanced initialization,*

then with probability at least $1 - \delta$, we have

$$\|a_N W_{N:1}(t) - W(t)\|_X^2 \leq D(\tau, q, t) \|a_N W_{N:1}(0) - W_*\|_X^2, \tag{10a}$$

$$|\mathcal{E}_{DLN}(t) - \mathcal{E}(t)| \leq \beta \left(q^{t/2} \sqrt{D(\tau, q, t)} + \frac{1}{2} D(\tau, q, t) \right) \|a_N W_{N:1}(0) - W_*\|_X^2, \tag{10b}$$

$$\mathcal{E}_{DLN}(t) \leq 3\beta(q + \tau)^t \|a_N W_{N:1}(0) - W_*\|_X^2, \tag{10c}$$

where $D(\tau, q, t) = \min\left\{\frac{\tau}{1-q}, 2(q + \tau)^t\right\}$, with $0 < q < 1$ defined in (15).

Remark 5. To our knowledge, we are the first who showed that the trajectory of the overparameterized deep linear neural networks is close to the original convex problem with appropriately rescaled learning rate.

Corollary 1. *Under the setting of Theorem 3.2, if we set $\eta = \frac{2n_N}{(\alpha+\beta)N}$, the following inequality hold with high probability,*

$$\mathcal{E}_{DLN}(t) \leq 3\beta \left(1 - \frac{4\kappa}{(1 + \kappa)^2} + \tau\right)^t \|a_N W_{N:1}(0) - W_*\|_X^2. \tag{11}$$

We notice that the rate of convergence in (11) is better than that in Theorem 3.1. It is because if $\kappa > 1$, we can choose τ sufficiently small, so that the following inequality holds:

$$1 - \frac{4\kappa}{(1 + \kappa)^2} + \tau < 1 - \frac{1}{\kappa}.$$

Roughly speaking, Theorem 3.1, Theorem 3.2, Theorem B.1, together with Theorem B.2 illustrate that the implicit regularization induced by GD for a convex problem recovers the convex problem itself in terms of optimization, at the cost of linear convergence only with high probability for random initialization.

Remark 6. Recall the constants C_2, C_5, C_6 defined in appendix B. The term $\frac{rank(X)}{\delta}$ is not optimal, since our concentration inequality only depends on the second moment. By using stronger concentration inequalities for our Lemma 2.3, similar to the proof of proposition 6.5 (Du & Hu, 2019) and Lemma 4.2 (Hu et al., 2020), the $\frac{rank(X)}{\delta}$ can be improved to $1 + \log(\frac{rank(X)}{\delta})$. The C_2 is proportional to κ^2, which is slightly better than the constant in Du & Hu (2019) that is proportional to κ^3. The C_5 is also slightly better than the constant in Hu et al. (2020), since we do not have the extra term $\frac{\|X\|_F^2}{\|X\|^2}$. The improvement of the constant is mainly due to introducing the semi-norm $\|\cdot\|_X$.

4 INSIGHTS FOR THEOREM 3.2

Initialization and convergence region: Arora et al. (2018a) showed that if the initialization is approximately balanced, and the product matrix $W_{N:1}(0)$ is very close to a global minimizer, then GD linearly converges to the global minimum for the deep linear network, without any requirement on the width. However, the convergence region in (Arora et al., 2018a) is very small, since $W_{N:1}(0)$ has to be very close to W_*. Later, several papers by Du & Hu (2019), and Hu et al. (2020) successfully proved that the GD with Gaussian, or orthogonal initialization linearly converges to a global minimizer of the overparameterized deep linear neural networks with high probability. They introduced a technique to analyze trajectories of GD with large widths for any deterministic initialization.

We introduce the following lemma which describes the linear convergence result for the deep linear network with deterministic initialization.

Lemma 4.1. *Under the setting of Lemma D.1, the GD for the deep linear network satisfies*
$$\mathcal{E}_{DLN}(t) \le (1 - \eta\gamma)^t \mathcal{E}_{DLN}(0), t = 1, 2, \cdots.$$

Our convergence region (see (31) in Lemma D.1 and Definition D.1) originated from Du & Hu (2019), and Hu et al. (2020). This convergence region can be view as a neighbourhood of special balanced initialization, if $n_1 = n_2 = \cdots = n_{N-1}$. Both Gaussian and orthogonal initialization are approximately balanced.

For l_2 loss, without loss of generality assume X to be full rank and $L(W_*) = 0$, due to the decomposition method in claim B.1 from Du & Hu (2019). However, when considering general strongly convex loss, we have to directly confront low rank X in our analysis. Thus, the $\|\cdot\|_X$ appears naturally and helps us to achieve the sharp rate of convergence in our main theorems. In addition to the technique in Du & Hu (2019), and Hu et al. (2020), we also use classical convex optimization techniques (such as inequalities in Lemma C.1, and Polyak-Łojasiewicz inequality in (26)), as well as the classical concentration inequalities for beta distribution (such as the Chernoff type bound in Lemma F.3).

Why trajectories of GD for overparameterized deep linear neural networks with approximate balanced initialization are close to trajectories for convex problems? Even though the recent result (Ziyin et al., 2022) can describe the exact global minimizer for the deep linear network (with regularization term such as l_2), the evolution of each W_j is still hard to track. Instead, we consider the discrete dynamics for product matrices (see (41) and (42)):
$$a_N W_{N:1}(t+1) = a_N W_{N:1}(t) - \eta \cdot P(t)[\nabla L(a_N W_{N:1}(t) P_X)] + a_N E(t).$$
Du & Hu (2019) showed for their own linear operator P_t that $\lambda_{max}(P_t) \le O(\frac{N}{n_N}) \cdot \lambda_{max}(X^T X)$ and $\lambda_{min}(P_t) \ge \Omega(\frac{N}{n_N}) \cdot \lambda_{min}(X^T X)$. To our knowledge, we are the first who proved that for our operator $P(t)[\cdot] \approx \frac{N}{n_N} I$ (also see (44)), where I is the identity operator. $E(t)$ is negligible, which leads to the following result on discrete dynamics (see Lemma D.3).

Lemma 4.2. *Under the setting of Lemma D.3, we have* $a_N W_{N:1}(t + 1) = a_N W_{N:1}(t) - \frac{N}{n_N}\eta\nabla L(a_N W_{N:1}(t)) + R(t)$, *with* $\|R(t)\|_X \le \tau \|a_N W_{N:1}(t) - W_*\|_X$.

Without the $R(t)$ term, the discrete dynamics is exactly the GD for a convex function. To control the distance between the two trajectories, we introduce the following lemma (also see Lemma D.4).

Lemma 4.3. *Assume* $\tau \in [0, 1)$. *Consider a discrete dynamical system $V(t)$ such that,*
$$V(t+1) = V(t) - \eta_*\nabla L(V(t)) + R(t), \text{ where } \|R(t)\|_X \le \tau \|V(t) - W_*\|_X.$$
If $\eta_* \le 2/\beta$, we have $\|V(t) - W_*\|_X^2 \le (q + 7\tau)^t \|V(0) - W_*\|_X^2$, where q is defined in (15).

With the help of this lemma, we further obtain the following trajectories comparison lemma (also see Lemma D.5), which leads to the main conclusions in Theorem 3.2.

Lemma 4.4. *Under the setting of Lemma D.5, we have*

$$\|a_N W_{N:1}(t) - W(t)\|_X^2 \leq D(\tau, q, t) \|a_N W_{N:1}(0) - W_*\|_X^2 , \tag{12a}$$

$$|\mathcal{E}_{DLN}(t) - \mathcal{E}(t)| \leq \beta \left(q^{t/2} \sqrt{D(\tau, q, t)} + \frac{1}{2} D(\tau, q, t) \right) \|a_N W_{N:1}(0) - W_*\|_X^2 , \tag{12b}$$

$$\mathcal{E}_{DLN}(t) \leq 3\beta(q + \tau)^t \|a_N W_{N:1}(0) - W_*\|_X^2 , \tag{12c}$$

where $D(\tau, q, t) = \min \left\{ \frac{\tau}{1-q}, 2(q+\tau)^t \right\}$, with $0 < q < 1$ defined in (15).

Why do bad saddles not affect GD for overparameterized deep linear neural networks? A critical point x^* of f is a bad saddle if $\lambda_{min}(\nabla^2 f(x^*)) = 0$. Kawaguchi (2016) showed that the deep linear network has bad saddles, thus in general vanishing Hessian can hinder optimization.

Theorem 2.3 in Kawaguchi (2016) showed that all bad saddles satisfy that $W_{N-1:2}$ is a non-full rank matrix. Thus, to show that trajectories of GD are away from bad saddle points, it suffices to show that $\inf_t \sigma_{min}(W_{N-1:2}(t)) > 0$. In literature, there are two main ways to avoid bad saddles for GD to train the deep linear network.

On the one hand, under the setting in Arora et al. (2018b), it has been showed that if the approximate balanced initialization satisfies $\|W_{N:1}(0) - W_*\|_F \leq \sigma_{min}(W_*) - c$, for some $0 < c < \sigma_{min}(W_*)$, then $\sigma_{min}(W_{N:1}(t)) \geq c$ through the training as well as $\|W_1(t)\| \leq (4 \|W_*\|_F)^{1/N}$, and $\|W_N(t)\| \leq (4 \|W_*\|_F)^{1/N}$. Thus,

$$\sigma_{min}(W_{N-1:2}(t)) \geq \frac{\sigma_{min}(W_{N:1}(t))}{\|W_1(t)\| \|W_N(t)\|} \geq \frac{c}{(4 \|W_*\|)^{2/N}}.$$

On the other hand, if we assume our rescaled and overparameterized weight initialization falls into the convergence region (31), we can show that (see $\mathcal{B}(t)$ in the proof of Lemma D.1)

$$\sigma_{min}(W_{N-1:2}(t)) \geq \max \left\{ \frac{\sigma_{min}(W_{N:2}(t))}{\sigma_{max}(W_N(t))}, \frac{\sigma_{min}(W_{N-1:1}(t))}{\sigma_{max}(W_1(t))} \right\}.$$

Thus, $\sigma_{min}(\frac{W_{N-1:2}}{(n_{N-1:2})^{1/2}}) \geq e^{-c_1-c_2} \max\{ \frac{n_1}{n_{N-1}}, \frac{n_{N-1}}{n_1} \} \geq e^{-c_1-c_2} > 0$.

In conclusion, we first made a conjecture that for the non-overparameterized deep linear network under the setting in Arora et al. (2018b), there are no bad saddles satisfying $\|W_{N:1}(0) - W_*\|_F < \sigma_{min}(W_*)$. Thus, $\|W_{N:1}(0) - W_*\|_F < \sigma_{min}(W_*)$ is indeed a convergence region. But this region in general is very small, and can even be empty provided $\sigma_{min}(W_*) = 0$. For the overparameterized deep linear network, GD initialized in the convergence region will force the trajectories away from all bad saddles.

Why does the width have to be large? We will discuss overparameterization phenomenons in the deep linear network. For simplicity, we consider special balanced initialization. First of all, we know that $\|W_i(t) - W_i(0)\|_F = O(\frac{1}{N})$ (see $\mathcal{C}(t)$ in the proof of Lemma D.1), provided $\eta = h \frac{n_N}{\beta N}$ and $\gamma = O(\frac{\alpha N}{n_N})$, where $h \in (0, 2)$.

The overparameterized deep linear network around special balanced initialization is full of global minimizers, i.e. the trajectory limit $(W_1^*, W_2^*, \cdots, W_N^*)$ is in the $O(\frac{1}{N})$ neighborhood of the special balanced initialization $(W_1(0), W_2(0), \cdots, W_N(0))$. Notice that

$$\sigma_{min}(W_{N-1:2}(0)) = \prod_{j=2}^{N-1} \sigma_{min}(W_j(0)) = \sigma_{max}(W_{N-1:2}(0)) = \prod_{j=2}^{N-1} \sigma_{max}(W_j(0)) = n^{(N-2)/2},$$

as well as for any $(W_1, W_2, \cdots, W_N)$ around the $O(\frac{1}{N})$ neighborhood of any given initialization $(W_1(0), W_2(0), \cdots, W_N(0))$, we have (detailed argument can be found in the proof of $\mathcal{B}(t)$ in Lemma D.1):

$$\|W_{N-1:2} - W_{N-1:2}(0)\|_F \leq \sum_{s=1}^{N-2} \binom{N-1}{s} O(\frac{1}{N})^s (n^{(N-2-s)/2}) \leq n^{(N-2)/2} O\left(\frac{1}{\sqrt{n}} \right).$$

Thus in terms of landscape, we have

$$\frac{\sigma_{min}(W_{N-1:2})}{\sigma_{min}(W_{N-1:2}(0))} \geq \frac{\sigma_{min}(W_{N-1:2}(0)) - \|W_{N-1:2} - W_{N-1:2}(0)\|_F}{\sigma_{min}(W_{N-1:2}(0))} \geq 1 - O(\frac{1}{\sqrt{n}}), \quad (13)$$

which means that there are no bad saddles around the $O(\frac{1}{N})$ neighborhood of special balanced initialization, if the width n is large enough.

In terms of training, we have (see the proof of Lemma D.1),

$$\|W_i(t) - W_i(0)\|_F = O(\frac{1}{N}), \ \|W_i(0)\|_F = n, 2 \leq i \leq N - 1,$$

as well as $\frac{\|W_i(t) - W_i(0)\|_F}{\|W_i(0)\|_F} = O(\frac{1}{Nn})$.

Thus for the overparameterized deep linear network, the GD with approximate balanced initialization only trains W_1 and W_N, and other weight matrices are almost unmoved. We provide empirical evidence to support the argument above in appendix G.

SGD for the Deep Linear network. The convergence result of SGD for deep linear resnets is obtained by Zou et al. (2020), if the initial loss is close to the global minimum $L(W_*)$. For the large width deep linear network, our conjecture is that the trajectory of SGD behaves similar to the trajectory of SGD for a corresponding convex problem.

Numerical Experiments: We will discuss some empirical evidence in appendix H to support the main results in Section 3. Figure 1 and 2 in appendix H show plots of the logarithm of loss as a function of number of iterations. When n is small, the trajectories of loss for deep linear neural networks are not even decreasing in some iterations; when n is large, the trajectories of loss for deep linear neural networks are close to those for the corresponding convex problem.

5 OVERVIEW OF THE PROOFS OF MAIN THEOREMS AND LEMMAS

Now we will give an overview of the proofs for all theorems in the main results. Since Theorem 3.1 in the main results are special cases of general theorems with non-optimal learning rate (see Theorem B.1 and Theorem B.2), we only need to focus on the proofs of the general theorems (see Theorem B.1, Theorem B.2, Theorem B.3, and Theorem 3.2).

We start with the convergence region of deep linear neural networks. Basically, the convergence region is the set of initialization which lead to the convergence of the GD for the deep linear neural networks. The precise definition can be found in the appendix D. Lemma 4.1 and Lemma 4.4 (also see Lemma D.1 and Lemma D.5) prove that this convergence region satisfies the following properties: if the initialization falls into the convergence region, then

(i) the GD is guaranteed to converge to a global minimizer of the deep linear neural networks,

(ii) the worst-case rate of convergence of the GD for the deep linear neural networks, which is a non-convex problem, is almost the same as the corresponding convex problem with a corresponding learning rate, and,

(iii) the trajectories of the GD for the deep linear neural networks are arbitrarily close to those for the corresponding convex problem.

More precisely, Lemma 4.1 (also see Lemma D.1) establishes the convergence region for deterministic initialization, and it demonstrates the first two properties, (i) and (ii). Additionally, in appendix E and appendix F we also proved that the spectral properties of products of random matrices partially reveal the mystery of overparameterization, that is, overparameterization by adding width of each hidden layer guarantees that the random initialization will fall into the convergence region with high probability. These results provide a foundation to establish the main linear convergence theorem for random initialization (see Theorem B.1, B.2, and B.3).

On the other hand, Lemma 4.2 (also see Lemma D.3) shows that if the initialization falls into the convergence region, the update rule for the product of weight matrices in GD for deep linear neural networks is more or less that given in (2). This result can be used to establish both Lemma 4.4 (also see Lemma D.5), and Theorem 3.2, which is precisely the property (iii) of the convergence region for deterministic initialization and non-deterministic initialization, respectively.

REFERENCES

Sanjeev Arora, Nadav Cohen, Noah Golowich, and Wei Hu. A convergence analysis of gradient descent for deep linear neural networks. *CoRR*, abs/1810.02281, 2018a.

Sanjeev Arora, Nadav Cohen, and Elad Hazan. On the optimization of deep networks: Implicit acceleration by overparameterization. In *International Conference on Machine Learning*, pp. 244–253. PMLR, 2018b.

Sanjeev Arora, Simon Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In *International Conference on Machine Learning*, pp. 322–332. PMLR, 2019a.

Sanjeev Arora, Simon S Du, Wei Hu, Zhiyuan Li, Russ R Salakhutdinov, and Ruosong Wang. On exact computation with an infinitely wide neural net. *Advances in Neural Information Processing Systems*, 32, 2019b.

Yuan Cao and Quanquan Gu. Generalization bounds of stochastic gradient descent for wide and deep neural networks. *Advances in neural information processing systems*, 32, 2019.

Djalil Chafaı, Djalil Chafä, Olivier Guédon, Guillaume Lecue, and Alain Pajor. Singular values of random matrices. *Lecture Notes*, 2009.

Lénaïc Chizat, Edouard Oyallon, and Francis R. Bach. On lazy training in differentiable programming. In *NeurIPS*, 2019.

Anna Choromanska, Mikael Henaff, Michael Mathieu, Gérard Ben Arous, and Yann LeCun. The loss surfaces of multilayer networks. In *Artificial intelligence and statistics*, pp. 192–204. PMLR, 2015a.

Anna Choromanska, Yann LeCun, and Gérard Ben Arous. Open problem: The landscape of the loss surfaces of multilayer networks. In *Conference on Learning Theory*, pp. 1756–1760. PMLR, 2015b.

Amit Daniely. Sgd learns the conjugate kernel class of the network. *Advances in Neural Information Processing Systems*, 30, 2017.

Yann N Dauphin, Razvan Pascanu, Caglar Gulcehre, Kyunghyun Cho, Surya Ganguli, and Yoshua Bengio. Identifying and attacking the saddle point problem in high-dimensional non-convex optimization. *Advances in neural information processing systems*, 27, 2014.

Simon Du and Wei Hu. Width provably matters in optimization for deep linear neural networks. In *International Conference on Machine Learning*, pp. 1655–1664. PMLR, 2019.

Simon Du, Jason Lee, Haochuan Li, Liwei Wang, and Xiyu Zhai. Gradient descent finds global minima of deep neural networks. In *International conference on machine learning*, pp. 1675–1685. PMLR, 2019.

Simon S Du, Chi Jin, Jason D Lee, Michael I Jordan, Aarti Singh, and Barnabas Poczos. Gradient descent can take exponential time to escape saddle points. *Advances in neural information processing systems*, 30, 2017.

Simon S Du, Wei Hu, and Jason D Lee. Algorithmic regularization in learning deep homogeneous models: Layers are automatically balanced. *Advances in Neural Information Processing Systems*, 31, 2018.

Morris L. Eaton. Group invariance applications in statistics. *Regional Conference Series in Probability and Statistics*, 1:i–133, 1989. ISSN 19355912.

Sam Elder. Bayesian adaptive data analysis guarantees from subgaussianity. *CoRR*, abs/1611.00065, 2016.

Rong Ge, Furong Huang, Chi Jin, and Yang Yuan. Escaping from saddle points—online stochastic gradient for tensor decomposition. In *Conference on learning theory*, pp. 797–842. PMLR, 2015.

Wei Hu, Lechao Xiao, and Jeffrey Pennington. Provable benefit of orthogonal initialization in optimizing deep linear networks, 2020.

Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. *Advances in neural information processing systems*, 31, 2018.

GJO Jameson. Inequalities for gamma function ratios. *The American Mathematical Monthly*, 120 (10):936–940, 2013.

Chi Jin, Rong Ge, Praneeth Netrapalli, Sham M Kakade, and Michael I Jordan. How to escape saddle points efficiently. In *International Conference on Machine Learning*, pp. 1724–1732. PMLR, 2017.

Kenji Kawaguchi. Deep learning without poor local minima. *In Advances In Neural Information Processing Systems*, pp. 586–594, 2016.

Kenji Kawaguchi and Haihao Lu. Deep creates no bad local minima. *arXiv: 1702.08580*, 2017.

Thomas Laurent and James Brecht. Deep linear networks with arbitrary loss: All local minima are global. *International Conference on Machine Learning*, pp. 2908–2913, 2018.

Jaehoon Lee, Lechao Xiao, Samuel Schoenholz, Yasaman Bahri, Roman Novak, Jascha Sohl-Dickstein, and Jeffrey Pennington. Wide neural networks of any depth evolve as linear models under gradient descent. *Advances in neural information processing systems*, 32, 2019.

Jason D Lee, Max Simchowitz, Michael I Jordan, and Benjamin Recht. Gradient descent only converges to minimizers. In *Conference on learning theory*, pp. 1246–1257. PMLR, 2016.

Chaoyue Liu, Libin Zhu, and Misha Belkin. On the linearity of large non-linear models: when and why the tangent kernel is constant. *Advances in Neural Information Processing Systems*, 33: 15954–15964, 2020.

Olivier Marchal and Julyan Arbel. On the sub-gaussianity of the beta and dirichlet distributions. *Electronic Communications in Probability*, 22(none):1 – 14, 2017.

Feng Qi, Bai-Ni Guo, and C. Chen. The best bounds in gautschi-kershaw inequalities. *Mathematical Inequalities Applications*, pp. 427–436, 2006.

Blake Woodworth, Suriya Gunasekar, Jason D Lee, Edward Moroshko, Pedro Savarese, Itay Golan, Daniel Soudry, and Nathan Srebro. Kernel and rich regimes in overparametrized models. In *Conference on Learning Theory*, pp. 3635–3673. PMLR, 2020.

Liu Ziyin, Botao Li, and Xiangming Meng. Exact solutions of a deep linear network. *arXiv preprint arXiv:2202.04777*, 2022.

Difan Zou, Philip M Long, and Quanquan Gu. On the global convergence of training deep linear resnets. *arXiv preprint arXiv:2003.01094*, 2020.

A PROOFS OF BASIC PROPERTIES OF SEMI-NORM

Proof of Lemma 2.1. The first property is a direct consequence of the definition of the projection matrix P_X.

Notice that

$$\frac{1}{\varepsilon}(L(W + \varepsilon\Delta W) - L(W)) = \frac{1}{\varepsilon}(L(WP_X + \varepsilon\Delta W P_X) - L(WP_X)).$$

By letting $\varepsilon \to 0$, the definition of the directional derivative implies

$$\langle \nabla L(W), \Delta W \rangle_F = \langle \nabla L(WP_X), \Delta W P_X \rangle_F = \langle \nabla L(WP_X)P_X, \Delta W \rangle_F, \forall \Delta W \in \mathbb{R}^{n_y \times n_x},$$

since $P_X = P_X^T$. This completes the proof of the second property.

The third property is obtained based on the fact that the orthogonal projection matrix satisfies $P_X = P_X^T = P_X^2 = P_X^3$, since

$$\langle \nabla L(W), V \rangle_F = \langle \nabla L(WP_X)P_X, V \rangle_F$$
$$= \langle \nabla L(WP_X)P_X^2, V P_X \rangle_F = \langle \nabla L(WP_X)P_X, V \rangle_X = \langle \nabla L(W), V \rangle_X.$$

Set $V = \nabla L(W)$. Then the fourth property is implied by the third property.

For the last property, first recall that $\|W\|_X = \|WP_X\|_F$ and $P_X = X(X^TX)^\dagger X^T$. X is of full row rank if and only if P_X is identity matrix, which completes the proof.

$\square$

Proof of Lemma 2.2. Because X is not full row rank, we know that $I - P_X \neq 0$. There exists W such that $W(I - P_X) \neq 0$. Applying the first property in Lemma 2.1, we have

$$L(\frac{1}{2}W + \frac{1}{2}WP_X) = L((\frac{1}{2}W + \frac{1}{2}WP_X)P_X) = L(WP_X) = \frac{1}{2}L(W) + \frac{1}{2}L(WP_X),$$

provided $W \neq WP_X$.

Hence, L is not strictly convex, which implies L is not strongly convex.

To prove the second property, it suffices to show that $g(W) = L(W) - \frac{\alpha(l)\lambda_{min}(XX^T)}{m}\|W\|_X^2$ is convex. It is obvious that

$$g(W) = L(W) - \frac{\alpha(l)}{m}\sum_{i=1}^{m}\|Wx_i - y_i\|_2^2 + \frac{\alpha(l)}{m}(\|WX - Y\|_F^2 - \lambda_{min}(X^TX)\|W\|_X^2). \quad (14)$$

$L(W) - \frac{\alpha(l)}{m}\sum_{i=1}^{m}\|Wx_i, y_i\|_F^2$ is convex, since $l(\cdot, y_i)$ is strongly convex. The Hessian of $\|WX - Y\|_F^2 - \lambda_{min}(W^TW)\|WP_X\|_F^2$ has no negative eigenvalue, thus the second term in (14) is also convex. This completes the proof. $\square$

B THE EXACT STATEMENTS OF THE MAIN THEOREMS

Define some quantities as follows:

$$q = \begin{cases} 1 - \alpha\eta_*(2 - \eta_*\alpha), & 0 < \eta_* \leq \frac{2}{\alpha+\beta} \\ 1 - \beta\eta_*(2 - \eta_*\beta), & \frac{2}{(\alpha+\beta)} < \eta_* < \frac{2}{\beta}, \end{cases} \quad (15)$$

$$B_\delta = \left(\frac{2 \cdot rank(X)}{\delta} + \|W_*\|_X^2 \right),$$

$$C_1 = n_N \kappa^2 B_\delta \frac{C_0}{(\eta_0 - \eta)^2/\eta_0^2} + \ln N,$$

$$C_2 = n_N \kappa^2 B_\delta C_0 + \ln N,$$

$$C_3 = n_N \kappa^2 B_\delta \frac{C_0}{(\eta_0 - \eta)^2/\eta_0^2} + C_0 \ln(\underline{N}),$$

$$C_4 = n_N \kappa^2 B_\delta \frac{1}{(\eta_0 - \eta)^2/\eta_0^2},$$

$$C_5 = n_N \kappa^2 B_\delta C_0 + C_0 \ln(\underline{N}),$$

$$C_6 = n_N \kappa^2 B_\delta,$$

where $\underline{N}$ denotes the number of distinct elements in the set $\{n_1, \cdots, n_{N-1}\}$, $\eta_1 = \frac{2n_N}{N\beta}$, and $\eta_0 = \frac{2n_N}{e^{2c}N\beta}$ with $c > 0$.

Theorem B.1. *Given any $c > 0$, and $0 < \delta < 1/2$, define $\eta_0 = \frac{2n_N}{e^{2c}N\beta}$, and consider the learning rate $\eta < \eta_0$. There exists a constant $C := C(c)$, such that if*

$$n_{min} \geq C \cdot C_1 \cdot N, \tag{16}$$

then with probability at least $1 - \delta$ over the random Gaussian initialization, we have

$$\mathcal{E}_{DLN}(t) \leq \left(1 - 4e^{-c\frac{\frac{\eta}{\eta_0}(1 - \frac{\eta}{\eta_0})}{\kappa}} \right)^t \mathcal{E}_{DLN}(0).$$

Theorem B.2. *Given any $c > 0$, and $0 < \delta < 1/2$, define $\eta_0 = \frac{2n_N}{e^{2c}\beta N}$, and consider the learning rate $\eta < \eta_0$. There exists a constant $C := C(c)$, such that if*

$$n_{min} \geq C \cdot C_3, \tag{17}$$

then with probability at least $1 - \delta$ over the random one peak projections and embeddings initialization, we have

$$\mathcal{E}_{DLN}(t) \leq \left(1 - 4e^{-c\frac{\frac{\eta}{\eta_0}(1 - \frac{\eta}{\eta_0})}{\kappa}} \right)^t \mathcal{E}_{DLN}(0).$$

Specially, if $n_1 = n_2 = \cdots = n_{N-1} = n \geq \min\{n_N, n_0\}$, then the requirement (17) can be replaced by

$$n \geq C \cdot C_4. \tag{18}$$

Remark 7. Assume $L(a_N W_N \cdots W_1) = \frac{1}{2} \|a_N W_N \cdots W_1 X - Y\|_F^2$, and $n_1 = \cdots = n_{N-1} = n$. Then for Gaussian initialization, our Theorem B.1 leads to Theorem 4.1 in Du & Hu (2019). Similarly, for orthogonal initialization, our Theorem B.2 leads to Theorem 4.1 in Hu et al. (2020).

Next, we present a version of the theorem related to balanced initialization.

Theorem B.3. *Assume $n_1 = \cdots = n_{N-1} = n$. Given any $c > 0$, and $0 < \delta < 1/2$, define $\eta_0 = \frac{2n_N}{e^{2c}\beta N}$, and consider the learning rate $\eta < \eta_0$. There exists a constant $C := C(c)$, such that as long as*

$$n \geq C \cdot C_4. \tag{19}$$

then with probability at least $1 - \delta$ over special balanced initial, we have

$$\mathcal{E}_{DLN}(t) \leq \left(1 - 4e^{-c\frac{\frac{\eta}{\eta_0}(1 - \frac{\eta}{\eta_0})}{\kappa}} \right)^t \mathcal{E}_{DLN}(0).$$

C   INEQUALITIES IN CONVEX OPTIMIZATION

Convex optimization has been studied for about a century. Recall the definitions and basic inequalities for $\alpha-$strongly convex and $\beta-$Lipschitz functions.

Definition C.1. A continues differentiable function f is said to be $\beta-$ Lipschitz if the gradient ∇f is $\beta-$ Lipschitz, that is if for all x, y,

$$\|\nabla f(y) - \nabla f(x)\| \leq \beta \|y - x\|, \tag{20}$$

f is said to be $\alpha-$strongly convex if for all x, y, we have

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{\alpha}{2} \|y - x\|^2. \tag{21}$$

Proposition C.1. *If f is $\alpha-$strongly convex and ∇f is $\beta-$Lipschitz with respect to a (semi-)norm, then $\alpha \leq \beta$ and*

$$\langle \nabla f(x), y - x \rangle + \frac{\alpha}{2} \|y - x\|^2 \leq f(y) - f(x) \leq \langle \nabla f(x), y - x \rangle + \frac{\beta}{2} \|y - x\|^2, \tag{22}$$

$$\langle \nabla f(x) - \nabla f(y), x - y \rangle \geq \frac{\alpha\beta}{\alpha + \beta} \|x - y\|^2 + \frac{1}{\alpha + \beta} \|\nabla f(x) - \nabla f(y)\|^2, \tag{23}$$

$$\|\nabla f(x) - \nabla f(y)\| \geq \alpha \|x - y\|, \tag{24}$$

$$f(x) - f(y) \leq \langle \nabla f(x), x - y \rangle - \frac{1}{2\beta} \|\nabla f(x) - \nabla f(y)\|^2. \tag{25}$$

Proof of Proposition C.1.   We only proof the last inequality.
Let $z = y - \frac{1}{\beta}(\nabla f(y) - \nabla f(x))$. Since f is convex $\beta-$Lipschitz, we have

$$f(z) - f(x) \geq \langle \nabla f(x), z - x \rangle$$

and

$$f(z) - f(y) \leq \langle \nabla f(y), z - y \rangle + \frac{\beta}{2} \|z - y\|^2.$$

Thus,

$$\begin{aligned}
f(x) - f(y) =& f(x) - f(z) + f(z) - f(y) \\
\leq & \langle \nabla f(x), x - z \rangle + \langle \nabla f(y), z - y \rangle + \frac{\beta}{2} \|z - y\|^2 \\
=& \langle \nabla f(x), x - y \rangle - \frac{1}{2\beta} \|\nabla f(x) - \nabla f(y)\|^2.
\end{aligned}$$

$\square$

Before we start to prove Lemma D.1, let us first include and prove the following result.

Lemma C.2. *1. Assume L is $\alpha-$strongly convex, $\alpha > 0$. Denote a global minimizer of L by W_*. Then for any W,*

$$L(W_*) - L(W) \geq -\frac{1}{2\alpha} \|\nabla L(W)\|_X^2. \tag{26}$$

2. Assume ∇L is $\beta-$Lipschitz, then

$$L(W_*) - L(W) \leq -\frac{1}{2\beta} \|\nabla L(W)\|_X^2. \tag{27}$$

Proof of Lemma C.2.   **1.** First, we know that $\nabla L(W_*) = 0$. L is $\alpha-$strongly convex, which implies the inequality (22) holds. Thus

$$L(V) - L(W) \geq \langle \nabla L(W), V - W \rangle_X + \frac{\alpha}{2} \|V - W\|_X^2 =: g(V).$$

Minimizing both sides in terms of V gives (26).

Now we focus on minimizing $g(V)$. Since $g(V) \in C^1$ and the global minimizer exits, we have
$$\nabla g(V^*) = \nabla L(W)P_X + \alpha(V^* - W)P_X = 0,$$
where V^* is a global minimizer for $g(V)$. Thus,
$$g(V^*) = -\frac{1}{2\alpha} \|\nabla L(W)\|_X^2. \tag{28}$$

2. Applying proposition C.1 to a $\beta-$Lipschitz function ∇L, we obtain
$$L(W_*) - L(W)$$
$$\leq \langle \nabla L(W_*), W_* - W \rangle_X - \frac{1}{2\beta} \|\nabla L(W) - \nabla L(W_*)\|_X^2$$
$$= -\frac{1}{2\beta} \|\nabla L(W)\|_X^2.$$

$\square$

D CONVERGENCE REGION

In this section, we study a class of the convergence region for deep linear neural networks, which works for deterministic initialization. Define $A|_{\mathcal{R}(X)} = AX^T(XX^T)^- X = AP_X$, and view $A|_{\mathcal{R}(X)}$ as a linear operator on $\mathcal{R}(X)$.

Recall the optimization problem
$$\underset{W_1, \cdots, W_N}{\text{minimize}} \quad L^N(W_1, \cdots W_N) := \frac{1}{m} \sum_{i=1}^m l(a_N W_{N:1} x_i, y_i) = L(a_N W_{N:1}), \tag{29}$$

and GD
$$\begin{cases} W_j(t+1) = W_j(t) - \eta \frac{\partial L^N}{\partial W_j}(W_1(t), \cdots, W_N(t)), j = 1, \cdots, N, \\ \text{where } \frac{\partial L^N}{\partial W_j}(W_1, \cdots, W_N) = a_N(W_{N:j+1})^T \nabla L(a_N W_{N:1})(W_{j-1:1})^T, \end{cases} \tag{30}$$
where the normalization factor $a_N = \frac{1}{\sqrt{n_1 n_2 \cdots n_{N-1} n_N}}$.

The following theorem generalizes the idea from the recent work (Du & Hu, 2019; Hu et al., 2020).

For notational convenience, we denote $W_{j:i}(t) = W_j(t) \cdots W_i(t), L_t = L(a_N W_{N:1}(t)), \nabla L_t = \nabla L(a_N W_{N:1}(t))$ etc.

Lemma D.1. *Assume the initialization satisfies the following conditions simultaneously:*
$$\begin{cases} \sigma_{max}(W_{N:i+1}(0)) \leq e^{c_1/2}(n_{N-1:i})^{1/2}, 1 \leq i \leq N-1, \\ \sigma_{min}(W_{N:i+1}(0)) \geq e^{-c_2/2}(n_{N-1:i})^{1/2}, 1 \leq i \leq N-1, \\ \sigma_{max}(W_{i-1:1}(0)|_{\mathcal{R}(X)}) \leq e^{c_1/2}(n_{i-1:1})^{1/2}, 2 \leq i \leq N, \\ \sigma_{min}(W_{i-1:1}(0)|_{\mathcal{R}(X)}) \geq e^{-c_2/2}(n_{i-1:1})^{1/2}, 2 \leq i \leq N, \\ \|W_{j:i}(0)\| \leq M/2 \cdot N^\theta(\prod_{i \leq k \leq j-1} n_k \cdot \max\{n_{i-1}, n_j\})^{1/2}, 1 < i \leq j < N, \\ L_0 - L(W_*) \leq \beta B_0 =: B, \end{cases} \tag{31}$$
where c_1, c_2, M are positive constant and $\theta \geq 0$.
Notice that B_0 is a proper upper bound for $\|a_N W_{N:1}(0)\|_X^2 + \|W_*\|_X^2$.

Set the learning rate $\eta = \frac{(1-\varepsilon)2n_N}{e^{6c_1 + 3c_2}\beta N}$, where $0 < \varepsilon < 1$. Define $\gamma = \frac{2e^{6c_1}\varepsilon\alpha N}{n_N}$.

Assume that
$$n_{min} \geq \frac{C(c_1, c_2)M^2\kappa^2 B_0}{\varepsilon^2} N^{2\theta} n_N. \tag{32}$$

Then the GD (30) satisfies
$$L_t - L(W_*) \leq (1 - \eta\gamma)^t (L_0 - L(W_*)), t = 1, 2, \cdots.$$

Definition D.1. For given $c_1, c_2, M, B_0 > 0$, and $\theta \geq 0$, we define the convergence region $\mathcal{R}(c_1, c_2, \theta, M, B_0)$ by the set of initialization that satisfies the inequality system (31).

Remark 8. The condition (31) describes the convergence region for initialization and the condition (32) describes the overparameterization for deep linear neural networks. At this time, it is not clear how large this convergence region is. Later, we will show that the properly scaled random initialization with some extra mild overparameterization conditions will fall into this convergence region with high probability.

Proof of Lemme D.1. To prove Lemma D.1, it suffices to show that the following three properties hold $\mathcal{A}(t), \mathcal{B}(t)$, and $\mathcal{C}(t)$ for all $t = 0, 1, \cdots$.

1. $\mathcal{A}(t)$:
$$L_t - L(W_*) \leq (1 - \eta\gamma)^t (L_0 - L(W_*)).$$

2. $\mathcal{B}(t)$:
$$\begin{cases} \sigma_{max}(W_{N:i+1}(t)) \leq e^{c_1}(n_{N-1:i})^{1/2}, 1 \leq i \leq N-1, \\ \sigma_{min}(W_{N:i+1}(t)) \geq e^{-c_2}(n_{N-1:i})^{1/2}, 1 \leq i \leq N-1, \\ \sigma_{max}(W_{i-1:1}(t)|_{\mathcal{R}(X)}) \leq e^{c_1}(n_{i-1:1})^{1/2}, 2 \leq i \leq N, \\ \sigma_{min}(W_{i-1:1}(t)|_{\mathcal{R}(X)}) \geq e^{-c_2}(n_{i-1:1})^{1/2}, 2 \leq i \leq N, \\ \|W_{j:i}(t)\| \leq M \cdot N^\theta (\frac{1}{n_{min}} \prod_{i-1 \leq k \leq j} n_k)^{1/2}, 1 < i \leq j < N. \end{cases}$$

3. $\mathcal{C}(t)$:
$$\|W_i(t) - W_i(0)\|_F \leq \frac{2e^{2c_1}\sqrt{2\beta B}}{\sqrt{n_N}\gamma} =: R, 1 \leq i \leq N.$$

Using simultaneous induction, the proof of Lemma D.1 is divided into the following 3 claims.

Claim 1. $\mathcal{A}(0), \cdots, \mathcal{A}(t), \mathcal{B}(0), \cdots, \mathcal{B}(t) \implies \mathcal{C}(t+1)$.

Claim 2. $\mathcal{C}(t) \implies \mathcal{B}(t)$, if $n_{min} \geq \frac{C(c_1,c_2)M^2\kappa^2B_0}{\varepsilon^2}N^{2\theta}n_N$, where $C(c_1, c_2)$ is a positive constant only depend on c_1, c_2.

Claim 3. $\mathcal{A}(t), \mathcal{B}(t) \implies \mathcal{A}(t+1)$, if $n_{min} \geq C(c_1,c_2)M^2B_0N^{2\theta}n_N$, where $C(c_1, c_2)$ is a positive constant only depend on c_1, c_2.

$\square$

Proof of Claim 1. As a consequence of Lemma C.2 and Lemma 2.1, and $\mathcal{A}(s), s \leq t$, we have

$$\begin{aligned} \|\nabla L(a_N W_{N:1}(s))\|_F^2 &= \|\nabla L_s - \nabla L(W_* P_X)\|_X^2 \\ &\leq 2\beta[L_s - L(W_*)] \\ &\leq 2\beta(1 - \eta\gamma)^s B. \end{aligned} \tag{33}$$

From $\mathcal{A}(0), \cdots, \mathcal{A}(t), \mathcal{B}(0), \cdots, \mathcal{B}(t)$, we have for any $0 \leq s \leq t$,

$$\begin{aligned} \left\|\frac{\partial L}{\partial W_i}(s)\right\|_F &\leq a_N \|W_{N:i+1}(s)\| \|\nabla L(a_N W_{N:1}(s))\|_F \|W_{i-1:1}(s)|_{\mathcal{R}(X)}\| \\ &\leq \frac{e^{2c_1}}{\sqrt{n_N}} \|\nabla L(a_N W_{N:1}(s))\|_F \\ &\leq \frac{e^{2c_1}}{\sqrt{n_N}} \sqrt{2\beta(1 - \eta\gamma)^s B}. \end{aligned} \tag{34}$$

Then,

$$
\begin{aligned}
\|W_i(t+1) - W_i(0)\|_F &\leq \sum_{s=0}^{t} \|W_i(s+1) - W_i(s)\|_F \\
&= \sum_{s=0}^{t} \left\| \eta \frac{\partial L}{\partial W_i}(s) \right\|_F \\
&\leq \eta \frac{e^{2c_1}}{\sqrt{n_N}} \sqrt{2\beta B} \sum_{s=0}^{t} (1 - \eta\gamma)^{s/2} \\
&\leq \eta \frac{e^{2c_1}}{\sqrt{n_N}} \sqrt{2\beta B} \sum_{s=0}^{t} (1 - \eta\gamma/2)^{s} \\
&\leq \frac{2e^{2c_1}\sqrt{2\beta B}}{\sqrt{n_N}\gamma} = R.
\end{aligned}
$$

This proves $\mathcal{C}(t+1)$. $\qquad\square$

Proof of Claim 2. Let $\delta_i = W_i(t) - W_i(0), 1 \leq i \leq N$. Using $\mathcal{C}(t)$, we have $\|\delta_i\|_F \leq R, 1 \leq i \leq N$. Set $\varepsilon_1 = e^{-c_1/2}\min\{e^{c_1} - e^{c_1/2}, e^{-c_2/2} - e^{-c_2}, 1/2\}$.
It is suffices to show that

$$\|W_{N:i}(t) - W_{N:i}(0)\| \leq e^{c_1/2}\varepsilon_1(n_{N-1}n_{N-1}\cdots n_{i-1})^{1/2}, 1 < i \leq N, \tag{35}$$

$$\left\|(W_{i:1}(t) - W_{i:1}(0))|_{\mathcal{R}(X)}\right\| \leq e^{c_1/2}\varepsilon_1(n_1 n_2 \cdots n_{i-1})^{1/2}, 1 \leq i < N, \tag{36}$$

and

$$\|W_{j:i}(t) - W_{j:i}(0)\| \leq M/2 \cdot N^{\theta} \left(\frac{1}{n_{min}} \prod_{i-1 \leq k \leq j} n_k \right)^{1/2}, 1 < i \leq j < N, \tag{37}$$

because $\sigma_{min}(A+B) \geq \sigma_{min}(A) - \sigma_{max}(B) = \sigma_{min}(A) - \|B\|$ and $\sigma_{max}(A+B) \leq \sigma_{max}(A) + \sigma_{max}(B) = \|A\| + \|B\|$ (e.g. see Theorem 1.3 in Chafaı̈ et al. (2009)).
Case 1. We first prove (37).
For $1 \leq i < j \leq N$, we can write $W_{j:i}(t) = (W_j(0) + \delta_j)\cdots(W_i(0) + \delta_i)$.
Expanding the above product, each term has the form:

$$W_{j:(k_s+1)}(0) \cdot \delta_{k_s} \cdot W_{(k_s-1):(k_{s-1}+1)}(0) \cdot \delta_{k_{s-1}} \cdots \delta_{k_1} \cdot W_{(k_1-1):i}(0), \tag{38}$$

where $i \leq k_1 < \cdots < k_s \leq j$ are positions at which perturbation terms δ_{k_l} are taken out.
Notice that the convergence region assumptions (31) implies that for any $1 < i \leq j < N$,

$$\|W_{j:i}(0)\| \leq M/2 \cdot N^{\theta} \left(\prod_{i \leq k \leq j-1} n_k \cdot \max\{n_{i-1}, n_j\} \right)^{1/2} \leq M \cdot N^{\theta} \left(\frac{\prod_{i-1 \leq k \leq j} n_k}{n_{min}} \right)^{1/2}. \tag{39}$$

WLOG, assume $M \geq 1$. If $i = j+1$, then

$$\|W_{j:i}(0)\| = \|I\| \leq M \cdot N^{\theta}(n_j/n_{min})^{1/2}.$$

Assume $i > 1, j < N$, applying inequality (39) as well as the following inequality

$$\sum_{s=1}^{j-i+1} \binom{j-i+1}{s} x^s = (1+x)^{j-i+1} - 1 \leq (1+x)^N - 1, \forall x \geq 0,$$

we obtain that

$$
\begin{aligned}
&\|W_{j:i}(t) - W_{j:i}(0)\| \\
&\leq \sum_{s=1}^{j-i+1} \binom{j-i+1}{s} R^s (M \cdot N^{\theta})^{s+1} n_{min}^{-s/2}(n_{i-1}\cdots n_j/n_{min})^{1/2} \\
&\leq M \cdot N^{\theta}(n_{i-1}\cdots n_j/n_{min})^{1/2}[(1 + R \cdot M \cdot N^{\theta}/\sqrt{n_{min}})^N - 1] \\
&\leq \varepsilon_1 M \cdot N^{\theta}(n_{i-1}\cdots n_j/n_{min})^{1/2}.
\end{aligned}
$$

The last line holds due to the following reasons:
there exists absolute constant $A_1, A_2 > 0$ such that

$$(1 + x)^N - 1 \le A_2 x N,$$

if $x \ge 0$, $N \ge 1$, and $xN \le A_1$. Since there exists positive constant $C(c_1, c_2)$, which only depends on c_1, c_2, such that when

$$n_{min} \ge \frac{C(c_1, c_2) M^2 \kappa^2 B_0}{\varepsilon^2} N^{2\theta} n_N \tag{40}$$

we can have

$$R \cdot M \cdot N^{\theta+1} / \sqrt{n_{min}} \le A_1,$$

as well as

$$[(1 + R \cdot M \cdot N^\theta / \sqrt{n_{min}})^N - 1] \le A_2 \cdot M \cdot R \cdot N^{\theta+1} / \sqrt{n_{min}} \le \varepsilon_1 = \varepsilon_1(c_1, c_2).$$

Case 2. The proof of (35) is similar. Set $j = N$, we can save the factor $M \cdot N^\theta$ from previous calculation, which means

$$\|W_{N:i}(t) - W_{N:i}(0)\|$$
$$\le e^{c_1/2} \sum_{s=1}^{N-i+1} \binom{N-i+1}{s} R^s (M \cdot N^\theta)^s n_{min}^{-s/2} (n_{i-1} \cdots n_{N-1})^{1/2}$$
$$\le e^{c_1/2} (n_{i-1} \cdots n_{N-1})^{1/2} [(1 + R \cdot M \cdot N^\theta / \sqrt{n_{min}})^N - 1]$$
$$\le e^{c_1/2} \varepsilon_1 (n_{i-1} \cdots n_{N-1})^{1/2}, i \ge 2,$$

where the last line is implied by equation (40).
Case 3. Similarly, we have

$$\left\|W_{j:1}(t)|_{\mathcal{R}(X)} - W_{j:1}(0)|_{\mathcal{R}(X)}\right\|$$
$$\le e^{c_1/2} \sum_{s=1}^{j} \binom{j}{s} R^s (M \cdot N^\theta)^s n_{min}^{-s/2} (n_1 \cdots n_j)^{1/2}$$
$$\le e^{c_1/2} (n_1 \cdots n_j)^{1/2} [(1 + R \cdot M \cdot N^\theta / \sqrt{n_{min}})^N - 1]$$
$$\le e^{c_1/2} \varepsilon_1 (n_1 \cdots n_j)^{1/2}, j \le N - 1$$

This proves $\mathcal{B}(t)$.

$\square$

Proof of Claim 3. The GD (7) implies

$$W_{N:1}(t+1)$$
$$= \left(W_N(t) - \eta \frac{\partial L^N}{\partial W_N}(t)\right) \left(W_{N-1}(t) - \eta \frac{\partial L^N}{\partial W_{N-1}}(t)\right) \cdots \left(W_1(t) - \eta \frac{\partial L^N}{\partial W_1}(t)\right)$$
$$= W_{N:1}(t) - \eta \cdot a_N \sum_{i=1}^{N} W_{N:i+1}(t) W_{N:i+1}^T(t) \nabla L(a_N W_{N:1}(t))(W_{i-1:1}(t))^T (W_{i-1:1}(t)) + E(t),$$

where $E(t)$ contains all high-order terms (those with η^2 or higher). Define a linear operator

$$P(t)[A] = a_N^2 \sum_{i=1}^{N} W_{N:i+1}(t) W_{N:i+1}^T(t)(A P_X)(W_{i-1:1}(t)|_{\mathcal{R}(X)})^T W_{i-1:1}(t)|_{\mathcal{R}(X)}, \tag{41}$$

for any $A \in \mathbb{R}^{n_N \times n_0}$.

Now we have

$$a_N W_{N:1}(t+1) = a_N W_{N:1}(t) - \eta \cdot P(t)[\nabla L(a_N W_{N:1}(t) P_X)] + a_N E(t). \tag{42}$$

Easy to check that $P(t)[\cdot]$ is a sum of positive semidefinite linear operator.

The following proposition describes the eigenvalues of the linear operator $P(t)[\cdot]$.

Proposition D.2. *Let S_1, S_2 be symmetric matrices. Suppose $S_1 = U\Lambda_1 U^T$, $S_2 = V\Lambda_2 V^T$, where $U = [u_1, u_2, \cdots, u_m]$, and $V = [v_1, v_2, \cdots, v_n]$ are othogonal matrices, and $\Lambda_1 = diag(\lambda_1, \lambda_2, \cdots, \lambda_m)$ and $\Lambda_2 = diag(\mu_1, \mu_2, \cdots, \mu_n)$ are diagonal matrices. Then the linear operator $L(A) := S_1 A S_2$ is orthogonally diagonalizable, and $L(A_{ij}) = \lambda_i \mu_j A_{ij}$, where $\lambda_i \mu_j$ represent all eigenvalues corresponding to their eigenvectors $A_{ij} = u_i v_j^T$.*

Applying this proposition and the assumption $\mathcal{B}(t)$, we obtain the upper bound and lower bound for the maximum and minimum eigenvalues of positive definite operator $P(t)$, respectively,

$$\lambda_{max}(P(t)) \leq a_N^2 \sum_{i=1}^{N} \sigma_{max}^2(W_{i-1:1}(t)|_{\mathcal{R}(X)}) \cdot \sigma_{max}^2(W_{N:i+1}(t)) \leq \frac{N}{n_N} e^{2c_1},$$

and

$$\lambda_{min}(P(t)) \geq a_N^2 \sum_{i=1}^{N} \sigma_{min}^2(W_{i-1:1}(t)|_{\mathcal{R}(X)}) \cdot \sigma_{min}^2(W_{N:i+1}(t)) \geq \frac{N}{n_N} e^{-2c_2}. \quad (43)$$

In conclusion, we have

$$\lambda_{max}(P(t)) \leq \frac{N}{n_N} e^{2c_1}, \text{ and } \lambda_{min}(P(t)) \geq \frac{N}{n_N} e^{-2c_2}. \quad (44)$$

With learning rate $\eta = \eta_\varepsilon = \frac{(1-\varepsilon)2n_N}{e^{6c_1+3c_2}\beta N}, 0 < \varepsilon < 1$, we have

$$\begin{aligned}
&L_{t+1} - L_t \\
&\leq \langle \nabla L_t, -\eta P(t)[\nabla L_t]\rangle_X + \langle \nabla L_t, a_N E(t)\rangle_X + \frac{\beta}{2} \|\eta P(t)[\nabla L_t] - a_N E(t)\|_X^2 \\
&= \langle \nabla L_t, -\eta P(t)[\nabla L_t]\rangle + \frac{\beta}{2}\eta^2 \|P(t)[\nabla L_t]\|_X^2 + F(t) \\
&\leq -\left(\eta\lambda_{min}(P(t)) - \frac{\beta}{2}\eta^2\lambda_{max}^2(P(t))\right)\|\nabla L_t\|_X^2 + F(t) \\
&\leq -e^{-2c_2}\frac{N}{n_N}\eta\left(1 - e^{4c_1+2c_2}\frac{\beta}{2}\eta\frac{N}{n_N}\right)\|\nabla L_t\|_X^2 + F(t),
\end{aligned} \quad (45)$$

where

$$F(t) = \langle \nabla L_t, a_N E(t)\rangle_X + \frac{\beta}{2}\|\eta P(t)[\nabla L_t] - a_N E(t)\|_X^2 - \frac{\beta}{2}\eta^2\|P(t)[\nabla L_t]\|_X^2.$$

We claim that $F(t)$ is small enough, such that

$$\begin{aligned}
&L_{t+1} - L_t \\
&\leq -e^{-2c_2}\frac{N}{n_N}\eta\left(1 - e^{4c_1+2c_2}\frac{\beta}{2}\eta\frac{N}{n_N}\right)\|\nabla L_t\|_X^2 + F(t) \\
&\leq -e^{-3c_2}\frac{N}{n_N}\eta\left(1 - e^{6c_1+3c_2}\frac{\beta}{2}\eta\frac{N}{n_N}\right)\|\nabla L_t\|_X^2 \\
&= -e^{-6(c_1+c_2)}\frac{2\varepsilon(1-\varepsilon)}{\beta}\|\nabla L_t\|_X^2.
\end{aligned} \quad (46)$$

Assuming this claim for the moment, we complete the proof. Combining (26) and (46), we have

$$\begin{cases}
L_{t+1} - L_t \leq -e^{-6(c_1+c_2)}\frac{2\varepsilon(1-\varepsilon)}{\beta}\|\nabla L_t\|_X^2, \\
L(W_*) - L_t \geq -\frac{1}{2\alpha}\|\nabla L_t\|_X^2,
\end{cases}$$

which implies

$$L_{t+1} - L(W_*) \leq \left(1 - e^{-6(c_1+c_2)}\frac{4\varepsilon(1-\varepsilon)}{\kappa}\right)(L_t - L(W_*)), \quad (47)$$

that is

$$L_t - L(W_*) \le \left(1 - e^{-6(c_1+c_2)}\frac{4\varepsilon(1-\varepsilon)}{\kappa}\right)^t (L_0 - L(W_*)) = (1 - \eta\gamma)^t (L_0 - L(W_*)). \quad (48)$$

Estimate $F(t)$

Notice that

$$|F(t)|$$
$$\le \|\nabla L_t\|_X \|a_N E(t)\|_X + \frac{\beta}{2}(2\eta\lambda_{max}(P(t))\|\nabla L_t\|_X \|a_N E(t)\|_X + \|a_N E(t)\|_X^2)$$
$$=: I_1 + I_2.$$

From (34), we have

$$\left\|\frac{\partial L}{\partial W_i}(t)\right\|_F \le \frac{e^{2c_1}}{\sqrt{n_N}}\|\nabla L(a_N W_{N:1}(t))\|_F = \frac{e^{2c_1}}{\sqrt{n_N}}\|\nabla L(a_N W_{N:1}(t))\|_X =: K.$$

Expanding the product

$$W_{N:1}(t+1) = \left(W_N(t) - \eta\frac{\partial L^N}{\partial W_N}(t)\right)\left(W_{N-1}(t) - \eta\frac{\partial L^N}{\partial W_{N-1}}(t)\right)\cdots\left(W_1(t) - \eta\frac{\partial L^N}{\partial W_1}(t)\right),$$

each term has the form:

$$\Delta = W_{N:(k_s+1)}(t) \cdot \eta\frac{\partial L}{\partial W_{k_s}}(t) \cdot W_{(k_s-1):(k_{s-1}+1)}(t) \cdot \eta\frac{\partial L}{\partial W_{k_{s-1}}}(t)\cdots\eta\frac{\partial L}{W_{k_1}}(t) \cdot W_{(k_1-1):1}(t),$$

where $1 \le k_1 < k_2 < \cdots < k_s \le N$.

As a direct consequence of inequality $\mathcal{B}(t)$ and inequality (39), we obtain

$$\|\Delta\|_X = \|\Delta P_X\|_F \le \frac{1}{a_N\sqrt{n_N}}e^{2c_1}(\eta K)^s\left(\frac{M \cdot N^\theta}{\sqrt{n_{min}}}\right)^{s-1},$$

Recall that $E(t)$ contains all high-order terms (those with η^2 or higher) in the expansion of the product. Thus, $E(t)$ can be expressed as follows:

$$\sum_{s=2}^N \sum_{1 \le k_1 < k_2 < \cdots < k_s \le N} W_{N:(k_s+1)}(t) \cdot \eta\frac{\partial L}{\partial W_{k_s}}(t) \cdot W_{(k_s-1):(k_{s-1}+1)}(t) \cdot \eta\frac{\partial L}{\partial W_{k_{s-1}}}(t)\cdots\eta\frac{\partial L}{W_{k_1}}(t) \cdot W_{(k_1-1):1}(t).$$

Set $\xi = \min\{(e^{-2c_2} - e^{-3c_2})/e^{4c_1+1}, \frac{1}{4}(e^{6c_1} - e^{4c_1})/e^{6c_1+1}, \frac{1}{2}(e^{6c_1} - e^{4c_1})^{1/2}/e^{4c_1+1}, 1\}$.

Recall the inequality $\binom{N}{s} \le (eN)^s$. Thus, we have

$$a_N\|E(t)\|_X$$
$$\le \frac{1}{\sqrt{n_N}}e^{2c_1}\sum_{s=2}^N \binom{N}{s}(\eta K)^s\left(\frac{M \cdot N^\theta}{\sqrt{n_{min}}}\right)^{s-1}$$
$$\le \frac{1}{\sqrt{n_N}}\left(\frac{M \cdot N^\theta}{\sqrt{n_{min}}}\right)^{-1}e^{2c_1}\sum_{s=2}^N (eN)^s(\eta K)^s\left(\frac{M \cdot N^\theta}{\sqrt{n_{min}}}\right)^s$$
$$\quad (49)$$
$$\le \frac{1}{\sqrt{n_N}}e^{2c_1}(\eta eKN)\frac{\eta eKM \cdot N^{\theta+1}/\sqrt{n_{min}}}{1 - \eta eKM \cdot N^{\theta+1}/\sqrt{n_{min}}}$$
$$\le \xi\frac{N}{n_N}\eta \cdot e^{4c_1+1}\|\nabla L(a_N W_{N:1}(t))\|_X \ (\text{if } \eta eKM \cdot N^{\theta+1}/\sqrt{n_{min}} < \xi/(1+\xi))$$
$$= \xi \cdot e^{4c_1+1}\left(\eta\frac{N}{n_N}\right)\|\nabla L(a_N W_{N:1}(t))\|_X.$$

Using (33) and the upper bound of η, we know that there exists constant $C(c_1, c_2)$, such that

$$n_{min} \ge C(c_1, c_2)M^2 \cdot B_0 N^{2\theta}n_N,$$

and

$$\eta e K M \cdot N^{\theta+1}/\sqrt{n_{min}} \leq \frac{2\sqrt{2}M \cdot e^{1+2c_1}\sqrt{B_0}N^\theta\sqrt{n_N}}{\sqrt{n_{min}}} = \frac{1}{C'(c_1,c_2)} \leq \frac{\xi}{2} \leq \frac{\xi}{1+\xi}.$$

Using (49), we have

$$I_1 \leq \xi \cdot e^{4c_1+1}\left(\eta\frac{N}{n_N}\right)\|\nabla L_t\|_X^2 \leq (e^{-2c_2} - e^{-3c_2})\left(\eta\frac{N}{n_N}\right)\|\nabla L_t\|_X^2, \qquad (50)$$

and

$$\begin{aligned}
&I_2\\
\leq&\frac{\beta}{2}\left(2\xi \cdot e^{6c_1+1}\left(\eta^2\frac{N^2}{n_N^2}\right)\|\nabla L_t\|_X^2 + \xi^2 \cdot e^{8c_1+2}\left(\eta^2\frac{N^2}{n_N^2}\right)\|\nabla L_t\|_X^2\right)\\
\leq&(e^{6c_1} - e^{4c_1})\frac{\beta}{2}\eta^2\frac{N^2}{n_N^2}\|\nabla L_t\|_X^2 .
\end{aligned}$$

Thus, (46) valid.
This proves $\mathcal{A}(t)$.

$\square$

As a direct consequence of the proof Lemma D.1, we can obtain the following lemma.

Lemma D.3. *Assume all assumptions in Lemma D.1 hold. For any $\tau > 0$, we can choose new constants c_1, c_2 as well as $C := C(c_1, c_2)$ such that the overparameterization assumption (32) in Lemma D.1 hold and*

$$\|R(t)\|_X \leq \tau \|a_N W_{N:1}(t) - W_*\|_X, \qquad (51)$$

where

$$a_N W_{N:1}(t+1) = a_N W_{N:1}(t) - \frac{N}{n_N}\eta\nabla L(a_N W_{N:1}(t)) + R(t).$$

Proof of Lemma D.3. Due to (33), (42), (44), (49), and lemma C.2, we have

$$\begin{aligned}
\|R(t)\|_X &= \left\|a_N E(t) + \eta\left(\frac{N}{n_N}\nabla L_t - P(t)[\nabla L_t]\right)\right\|_X\\
&\leq \|a_N E(t)\|_X + \eta\max\left\{\lambda_{max}(P(t)) - \frac{N}{n_N}, \frac{N}{n_N} - \lambda_{min}(P(t))\right\}\|\nabla L_t\|_X\\
&\leq(C' \cdot \xi + \max\{e^{2c_1} - 1, 1 - e^{-2c_2}\}) \cdot \eta\frac{N}{n_N} \cdot \|\nabla L_t\|_X\\
&\leq\frac{2\sqrt{2\beta(L_t - L(W_*))}}{e^{6c_1+3c_2} \cdot \beta} \cdot (C' \cdot \xi + \max\{e^{2c_1} - 1, 1 - e^{-2c_2}\}).
\end{aligned}$$

Due to the fact that $L_t - L(W_*)$ is non-increasing in t, and C' is a constant only depend on c_1, c_2, we can choose small enough positive c_1, c_2 and ξ, which depends on τ, such that

$$\|R(t)\|_X \leq \tau\frac{\sqrt{2\beta(L_t - L(W_*))}}{\beta} \leq \tau\|a_N W_{N:1}(t) - W_*\|_X.$$

$\square$

Lemma D.4. *Assume $\tau \in [0, 1)$. Consider a discrete dynamical system $V(t)$ such that,*

$$V(t+1) = V(t) - \eta_*\nabla L(V(t)) + R(t),$$

where $\|R(t)\|_X \leq \tau\|V(t) - W_*\|_X$. If $\eta_* \leq 2/\beta$, we have

$$\|V(t) - W_*\|_X^2 \leq (q + 7\tau)^t \|V(0) - W_*\|_X^2,$$

where q is defined in (15).

Proof of Lemma D.4. Set $\Delta(t) = V(t) - W_*$ and $\tau' = \tau \|\Delta(t)\|_X$. Notice that

$$\Delta(t+1) = \Delta(t) - \eta_*(\nabla L(V(t)) - \nabla L(W_*)) + R(t),$$

and

$$\begin{aligned}
&\|\Delta(t+1)\|_X^2\\
&\leq \eta_*^2 \|\nabla L(V(t)) - \nabla L(W_*)\|_X^2 - 2\eta_*\langle\Delta(t), \nabla L(V(t)) - \nabla L(W_*)\rangle_X\\
&\quad + \|\Delta(t)\|_X^2 + (2\|\Delta(t)\|_X + 2\eta_*\|\nabla L(V(t)) - \nabla L(W_*)\|_X + \tau')\tau'.
\end{aligned}$$

By inequality (23),

$$\begin{aligned}
&\|\Delta(t+1)\|_X^2\\
&\leq \|\Delta(t)\|_X^2 - 2\eta_*\langle\Delta(t), \nabla L(V(t)) - \nabla L(W_*)\rangle_X\\
&\quad + \eta_*^2 \|\nabla L(V(t)) - \nabla L(W_*)\|_X^2 + 7\tau\|\Delta(t)\|_X^2\\
&= (1+7\tau)\|\Delta(t)\|_X^2 - 2\eta_*\langle\Delta(t), \nabla L(V(t)) - \nabla L(W_*)\rangle_X\\
&\quad + \eta_*^2 \|\nabla L(V(t)) - \nabla L(W_*)\|_X^2\\
&\leq (1+7\tau)\|\Delta(t)\|_X^2 - 2\eta_*\frac{\alpha\beta}{\alpha+\beta}\|\Delta(t)\|_X^2\\
&\quad + \left(\eta_*^2 - \frac{2\eta_*}{\alpha+\beta}\right)\|\nabla L(V(t)) - \nabla L(W_*)\|_X^2.
\end{aligned}$$

Case 1: $\frac{2}{\alpha+\beta} < \eta_* < \frac{2}{\beta}$.

In this case, we have

$$\begin{aligned}
&\|\Delta(t+1)\|_X^2\\
&\leq (1+7\tau)\|\Delta(t)\|_X^2 - 2\eta_*\frac{\alpha\beta}{\alpha+\beta}\|\Delta(t)\|_X^2 + \left(\eta_*^2 - \frac{2\eta_*}{\alpha+\beta}\right)\|\nabla L(V(t)) - \nabla L(W_*)\|_X^2\\
&\leq (1+7\tau)\|\Delta(t)\|_X^2 - 2\eta_*\frac{\alpha\beta}{\alpha+\beta}\|\Delta(t)\|_X^2 + \left(\eta_*^2 - \frac{2\eta_*}{\alpha+\beta}\right)\beta^2\|\Delta(t)\|_X^2\\
&\leq (1 + 7\tau - \beta\eta_*(2 - \eta_*\beta))\|\Delta(t)\|_X^2\\
&= (q + 7\tau)\|\Delta(t)\|_X^2.
\end{aligned}$$

Case 2: $0 < \eta_* \leq \frac{2}{\alpha+\beta}$.

Similarly, we have

$$\|\Delta(t+1)\|_X^2 \leq (1 + 7\tau - \alpha\eta_*(2 - \eta_*\alpha))\|\Delta(t)\|_X^2 = (q + 7\tau)\|\Delta(t)\|_X^2.$$

In both cases, we have $\|\Delta(t+1)\|_X^2 \leq (q + 7\tau)\|\Delta(t)\|_X^2$.

Thus, $\|\Delta(t)\|_X^2 \leq (q + 7\tau)^t \|\Delta(0)\|_X^2$.

$\square$

Next, we will show that the trajectories of the GD (30) for deep linear neural networks (29) are close to those of GD (2) for the corresponding convex problem (1).

Lemma D.5. *Consider the GD for the deep linear neural networks (30) with learning rate $\eta < \eta_1$ for $a_N W_{N:1}(t), t = 0, 1, \cdots$, and the GD (2) with learning rate $\eta_* = \frac{N}{n_N}\eta$ for $W(t), t = 0, 1, \cdots$.*

Assume $C(c_1, c_2)$ exists in Lemma D.1 for any $c_1, c_2 > 0$. For any $\tau \in (0, 1)$, $\eta < \eta_1$ (η_1 defined in B), we can choose $c_1, c_2 > 0$ and the constant $C = C(c_1, c_2) = C'(\tau, \eta/\eta_1)$, such that inequality (51) holds, given initialization condition (31), and overparameterization condition

$$n_{min} \geq C M^2 \kappa^2 B_0 N^{2\theta} n_N. \tag{52}$$

Furthermore, we have

$$\|a_N W_{N:1}(t) - W(t)\|_X^2 \le D(\tau, q, t) \|a_N W_{N:1}(0) - W_*\|_X^2, \tag{53a}$$

$$|\mathcal{E}_{DLN}(t) - \mathcal{E}(t)| \le \beta \left(q^{t/2} \sqrt{D(\tau, q, t)} + \frac{1}{2} D(\tau, q, t) \right) \|a_N W_{N:1}(0) - W_*\|_X^2, \tag{53b}$$

$$\mathcal{E}_{DLN}(t) \le 3\beta(q + \tau)^t \|a_N W_{N:1}(0) - W_*\|_X^2, \tag{53c}$$

where $D(\tau, q, t) = \min \left\{ \frac{\tau}{1-q}, 2(q + \tau)^t \right\}$, with q defined in (15).

Proof of Lemma D.5. Using Lemma D.3, we obtain that for any $\tau \in (0, 1)$ and $\eta < \eta_1$, we can find small enough positive constant c_1, c_2, which are only depend on $\tau, \eta/\eta_1$, and constant $C = C(c_1, c_2) = C''(\tau, \eta/\eta_1)$ mentioned in Lemma D.3, such that

$$\eta = \frac{(1 - \varepsilon)2n_N}{e^{6c_1 + 3c_2}\beta N},$$

where $0 < \varepsilon < 1$, as well as

$$V(t + 1) = V(t) - \eta_* \nabla L(V(t)) + R(t),$$

where $V(t) = a_N W_{N:1}(t)$, $\eta_* = \frac{N}{n_N}\eta$, and $\|R(t)\|_X \le \tau' = \tau \|V(t) - W_*\|_X$.

Notice that $\theta_0 := \eta/\eta_1 = \frac{1-\varepsilon}{e^{6c_1 + 3c_2}}$ and $\eta/\eta_0 = 1 - \varepsilon$, where $\eta_0 = \frac{2n_N}{e^{6c_1 + 3c_2}\beta N}$.

For the right hand side of inequality (32), we have

$$\frac{C(c_1, c_2)M^2 \kappa^2 B_0}{\varepsilon^2} N^{2\theta} n_N = \frac{C''(\tau, \eta/\eta_1)M^2 \kappa^2 B_0}{\varepsilon^2} N^{2\theta} n_N.$$

To show that inequality (32) is equivalent to inequality (52), it suffices to show that ε only depend on $\tau, \eta/\eta_1$. Notice that

$$\varepsilon = 1 - \eta/\eta_0 = 1 - \theta_0 e^{6c_1 + 3c_2},$$

and c_1, c_2 only depend on τ and η/η_1, which implies ε only depend on $\tau, \eta/\eta_1$.

Now, we start to prove the three inequalities in (53).

Recall the GD (2) for $W(t)$. Define $\Delta(t) = V(t) - W(t) = a_N W_{N:1}(t) - W(t)$. Notice that

$$\Delta(t + 1) = \Delta(t) - \eta_*(\nabla L(V(t)) - \nabla L(W(t))) + R(t),$$

and

$$\begin{aligned}
&\|\Delta(t + 1)\|_X^2 \\
\le & \eta_*^2 \|\nabla L(V(t)) - \nabla L(W(t))\|_X^2 - 2\eta_* \langle \Delta(t), \nabla L(V(t)) - \nabla L(W(t)) \rangle_X \\
& + \|\Delta(t)\|_X^2 + (2 \|\Delta(t)\|_X + 2\eta_* \|\nabla L(V(t)) - \nabla L(W(t))\|_X + \tau')\tau'.
\end{aligned}$$

Let $l_t = 2 \|\Delta(t)\|_X + 2\eta_* \|\nabla L(V(t)) - \nabla L(W(t))\|_X + \tau'$.

Now, we aim to find an upper bound for l_t.

Applying lemma C.2 with the assumption $0 < \eta_* = \frac{N}{n_N}\eta < \frac{2}{\beta}$, we know that

$$l_t \le (6 \|\Delta(t)\|_X + \tau') \le 7(\|W(t) - W_*\|_X + \|V(t) - W_*\|_X). \tag{54}$$

Thus

$$l_t \tau' \le 7\tau \|V(t) - W_*\|_X (\|V(t) - W_*\|_X + \|W(t) - W_*\|_X) =: U_t \tau.$$

By inequality (23),

$$\|\Delta(t+1)\|_X^2$$
$$\leq \|\Delta(t)\|_X^2 - 2\eta_* \langle \Delta(t), \nabla L(V(t)) - \nabla L(W(t)) \rangle_X$$
$$+ \eta_*^2 \|\nabla L(V(t)) - \nabla L(W(t))\|_X^2 + U_t \tau$$
$$= \|\Delta(t)\|_X^2 - 2\eta_* \langle V(t) - W(t), \nabla L(V(t)) - \nabla L(W(t)) \rangle_X$$
$$+ \eta_*^2 \|\nabla L(V(t)) - \nabla L(W(t))\|_X^2 + U_t \tau$$
$$\leq \|\Delta(t)\|_X^2 - 2\eta_* \frac{\alpha\beta}{\alpha+\beta} \|\Delta(t)\|_X^2$$
$$+ \left(\eta_*^2 - \frac{2\eta_*}{\alpha+\beta}\right) \|\nabla L(V(t)) - \nabla L(W(t))\|_X^2 + U_t \tau.$$

Case 1: $\frac{2}{\alpha+\beta} < \eta_* < \frac{2}{\beta}$.
In this case, we have

$$\|\Delta(t+1)\|_X^2$$
$$\leq \|\Delta(t)\|_X^2 - 2\eta_* \frac{\alpha\beta}{\alpha+\beta} \|\Delta(t)\|_X^2 + \left(\eta_*^2 - \frac{2\eta_*}{\alpha+\beta}\right) \|\nabla L(V(t)) - \nabla L(W(t))\|_X^2 + U_t \tau$$
$$\leq \|\Delta(t)\|_X^2 - 2\eta_* \frac{\alpha\beta}{\alpha+\beta} \|\Delta(t)\|_X^2 + \left(\eta_*^2 - \frac{2\eta_*}{\alpha+\beta}\right) \beta^2 \|\Delta(t)\|_X^2 + U_t \tau$$
$$\leq (1 - \beta\eta_*(2 - \eta_*\beta)) \|\Delta(t)\|_X^2 + U_t \tau$$
$$=: q \|\Delta(t)\|_X^2 + U_t \tau.$$

Case 2: $0 < \eta_* \leq \frac{2}{\alpha+\beta}$.
Similarly, we have

$$\|\Delta(t+1)\|_X^2 \leq (1 - \alpha\eta_*(2 - \eta_*\alpha)) \|\Delta(t)\|_X^2 + U_t \tau =: q \|\Delta(t)\|_X^2 + U_t \tau. \tag{55}$$

In both cases, we have $0 < q < 1$.

First of all, since $U_t \leq U_0$ and $\|\Delta(0)\|_X = 0$, we obtain that

$$\|\Delta(t)\|_X^2 \leq \frac{U_0\tau}{1-q} + q^t \left(\|\Delta(0)\|_X^2 - \frac{U_0\tau}{1-q}\right) \leq \frac{U_0\tau}{1-q} \leq \frac{14\tau}{1-q} \|V(0) - W_*\|_X^2.$$

Applying Lemma D.4 for $V(t)$ and $W(t)$, we obtain $\|V(t) - W_*\|_X^2 \leq (1+\varepsilon)^t q^t \|V(0) - W_*\|_X^2$ and $\|W(t) - W_*\|_X^2 \leq q^t \|W(0) - W_*\|_X^2$, respectively. Thus,

$$|L(W(t)) - L(a_N W_{N:1}(t))|$$
$$\leq |\langle \nabla L(W(t)), \Delta(t) \rangle_X| + \frac{\beta}{2} \|\Delta(t)\|_X^2$$
$$\leq \beta \|W(t) - W_*\|_X \cdot \|\Delta(t)\|_X + \frac{\beta}{2} \|\Delta(t)\|_X^2$$
$$\leq \beta \left(q^{t/2} \sqrt{\frac{14\tau}{1-q}} + \frac{7\tau}{1-q}\right) \|V(0) - W_*\|_X^2.$$

Generally speaking, (55) implies

$$\|\Delta(t)\|_X^2 \leq \tau \sum_{j=0}^{t-1} q^{t-1-j} U_j.$$

We have

$$\|\Delta(t)\|_X^2 \leq 14\tau \sum_{j=0}^{t-1} (q + 7\tau)^j q^{t-1-j} \|V(0) - W_*\|_X^2$$
$$\leq 2(q + 7\tau)^t \left(1 - \left(\frac{q}{q + 7\tau}\right)^t\right) \|V(0) - W_*\|_X^2$$

Thus, we have

$$\|a_N W_{N:1}(t) - W(t)\|_X^2 \leq \min\left\{\frac{14\tau}{1-q}, 2(q+7\tau)^t\right\} \|V(0) - W_*\|_X^2,$$

as well as

$$|L(W(t)) - L(a_N W_{N:1}(t))|$$

$$\leq \beta \|W(t) - W_*\|_X \cdot \|\Delta(t)\|_X + \frac{\beta}{2}\|\Delta(t)\|_X^2$$

$$\leq \beta \left(\sqrt{\min\left\{\frac{14\tau}{1-q}, 2(q+7\tau)^t\right\}} \cdot q^{t/2} + \frac{1}{2}\min\left\{\frac{14\tau}{1-q}, 2(q+7\tau)^t\right\}\right) \|V(0) - W_*\|_X^2.$$

By triangle inequality as well as $L(W(t)) - L(W_*) \leq \frac{\beta}{2}q^t \|V(0) - W_*\|_X^2$, we have

$$|L(a_N W_{N:1}(t)) - L(W_*)| \leq 3\beta(q+7\tau)^t \|V(0) - W_*\|_X^2.$$

Without loss of generality, we replace all 14τ and 7τ by τ, which completes the proof.

$\square$

E GAUSSIAN INITIALIZATION FALL INTO THE CONVERGENCE REGION

In this section, we first establish some spectral properties of the products of random Gaussian matrices. The spectral properties lead to the conclusion that overparameterization guarantees that the random initialization will fall into the convergence region with high probability.

Gaussian initialization:
Denote by $N(0,1)$ the standard Gaussian distribution, and χ_k^2 the chi square distribution with k degrees of freedom. Let $S^{d-1} = \{x \in \mathbb{R}^d; \|x\|_2 = 1\}$ be the unit sphere in $\mathbb{R}^d$.
The scaling factor $a_N = \frac{1}{\sqrt{n_1 n_2 \cdots n_N}}$ ensures that the networks at initialization preserves the norm of every input in expectation.

Lemma E.1. *For any $x \in \mathbb{R}^{n_0}$, the Gaussian initialization satisfies*

$$\mathbb{E}\left[\|a_N W_{N:1}(0)x\|_2^2\right] = \|x\|_2^2.$$

Proof of Lemma E.1. For random matrix $A \in \mathbb{R}^{n_i \times n_{i-1}}$ with i.i.d $N(0,1)$ entries and any vector $0 \neq v \in \mathbb{R}^{n_{i-1}}$, the distribution of $\frac{\|Av\|_2^2}{\|v\|_2^2}$ is $\chi_{n_i^2}$. We rewrite

$$\|W_{N:1}(0)x\|_2^2 / \|x\|_2^2 = Z_N Z_{N-1} \cdots Z_1,$$

where $Z_i = \|W_{i:1}(0)x\|^2 / \|W_{i-1:1}(0)x\|^2$.
Then we know that the distribution of random variable $Z_1 \sim \chi_{n_1}^2$, and conditional distribution of random variables $Z_i|(Z_1, \cdots, Z_{i-1}) \sim \chi_{n_i}^2 (1 < i \leq N)$. Thus, $Z_1, \cdots, Z_{n_i}$ are independent. By law of iterated expectations, we have

$$\mathbb{E}[\|W_{N:1}(0)x\|_2^2 / \|x\|_2^2] = \prod_{j=1}^{N} n_j.$$

$\square$

Define $\Delta_1 = \sum_{j=1}^{N-1} 1/n_j$. Now, we introduce a new notation $\Omega\left(\frac{1}{\Delta_1}\right)$, which means that there exists $k > 0$, such that $\Omega\left(\frac{1}{\Delta_1}\right) \geq \frac{k}{\Delta_1}$.

Lemma E.2. *Consider real random matrix $A_j \in \mathbb{R}^{n_j \times n_{j-1}}, 1 \leq j \leq q$ with i.i.d $N(0,1)$ entries and any vector $0 \neq x \in \mathbb{R}^{n_1}$.*
Define $\Delta_1(q) = \sum_{j=1}^{q} \frac{1}{n_j}$ and $n_{min} = \min_{1 \leq j \leq q} n_j$. Then

$$\mathbb{P}(\|A_q A_{q-1} \cdots A_1 x\|_2^2 / \|x\|_2^2 > e^c n_1 \cdots n_q) \leq \exp\left\{-\frac{c^2}{8\Delta_1(q)}\right\} =: f_1(c), \forall c > 0. \quad (56)$$

When $0 < c \leq 3 \ln 2$, $\Delta_1(q) \leq c/(12 \ln 2)$, we have

$$\mathbb{P}(\|A_q A_{q-1} \cdots A_1 x\|_2^2 / \|x\|_2^2 < e^{-c} n_1 \cdots n_q) \leq \exp\left\{-\frac{c^2}{36 \ln(2)\Delta_1(q)}\right\} =: f_2(c). \quad (57)$$

Hence, for any $x \in S^{n_0 - 1}$ with probability at least $1 - e^{-\Omega(\frac{1}{\Delta_1(q)})}$, we have

$$e^{-c_2/2}(n_1 \cdots n_q)^{1/2} \leq \|A_q \cdots A_1 x\|_2 \leq e^{c_1/2}(n_1 \cdots n_q)^{1/2},$$

when $0 < c_2 \leq 3 \ln 2$, $\Delta_1(q) \leq c_2/(12 \ln 2)$.

Proof of Lemma E.2. For random matrix $A_i \in \mathbb{R}^{n_i \times n_{i-1}}$ with i.i.d $N(0,1)$ entries and any vector $0 \neq v \in \mathbb{R}^{n_{i-1}}$, the random variable $\frac{\|A_i v\|_2^2}{\|v\|_2^2}$ is distributed as $\chi_{n_i}^2$. We rewrite

$$\|A_q \cdots A_1 x\|_2^2 / \|x\|_2^2 = Z_q Z_{q-1} \cdots Z_1,$$

where $Z_i = \|A_{i:1}x\|^2 / \|A_{i-1:1}x\|^2$. We have $Z_1 \sim \chi_{n_1}^2$, $Z_i | (Z_1, \cdots, Z_{i-1}) \sim \chi_{n_i}^2 (1 < i \leq q)$. Recall the moments of $Z \sim \chi_m^2$:

$$\mathbb{E}[Z^\lambda] = \frac{2^\lambda \Gamma(\frac{m}{2} + \lambda)}{\Gamma(\frac{m}{2})}, \forall \lambda > -\frac{m}{2}.$$

Now, we aim to find the Chernoff type bound.

Case 1: We define ratio of Gamma function

$$R(x, \lambda) = \frac{\Gamma(x + \lambda)}{\Gamma(x)}, \lambda > 0, x > 0.$$

In Jameson (2013), we have

$$R(x, \lambda) \leq x(x + \lambda)^{\lambda - 1} \leq (x + \lambda)^\lambda, \lambda > 0, x > 0. \quad (58)$$

Fixed $c > 0$, for any $\lambda > 0$ we have

$$
\begin{aligned}
\mathbb{P}(Z_q \cdots Z_1 > e^c n_1 \cdots n_q) &\leq \mathbb{P}((Z_q \cdots Z_1)^\lambda > e^{\lambda c}(n_1 \cdots n_q)^\lambda) \\
&\leq e^{-\lambda c}(n_1 \cdots n_q)^{-\lambda} \mathbb{E}[(Z_q \cdots Z_1)^\lambda] && \text{(Markov inequality)} \\
&= \exp\{-\lambda(c + \ln(n_1 \cdots n_q))\} \prod_{j=1}^q 2^\lambda R(n_j/2, \lambda) && \text{(Law of total expectation)} \\
&\leq \exp\{-\lambda(c + \ln(n_1 \cdots n_q)) + q\lambda \ln 2 + \sum_{j=1}^q \lambda \ln(\frac{n_j}{2} + \lambda)\} && \text{(Inequality (58))} \\
&= \exp\{-\lambda c + \lambda \sum_{j=1}^q \ln(1 + \frac{2\lambda}{n_j})\} \\
&\leq \exp\{-\lambda c + 2\lambda^2 \sum_{j=1}^q \frac{1}{n_j}\}.
\end{aligned}
$$

Define constant $\Delta_1(q) = \sum_{j=1}^q \frac{1}{n_j}$. Set $\lambda = \frac{c}{4\Delta_1(q)}$, we obtain (56).

Case 2: Let $n_{min} = \min_{1 \leq j \leq q} n_j$.

$$
\begin{aligned}
\mathbb{P}(Z_q \cdots Z_1 < e^{-c} n_1 \cdots n_q) &\leq \mathbb{P}((Z_q \cdots Z_1)^\lambda > e^{-\lambda c}(n_1 \cdots n_q)^\lambda) \\
&\leq \exp\{\lambda(c - \ln(n_1 \cdots n_q)) + q\lambda \ln 2 + \sum_{j=1}^q \ln R(\frac{n_j}{2}, \lambda)\}.
\end{aligned}
$$

Define

$$f(\lambda) = \lambda(c - \ln(n_1 \cdots n_q)) + q\lambda \ln 2 + \sum_{j=1}^{q} \ln R(\frac{n_j}{2}, \lambda), -\frac{n_{min}}{2} < \lambda \leq 0.$$

Notice that $f(0) = 0$. Define digamma function,

$$\psi(x) = \frac{d}{dx} \ln(\Gamma(x)) = \frac{\Gamma'(x)}{\Gamma(x)}.$$

Qi et al. (2006) proved the following sharp inequality of digamma function,

$$\ln(x + \frac{1}{2}) - \frac{1}{x} < \psi(x) < \ln(x + e^{-\gamma}) - \frac{1}{x}, x > 0,$$

where γ is the Euler-Mascheroni constant, and $e^{-\gamma} \approx 0.561459$.
Thus,

$$f'(\lambda) = c + \sum_{j=1}^{q} \left[-\ln(\frac{n_j}{2}) + \psi(\frac{n_j}{2} + \lambda) \right] \geq c + \sum_{j=1}^{q} \ln(1 + \frac{\lambda + 1/2}{n_j/2}) - \sum_{j=1}^{q} \frac{1}{n_j/2 + \lambda}.$$

Since $\ln(1 + x)$ is concave, we have

$$\ln(1 + x) \geq 2\ln(2)x, x \in [-1/2, 0].$$

If $-\frac{n_{min}}{4} \leq \lambda \leq 0$, then

$$f(\lambda) = f(0) - \int_{\lambda}^{0} f'(x)dx$$

$$\leq c\lambda + \int_{0}^{\lambda} \left[\sum_{j=1}^{q} \ln(1 + \frac{x + 1/2}{n_j/2}) - \sum_{j=1}^{q} \frac{1}{n_j/2 + x} \right] dx$$

$$= c\lambda + \sum_{j=1}^{q} \left[\lambda \ln(1 + \frac{\lambda + 1/2}{n_j/2}) + (n_j/2 + 1/2) \ln(1 + \frac{\lambda}{n_j/2 + 1/2}) - \lambda - \ln(1 + \frac{\lambda}{n_j/2}) \right]$$

$$\leq c\lambda + \sum_{j=1}^{q} (\lambda - 1) \ln(1 + \frac{\lambda}{n_j/2})$$

$$\leq c\lambda + 4\ln(2)\lambda(\lambda - 1)\Delta_1(q).$$

Assume $0 < c \leq 3\ln 2$. Let $A = 12\ln 2$, and $\lambda^* = -\frac{c}{A\Delta_1(q)}$. Since $n_{min}\Delta_1(q) \geq 1$, we have $\lambda^* \geq -n_{min}/4$.
Assume $\Delta_1(q) \leq c/(12\ln 2)$.
Thus

$$f(\lambda^*) \leq -\frac{c^2}{A\Delta_1(q)} + 4\ln 2 \frac{c^2}{A\Delta_1(q)} \left(\frac{\Delta_1(q)}{c} + \frac{1}{A} \right) \leq -\frac{c^2}{36\ln(2)\Delta_1(q)}. \tag{59}$$

Thus, we obtain (57). $\qquad\square$

Lemma E.3. *There exists a positive constant $C(c_1, c_2)$ which only depends on c_1, c_2, such that if $n_N\Delta_1 \leq C(c_1, c_2)$, then for any fixed $1 < i \leq N$, with probability at least $1 - \exp\left\{ -\Omega\left(\frac{1}{\Delta_1} \right) \right\}$ we have*

$$\sigma_{\max}(W_{N:i}(0)) \leq e^{c_1}(n_{i-1}n_i \cdots n_{N-1})^{1/2}, \tag{60}$$

and

$$\sigma_{\min}(W_{N:i}(0)) \geq e^{-c_2}(n_{i-1}n_i \cdots n_{N-1})^{1/2}. \tag{61}$$

Proof of Lemma E.3. Let $A = W_{N:i}^T(0)$. We know that

$$\sigma_{max}(A) = \|A\| = \sup_{v \in S^{n_N - 1}} \|Av\|_2$$

and

$$\sigma_{min}(A) = \inf_{v \in S^{n_N-1}} \|Av\|_2.$$

Applying lemma E.2, we know that with probability at least $1 - \exp\left\{-\Omega\left(\frac{1}{\Delta_1}\right)\right\}$,

$$\|Av\|_2 / \|v\|_2 \in [e^{-c_2/2}P, e^{c_1/2}P],$$

where $P = (n_{i-1} \cdots n_{N-1})^{1/2}$.

Set $\phi = \min\{1 - e^{-c_1/2}, (e^{-c_2/2} - e^{-c_2})/(e^{-c_2/2} + e^{c_1})\}$. Take a ϕ-net $\mathcal{N}_\phi$ for S^{n_N-1} with size $|\mathcal{N}_\phi| \le (3/\phi)^{n_N}$. Notice that with this size we can actually cover the unit ball, not only the unit sphere.

Thus, with probability at least $1 - |\mathcal{N}_\phi| \exp\left\{-\Omega\left(\frac{1}{\Delta_1}\right)\right\}$, for all $u \in \mathcal{N}_\phi$ simultaneously we have

$$\|Au\|_2 / \|u\|_2 \in [e^{-c_2/2}P, e^{c_1/2}P].$$

Fixed $v \in S^{n_N-1}$, there exists $u \in \mathcal{N}_\phi$ such that $\|u - v\|_2 \le \phi$. WLOG, we assume $1 - \phi \le \|u\|_2 \le 1$. We obtain

$$\|Av\|_2 \le \|Au\|_2 + \|A(u - v)\|_2 \le e^{c_1/2}P + \phi\|A\|.$$

Taking supereme over $\|v\|_2 = 1$, we obtain

$$\sigma_{max}(A) = \|A\| \le \frac{e^{c_1/2}}{1 - \phi}P \le e^{c_1}P.$$

For the lower bound, we have

$$\|Av\|_2 \ge \|Au\|_2 - \|A(u - v)\|_2 \ge e^{-c_2/2}P\|u\| - \phi\|A\| \ge \left[(1 - \phi)e^{-c_2/2} - \phi e^{c_1}\right]P \ge e^{-c_2}P.$$

Taking the infimum over $\|v\|_2 = 1$, we get

$$\sigma_{min}(A) \ge e^{-c_2}P.$$

The conclusions hold with probability at least

$$1 - |\mathcal{N}_\phi| \exp\left\{-\Omega\left(\frac{1}{\Delta_1}\right)\right\}$$

$$\ge 1 - \exp\{n_N \ln(3/\phi)\} \exp\left\{-\Omega\left(\frac{1}{\Delta_1}\right)\right\}$$

$$\ge 1 - \exp\left\{-\Omega\left(\frac{1}{\Delta_1}\right)\right\},$$

since $n_N \Delta_1 \le C(c_1, c_2)$. $\qquad\square$

Lemma E.4. *There exists a positive constant $C(c_1, c_2)$ which only depends on c_1, c_2, such that if $rank(X)\Delta_1 \le C(c_1, c_2)$, then for any fixed $1 \le j < N$, with probability at least $1 - \exp\{-\Omega\left(\frac{1}{\Delta_1}\right)\}$ we have*

$$\sigma_{\max}(W_{j:1}(0)|_{\mathcal{R}(X)}) \le e^{c_1}(n_1 n_2 \cdots n_j)^{1/2}, \tag{62}$$

and

$$\sigma_{\min}(W_{j:1}(0)|_{\mathcal{R}(X)}) \ge e^{-c_2}(n_1 n_2 \cdots n_j)^{1/2}. \tag{63}$$

Proof of Lemma E.4. The proof is similar to that of previous lemma. The only difference is that now we consider the ϕ−net to cover the unit sphere in $\mathcal{R}(X) \cap \mathbb{R}^{n_0}$, with $\dim \mathcal{R}(X) \cap \mathbb{R}^{n_0} = rank(X)$, where $\mathcal{R}(X)$ represents the column space of X. $\qquad\square$

Lemma E.5. *Set $C = n_{max}/n_{min} < \infty$, $\theta = 1/2$. Assume $\Omega(1/\Delta_1) \geq \frac{k}{\Delta_1}$, where $0 < k < 1$ is a constant and Δ_1 satisfies*

$$\begin{cases} \Delta_1 \leq \min\left\{\frac{k}{5\ln(6)}, \frac{k}{5\ln(5\ln(6)e/k)}\right\} \\ \Delta_1 \ln(C) \leq \min\left\{\frac{k}{5\ln(5\ln(6)e/k)}, \frac{k}{5}\right\} \\ \Delta_1 \ln(N^{2\theta}) \leq k/5. \end{cases}$$

Given $1 < i \leq j < N$, with probability at least $1 - 2e^{-k/(5\Delta_1)} = 1 - e^{-\Omega(1/\Delta_1)}$ we have

$$\|W_{j:i}(0)\| \leq M_k\sqrt{C}N^\theta(n_i \cdots n_{j-1} \cdot \max\{n_{i-1}, n_j\})^{1/2},$$

where M_k is a positive constant that only depends on k.

Proof of Lemma E.5. WLOG, assume $n_{i-1} \leq n_j$. Let $A = W_{j:i}(0)$. From lemma E.2, we know that fixed $v \in S^{n_{i-1}-1}$, with probability at least $1 - e^{-\Omega(1/\Delta_1)}$ we have $\|Av\|_2 \leq 4/3(n_i \cdots n_j)^{1/2}$.
.
Take a small constant $c = \frac{kN^{2\theta}}{5\ln(6)\Delta_1 n_{i-1}} \geq \frac{k}{5\ln(6)C}$. Let $v_1, \cdots, v_{n_{i-1}}$ be an orthonormal basis for $R^{n_{i-1}}$. Partition the index set $\{1, 2, \cdots, v_{n_{i-1}}\} = S_1 \cup S_2 \cup \cdots \cup S_{\lceil N^{2\theta}/c\rceil}$, where $|S_l| \leq \lceil cn_{i-1}/N^{2\theta}\rceil$ for each $1 \leq l \leq \lceil N^{2\theta}/c\rceil$.
The following discussion is similar to the proof of lemma E.3, hence we omit some details. For each l, taking a $1/2-$ net $\mathcal{N}_l$ for the set $V_{S_l} = \{v \in S^{n_{i-1}-1}; v \in span\{v_i; i \in S_l\}\}$, we can get

$$\|Au\|_2 \leq 4(n_i \cdots n_j)^{1/2}, u \in V_{S_l},$$

with probability at least

$$1 - |\mathcal{N}_l|e^{-k/\Delta_1} \geq 1 - \exp\{-k/\Delta_1 + (cn_{i-1}/N + 1)\ln 6\} \geq 1 - e^{-3k/(5\Delta_1)},$$

since $\Delta_1 \leq \frac{k}{5\ln(6)}$.
Therefore, for any $v \in \mathbb{R}^{n_{i-1}}$, we can write it as the sum $v = \sum_l a_l v_l$, where $\alpha_l \in \mathbb{R}$ and $v_l \in V_{S_l}$ for each l. We also know that $\|v\|_2^2 = \sum_{l\geq 1} |\alpha_l|^2$.
Then we have

$$\|Av\|_2 \leq \sum_l |\alpha_l|\|Av_l\|_2 \leq 4(n_i \cdots n_j)^{1/2}\sqrt{\lceil N^{2\theta}/c\rceil \sum_l |a_l|^2} \leq M_k\sqrt{C}N^\theta(n_i \cdots n_j)^{1/2}\|v\|_2.$$

Thus,

$$\|A\| \leq M_k\sqrt{C}N^\theta(n_i \cdots n_j)^{1/2}.$$

Notice that when $C \leq e$, $\Delta_1 \leq \frac{k}{5\ln(5\ln(6)e/k)} \leq \frac{k}{5\ln(5\ln(6)\cdot C/k)}$, and when $C > e$, we have

$$\Delta_1 \ln(C) \leq \min\left\{\frac{k}{5\ln(5\ln(6)e/k)}, k/5\right\} \leq \frac{k\ln(C)}{5\ln(5\ln(6)\cdot C/k)}.$$

The success probability is at least

$$1 - \lceil N^{2\theta}/c\rceil \cdot e^{-3k/(5\Delta_1)}$$
$$\geq 1 - \exp\left\{\ln\left(\frac{5\ln(6)\cdot C}{k}\right) + \ln(N^{2\theta}) - 3k/(5\Delta_1)\right\} - e^{-3k/(5\Delta_1)}$$
$$\geq 1 - 2e^{-k/(5\Delta_1)},$$

since

$$\Delta_1 \leq \frac{k}{5\ln\left(5\ln(6)\cdot C/k\right)} \text{ and } \Delta_1 \ln(N^{2\theta}) \leq k/5.$$

$\square$

Proof of Lemma 2.3. Set $r = rank(X)$, and $u_1, \cdots, u_r$ be an orthonormal basis of the column space of X.

Then, $P_X = \sum_{i=1}^{r} u_i u_i^T$.

Notice that

$$\|a_n W_{N:1}(0)\|_X^2 = \|a_n W_{N:1}(0) P_X\|_F^2 = \sum_{i=1}^{r} \|a_n W_{N:1}(0) u_i\|_2^2.$$

By assumption, we have

$$\mathbb{E} \|a_n W_{N:1}(0)\|_X^2 = \mathbb{E} \sum_{i=1}^{r} \|a_n W_{N:1}(0) u_i\|_2^2 = r.$$

The Markov inequality implies

$$\mathbb{P}(\|a_n W_{N:1}(0)\|_X^2 \geq \frac{2r}{\delta}) \leq \frac{\delta}{2}.$$

Therefore, we can bound the initial loss value as

$$L_0 - L(W_*) \leq \langle \nabla L(W_*), a_N W_{N:1}(0) X - W_* \rangle + \frac{\beta}{2} \|a_N W_{N:1}(0) - W_*\|_X^2$$

$$= \frac{\beta}{2} \|a_N W_{N:1}(0) - W_*\|_X^2$$

$$\leq \beta(\|a_N W_{N:1}(0)\|_X^2 + \|W_*\|_X^2)$$

$$\leq \beta(\frac{2r}{\delta} + \|W_*\|_X^2),$$

with probability at least $1 - \delta/2$.

$\square$

Proof of Theorem B.1. The requirement on size $\{n_1, n_2, \cdots, n_{N-1}, N\}$ in (16) makes sure that lemma E.3, E.4, E.5, 2.3, and D.1 hold.
WLOG, we set $c_1 = c/6, c_2 = c/3, M = 2M_k\sqrt{C_0}, B_0 = B_\delta$, and $\eta =: \frac{(1-\varepsilon)2n_N}{e^{2c}\beta N}$, then with probability at least

$$1 - N^2 e^{-\Omega(1/\Delta_1)} - \delta/2 \geq 1 - \delta, \text{ since } \Delta_1 \leq \frac{1}{C(c)} \min \left\{ \frac{1}{\ln N}, \frac{1}{\ln(1/\delta)} \right\},$$

the random initialization satisfies the initialization assumption (31) and the overparameterization assumption (32). Applying Lemma D.1, we complete the proof.

$\square$

F ORTHOGONAL INITIALIZATION FALL INTO THE CONVERGENCE REGION

There are some basic facts for random projections and embeddings. Most of the following properties can be found in Eaton (1989).

Proposition F.1.

1. *A is a random embedding if and only if A^T is a random projection.*

2. *If A is a square matrix, then random projection, random embedding and random orthogonal matrix are equivalent.*

3. *The uniform distribution on the group is a left and right invariant probability measure, that is, if A is a random orthogonal matrix, then A, UA, AU are all random orthogonal matrix, where U is a non-random orthogonal matrix.*

4. *Assume X is a $n \times q(q \leq n)$ random matrix whose entries are i.i.d. $N(0,1)$ random variables. Then $A := X(X^T X)^{-1/2}$ is a random embedding, since $A^T A = I_q$ and the distribution of A is left invariant, which means that A and UA have the same distribution, where U is a non-random orthogonal matrix.*

5. *If A is a uniform distribution over an orthogonal group of order n and A is partitioned as $A = (A_1, A_2)$, where A_1 is $n \times q$ and A_2 is $n \times (n-q)$, then A_1^T and A_2^T are both random orthogonal matrix.*

6. *The columns of uniform distribution over orthogonal group of order n, and*

$$\frac{(\xi_1, \cdots, \xi_n)}{\sqrt{\xi_1^2 + \xi_2^2 + \cdots + \xi_n^2}}$$

   have the same distribution, where $\xi_1, \cdots, \xi_n$ are i.i.d. $N(0,1)$ random variables.

7. *Assume $A = A_{n \times p}, n \leq p$ is a random orthogonal projection. For any $v \in S^{p-1}, \|Av\|_2^2$ and $(\sum_{i=1}^n \xi_i^2)/(\sum_{j=1}^p \xi_j^2)$ are both following beta distribution with $\alpha = n/2, \beta = (p - n)/2$, where $\xi_1, \cdots, \xi_n$ are i.i.d. $N(0,1)$ random variables.*

Remark 9. There are several ways to construct random matrix $A = (a_{ij})_{q \times n}, q \leq n$, which is uniformly distributed over rectangular matrices with $AA^T = c^2 I_q, c > 0$. Let O_n be uniformly distributed over real orthogonal group of order n, and O_n is partitioned as $O_n = (A_1^T, A_2^T)^T$, where A_1 is $q \times n$. Assume $X = (x_{ij})_{q \times n}$, and x_{ij} are independent standard normal random variables. Then A, cA_1, and $c(XX^T)^{-1/2}X$ have the same distribution.

Lemma F.2. *For any $x \in \mathbb{R}^{n_0}$, the one peak random projections and embedding initiation satisfies*

$$\mathbb{E}\left[\|a_N W_{N:1}(0)x\|_2^2\right] = \|x\|_2^2.$$

Proof. Let $D = W_{p:1}(0)/\sqrt{n_1 n_2 \cdots n_p}$. Then D is an embedding matrix. Thus, $\|Dx\|_2^2 = \|x\|_2^2$. Let $A_i = W_{i:p+1}(0)/\sqrt{n_p n_{p+1} \cdots n_{i-1}}$, where $i \geq p + 1$, and $A_p = I$.

Set $B_i = \|A_i Dx\|_2^2 / \|A_{i-1} Dx\|_2^2, i \geq p + 1$. Then, B_i follows beta distribution $B(n_i/2, (n_{i-1} - n_i)/2)$ given $B_{i-1}, B_{i-2}, \cdots, B_{p+1}, i \geq p + 1$. If $n_i = n_{i-1}$, then $B_i|(B_{i-1}, B_{i-2}, \cdots, B_{p+1}) = 1$, a.s.

If $B \sim B(a, b)$, then the expectation is given by the following equation,

$$\mathbb{E}B = \frac{a}{a + b}.$$

Thus, by law of total expectation, we have

$$\frac{n_N}{n_p}\mathbb{E}\|a_N W_{N:1}(0)x\|_2^2 = \mathbb{E}\|A_N Dx\|_2^2 = \mathbb{E}B_N B_{N-1} \cdots B_{p+1}\|Dx\|_2^2 = \frac{n_N}{n_p}\|x\|_2^2.$$

This completes the proof. $\qquad\square$

Next, we introduce sub-Gaussian random variables, associated with bounds on how a random variables deviate their expected value.

Definition F.1. A random variable X with finite mean $\mu = \mathbb{E}X$ is sub-Gaussian if there is a positive number σ such that:

$$\mathbb{E}[\exp(\lambda(X - \mu))] \leq \exp\left(\frac{\lambda^2 \sigma^2}{2}\right) \text{ for all } \lambda \in \mathbb{R} \tag{64}$$

Such a constant σ^2 is called a proxy variance, and we say that X is σ^2-sub-Gaussian, and we write $X \sim SG(\sigma^2)$.

Example F.1. Normal distribution $N(\mu, \sigma^2)$ of course is σ^2 sub-Gaussian.
For beta distribution, Elder (2016) showed that $B(a, b)$ is $\frac{1}{4(a+b)+2}$-sub-Gaussian and later, Marchal & Arbel (2017) concluded $\frac{1}{4(a+b+1)}$-sub-Gaussian.

The Hoeffding bound for random variable X with mean μ and sub-Gaussian parameter σ is given by,

$$\mathbb{P}\left[|X - \mu| \geq t\right] \leq 2\exp\left\{-\frac{t^2}{2\sigma^2}\right\}, \forall t \geq 0. \tag{65}$$

Simply applying the Chernoff bound for $B(a, b)$, we obtain the following lemma.

Lemma F.3. *Assume random variable B distributed as beta distribution $B(a, b)$ with two positive shape parameters a and b. Then*

$$\mathbb{P}(\left|B - \frac{a}{a+b}\right| \geq y) \leq 2\exp\left\{-2(a+b)y^2\right\}, y \geq 0.$$

Hence,

$$\mathbb{P}\left(\left|B - \frac{a}{a+b}\right| \leq \varepsilon\frac{a}{a+b}\right) \geq 1 - \exp\{-\Omega(a^2/(a+b))\},$$

where $\Omega(\cdot)$ only depend on ε.
For the upper tail, we can obtain a better bound,

$$\mathbb{P}\left(B \geq (1+\varepsilon)\frac{a}{a+b}\right) \leq \exp\left\{-(\varepsilon - \ln(\varepsilon+1))a\right\}. \tag{66}$$

Proof of Lemma F.3. We only need to prove the third inequality. Assume random variable $B \sim B(a, b)$. Set $v = a + b$, $(1+t)\frac{a}{v} \leq y < 1, t > 0$, and $r > 0$.
We are going to estimate the Chernoff bound for B, which is

$$\mathbb{P}(B \geq y) \leq e^{-(ry - \ln \mathbb{E}e^{rB})} =: e^{-I_r(y)}.$$

The moment generating function of B is given by

$$\mathbb{E}e^{rB} = 1 + \sum_{k=1}^{\infty} \frac{a(a+1)\cdots(a+k-1)}{v(v+1)\cdots(v+k-1)}\frac{r^k}{k!} \leq 1 + \sum_{k=1}^{\infty} \frac{a(a+1)\cdots(a+k-1)}{v^k}\frac{r^k}{k!}, r > 0.$$

Recall that the Maclaurin series of $(1 - r/v)^{-a}$ over $(-v, v)$, is given by equation

$$(1 - r/v)^{-a} = 1 + \sum_{k=1}^{\infty} \frac{a(a+1)\cdots(a+k-1)}{v^k}\frac{r^k}{k!}.$$

Thus,

$$I_r(y) = ry - \ln \mathbb{E}e^{rB} \geq ry + a\ln(1 - r/v).$$

Set $r = v - a/y \in (0, v)$. We obtain

$$\mathbb{P}(B \geq y) \leq \exp\{-(vy - a + a\ln(a/(vy)))\} =: \exp\{-vy \cdot g(a/(vy))\}, (1+t)\frac{a}{v} \leq y < 1$$

where $g(x) = 1 - x + x\ln(x)$, $x = a/(vy) \in (0, 1/(1+t)]$. Notice that $g(1) = 0$ and $g'(x) = \ln(x) < 0$ over $x \in (0, 1)$.
We know that

$$g(x) \geq g(1/(1+t)) = \frac{t - \ln(1+t)}{t+1}, t > 0.$$

Thus,

$$\mathbb{P}(B \geq y) \leq \exp\left\{-vy \cdot \frac{t - \ln(1+t)}{t+1}\right\} = \exp\left\{-(t - \ln(1+t))a\right\}, y = (1+t)\frac{a}{v} < 1.$$

Set $y = (1+\varepsilon)\frac{a}{a+b}$. We obtain the inequality (66). $\qquad\square$

Remark 10. It is trivial to check
$\|W_{j:i}(0)\| = (n_i n_{i+1}\cdots n_j)^{1/2}, 1 \leq i \leq j \leq p,$
$\|W_{j:i}(0)\| = (n_{i-1}n_i\cdots n_{j-1})^{1/2}, p+1 \leq i \leq j \leq N,$
$\|W_{j:i}(0)\| \leq (n_i n_{i+1}\cdots n_{j-1})^{1/2}(n_p)^{1/2}$

$$\leq \left(\frac{n_{max}}{n_{min}}\right)^{1/2}(n_i n_{i+1}\cdots n_{j-1}\cdot\max\{n_{i-1}, n_j\})^{1/2}, 1 \leq i < p < j \leq N, (i, j) \neq (1, N).$$

Remark 11. As a special case, if $n_1 = n_2 = \cdots = n_{N-1} = n$, we know that $\|W_{j:i}(0)\| = (n_{i-1}n_i \cdots n_{N-1})^{1/2} = n^{(N-i+1)/2}$.

Lemma F.4. *Assume* $n_p/\min\{n_1, n_{N-1}\} \leq C_0 < \infty$. *Set* $\varepsilon > 0$. *Let* $C(\varepsilon)$ *represent the constant depend only on* ε. *If* $n_1/C_0 \geq C(\varepsilon)n_N$, *then with probability at least* $1 - e^{-\Omega(n_1/C_0)}$

$$\sigma_{max}(W_{N:i}(0)) \leq (1+\varepsilon)(n_{i-1}n_i \cdots n_{N-1})^{1/2}, 2 \leq i \leq p$$

$$\sigma_{min}(W_{N:i}(0)) \geq (1-\varepsilon)(n_{i-1}n_i \cdots n_{N-1})^{1/2}, 2 \leq i \leq p.$$

Similarly, if $n_{N-1}/C_0 \geq C(\varepsilon)rank(X)$, *then with probability at least* $1 - e^{-\Omega(n_{N-1}/C_0)}$

$$\sigma_{max}(W_{j:1}(0)|_{\mathcal{R}(X)}) \leq (1+\varepsilon)(n_1 n_2 \cdots n_j)^{1/2}, p+1 \leq j \leq N$$

$$\sigma_{min}(W_{j:1}(0)|_{\mathcal{R}(X)}) \geq (1-\varepsilon)(n_1 n_2 \cdots n_j)^{1/2}, p+1 \leq j \leq N.$$

Proof of Lemma F.4. Let $D = (n_{N-1}n_{N-2} \cdots n_p)^{-1/2}W_{N:p+1}^T(0)$ and $A_i = (n_p n_{p-1} \cdots n_i)^{-1/2}W_{p:i}^T(0)$. Assume $v \in S^{n_N-1}$. Easy to see that A_i is a product of random orthogonal projections and D is a random embedding.

Let $e_1 = (1, 0, 0, \cdots, 0)^T \in \mathbb{R}^{n_p}$. There exists orthogonal matrix T such that $TDv = e_1$, $\|e_1\|_2 = \|TDv\|_2 = \|v\|_2 = 1$.

Since random orthogonal projections are right invariant, we have

$$\mathbb{P}(\|A_iDv\|_2 \geq y) = \mathbb{E}\left[\mathbb{E}\left(I_{\{\|A_iT^Te_1\|_2 \geq y\}}\Big| D\right)\right] = \mathbb{E}\left[\mathbb{E}\left(I_{\{\|A_ie_1\|_2 \geq y\}}\Big| D\right)\right] = \mathbb{P}(\|A_ie_1\|_2 \geq y).$$

This proves that $\|A_iDv\|_2^2$ and $\|A_ie_1\|_2^2$ have the same distribution.

Claim: If $v \neq 0$, then $\|A_iDv\|_2^2 / \|v\|_2^2 = \left\|(n_i n_{i+1} \cdots n_p^2 \cdots n_{N-1})^{-1/2}W_{N:i}^Tv\right\|_2^2 / \|v\|_2^2$ follows beta distribution $B(n_{i-1}/2, (n_p - n_{i-1})/2)$.

Define $B_p = \|A_pe_1\|_2^2$, $B_i = \|A_ie_1\|_2^2 / \|A_{i+1}e_1\|_2^2$, $i = p-1, p-2, \cdots, 1$. Then $B_p \sim B(n_{p-1}/2, (n_p - n_{p-1})/2)$, $B_{p-1}|B_p \sim B(n_{p-2}/2, (n_{p-1} - n_{p-2})/2)$, $\cdots$, $B_i|(B_p, \cdots, B_{i+1}) \sim B(n_{i-1}/2, (n_i - n_{i-1})/2)$.

If $n_{i+1} = n_i$, we know that $B_i|(B_p, \cdots, B_{i+1}) = 1, a.s.$

If $B \sim B(a, b)$, then the moments are given by the following equations,

$$\mathbb{E}B = \frac{a}{a+b}, \text{ and } \mathbb{E}B^k = \frac{a}{a+b}\frac{a+1}{a+b+1} \cdots \frac{a+k-1}{a+b+k-1}. \tag{67}$$

By law of total expectation, we have

$$\mathbb{E}B_iB_{i+1} \cdots B_p = \frac{n_{i-1}}{n_i}\frac{n_i}{n_{i+1}} \cdots \frac{n_{p-1}}{n_p} = \frac{n_{i-1}}{n_p},$$

as well as

$$\mathbb{E}(B_iB_{i+1} \cdots B_p)^k = \frac{n_{i-1}/2}{n_p/2}\frac{n_{i-1}/2+1}{n_p/2+1} \cdots \frac{n_{i-1}/2+k-1}{n_p/2+k-1}.$$

Notice that all integer moments of $B_iB_{i+1} \cdots B_p$ match those of $B(n_{i-1}/2, (n_p - n_{i-1})/2)$. We can verify that beta distribution satisfies Carleman's condition, which implies that $B_iB_{i+1} \cdots B_p \sim B(n_{i-1}/2, (n_p - n_{i-1})/2)$.

Thus, $\|A_iDv\|_2^2 / \|v\|_2^2 \sim B(n_{i-1}/2, (n_p - n_{i-1})/2)$, which proves the claim.

With probability at least $1 - \exp\{-\Omega(n_1/C_0)\}$, we have

$$(1-\varepsilon)^2\frac{n_{i-1}}{n_p} \leq \|ADv\|_2^2 \leq (1+\varepsilon)^2\frac{n_{i-1}}{n_p}, \|v\|_2 = 1.$$

Using the $\phi-$net technique which has already been used to prove lemma E.3, we know that

$$\sigma_{min}(AD) \geq (1-\varepsilon)\left(\frac{n_{i-1}}{n_p}\right)^{1/2},$$

and

$$\sigma_{max}(AD) \leq (1+\varepsilon)\left(\frac{n_{i-1}}{n_p}\right)^{1/2},$$

with probability at least $1 - \exp\{n_N \ln(3/\phi(\varepsilon))\} \exp\{-\Omega(n_1/C_0)\} \geq 1 - \exp\{-\Omega(n_1/C_0)\}$, since $n_1/C_0 \geq C(\varepsilon)n_N$, for $2 \leq i \leq p$.

Hence, with probability at least $1 - e^{-\Omega(n_1/C_0)}$, we have

$$\sigma_{min}(W_{N:i}(0)) \geq (1-\varepsilon)(n_{i-1}\cdots n_{N-1})^{1/2},$$

and

$$\sigma_{max}(W_{N:i}(0)) \leq (1+\varepsilon)(n_{i-1}\cdots n_{N-1})^{1/2}.$$

The other part of the proof is similar to that of lemma E.4, so we omit it.

$\square$

Proof of Theorem B.2. Set $c > 0$, $c_1 = c/6, c_2 = c/3$. In lemma F.4, we can pick a $\varepsilon > 0$, such that $1 + \varepsilon \leq e^{c_1/2}$ and $1 - \varepsilon \geq e^{-c_2/2}$. Set $M = 2\sqrt{C_0}, \theta = 0, B_0 = B_\delta$, and $\eta = \frac{(1-\varepsilon)2n_N}{e^{2c}\beta N}$.

The requirement on size $\{n_1, n_2, \cdots, n_{N-1}, N\}$ in (17) make sure that the remark 10, lemma F.4, lemma 2.3, and lemma D.1 all hold.

Notice that even though we need the conclusions in lemma F.4 simultaneously hold for $2 \leq i \leq p$, $p + 1 \leq j \leq N$, it suffices to apply lemma F.4 over $i \in I$ and $j \in J$, such that $\{n_i; i \in I\}$ and $\{n_j; j \in I\}$ both have distinct values. Since $|I| \leq \underline{N}$ and $|J| \leq \underline{N}$, with probability at least

$$1 - 2\underline{N}e^{-\Omega(n_{min}/C_0)} - \delta/2 \geq 1 - \delta,$$

the one peak random orthogonal projections and embeddings initialization satisfies the initialization assumption (31) and the overparameterization assumption (32).

Under assumption $n_1 = n_2 = \cdots = n_{N-1}$, we can use remark 11 to replace lemma F.4. Thus, with probability at least $1 - \delta/2 \geq 1 - \delta$, (31) holds. Applying lemma 2.3 and D.1, we complete the proof.

$\square$

Proof of Theorem B.3. Let $W_N(0) = \sqrt{n}U_N[I_{n_y}, 0]V_N^T, \cdots, W_i(0) = \sqrt{n}U_iI_nV_i^T, 2 \leq i \leq N - 1$, and $W_1(0) = \sqrt{n}U_1[I_{n_x}, 0]^T V_1^T$. Now, we want to verify (31). By simply calculation, we have

$$\begin{cases} \sigma_{max}(W_{N:i+1}(0)) = \sigma_{min}(W_{N:i+1}(0)) = n^{(N-i)/2}, 1 \leq i \leq N - 1, \\ \sigma_{max}(W_{i-1:1}(0)|_{\mathcal{R}(X)}) = \sigma_{max}(W_{i-1:1}(0)|_{\mathcal{R}(X)}) = n^{(i-1)/2}, 2 \leq i \leq N, \\ \|W_{j:i}(0)\| = n^{(j-i+1)/2}, 1 < i \leq j < N. \end{cases} \quad (68)$$

Notice that for any $1 \leq p \leq m$

$$\|a_N W_{N:1}(0)x\|_2^2 = \frac{n}{n_N}\left\|U_N[I_{n_y}, 0]V_N^T U_N[I_{n_x}, 0]^T V_1^T x\right\|_2^2 = \frac{n}{n_N}\left\|U_N[I_{n_y}, 0]V_N^T x'\right\|_2^2,$$

where $x' = U_N[I_{n_x}, 0]^T V_1^T x$, $\|x\|_2 = \|x'\|_2$.

Since the distribution of $U_N[I_{n_y}, 0]V_N^T$ is right invariant under multiplying orthogonal matrices, we have

$$\left\|U_N[I_{n_y}, 0]V_N^T x'\right\|_2^2 / \|x\|_2^2 \sim B(\frac{n_y}{2}, \frac{n - n_y}{2}).$$

Thus,

$$\mathbb{E}\left[\|a_N W_{N:1}(0)x\|_2^2\right] = \|x\|_2^2.$$

Applying lemma 2.3, we have

$$L_0 - L(W_*) \leq \beta\left(\frac{2 \cdot rank(X)}{\delta} + \|W_*\|_X^2\right),$$

with probability at least $1 - \delta/2$.

Applying Lemma D.1 with $c > 0, c_1 = c/6, c_2 = c/3, \theta = 0$, we complete the proof. $\square$

Proof of Theorem 3.1. Theorem 3.1 is a special case of Theorem B.1 and Theorem B.2. Hence, we omit the proof. $\square$

Proof of Theorem 3.2. In Theorem B.1, B.2, and B.3, we proved that for given constant $c_1, c_2 > 0$ and $0 < \varepsilon, \delta/2 < 1/2$ as well as learning rate η, there exists constant $C = C(c_1, c_2)$ such that all three kinds of random initializations will fall into the convergence region with probability at least $1 - \delta$. Applying Lemma 2.3, we complete the proof.

$\square$

G TABLES

In this section, we provide some empirical evidence to support the argument in Section 4:**Why do bad saddles not affect GD for overparameterized deep linear neural networks?** Consider the following procedures for tables of $\frac{\|W_i(t) - W_i(0)\|_F}{\|W_i(0)\|_F}$:

a) We consider $X \in \mathbb{R}^{128 \times 1000}$, and $W_* \in \mathbb{R}^{10 \times 128}$ and set $Y = W_* X + \varepsilon$, where the entries in X and ε are drawn i.i.d. from $N(0, 1)$.

b) We consider the loss function $\frac{1}{2} \|a_N W_{N:1} X - Y\|_F^2$.

c) For the given deep linear networks, we apply orthogonal initialization, which are denoted as $W_j(0), 1 \le j \le N$.

d) We set the learning rate $\eta = \frac{n_N}{N \cdot \|X\|^2}$ for the deep linear neural networks.

e) We make the tables for $\frac{\|W_i(t) - W_i(0)\|_F}{\|W_i(0)\|_F}$.

Let $n_1 = n_2 = n_3 = 2000, N = 4$. Assume W_* are drawn i.i.d. from $N(0, 25)$. We obtain the following table:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
$t = 1$	0.05161	0.00826	0.00826	0.18464
$t = 2$	0.08779	0.01389	0.01389	0.31396
$t = 3$	0.11335	0.01781	0.01779	0.40435
$t = 4$	0.12109	0.01894	0.01889	0.42920
$t = 5$	0.12527	0.01956	0.01948	0.44282
$t = 6$	0.12611	0.01967	0.01958	0.44476
$t = 7$	0.12755	0.01988	0.01978	0.44955
$t = 8$	0.12745	0.01986	0.01975	0.44876
$t = 9$	0.12819	0.01997	0.01987	0.45136
$t = 10$	0.12793	0.01992	0.01982	0.45018

Let $n_1 = n_2 = 10000, N = 2$. Assume W_* are drawn i.i.d. from $N(0, 4)$. We obtain the following table:

	$i = 1$	$i = 2$	$i = 3$
$t = 1$	0.02708	0.00153	0.04844
$t = 2$	0.04319	0.00244	0.07727
$t = 3$	0.05296	0.00299	0.09474
$t = 4$	0.05888	0.00333	0.10533
$t = 5$	0.06248	0.00353	0.11176
$t = 6$	0.06468	0.00365	0.11569
$t = 7$	0.06603	0.00373	0.11811
$t = 8$	0.06688	0.00377	0.11962
$t = 9$	0.06741	0.00380	0.12057
$t = 10$	0.06775	0.00382	0.12117

Let $n_1 = n_2 = 4000, N = 2$. Assume W_* are drawn i.i.d. from $N(0, 1)$. We obtain the following table:

	$i = 1$	$i = 2$	$i = 3$
$t = 1$	0.01622	0.00290	0.05802
$t = 2$	0.02684	0.00480	0.09601
$t = 3$	0.03411	0.00609	0.12202
$t = 4$	0.03919	0.00700	0.14018
$t = 5$	0.04280	0.00764	0.15306
$t = 6$	0.04539	0.00810	0.16232
$t = 7$	0.04729	0.00844	0.16908
$t = 8$	0.04869	0.00868	0.17408
$t = 9$	0.04974	0.00887	0.17782
$t = 10$	0.05054	0.00901	0.18066

Let $n_1 = n_2 = 8000, N = 2$. Assume W_* are drawn i.i.d. from $N(0, 1)$. We obtain the following table:

	$i = 1$	$i = 2$	$i = 3$
$t = 1$	0.01173	0.00148	0.04195
$t = 2$	0.01944	0.00246	0.06955
$t = 3$	0.02470	0.00312	0.08838
$t = 4$	0.02838	0.00358	0.10151
$t = 5$	0.03098	0.00391	0.11083
$t = 6$	0.03287	0.00415	0.11758
$t = 7$	0.03426	0.00432	0.12253
$t = 8$	0.03530	0.00445	0.12624
$t = 9$	0.03608	0.00455	0.12904
$t = 10$	0.03668	0.00463	0.13118

Let $n_1 = n_2 = 12000, N = 2$. Assume W_* are drawn i.i.d. from $N(0, 1)$. We obtain the following table:

	$i = 1$	$i = 2$	$i = 3$
$t = 1$	0.00965	0.00099	0.03453
$t = 2$	0.01597	0.00164	0.05712
$t = 3$	0.02025	0.00208	0.07244
$t = 4$	0.02323	0.00239	0.08310
$t = 5$	0.02535	0.00261	0.09069
$t = 6$	0.02690	0.00277	0.09621
$t = 7$	0.02804	0.00289	0.10029
$t = 8$	0.02890	0.00297	0.10336
$t = 9$	0.02955	0.00304	0.10570
$t = 10$	0.03006	0.00309	0.10750

Let $n_1 = n_2 = 20000, N = 2$. Assume W_* are drawn i.i.d. from $N(0, 1)$. We obtain the following table:

	$i = 1$	$i = 2$	$i = 3$
$t = 1$	0.00713	0.00057	0.02551
$t = 2$	0.01181	0.00095	0.04225
$t = 3$	0.01499	0.00121	0.05362
$t = 4$	0.01720	0.00138	0.06154
$t = 5$	0.01878	0.00151	0.06720
$t = 6$	0.01994	0.00161	0.07132
$t = 7$	0.02079	0.00168	0.07438
$t = 8$	0.02144	0.00173	0.07668
$t = 9$	0.02193	0.00177	0.07844
$t = 10$	0.02231	0.00179	0.07981

H   FIGURES

In this section, we provide some empirical evidence to support the results in Section 4: **Numerical Experiments**. We will show how the trajectories of the non-convex deep linear neural networks are

related to a convex optimization problem for GD under different initialization schemes. Consider the following procedures for plots of the logarithm of loss as a function of number of iterations:

a) We choose $X \in \mathbb{R}^{128 \times 1000}$ and $W_* \in \mathbb{R}^{10 \times 128}$ and set $Y = W_* X + \varepsilon$, where the entries in X, W_* and ε are drawn i.i.d. from $N(0, 1)$.

b) We consider the loss function $\frac{1}{2} \|a_N W_{N:1} X - Y\|_F^2$.

c) For the given linear networks, we apply the Gaussian initialization and the one peak random orthogonal projections and embeddings initialization, which are denoted as $W_j(0), 1 \leq j \leq N$.

d) For the convex optimization problem (1), we set the initialization to be $W(0) = a_N W_N(0) \cdots W_1(0)$.

e) We set the learning rate $\eta = \frac{n_N}{N \cdot \|X\|^2}$ and $\eta_* = \frac{N}{n_N} \eta$ for the deep linear neural networks and the convex problem, respectively.

f) We draw the loss function through 25 iterations.

Figure 1: Plot of Loss as a function of number of iterations with $n_1 = n_2 = n_3 = 128$ (First), 200 (Second), 2000 (Third) for Gaussian initialization, respectively.

Figure 2: Plot of Loss as a function of number of iterations with $n_1 = n_2 = n_3 = 128$ (First), 200 (Second), 5000 (Third) for Orthogonal initialization, respectively.