Subspace Recovery from Heterogeneous Data with Non-isotropic Noise

Abstract

Recovering linear subspaces from data is a fundamental and important task in statistics and machine learning. Motivated by heterogeneity in Federated Learning settings, we study a basic formulation of this problem: the principal component analysis (PCA), with a focus on dealing with irregular noise. Our data come from n users with user i contributing data samples from a d-dimensional distribution with mean μ_i. Our goal is to recover the linear subspace shared by $\mu_1, \ldots, \mu_n$ using the data points from all users, where every data point from user i is formed by adding an independent mean-zero noise vector to μ_i. If we only have one data point from every user, subspace recovery is information-theoretically impossible when the covariance matrices of the noise vectors can be non-spherical, necessitating additional restrictive assumptions in previous work. We avoid these assumptions by leveraging at least two data points from each user, which allows us to design an efficiently-computable estimator under non-spherical and user-dependent noise. We prove an upper bound for the estimation error of our estimator in general scenarios where the number of data points and amount of noise can vary across users, and prove an information-theoretic error lower bound that not only matches the upper bound up to a constant factor, but also holds even for spherical Gaussian noise. This implies that our estimator does not introduce additional estimation error (up to a constant factor) due to irregularity in the noise. We show additional results for a linear regression problem in a similar setup.

1 Introduction

We study the problem of learning low-dimensional structure amongst data distributions, given multiple samples from each distribution. This problem arises naturally in settings such as federated learning, where we want to learn from data coming from a set of individuals, each of which has samples from their own distribution. These distributions however are related to each other, and in this work, we consider the setting when these distributions have means lying in a low-dimensional subspace. The goal is to learn this subspace, even when the distributions may have different (and potentially non-spherical) variances. This heterogeneity can manifest itself in practice as differing number of samples per user, or the variance differing across individuals, possibly depending on their mean. Recovery of the subspace containing the means can in turn help better estimate individual means. In other words, this can allow for learning good estimator for all individual means, by leveraging information from all the individuals.

The irregularity of the noise makes this task challenging even when we have sufficiently many individual distributions. For example, suppose we have n individuals and for every $i = 1, \ldots, n$, an unknown $\mu_i \in \mathbb{R}^d$. For simplicity, suppose that $\mu_1, \ldots, \mu_n$ are distributed independently as $N(0, \sigma^2 u u^{\mathsf{T}})$ for $\sigma \in \mathbb{R}_{\geq 0}$ and an unknown unit vector $u \in \mathbb{R}^d$. In this setting, our goal is to

recover the one-dimensional subspace, equivalently the vector u. For every i, we have a data point $x_i = \mu_i + z_i$ where $z_i \in \mathbb{R}^d$ is a mean-zero noise vector. If z_i is drawn independently from a spherical Gaussian $N(0, \alpha^2 I)$, we can recover the unknown subspace with arbitrary accuracy as n grows to infinity because $\frac{1}{n}\sum x_i x_i^\mathsf{T}$ concentrates to $\mathbb{E}[x_i x_i^\mathsf{T}] = \sigma^2 u u^\mathsf{T} + \alpha^2 I$, whose top eigenvector is $\pm u$. However, if the noise z_i is drawn from a non-spherical distribution, the top eigenvector of $\frac{1}{n}\sum x_i x_i^\mathsf{T}$ can deviate from $\pm u$ significantly, and to make things worse, if the noise z_i is drawn independently from a non-spherical Gaussian $N(0, \sigma^2(I - uu^\mathsf{T}) + \alpha^2 I)$, then our data points $x_i = \mu_i + z_i$ distribute independently as $N(0, (\sigma^2 + \alpha^2)I)$, giving no information about the vector u.

The information-theoretic impossibility in this example however disappears as soon as one has at least two samples from each distribution. Indeed, given two data points $x_{i1} = \mu_i + z_{i1}$ and $x_{i2} = \mu_i + z_{i2}$ from user i, as long as the noise z_{i1}, z_{i2} are independent and have zero mean, we always have $\mathbb{E}[x_{i1} x_{i2}^\mathsf{T}] = \sigma^2 u u^\mathsf{T}$ regardless of the specific distributions of z_{i1} and z_{i2}. This allows us to recover the subspace in this example, as long as we have sufficiently many users each contributing at least two examples.

As this is commonly the case in our motivating examples, we make this assumption of multiple data points per user, and show that this intuition extends well beyond this particular example. We design efficiently computable estimators for this subspace recovery problem given samples from multiple heteroscedastic distributions (see Section 1.1 for details). We prove upper bounds on the error of our estimator measured in the *maximum principal angle* (see Section 2 for definition). We also prove an information-theoretic error lower bound, showing that our estimator achieves the optimal error up to a constant factor in general scenarios where the number of data points and the amount of noise can vary across users. Somewhat surprisingly, our lower bound holds even when the noise distributes as spherical Gaussians. Thus non-spherical noise in setting does not lead to increased error.

We then show that our techniques extend beyond the mean estimation problem to a linear regression setting where for each μ_i, we get (at least two) samples $(x_{ij}, x_{ij}^\mathsf{T}\mu_i + z_{ij})$ where z_{ij} is zero-mean noise from some noise distribution that depends on i and x_{ij}. This turns out to be a model that was recently studied in the meta-learning literature under more restrictive assumptions (e.g. z_{ij} is independent of x_{ij}) [Kong et al., 2020, Tripuraneni et al., 2021, Collins et al., 2021, Thekumparampil et al., 2021]. We show a simple estimator achieving an error upper bound matching the ones in prior work without making these restrictive assumptions.

1.1 Our contributions

PCA with heterogeneous and non-isotropic noise: Upper Bounds. In the PCA setting, the data points from each user i are drawn from a user-specific distribution with mean $\mu_i \in \mathbb{R}^d$, and we assume that $\mu_1, \ldots, \mu_n$ lie in a shared k-dimensional subspace that we want to recover. Specifically, we have m_i data points $x_{ij} \in \mathbb{R}^d$ from user i for $j = 1, \ldots, m_i$, and each data point is determined by $x_{ij} = \mu_i + z_{ij}$ where $z_{ij} \in \mathbb{R}^d$ is a noise vector drawn independently from a mean zero distribution. We use $\eta_i \in \mathbb{R}_{\geq 0}$ to quantify the amount of noise in user i's data points by assuming that z_{ij} is an η_i-*sub-Gaussian* random variable.

As mentioned earlier, if we only have a single data point from each user, it is information-theoretically impossible to recover the subspace. Thus, we focus on the case where $m_i \geq 2$ for every $i = 1, \ldots, n$. In this setting, for appropriate weights $w_1, \ldots, w_n \in \mathbb{R}_{\geq 0}$, we compute a matrix A:

$$A = \sum_{i=1}^n \frac{w_i}{m_i(m_i - 1)} \sum_{j_1 \neq j_2} x_{ij_1} x_{ij_2}^\mathsf{T}, \tag{1}$$

where the inner summation is over all pairs $j_1, j_2 \in \{1, \ldots, m_i\}$ satisfying $j_1 \neq j_2$. Our estimator is then defined by the subspace spanned by the top-k eigenvectors of A.

The flexibility in the weights $w_1, \ldots, w_n$ allows us to deal with variations in m_i and η_i for different users i. In the special case where $\eta_1 = \cdots = \eta_n = \eta$ and $m_1 = \cdots = m_n = m$, we choose $w_1 = \cdots = w_n = 1/n$ and we show that our estimator achieves the following error upper bound with success probability at least $1 - \delta$:

$$\sin\theta = O\left(\left(\frac{\eta\sigma_1}{\sigma_k^2\sqrt{m}} + \frac{\eta^2}{\sigma_k^2 m}\right)\sqrt{\frac{d + \log(1/\delta)}{n}}\right).$$

Here, θ is the maximum principal angle between our estimator and the true subspace shared by $\mu_1, \dots, \mu_n$, and σ_ℓ^2 is the ℓ-th largest eigenvalue of $\frac{1}{n} \sum_{i=1}^n \mu_i \mu_i^\mathsf{T}$. Our error upper bound for general m_i, η_i, w_i is given in Theorem 3.1.

We instantiate our error upper bound to the case where $\mu_1, \dots, \mu_n$ are drawn iid from a Gaussian distribution $N(0, \sigma^2 U U^\mathsf{T})$, where the columns of $U \in \mathbb{R}^{d \times k}$ form an orthonormal basis of the subspace containing $\mu_1, \dots, \mu_n$. By choosing the weights $w_1, \dots, w_n$ according to $m_1, \dots, m_n$ and $\eta_1, \dots, \eta_n$, our estimator achieves the error upper bound

$$\sin \theta \le O\left(\sqrt{\frac{d + \log(1/\delta)}{\sum_{i=1}^n \gamma_i'}}\right) \tag{2}$$

under a mild assumption (Assumption 3.2), where γ_i' is defined in Definition 3.1 and often equals $\left(\frac{\eta_i^2}{\sigma^2 m_i} + \frac{\eta_i^4}{\sigma^4 m_i^2}\right)^{-1}$.

PCA: Lower Bounds. We show that the error upper bound (2) is optimal up to a constant factor by proving a matching information-theoretic lower bound (Theorem 3.7). Our lower bound holds for general m_i and η_i that can vary among users i, and it holds even when the noise vectors z_{ij} are drawn from spherical Gaussians, showing that our estimator essentially pays no additional cost in error or sample complexity due to non-isotropic noise.

We prove the lower bound using Fano's method on a local packing over the Grassmannian manifold. We carefully select a non-trivial hard distribution so that the strength of our lower bound is not affected by a group of fewer than k users each having a huge amount of data points with little noise.

Linear Models. While the PCA setting is the main focus of our paper, we extend our research to a related linear models setting that has recently been well studied in the meta-learning and federated learning literature [Kong et al., 2020, Tripuraneni et al., 2021, Collins et al., 2021, Thekumparampil et al., 2021]. Here, the user-specific distribution of each user i is parameterized by $\beta_i \in \mathbb{R}^d$, and we again assume that $\beta_1, \dots, \beta_n$ lie in a linear subspace that we want to recover. From each user i we observe m_i data points $(x_{ij}, y_{ij}) \in \mathbb{R}^d \times \mathbb{R}$ for $j = 1, \dots, m_i$ drawn from the user-specific distribution satisfying $y_{ij} = x_{ij}^\mathsf{T} \beta_i + z_{ij}$ for an $O(1)$-sub-Gaussian measurement vector $x_{ij} \in \mathbb{R}^d$ with zero mean and identity covariance and an η_i-sub-Gaussian mean-zero noise term $z_{ij} \in \mathbb{R}$. While it may seem that non-isotropic noise is less of a challenge in this setting since each noise term z_{ij} is a scalar, our goal is to handle a challenging scenario where the variances of the noise terms z_{ij} can depend on the *realized* measurements x_{ij}, which is a more general and widely applicable setting compared to those in prior work. Similarly to the PCA setting, our relaxed assumptions on the noise make it information-theoretically impossible to do subspace recovery if we only have one data point from each user (see Section 4), and thus we assume each user contributes at least two data points. We use the subspace spanned by the top-k eigenvectors of the following matrix A as our estimator:

$$A = \sum_{i=1}^n \frac{w_i}{m_i(m_i - 1)} \sum_{j_1 \ne j_2} (x_{ij_1} y_{ij_1})(x_{ij_2} y_{ij_2})^\mathsf{T}. \tag{3}$$

In the special case where $\eta_1 = \dots = \eta_n = \eta, m_1 = \dots = m_n = m$, and $\|\beta_i\|_2 \le r$ for all i, our estimator achieves the following error upper bound:

$$\sin \theta \le O\left(\log^3(nd/\delta)\sqrt{\frac{d(r^4 + r^2 \eta^2 + \eta^4/m)}{mn\sigma_k^4}}\right), \tag{4}$$

where σ_k^2 is the k-th largest eigenvalue of $\frac{1}{n} \sum_{i=1}^n \beta_i \beta_i^\mathsf{T}$ (Corollary L.2). Our error upper bound extends smoothly to more general cases where η_i and m_i vary among users (Theorem L.1). Moreover, our upper bound matches the ones in prior work [e.g. Tripuraneni et al., 2021, Theorem 3] despite requiring less restrictive assumptions.

1.2 Related Work

Principal component analysis under non-isotropic noise has been studied by Vaswani and Narayanamurthy [2017], Zhang et al. [2018] and Narayanamurthy and Vaswani [2020]. When translated to

our setting, these papers focus on having only one data point from each user and thus they require additional assumptions—either the level of non-isotropy is low, or the noise is coordinate-wise independent and the subspace is incoherent. The estimation error guarantees in these papers depend crucially on how well these additional assumptions are satisfied. Zhu et al. [2019] and Cai et al. [2021] study PCA with noise and missing data, and Chen et al. [2021] and Cheng et al. [2021] study eigenvalue and eigenvector estimation under heteroscedastic noise. These four papers all assume that the noise is coordinate-wise independent and the subspace/eigenspace is incoherent.

The linear models setting we consider has recently been studied as a basic setting of meta-learning and federated learning by Kong et al. [2020], Tripuraneni et al. [2021], Collins et al. [2021], and Thekumparampil et al. [2021]. These papers all make the assumption that the noise terms z_{ij} are independent of the measurements x_{ij}, an assumption that we relax in this paper. Collins et al. [2021] and Thekumparampil et al. [2021] make improvements in sample complexity and error guarantees compared to earlier work by Kong et al. [2020] and Tripuraneni et al. [2021], but Collins et al. [2021] focus on the noiseless setting ($z_{ij} = 0$) and Thekumparampil et al. [2021] require at least $\Omega(k^2)$ examples per user. Tripuraneni et al. [2021] and Thekumparampil et al. [2021] assume that the measurements x_{ij} are drawn from the standard (multivariate) Gaussian distribution, where as Kong et al. [2020], Collins et al. [2021] and our work make the relaxed assumption that x_{ij} are sub-Gaussian with identity covariance, which, in particular, allows the fourth-order moments of x_{ij} to be non-isotropic. There is a large body of prior work on meta-learning beyond the linear setting [see e.g. Maurer et al., 2016, Tripuraneni et al., 2020, Du et al., 2020].

When collecting data from users, it is often important to ensure that private information about users is not revealed through the release of the learned estimator. Many recent works proposed and analyzed estimators that achieve *user-level differential privacy* in settings including mean estimation [Levy et al., 2021, Esfandiari et al., 2021], meta-learning [Jain et al., 2021] and PAC learning [Ghazi et al., 2021]. Recently, Cummings et al. [2021] study one-dimensional mean estimation in a setting similar to ours, under a differential privacy constraint.

1.3 Paper Organization

In Section 2, we formally define the maximum principal angle and other notions we use throughout the paper. Our results in the PCA setting and the linear models setting are presented in Sections 3 and 4, respectively. We defer most technical proofs to the appendices.

2 Preliminaries

We use $\|A\|$ to denote the spectral norm of a matrix A, and use $\|u\|_2$ to denote the ℓ_2 norm of a vector u. For positive integers $k \le d$, we use $\mathbb{O}_{d,k}$ to denote the set of matrices $A \in \mathbb{R}^{d \times k}$ satisfying $A^\mathsf{T} A = I_k$, where I_k is the $k \times k$ identity matrix. We use $\mathbb{O}_d$ to denote $\mathbb{O}_{d,d}$, which is the set of $d \times d$ orthogonal matrices. We use $\mathrm{col}(A)$ to denote the linear subspace spanned by the columns of a matrix A. We use the base-e logarithm throughout the paper.

Maximum Principal Angle. Let $U, \hat{U} \in \mathbb{O}_d$ be two orthogonal matrices. Suppose the columns of U and $\hat{U}$ are partitioned as $U = [U_1 \ U_2], \hat{U} = [\hat{U}_1 \ \hat{U}_2]$ where $U_1, \hat{U}_1 \in \mathbb{O}_{d,k}$ for an integer k satisfying $0 < k < d$. Let Γ (resp. $\hat{\Gamma}$) be the k-dimensional linear subspace spanned by the columns of U_1 (resp. $\hat{U}_1$). The *maximum principal angle* $\theta \in [0, \pi/2]$ between Γ and $\hat{\Gamma}$, denoted by $\angle(\Gamma, \hat{\Gamma})$ or $\angle(U_1, \hat{U}_1)$, is defined by $\sin\theta = \|U_1 U_1^\mathsf{T} - \hat{U}_1 \hat{U}_1^\mathsf{T}\| = \|U_1^\mathsf{T} \hat{U}_2\| = \|U_2^\mathsf{T} \hat{U}_1\|$. It is not hard to see that the maximum principal angle depend only on the subspaces $\Gamma, \hat{\Gamma}$ and not on the choices of U and $\hat{U}$, and $\sin\angle(\Gamma, \hat{\Gamma})$ is a natural metric between k-dimensional subspaces (see Appendix A for more details where we define principal angles for any two subspaces with possibly different dimensions).

With the definition of the maximum principal angle, we can now state a variant of the Davis–Kahan $\sin\theta$ theorem [Davis and Kahan, 1970] that will be useful in our analysis (see Appendix E for proof):

Theorem 2.1 (Variant of Davis–Kahan $\sin\theta$ theorem). *Let $A, \hat{A} \in \mathbb{R}^{d \times d}$ be symmetric matrices. Let λ_i denote the i-th largest eigenvalue of A. For a positive integer k smaller than d, let θ denote the maximum principal angle between the subspaces spanned by the top-k eigenvectors of A and $\hat{A}$.*

Assuming $\lambda_k > \lambda_{k+1}$,

$$\sin\theta \le \frac{2\|A - \hat{A}\|}{\lambda_k - \lambda_{k+1}}.$$

Sub-Gaussian and sub-exponential distributions. We say a random variable $x \in \mathbb{R}$ with expectation $\mathbb{E}[x] \in \mathbb{R}$ has *sub-Gaussian* constant $b \in \mathbb{R}_{\ge 0}$ if $\mathbb{E}[|x - \mathbb{E}[x]|^p]^{1/p} \le b\sqrt{p}$ for every $p \ge 1$. We say x has *sub-exponential* constant $b \in \mathbb{R}_{\ge 0}$ if $\mathbb{E}[|x - \mathbb{E}[x]|^p]^{1/p} \le bp$ for every $p \ge 1$. We say a random vector $y \in \mathbb{R}^d$ has sub-Gaussian (resp. sub-exponential) constant $b \in \mathbb{R}_{\ge 0}$ if for every unit vector $u \in \mathbb{R}^d$ (i.e., $\|u\|_2 = 1$), the random variable $u^\mathsf{T} y \in \mathbb{R}$ has sub-Gaussian (resp. sub-exponential) constant b. We say y is b-*sub-Gaussian* (resp. b-*sub-exponential*) if it has sub-Gaussian (resp. sub-exponential) constant b.

3 Principal Component Analysis

In the principal component analysis (PCA) setting, our goal is to recover the k-dimensional subspace Γ spanned by the user-specific means $\mu_1, \ldots, \mu_n \in \mathbb{R}^d$ of the n users. From each user i, we have $m_i \ge 2$ data points

$$x_{ij} = \mu_i + z_{ij} \quad \text{for } j = 1, \ldots, m_i.$$

We assume the noise $z_{ij} \in \mathbb{R}^d$ is drawn independently from a mean zero distribution with sub-Gaussian constant η_i. We do *not* assume that the variance of z_{ij} is the same along every direction, *nor* do we assume that the distribution of z_{ij} is the same for different (i, j). We first show an error upper bound for our estimator when the user-specific means $\mu_1, \ldots, \mu_n$ are deterministic vectors (Section 3.1) and then apply this result to the case where $\mu_1, \ldots, \mu_n$ are drawn from a sub-Gaussian distribution (Section 3.2). In Section 3.3 we prove an information-theoretic error lower bound matching our upper bound.

3.1 Fixed User-Specific Means

We first focus on the case where $\mu_1, \ldots, \mu_n$ are deterministic vectors. In this case, all the randomness in the data comes from the noise z_{ij}. Our estimator is the subspace $\hat{\Gamma}$ spanned by the top-k eigenvectors of A defined in (1). For $\ell = 1, \ldots, d$, we define σ_ℓ^2 to be the ℓ-th largest eigenvalue of $\sum_{i=1}^n w_i \mu_i \mu_i^\mathsf{T}$. Since $\mu_1, \ldots, \mu_n$ share a k-dimensional subspace, $\sigma_\ell = 0$ for $\ell > k$. We prove the following general theorem on the error guarantee of our estimator:

Theorem 3.1. *Define $\xi^2 = \|\sum_{i=1}^n w_i^2 \mu_i \mu_i^\mathsf{T} \eta_i^2 / m_i\|$ and let θ denote the maximum principal angle between our estimator $\hat{\Gamma}$ and the true subspace Γ spanned by $\mu_1, \ldots, \mu_n$. For any $\delta \in (0, 1/2)$, with probability at least $1 - \delta$,*

$$\sin\theta = O\left(\sigma_k^{-2}\sqrt{(d + \log(1/\delta))\left(\xi^2 + \sum_{i=1}^n \frac{w_i^2 \eta_i^4}{m_i^2}\right)} + \sigma_k^{-2}(d + \log(1/\delta))\max_i \frac{w_i \eta_i^2}{m_i}\right). \quad (5)$$

We can simplify the bound in Theorem 3.1 by considering special cases:

Corollary 3.2. *Assume $\max\{\eta_1/\sqrt{m_1}, \ldots, \eta_n/\sqrt{m_n}\} = t$ and we choose $w_1 = \cdots = w_n = 1/n$. For any $\delta \in (0, 1/2)$, with probability at least $1 - \delta$,*

$$\sin\theta = O\left(\frac{t\sigma_1 + t^2}{\sigma_k^2}\sqrt{\frac{d + \log(1/\delta)}{n}}\right). \quad (6)$$

In particular, when $\eta_1 = \cdots = \eta_n = \eta$, and $m_1 = \cdots = m_n = m$, error bound (6) becomes

$$\sin\theta = O\left(\left(\frac{\eta\sigma_1}{\sigma_k^2\sqrt{m}} + \frac{\eta^2}{\sigma_k^2 m}\right)\sqrt{\frac{d + \log(1/\delta)}{n}}\right).$$

We defer the complete proof of Theorem 3.1 and Corollary 3.2 to Appendices F and G. Our proof is based on the Davis-Kahan $\sin\theta$ theorem (Theorem 2.1). Since $\sigma_{k+1}^2 = 0$, Theorem 2.1 implies

$$\sin\theta \le \frac{2\|A - \sum_{i=1}^n w_i \mu_i \mu_i^\mathsf{T}\|}{\sigma_k^2}. \quad (7)$$

This reduces our goal to proving an upper bound on the spectral norm of $A - \sum_{i=1}^{n} w_i \mu_i \mu_i^\mathsf{T}$. Since for distinct j_1 and j_2 in $\{1, \ldots, m_i\}$ we have $\mathbb{E}[x_{ij_1} x_{ij_2}^\mathsf{T}] = \mu_i \mu_i^\mathsf{T}$, our construction of A in (1) guarantees $\mathbb{E}[A] = \sum_{i=1}^{n} w_i \mu_i \mu_i^\mathsf{T}$. Therefore, our goal becomes controlling the deviation of A from its expectation, and we achieve this goal using techniques for matrix concentration inequalities.

3.2 Sub-Gaussian User-Specific Means

We apply our error upper bound in Theorem 3.1 to the case where $\mu_1, \ldots, \mu_n$ are drawn iid from $N(0, \sigma^2 U U^\mathsf{T})$ for an unknown $U \in \mathbb{O}_{d,k}$. We use the same estimator $\hat{\Gamma}$ as before: $\hat{\Gamma}$ is the subspace spanned by the top-k eigenvectors of A defined in (1). We determine the optimal weights $w_1, \ldots, w_n$ in (1) as long as $m_1, \ldots, m_n$ and $\eta_1, \ldots, \eta_n$ satisfy a mild assumption (Assumption 3.2), achieving an error upper bound in Theorem 3.4. In the next subsection, we prove an error lower bound (Theorem 3.7) that matches our upper bound (Theorem 3.4) up to a constant factor, assuming $d \geq (1 + \Omega(1))k$ and $\delta = \Theta(1)$.

We prove our error upper bound in a slightly more general setting than $\mu_1, \ldots, \mu_n$ drawn iid from $N(0, \sigma^2 U U^\mathsf{T})$. Specifically, we make the following assumption on the distribution of $\mu_1, \ldots, \mu_n$:

Assumption 3.1. *The user-specific means $\mu_1, \ldots, \mu_n \in \mathbb{R}^d$ are mean-zero independent random vectors supported on an unknown k-dimensional subspace Γ. Moreover, for a parameter $\sigma > 0$, for every $i = 1, \ldots, n$, μ_i has sub-Gaussian constant $O(\sigma)$, and the k-th largest eigenvalue of $\mathbb{E}[\mu_i \mu_i^\mathsf{T}]$ is at least σ^2.*

Under this assumption, we have the following lower bound on the σ_k^2 in Theorem 3.1 (see Appendix H for proof):

Claim 3.3. *Under Assumption 3.1, let $w_1, \ldots, w_n \in \mathbb{R}_{\geq 0}$ be user weights satisfying $w_1 + \cdots + w_n = 1$ and σ_k^2 be the k-th largest eigenvalue of $\sum_{i=1}^{n} w_i \mu_i \mu_i^\mathsf{T}$. There exists an absolute constant $C_* > 1$ such that for any $\delta \in (0, 1/2)$, as long as $\max_{1 \leq i \leq n} w_i \leq 1/C_*(k + \log(1/\delta))$, then $\sigma_k^2 \geq \sigma^2/2$ with probability at least $1 - \delta/2$.*

The following definition is important for us to choose the weights $w_1, \ldots, w_n$ in (1) optimally:

Definition 3.1. *Define $\gamma_i = \left(\frac{\eta_i^2}{\sigma^2 m_i} + \frac{\eta_i^4}{\sigma^4 m_i^2} \right)^{-1}$ and assume w.l.o.g. that $\gamma_1 \geq \cdots \geq \gamma_n$. Define $\gamma_i' = \gamma_i$ if $i \geq k$, and $\gamma_i' = \gamma_k$ if $i < k$.*

We make the following mild assumption on γ_i' under which we achieve optimal estimation error:

Assumption 3.2. $\sum_{i=1}^{n} \gamma_i' \geq C_*(k + \log(1/\delta)) \gamma_1'$ *for C_* defined in Claim 3.3.*

By the definition of γ_i', it is easy to show that Assumption 3.2 is equivalent to $\sum_{i=k+1}^{n} \gamma_i \geq ((C_* - 1)k + C_* \log(1/\delta)) \gamma_k$. Therefore, Assumption 3.2 holds as long as γ_k is not too large relative to $\gamma_{k+1} + \cdots + \gamma_n$. For example, when $\gamma_k = \cdots = \gamma_n$, Assumption 3.2 holds as long as $n \geq C_*(k + \log(1/\delta))$. Also, since $\gamma_1' = \cdots = \gamma_k' \geq \gamma_{k+1}' \geq \cdots \geq \gamma_n' \geq 0$, it trivially holds that $\sum_{i=1}^{n} \gamma_i' \geq k \gamma_1'$. Assumption 3.2 is relatively mild when compared to this trivial inequality.

Under Assumption 3.2, we show that it is optimal to choose the weights $w_1, \ldots, w_n$ as

$$w_i = \frac{\gamma_i'}{\sum_{\ell=1}^{n} \gamma_\ell'}. \tag{8}$$

Specifically, if we plug (8) into Theorem 3.1 and bound ξ and σ_k based on the distribution of $\mu_1, \ldots, \mu_n$, we get the following error upper bound which matches our lower bound (Theorem 3.7) in Section 3.3. We defer its proof to Appendix I.

Theorem 3.4. *Under Assumptions 3.1 and 3.2, if we choose $w_1, \ldots, w_n$ as in (8) and define $\theta = \angle(\Gamma, \hat{\Gamma})$, for $\delta \in (0, 1/2)$, with probability at least $1 - \delta$,*

$$\sin \theta \leq O \left(\sqrt{\frac{d + \log(1/\delta)}{\sum_{i=1}^{n} \gamma_i'}} \right). \tag{9}$$

For comparison, consider the setting when $\sigma = \eta_i = 1$. The result then says that $\sin \theta$ is bounded by approximately $\sqrt{\frac{d}{\sum_i m_i}}$. This is the same rate as we would get if we have $\sum_i m_i$ independent

samples with homogeneous spherical noise. Thus as long as the samples are not too concentrated on fewer than k users, the heterogeneity comes at no additional cost.

3.3 Lower Bound

We prove a lower bound matching the upper bound in Theorem 3.4 up to constant in the setting where $\delta = \Theta(1), d \geq (1 + \Omega(1))k$.

For every positive integer d, there is a natural "uniform" distribution over $\mathbb{O}_d$ given by Haar's theorem. We denote this distribution by $\mathsf{Haar}(\mathbb{O}_d)$. A random matrix A drawn from $\mathsf{Haar}(\mathbb{O}_d)$ has the following invariance property: for any deterministic matrix $B \in \mathbb{O}_d$, the random matrices A, AB and BA all have the same distribution. For an integer $k \leq d$, we can construct a random matrix $A_1 \in \mathbb{O}_{d,k}$ by first drawing $A \in \mathbb{R}^{d \times d}$ from $\mathsf{Haar}(\mathbb{O}_d)$ and then take the first k columns of A. We denote the distribution of A_1 by $\mathsf{Haar}(\mathbb{O}_{d,k})$. The invariance property of $\mathsf{Haar}(\mathbb{O}_d)$ immediately implies the following claims:

Claim 3.5. *Let $A \in \mathbb{O}_d$ be a random matrix drawn from $\mathsf{Haar}(\mathbb{O}_d)$ and let $B \in \mathbb{O}_{d,k}$ be a fixed matrix. Then AB distributes as $\mathsf{Haar}(\mathbb{O}_{d,k})$.*

Proof. The matrix B can be written as the first k columns of a matrix $C \in \mathbb{O}_d$. Now AB is the first k columns of AC, where AC distributes as $\mathsf{Haar}(\mathbb{O}_d)$ by the invariance property. This implies that AB distributes as $\mathsf{Haar}(\mathbb{O}_{d,k})$. □

Claim 3.6. *Let $B \in \mathbb{O}_{d,k}$ be a random matrix. Assume for every fixed matrix $A \in \mathbb{O}_d$, the random matrices B and AB have the same distribution. Then $B \sim \mathsf{Haar}(\mathbb{O}_{d,k})$.*

Proof. If we draw A independently from $\mathsf{Haar}(\mathbb{O}_d)$, the random matrices B and AB still have the same distribution. By Claim 3.5, AB distributes as $\mathsf{Haar}(\mathbb{O}_{d,k})$, so B must also distribute as $\mathsf{Haar}(\mathbb{O}_{d,k})$. □

With the definition of $\mathsf{Haar}(\mathbb{O}_{d,k})$, we state our lower bound in the following theorem:

Theorem 3.7. *Let k, d, n be positive integers satisfying $k < d$ and $k \leq n$. Let $m_1, \ldots, m_n$ be positive integers and $\sigma, \eta_1, \ldots, \eta_n$ be positive real numbers. Suppose we draw $U \in \mathbb{O}_{d,k}$ from $\mathsf{Haar}(\mathbb{O}_{d,k})$ and then draw $\mu_1, \ldots, \mu_n$ independently from $N(0, \sigma^2 UU^\mathsf{T})$. For every $i = 1, \ldots, n$, we draw m_i data points x_{ij} for $j = 1, \ldots, m_i$ as $x_{ij} = \mu_i + z_{ij}$, where each z_{ij} is drawn independently from the spherical Gaussian $N(0, \eta_i^2 I)$. Let $\hat{\Gamma}$ be any estimator mapping $(x_{ij})_{1 \leq i \leq n, 1 \leq j \leq m_i}$ to a (possibly randomized) k-dimensional subspace of $\mathbb{R}^d$. Let θ denote the maximum principal angle between $\hat{\Gamma}((x_{ij})_{1 \leq i \leq n, 1 \leq j \leq m_i})$ and the true subspace $\Gamma = \mathsf{col}(U)$. If real numbers $t \geq 0$ and $\delta \in [0, 1/2)$ satisfy $\Pr[\sin \theta \leq t] \geq 1 - \delta$, then*

$$t \geq \Omega\left(\min\left\{1, \sqrt{\frac{(d-k)(1-\delta)}{\sum_{i=k}^n \gamma_i}}\right\}\right), \tag{10}$$

where $\gamma_1, \ldots, \gamma_n$ are defined in Definition 3.1.

Note that $\gamma_i' = \gamma_i$ for $i \geq k$, so our upper bound in (9) matches the lower bound (10) up to a constant factor assuming $\delta = \Theta(1)$ and $d \geq (1 + \Omega(1))k$.

We use the local Fano method to prove the lower bound using the technical lemmas in Appendix D. In particular, we reduce our goal to proving an upper bound on the KL divergence between Gaussian distributions whose covariance matrices are defined based on matrices $U, \hat{U} \in \mathbb{O}_{d,k}$ with $\|UU^\mathsf{T} - \hat{U}\hat{U}^\mathsf{T}\|_F$ bounded. We prove the following lemma in Appendix J that upper bounds the KL divergence using $\|UU^\mathsf{T} - \hat{U}\hat{U}^\mathsf{T}\|_F$:

Lemma 3.8. *For $\sigma \in \mathbb{R}_{\geq 0}, \eta \in \mathbb{R}_{>0}, U, \hat{U} \in \mathbb{O}_{d,k}$, define $\Sigma = \sigma^2 UU^\mathsf{T} + \eta^2 I$ and $\hat{\Sigma} = \sigma^2 \hat{U}\hat{U}^\mathsf{T} + \eta^2 I$. Then,*

$$D_{\mathrm{kl}}(N(0, \hat{\Sigma}) \| N(0, \Sigma)) = \frac{\sigma^4 \|UU^\mathsf{T} - \hat{U}\hat{U}^\mathsf{T}\|_F^2}{4(\sigma^2 \eta^2 + \eta^4)}.$$

Lemma 3.8 and the results in Appendix D allow us to prove a version of (10) in which the sum in the demoninator is over $i = 1, \ldots, n$. This, however, is weaker and less useful than (10) in which the sum in the denominator is over $i = k, k+1, \ldots, n$. To prove Theorem 3.7, we extract a hard distribution in which the data points from users $1, \ldots, k-1$ are "useless" in terms of subspace recovery.

Let Γ_1 be the $(k-1)$-dimensional subspace spanned by $\mu_1, \ldots, \mu_{k-1}$. We let $v_1, \ldots, v_{k-1}$ be a random orthonormal basis of Γ_1, and we append another vector $v_k \in \Gamma$ to form an orthonormal basis $v_1, \ldots, v_k$ of Γ. We define $V_1 = [v_1 \cdots v_{k-1}] \in \mathbb{O}_{d,k-1}$ and $V = [v_1 \cdots v_k] \in \mathbb{O}_{d,k}$. In Figure 1 we show a graphical model demonstrating the dependency among the random objects we defined.

$$U \longrightarrow (\mu_1, \ldots, \mu_{k-1}) \longrightarrow \Gamma_1 \longrightarrow V_1 \searrow$$
$$U \xrightarrow{\hspace{3cm}} \Gamma \longrightarrow V$$
$$U \searrow (\mu_k, \ldots, \mu_n)$$

Figure 1: Graphical model A.

Let us focus on the joint distribution of $(V_1, V, (\mu_1, \ldots, \mu_{k-1}))$. By the invariance property, for any matrices $\tilde{V}_1 \in \mathbb{O}_{d,k-1}, \tilde{V} \in \mathbb{O}_{d,k}$, measurable set $S \subseteq (\mathbb{R}^d)^{k-1}$, and $G \in \mathbb{O}_d$,

$$\Pr[(\mu_1, \ldots, \mu_{k-1}) \in S | V = \tilde{V}, V_1 = \tilde{V}_1] = \Pr[(\mu_1, \ldots, \mu_{k-1}) \in S_G | V = G\tilde{V}, V_1 = G\tilde{V}_1],$$

where $S_G = \{(G\mu_1, \ldots, G\mu_{k-1}) : (\mu_1, \ldots, \mu_{k-1}) \in S\}$. For any $\tilde{V}, \tilde{V}' \in \mathbb{O}_{d,k}$ whose first $k-1$ columns are both $\tilde{V}_1$, there exists $G \in \mathbb{O}_d$ such that $\tilde{V}' = G\tilde{V}$ and thus $\tilde{V}_1 = G\tilde{V}_1$. This implies that for any $\mu \in \text{col}(V_1) = \Gamma_1$, we have $G\mu = \mu$, and thus $(S \cap \Gamma_1)_G = S \cap \Gamma_1$ for any measurable $S \subseteq (\mathbb{R}^d)^{k-1}$. Therefore,

$$\Pr[(\mu_1, \ldots, \mu_{k-1}) \in S | V = \tilde{V}, V_1 = \tilde{V}_1]$$
$$= \Pr[(\mu_1, \ldots, \mu_{k-1}) \in S \cap \Gamma_1 | V = \tilde{V}, V_1 = \tilde{V}_1]$$
$$= \Pr[(\mu_1, \ldots, \mu_{k-1}) \in (S \cap \Gamma_1)_G | V = G\tilde{V}, V_1 = G\tilde{V}_1]$$
$$= \Pr[(\mu_1, \ldots, \mu_{k-1}) \in S \cap \Gamma_1 | V = \tilde{V}', V_1 = \tilde{V}_1]$$
$$= \Pr[(\mu_1, \ldots, \mu_{k-1}) \in S | V = \tilde{V}', V_1 = \tilde{V}_1].$$

This implies that $(\mu_1, \ldots, \mu_{k-1})$ and V are conditionally independent given V_1. Therefore, the joint distribution of $(V_1, V, (\mu_1, \ldots, \mu_{k-1}))$ can be formed by first drawing V and V_1, and then drawing $\mu_1, \ldots, \mu_{k-1}$ based only on V_1 and not on V. Since $\mu_k, \ldots, \mu_n$ are drawn iid from $N(0, \sigma^2 UU^\mathsf{T}) = N(0, \sigma^2 VV^\mathsf{T})$, we have the graphical model shown in Figure 2.

$$V \longrightarrow V_1 \longrightarrow (\mu_1, \ldots, \mu_{k-1})$$
$$V \searrow (\mu_k, \ldots, \mu_n)$$

Figure 2: Graphical model B.

By Claim 3.6, the marginal distribution of V is $\mathsf{Haar}(\mathbb{O}_{d,k})$. By Claim 3.5, we can implement this distribution by first drawing $W \sim \mathsf{Haar}(\mathbb{O}_d)$ and then drawing E independently from any distribution over $\mathbb{O}_{d,k}$ and let $V = WE$. We choose the distribution of E later, where we ensure that the first $k-1$ columns of E is always $\begin{bmatrix} I_{k-1} \\ 0 \end{bmatrix}$. This guarantees that the first $k-1$ columns of W and V are the same, and thus V_1 is exactly the first $k-1$ columns of W, resulting in the graphical model shown in Figure 3.

Note that in Figure 3 there is no directed path from E to $(\mu_1, \ldots, \mu_{k-1})$. Intuitively, this means that knowing $(\mu_1, \ldots, \mu_{k-1})$ gives us no information about E. Now by choosing the distribution of E appropriately, we can prove (10) in which the denominator does not contain $\gamma_1, \ldots, \gamma_{k-1}$. We defer the complete proof of Theorem 3.7 to Appendix K.

$$W \longrightarrow V_1 \longrightarrow (\mu_1, \ldots, \mu_{k-1})$$
$$E \longrightarrow V \longrightarrow (\mu_k, \ldots, \mu_n)$$

Figure 3: Graphical model C.

4   Linear Models

In the linear models setting, the data distribution of user i is parameterized by an unknown vector $\beta_i \in \mathbb{R}^d$. As before, we assume that the vectors $\beta_1, \ldots, \beta_n$ from the n users lie in an unknown k-dimensional subspace Γ. Our goal is to recover the subspace using the following data. For every $i = 1, \ldots, n$, we have m_i data points from user i: $(x_{i1}, y_{i1}), \ldots, (x_{im_i}, y_{im_i}) \in \mathbb{R}^d \times \mathbb{R}$. For every $j = 1, \ldots, m_i$, we assume the *measurement* $x_{ij} \in \mathbb{R}^d$ is a random vector drawn independently from an $O(1)$-sub-Gaussian distribution with zero mean and identity covariance matrix. The measurement *outcome* y_{ij} is determined by $y_{ij} = x_{ij}^\mathsf{T} \beta_i + z_{ij}$, where the random *noise* $z_{ij} \in \mathbb{R}$ can depend on the measurements $x_{i1}, \ldots, x_{im_i}$. When conditioned on $x_{i1}, \ldots, x_{im_i}$, we assume every z_{ij} for $j = 1, \ldots, m_i$ is independently drawn from an η_i-sub-Gaussian distribution with zero mean, but we do *not* assume that the conditional distribution of z_{ij} is the same for every $j = 1, \ldots, m_i$. The (in)dependence among x_{ij} and z_{ij} for $i = 1, \ldots, n$ and $j = 1, \ldots, m_i$ can be summarized by the example graphical model in Figure 4.

Figure 4: An example for $n = 3, m_1 = 2, m_2 = 3, m_3 = 2$.

Since we allow the noise z_{ij} to depend on the measurements x_{ij}, it is information-theoretically impossible to recover the subspace if we only have one data point from every user. Consider the scenario where every β_i is drawn independently from $N(0, \sigma^2 uu^\mathsf{T})$ for an unknown unit vector $u \in \mathbb{R}^d$ and every x_{ij} is drawn independently and uniformly from $\{-1, 1\}^d$. If we set z_{ij} to be $z_{ij} = x_{ij}^\mathsf{T} \nu_{ij}$ where ν_{ij} is independently drawn from $N(0, \sigma^2(I - uu^\mathsf{T}))$, then every y_{ij} satisfies $y_{ij} = x_{ij}^\mathsf{T}(\beta_i + \nu_{ij})$ where $\beta_i + \nu_{ij}$ distributes as $N(0, \sigma^2 I)$ independently from x_{ij}. This implies that the joint distribution of $((x_{i1}, y_{i1}))_{i=1,\ldots,n}$ does not change with u, i.e., we get no information about u from one data point per user.

Thus, we assume $m_i \geq 2$ for every user i. In this case, we achieve error upper bounds that match the ones in [Tripuraneni et al., 2021] despite our relaxed assumptions on the noise. Our estimator is the subspace $\hat{\Gamma}$ spanned by the top-k eigenvectors of A defined in (3). We defer the analysis of our estimator to Appendix L.

References

Changxiao Cai, Gen Li, Yuejie Chi, H. Vincent Poor, and Yuxin Chen. Subspace estimation from unbalanced and incomplete data matrices: $\ell_{2,\infty}$ statistical guarantees. *Ann. Statist.*, 49(2):944–967, 2021. ISSN 0090-5364. doi: 10.1214/20-aos1986. URL https://doi.org/10.1214/20-aos1986.

T. Tony Cai, Zongming Ma, and Yihong Wu. Sparse PCA: optimal rates and adaptive estimation. *Ann. Statist.*, 41(6):3074–3110, 2013. ISSN 0090-5364. doi: 10.1214/13-AOS1178. URL https://doi.org/10.1214/13-AOS1178.

Yuxin Chen, Chen Cheng, and Jianqing Fan. Asymmetry helps: eigenvalue and eigenvector analyses of asymmetrically perturbed low-rank matrices. *Ann. Statist.*, 49(1):435–458, 2021. ISSN 0090-5364. doi: 10.1214/20-AOS1963. URL https://doi.org/10.1214/20-AOS1963.

Chen Cheng, Yuting Wei, and Yuxin Chen. Tackling small eigen-gaps: fine-grained eigenvector estimation and inference under heteroscedastic noise. *IEEE Trans. Inform. Theory*, 67(11):7380–7419, 2021. ISSN 0018-9448. doi: 10.1109/TIT.2021.3111828. URL https://doi.org/10.1109/TIT.2021.3111828.

Liam Collins, Hamed Hassani, Aryan Mokhtari, and Sanjay Shakkottai. Exploiting shared representations for personalized federated learning. In *International Conference on Machine Learning*, pages 2089–2099. PMLR, 2021.

Rachel Cummings, Vitaly Feldman, Audra McMillan, and Kunal Talwar. Mean estimation with user-level privacy under data heterogeneity, 2021. TPDP.

Chandler Davis and William Morton Kahan. The rotation of eigenvectors by a perturbation. iii. *SIAM Journal on Numerical Analysis*, 7(1):1–46, 1970.

Simon Shaolei Du, Wei Hu, Sham M Kakade, Jason D Lee, and Qi Lei. Few-shot learning via learning the representation, provably. In *International Conference on Learning Representations*, 2020.

Hossein Esfandiari, Vahab Mirrokni, and Shyam Narayanan. Tight and robust private mean estimation with few users. *arXiv preprint arXiv:2110.11876*, 2021.

Badih Ghazi, Ravi Kumar, and Pasin Manurangsi. User-level private learning via correlated sampling. *Advances in Neural Information Processing Systems*, 34, 2021.

Prateek Jain, John Rush, Adam Smith, Shuang Song, and Abhradeep Guha Thakurta. Differentially private model personalization. *Advances in Neural Information Processing Systems*, 34, 2021.

Weihao Kong, Raghav Somani, Zhao Song, Sham Kakade, and Sewoong Oh. Meta-learning for mixed linear regression. In *International Conference on Machine Learning*, pages 5394–5404. PMLR, 2020.

Daniel Levy, Ziteng Sun, Kareem Amin, Satyen Kale, Alex Kulesza, Mehryar Mohri, and Ananda Theertha Suresh. Learning with user-level privacy. *Advances in Neural Information Processing Systems*, 34, 2021.

Andreas Maurer, Massimiliano Pontil, and Bernardino Romera-Paredes. The benefit of multitask representation learning. *J. Mach. Learn. Res.*, 17:Paper No. 81, 32, 2016. ISSN 1532-4435.

Praneeth Narayanamurthy and Namrata Vaswani. Fast robust subspace tracking via pca in sparse data-dependent noise. *IEEE Journal on Selected Areas in Information Theory*, 1(3):723–744, 2020.

Kiran Koshy Thekumparampil, Prateek Jain, Praneeth Netrapalli, and Sewoong Oh. Sample efficient linear meta-learning by alternating minimization. *arXiv preprint arXiv:2105.08306*, 2021.

Nilesh Tripuraneni, Michael Jordan, and Chi Jin. On the theory of transfer learning: The importance of task diversity. *Advances in Neural Information Processing Systems*, 33:7852–7862, 2020.

Nilesh Tripuraneni, Chi Jin, and Michael Jordan. Provable meta-learning of linear representations. In *International Conference on Machine Learning*, pages 10434–10443. PMLR, 2021.

Namrata Vaswani and Praneeth Narayanamurthy. Finite sample guarantees for PCA in non-isotropic and data-dependent noise. In *2017 55th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 783–789. IEEE, 2017.

Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.

Anru R Zhang, T Tony Cai, and Yihong Wu. Heteroskedastic PCA: Algorithm, optimality, and applications. *arXiv preprint arXiv:1810.08316*, 2018.

Ziwei Zhu, Tengyao Wang, and Richard J Samworth. High-dimensional principal component analysis with heterogeneous missingness. *arXiv preprint arXiv:1906.12125*, 2019.