Causal Bandits: Online Decision-Making in Endogenous Settings

Abstract

The deployment of Multi-Armed Bandits (MAB) has become commonplace in
many economic applications. However, regret guarantees for even state-of-the-art
linear bandit algorithms (such as Optimism in the Face of Uncertainty Linear
bandit (OFUL)) make strong exogeneity assumptions w.r.t. arm covariates. This
assumption is very often violated in many economic contexts and using such
algorithms can lead to sub-optimal decisions. In this paper, we consider the
problem of online learning in linear stochastic multi-armed bandit problems with
endogenous covariates. We propose an algorithm we term *BanditIV*, that uses
instrumental variables to correct for this bias, and prove an $\tilde{\mathcal{O}}(k\sqrt{T})$ upper bound
for the expected regret of the algorithm. Further, in economic contexts, it is also
important to understand how the model parameters behave asymptotically. To this
end, we additionally propose ϵ-*BanditIV* algorithm and demonstrate its asymptotic
consistency and normality while ensuring the same regret bound. Finally, we carry
out extensive Monte Carlo simulations to demonstrate the performance of our
algorithms compared to other methods. We show that *BanditIV* and ϵ-*BanditIV*
significantly outperform other existing methods.

1 Introduction

The proliferation of user-level data presents at the same time a unique opportunity and challenge in
front of decision-makers. Decision-makers want to use individual-level data to tailor their decisions
for each user. Further, given the dynamic nature of the platform, decision-makers want to be able
to adopt these decisions with incrementally available data. Woodroofe 1979 first proposed a simple
model to solve such sequential decision-making problems with covariates. Langford and T. Zhang
2007 later named the model "contextual bandit". Contextual linear bandits have been adopted across
a wide variety of applications from advertising (Tang et al. 2015, Aramayo, Schiappacasse, and
Goic 2022) to healthcare (Durand et al. 2018), dialogue systems (Liu et al. 2018), and personalized
product recommendations (L. Li et al. 2010, Qin, S. Chen, and Zhu 2014). In the setting of a
contextual linear bandit problem, in each round, the decision-maker observes a set of actions with
each action characterized by a set of features. Decision-maker selects an action and observes a reward
corresponding to that action. The objective of the decision-maker is to achieve cumulative reward
close to that of optimal policy in hindsight.

Traditional formulations of contextual bandits make the unconfoundedness assumption i.e., the arm
covariates are exogenous. However, in many economic settings, arm features can be correlated with
the unobserved noise. For instance, consider the problem of generating product recommendations
for consumers. Platform operators would usually run online experiments to uncover the relationship
between product features and consumer demand. Generally in such settings, product observable

features like price are controlled by product owners (different from the platform operator) which could be set in anticipation of consumer demand and hence be correlated with the demand shocks unobserved by the platform operator. In such settings, traditional bandit algorithms might lead to sub-optimal decisions. A common approach to correct the endogeneity bias in the offline setting is to use instrumental variables. Instruments are correlated with the endogenous variable but are otherwise not associated with the outcome variable. The method of instrumental variables uses the variation in the exogenous component of the endogenous variable induced by the variation in the instrumental variable to make inference of causal effects. Thus, to address the issues induced by endogenous features in online settings, we propose an online estimator we term *BanditIV* that uses instrumental variables to learn the relationship between rewards and features.

Next, literature in contextual linear bandit has also primarily focused on minimizing the expected regret of the algorithm (Auer 2002, Dani, Hayes, and Kakade 2008, Chu et al. 2011, and Abbasi-Yadkori, Pál, and Szepesvári 2011). However, in a variety of practical situations, one may also be interested in the inference of model parameters, and algorithms that minimize regret may not guarantee the consistency of the model parameter estimator. Further, knowing the asymptotic distributions, one can easily test the significance of features, and offers a traditional way to select variables. For instance, in the product recommendation example, an online platform might also be interested in understanding the effect of various marketing-mix variables (like price, and promotion) on consumer demand. To this end, we propose a linear bandit algorithm that pursues both regret minimization and consistent estimation of model parameters in endogenous settings. We use the martingale central limit theorem to show that our estimator of model parameters is asymptotically normal.

Finally, we compare our method with existing online algorithms including Optimism in the Face of Uncertainty Linear bandit (OFUL) and Thompson Sampling (TS) through numerical experiments. We find significant advantages of our algorithm in both expected regret and parameter inference.

To summarize, our paper makes the following contributions –

- We study the stochastic linear bandit problem with endogenous features. We propose a new algorithm, we term *BanditIV*, which incorporates instrumental variables to correct for the bias induced by endogenous features and show that the total expected regret of the algorithm is upper bounded by $\tilde{\mathcal{O}}(k\sqrt{T})$.

- Next, as in many economic contexts, researchers might not only be interested in minimizing the regret over the outcomes but also in conducting inference over estimated parameters. To this end, we propose the ϵ-*BanditIV* algorithm with the same upper bound and establish the asymptotic consistency and normality of the estimator.

- Finally, we conduct numerical experiments of *BanditIV* and ϵ-*BanditIV* algorithms on synthetic data. We show *BanditIV* and ϵ-*BanditIV* always outperform other existing linear bandit algorithms including OFUL and TS in terms of both total expected regret and inference.

To see a complete version of literature review, please refer Appendix A.

Notation. Throughout this paper, we use $||\cdot||_p$ to denote the p-norm of a vector or a matrix. We use $||\cdot||_F$ to denote the Frobenius norm for a matrix. For a vector x and positive definite matrix A, we denote $\sqrt{x'Ax}$ as $||x||_A$. We use $\langle\cdot,\cdot\rangle$ as the inner product. We denote the sequence $a_0, a_1, \cdots, a_\infty$ as $\{a_t\}_{t=0}^\infty$. For a matrix X, the i^{th} column is denoted as $X^{(i)}$ and the j^{th} element of the i^{th} column is denoted as $X^{(i)(j)}$.

2 Problem setting

Let T be the number of rounds and K the number of arms in each round. In each round t, the learner observes K feature vectors $x_{t,a} \in \mathbb{R}^d$, $a \in \{1, \cdots, K\}$, with $||x_{t,a}||_2 \leq L_x$ for samples (one feature vector for each sample), where the subscript a represents the a^{th} arm or sample from the K-sample set. Let $y_{t,a} \in \mathbb{R}$, s.t. $|y_{t,a}| \leq L_y$, be the reward of arm a on round t. After observing the feature vectors, the learner selects an arm a_t and receives reward $y_{t,a}$. Under linear realizability assumption,

86 we have $\mathbb{E}[y_{t,a}|x_{t,a}] = x'_{t,a}\beta_0$ for all t and a, where $\beta_0 \in \mathbb{R}^d$ is an unknown true coefficient vector.
87 We assume $y_{t,a}$ are independent random variables with expectation $x'_{t,a}\beta_0$, as the following.

$$y_{t,a} = \beta'_0 x_{t,a} + e_{t,a} \tag{1}$$

88 where $e_{t,a} \in \mathbb{R}$ is a 1-subgaussian error term, s.t. $\mathbb{E}[e_{t,a}] = 0$. Standard Contextual Linear Bandit
89 settings have $\mathbb{E}[e_{t,a}x_{t,a}] = 0$, which assumes that the error term is independent with the feature
90 vector. However, this assumption can be violated in cases such as when we have omitted variable
91 in the error term, measurement error in the regressor, simultaneous equation estimation, and etc.
92 Mathematically, when we have the following holds,

$$\mathbb{E}[e_{t,a}x_{t,a}] \neq 0$$

93 the endogeneity problem occurs. Ordinary Least Square (OLS) estimator is inconsistent in en-
94 dogenous settings. Econometrics literature uses Instrumental Variable (IV) method to address the
95 endogeneity problem. A variable $z_{t,a}$ is a valid instrumental variable if it satisfies following three
96 conditions (M Wooldridge 2014): (i) it is uncorrelated with the error term $e_{t,a}$, i.e. $\mathbb{E}[z_{t,a}e_{t,a}] = 0$,
97 (ii) it is correlated with the endogenous covariates $x_{t,a}$, (iii) it has no direct effect on the reward $y_{t,a}$.
98 Consider a valid instrumental variable $z_{t,a} \in \mathbb{R}^k$, with $||z_{t,a}||_2 \leq L_z$, and the minimum eigenvalue
99 of $\mathbb{E}[z_{t,a}z'_{t,a}]$ is positive. We assume $\mathbb{E}[z_{t,a}x'_{t,a}] \in \mathbb{R}^{k \times d}$ has full column rank d.[1]

$$x_{t,a} = \Gamma'_0 z_{t,a} + u_{t,a} \tag{2}$$

100 where $\Gamma_0 \in \mathbb{R}^{k \times d}$ is an unknown true coefficient vector, $u_{t,a} \in \mathbb{R}^d$ is a 1-subgaussian error term
101 with mean zero and the independency condition $\mathbb{E}[z_{t,a}u'_{t,a}] = 0$ is satisfied by construction. By
102 plugging the equation of $x_{t,a}$ (Equation 2) into Equation 1, we have $y_{t,a} = (\Gamma_0\beta_0)'z_{t,a} + v_{t,a}$ where
103 $v_{t,a} = \beta'_0 u_{t,a} + e_{t,a}$. We denote $\Gamma_0\beta_0$ as δ_0.

104 Recall that in offline settings, we can use instrumental variables to solve endogeneity problem
105 through methods like Two-Stage Least-Squares (TSLS). Consider vector $Y \in \mathbb{R}^T$ with $y_{t,a}$ as its
106 elements, matrix $X \in \mathbb{R}^{T \times d}$ with $x'_{t,a}$ as its rows, matrix $Z \in \mathbb{R}^{T \times k}$ with $z'_{t,a}$ as its rows, where
107 $t \in \{1, \cdots, T\}$. We briefly illustrate TSLS estimation procedure in offline settings as the following:
108 (i) First, regress the set of feature vectors X on the set of instrumental variable vectors Z using OLS
109 method to obtain an estimated sample features $\hat{X}$; (ii) Then regress Y on the estimated feature vectors
110 $\hat{X}$ to obtain the estimator $\hat{\beta}_{\hat{X},Y}$ using OLS method again.

111 **Definition 1.** (Two-Stage Least Squares Estimator)

$$\hat{X} = P_Z X, P_Z = Z(Z'Z)^{-1}Z'$$
$$\hat{\beta}_{\hat{X},Y} = (\hat{X}'\hat{X})^{-1}\hat{X}'Y$$

112 To facilitate further analysis on the TSLS estimator, we also define two related OLS estimators,
113 $\hat{\Gamma} = (Z'Z)^{-1}Z'X$ and $\hat{\delta} = (Z'Z)^{-1}Z'Y$ where $\hat{\delta} = \hat{\Gamma}\hat{\beta}_{\hat{X},Y}$ (to see this equation, refer Appendix B
114 for the proof).

115 It's known that TSLS estimator is consistent in offline settings, while it remains unknown for the
116 consistency of the TSLS estimator in online settings. As far as we know, we are the first who combine
117 the contextual linear bandit algorithm with TSLS and prove the consistency of TSLS estimator in
118 online settings. We provide the asymptotic properties in Section 4.

119 We aim to design an online decision making algorithm that learns the coefficient of main interest β_0
120 so that we can maximize the total expected reward after pulling arms. We define the total expected
121 regret of an algorithm $\mathcal{A}$ after T rounds as

$$R_T = \sum_{t=1}^{T} \mathbb{E}[y_{t,a_t^*} - y_{t,a_t}] = \sum_{t=1}^{T} \mathbb{E}[\langle \Gamma'_0 z_{t,a_t^*}, \beta_0 \rangle - \langle \Gamma'_0 z_{t,a_t}, \beta_0 \rangle]$$

122 where $a_t^* = \arg\max_a \langle \Gamma'_0 z_{t,a}, \beta_0 \rangle$ is the best arm at round t according to the true coefficient vectors
123 β_0 and Γ_0, and a_t is the arm selected by the algorithm $\mathcal{A}$ at round t.

124 We relist key assumptions in this paper as the following,

---

[1]The assumption $\mathbb{E}[z_{t,a}x'_{t,a}] \in \mathbb{R}^{k \times d}$ has full column rank d implies that $k \geq d$

125   *Assumption* 1. $||z_{t,a}||_2 \leq L_z$, $||x_{t,a}||_2 \leq L_x$, $||y_{t,a}||_2 \leq L_y$ for all t and a.

126   *Assumption* 2. The minimum eigenvalue of $\mathbb{E}[z_{t,a}z'_{t,a}]$ is larger than a positive constant λ.

127   For the simplicity of notations, we omit the subscript a of $z_{t,a}$, $x_{t,a}$, and $y_{t,a}$ from below. By x_t, we
128   mean one sample or arm chosen from the arm set $\mathcal{X}_t$ at time t; z_t is the instrumental variable related
129   to x_t; y_t is the reward generated from the arm x_t.

130 3   BanditIV algorithm

131   To address the endogeneity problem in online settings, we propose the following *BanditIV* Algorithm
132   based on existing linear bandit algorithms, especially the OFUL algorithm proposed by Abbasi-
133   Yadkori, Pál, and Szepesvári 2011.

134   The *BanditIV* algorithm[2] takes as input initial regularization parameters $\gamma_z, \gamma_x > 0$, as well as
135   confidence interval parameters $\{G_t\}_{t=1}^{\infty}$, $\{B_t\}_{t=1}^{\infty}$. We construct two confidence sets $C_{1,s}$, $C_{2,s}$,
136   $s \in \{1, \cdots, T\}$ for the first stage estimation and the second stage estimation respectively in each
137   round. The idea for the confidence sets are to make the estimation optimistic with the condition
138   that "with high probability" the true coefficients are in the confidence sets (Abbasi-Yadkori, Pál, and
139   Szepesvári 2011). The best arm generated by the algorithm is a sample x_t which is related to an
140   instrumental variable z_t that can maximize the estimated reward jointly with a pair of optimistic
141   estimates of two-stage coefficients in the confidence sets. If the current time t is after the first round,
142   we utilize past choices X_t and observations of rewards Y_t to estimate the coefficients Γ_0 and β_0 by
143   the TSLS estimator in each round. We denote the estimator $\hat{\beta}_t$ used in this algorithm as the *BanditIV*
144   *estimator*. Based on the estimation and confidence sets, we choose the best arm and then observe the
145   reward.

146   Taking the *BanditIV* algorithm as a special case, we further propose the ϵ-*BanditIV* algorithm (which
147   we show in Algorithm 1). The ϵ-*BanditIV* algorithm additionally takes a sequence of non-increasing
148   exploration parameters $\{\epsilon_t\}_{t=1}^{\infty}$ as input. In each round, instead of choosing the best arm, we conduct
149   a stochastic decision with probability ϵ_t for choosing a random arm and probability $1 - \epsilon_t$ for choosing
150   the estimated best arm. Notice that when $\epsilon_t = 0$ for $t \in \{1, 2 \cdots, \infty\}$, the ϵ-*BanditIV* algorithm
151   degenerates to the *BanditIV*. For the simplicity of notations, we omit the subscript a of $z_{t,a}$, $x_{t,a}$, and
152   $y_{t,a}$ from below. By x_t, we mean one sample or action chosen from the action set $\mathcal{X}_t$ at time t; z_t is
153   the IV related with x_t; y_t is the reward generated from the action x_t.

Algorithm 1 ϵ-BanditIV

Input: $\gamma_z, \gamma_x > 0$, $\{G_t\}_{t=1}^{\infty}, \{B_t\}_{t=1}^{\infty}, \{\epsilon_t\}_{t=1}^{\infty}$
Set $U_0 = \gamma_z I \in \mathbb{R}^{k \times k}, V_0 = 0 \in \mathbb{R}^{k \times d}, W_0 = \gamma_x I \in \mathbb{R}^{d \times d}, Q_0 = 0 \in \mathbb{R}^{d \times 1}$
Set $C_{1,s} = \{\Gamma : ||\Gamma^{(i)} - \hat{\Gamma}_s^{(i)}||_{U_s} \leq G_s\}, C_{2,s} = \{\beta : ||\beta - \hat{\beta}_s||_{W_s} \leq B_s\}, \forall s \in \{1, 2, \cdots T\},$
$\forall i \in \{1, 2, \cdots d\}$
Nature reveals $\mathcal{Z}_0$. We randomly choose $z_0 \in \mathcal{Z}_0$ and set $Z_0 = z_0$
Nature reveals $\mathcal{X}_0$. From the set $\mathcal{X}_0$, we play x_0 which is related with z_0 and then observe the reward y_0. We
set $X_0 = x_0$ and $Y_0 = y_0$
for $t := 1, 2, \cdots, T,$ **do**
    $U_t = U_{t-1} + z_{t-1}z'_{t-1}, V_t = V_{t-1} + z_{t-1}x'_{t-1}$
    $\hat{\Gamma}_t = U_t^{-1}V_t, \hat{X}_{t-1} = Z_{t-1}\hat{\Gamma}_t$
    $W_t = W_0 + \hat{X}'_{t-1}\hat{X}_{t-1}, Q_t = Q_0 + \hat{X}'_{t-1}Y_{t-1}$
    $\hat{\beta}_t = W_t^{-1}Q_t$
    Nature reveals $\mathcal{Z}_t$. With probability ϵ_t, we uniformly choose a random $z_t \in \mathcal{Z}_t$. With probability $1 - \epsilon_t$,
we choose $z_t = \arg\max_{z \in \mathcal{Z}_t} \max_{\Gamma \in C_{1,t}} \max_{\beta \in C_{2,t}} \langle \Gamma'z, \beta \rangle$ and update $Z_t = [Z'_{t-1} \quad z_t]'$
    Nature reveals the set of arms $\mathcal{X}_t$. We play x_t which is related to z_t and observe the reward y_t. Update
$X_t = [X'_{t-1} \quad x_t]', Y_t = [Y'_{t-1} \quad y_t]'$
end for

---

           [2]Please refer Appendix A to see more details for the algorithm.

3.1 Regret analysis

In this section, we give upper bounds on the regret of the *BanditIV* algorithm and ϵ-*BanditIV* algorithm. The proofs can be found in Appendix A. We show an $\tilde{\mathcal{O}}(k\sqrt{T})$ upper bound for the total expected regret of both algorithms with parameters of confidence set by Theorem 1.

Theorem 1. *The expected cumulative regret of the ϵ-BanditIV at time T, with probability at least $1 - \delta$, is upper-bounded by*

$$R_T \le B_T \sqrt{2Td \log(\frac{T+d}{d})} + (\frac{2}{\gamma_x} + \|\beta_0\|_2) G_T \sqrt{2Tk \log(\frac{T+k}{k})} + 2\epsilon_0 T L_y$$

where $B_T = \sqrt{\gamma_x}\|\beta_0\|_2 + \sqrt{2\log(\frac{2T}{\delta}) + d\log\left(\frac{5TL_x^2}{d}\right)}$,

$G_T = \sqrt{\gamma_z}\|\Gamma_0^{(i)}\|_2 + \sqrt{2\log(\frac{2T}{\delta}) + k\log\left(\frac{5TL_y^2}{k}\right)}.$

By setting $\epsilon_t = 0$, we can derive the upper bound for the *BanditIV* directly.

Remark 1. Notice that in order to guarantee an $\tilde{\mathcal{O}}(K\sqrt{T})$ upper bound for the total expected regret of the ϵ-BanditIV algorithm, we need a small enough ϵ_t, such as an $\epsilon_t \le \frac{\sqrt{\log(t)}}{\sqrt{t}}$. The *BanditIV* algorithm which is a special case with $\epsilon_t = 0$ in ϵ-BanditIV algorithm, naturally satisfies this condition.

4 Inference: tail inequality and normality

In this section, we show asymptotic properties of the *BanditIV* estimator following H. Chen, Lu, and Song 2021 and Bastani and Bayati 2020. We first present the consistency and then the normality of the *BanditIV* estimator. The proofs are presented in Appendix A. Additionally, we calculate confidence intervals and show the results in A due to limited space.

Proposition 1. *(Tail bound for the BanditIV estimator) In the online decision-making model with ϵ-greedy policy, if the Assumptions 1 and 2 are satisfied, and ϵ_t is non-increasing, then for any $\eta_1, \eta_2 > 0$, $\eta_2 \ne \frac{\lambda_{min}(\Gamma_0)}{(kd)^{\frac{1}{2}}}$,*

$$P(\|\hat{\beta}_t - \beta_0\|_1 \le C_\beta) \ge p_1 p_2^d$$

where $C_\beta = \dfrac{\eta_1 + \eta_2 \|\beta_0\|_1}{\lambda_{min}(\Gamma_0) d^{\frac{-1}{2}} - \eta_2 (k)^{\frac{1}{2}}}$,

$$p_1 = 1 - \exp\{-\frac{t\epsilon_t}{8}\} - k\exp\{-\frac{t\epsilon_t \lambda}{32L_z^2}\} - 2k\exp\{-\frac{t\epsilon_t^2 \lambda^2 \eta_1^2}{128k^2\sigma_v^2 L_z^2}\} + 2k^2 \exp\{-\frac{t\epsilon_t^2 \lambda^2 \eta_1^2 + 4t\epsilon_t \lambda k^2 \sigma_v^2}{128k^2\sigma_v^2 L_z^2}\},$$

$$p_2 = 1 - \exp\{-\frac{t\epsilon_t}{8}\} - k\exp\{-\frac{t\epsilon_t \lambda}{32L_z^2}\} - 2k\exp\{-\frac{t\epsilon_t^2 \lambda^2 \eta_1^2}{128k^2\sigma_u^2 L_z^2}\} + 2k^2 \exp\{-\frac{t\epsilon_t^2 \lambda^2 \eta_1^2 + 4t\epsilon_t \lambda k^2 \sigma_v^2}{128k^2\sigma_u^2 L_z^2}\}$$

Remark 2. If $t\epsilon_t^2 \to \infty$ as $t \to \infty$, then the probability of $\|\hat{\beta}_t - \beta_0\| \le C_\beta$ goes to 1 for any $\eta_1, \eta_2 > 0$, $\eta_2 \ne \frac{\lambda_{min}(\Gamma_0)}{(kd)^{\frac{1}{2}}}$. We need carefully choose the value for ϵ_t in order to guarantee both the regret bound and the tail bound. For example, an $\epsilon_t = \frac{\sqrt{\log(t)}}{\sqrt{t}}$ or $\epsilon_t = \frac{\sqrt{\log\log(t)}}{\sqrt{t}}$ satisfies both the conditions for the regret bound and the tail bound.

Following from the Propositions 1, we obtain the consistency of the BanditIV estimator easily.

Corollary 2. *(Consistency of the online BanditIV estimator). If Assumptions 1 and 2 are satisfied, ϵ_t is non-increasing and $t\epsilon_t^2 \to \infty$ as $t \to \infty$, then the online BanditIV estimator $\hat{\beta}_t$ is a consistent estimator for β_0.*

Theorem 3. *(Aymptotic normality of the online BanditIV estimator) If Assumptions 1 and 2 are satisfied, ϵ_t is non-increasing and $t\epsilon_t^2 \to \infty$ as $t \to \infty$, the non-increasing function $\epsilon_t \to \epsilon_\infty$ and $t\epsilon_t^2 \to \infty$ as $t \to \infty$. Then*

$$\sqrt{t}(\hat{\beta}_t - \beta_0) \xrightarrow{d} \mathcal{N}_d(0, \mathbb{E}[v_t^2](\Gamma_0'\Gamma_0)^{-1}\Gamma_0'(\int zz'd\mathcal{P}_z)^{-1}\Gamma_0(\Gamma_0'\Gamma_0)^{-1})$$

Theorem 3 provides a theoretical guarantee that the *BanditIV* estimator asymptotically follows a normal distribution with mean zero and variance $\mathbb{E}[v_t^2](\Gamma_0'\Gamma_0)^{-1}\Gamma_0'(\int zz'd\mathcal{P}_z)^{-1}\Gamma_0(\Gamma_0'\Gamma_0)^{-1}$. We can see that this variance depends on the expectation of the error term v_t, which includes the errors in both first stage and the second stage. The variance also depends on the distribution of the instrumental variable, $\mathcal{P}_z$. Notice that, although we need the assumption about ϵ_t to guarantee the consistency of the estimator, the asymptotic variance of the *BanditIV* estimator does not depend on ϵ_t.

5   Numerical experiments

In this section, we construct synthetic data to further validate our algorithm. Referring the simulation set up in Bakhitov and Singh 2021, we consider the model as follows, for $t = 1, \cdots, T$,

$$Y_t = X_t\beta_0 + e_t\rho + \varepsilon_t$$
$$X_t = Z_t\Gamma_0 + e_t + u_t$$

Suppose that all elements of the instrument $Z_t \in \mathbb{R}^{k \times 1}$ are uniformly distributed on the support $[-3, 3]$. $e_t \in \mathbb{R}^{n_t \times d} \sim \mathbb{N}(0, 1)$ is the confounder. $\rho \in \mathbb{R}^{d \times 1}$, is the parameter measuring the degree of endogeneity. When $\rho = 0$, endogeneity disappears, which we can see from Equation (3). $u_t \in \mathbb{R}^{n_t \times d}$, and $\varepsilon_t \in \mathbb{R}^{n_t \times 1}$ are i.i.d. normally distributed additional noise components, where n_t is the number of arms at time t.

$$\mathbb{E}[(e_t\rho + \varepsilon_t)'X_t] = \mathbb{E}[\rho'e_t'X_t] = \mathbb{E}[\rho'e_t'\mathbb{E}[X_t|e_t]] = \rho'\mathbb{E}[e_t'e_t] \tag{3}$$

In our simulation, we set $d = 3, k = 5, n_t = 50$ for $t = 1, \cdots, T$. We run $T = 2000$ timesteps for each algorithm. Figures 1 shows the results of regret variation and estimation bias for four algorithms including BanditIV, ϵ-BanditIV, TS and OFUL. We compare algorithm performance under endogeneity case when $\rho = [0.5 \quad 0.5 \quad 0.5]'$ (shown in Figure 1a and 1b) as well as non-endogeneity case when $\rho = [0. \quad 0. \quad 0.]'$ (shown in Figure 1c and 1d). We find that BanditIV and epsilon-greedy BanditIV outperform other two classic algorithms both on regret and inference. This outperformance is more significant under endogeneity cases. We can see that when the exogeneous assumption wrt covariates are satisfied, all the four algorithms achieve approximately zero bias in estimation. However, when endogenous covariates exist, only BanditIV and ϵ-BanditIV can achieve approximately zero bias in inference. Consistent with our theoretical analysis in Section 3.1, BanditIV reaches lower expected regret than ϵ-BanditIV, but ϵ-BanditIV can obtain a less biased estimation.

6   Conclusion

In this paper, we study the endogeneity problem in online decision making settings where we formulate the decision making process as a Contextual Linear Bandit model. The existence of endogeneity can lead to estimation bias and sub-optimal decision. To correct the bias and optimize the decision, we propose the *BanditIV* algorithm by utilizing both existing linear bandit algorithms in computer science literature and TSLS method in econometric literature. We present various theoretical properties of the algorithm. We first show an upper bound for the total expected regret, and then show the consistency, asymptotic normality of the estimator in the algorithm. On the applied side, we conduct simulations on synthetic data. We find that the *BanditIV* algorithm outperforms several benchmark linear bandit algorithms, especially when endogeneity problem occurs.

References

Abbasi-Yadkori, Yasin, Dávid Pál, and Csaba Szepesvári (2011). "Improved algorithms for linear stochastic bandits". In: Advances in neural information processing systems 24.

Ai, Chunrong and Xiaohong Chen (2003). "Efficient estimation of models with conditional moment restrictions containing unknown functions". In: Econometrica 71.6, pp. 1795–1843.

Angrist, Joshua and Guido Imbens (1995). Identification and estimation of local average treatment effects.

Aramayo, Nicolás, Mario Schiappacasse, and Marcel Goic (2022). "A Multi-Armed Bandit Approach for House Ads Recommendations". In: Available at SSRN 4107976.

Auer, Peter (2002). "Using confidence bounds for exploitation-exploration trade-offs". In: Journal of Machine Learning Research 3.Nov, pp. 397–422.

Bakhitov, Edvard and Amandeep Singh (2021). "Causal Gradient Boosting: Boosted Instrumental Variable Regression". In: arXiv preprint arXiv:2101.06078.

Bastani, Hamsa and Mohsen Bayati (2020). "Online decision making with high-dimensional covariates". In: Operations Research 68.1, pp. 276–294.

Bastani, Hamsa, Mohsen Bayati, and Khashayar Khosravi (2021). "Mostly exploration-free algorithms for contextual bandits". In: Management Science 67.3, pp. 1329–1349.

Bibaut, Aurélien et al. (2021). "Post-contextual-bandit inference". In: Advances in Neural Information Processing Systems 34, pp. 28548–28559.

Bojinov, Iavor, David Simchi-Levi, and Jinglong Zhao (2020). "Design and analysis of switchback experiments". In: arXiv preprint arXiv:2009.00148.

Carpentier, Alexandra, Claire Vernade, and Yasin Abbasi-Yadkori (2020). "The elliptical potential lemma revisited". In: arXiv preprint arXiv:2010.10182.

Chen, Haoyu, Wenbin Lu, and Rui Song (2021). "Statistical inference for online decision making: In a contextual bandit setting". In: Journal of the American Statistical Association 116.533, pp. 240–255.

Chen, Xiaohong, Han Hong, and Elie Tamer (2005). "Measurement error models with auxiliary data". In: The Review of Economic Studies 72.2, pp. 343–366.

Chernozhukov, Victor, Guido W Imbens, and Whitney K Newey (2007). "Instrumental variable estimation of nonseparable models". In: Journal of Econometrics 139.1, pp. 4–14.

Chu, Wei et al. (2011). "Contextual bandits with linear payoff functions". In: Proceedings of the Fourteenth International Conference on Artificial Intelligence and Statistics. JMLR Workshop and Conference Proceedings, pp. 208–214.

Dani, Varsha, Thomas P Hayes, and Sham M Kakade (2008). "Stochastic linear optimization under bandit feedback". In.

Durand, Audrey et al. (2018). "Contextual bandits for adapting treatment in a mouse model of de novo carcinogenesis". In: Machine learning for healthcare conference. PMLR, pp. 67–82.

Farias, Vivek et al. (2022). "Synthetically Controlled Bandits". In: arXiv preprint arXiv:2202.07079.

Griliches, Zvi (1977). "Estimating the returns to schooling: Some econometric problems". In: Econometrica: Journal of the Econometric Society, pp. 1–22.

Hadad, Vitor et al. (2021). "Confidence intervals for policy evaluation in adaptive experiments". In: Proceedings of the National Academy of Sciences 118.15, e2014602118.

Hall, Peter and Christopher C Heyde (2014). Martingale limit theory and its application. Academic press.

Hansen, Lars Peter (1982). "Large sample properties of generalized method of moments estimators". In: Econometrica: Journal of the econometric society, pp. 1029–1054.

Hausman, Jerry A (1983). "Specification and estimation of simultaneous equation models". In: Handbook of econometrics 1, pp. 391–448.

Imbens, Guido (2014). Instrumental variables: an econometrician's perspective. Tech. rep. National Bureau of Economic Research.

Imbens, Guido W and Donald B Rubin (2015). Causal inference in statistics, social, and biomedical sciences. Cambridge University Press.

Langford, John and Tong Zhang (2007). "The epoch-greedy algorithm for contextual multi-armed bandits". In: Advances in neural information processing systems 20.1, pp. 96–1.

Lattimore, Tor and Csaba Szepesvári (2020). Bandit algorithms. Cambridge University Press.

Li, Jin, Ye Luo, and Xiaowei Zhang (2021). "Causal reinforcement learning: An instrumental variable approach". In: arXiv preprint arXiv:2103.04021.

Li, Lihong et al. (2010). "A contextual-bandit approach to personalized news article recommendation". In: Proceedings of the 19th international conference on World wide web, pp. 661–670.

Liu, Bing et al. (2018). "Customized nonlinear bandits for online response selection in neural conversation models". In: Thirty-Second AAAI Conference on Artificial Intelligence.

M Wooldridge, Jeffrey (2014). Introductory econometrics.

Newey, Whitney K and James L Powell (2003). "Instrumental variable estimation of nonparametric models". In: Econometrica 71.5, pp. 1565–1578.

Nie, Xinkun et al. (2018). "Why adaptively collected data have negative bias and how to correct for it". In: International Conference on Artificial Intelligence and Statistics. PMLR, pp. 1261–1269.

Qin, Lijing, Shouyuan Chen, and Xiaoyan Zhu (2014). "Contextual combinatorial bandit and its application on diversified online recommendation". In: Proceedings of the 2014 SIAM International Conference on Data Mining. SIAM, pp. 461–469.

Tang, Liang et al. (2015). "Personalized recommendation via parameter-free contextual bandits". In: Proceedings of the 38th international ACM SIGIR conference on research and development in information retrieval, pp. 323–332.

Wagenmaker, Andrew et al. (2021). "First-Order Regret in Reinforcement Learning with Linear Function Approximation: A Robust Estimation Approach". In: arXiv preprint arXiv:2112.03432.

Woodroofe, Michael (1979). "A one-armed bandit problem with a concomitant variable". In: Journal of the American Statistical Association 74.368, pp. 799–806.

A   Appendix

Appendix A.1 Figures

Figure 1: Simulation results on synthetic data

Appendix A.2 Literature review

This paper focuses on Contextual Multi-Armed Bandits (CMAB) approach. Auer 2002 first introduced the contextual bandit setting through the Linear Reinforcement Learning (LinRel) algorithm with linear value functions. Subsequently, the contextual framework was improved by Dani, Hayes,

and Kakade 2008 and Chu et al. 2011 through Upper Confidence Bound (UCB) type algorithms. Abbasi-Yadkori, Pál, and Szepesvári 2011 further modified the analysis of the linear bandit problem and improved the regret bound by a logarithmic factor. More recently, Bastani, Bayati, and Khosravi 2021 studied performance of exploration-free policies in contextual bandit settings and proposed the Greedy-First algorithm which only utilizes observed contexts and rewards to decide whether to follow a greedy algorithm or to explore. This literature often requires independency between contexts and the random error term, which however cannot be always satisfied in the reality. We consider a setting which allows the dependency between contexts and error term. This more general setting is well suited to real-world applications where endogeneity problems happen.

This paper contributes to the stream of literature in inference with MAB. The adaptive nature of data generation in online settings complicates the inference of unknown parameters. Nie et al. 2018 showed that estimates of arm-specific expected rewards in UCB and TS algorithms are biased downwards. This downward bias is due to that arms with random upward fluctuation are sampled more, while those arms with downward fluctuation are sampled less (Hadad et al. 2021). Bibaut et al. 2021 presented that standard estimators no longer follow normal distribution asymptotically so that classic confidence interval fail to provide correct coverage. To address the inference problem in online settings, H. Chen, Lu, and Song 2021 studied the consistency and asymptotic distribution of the online ordinary least square estimator under epsilon-greedy policy. We extend their approach to two stage least square estimation in online settings and contribute to the literature by proving the consistency and asymptotic normality of our estimator.

This paper is also closely related to the stream of literature in instrumental variable methods for endogeneity problems. Griliches 1977, Hausman 1983, Angrist and G. Imbens 1995, and X. Chen, Hong, and Tamer 2005 provided theoretical analysis of instrumental variables in linear models, while Hansen 1982, Ai and X. Chen 2003, Newey and Powell 2003 and Chernozhukov, G. W. Imbens, and Newey 2007 studied instrumental variables in nonlinear models. G. Imbens 2014 reviewed work on instrumental variable methods and discussed applications and restrictions. Different from the standard data analysis framework in offline settings, the endogeneity problem can be exacerbated during the dynamic interaction between data generation and data analysis in online settings (J. Li, Luo, and X. Zhang 2021). This paper builds on existing instrumental variable methods in offline settings and considers the dynamic interaction by utilizing the CMAB framework. For these reasons, the paper contributes to the literature by combining both CMAB models and instrumental variable methods to improve the estimation of causal effect of treatments and hence achieve better performance of decision policies.

Appendix A.3 Further explanation for algorithm

Estimation: The algorithm maintains four matrices U_t, V_t, W_t, Q_t to calculate estimated parameters in each round as described in the pseudo code of Algorithm 2. If the current time t is after the first round, the algorithm utilizes past-choice-related instrumental variables Z_t and past choices X_t to estimate Γ_0 through OLS and obtain an estimated X_t, which is $\hat{X}_t$. Then, based on $\hat{X}_t$ and past observations of rewards Y_t, the algorithm estimates β_0 by OLS again. We denote the estimator for β_0, which is $\hat{\beta}_t$, in the *BanditIV* algorithm as the *BanditIV estimator*.

Confidence Sets: Following the *Optimisim in the Face of Uncertainty principle* (OFU), we need maintain confidence sets for all unknown parameters in the model. As described in Section 2, we have two unknown parameters, Γ_0 and β_0. Thus, we construct two confidence sets $C_{1,s}$, $C_{2,s}$, $s \in \{1, \cdots, T\}$ for the first and the second stage estimation respectively in each round. The idea for the confidence sets are to make the estimation optimistic with the conditions that "with high probability" the true coefficients are in the confidence sets and we can calculate the confidence sets from the past chosen arms X_t, related instrumental variables Z_t, and rewards Y_t.

Execution: The best arm generated by the algorithm is a sample x_t which is related to an instrumental variable z_t that can maximize the estimated reward jointly with a pair of optimistic estimates of two-stage coefficients in the confidence sets. After an arm is chosen, we observe the reward y_t as stated in Equation (1).

Algorithm 2 BanditIV

Input: $\gamma_z, \gamma_x, \{G_t\}_{t=1}^{\infty}, \{B_t\}_{t=1}^{\infty}$
Set $U_0 = \gamma_z I \in \mathbb{R}^{k \times k}, V_0 = 0 \in \mathbb{R}^{k \times d}, W_0 = \gamma_x I \in \mathbb{R}^{d \times d}, Q_0 = 0 \in \mathbb{R}^{d \times 1}$
Set $C_{1,s} = \{\Gamma : ||\Gamma^{(i)} - \hat{\Gamma}_s^{(i)}||_{U_s} \leq G_s\}, C_{2,s} = \{\beta : ||\beta - \hat{\beta}_s||_{W_s} \leq B_s\}, \forall s \in \{1, 2, \cdots T\},$
$\forall i \in \{1, 2, \cdots d\}$
Nature reveals $\mathcal{Z}_0$. We randomly choose $z_0 \in \mathcal{Z}_0$ and set $Z_0 = z_0$
Nature reveals $\mathcal{X}_0$. From the set $\mathcal{X}_0$, we play x_0 which is related with z_0 and then observe the reward y_0. We
set $X_0 = x_0$ and $Y_0 = y_0$
for $t := 1, 2, \cdots, T$, **do**
    $U_t = U_{t-1} + z_{t-1} z_{t-1}', V_t = V_{t-1} + z_{t-1} x_{t-1}'$
    $\hat{\Gamma}_t = U_t^{-1} V_t, \hat{X}_{t-1} = Z_{t-1} \hat{\Gamma}_t$
    $W_t = W_0 + \hat{X}_{t-1}' \hat{X}_{t-1}, Q_t = Q_0 + \hat{X}_{t-1}' Y_{t-1}$
    $\hat{\beta}_t = W_t^{-1} Q_t$
    Nature reveals $\mathcal{Z}_t$. We choose $z_t = \arg\max_{z \in \mathcal{Z}_t} \max_{\Gamma \in C_{1,t}} \max_{\beta \in C_{2,t}} \langle \Gamma' z, \beta \rangle$ and update $Z_t = [Z_{t-1}' \quad z_t]'$
    Nature reveals the set of arms $\mathcal{X}_t$. We play x_t which is related with z_t and observe the reward y_t. Update
$X_t = [X_{t-1}' \quad x_t]', Y_t = [Y_{t-1}' \quad y_t]'$
end for

Appendix A.4 Confidence intervals

To compute confidence intervals of our estimate of β_0 post running our bandits, we use a re-randomization test similar to that of Bojinov, Simchi-Levi, and Zhao 2020 and Farias et al. 2022. For the estimate $\hat{\beta}$, we test the sharp null hypothesis that the $\beta_0 = \tau$ for all t. The sharp null hypothesis implies that the new outcome is $y_t + \tau'(x_t^H - x_t)$ where x_t^H is a newly pulled arm and x_t is a pulled arm by our bandit algorithm.

We conduct exact tests by using the known assignment mechanism to simulate new assignment paths. Algorithm 3 provides the details of how to implement it. In particular, we propose τ by a downward search method based on the estimate from the bandit algorithm. Under the sharp null hypothesis of $\beta_0 = \tau$, a new arm assignment path $1 : T$ leads to a sequence of observed outcomes $y_t + \tau'(x_t^H - x_t)$ for $t \in \{1, \cdots, T\}$. To obtain a confidence interval, we invert a sequence of exact hypothesis tests to identify the region outside where the null hypothesis is violated at the prespecified significance level (G. W. Imbens and Rubin 2015, Bojinov, Simchi-Levi, and Zhao 2020).

Algorithm 3 Sharp Null Hypothesis Test

Require: Fix N_H, total number of samples drawn; Given $\hat{\Gamma}$, X, and Y from the bandit algorithm;
Given a prespecified significance level α
    for i in $1 : N_H$ **do**
        Sample a new assignment path, x_t^H, for $t \in \{1, \cdots, T\}$ according to the assignment mechanism
under the null hypothesis $\beta_0 = \tau$
        Change the sequence of original outcomes y_t to $y_t + \langle \tau, (x_t^H - x_t) \rangle$ for $t \in \{1, \cdots, T\}$
        Compute $\hat{\tau}^{[i]}$ as the original algorithm does, i.e.

$$\hat{X}^H = Z^H \hat{\Gamma}$$

$$\hat{\tau}^{[i]} = \gamma I + ((\hat{X}^H)' \hat{X}^H)^{-1} (\hat{X}^H)' Y^H$$

    end for
    Compute $\hat{p} = N_H^{-1} \sum_{i=1}^{N_H} \mathbf{1}\{|\hat{\tau}^{[i]}| > |\hat{\beta}|\}$
    Reject the null hypothesis if $\hat{p} < \alpha$

All of these experiments are run on synthetic data constructed as in Section 5. We run 50 rounds in total. In each round, we propose 20 hypotheses for β_0 based on the estimated coefficient of our main interest, $\hat{\beta}$ from a pre-run BanditIV or ϵ-greedy BanditIV and construct confidence intervals for all dimensions of β_0. For each hypothesis of β_0, we run the sharp null hypothesis test where we set N_H as 200. We choose the significance level to be $\alpha = 0.05$. Under BanditIV, the coverage of

the test for each dimension of the estimate is listed as follows: 94%, 96%, 96% ; Under ϵ-greedy BanditIV, the coverage of the test for each dimension of the estimate is listed as follows: 96%, 92%, 94%. We can see that, given the significance level as 0.05, the coverage of the test is close to ideal across both BanditIV and ϵ-greedy BanditIV. Thus, the re-randomization tests and corresponding confidence intervals reported here are adequate for inference of the main treatment effect.

Appendix A.5 Proof of Main Results

Proof for Theorem 1

Proof. This regret can be decomposed into

$$
R_T = \sum_{t=1}^{T}(1-\epsilon_t)\mathbb{E}_{u_t}[\langle x_t^*, \beta_0\rangle - \langle x_t, \beta_0\rangle]\mathbf{1}[x_t = \Gamma_0' z_t] + \sum_{t=1}^{T}\epsilon_t\mathbb{E}_{u_t}[\langle x_t^*, \beta_0\rangle - \langle x_t, \beta_0\rangle]\mathbf{1}[x_t \neq \Gamma_0' z_t]
$$

$$
\leq \sum_{t=1}^{T}\langle \Gamma_0' z_t^*, \beta_0\rangle - \langle \Gamma_0' z_t, \beta_0\rangle + 2\epsilon_0 T L_y
$$

Next we will upper bound the first term. Let $(\tilde{\Gamma}_t, \tilde{\beta}_t) = \arg\max_{\Gamma \in G_{t-1}, \beta \in C_{t-1}}\langle \Gamma' z_t, \beta\rangle$. By using the *two-stage* optimism we can decompose our regret as follows,

$$
\sum_{t=1}^{T}\langle \Gamma_0' z_t^*, \beta_0\rangle - \langle \Gamma_0' z_t, \beta_0\rangle
$$

$$
\leq \sum_{t=1}^{T}\max_{\Gamma \in \mathcal{C}_{2,t-1}}\max_{\beta \in \mathcal{C}_{2,t-1}}\langle \Gamma' z_t^*, \beta\rangle - \langle \Gamma_0' z_t, \beta_0\rangle
$$

$$
\leq \sum_{t=1}^{T}\langle \tilde{\Gamma}_t' z_t, \tilde{\beta}_t\rangle - \langle \Gamma_0' z_t, \beta_0\rangle
$$

$$
= \sum_{t=1}^{T}\langle (\tilde{\Gamma}_t - \Gamma_0)' z_t, (\tilde{\beta}_t - \beta_0) + \beta_0\rangle + \langle \hat{x}_t + (\Gamma_0 - \hat{\Gamma}_t)' z_t, \tilde{\beta}_t - \beta_0\rangle
$$

$$
\leq \sum_{t=1}^{T}||\hat{x}_t||_{W_{t-1}^{-1}}||\tilde{\beta}_t - \beta_0||_{W_{t-1}} + \sum_{t=1}^{T}||(\Gamma_0 - \hat{\Gamma}_t)' z_t||_2||\tilde{\beta}_t - \beta_0||_2
$$

$$
+ ||\beta_0||_2 \sum_{t=1}^{T}||(\tilde{\Gamma}_t - \Gamma_0)' z_t||_2 + \sum_{t=1}^{T}||(\tilde{\Gamma}_t - \Gamma_0)' z_t||_2||\tilde{\beta}_t - \beta_0||_2
$$

$$
= \sum_{t=1}^{T}||\hat{x}_t||_{W_{t-1}^{-1}}||\tilde{\beta}_t - \beta_0||_{W_{t-1}} + ||\beta_0||_2 \sum_{t=1}^{T}||(\tilde{\Gamma}_t - \Gamma_0)' z_t||_2 + 2\sum_{t=1}^{T}||(\Gamma_0 - \hat{\Gamma}_t)' z_t||_2||\tilde{\beta}_t - \beta_0||_2
$$

where the first inequality comes from the Lemma 1, 2 that $\Gamma_0 \in \mathcal{C}_{2,t}, \beta_0 \in \mathcal{C}_{1,t}$ for all t, the second inequality comes from the definition of z_t, the second equality comes from the definition of $\hat{x}_t$ and some careful decomposition, and the last inequality comes from cauchy-swartz inequality.

Now our goal is to upper bounded these three terms, which mainly comes form the estimation error of Γ_0 in the first stage and the estimation error of β_0 in the second stage. For the first term, by applying Cauchy-Schwartz inequality and the standard elliptical potential lemma in Carpentier, Vernade, and Abbasi-Yadkori 2020, we have

$$
\sum_{t=1}^{T}||\hat{x}_t||_{W_{t-1}^{-1}}||\tilde{\beta}_t - \beta_0||_{W_{t-1}} \leq \max_t ||\tilde{\beta}_t - \beta_0||_{W_{t-1}} \sum_{t=1}^{T}||\hat{x}_t||_{W_{t-1}^{-1}}
$$

$$
\leq \max_t ||\tilde{\beta}_t - \beta_0||_{W_{t-1}} \sqrt{2Td\log(\frac{T+d}{d})}
$$

For the second term, by using similar arguments, we have that

$$\|\beta_0\|_2 \sum_{t=1}^{T} \|(\tilde{\Gamma}_t - \Gamma_0)'z_t\|_2 \leq \|\beta_0\|_2 \sum_{t=1}^{T} \sum_{i=1}^{d} |(\tilde{\Gamma}_t^{(i)} - \Gamma_0^{(i)})'z_t|$$

$$\leq \|\beta_0\|_2 \sum_{t=1}^{T} \sum_{i=1}^{d} \|\tilde{\Gamma}_t^{(i)} - \Gamma_0^{(i)})\|_{U_{t-1}} \|z_t\|_{U_{t-1}^{-1}}$$

$$\leq \|\beta_0\|_2 d \max_{t\in[T],i\in[d]} \|\tilde{\Gamma}_t^{(i)} - \Gamma_0^{(i)})\|_{U_{t-1}} \sum_{t=1}^{T} \|z_t\|_{U_{t-1}^{-1}}$$

$$\leq \|\beta_0\|_2 d \max_{t\in[T],i\in[d]} \|\tilde{\Gamma}_t^{(i)} - \Gamma_0^{(i)})\|_{U_{t-1}} \sqrt{2Tk\log(\frac{T+k}{k})}$$

where the last inequality again comes from standard ellipse potential lemma.

Finally, for the third term, by the definition of W_t, we have $\|\tilde{\beta}_t - \beta_0\|_2 \leq \frac{1}{\gamma}\|\tilde{\beta}_t - \beta_0\|_{W_{t-1}}$. Therefore, by repeating the proof in the second term, we have that,

$$2\sum_{t=1}^{T} \|(\Gamma_0 - \hat{\Gamma}_t)'z_t\|_2 \|\tilde{\beta}_t - \beta_0\|_2$$

$$\leq \frac{2}{\gamma_x} \max_t \|\tilde{\beta}_t - \beta_0\|_{W_{t-1}} \max_{t\in[T],i\in[d]} \|\tilde{\Gamma}_t^{(i)} - \Gamma_0^{(i)})\|_{U_{t-1}} \sqrt{2Tk\log(\frac{T+k}{k})}$$

Therefore, by combing these three terms and the definition of $\mathcal{C}_{1,t}, \mathcal{C}_{2,t}$, we can upper bound our regret as

$$R_T \leq B_T \sqrt{2Td\log(\frac{T+d}{d})} + (\frac{2}{\gamma_x} + \|\beta_0\|_2)G_T \sqrt{2Tk\log(\frac{T+k}{k})}$$

$\square$

Lemma 1 (Optimistic estimation of β_0). *With high prob* $1 - \delta/2$, *for all* $t \in [T]$,

$$\|\hat{\beta}_t - \beta_0\|_{W_t} \leq B_t$$

where $B_t = \sqrt{\gamma_x}\|\beta_0\|_2 + \sqrt{2\log(\frac{2T}{\delta}) + d\log\left(\frac{5TL_x^2}{d}\right)}$,

Proof. At any fixed time t, we denote $\hat{x}_s^t = \Gamma_t'z_s$ for all $s \leq t$, so the $\hat{X}_t$ defined in the algorithm is a collection of all $\{\hat{x}_s^t\}_{s\leq t}$. For convenience, we drop the superscript t here. We also denote $\mathbf{e}_t \in \mathbb{R}^t$ as a collection of $\{e_s\}_{s\leq t}$. Now we get the closed-form for $\hat{\beta}_t$ by ridge 2SLS estimator as follows,

$$\hat{\beta}_t = W_t^{-1}Q_t$$
$$= W_t^{-1}\hat{X}_t'(X_t\beta_0 + \mathbf{e}_t)$$
$$= (X_t'P_{Z_t}X_t + \gamma_xI)^{-1}X_t'P_{Z_t}X_t\beta_0 + W_t^{-1}\hat{X}_t'\mathbf{e}_t$$
$$= \beta_0 - \gamma_xW_t^{-1}\beta_0 + W_t^{-1}\hat{X}_t'\mathbf{e}_t$$

where we denote $Z_t(Z_t'Z_t)^{-1}Z_t$ as P_{Z_t}. The third equality comes from the definition of $\hat{W}_t$ and the upper bound of $\hat{X}_t$ in Lemma 8 Therefore, we can write the estimation error of $\hat{\beta}$ compared to β_0 as

$$\|\hat{\beta}_t - \beta_0\|_{W_t} = \|W_t^{-1}\hat{X}_t'\mathbf{e}_t - \gamma_xW_t^{-1}\beta_0\|_{W_t}$$
$$= \|\hat{X}_t'\mathbf{e}_t - \gamma_x\beta_0\|_{W_t^{-1}}$$
$$\leq \|\hat{X}_t'\mathbf{e}_t\|_{W_t^{-1}} + \gamma_x\|\beta_0\|_{W_t^{-1}}$$
$$\leq \|\hat{X}_t'\mathbf{e}_t\|_{W_t^{-1}} + \gamma_x\|\beta_0\|_{(\gamma_xI)^{-1}}$$
$$= \|\hat{X}_t'\mathbf{e}_t\|_{W_t^{-1}} + \sqrt{\gamma_x}\|\beta_0\|_2$$

Finally, by noticing that $\mathbf{e}_t$ is 1-subgaussian and the that $\exp(\langle q, \hat{X}'_t \mathbf{e}_t \rangle - \frac{||q||^2_{\hat{X}'_t \hat{X}'_t}}{2})$ for all $q \in \mathbb{R}^d$ is a supermartingale, according to the Section 20.1 in Lattimore and Szepesvári 2020, we have that, with probability $1 - \delta/2T$,

$$||\hat{X}'_t \mathbf{e}_t||_{W_t^{-1}} \leq \sqrt{2 \log(\frac{2T}{\delta}) + \log(\frac{\det(W_t)}{\gamma_x^d}))} \leq B_t.$$

where the last inequality comes from the explicit calculation of W_t detailed in Lemma 8. $\qquad\square$

Lemma 2 (Optimistic estimation of Γ_0). *With high prob $1 - \delta/2$, for all $t \in [T]$ and all $i \in [d]$,*

$$||\hat{\Gamma}_t^{(i)} - \Gamma_0^{(i)}||_{U_t} \leq G_t$$

where $G_t = \sqrt{\gamma_z}||\Gamma_0^{(i)}||_2 + \sqrt{2 \log(\frac{2T}{\delta}) + k \log\left(\frac{5TL_y^2}{k}\right)}.$

Proof. The proof steps are similar to the previous lemma. At any fixed time t and dimension $i \in [d]$, we denote $\mathbf{u}_t^i \in \mathbb{R}^t$ as a collection of $\{e_s\}_{s \leq t}$. We again get the closed-form of $\hat{\Gamma}_t$ as

$$\hat{\Gamma}_t = U_t^{-1} V_t = U_t^{-1}(U_t - \gamma_z I)\Gamma_0 + U_t^{-1} Z'_t \mathbf{u}_t^i = \Gamma_0 - \gamma_z U_t^{-1}\Gamma_0 + U_t^{-1} Z'_t \mathbf{u}_t^i$$

And therefore, $||\hat{\Gamma}_t^{(i)} - \Gamma_0^{(i)}||_{U_t} \leq ||Z'_t \mathbf{u}_t^i||_{U_t^{-1}} + \sqrt{\gamma_z}||\Gamma_0^{(i)}||_2.$

Finally, again by noticing that $\mathbf{u}_t^i$ is 1-subgaussian and the that $\exp(\langle q, \hat{X}'_t \mathbf{u}_t^i \rangle - \frac{||q||^2_{\hat{X}'_t \hat{X}'_t}}{2})$ for all $q \in \mathbb{R}^d$ is a supermartingale, we have that, with probability $1 - \delta/2T$,

$$||\hat{\Gamma}_t^{(i)} - \Gamma_0^{(i)}||_{U_t} \leq \sqrt{\gamma_z}||\Gamma_0^{(i)}||_2 + \sqrt{2 \log(\frac{2T}{\delta}) + \log(\frac{\det(U_t)}{\gamma_z^k})} \leq G_t$$

where the last inequality comes from the explicit calculation of U_t detailed in Lemma 8 and the union bound over all $t \in [T]$.

$\qquad\square$

Lemma 3. *Suppose $\{\mathcal{F}_t : t = 1, \cdots, T\}$ is an increasing filtration of σ-fields. Let $\{W_t : t = 1, \cdots, T\}$ be a sequence of variables such that W_t is $\mathcal{F}_{t-1}$ measurable and $|W_t| \leq L_w$ almost surely for all t. Let $\{v_t : t = 1, \cdots, T\}$ be independent σ_v-subgaussian, and $v_t \perp \mathcal{F}_{t-1}$ for all t. Let $\mathcal{S} = \{s_1, \cdots, s_{|\mathcal{S}|}\} \subseteq \{1, \cdots, T\}$ be an index set where $|\mathcal{S}|$ is the number of elements in $\mathcal{S}$. Then for $\kappa > 0$,*

$$P(\sum_{s \in \mathcal{S}} W_s v_s \geq \kappa) \leq \exp\{-\frac{\kappa^2}{2|\mathcal{S}|\sigma_v^2 L_w^2}\}$$

The proof of this lemma is provided in Lemma 1 of H. Chen, Lu, and Song 2021.

Lemma 4. *(Dependent OLS Tail Inequality). For the online decision making model, if all realizations of z_t satisfy $||z_t||_\infty \leq L_z$ for all t, and $\hat{\Sigma} = \frac{1}{t}\sum_{s=1}^t z_s z'_s$ has minimum eigenvalue $\lambda_{min}(\hat{\Sigma}) > \lambda$ for some $\lambda > 0$ almost surely. Then for any $\eta > 0$,*

$$P(||\hat{\delta}_t - \delta_0||_1 \leq \eta) \geq 1 - 2k \exp\left(-\frac{t\lambda^2\eta^2}{2k^2\sigma_v^2 L_z^2}\right)$$

Proof. Based on the proofs provided by H. Chen, Lu, and Song 2021 and Bastani and Bayati 2020, we make some minor changes. The relation between eigenvalue and l_2 norm of symmetric matrix gives $||\hat{\Sigma}^{-1}||_2 = \lambda_{max}(\hat{\Sigma}^{-1}) = (\lambda_{min}(\hat{\Sigma}))^{-1}$. Therefore,

$$||\hat{\delta}_t - \delta_0||_2 = ||\hat{\Sigma}^{-1}(\frac{1}{t}\sum_{s=1}^t z_s v_s)||_2 \leq \frac{1}{t}||\hat{\Sigma}^{-1}||_2||\sum_{s=1}^t z_s v_s||_2 \leq \frac{1}{t\lambda}||\sum_{s=1}^t z_s v_s||_2$$

Hence, we have

$$P(||\hat{\delta}_t - \delta_0||_2 \leq \eta) \geq P(||\sum_{s=1}^{t} z_s v_s||_2 \leq t\lambda\eta)$$

$$\geq P(|\sum_{s=1}^{t} Z_{j,s} v_s| \leq \frac{t\lambda\eta}{\sqrt{k}}, \cdots, |\sum_{s=1}^{t} Z_{k,s} v_s| \leq \frac{t\lambda\eta}{\sqrt{k}})$$

$$= 1 - P(\bigcup_{j=1}^{k} \{|\sum_{s=1}^{t} Z_{j,s} v_s| > \frac{t\lambda\eta}{\sqrt{k}}\})$$

$$\geq 1 - \sum_{j=1}^{k} P(|\sum_{s=1}^{t} Z_{j,s} v_s| > \frac{t\lambda\eta}{\sqrt{k}})$$

Because we know that v_s in the above inequality are i.i.d. subgaussian and $v_s \perp z_s$, we can apply Lemma 3 and have the following

$$P(|\sum_{s=1}^{t} Z_{j,s} v_s| > \frac{t\lambda\eta}{\sqrt{k}}) \leq 2\exp\{-\frac{t\lambda^2\eta^2}{2k\sigma_v^2 L_z^2}\}$$

Thus,

$$P(||\hat{\delta}_t - \delta_0||_1 \leq \eta) \geq P(||\hat{\delta}_t - \delta_0||_2 \leq \frac{\eta}{\sqrt{k}}) \geq 1 - 2k\exp\{-\frac{t\lambda^2\eta^2}{2k^2\sigma_v^2 L_z^2}\}$$

$\square$

Lemma 5. *Let $\{z_t : t = 1, \cdots, T\}$ be a sequence of i.i.d. k-dimension random vectors such that all realizations of z_t satisfy $||z_t||_\infty \leq L_z$ for all t. Denote $\hat{\Sigma} = \frac{1}{T}\sum_{t=1}^{T} z_t z_t'$. If $\Sigma = \mathbb{E}[z_t z_t']$ has minimum eigenvalue $\lambda_{min}(\Sigma) > \lambda$ for some $\lambda > 0$, then*

$$P(\lambda_{min}(\hat{\Sigma}) \leq \frac{\lambda}{2}) \leq k\exp\{-\frac{T\lambda}{8L_z^2}\}$$

Proof. This proof is based on the proof of Lemma 3 in H. Chen, Lu, and Song 2021 with minor changes. First, we have

$$\lambda_{max}(\frac{z_t z_t'}{T}) = \max_{||a||_2=1} a'(\frac{z_t z_t'}{T})a = \frac{1}{T}\max_{||a||_2=1}(a'z_t)^2 \leq \frac{L_z^2}{T}$$

$$\mu_{min} \equiv \lambda_{min}(\mathbb{E}\hat{\Sigma}) = \lambda_{min}(\frac{1}{T}\sum_{t=1}^{t}\mathbb{E}[z_t z_t']) = \lambda_{min}(\Sigma) > \lambda$$

Then, using the Matrix Chernoff bound,

$$P(\lambda_{min}(\hat{\Sigma}) \leq \frac{\lambda}{2}) \leq P(\lambda_{min}(\hat{\Sigma}) \leq \frac{\mu_{min}}{2}) \leq k\exp\{-\frac{T\mu_{min}}{8L_z^2}\} \leq k\exp\{-\frac{T\lambda}{8L_z^2}\}$$

$\square$

Lemma 6.

$$||\hat{\Gamma}(\hat{\beta} - \beta_0)||_1 \geq \frac{\lambda_{min}(\hat{\Gamma})}{\sqrt{d}}||\hat{\beta} - \beta_0||_1$$

where $\lambda_{min}(\hat{\Gamma})$ is the smallest magnitude of a singular value of this matrix.

Proof. By doing Singular Value Decomposition for $\hat{\Gamma}$, we have the following, where the second equality is due to that U is orthonormal.

$$||\hat{\Gamma}(\hat{\beta} - \beta_0)||_2 = ||U\Sigma V'(\hat{\beta} - \beta_0)||_2 = ||\Sigma V'(\hat{\beta} - \beta_0)||_2$$

We assume $rank(\hat{\Gamma}) = min\{k,d\} = d$. Using matrix form, we have

$$
\Sigma V'(\hat{\beta} - \beta_0) = \begin{bmatrix} \sigma_1 & 0 & \cdots & 0 \\ 0 & \sigma_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & \sigma_d \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix} \begin{bmatrix} V^{(1)(1)} & V^{(2)(1)} & \cdots & V^{(d)(1)} \\ V^{(1)(2)} & V^{(2)(2)} & \cdots & V^{(d)(2)} \\ \vdots & \vdots & \vdots & \vdots \\ V^{(1)(d)} & V^{(2)(d)} & \cdots & V^{(d)(d)} \end{bmatrix} \begin{bmatrix} \hat{\beta}^{(1)} - \beta_0^{(1)} \\ \hat{\beta}^{(2)} - \beta_0^{(2)} \\ \vdots \\ \hat{\beta}^{(d)} - \beta_0^{(d)} \end{bmatrix}
$$

$$
= \begin{bmatrix} \sigma_1 \sum_{i=1}^{d} V^{(i)(1)}(\hat{\beta}^{(i)} - \beta_0^{(i)}) \\ \sigma_2 \sum_{i=1}^{d} V^{(i)(2)}(\hat{\beta}^{(i)} - \beta_0^{(i)}) \\ \vdots \\ \sigma_d \sum_{i=1}^{d} V^{(i)(d)}(\hat{\beta}^{(i)} - \beta_0^{(i)}) \\ 0 \\ \vdots \\ 0 \end{bmatrix}
$$

Hence,

$$
||\Sigma V'(\hat{\beta} - \beta_0)||_1 \geq ||\Sigma V'(\hat{\beta} - \beta_0)||_2 = \sqrt{\sum_{j=1}^{d} |\sigma_j \sum_{i=1}^{d} V^{(i)(j)}(\hat{\beta}^{(i)} - \beta_0^{(i)})|^2}
$$

$$
\geq \lambda_{min}(\hat{\Gamma}) \sqrt{\sum_{j=1}^{d} |\sum_{i=1}^{d} V^{(i)(j)}(\hat{\beta}^{(i)} - \beta_0^{(i)})|^2}
$$

$$
= \lambda_{min}(\hat{\Gamma}) ||V'(\hat{\beta} - \beta_0)||_2
$$

$$
= \lambda_{min}(\hat{\Gamma}) ||(\hat{\beta} - \beta_0)||_2 \tag{4}
$$

where the last equality is due to that V is orthonormal.

Applying Cauchy–Schwarz inequality, we obtain

$$
||(\hat{\beta} - \beta_0)||_2 \geq \frac{1}{\sqrt{d}} ||(\hat{\beta} - \beta_0)||_1 \tag{5}
$$

Combing inequalities (4) and (5), we complete the proof. $\square$

Lemma 7.

$$
\lambda_{min}(\hat{\Gamma}) \geq \lambda_{min}(\Gamma_0) - \eta_2(kd)^{\frac{1}{2}} \tag{6}
$$

Proof. On the event that $||\hat{\Gamma} - \Gamma_0|| \leq \eta_2$, we can rewrite the equation for each element in $\hat{\Gamma}$ using a newly defined matrix $A \in \mathbb{R}^{k \times d} : |A^{(i)(l)}| \leq 1, \forall i \in i \in \{1, 2, \cdots, d\}, \forall l \in \{1, 2, \cdots, k\}$.

$$
\hat{\Gamma}^{(i)(l)} = \Gamma_0^{(i)(l)} + \eta_2 A^{(i)(l)} \tag{7}
$$

We know that all singular values are non-negative. Here, we assume the singular values of Γ_0 are strictly positive. We can write the singular values of Γ_0 and $\hat{\Gamma}$ as two increasing sequences, $0 < \sigma_\Gamma^{(1)} \leq \sigma_\Gamma^{(2)} \leq \cdots \leq \sigma_\Gamma^{(d)}$ and $\sigma^{(1)} \leq \sigma^{(2)} \leq \cdots \leq \sigma^{(d)}$, respectively.

Using the min-max principle for singular values, we have

$$
\sigma_\Gamma^{(1)} = \min_{S:dim(S)=1} \max_{\substack{a \in S \\ ||a||_2=1}} ||\Gamma_0 a||_2 = \min_{\substack{a \in \mathbb{R}^d \\ ||a||_2=1}} ||\Gamma_0 a||_2
$$

where the first equality is directly derived by applying the min-max theorem. The second equality is from the two facts. First, a should be of an order $d \times 1$ to fit the order of Γ_0. Second, there are only two unique vectors fitting the constraint $||a||_2 = 1$ in the given space S and they generate the same objective values $||\Gamma_0 a||_2$.

Similarly, we can rewrite the smallest singular value of $\hat{\Gamma}$ as the following

$$\sigma^{(1)} = \min_{\substack{a \in \mathbb{R}^d \\ ||a||_2 = 1}} ||\hat{\Gamma} a||_2 \tag{8}$$

$$
\begin{aligned}
\min_{\substack{a \in \mathbb{R}^d \\ ||a||_2 = 1}} ||\hat{\Gamma} a||_2 &\geq \min_{\substack{a \in \mathbb{R}^d \\ ||a||_2 = 1}} ||\Gamma_0 a||_2 - \eta_2 ||Aa||_2 \\
&\geq \min_{\substack{a \in \mathbb{R}^d \\ ||a||_2 = 1}} ||\Gamma_0 a||_2 - \eta_2 ||A||_2 ||a||_2 \\
&= \min_{\substack{a \in \mathbb{R}^d \\ ||a||_2 = 1}} ||\Gamma_0 a||_2 - \eta_2 ||A||_2 \\
&\geq \min_{\substack{a \in \mathbb{R}^d \\ ||a||_2 = 1}} ||\Gamma_0 a||_2 - \eta_2 (kd)^{\frac{1}{2}} \\
&= \sigma_\Gamma^{(1)} - \eta_2 (kd)^{\frac{1}{2}}
\end{aligned}
$$

where the first inequality is derived from the equation (7) and triangle inequality. The third equality is from the constraint that $||a||_2 = 1$. The fourth inequality is from the property of Frobenius norm, which we illustrate as the following

$$||A||_2 \leq ||A||_F = \left(\sum_{i=1}^{d} \sum_{l=1}^{k} |A^{(i)(l)}|^2\right)^{\frac{1}{2}} \leq \left(\sum_{i=1}^{d} \sum_{l=1}^{k} 1\right)^{\frac{1}{2}} = (kd)^{\frac{1}{2}}$$

$\square$

Proposition 2. *(Tail bounds for the online OLS estimators). In the online decision making model with the ϵ-greedy policy, if Assumptions 1 and 2 are satisfied, and ϵ_t is non-increasing, then for any $\eta_1, \eta_2 > 0$, any $i \in \{1, \cdots, d\}$,*

$$
\begin{aligned}
P(||\hat{\delta}_t - \delta_0||_1 \leq \eta_1) \geq {} & 1 - \exp\left\{-\frac{t\epsilon_t}{8}\right\} - k \exp\left\{-\frac{t\epsilon_t \lambda}{32 L_z^2}\right\} - 2k \exp\left\{-\frac{t\epsilon_t^2 \lambda^2 \eta_1^2}{128 k^2 \sigma_v^2 L_z^2}\right\} \\
& + 2k^2 \exp\left\{-\frac{t\epsilon_t^2 \lambda^2 \eta_1^2 + 4 t\epsilon_t \lambda k^2 \sigma_v^2}{128 k^2 \sigma_v^2 L_z^2}\right\}
\end{aligned}
$$

$$
\begin{aligned}
P(||\hat{\Gamma}_t^{(i)} - \Gamma_0^{(i)}||_1 \leq \eta_2) \geq {} & 1 - \exp\left\{-\frac{t\epsilon_t}{8}\right\} - k \exp\left\{-\frac{t\epsilon_t \lambda}{32 L_z^2}\right\} - 2k \exp\left\{-\frac{t\epsilon_t^2 \lambda^2 \eta_2^2}{128 k^2 \sigma_u^2 L_z^2}\right\} \\
& + 2k^2 \exp\left\{-\frac{t\epsilon_t^2 \lambda^2 \eta_2^2 + 4 t\epsilon_t \lambda k^2 \sigma_u^2}{128 k^2 \sigma_u^2 L_z^2}\right\}
\end{aligned}
$$

Proof for Proposition 2

Proof. Following the proof of Proposition 3.1 in H. Chen, Lu, and Song 2021, we provide this proof adapting to our setting. Denote $\mathcal{S}_t = \{1, \cdots, t\}$ and define $\hat{\Sigma}(\mathcal{I}) = |\mathcal{I}|^{-1} \sum_{s \in \mathcal{I}} z_s z_s'$ for any $\mathcal{I} \subseteq \mathcal{S}_t$, where $|\mathcal{I}|$ is the number of element in the set $\mathcal{I}$ and $|\mathcal{S}_t| = t$. We have

$$\hat{\delta}_t - \delta_0 = \left(\frac{1}{t} \sum_{s=1}^{t} z_s z_s'\right)^{-1} \frac{1}{t} \sum_{s=1}^{t} z_s v_s = \{\hat{\Sigma}(\mathcal{S}_t)\}^{-1} \frac{1}{t} \sum_{s=1}^{t} z_s v_s$$

523 Under ϵ-greedy policy, we pull the estimated optimal arm with probability ϵ_s at each time point s.
524 We denote the collection of time points up to time t when the estimated optimal arm is chosen as
525 $\mathcal{R}_t$. We will first bound the minimum eigenvalue of $\hat{\Sigma}(\mathcal{R}_t)$ and then use it to infer the bound of the
526 minimum eigenvalue of $\hat{\Sigma}(\mathcal{S}_t)$.

527 We denote the following event,

$$E := \{\lambda_{min}(\hat{\Sigma}(\mathcal{R}_t)) > \frac{\lambda}{4}\}$$

528 Applying Lemma 5, if $\lambda_{min}(\Sigma) > \lambda$, then we have

$$P(E) \geq 1 - k \exp\{-\frac{|\mathcal{R}_t|\lambda}{16L_z^2}\}$$

529 Meanwhile, we know that

$$\hat{\Sigma}(\mathcal{S}_t) = \frac{|\mathcal{R}_t|}{|\mathcal{S}_t|}\hat{\Sigma}(\mathcal{R}_t) + \frac{|\mathcal{S}_t| - |\mathcal{R}_t|}{|\mathcal{S}_t|}\hat{\Sigma}(\mathcal{S}_t \backslash \mathcal{R}_t)$$

530 By Weyl's inequality, on event E,

$$\lambda_{min}(\hat{\Sigma}(\mathcal{S}_t)) \geq \lambda_{min}(\frac{|\mathcal{R}_t|}{|\mathcal{S}_t|}\hat{\Sigma}(\mathcal{R}_t)) + \lambda_{min}(\frac{|\mathcal{S}_t| - |\mathcal{R}_t|}{|\mathcal{S}_t|}\hat{\Sigma}(\mathcal{S}_t \backslash \mathcal{R}_t)) \geq \frac{|\mathcal{R}_t|}{|\mathcal{S}_t|}\lambda_{min}(\hat{\Sigma}(\mathcal{R}_t)) \geq \frac{\lambda|\mathcal{R}_t|}{4t}$$

531 Applying Lemma 4, we have

$$P(||\hat{\delta}_t - \delta_0||_1 \leq \eta|E) \geq 1 - 2k \exp\left(-\frac{|\mathcal{R}_t|^2\lambda^2\eta^2}{32tk^2\sigma_v^2 L_z^2}\right)$$

532 Hence,

$$P(||\hat{\delta}_t - \delta_0||_1 \leq \eta) \geq P(||\hat{\delta}_t - \delta_0||_1 \leq \eta|E)P(E)$$
$$\geq 1 - 2k \exp\left(-\frac{|\mathcal{R}_t|^2\lambda^2\eta^2}{32tk^2\sigma_v^2 L_z^2}\right) - k \exp\{-\frac{|\mathcal{R}_t|\lambda}{16L_z^2}\} + 2k^2 \exp\{-\frac{|\mathcal{R}_t|\lambda}{16L_z^2} - \frac{|\mathcal{R}_t|^2\lambda^2\eta^2}{32tk^2\sigma_v^2 L_z^2}\}$$

533 The checking step for that $|\mathcal{R}_t|$ is large enough is exactly the same as that in H. Chen, Lu, and Song
534 2021. We complete the proof by combining all the results above,

$$P(||\hat{\delta}_t - \delta_0||_1 \leq \eta) \geq 1 - \exp\{-\frac{t\epsilon_t}{8}\} - k \exp\{-\frac{t\epsilon_t\lambda}{32L_z^2}\} - 2k \exp\{-\frac{t\epsilon_t^2\lambda^2\eta^2}{128k^2\sigma_v^2 L_z^2}\}$$
$$+ 2k^2 \exp\{-\frac{t\epsilon_t^2\lambda^2\eta^2 + 4t\epsilon_t\lambda k^2\sigma_v^2}{128k^2\sigma_v^2 L_z^2}\}$$

535 We can prove the inequality for $P(||\hat{\Gamma}_t^{(i)} - \Gamma_0^{(i)}||_1 \leq \eta_2)$ by similar steps. $\qquad \square$

536 Following from the Proposition 2, we can easily obtain the following corollary.
537 **Corollary 4.** *(Consistency of the online OLS estimators). If Assumptions 1 and 2 are satisfied, ϵ_t is*
538 *non-increasing and $t\epsilon_t^2 \to \infty$ as $t \to \infty$, then the online OLS estimators $\hat{\Gamma}_t$ is a consistent estimator*
539 *for Γ_0 and $\hat{\delta}_t$ is a consistent estimator for δ_0.*

540 **Proof for Proposition 1**

541 *Proof.* By Proposition 2, we have $||\hat{\Gamma} - \Gamma_0||_1 = \max_{1 \leq i \leq d}||\hat{\Gamma}^{(i)} - \Gamma_0^{(i)}||_1 \leq \eta_2$ with a probability
542 larger than p_2 and $||\hat{\delta} - \delta_0||_1 \leq \eta_1$ with a probability larger than p_1. On the events that $||\hat{\Gamma} - \Gamma_0||_1 \leq \eta_2$
543 and $||\hat{\delta} - \delta_0||_1 \leq \eta_1$, we can prove the upper bound of $||\hat{\beta} - \beta_0||_1$ as the following by utilizing
544 triangle inequalities.

$$||\hat{\Gamma}(\hat{\beta} - \beta_0)||_1 - ||(\hat{\Gamma} - \Gamma_0)\beta_0||_1 \leq ||\hat{\Gamma}(\hat{\beta} - \beta_0) + (\hat{\Gamma} - \Gamma_0)\beta_0||_1 = ||\hat{\Gamma}\hat{\beta} - \Gamma_0\beta_0||_1 = ||\hat{\delta} - \delta_0||_1 \leq \eta_1$$

545 Hence,

$$||\hat{\Gamma}(\hat{\beta} - \beta_0)||_1 \leq \eta_1 + ||(\hat{\Gamma} - \Gamma_0)\beta_0||_1 \leq \eta_1 + ||\hat{\Gamma} - \Gamma_0||_1||\beta_0||_1 \leq \eta_1 + \eta_2||\beta_0||_1 \qquad (9)$$

546 Combining inequality (9), Lemma 6 and Lemma 7, we complete the proof.

547 $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \square$

548   **Theorem 5.** *(Asymptotic normality of the online OLS estimator) If Assumptions 1 and 2 are satisfied,*
549   ϵ_t *is non-increasing and* $t\epsilon_t^2 \to \infty$ *as* $t \to \infty$, *then*

$$\sqrt{t}(\hat{\delta}_t - \delta_0) \xrightarrow{d} \mathcal{N}_k(0, \mathbb{E}[v_t^2](\int zz' d\mathcal{P}_z)^{-1})$$

550   **Theorem** 5

551   *Proof.* In this proof, we mainly use martingale central limit theorem.

552   By definition of the OLS estimator $\hat{\delta}_t$, we have

$$\sqrt{t}(\hat{\delta}_t - \delta_0) = (\frac{1}{t}\sum_{s=1}^{t} z_s z_s')^{-1}(\frac{1}{\sqrt{t}}\sum_{s=1}^{t} z_s v_s)$$

553   We define for $1 \le j \le t, t \ge 1$, and any $q \in \mathbb{R}^k$, the σ-field $\mathcal{F}_{tj}$ as $\mathcal{F}_j$,

$$H_{tj} = \frac{1}{\sqrt{t}}\sum_{s=1}^{j} q' z_s v_s.$$

554   We have $\mathbb{E}[q' z_s v_s | \mathcal{F}_{s-1}] = 0$. We also notice that $\{H_{tj}, \mathcal{F}_{tj}, 1 \le j \le t, t \ge 1\}$ is a martingale array.
555   Hence we are going to prove the convergence of H_{tt}.

556   We first check the Linderberg condition as follows, for each fixed $\zeta > 0$,

$$\sum_{s=1}^{t} \mathbb{E}[\frac{1}{t}(q' z_s)^2 v_s^2 I\{|q' z_s v_s| > \zeta\sqrt{t}\} | \mathcal{F}_{t,s-1}]$$

$$\le \frac{||q||_2^2 L_z^2}{t} \sum_{s=1}^{t} \mathbb{E}[v_s^2 I\{v_s^2 > \frac{\zeta^2 t}{||q||_2^2 L_z^2}\} | \mathcal{F}_{s-1}]$$

$$\le ||q||_2^2 L_z^2 \mathbb{E}[v_1^2 I\{v_1^2 > \frac{\zeta^2 t}{||q||_2^2 L_z^2}\}]$$

557   where $v_s, s = 1, \cdots, t$ are i.i.d. from $\mathcal{P}_v$. Notice that $v_1^2 I\{v_1^2 > \frac{\zeta^2 t}{||q||_2^2 L_z^2}\}$ is dominated by v_1^2 with
558   $\mathbb{E}[v_1^2] \le ||\beta_0||_2^2 + 1$. Also, $v_1^2 I\{v_1^2 > \frac{\zeta^2 t}{||q||_2^2 L_z^2}\}$ converges to 0 almost surely when $t \to \infty$. Thus,
559   applying dominated convergence theorem, we can get, as $t \to \infty$,

$$\sum_{s=1}^{t} \mathbb{E}[\frac{1}{t}(q' z_s)^2 v_s^2 I\{|q' z_s v_s| > \zeta\sqrt{t}\} | \mathcal{F}_{t,s-1}] \to 0$$

560   Then we are to find the limit of the conditional variance for H_{tt}. Notice that

$$\mathbb{E}[(q' z_s)^2 | \mathcal{F}_{s-1}] = \int q' z_s z_s' q d\mathcal{P}_z$$

561   and $q' \mathbb{E}[z_s z_s'] q \le ||q||_2^2 L_z^2$. Therefore,

$$\sum_{s=1}^{t} \mathbb{E}[\frac{1}{t}(q' z_s)^2 v_s^2 | \mathcal{F}_{t,s-1}]$$

$$= \frac{\mathbb{E}[v_1^2]}{t} \sum_{s=1}^{t} \mathbb{E}[(q' z_s)^2 | \mathcal{F}_{s-1}]$$

$$\xrightarrow{p} \mathbb{E}[v_1^2] q'(\int z_s z_s' d\mathcal{P}_z) q \qquad (10)$$

562   Applying Martingale Limit Theorem, we have

$$\frac{1}{\sqrt{t}}\sum_{s=1}^{t} z_s v_s \xrightarrow{d} \mathcal{N}_k(0, \mathbb{E}[v_1^2] \int z_s z_s' d\mathcal{P}_z) \qquad (11)$$

Now we are to find the limit of $\frac{1}{t}\sum_{s=1}^{t} q'z_s z_s' q$ for any $q \in \mathbb{R}^k$. We know that $q'z_s z_s' q$ is $\mathcal{F}_s$-measurable for each s and $\mathbb{E}[q'z_s z_s' q] \le q'\mathbf{1}\mathbf{1}'q L_z^2 < \infty$. Thus, by Theorem 2.19 in Hall and Heyde 2014, we have

$$\frac{1}{t}\sum_{s=1}^{t} (q'z_s z_s' q - \mathbb{E}[q'z_s z_s' q|\mathcal{F}_{s-1}]) \xrightarrow{p} 0 \tag{12}$$

From the result in (10) and (12), we have $\frac{1}{t}\sum_{s=1}^{t} q'z_s z_s' q \xrightarrow{p} q'(\int z_s z_s' d\mathcal{P}_z)q$. Applying Lemma 6 in H. Chen, Lu, and Song 2021 and Continuous Mapping Theorem, we obtain

$$(\frac{1}{t}\sum_{s=1}^{t} z_s z_s')^{-1} \xrightarrow{p} (\int z_s z_s' d\mathcal{P}_z)^{-1} \tag{13}$$

Combining the results in (11) and (13), we complete the proof.

$\square$

Proof for Theorem 3: Combining the Theorem 5 and Corollary 4, we derive the normality of the BanditIV estimator by applying Slutsky Theorem directly.

Appendix B. Auxiliary Lemmas

Lemma 8. *For each t, we can explicitly upper bound the following terms,*

$$W_t \le X_t' Z_t' (Z_t' Z_t)^{-1} Z_t X_t, \quad \log(\det(W_t)) \le d\log\left(\frac{5TL_x^2}{d}\right)$$

$$U_t \le Z_t' Z_t + \gamma_z I, \quad \log(\det(U_t)) \le k\log\left(\frac{5TL_y^2}{k}\right)$$

$$\hat{X}_t \le P_{Z_t} X_t$$

Proof. For the first inequality,

$$\hat{X}_t = Z_t \hat{\Gamma}_t = Z_t (Z_t' Z_t + \gamma_z I)^{-1} Z_t' X_t \le Z_t (Z_t' Z_t)^{-1} Z_t' X_t.$$

Then for the second inequality,

$$\begin{aligned}
W_t &= \hat{X}_t' \hat{X}_t \\
&= (Z_t \hat{\Gamma}_t)'(Z_t \hat{\Gamma}_t) \\
&= (Z_t (Z_t' Z_t + \gamma_z I)^{-1} Z_t' X_t)'(Z_t (Z_t' Z_t + \gamma_z I)^{-1} Z_t' X_t) \\
&= X_t' Z_t (Z_t' Z_t + \gamma_z I)^{-1} Z_t' Z_t (Z_t' Z_t + \gamma_z I)^{-1} Z_t' X_t \\
&\le X_t' Z_t (Z_t' Z_t)^{-1} Z_t' X_t
\end{aligned}$$

Recall that $P_{Z_t} = Z_t'(Z_t' Z_t)^{-1} Z_t$, we finish the upper bound of W_t. Then we have

$$\begin{aligned}
\log(\det(W_t)) &\le d\log\left(\frac{1}{d}\text{trace}(W_t)\right) \\
&\le d\log\left(\frac{1}{d}\text{trace}((Z_t' Z_t)^{-1} Z_t' X_t X_t' Z_t)\right) \\
&= d\log\left(\frac{1}{d}\text{trace}((Z_t' Z_t)^{-1} Z_t'(Z_t \Gamma_0 + \boldsymbol{u}_t)(Z_t \Gamma_0 + \boldsymbol{u}_t)' Z_t)\right) \\
&= d\log\left(\frac{1}{d}\left(\text{trace}(\Gamma_0 \Gamma_0' Z_t' Z_t) + \text{trace}(\Gamma_0 \Gamma_0' Z_t' \boldsymbol{u}_t \boldsymbol{u}_t' Z_t) + 2\text{trace}(Z_t' \boldsymbol{u}\Gamma_0')\right)\right)
\end{aligned}$$

Now because u_t is 1-subgaussian, then we have with high probability, $\boldsymbol{u}_t\boldsymbol{u}_t' < 2I$. Then the above is upper bounded by

$$d\log\left(\frac{1}{d}\left(3\|Z_t\Gamma_0\|_F^2 + 2\|Z_t\Gamma_0\|_F\|\boldsymbol{u}_t\|_F\right)\right) \le d\log\left(\frac{5TL_x^2}{d}\right)$$

and similarly

$$\log(\det(U_t)) \le k\log\left(\frac{5TL_y^2}{k}\right)$$

$\square$

Lemma 9. *Consider a sequence of vectors $(x_t)_{t=1}^T, x_t \in \mathbb{R}^d$, and assume that $\|x_t\|_2 \le a$ for all t. Let $V_t = \lambda I + \sum_{s=1}^t x_s x_s'$ for some $\lambda > 0$. Then, we will have that $\|x_t\|_{V_{t-1}^{-1}} > b$ at most*

$$d\log(1 + a^2T/\lambda)/\log(1 + b)$$

times.

Lemma 9 is directly from lemma 6.2 in Wagenmaker et al. 2021.

Lemma 10. *$a_1, a_2, \cdots \in \mathbb{R}^d$ is $\mathcal{F}_t$ adapted and $\mathbb{E}[a_t|\mathcal{F}_{t-1}] = 0$ and let $L_t \in \mathbb{R}^{d\times d}$ be a predictable sequence such that for any $\lambda \in \mathbb{R}^d$,*

$$\mathbb{E}[\exp\left(\langle\lambda, a_t\rangle\right)|\mathcal{F}_{t-1}] \le \exp\left(\|\lambda\|_{L_t}^2/2\right)$$

Let $H_t = \sum_{s=1}^t a_s$, $J_t = \sum_{s=1}^t L_t$.

$$K_t(\lambda) = \exp\left(\langle\lambda, H_t\rangle - \|\lambda\|_{J_t}^2/2\right)$$

is a super martingale. Moreover, we have that

$$\mathbb{P}(\exists t : \|H_t\|_{(J_t+\gamma_x I)^{-1}} \ge \sqrt{2\log(1/\delta) + \log(\frac{\det(J_t + \gamma_x I)}{\gamma_x^d})}) \le \delta$$

Lemma 10 is from the lemma for vector-valued martingale in Kevin's lecture notes (Proposition 3 in Section 3.3.5).

Proof for "$v^{(i)}$ is $(\|\beta_0\|_2^2 + 1)$-subgaussian"

Proof.

$$\mathbb{E}[exp(\lambda v^{(i)})] = \mathbb{E}[exp(\lambda(\sum_{l=1}^d u^{(i)(l)}\beta_0^{(l)} + e^{(i)}))]$$

$$= \prod_{l=1}^d \mathbb{E}[exp(\lambda u^{(i)(l)}\beta_0^{(l)})]\mathbb{E}[exp(\lambda e^{(i)})]$$

$$\le \prod_{l=1}^d \mathbb{E}[exp(\frac{\lambda^2\beta_0^{(l)^2}}{2})]\mathbb{E}[exp(\frac{\lambda^2}{2})]$$

$$= exp(\frac{\lambda^2}{2}(\sum_{l=1}^d \beta_0^{(l)^2} + 1))$$

$$= exp(\frac{\lambda^2}{2}(\|\beta_0\|_2^2 + 1))$$

where the second equality is due to the independency of distributions of $u^{(i)(l)}$ and $e^{(i)}$. The third inequality is because that $u^{(i)(l)}$ and $e^{(i)}$ are 1-subgaussian with mean zero. $\square$