Scalable Quasi-Bayesian Inference for Instrumental Variable Regression

Abstract

Recent years have witnessed an upsurge of interest in employing flexible machine learning models for instrumental variable (IV) regression, but the development of uncertainty quantification methodology is still lacking. In this work we present a scalable quasi-Bayesian procedure for IV regression, building upon the recently developed kernelized IV models. Contrary to Bayesian modeling for IV, our approach does not require additional assumptions on the data generating process, and leads to a scalable approximate inference algorithm with time cost comparable to the corresponding point estimation methods. Our algorithm can be further extended to work with neural network models. We analyze the theoretical properties of the proposed quasi-posterior, and demonstrate through empirical evaluation the competitive performance of our method.

1 Introduction

Instrumental variable (IV) regression is a standard approach for estimating causal effect from confounded observational data. In the presence of confounding, any regression method estimating $\mathbb{E}(\mathbf{y} \mid \mathbf{x})$ cannot recover the causal relation $f^{\dagger}$ between the outcome $\mathbf{y}$ and the treatment $\mathbf{x}$, since the residual $\mathbf{u} = \mathbf{y} - f^{\dagger}(\mathbf{x})$ is correlated with $\mathbf{x}$ due to the unobserved confounders. IV regression enables identification of the causal effect through the introduction of *instruments*, variables $\mathbf{z}$ that are known to influence $\mathbf{y}$ only through $\mathbf{x}$.

IV regression is widely used in economics [1], epidemiology [2] and clinical research [3], but modeling nonlinear effect in IV regression can be challenging. Recent years have seen great development in adopting modern machine learning models for IV regression [4–8]. However, there is still a lack of uncertainty quantification measures for these flexible IV models. Uncertainty quantification is especially important for IV analysis, since unlike in standard supervised learning scenarios, we do not have (unconfounded) validation data, from which we could deduce the error pattern of the estimated model. Moreover, the instrument of choice may be weak, meaning that it only provides limited information for $\mathbf{x}$; in such cases point estimators suffer from high variance [9]. This problem is exacerbated in the nonparametric setting, where IV estimation is known to be ill-posed [10, 11].

Unfortunately, the IV setting brings unique challenges for uncertainty quantification. For example, while it is natural to consider a Bayesian approach, specification of the likelihood requires knowledge of the entire data generating process, which is typically unavailable in IV regression. Consequently, most, if not all, existing work on Bayesian IV [12–16] assumes the following data generating process: $\mathbf{x} = g(\mathbf{z}) + \mathbf{u}_1$, $\mathbf{y} = f(\mathbf{x}) + \mathbf{u}_2$, where $(\mathbf{u}_1, \mathbf{u}_2)$ are correlated and independent of $\mathbf{z}$. These methods then conduct posterior inference on g, f as well as the *generative model* for the unobserved confounders $(\mathbf{u}_1, \mathbf{u}_2)$. However, the additive error in $\mathbf{x}$ is an unnecessary assumption for most point estimation procedures, and is difficult to check in high dimensions. Furthermore, the need to model the generating process of $(\mathbf{u}_1, \mathbf{u}_2)$ introduces an extra risk of model misspecifica-

tion, and Bayesian inference on the generative model is computationally expensive, especially on complex high-dimensional datasets. None of these issues present if only point estimation is needed.

For the above reasons, it is appealing to turn to an alternative *quasi-Bayesian* approach [17–19]. Quasi-Bayesian analysis views IV estimation as a generalized method-of-moments (GMM) procedure, and defines the quasi-posterior as a Gibbs distribution constructed from a chosen prior and violation of the moment constraints. It does not require full knowledge of the data generating process, and thus does not suffer from the aforementioned drawbacks of Bayesian inference. However, computation of the quasi-posterior is non-trivial, as its density contains a conditional expectation term $\mathbb{E}(\mathbf{y} - f(\mathbf{x}) \mid \mathbf{z})$, which itself needs to be estimated from data. This makes both approximate inference and theoretical analysis difficult. So far, quasi-Bayesian analysis for nonparametric IV is only developed on classical models such as wavelet basis [18, 19] or Nadaraya-Watson smoothing [20], while adoption of flexible machine learning models such as kernel machines or neural networks remains an open challenge. Moreover, numerical study has been limited in previous work, so little is known about the empirical performance of the quasi-Bayesian approach.

In this work, we present a novel quasi-Bayesian procedure for IV regression, building upon the recent development in kernelized IV models [7, 21]. We employ a Gaussian process (GP) prior and construct a quasi-likelihood using a kernel conditional expectation estimator. We establish theoretical properties of the resultant quasi-posterior, proving its consistency and showing that it may quantify instrument strength. Furthermore, inspired from the minimax formulation of IV estimation [5, 8, 21, 22], we design a principled approximate inference algorithm using random feature expansion and a novel adaptation of the "randomized prior trick" [23]. The algorithm has the form of stochastic gradient descent-ascent and is thus scalable and easy to implement. It can also be adapted to work with flexible neural network models, in which case its behavior can be formally justified by analyzing the neural networks in the kernel regime. Empirical evaluation shows that the proposed method produces informative uncertainty estimates, scales to high-dimensional and complex nonlinear problems, and is especially advantageous when the instrument is weak.

The rest of the paper is organized as follows: In Section 2 we set up the problem. We then derive the quasi-posterior and analyze its theoretical properties in Section 3, and present the approximate inference algorithm in Section 4. Section 5 reviews related work, and Section 6 presents numerical studies. Finally, we discuss conclusion and future work in Section 7.

2 Notations and Setup

Notations   We use boldface $(\mathbf{x}, \mathbf{y}, \mathbf{z})$ to represent random variables on the space $\mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$, regular font (x, y, z) to denote deterministic values. $[n]$ denotes the set $\{1, 2, \cdots, n\}$. $\{(x_i, y_i, z_i) : i \in [n]\}$ indicates the training data. We use the notations $X := (x_1, \ldots, x_n) \in \mathcal{X}^n$, $f(X) := (f(x_1), \ldots, f(x_n))$; likewise for Y, Z. For finite-dimensional vectors $\theta, \theta' \in \mathbb{R}^m$, we use $\|\theta\|_2, \langle \theta, \theta' \rangle_2$ to denote the Euclidean norm and inner product, respectively. For any operator $A : H_1 \to H_2$ between Hilbert spaces H_1 and H_2, we denote its adjoint by $A^* : H_2 \to H_1$. When $H_1 = H_2$ and $\lambda \in \mathbb{R}$, we use the notation $A_\lambda := A + \lambda I$ for simplicity.

Instrumental variable regression   Denote the treatment and response variables as $\mathbf{x}, \mathbf{y}$, and the IV as $\mathbf{z}$. Let the true *structural function* of interest be $f^\dagger$. We consider the data generating process $\mathbf{y} = f^\dagger(\mathbf{x}) + \mathbf{u}$, $\mathbb{E}(\mathbf{u} \mid \mathbf{z}) = 0$, where $\mathbf{u}$ is unobserved and correlated with $\mathbf{x}$. Hence $f^\dagger$ satisfies

$$\mathbb{E}(\mathbf{y} - f^\dagger(\mathbf{x}) \mid \mathbf{z}) = 0 \text{ a.s. } [p(\mathbf{z})], \tag{1}$$

where p denotes the data distribution. (1) is a continuum version of generalized moment constraints on the structural function $f^\dagger$, and any f in our hypothesis space that satisfies these constraints is considered as a solution to the IV regression problem. This is the standard definition in the nonparametric IV literature (e.g., [24, 19]), and is also used in recent work on machine learning models for IV (e.g., [7, 25]). Note that it does not place any structural constraint on the conditional distribution $p(\mathbf{x} \mid \mathbf{z})$, such as additive noise. As discussed in [4], the setup can also be extended to incorporate observed confounders $\mathbf{v}$, by including $\mathbf{v}$ in both $\mathbf{x}$ and $\mathbf{z}$.

Kernelized IV and a dual view   Let $\mathcal{H}, \mathcal{I}$ be suitably chosen function spaces on $\mathcal{X}, \mathcal{Z}$, respectively. The generalized moment constraint (1) motivates the use of the following objective

$$\min_{f \in \mathcal{H}} \quad \mathcal{L}(f) := d_n^2(\hat{E}_n f - \hat{b}) + \bar{\lambda} \Omega(f), \tag{2}$$

where $\hat{E}_n : \mathcal{H} \to \mathcal{I}$ is an empirical approximation to the conditional expectation operator $E : f \mapsto \mathbb{E}(f(\mathbf{x}) \mid \mathbf{z} = \cdot)$, $\hat{b}$ is an estimator of $\mathbb{E}(\mathbf{y} \mid \mathbf{z} = \cdot)$, $\{d_n\}$ is a sequence of suitable (semi-)norm on $\mathcal{I}$, and $\Omega : \mathcal{H} \to \mathbb{R}$ is a regularization term.

When $\mathcal{H}, \mathcal{I}$ are reproducing kernel Hilbert spaces (RKHS) with corresponding kernels k_x, k_z, it is natural to set $\Omega(f) = \frac{1}{2}\|f\|_{\mathcal{H}}^2$ and define $\hat{b}$ as the result of kernel ridge regression on the response variable $\mathbf{y}$ with respect to the instrumental variable $\mathbf{z}$. In this case $\hat{E}_n$ can be defined with the empirical *kernel conditional expectation* operator: let $C_{zx} = \mathbb{E}(k(\mathbf{x}, \cdot) \otimes k(\mathbf{z}, \cdot)), C_{zz} = \mathbb{E}(k(\mathbf{z}, \cdot) \otimes k(\mathbf{z}, \cdot))$. Assuming E maps all $f \in \mathcal{H}$ to $Ef \in \mathcal{I}$, we have [26]

$$C_{zz}Ef = C_{zx}f, \quad \forall f \in \mathcal{H}.$$

This motivates the use of $\hat{E}_n := \hat{C}_{zz,\bar{\nu}}^{-1}\hat{C}_{zx}$ where $\hat{C}_{zz} := \frac{1}{n}\sum_{i=1}^n k(z_i, \cdot) \otimes k(z_i, \cdot)$, and $\hat{C}_{zx}$ is defined similarly using the empirical data distribution.[1] The choice of d_n is flexible; the dual IV formulation uses $d_n^2(g) := \frac{1}{2n}\sum_{j=1}^n g(z_j)^2 + \frac{\bar{\nu}}{2}\|g\|_{\mathcal{I}}^2 = \frac{1}{2}\langle g, \hat{C}_{zz,\bar{\nu}}g\rangle_{\mathcal{I}}$. Introducing the evaluation operator $S_z : \mathcal{I} \to \mathbb{R}^n, S_z g := (g(z_1), \ldots, g(z_n))$, we have

$$\mathcal{L}(f) = \frac{1}{2}\left\| \hat{C}_{zz,\bar{\nu}}^{-1/2}\left(\hat{C}_{zx}f - \frac{S_z^* Y}{n}\right)\right\|_{\mathcal{I}}^2 + \frac{\bar{\lambda}}{2}\|f\|_{\mathcal{H}}^2 \tag{3}$$

$$= \max_{g \in \mathcal{I}} \frac{1}{n}\sum_{j=1}^n \left((f(x_j) - y_j)g(z_j) - \frac{g(z_j)^2}{2}\right) - \frac{\bar{\nu}}{2}\|g\|_{\mathcal{I}}^2 + \frac{\bar{\lambda}}{2}\|f\|_{\mathcal{H}}^2, \tag{4}$$

where (4) holds because of the Fenchel duality $\frac{1}{2}u^2 = \sup_{v \in \mathbb{R}}\left(uv - \frac{1}{2}v^2\right)$ and the equality $\mathbb{E}(u(\mathbf{x}, \mathbf{y})g(\mathbf{z})) = \mathbb{E}(\mathbb{E}(u(\mathbf{x}, \mathbf{y}) \mid \mathbf{z})g(\mathbf{z}))$ [27, 8, 22, 25]. The dual formulation (4) circumvents the need to compute $\hat{E}_n$ directly, an operator between the typically infinite-dimensional spaces $\mathcal{H}$ and $\mathcal{I}$, and leads to a scalable estimation procedure based on stochastic gradient descent-ascent (SGDA). It can also be generalized to work with deep neural networks (DNNs) instead of kernel machines, by replacing the RKHS regularizer with a suitable regularizer for DNNs, although theoretical analysis for the resulted algorithm requires separate effort [8, 25]. As we shall see, the formulation (4) will also enable the construction of a scalable approximate inference algorithm.

3 Quasi-Bayesian Analysis of Dual IV

Introducing the notations $S_x : \mathcal{H} \to \mathbb{R}^n, S_x f := (f(x_1), \ldots, f(x_n))$, so that $\hat{C}_{zx} = \frac{1}{n}S_z^* S_x, \hat{C}_{zz} = \frac{1}{n}S_z^* S_z$, and $\lambda := n\bar{\lambda}$, we can re-express (3) in an equivalent form as:

$$\bar{\mathcal{L}}(f) := \frac{n}{\lambda}\mathcal{L}(f) = \frac{1}{2}(f(X) - Y)^\top (\lambda^{-1}L)(f(X) - Y) + \frac{1}{2}\|f\|_{\mathcal{H}}^2, \tag{5}$$

where $L = \frac{1}{n}S_z \hat{C}_{zz,\bar{\nu}}^{-1}S_z^*$ is a linear map from $\mathbb{R}^n$ to $\mathbb{R}^n$, and thus can be identified with an $n \times n$ matrix. Since the first term above is equivalent to the log density of the multivariate normal distribution $\mathcal{N}(Y \mid f(X), \lambda L^{-1})$, we can view (5) as the objective of a kernel ridge regression problem, which has a data-dependent noise covariance λL^{-1}.[2] The connection between kernel ridge regression and Gaussian process regression [29] thus motivates the use of the *quasi-posterior* $\Pi(df \mid \mathcal{D}^{(n)})$, defined through the following Radon-Nikodym derivative with respect to the standard Gaussian process prior $\Pi = \mathcal{GP}(0, k_x)$:

$$\frac{d\Pi(\cdot \mid \mathcal{D}^{(n)})}{d\Pi}(f) \propto \exp\left(-\frac{1}{2}(f(X) - Y)^\top (\lambda^{-1}L)(f(X) - Y)\right) = \exp\left(-\frac{n}{\lambda}d_n^2(\hat{E}_n f - \hat{b})\right). \tag{6}$$

Note that contrary to standard Bayesian modeling, we do not assume $Y \sim \mathcal{N}(f(X), \lambda L^{-1})$ is part of the true data generating process, nor does the theoretical analysis below rely on it. Instead, the quasi-posterior (6) should be interpreted as a Gibbs distribution which trades off between the

---

[1]With an abuse of notation, k refers to the reproducing kernel of the corresponding RKHS (k_x for $\mathcal{H}$ or k_z for $\mathcal{I}$) whenever the denotation is clear.

[2]Here we assume the invertibility of L for brevity. Alternatively, observe that (5) defines a linear inverse problem with the finite-dimensional observation operator $f \mapsto \sqrt{L}f(X)$ and noise variance $\lambda^{-1}I$. Thus we can follow [28, Chapter 6] to derive the same quasi-posterior.

properly scaled *evidence* $\lambda^{-1} n d_n^2(\hat{E}_n f - \hat{b})$, which characterizes the estimated violation of the GMM constraint (1), and our *prior belief* $\Pi(df)$. This trade-off is most clear from the well-known variational characterization of the Gibbs distribution [30],

$$\Pi(\cdot \mid \mathcal{D}^{(n)}) = \arg\min_{\Psi} \mathbb{E}_{f \sim \Psi}[\lambda^{-1} n d_n^2(\hat{E}_n f - \hat{b})] + \mathrm{KL}(\Psi \parallel \Pi). \tag{7}$$

Nonetheless, the fictitious data generating process $f \sim \mathcal{GP}(0, k), Y \sim \mathcal{N}(f(X), \lambda L^{-1})$ is useful for deriving the quasi-posterior, since its conditional distribution $p_{\mathrm{fic}}(df \mid Y)$ coincides with (6) [28, Chapter 6]. In the probability space of this fictitious data generating process, for any finite set of test inputs x_*, we have

$$p_{\mathrm{fic}}(f(x_*), Y) \sim \mathcal{N}\left(0, \begin{bmatrix} K_{**} & K_{*x} \\ K_{x*} & K_{xx} + \lambda L^{-1} \end{bmatrix}\right),$$

where $K_{(\cdot)}$ denote the corresponding Gram matrices with subscript $*$ denoting x_* and x denoting X (so, e.g., $K_{*x} := k(x_*, X)$). Thus by the Gaussian conditioning formula, we have

$$\Pi(f(x_*) \mid \mathcal{D}^{(n)}) = p_{\mathrm{fic}}(f(x_*) \mid Y) = \mathcal{N}(m, S), \tag{8}$$

$$\text{where} \quad m := K_{*x}(\lambda I + L K_{xx})^{-1} L Y, \tag{9}$$

$$S := K_{**} - K_{*x} L(\lambda I + K_{xx} L)^{-1} K_{x*}, \tag{10}$$

$$L = K_{zz}(K_{zz} + \nu I)^{-1}. \tag{11}$$

In the above $K_{zz} := k(Z, Z)$ denotes the Gram matrix, and (11) follows from the Woodbury identity.

Theoretical Analysis Intuitively, for the quasi-posterior $\Pi(\cdot \mid \mathcal{D}^{(n)})$ to be a useful measure of uncertainty, it needs to satisfy the following informal criteria: as $n \to \infty$, we expect

(C1) $\Pi(\cdot \mid \mathcal{D}^{(n)})$ will exclude incorrect solutions;

(C2) In cases of non-identification, $\Pi(\cdot \mid \mathcal{D}^{(n)})$ will not exclude any valid solution in the hypothesis space.

In the following, we formalize these criteria, and demonstrate that with appropriately chosen hyperparameters, the quasi-posterior satisfies both criteria. We will work with the following assumptions:

Assumption 3.1. *The restriction of the conditional expectation operator $f \mapsto \mathbb{E}(f(\mathbf{x}) \mid \mathbf{z} = \cdot)$ on $\mathcal{H}$, denoted as E, has its image contained in $\mathcal{I}$. Moreover, $E : \mathcal{H} \to \mathcal{I}$ is bounded.*

Assumption 3.1 is intuitive. It is also slightly more general than some previous work [21, 25] that require the conditional expectation operator to be bounded on the entire hypothesis space, which typically corresponds to the "sample space" of the Gaussian process prior in our setting, and is much larger than $\mathcal{H}$ (see Appendix A). Nonetheless, we note that Hypothesis 4 in Singh et al. [7] may be more general, although they impose extra smoothness assumptions on E.

Assumption 3.2. *The true structural function $f^\dagger(x)$ is such that, there exists a sequence $\{\tau_m : m \in \mathbb{N}\}$ satisfying $\tau_m \to 0$, and*

$$m\tau_m^2 \geq \inf_{f_m^\dagger \in \mathcal{H}: \|f^\dagger - f_m^\dagger\| \leq \tau_m} \|f_m^\dagger\|_{\mathcal{H}}^2 - \log \Pi(\{f : \|f\| \leq \tau_m\}),$$

where $\|\cdot\|$ denotes the sup norm.

Assumption 3.2 requires that $f^\dagger$ can be well approximated in $\mathcal{H}$; this is more general than previous work [e.g., 21, 7] that require $f^\dagger \in \mathcal{H}$. The sequence $\{\tau_m\}$ is determined by the complexity of $\mathcal{H}$ and its ability for approximating $f^\dagger$, and is usually the optimal posterior contraction rate for Gaussian process regression on the unconfounded dataset $\{(x_i, f^\dagger(x_i) + \epsilon_i) : i \in [m]\}$ [31, 32].

Assumption 3.3. *$\mathcal{X}$ and $\mathcal{Z}$ are Polish spaces. The kernels k_x, k_z are continuous, $\sup_{x \in \mathcal{X}} k(x, x) + \sup_{z \in \mathcal{Z}} k(z, z) \leq \kappa^2$, and Mercer's representations [33] of k_x and k_z exist. The random variable $\mathbf{y} - f^\dagger(\mathbf{x})$ is sub-exponential.*

The conditions in Assumption 3.3 are technical and frequently assumed in literature [e.g., 34]. The requirements on the kernels can be satisfied by e.g. continuous kernels on compact subsets of $\mathbb{R}^d$.

Given the assumptions above, we characterize (C1) by showing that as $n \to \infty$, the posterior places vanishing mass on the region of functions that violates the GMM constraints (1). Concretely,

Theorem 3.1 (Proof in Appendix B). *There exist a constant $M > 0$ depending on the data distribution and the kernels of choice, and a sufficiently slowly growing sequence $\{\gamma_n\} \to \infty$ (e.g., we can always have $\gamma_n \leq \log \log n$) such that when taking $\bar{\lambda} = n^{-1/2}, \bar{\nu} = \min\{\tau^2_{\sqrt{n}}, n^{-1/2}\gamma_n\}$, it has*

$$\mathbb{E}_{\mathcal{D}^{(n)}}\Pi(\{f : \mathbb{E}^2(f(\mathbf{x}) - \mathbf{y} \mid \mathbf{z}) > M\tau^2_{\sqrt{n}}\} \mid \mathcal{D}^{(n)}) \to 0. \tag{12}$$

Remark 3.1. The above result provides a posterior contraction rate [35] in the order of $\tau_{\sqrt{n}}$ for the reduced form $\mathbb{E}(f(\mathbf{x}) \mid \mathbf{z})$ where $\{\tau_m\}$ is defined in Assumption 3.2. While the analysis of IV regression is obviously more difficult than that for least square regression, our analysis here is somewhat crude, and should be viewed as upper bounding the performance of our method. The rate is also likely improvable if we switch to using separate sets of samples for estimation of $\hat{E}_n$ and inference on f, with different sample sizes, as in [7]. We leave a refined analysis as future work.

Nonetheless, the contraction rate established here is often informative: for example, suppose that, roughly speaking, the regularity of $f^\dagger$ is similar to functions in the Matérn-$1/2$ RKHS. This is a very basic requirement under our assumption, and for suitable choice of the kernels, we can have $\tau^2_{\sqrt{n}} = O(n^{-1/4})$; see Remark A.3 for details. Moreover, assuming identifiability of $f^\dagger$, the theorem always implies the consistency of the quasi-posterior.

Remark 3.2. While the use of an increasing $\lambda = n\bar{\lambda}$ might be strange from a Bayesian view, it is common to impose extra regularization in nonparametric IV analysis, due to the need to estimate E from data: [19, 18] regularize by using priors with increasing complexity; previous work in kernelized IV [7, 21, 25] also uses $\bar{\lambda} \gg n^{-1}$. However, while a slowly growing λ may be necessary, the choice of $\lambda = \sqrt{n}$ is likely conservative. In practice, our hyperparameter selection procedure chooses λ with slower growth.

The following proposition characterizes (C2):

Proposition 3.1 (Proof in Appendix B). *For almost every $f \sim \Pi$, the scaled log quasi-likelihood estimates converge in probability to the true log quasi-likelihood, i.e.,*

$$\frac{1}{n}\sum_{i=1}^{n}\hat{E}_n(f(\mathbf{x}) - \mathbf{y} \mid \mathbf{z} = z_i)^2 \xrightarrow{p} \mathbb{E}(\mathbb{E}(f(\mathbf{x}) - \mathbf{y} \mid \mathbf{z})^2).$$

Remark 3.3. The proposition characterizes (C2), since when there are multiple f that satisfy (1), all of them will eventually have significantly higher log likelihood than functions violating (1): the difference is $\Theta(\lambda^{-1}n)$. Interpolating from such non-identified settings to scenarios where the IV is more informative, we can also see that this result suggests the quasi-posterior reflects IV strength.

Remark 3.4. To see that in the weak IV setting, the behavior of the quasi-posterior may be qualitatively different from that of bootstrap, we can also consider the following extreme example: suppose $\mathbf{z}$ is completely non-informative so that $\mathbb{E}(f(\mathbf{x}) \mid \mathbf{z}) \equiv \mathbb{E}f(\mathbf{x})$ for all f; and suppose that the estimated conditional expectation $\hat{E}_n$ is sufficiently accurate, so that we replace it with E.[3] In this case bootstrap on (3) will always return the point estimator $f \equiv 0$ due to the regularization on $\|f\|_{\mathcal{H}}$, while the quasi-posterior behaves like the prior, correctly reflecting the complete lack of evidence in data. While the example is clearly oversimplified, and in practice the estimation error of E may play an important role, it is known that bootstrap uncertainty estimates can be problematic given weak IVs [36–38]. Moreover, we observe similar failures for bootstrap in experiments (see Section 6.1).

4 Scalable Approximate Inference via a Randomized Prior Trick

We now turn to approximate inference with parametric models such as random feature expansion or wide NNs. Scalable inference for the IV quasi-posterior appears difficult, since for any f, computing the quasi-likelihood involves computing $\hat{E}_n f$, which in turn requires either inverting an $n \times n$ Gram matrix, judging from (5) and (11), or solving an optimization problem specific to f from (4).

---

[3]This setting could be realistic in the sample splitting setup considered in [7], where we observe separate sets of samples $(Z^{(1)}, X^{(1)})$ and $(X^{(2)}, Y^{(2)})$, and the sample size for the first stage (i.e., estimation of $\hat{E}_n$) is sufficiently large. Note that as long as we have finite samples for the second stage (i.e., estimation of f), we still need a non-zero regularizer $\bar{\lambda} > 0$ on f.

Nonetheless, we show that it is possible, by extending the "randomized prior" trick for Gaussian process regression [23] to work with (quasi-)likelihoods with an optimization formulation as in (4).

Our algorithm works with random feature models. A random feature model for k_z approximates $k_z(z, z') \approx \tilde{k}_{z,m}(z, z') := \frac{1}{m}\phi_{z,m}(z)^\top \phi_{z,m}(z')$, where $\phi_{z,m}$ takes value in $\mathbb{R}^m$. Then the map $\varphi \mapsto \frac{1}{\sqrt{m}}\varphi^\top \phi_{z,m}(\cdot) =: g(\cdot; \varphi)$ parameterizes an approximate RKHS $\tilde{\mathcal{H}}$; and for all $c > 0$, the random function $g(\cdot; \varphi)$, where $\varphi \sim \mathcal{N}(0, cI)$, is distributed as $\mathcal{GP}(0, c\tilde{k}_{z,m})$. The notations related to k_x are similar and thus omitted. Now we can state the objective function:

Proposition 4.1 (Proof in Appendix C.1)**.** *Let $\phi_0 \sim \mathcal{N}(0, \lambda\nu^{-1}I), \theta_0 \sim \mathcal{N}(0, I), \tilde{y}_i \sim \mathcal{N}(y_i, \lambda)$.* *Then the optima θ^* of*

$$\min_{\theta \in \mathbb{R}^m} \max_{\phi \in \mathbb{R}^m} \sum_{i=1}^{n} \left((f(x_i; \theta) - \tilde{y}_i)g(z_i; \phi) - \frac{g(z_i; \phi)^2}{2} \right) - \frac{\nu}{2}\|\phi - \phi_0\|_2^2 + \frac{\lambda}{2}\|\theta - \theta_0\|_2^2 \quad (13)$$

parameterizes a random function which follows the quasi-posterior distribution (6)*, where the kernels are replaced by the random feature approximations.*

Given the above proposition, we can sample from the random feature-approximated quasi-posterior by solving (13) with stochastic gradient descent-ascent; the approximation errors will be analyzed in the following. The objective (13) is closely related to (4); as we show in Appendix C.1.1, it is equivalent to

$$\min_{f \in \tilde{\mathcal{H}}} \max_{g \in \tilde{\mathcal{I}}} \sum_{i=1}^{n} \left((f(x_i) - \tilde{y}_i)g(z_i) - \frac{g(z_i)^2}{2} \right) - \frac{\nu}{2}\|g - g_0\|_{\tilde{\mathcal{I}}}^2 + \frac{\lambda}{2}\|f - f_0\|_{\tilde{\mathcal{H}}}^2, \quad (14)$$

which differs from (4) only in the regularizers: instead of regularizing the norm of f and g, (14) encourages the functions to stay close to randomly sampled *anchors* [39]. Alternatively, we can view (14) as *perturbing* the point estimator (4), so that it has a covariance matching that of the quasi-posterior. A similar relation is also observed in the original randomized prior trick, which transforms GP regression to the optimization problem $\min_{f \in \tilde{\mathcal{H}}} \sum_{i=1}^{n}(f(x_i) - \tilde{y}_i)^2 + \lambda\|f - f_0\|_2^2$. In both cases, the resultant algorithm for approximate inference has the same time complexity as ensemble training for point estimation.

While the algorithm can be directly applied to neural network models as in [23], we follow [40] and modify the objective, to account for the difference between the neural tangent kernel (NTK) [41] of a wide neural network architecture and the NNGP kernel of the corresponding infinite-width Bayesian neural network [42–44]. Concretely, we modify (13) as

$$\min_{\theta} \max_{\phi} \sum_{i=1}^{n} \left((\tilde{f}_\theta(x_i) - \tilde{y}_i)\tilde{g}_\phi(z_i) - \frac{\tilde{g}_\phi(z_i)^2}{2} \right) - \frac{\nu}{2}\|\phi - \phi_0\|_2^2 + \frac{\lambda}{2}\|\theta - \theta_0\|_2^2, \quad (15)$$

$$\text{where } \tilde{g}_\phi(z) := g(z; \phi) - g(z; \phi_0) + \tilde{g}_0(z), \ \tilde{g}_0(z) := \sqrt{\frac{\lambda}{\nu}}\langle \bar{\phi}_0, \frac{\partial g}{\partial \phi}|_{\phi=\phi_0}(z)\rangle,$$

and ϕ_0 denotes the initial value of ϕ, and $\bar{\phi}_0 \sim \mathcal{N}(0, I)$ is a set of randomly initialized NN parameters independent of ϕ_0; and $\tilde{f}_\theta$ is defined similarly.

We only give a formal justification for the modification, *under the assumption*[4] that the NNs remain in the kernel regime throughout training, so that $g(z; \phi) - g(z; \phi_0) = \langle \phi - \phi_0, \frac{\partial g(z)}{\partial \phi}|_{\phi_0}\rangle_2$ [47]. Thus for the purpose of analyzing $g(\cdot; \phi) - g(\cdot; \phi_0)$, we can view g as a random feature model with the parameterization $\phi \mapsto \langle \phi, \frac{\partial g(z)}{\partial \phi}|_{\phi_0}\rangle_2$. Thus by the argument in Appendix C.1.1, we can show that the weight regularizer $\|\phi - \phi_0\|_2$ is equivalent to $\|g(\cdot; \phi) - g(\cdot; \phi_0)\|_{\tilde{\mathcal{I}}} = \|\tilde{g}_\phi - \tilde{g}_0\|_{\tilde{\mathcal{I}}}$, where $\tilde{\mathcal{I}}$ is determined by the NTK $k_{g,ntk}(z, z') := \langle \frac{\partial g(z)}{\partial \phi}|_{\phi_0}, \frac{\partial g(z')}{\partial \phi}|_{\phi_0}\rangle_2$. Similar arguments can be made for $\tilde{f}_\theta$ and $\tilde{f}_0$. Consequently, (15) is equivalent to an instance of (14) with $\tilde{\mathcal{H}}, \tilde{\mathcal{I}}$ defined by the NTKs.

Implementation details for the algorithm, including hyperparameter selection, are discussed in Appendix D.

---

[4]The same linearization assumption has been employed in [40, 45]. We refer readers to [46, 22] for analysis of the linearization error in various settings similar to ours.

Convergence analysis   We now complete the analysis of the inference algorithm, by showing that for any fixed set of test points x_*, SGDA can approximate the marginal distribution $\Pi(f(x_*) \mid \mathcal{D}^{(n)})$ arbitrarily well given a sufficient computational budget. This implies that the approximate posterior is good enough for prediction purposes.

We place several mild assumptions on the random feature model, listed in Appendix C.2; they are satisfied by common approximations such as the random Fourier features [48]. The SGDA algorithm is described in detail in Appendix C.4. Under this setup, we have

Proposition 4.2 (Proof in Appendix C.5). *Fix the training data $\mathcal{D}^{(n)}$ and hyperparameters $\lambda, \nu > 0$. Then there exist a sequence of choices of m and SGDA step-size schemes, such that for any $l \in \mathbb{N}$, we have*

$$\sup_{x^* \in \mathcal{X}^l} \max(\|\hat{\mu}_m - \mu_m\|_2, \|\hat{S}_m - S_m\|_F) \xrightarrow{p} 0.$$

In the above, $\hat{\mu}_m, \hat{S}_m$ denote the mean and covariance of the approximate marginal posterior for $f(x_*)$, μ, S correspond to the true posterior, $\|\cdot\|_F$ denotes the Frobenius norm, and the convergence in probability is defined with respect to the sampling of random feature basis.

5   Related Work

Quasi-Bayesian analysis for GMM estimation problems was first developed in [49, 50, 17], which provided theoretical analysis for parametric models. The use of the quasi-posterior is motivated from the maximum entropy principle, based on which similar ideas have been developed in the machine learning literature [51–53]. [18, 19] first analyzed the use of nonparametric models for quasi-Bayesian IV; no numerical study was presented. Closer to our work is [20] which constructed a quasi-posterior using Nadaraya-Watson smoothing as $\hat{E}_n$. While similar in form to our method, the smoothing approach relies on stronger assumptions, especially for high-dimensional data; see the discussion in [7, Appendix A.2.1] which compared the corresponding point estimators. Their approach is also harder to scale to large datasets.

Our quasi-Bayesian procedure builds upon the kernelized IV models [7, 21] and the dual formulation of IV regression [6, 21, 22, 25]. The kernelized dual IV formulation (3)-(4) is from [21, 25]. [7] proposes a similar method; although it was motivated differently as a kernelized two-stage least squares (2SLS) estimator, its objective is asymptotically equivalent to (3); however, the slight difference in regularization prevents the use of estimation procedures similar to (4), as we discuss in Appendix C.1.4. Other recent work on nonlinear IV include [4, 54–56]. It remains interesting future work to develop scalable quasi-Bayesian procedures for these methods, although we note that the mean estimator derived from our quasi-posterior implementation has competitive performance.

Alternative approaches for uncertainty quantification include Bayesian inference and bootstrap. We have discussed previous works on Bayesian IV and their limitations in Section 1. Bootstrap is typically justified in the asymptotic regime, which does not cover many scenarios where uncertainty is most needed; this is different from the quasi-Bayesian approach which can always be justified through (7). Moreover, standard bootstrap inference on 2SLS is known to be unreliable when the instrument strength is weak [36–38]; while remedies exist (e.g., [38]), they heavily rely on the linearity and additive noise assumptions, and are thus difficult to generalize to the nonlinear setting we are interested in. As the kernelized IV methods generalize 2SLS, we expect similar issues to exist in our setting.

6   Experiments

6.1   1D Simulation

We first experiment on a variety of 1D synthetic datasets, constructed by modifying the setup in [5] to incorporate a nonlinear first stage, in a way similar to [7, 57]:

$$z := \text{sigmoid}(w), \quad x := \text{sigmoid}\left(\frac{\alpha w + (1 - \alpha)u'}{\sqrt{\alpha^2 + (1 - \alpha)^2}}\right), \quad y_i \sim \mathcal{N}(f_0(2x - 1) + 2u, 0.1),$$

where (u, u') are normal random variables with unit variance and a correlation of 0.5, $w \sim \mathcal{N}(0, 1)$ is independent of (u, u'), α is a parameter controlling the instrument strength, and f_0 is constructed from the `sine`, `step`, `abs` or a `linear` function. We choose $N \in \{200, 1000\}$ and $\alpha \in \{0.05, 0.5\}$.

267 Our baselines include BayesIV [16], a state-of-the-art Bayesian model based on B-splines and
268 Dirichlet process mixture; we also include bootstrap on 2SLS with ridge regularization, either ap-
269 plied directly to the input features (Linear), on their polynomial expansion (Poly), or on the same
270 kernelized models (KIV) as ours.[5] Hyperparameter for the kernelized IV methods are selected by
271 cross validation based on the observable first-stage and second-stage losses as in previous work
272 [7, 21]; see Appendix D.1. For kernels we choose the RBF and Matérn kernels, although results for
273 Matérn kernels are deferred to appendix for brevity. See Appendix D.3 for the detailed setup.

Figure 1: Test MSE and CI coverage on the `sine` dataset. The left two plots correspond to $\alpha = 0.05$, while the right two correspond to $\alpha = 0.5$. `bs` denotes bootstrap, `qb` denotes quasi-Bayesian.

274 Normalized MSE and coverage rate of 95% credible intervals (CI) on the `sine` datasets are plotted in
275 Figure 1. We report results on 10 independently generated datasets. As we can see, quasi-Bayesian
276 inference provides the most reliable uncertainty estimates, and is especially advantageous in the
277 weak IV setting ($\alpha = 0.05$). While its CI can be conservative, we note that it is still informative,
278 and properly reflects the sample size and instrument strength; see Appendix D.4 for visualizations.
279 These results connect to previous work showing (in a different setting) that the radius of credible set
280 produced by a GP posterior can have the correct order of magnitude [58].

281 Full results on all datasets, visualizations and additional experiments are deferred to Appendix D.4.
282 As a summary, (i) on the `abs` and `linear` datasets where all modeling assumptions hold, the results
283 are qualitatively similar to the sin dataset. Moreover, the over-smoothed RBF kernel appears to have
284 similar coverage comparing with the optimal kernel, and follow a similar contraction rate, as the pre-
285 vious work on GP regression [59] suggests. (ii) On the `step` dataset which violates Assumption 3.2,
286 the quasi-posterior still provides more coverage than the baselines. (iii) Uncertainty estimates pro-
287 duced by the approximate inference algorithm are similar to that from the exact quasi-posterior.

6.2 Airline Demand

We now turn to the more challenging demand simulation first proposed by [4]. The dataset simulates
a scenario where we need to predict the demand of airline tickets y, as a function of the price x, and
two observed confounders: customer type s, and time of year t. The data generating process is

$$x := (z + 3)\psi(t) + 25 + u', \;\; y := f_0(x, t, s) + u, \;\; f_0(x, t, s) := 100 + (10 + x) \cdot s \cdot \psi(t) - 2x$$

289 where (u, u') are standard normal variables with correlation ρ, $z \sim \mathcal{N}(0, 1)$ is independent of
290 (u, u'), and ψ is a nonlinear function whose shape is given in Figure 2. The variable s either
291 varies across $\{0, \ldots, 6\}$ (the lower-dimensional setting), or is observed as an MNIST image repre-
292 senting the corresponding digit; the latter case represents the real-world scenario where only high-
293 dimensional surrogates of the true confounder is observed. We only report results for $\rho = 0.5$,
294 noting that results using other choices of ρ have been similar. We use $n \in \{1000, 10000\}$ for the
295 lower-dimensional setting, and use $n = 50000$ for the image-based setting.

296 We compare our method with bootstrap on the same model, BayesIV, and bootstrap on linear or
297 polynomial models. Performance of other point estimators on this dataset has been reported in
298 [7, 21, 56], compared with which our method is generally competitive. We implement the dual IV
299 model using both an RBF kernel and DNN models. See Appendix D.5 for details.

300 We report the test MSE and CI coverage for $N = 1000$ in Table 1, and visualize all approxi-
301 mate quasi-posteriors for the NN models in Figure 2. As we can see, when implemented with
302 DNNs, our method produces uncertainty estimates with excellent coverage, which also correctly

---

[5] We do not compare with [20] since their source code is unavailable. Note that their smoothing-based method does not scale to large datasets, and there is no numerical study on the credible interval in [20].

(a) lower-dim, $N = 10^3$ (b) lower-dim, $N = 10^4$ (c) image setup

Figure 2: Approximate quasi-posteriors in the demand simulation. We plot a cross-section by fixing s, x to their mean values and varying t.

reflects the information available in the dataset: the CI is wider when N is smaller, or in the high-dimensional experiment where estimation is harder. Bootstrap has a noticeably worse performance when $N = 1000$. Still, it performs well in the (arguably less interesting) large-sample setting, with a CI coverage similar to our method; see Appendix D.6. This is because on this dataset, the total instrument strength is stronger due to the presence of observed confounders, and the NN model is a good fit. Consequently, the asymptotic behavior of bootstrap can be observed when N is large.

Both methods have poorer performances when we switch to the RBF kernel, although the quasi-posterior is still more reliable. It is natural that uncertainty estimates are only as good as the model allows; the inflexibility of the fixed-form RBF kernel affects the contraction rate, through both τ_m and the kernel-dependent constant M. For the latter, we can see from the proof that M depends on the operator norm $\|E\|$, which reflects the difficulty of conditional expectation estimation using the models of choice. These results highlight the importance of using flexible NN models, which our inference algorithm naturally supports.

The other baselines perform poorly due to their inflexibility; in particular, note that BayesIV uses additive models for both stages (e.g., $f(x, t, s) = f_1(x) + f_2(t) + f_3(s)$) which do not approximate this data generating process well. Full results and visualizations are deferred to Appendix D.6.

Table 1: Test normalized MSE and CI coverage on the demand dataset with $N = 1000$, averaged over 10 trials. **Boldface** indicates the best result.

Method	BS-Linear	BS-Poly	BayesIV	BS-RBF	QB-RBF	BS-NN	QB-NN
NMSE	$.37 \pm .01$	$.31 \pm .06$	$.28 \pm .04$	$.17 \pm .01$	$.17 \pm .01$	$.06 \pm .03$	$\mathbf{.04 \pm .00}$
CI Cvg.	$.09 \pm .01$	$.15 \pm .03$	$.27 \pm .06$	$.45 \pm .02$	$.77 \pm .02$	$.85 \pm .02$	$\mathbf{.94 \pm .01}$

7   Conclusion

In this work we propose a scalable quasi-Bayesian procedure for IV regression. We analyze the theoretical properties of the proposed quasi-posterior, and derive a scalable algorithm for approximate inference. Empirical evaluations show that the proposed method scales to large and high-dimensional datasets, and can be particularly advantageous when the instrument strength is weak.

Beyond IV regression, formulations like (1) also appear in various other problems in causal inference and statistics, as discussed in [22]; our method can be readily applied to these problems. Future work includes extension to more general conditional moment restriction problems with nonlinear constraints [60, 18, 21], and refining the theoretical analysis of the quasi-posterior.

Broader Impact

IV analysis is widely used in social science and clinical research, where it can be highly undesirable to have a systematic error or increased prediction variance on subpopulations which are underrepresented in the training dataset, or on which the instrument is locally weak. Uncertainty quantification is an important first step in detecting such issues, and the quasi-Bayesian approach we take can be preferable since it has a clear interpretation in a non-asymptotic setting, and when the instrument strength is weak. However, epistemic uncertainty estimates are only as good as the model allows, and it should not create a false sense of complete security. Caution must be taken when employing complex models in scenarios with potential fairness implications.

References

[1] J. D. Angrist and J.-S. Pischke, *Mostly harmless econometrics: An empiricist's companion.* Princeton university press, 2008.

[2] S. Greenland, "An introduction to instrumental variables for epidemiologists," *International Journal of Epidemiology*, vol. 29, no. 4, pp. 722–729, 2000.

[3] J. Cuzick, R. Edwards, and N. Segnan, "Adjusting for non-compliance and contamination in randomized clinical trials," *Statistics in Medicine*, vol. 16, no. 9, pp. 1017–1029, 1997.

[4] J. Hartford, G. Lewis, K. Leyton-Brown, and M. Taddy, "Deep IV: A flexible approach for counterfactual prediction," in *International Conference on Machine Learning*. PMLR, 2017, pp. 1414–1423.

[5] G. Lewis and V. Syrgkanis, "Adversarial generalized method of moments," *arXiv preprint arXiv:1803.07164*, 2018.

[6] A. Bennett, N. Kallus, and T. Schnabel, "Deep generalized method of moments for instrumental variable analysis," *arXiv preprint arXiv:1905.12495*, 2019.

[7] R. Singh, M. Sahani, and A. Gretton, "Kernel Instrumental Variable Regression," *arXiv:1906.00232 [cs, econ, math, stat]*, Jul. 2020, arXiv: 1906.00232.

[8] A. Bennett and N. Kallus, "The Variational Method of Moments," *arXiv:2012.09422 [cs, econ, math, stat]*, Dec. 2020, arXiv: 2012.09422.

[9] J. H. Stock, J. H. Wright, and M. Yogo, "A survey of weak instruments and weak identification in generalized method of moments," *Journal of Business & Economic Statistics*, vol. 20, no. 4, pp. 518–529, 2002.

[10] X. Chen and M. Reiss, "On rate optimality for ill-posed inverse problems in econometrics," *Econometric Theory*, pp. 497–521, 2011.

[11] W. K. Newey, "Nonparametric instrumental variables estimation," *American Economic Review*, vol. 103, no. 3, pp. 550–56, 2013.

[12] F. Kleibergen and H. K. Van Dijk, "Bayesian simultaneous equations analysis using reduced rank structures," *Econometric theory*, pp. 701–743, 1998.

[13] F. Kleibergen and E. Zivot, "Bayesian and classical approaches to instrumental variable regression," *Journal of Econometrics*, vol. 114, no. 1, pp. 29–72, 2003.

[14] T. G. Conley, C. B. Hansen, R. E. McCulloch, and P. E. Rossi, "A semi-parametric Bayesian approach to the instrumental variable problem," *Journal of Econometrics*, vol. 144, no. 1, pp. 276–305, 2008.

[15] H. F. Lopes and N. G. Polson, "Bayesian instrumental variables: priors and likelihoods," *Econometric Reviews*, vol. 33, no. 1-4, pp. 100–121, 2014.

[16] M. Wiesenfarth, C. M. Hisgen, T. Kneib, and C. Cadarso-Suarez, "Bayesian nonparametric instrumental variables regression based on penalized splines and dirichlet process mixtures," *Journal of Business & Economic Statistics*, vol. 32, no. 3, pp. 468–482, 2014.

[17] V. Chernozhukov and H. Hong, "An MCMC approach to classical estimation," *Journal of Econometrics*, vol. 115, no. 2, pp. 293–346, Aug. 2003.

[18] Y. Liao and W. Jiang, "Posterior consistency of nonparametric conditional moment restricted models," *The Annals of Statistics*, vol. 39, no. 6, pp. 3003–3031, Dec. 2011.

[19] K. Kato, "Quasi-Bayesian analysis of nonparametric instrumental variables models," *The Annals of Statistics*, vol. 41, no. 5, pp. 2359–2390, Oct. 2013.

[20] J.-P. Florens and A. Simoni, "Nonparametric estimation of an instrumental regression: A quasi-Bayesian approach based on regularized posterior," *Journal of Econometrics*, vol. 170, no. 2, pp. 458–475, Oct. 2012.

[21] K. Muandet, A. Mehrjou, S. K. Lee, and A. Raj, "Dual Instrumental Variable Regression," *arXiv:1910.12358 [cs, econ, stat]*, Oct. 2020, arXiv: 1910.12358.

[22] L. Liao, Y.-L. Chen, Z. Yang, B. Dai, Z. Wang, and M. Kolar, "Provably efficient neural estimation of structural equation model: An adversarial approach," *arXiv preprint arXiv:2007.01290*, 2020.

[23] I. Osband, J. Aslanides, and A. Cassirer, "Randomized prior functions for deep reinforcement learning," in *Proceedings of the 32nd International Conference on Neural Information Processing Systems*, 2018, pp. 8626–8638.

[24] W. K. Newey and J. L. Powell, "Instrumental variable estimation of nonparametric models," *Econometrica*, vol. 71, no. 5, pp. 1565–1578, 2003.

[25] N. Dikkala, G. Lewis, L. Mackey, and V. Syrgkanis, "Minimax estimation of conditional moment models," in *Advances in Neural Information Processing Systems*, H. Larochelle, M. Ranzato, R. Hadsell, M. F. Balcan, and H. Lin, Eds., vol. 33. Curran Associates, Inc., 2020, pp. 12 248–12 262.

[26] K. Fukumizu, F. R. Bach, and A. Gretton, "Statistical Consistency of Kernel Canonical Correlation Analysis," *Journal of Machine Learning Research*, p. 23, 2004.

[27] B. Dai, N. He, Y. Pan, B. Boots, and L. Song, "Learning from Conditional Distributions via Dual Embeddings," *arXiv:1607.04579 [cs, math, stat]*, Dec. 2016, arXiv: 1607.04579.

[28] A. M. Stuart, "Inverse problems: a Bayesian perspective," *Acta numerica*, vol. 19, pp. 451–559, 2010.

[29] M. Kanagawa, P. Hennig, D. Sejdinovic, and B. K. Sriperumbudur, "Gaussian processes and kernel methods: A review on connections and equivalences," *arXiv preprint arXiv:1807.02582*, 2018.

[30] T. Zhang *et al.*, "From ϵ-entropy to KL-entropy: Analysis of minimum information complexity density estimation," *The Annals of Statistics*, vol. 34, no. 5, pp. 2180–2210, 2006.

[31] I. Castillo *et al.*, "Lower bounds for posterior rates with gaussian process priors," *Electronic Journal of Statistics*, vol. 2, pp. 1281–1299, 2008.

[32] A. W. van der Vaart and J. H. van Zanten, "Information Rates of Nonparametric Gaussian Process Regression," *Journal of Machine Learning Research*, vol. 12, pp. 2095–2119, 2011.

[33] I. Steinwart and C. Scovel, "Mercer's Theorem on General Domains: On the Interaction between Measures, Kernels, and RKHSs," *Constructive Approximation*, vol. 35, no. 3, pp. 363–417, Jun. 2012.

[34] L. H. Dicker, D. P. Foster, and D. Hsu, "Kernel ridge vs. principal component regression: Minimax bounds and the qualification of regularization operators," *Electronic Journal of Statistics*, vol. 11, no. 1, Jan. 2017.

[35] S. Ghosal and A. Van der Vaart, *Fundamentals of nonparametric Bayesian inference*. Cambridge University Press, 2017, vol. 44.

[36] M. Moreira, J. R. Porter, and G. A. Suarez, "Bootstrap and higher-order expansion validity when instruments may be weak," 2004.

[37] A. Flores-Lagunes, "Finite sample evidence of IV estimators under weak instruments," *Journal of Applied Econometrics*, vol. 22, no. 3, pp. 677–694, Apr. 2007.

[38] R. Davidson and J. G. Mackinnon, "Wild Bootstrap Tests for IV Regression," *Journal of Business & Economic Statistics*, vol. 28, no. 1, pp. 128–144, 2010, publisher: [American Statistical Association, Taylor & Francis, Ltd.].

[39] T. Pearce, F. Leibfried, and A. Brintrup, "Uncertainty in neural networks: Approximately Bayesian ensembling," in *Proceedings of the Twenty Third International Conference on Artificial Intelligence and Statistics*, ser. Proceedings of Machine Learning Research, S. Chiappa and R. Calandra, Eds., vol. 108.   PMLR, 26–28 Aug 2020, pp. 234–244.

[40] B. He, B. Lakshminarayanan, and Y. W. Teh, "Bayesian deep ensembles via the neural tangent kernel," *arXiv preprint arXiv:2007.05864*, 2020.

[41] A. Jacot, F. Gabriel, and C. Hongler, "Neural Tangent Kernel: Convergence and Generalization in Neural Networks," in *Advances in Neural Information Processing Systems*, S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, Eds., vol. 31.   Curran Associates, Inc., 2018.

[42] R. M. Neal, *Bayesian learning for neural networks*.   Springer Science & Business Media, 2012, vol. 118.

[43] J. Lee, Y. Bahri, R. Novak, S. S. Schoenholz, J. Pennington, and J. Sohl-Dickstein, "Deep neural networks as Gaussian processes," *arXiv preprint arXiv:1711.00165*, 2017.

[44] A. G. d. G. Matthews, M. Rowland, J. Hron, R. E. Turner, and Z. Ghahramani, "Gaussian process behaviour in wide deep neural networks," *arXiv preprint arXiv:1804.11271*, 2018.

[45] W. Hu, Z. Li, and D. Yu, "Simple and effective regularization methods for training on noisily labeled data with generalization guarantee," in *International Conference on Learning Representations*, 2020.

[46] T. Hu, W. Wang, C. Lin, and G. Cheng, "Regularization matters: A nonparametric perspective on overparametrized neural network," in *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics*, ser. Proceedings of Machine Learning Research, A. Banerjee and K. Fukumizu, Eds., vol. 130.   PMLR, 13–15 Apr 2021, pp. 829–837.

[47] J. Lee, L. Xiao, S. Schoenholz, Y. Bahri, R. Novak, J. Sohl-Dickstein, and J. Pennington, "Wide Neural Networks of Any Depth Evolve as Linear Models Under Gradient Descent," in *Advances in Neural Information Processing Systems*, H. Wallach, H. Larochelle, A. Beygelzimer, F. d. Alché-Buc, E. Fox, and R. Garnett, Eds., vol. 32.   Curran Associates, Inc., 2019.

[48] A. Rahimi, B. Recht *et al.*, "Random features for large-scale kernel machines." in *NIPS*, vol. 3, no. 4.   Citeseer, 2007, p. 5.

[49] A. Zellner, "Bayesian Method of Moments (BMOM) Analysis of Mean and Regression Models," *arXiv:bayes-an/9511001*, Dec. 1995, arXiv: bayes-an/9511001.

[50] J.-Y. Kim, "Limited information likelihood and Bayesian analysis," *Journal of Econometrics*, p. 19, 2002.

[51] T. Jaakkola, M. Meila, and T. Jebara, "Maximum entropy discrimination," in *Proceedings of the 12th International Conference on Neural Information Processing Systems*, 1999, pp. 470–476.

[52] M. Dudík, S. J. Phillips, and R. E. Schapire, "Maximum entropy density estimation with generalized regularization and an application to species distribution modeling," *Journal of Machine Learning Research*, 2007.

[53] J. Zhu and E. P. Xing, "Maximum entropy discrimination Markov networks." *Journal of Machine Learning Research*, vol. 10, no. 11, 2009.

[54] R. Zhang, M. Imaizumi, B. Schölkopf, and K. Muandet, "Maximum Moment Restriction for Instrumental Variable Regression," *arXiv:2010.07684 [cs]*, Oct. 2020, arXiv: 2010.07684.

[55] A. Puli and R. Ranganath, "General control functions for causal effect estimation from instrumental variables," *Advances in Neural Information Processing Systems*, vol. 33, p. 8440, 2020.

[56] L. Xu, Y. Chen, S. Srinivasan, N. de Freitas, A. Doucet, and A. Gretton, "Learning Deep Features in Instrumental Variable Regression," *arXiv:2010.07154 [cs, stat]*, Oct. 2020, arXiv: 2010.07154.

[57] X. Chen and T. M. Christensen, "Optimal sup-norm rates and uniform inference on nonlinear functionals of nonparametric iv regression," *Quantitative Economics*, vol. 9, no. 1, pp. 39–84, 2018.

[58] B. T. Knapik, A. W. van der Vaart, and J. H. van Zanten, "Bayesian inverse problems with Gaussian priors," *The Annals of Statistics*, vol. 39, no. 5, pp. 2626–2657, Oct. 2011, arXiv: 1103.2692.

[59] A. Van der Vaart, H. Van Zanten *et al.*, "Bayesian inference with rescaled Gaussian process priors," *Electronic Journal of Statistics*, vol. 1, pp. 433–448, 2007.

[60] C. Ai and X. Chen, "Efficient estimation of models with conditional moment restrictions containing unknown functions," *Econometrica*, vol. 71, no. 6, pp. 1795–1843, 2003.

[61] A. W. van der Vaart and J. H. van Zanten, "Rates of contraction of posterior distributions based on Gaussian process priors," *The Annals of Statistics*, vol. 36, no. 3, Jun. 2008.

[62] S. Ghosal, J. K. Ghosh, and A. W. Van Der Vaart, "Convergence rates of posterior distributions," *Annals of Statistics*, pp. 500–531, 2000.

[63] A. Caponnetto and E. De Vito, "Optimal rates for the regularized least-squares algorithm," *Foundations of Computational Mathematics*, vol. 7, no. 3, pp. 331–368, 2007.

[64] K. Fukumizu, "Nonparametric Bayesian inference with kernel mean embedding," in *Modern Methodology and Applications in Spatial-Temporal Modeling*. Springer, 2015, pp. 1–24.

[65] E. De Vito, L. Rosasco, A. Caponnetto, U. De Giovannini, F. Odone, and P. Bartlett, "Learning from examples as an inverse problem." *Journal of Machine Learning Research*, vol. 6, no. 5, 2005.

[66] R. Vershynin, *High-dimensional probability: An introduction with applications in data science*. Cambridge University Press, 2018, vol. 47.