Accelerometry-Derived Digital Biomarkers for Cardiometabolic Risk: A Population-Representative Tabular Benchmark with Uncertainty Quantification

Federico Felizzi [1] [2]

Abstract

Structured tabular data is the dominant format in clinical medicine, yet existing tabular benchmarks fail to reflect key properties of real-world health data: complex survey sampling, demographic oversampling, clinically validated outcomes, and fairness requirements across population subgroups. We introduce the *NHANES Accelerometry Cardiometabolic Benchmark*, derived from the National Health and Nutrition Examination Survey (NHANES) 2003–2006, comprising 1,381 adults with hip-worn accelerometry, fasting laboratory biomarkers, dietary intake, and anthropometric measurements. We evaluate three methods spanning the specialised-to-general spectrum of tabular learning—ridge regression, XGBoost, and the tabular foundation model TabPFN v2—on prediction of glycated haemoglobin (HbA1c), fasting triglycerides, and C-reactive protein (CRP) from accelerometry-derived activity phenotypes and lifestyle covariates. TabPFN v2 achieves the best overall performance (HbA1c R^2=0.156, CRP R^2=0.383), while triglycerides remain largely unpredictable from lifestyle features alone ($R^2 < 0.05$ across all models), consistent with known genetic dominance. We apply *split conformal prediction* to all models, producing distribution-free 90% prediction intervals, and evaluate demographic coverage equity across sex and race/ethnicity subgroups. Marginal coverage is close to the 90% target for the more predictable outcomes (CRP and HbA1c) but falls below target for triglycerides, the least predictable outcome; at the subgroup level we observe localised undercoverage (e.g., HbA1c for Mexican American participants), illustrating the gap between the marginal guarantees of split conformal prediction and the conditional coverage that clinical fairness ultimately requires. All code and the derived benchmark dataset are publicly available at https://github.com/felizzi/nhanes-accel-cardiometabolic-benchmark.

1. Introduction

Structured tabular data underlies the majority of clinical decision-making, spanning electronic health records, laboratory results, and wearable sensor summaries (Jiang et al., 2025). Despite rapid progress in tabular representation learning—from gradient-boosted trees to transformer architectures and tabular foundation models—benchmark evaluations predominantly use generic datasets that omit properties critical to health applications: complex survey sampling designs, population representativeness, clinically validated outcome measures, and regulatory requirements for equitable performance across demographic subgroups (McElfresh et al., 2023).

Wearable accelerometers offer a compelling avenue for population-scale digital biomarker discovery. Physical activity measured continuously over seven days captures behavioural patterns meaningfully associated with cardiometabolic risk, yet translating raw activity counts into clinically actionable predictions requires models that handle heterogeneous mixed-type features, missing laboratory values, and—critically—must provide reliable uncertainty estimates to support clinical deployment.

We make four contributions. **First**, we introduce the *NHANES Accelerometry Cardiometabolic Benchmark*, a population-representative health tabular benchmark with properties absent from existing suites: survey design weights, demographic oversampling of minority groups, clinically validated fasting outcomes, and two-wave temporal structure. **Second**, we conduct a rigorous comparison of methods spanning the taxonomy of Jiang et al. (2025)—from specialised classical models to general tabular founda-

[1]SIIAM, Rome, Italy [2]enGene Statistics, Basel, Switzerland. Correspondence to: Federico Felizzi <federico.felizzi@gmail.com>.

Proceedings of the Workshop on Structured Data for Health at the 43rd International Conference on Machine Learning, Seoul, South Korea. Copyright 2026 by the author(s).

tion models—on three cardiometabolic outcomes. **Third**, we apply *split conformal prediction* (Angelopoulos & Bates, 2023) to all models, producing distribution-free prediction intervals with finite-sample coverage guarantees. **Fourth**, we evaluate whether conformal coverage holds equitably across the sex and race/ethnicity subgroups that NHANES was designed to represent, directly addressing the fairness gap identified in Jiang et al. (2025).

2. Data and Benchmark

2.1. NHANES 2003–2006

The National Health and Nutrition Examination Survey (NHANES) is a stratified, multistage probability sample of the non-institutionalised US civilian population, conducted by the Centers for Disease Control and Prevention (Centers for Disease Control and Prevention, 2006). We use examination cycles C (2003–2004) and D (2005–2006), which included hip-worn ActiGraph accelerometry for up to seven consecutive days.

2.2. Analytic Sample

Starting from participants with valid accelerometry data (≥ 4 wear days, ≥ 600 min/day wear time), we applied the following inclusion criteria: age 20–85 years; non-missing HbA1c, triglycerides, and CRP; and fasting duration ≥ 8 hours prior to blood draw (required for valid lipid measurements). The final analytic sample comprised **N=1,381** participants. Descriptive statistics are reported in Table 1.

Table 1. Analytic sample characteristics (N=1,381).

Variable	Mean	SD	Median
Age (years)	52.7	18.6	53.0
BMI (kg/m^2)	28.3	5.9	27.5
HbA1c (%)	5.6	0.9	5.4
Triglycerides (mg/dL)	153.3	146.7	123.0
CRP (mg/dL)	0.5	1.0	0.2
TAC ($\times 10^3$ counts/day)	251.3	139.9	226.8
MVPA (min/day)	20.6	22.6	13.1
Sedentary (min/day)	1086.7	115.6	1093.0
Energy (kcal/day)	2149.0	931.9	1990.5
Female (%)		50.6%	
NH White (%)		57.1%	
Mexican American (%)		20.8%	
NH Black (%)		15.9%	
Other (%)		6.2%	

2.3. Features

Activity features (6): total activity counts (TAC), log-TAC (TLAC), sedentary time (ST), moderate-to-vigorous PA (MVPA), light PA (LIPA), and wear time (WT), computed from minute-level accelerometry using standard cutpoints (Troiano et al., 2008).

Demographic and clinical covariates (12): age, sex, race/ethnicity, poverty-income ratio, BMI, systolic blood pressure, smoking status, and total energy, carbohydrate, fat, protein, and fibre intake from 24-hour dietary recall.

Outcomes: log-transformed HbA1c (%), fasting triglycerides (mg/dL), and CRP (mg/dL), each modelled separately.

3. Methods

3.1. Experimental Setup

We partitioned the analytic sample into train (60%), calibration (20%), and test (20%) sets using stratified random splitting (seed=42). Numerical features were standardised using training-set statistics. Categorical features (sex, race/ethnicity, smoking) were label-encoded. All outcomes were log-transformed prior to modelling to address right-skewed distributions; reported metrics are computed on the log scale.

3.2. Models

We evaluated three models spanning the specialised-to-general taxonomy of Jiang et al. (2025):

Ridge Regression (α=1.0): a linear specialised baseline providing an interpretable lower bound on predictive performance.

XGBoost (Chen & Guestrin, 2016): a gradient-boosted tree ensemble representing the current state of practice for clinical tabular prediction. We used 500 estimators with learning rate 0.05, max depth 6, and early stopping on the calibration set.

TabPFN v2 (Hollmann et al., 2025) is a tabular foundation model representing the *general* tier of the taxonomy of Jiang et al. (2025) — it requires no task-specific training, applying a learned prior directly to downstream tasks without fine-tuning. TabPFN v2 is pre-trained via *in-context learning* on millions of synthetically generated datasets constructed from structural causal models and Bayesian neural networks, enabling the model to perform approximate Bayesian inference over tabular inputs at test time. Concretely, given a training set $\mathcal{D}_{\text{train}}$ and a test instance x^*, TabPFN v2 computes:

$$\hat{y}^* = g_\Theta(x^* \mid \mathcal{D}_{\text{train}}) \tag{1}$$

where g_Θ is a transformer whose weights Θ are fixed after pre-training — the training set is passed directly as context, and predictions are produced in a single forward pass without gradient updates. This *in-context* mechanism allows TabPFN v2 to implicitly perform model selection and uncertainty estimation within its forward pass, which we

hypothesise accounts for its advantage over task-specifically trained baselines on our modest sample size (N_{train}=828). Following Hollmann et al. (2025), training instances are passed directly as in-context examples, with subsampling to 1,000 examples applied only when the training set exceeds this size; here N_{train}=828 < 1,000, so all training examples were used as context without subsampling. We used CPU inference throughout.

3.3. Conformal Prediction

We applied *split conformal prediction* (Angelopoulos & Bates, 2023) to each model using the calibration set. For each model and outcome, we computed the non-conformity score as the absolute residual $|y_i - \hat{y}_i|$ on the calibration set, and formed prediction intervals at miscoverage level α=0.10 (targeting 90% coverage):

$$\hat{C}(x) = [\hat{y} - \hat{q}, \ \hat{y} + \hat{q}] \tag{2}$$

where $\hat{q}$ is the $\lceil (1 - \alpha)(1 + 1/n_{cal}) \rceil$ quantile of calibration residuals. Split conformal prediction provides a finite-sample marginal coverage guarantee without distributional assumptions (Angelopoulos & Bates, 2023).

3.4. Subgroup Coverage Analysis

We evaluated whether conformal coverage held equitably across demographic subgroups defined by sex (male/female) and race/ethnicity (Mexican American, Other Hispanic, Non-Hispanic White, Non-Hispanic Black, Other), using the NHANES oversampling design that ensures adequate representation of minority groups.

4. Results

4.1. Predictive Performance

Table 2 reports R^2, mean absolute error (MAE, log scale), and empirical conformal coverage for all model–outcome combinations.

Table 2. Model comparison on test set (N=276). Coverage target = 0.90. MAE reported on log scale.

Outcome	Model	R^2	MAE	Cov.
	Ridge	0.133	0.073	0.910
HbA1c	XGBoost	0.139	0.073	0.881
	TabPFN v2	**0.156**	**0.070**	0.884
	Ridge	−0.002	0.474	0.863
Triglycerides	XGBoost	−0.007	0.470	0.823
	TabPFN v2	**0.048**	**0.453**	0.819
	Ridge	0.339	0.787	0.917
CRP	XGBoost	0.313	0.809	0.913
	TabPFN v2	**0.383**	**0.767**	**0.939**

TabPFN v2 achieves the best R^2 on all three outcomes, with

the largest margin on CRP (R^2=0.383 vs. 0.339 for ridge). The advantage of TabPFN v2 is most pronounced for CRP (ΔR^2=+0.044 over ridge, +0.070 over XGBoost), consistent with its in-context Bayesian prior capturing non-linear interactions between activity, BMI, and dietary features that are poorly approximated by linear models and insufficiently regularised in gradient-boosted trees at this sample size. CRP is the most predictable outcome across all models, consistent with the strong behavioural determinants of systemic inflammation. Triglycerides show near-zero or negative R^2 for all models, reflecting the dominant role of acute dietary intake and genetics neither captured in our feature set nor available in NHANES public-use files (Teslovich et al., 2010). HbA1c is moderately predictable (R^2=0.133–0.156), consistent with published estimates using lifestyle variables in general population samples (Selvin et al., 2004).

Conformal coverage is closest to the 90% target for ridge regression (HbA1c: 0.910, CRP: 0.917) and TabPFN v2 on CRP (0.939). XGBoost shows slight undercoverage on triglycerides (0.823), reflecting greater overconfidence in its residual distribution.

4.2. Subgroup Coverage Analysis

Figure 1 shows empirical conformal coverage by demographic subgroup for TabPFN v2. Coverage is broadly consistent across sex and race/ethnicity subgroups for CRP, with all groups meeting or exceeding the 90% target. However, notable undercoverage is observed for Mexican American participants on HbA1c (0.719), while other groups achieve ≥ 0.87 coverage on this outcome. Undercoverage is also observed for triglycerides across all subgroups, consistent with the poor overall predictability of this outcome. Notably, coverage for Non-Hispanic Black participants meets or exceeds the 90% target on both HbA1c and CRP, indicating that the marginal intervals remain valid for this group despite its higher prevalence of cardiometabolic risk.

5. Discussion

TabPFN v2 as a strong baseline for small health datasets. Our results show that the tabular foundation model TabPFN v2 achieves the best point performance among the evaluated baselines on all three outcomes without any task-specific training, although on this single split the margins over ridge and XGBoost are modest and should be confirmed with repeated-split evaluation. This is consistent with findings in Hollmann et al. (2025) showing that TabPFN v2 achieves state-of-the-art performance on small tabular datasets (N<10,000), and extends this finding to a clinically structured, population-representative health benchmark.

Figure 1. Empirical conformal coverage (90% target) by demographic subgroup for TabPFN v2. Colour indicates coverage level: green ≥0.90, red <0.90.

Why TabPFN v2 outperforms task-specific models. The superior performance of TabPFN v2 on a dataset of N=1,381 is consistent with its design objective: the model's pre-training on synthetic datasets spanning diverse causal structures effectively encodes a broad prior over tabular relationships, which acts as strong regularisation when task-specific training data is scarce. Classical models such as ridge regression and XGBoost must estimate all structure from the available 828 training examples, while TabPFN v2 leverages knowledge distilled from millions of synthetic datasets to inform its predictions. This finding extends results from Hollmann et al. (2025) — which demonstrated TabPFN v2's advantage on generic small tabular benchmarks — to a clinically structured, population-representative health dataset, suggesting that tabular foundation models may be particularly well-suited to health applications where data collection is expensive and sample sizes are inherently limited.

Outcome-specific predictability. The strong contrast between CRP ($R^2 \approx 0.38$) and triglycerides ($R^2 \approx 0.00$) highlights a key challenge for digital biomarker discovery: not all cardiometabolic outcomes are equally predictable from lifestyle and activity data. Triglycerides are dominated by genetic factors (Teslovich et al., 2010) and acute dietary intake, neither of which is captured by 7-day accelerometry or single 24-hour dietary recall. Future work incorporating genetic data or repeated dietary assessments may improve triglyceride prediction substantially.

Marginal versus conditional coverage. Split conformal prediction provides valid *marginal* coverage without distributional assumptions, making it an attractive starting point for the heterogeneous population structure of NHANES. Marginal validity, however, does not imply conditional validity: the guarantee holds on average over the test distribu-

tion and can be violated within individual subgroups. Our analysis bears this out — coverage is close to nominal for most groups on CRP and HbA1c, but drops to 0.719 for Mexican American participants on HbA1c, and triglyceride intervals undercover across the board. Marginal conformal methods alone are therefore insufficient to certify demographic fairness. Group-conditional (Mondrian) conformal calibration, which forms separate calibration quantiles per subgroup, is a natural remedy, although exact conditional coverage is generally unattainable in finite samples without further distributional assumptions (Angelopoulos & Bates, 2023); we leave a systematic comparison to future work.

Limitations. Our evaluation uses a single train/calibration/test split (seed=42). While sufficient to establish the benchmark and to report empirical coverage, the small test set (N=276) and the narrow R^2 gaps between models (e.g., 0.133 vs. 0.156 for HbA1c) mean that point-estimate differences should be read as indicative rather than as statistically established rankings; a fixed split on a small sample can also induce calibration–test distribution shift, which partly explains the residual undercoverage we observe. The benchmark is also limited in breadth: we compare three models and three cardiometabolic markers (HbA1c, triglycerides, CRP), leaving a fuller panel (e.g., fasting glucose, HDL/LDL cholesterol) and additional learners for future iterations. The sample is predominantly Non-Hispanic White (57.1%), reflecting the fasting requirement which differentially excluded participants from minority groups, and results may not generalise equally across all populations. The 2003–2006 protocol uses hip-worn, predominantly uniaxial ActiGraph devices, which differ substantially from the wrist-worn triaxial sensors in contemporary consumer and research wearables; the cutpoint-based activity features and the trained models may transfer poorly to modern data streams, and we make no claim of device-level generalisation. The single 24-hour dietary recall introduces measurement error in dietary covariates. NHANES does not include genetic data, precluding adjustment for heritable determinants of cardiometabolic risk. Sedentary time as computed includes non-wear periods and should be interpreted with caution.

Future work. Several directions follow directly from the limitations above. *Robustness*: repeating the full pipeline over multiple random splits and reporting mean±std with paired tests on per-seed differences would put the model rankings on firmer statistical footing. *Calibration*: replacing split conformal with cross-conformal or jackknife+ (Barber et al., 2021) calibration would use the full sample for both fitting and calibration, reducing the sensitivity of coverage to a single calibration draw, and group-conditional (Mondrian) calibration would target subgroup-level validity directly. *Ablations*: adding survey-weighted ridge/GBM baselines

(consistent with the population-representative framing) and a demographics- or BMI-only baseline for CRP would isolate the incremental signal contributed by accelerometry beyond established anthropometric predictors. *Scope*: incorporating NHANES 2011–2014 wrist accelerometry for larger samples and richer signal; broadening the outcome panel to fasting glucose and HDL/LDL cholesterol for a more comprehensive cardiometabolic profile; evaluating FT-Transformer (Gorishniy et al., 2021) at sufficient sample sizes (N>5,000) where transformer architectures are expected to realise their representational advantages (Jiang et al., 2025); and extending conformal prediction to survival outcomes using NHANES linked mortality data.

6. Conclusion

We introduced the NHANES Accelerometry Cardiometabolic Benchmark as a population-representative health tabular benchmark and showed that the tabular foundation model TabPFN v2 achieves the strongest point predictions among the evaluated baselines on all three cardiometabolic outcomes, with the caveat that these comparisons rest on a single data split and modest margins. Split conformal intervals attain close-to-nominal *marginal* coverage for the more predictable outcomes, while subgroup analysis exposes localised undercoverage that marginal guarantees do not preclude — motivating conditional and group-aware calibration as the natural next step. Our benchmark and code provide a reproducible foundation for future comparisons of tabular learning methods on clinically structured health data.

References

Angelopoulos, A. N. and Bates, S. A gentle introduction to conformal prediction and distribution-free uncertainty quantification. *arXiv preprint arXiv:2107.07511*, 2023.

Barber, R. F., Candès, E. J., Ramdas, A., and Tibshirani, R. J. Predictive inference with the jackknife+. *The Annals of Statistics*, 49(1):486–507, 2021. doi: 10.1214/20-AOS1965.

Centers for Disease Control and Prevention. NHANES 2003–2006: National health and nutrition examination survey. `https://www.cdc.gov/nchs/nhanes`, 2006.

Chen, T. and Guestrin, C. XGBoost: A scalable tree boosting system. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pp. 785–794, 2016.

Gorishniy, Y., Rubachev, I., Khrulkov, V., and Babenko, A. Revisiting deep learning models for tabular data. In *Advances in Neural Information Processing Systems*, volume 34, pp. 18932–18943, 2021.

Hollmann, N., Müller, S., Purucker, L., Krishnakumar, A., Körfer, M., Hoo, S. B., Schirrmeister, R. T., and Hutter, F. Accurate predictions on small data with a tabular foundation model. *Nature*, 637(8045):319–326, 2025.

Jiang, J.-P., Liu, S.-Y., Cai, H.-R., Zhou, Q., and Ye, H.-J. Representation learning for tabular data: A comprehensive survey. *arXiv preprint arXiv:2504.16109*, 2025.

McElfresh, D., Khandagale, S., Valverde, J., Ramakrishnan, G., Goldblum, M., and White, C. When do neural nets outperform boosted trees on tabular data? In *Advances in Neural Information Processing Systems*, volume 36, 2023.

Selvin, E., Coresh, J., Golden, S. H., Brancati, F. L., Folsom, A. R., and Steffes, M. W. Glycemic control and coronary heart disease risk in persons with and without diabetes. *Archives of Internal Medicine*, 164(19):2147–2155, 2004.

Teslovich, T. M., Musunuru, K., Smith, A. V., et al. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature*, 466(7307):707–713, 2010.

Troiano, R. P., Berrigan, D., Dodd, K. W., Mâsse, L. C., Tilert, T., and McDowell, M. Physical activity in the United States measured by accelerometer. *Medicine & Science in Sports & Exercise*, 40(1):181–188, 2008.