Riemannian Score-Based Generative Modelling

Abstract

Score-based generative models (SGMs) are a powerful class of generative models that exhibit remarkable empirical performance. Score-based generative modelling (SGM) consists of a "noising" stage, whereby a diffusion is used to gradually add Gaussian noise to data, and a generative model, which entails a "denoising" process defined by approximating the time-reversal of the diffusion. Existing SGMs assume that data is supported on a Euclidean space, i.e. a manifold with flat geometry. In many domains such as robotics, geoscience or protein modelling, data is often naturally described by distributions living on Riemannian manifolds and current SGM techniques are not appropriate. We introduce here *Riemannian Score-based Generative Models* (RSGMs), a class of generative models extending SGMs to Riemannian manifolds. We demonstrate our approach on a variety of compact manifolds, and in particular with earth and climate science spherical data.

1 Introduction

Score-based Generative Models (SGMs) (Song and Ermon, 2019; Song et al., 2021; Ho et al., 2020; Dhariwal and Nichol, 2021) formulate generative modelling as a denoising process. Noise is incrementally added to data using a diffusion process until it gets approximately Gaussian. The generative model is then obtained by simulating an approximation of the corresponding time-reversal process, which progressively denoises a Gaussian sample to obtain a data sample. This process is also a diffusion whose drift depends on the logarithmic gradients of the noised data densities, i.e. the Stein scores, estimated using a neural network via score matching (Hyvärinen, 2005; Vincent, 2011).

SGMs have been primarily applied to data living on Euclidean spaces, i.e. manifolds with flat geometry. However, in a large number of scientific domains the distributions of interest are supported on Riemannian manifolds. These include, to name a few, protein modelling (Shapovalov and Dunbrack Jr, 2011), cell development (Klimovskaia et al., 2020), image recognition (Lui, 2012), geological sciences (Karpatne et al., 2018; Peel et al., 2001), graph-structured and hierarchical data (Roy et al., 2007; Steyvers and Tenenbaum, 2005), robotics (Feiten et al., 2013; Senanayake and Ramos, 2018) and high-energy physics (Brehmer and Cranmer, 2020).

We introduce in this work *Riemannian Score-based Generative Models* (RSGMs), an extension of SGMs to Riemannian manifolds which incorporate the geometry of the data by defining the forward diffusion process directly on the Riemannian manifold, inducing a manifold-valued reverse process. This requires constructing a noising process on the manifold that converges to an easy-to-sample reference distribution. We focus on the Brownian motion which converges with geometric rate to the uniform distribution on compact Riemannian manifolds (Saloff-Coste, 1994). We establish that, as in the Euclidean case, the corresponding time-reversal process is also a diffusion whose drift includes the Stein score which is intractable but can similarly be estimated via score matching. Methodological extensions are required as the transition kernel of the Brownian motion cannot be sampled exactly and is typically only available as an infinite sum through the Sturm–Liouville decomposition (Chavel, 1984) on compact Riemmanian manifolds. To this end, we develop non-standard techniques for score estimation and rely on the use of Geodesic Random Walks for sampling (Jørgensen, 1975).

We provide theoretical convergence bounds for RSGMs on compact manifolds and demonstrate our approach on a range of manifolds and tasks, including modelling a number of natural disaster occurrence datasets collected by Mathieu and Nickel (2020). We show that RGSMs achieve better performance than recent baselines (Mathieu and Nickel, 2020; Rozen et al., 2021) and scale better to high-dimensional manifolds.

2 Euclidean Score-based Generative Modelling

We recall here briefly the key concepts behind SGMs on the Euclidean space $\mathbb{R}^d$ and refer the readers to Song et al. (2021) for a more detailed introduction. We consider a forward *noising* process $(\mathbf{X}_t)_{t \geq 0}$ defined by the following Stochastic Differential Equation (SDE)

$$d\mathbf{X}_t = -\mathbf{X}_t dt + \sqrt{2}d\mathbf{B}_t, \quad \mathbf{X}_0 \sim p_0, \tag{1}$$

where $(\mathbf{B}_t)_{t \geq 0}$ is a d-dimensional Brownian motion and p_0 is the data distribution. The available data gives us an empirical approximation of p_0. The process $(\mathbf{X}_t)_{t \geq 0}$ is simply an Ornstein–Ulhenbeck (OU) process which converges with geometric rate to $\mathrm{N}(0, \mathrm{Id})$. Under mild conditions on p_0, the time-reversed process $(\mathbf{Y}_t)_{t \geq 0} = (\mathbf{X}_{T-t})_{t \in [0,T]}$ also satisfies an SDE (Cattiaux et al., 2021; Haussmann and Pardoux, 1986) given by

$$d\mathbf{Y}_t = \{\mathbf{Y}_t + 2\nabla \log p_{T-t}(\mathbf{Y}_t)\}dt + \sqrt{2}d\mathbf{B}_t, \quad \mathbf{Y}_0 \sim p_T, \tag{2}$$

where p_t denotes the density of $\mathbf{X}_t$. By construction, the law of $\mathbf{Y}_{T-t}$ is equal to the law of $\mathbf{X}_t$ for $t \in [0, T]$ and in particular $\mathbf{Y}_T \sim p_0$. Hence, if one could sample from $(\mathbf{Y}_t)_{t \in [0,T]}$ then its final distribution would be the data distribution p_0. Unfortunately we cannot sample exactly from (2) as p_T and the scores $(\nabla \log p_t(x))_{t \in [0,T]}$ are intractable. Hence SGMs rely on a few approximations. First, p_T is replaced by the reference distribution $\mathrm{N}(0, \mathrm{Id})$ as we know that p_T converges geometrically towards it. Second, the following denoising score matching identity is exploited to estimate the scores

$$\nabla_{x_t} \log p_t(x_t) = \int_{\mathbb{R}^d} \nabla_{x_t} \log p_{t|0}(x_t|x_0) \, p_{0|t}(x_0|x_t) dx_0,$$

where $p_{t|0}(x_t|x_0)$ is the transition density of the OU process (1) which is available in closed-form. It follows directly that $\nabla \log p_t$ is the minimizer of $\ell_t(\mathbf{s}) = \mathbb{E}[\|\mathbf{s}(\mathbf{X}_t) - \nabla_{x_t} \log p_{t|0}(\mathbf{X}_t|\mathbf{X}_0)\|^2]$ over functions $\mathbf{s}$ where the expectation is over the joint distribution of $\mathbf{X}_0, \mathbf{X}_t$. This result can be leveraged by considering a neural network $\mathbf{s}_\theta : [0,T] \times \mathbb{R}^d \to \mathbb{R}^d$ trained by minimizing the loss function $\ell(\theta) = \int_0^T \lambda_t \ell_t(\mathbf{s}_\theta(t, \cdot)) dt$ for some weighting function $\lambda_t > 0$. Finally, an Euler–Maruyama discretization of (2) is performed using a discretization step γ such that $T = \gamma N$ for $N \in \mathbb{N}$

$$Y_{n+1} = Y_n + \gamma\{Y_n + 2\mathbf{s}_\theta(T - n\gamma, Y_n)\} + \sqrt{2\gamma}Z_{n+1}, \quad Y_0 \sim \mathrm{N}(0, \mathrm{Id}), \quad Z_n \overset{\text{i.i.d.}}{\sim} \mathrm{N}(0, \mathrm{Id}).$$

The above showcases the basics of SGMs but we highlight that many improvements have been proposed; see (e.g. Song and Ermon, 2020; Jolicoeur-Martineau et al., 2021; Dhariwal and Nichol, 2021). In particular, selecting an adaptive stepsize $(\gamma_n)_{n \in \mathbb{N}}$ (Bao et al., 2022; Watson et al., 2021) and using a predictor-corrector scheme (Song et al., 2021) instead of a simple Euler–Maruyama discretization drastically improves performance.

3 Riemannian Score-based Generative Modelling

We now move to the manifold setting, and more specially assume that $\mathcal{M}$ is a complete and connected Riemannian manifold. As discussed in Section 1, three components are required to extend SGM: i) a time-reversal formula on $\mathcal{M}$ which defines a backward generative process, ii) a forward *noising* process on $\mathcal{M}$ which converges to an easy-to-sample reference distribution, iii) a method to efficiently approximate the drift of the time-reversal process.

3.1 Time-reversal on Riemannian manifolds

First we prove the time-reversal formula for manifolds, a generalisation of the results in the Euclidean case, e.g. see Cattiaux et al. (2021, Theorem 4.9). Consider an SDE of the form $d\mathbf{X}_t = b(\mathbf{X}_t)dt + d\mathbf{B}_t^{\mathcal{M}}$ where $\mathbf{B}_t^{\mathcal{M}}$ is a Brownian motion on $\mathcal{M}$. We refer to Appendix B.3 for an introduction to Brownian motions on manifolds. This result shows that if $(\mathbf{X}_t)_{t \in [0,T]}$ is a diffusion process then

---

Algorithm 1 GRW (Geodesic Random Walk)

Require: $T, N, X_0^\gamma, b, \sigma, \mathrm{P}$

1: $\gamma = T/N$ ▷ Step-size

2: **for** $k \in \{0, \ldots, N-1\}$ **do**

3:      $\bar{Z}_{k+1} \sim \mathrm{N}(0, \mathrm{Id})$ ▷ Standard Gaussian in ambient space $\mathbb{R}^p$

4:      $Z_{k+1} = \mathrm{P}(X_k^\gamma)\bar{Z}_{k+1}$ ▷ Projection in the tangent space $\mathrm{T}_x\mathcal{M}$

5:      $W_{k+1} = \gamma b(k\gamma, X_k^\gamma) + \sqrt{\gamma}\sigma(k\gamma, X_k^\gamma)Z_{k+1}$ ▷ Euler–Maruyama step on tangent space

6:      $X_{k+1}^\gamma = \exp_{X_k^\gamma}[W_{k+1}]$ ▷ Geodesic projection onto $\mathcal{M}$

7: **end for**

8: **return** $\{X_k^\gamma\}_{k=0}^N$

$(\mathbf{X}_{T-t})_{t\in[0,T]}$ is also a diffusion process w.r.t. the backward filtration whose coefficients can be computed. The proof relies on an extension of Cattiaux et al. (2021, Theorem 4.9) to the Riemannian manifold case and is postponed to Appendix G.

Theorem 1 (Time-reversed diffusion). *Let $T \geq 0$ and $(\mathbf{B}_t^{\mathcal{M}})_{t\geq 0}$ be a Brownian motion on $\mathcal{M}$ such that $\mathbf{B}_0^{\mathcal{M}}$ has distribution the volume form p_{ref}[1]. Let $(\mathbf{X}_t)_{t\in[0,T]}$ associated with the SDE $\mathrm{d}\mathbf{X}_t = b(\mathbf{X}_t)\mathrm{d}t + \mathrm{d}\mathbf{B}_t^{\mathcal{M}}$. Let $(\mathbf{Y}_t)_{t\in[0,T]} = (\mathbf{X}_{T-t})_{t\in[0,T]}$ and assume that $\mathrm{KL}(\mathbb{P}|\mathbb{Q}) < +\infty$, where $\mathbb{Q}$ is the distribution of $(\mathbf{B}_t^{\mathcal{M}})_{t\in[0,T]}$ and $\mathbb{P}$ the distribution of $(\mathbf{X}_t)_{t\in[0,T]}$. In addition, assume that $\mathbb{P}_t = \mathcal{L}(\mathbf{X}_t)$, the distribution of $\mathbf{X}_t$, admits a smooth positive density p_t w.r.t. p_{ref} for any $t \in [0,T]$. Then, $(\mathbf{Y}_t)_{t\in[0,T]}$ is associated with the SDE*

$$\mathrm{d}\mathbf{Y}_t = \{-b(\mathbf{Y}_t) + \nabla \log p_{T-t}(\mathbf{Y}_t)\}\mathrm{d}t + \mathrm{d}\mathbf{B}_t^{\mathcal{M}}. \tag{3}$$

3.2 Approximate sampling of diffusions

Obtaining samples from a Brownian motion on a manifold is non-trivial in general. If $\mathcal{M}$ is isometrically embedded into $\mathbb{R}^p$ (with $p \geq d$) one can define $(\mathbf{B}_t^{\mathcal{M}})_{t\geq 0}$ as a $\mathbb{R}^p$-valued process, see Appendix B.3. However, this approach is *extrinsic* and requires the knowledge of the projection operator. Here we consider an *intrisic* approach based on Geodesic Random Walks (GRWs), see Jørgensen (1975) for a review of their properties. GRWs are not restricted to approximating the Brownian motion on $\mathcal{M}$ but in fact can approximate *any* well-behaved diffusion on $\mathcal{M}$. Hence, we introduce GRWs in a general framework and consider a discrete-time process $(X_n^\gamma)_{n\in\mathbb{N}}$ which approximates the diffusion $(\mathbf{X}_t)_{t\geq 0}$ defined by

$$\mathrm{d}\mathbf{X}_t = b(t, \mathbf{X}_t)\mathrm{d}t + \sigma(t, \mathbf{X}_t)\mathrm{d}\mathbf{B}_t^{\mathcal{M}}. \tag{4}$$

This generalization will be key when sampling the backward diffusion process defined in Theorem 1.

Definition 2 (Geodesic Random Walk). *Let X_0^γ be a $\mathcal{M}$-valued random variable. For any $\gamma > 0$, we define $(X_n^\gamma)_{n\in\mathbb{N}}$ such that for any $n \in \mathbb{N}$, $X_{n+1}^\gamma = \exp_{X_n^\gamma}[\gamma\{b(X_n^\gamma) + (1/\sqrt{\gamma})(V_{n+1} - b(X_n^\gamma))\}]$[2], where $(V_n)_{n\in\mathbb{N}}$ is a sequence of $\mathrm{T}\mathcal{M}$-valued random variables such that for any $n \in \mathbb{N}$, $\mathbb{E}[V_{n+1}|\mathcal{F}_n] = b(X_n^\gamma)$ and $\mathbb{E}[V_{n+1}V_{n+1}^\top|\mathcal{F}_n] = \sigma\sigma^\top(X_n^\gamma)$, where $\mathcal{F}_n$ is the filtration generated by $\{X_k^\gamma\}_{k=0}^n$. We say that the $\mathcal{M}$-valued process $(X_n^\gamma)_{n\in\mathbb{N}}$ is a Geodesic Random Walk.*

Algorithm 1 approximately simulates the diffusion $(\mathbf{X}_t)_{t\in[0,T]}$ defined in Equation (4) using GRWs; see Kuwada (2012); Cheng et al. (2022) for quantitative error bounds in the time-homogeneous case and Appendix H.2 for the time-inhomogeneous case.

3.3 Score approximation on Riemannian manifolds

Score-matching and loss functions. The reverse process from Equation (3) involves the Stein score $\nabla \log p_t$ which is unfortunately intractable. To derive an approximation, we first remark that for any $s, t \in (0, T]$ with $t > s$ and $x_t \in \mathcal{M}$, $p_t(x_t) = \int_{\mathcal{M}} p_{t|s}(x_t|x_s)\mathrm{d}\mathbb{P}_s(x_s)$, where $\mathbb{P}_s = \mathcal{L}(\mathbf{X}_s)$. Thus, we have that for any $s, t \in [0, T]$ with $t > s$ and $x_t \in \mathcal{M}$

$$\nabla_{x_t} \log p_t(x_t) = \int_{\mathcal{M}} \nabla_{x_t} \log p_{t|s}(x_t|x_s)\mathbb{P}_{s|t}(x_t, \mathrm{d}x_s).$$

Hence, for any $s, t \in [0, T]$ with $t > s$ we have that $\nabla \log p_t = \arg\min\{\ell_{t|s}(\mathbf{s}_t) : \mathbf{s}_t \in \mathrm{L}^2(\mathbb{P}_t)\}$,

---

[1]Note that in the case of a non-compact manifold p_{ref} is only a measure and not a probability measure.

[2]$\exp_x : \mathrm{T}_x\mathcal{M} \to \mathcal{M}$ denotes the exponential mapping on the manifold, see e.g. Lee (2013, Chapter 20).

Algorithm 2 RSGM (Riemannian Score-Based Generative Model)

Require: $\varepsilon, T, N, \{X_0^m\}_{m=1}^M, \text{loss}, \mathbf{s}, \theta_0, N_{\text{iter}}, p_{\text{ref}}, \mathrm{P}$
1: /// TRAINING ///
2: **for** $n \in \{0, \dots, N_{\text{iter}} - 1\}$ **do**
3:     $X_0 \sim (1/M) \sum_{m=1}^M \delta_{X_0^m}$ ▷ Random mini-batch from dataset
4:     $t \sim U([\varepsilon, T])$ ▷ Uniform sampling between ε and T
5:     $\mathbf{X}_t = \text{GRW}(t, N, X_0, b, \text{Id}, \mathrm{P})$ ▷ Approximate forward diffusion with Algorithm 1
6:     $\ell(\theta_n) = \ell_t(T, N, X_0, \mathbf{X}_t, \text{loss}, \mathbf{s}_{\theta_n})$ ▷ Compute score matching loss from Table 2
7:     $\theta_{n+1} = \texttt{optimizer\_update}(\theta_n, \ell(\theta_n))$ ▷ ADAM optimizer step
8: **end for**
9: $\theta^\star = \theta_{N_{\text{epoch}}}$
10: /// SAMPLING ///
11: $Y_0 \sim p_{\text{ref}}$ ▷ Sample from uniform distribution
12: $b_\theta^\star(t, x) = \mathbf{s}_{\theta^\star}(T - t, x)$ for any $t \in [0, T], x \in \mathcal{M}$ ▷ Reverse process drift
13: $\{Y_k\}_{k=0}^N = \text{GRW}(T, N, Y_0, b_{\theta^\star}, \text{Id}, \mathrm{P})$ ▷ Approximate reverse diffusion with Algorithm 1
14: **return** $\theta^\star, \{Y_k\}_{k=0}^N$

---

where $\ell_{t|s}(\mathbf{s}_t) = \int_{\mathcal{M}^2} \|\nabla_x \log p_{t|s}(x_t|x_s) - \mathbf{s}_t(x_t)\|^2 \mathrm{d}\mathbb{P}_{s,t}(x_s, x_t)$, which is referred as the Denoising Score Matching (DSM) loss. It can also be written in an *implicit* fashion.

Proposition 3. *Let $t, s \in (0, T)$ with $t > s$. Then, for any $\mathbf{s}_t \in C^\infty(\mathcal{M})$, $\ell_{t|s}(\mathbf{s}_t) = 2\ell_t^{\text{im}}(\mathbf{s}_t) + \int_{\mathcal{M}^2} \|\nabla_{x_t} \log p_{t|s}(x_t|x_s)\|^2 \mathrm{d}\mathbb{P}_{s,t}(x_s, x_t)$, where $\ell_t^{\text{im}}(\mathbf{s}_t) = \int_{\mathcal{M}} \{\frac{1}{2}\|\mathbf{s}_t(x_t)\|^2 + \text{div}(\mathbf{s}_t)(x_t)\} \mathrm{d}\mathbb{P}_t(x_t)$.*

The proof is postponed to Appendix I. For any $t \in (0, T)$ the minimizers of the loss ℓ_t^{im} on $\mathcal{X}(\mathcal{M})$ (where $\mathcal{X}(\mathcal{M})$ is the set of vector fields on $\mathcal{M}$) are the same as the ones for $\ell_{t|s}$. The loss ℓ_t^{im} is referred to as the *implicit* score matching (ISM) loss (Hyvärinen, 2005).

In the case that we have access to $\{\nabla \log p_{t|s} : T \leq t > s \geq 0\}$ or an approximation of this family, then we can use the DSM loss to learn $\{\mathbf{s}_t \in \mathcal{X}(\mathcal{M}) : t \in [0, t]\}$. If this is not the case then we turn to ℓ_t^{im}. Note that ℓ_t^{im} requires the computation of a divergence term which requires d Jacobian-vector calls. In high dimension, a stochastic estimator is necessary (Hutchinson, 1989). Following Song and Ermon (2020); Nichol and Dhariwal (2021) the loss can be weighted with a term $\lambda_t > 0$.

Parametric family of vector fields. We approximate $(\nabla \log p_t)_{t \in [0,T]}$ by a family of functions $\{\mathbf{s}_\theta\}_{\theta \in \Theta}$ where Θ is a set of parameters and $\mathbf{s}_\theta : [0, T] \to \mathcal{X}(\mathcal{M})$. In a Euclidean space, vector fields are simply functions $\mathbf{s}_\theta : \mathbb{R}^d \to \mathbb{R}^d$. In manifolds, although for any $x \in \mathcal{M}$, $T_x\mathcal{M} \cong \mathbb{R}^d$, there does not necessarily exist a set of d smooth vector fields $\{E_i\}_{i=1}^d$ such that $\text{span}\left(\{E_i(x)\}_{i=1}^d\right) = T_x\mathcal{M}$ (Chapter 8, page 179, Lee, 2006) [3]. Fortunately, one can rely on a larger set of smooth vector fields $\{E_i(x)\}_{i=1}^n$ with $n > d$ that *does* span the tangent bundle. Then it suffices to construct a neural network $\mathbf{s}_\theta : [0, T] \times \mathcal{M} \to \mathbb{R}^n$ to parametrize the score network as $\mathbf{s}_\theta(t, x) = \sum_{i=1}^n \mathbf{s}_\theta^i(t, x) E_i(x)$. See Appendix D for a discussion on the different choices of generating sets $\{E_i(x)\}_{i=1}^n$.

Combining this parameterization with the score matching losses, the time-reversal formula of Theorem 1 and the sampling of forward and backward processes described in Section 3.2, we define our RGSM algorithm in Algorithm 2. This algorithm can also benefit from a predictor-corrector scheme as in (Song et al., 2021), see Appendix F.

To implement RSGM on a specific manifold, one finally needs choosing a forward noising process. There is no unique choice and in Appendix M we discuss different choices of forward drift for potentially non-compact manifolds. In general, transition kernels may not be available in exact or approximate form, so one may need to rely on the ISM loss.

3.4 The compact manifold framework

Assuming compactness of the manifold $\mathcal{M}$, we can leverage a number of special properties to implement our algorithm. In particular we can choose a generic sensible noising process, and have access to a variety of approximations of the heat kernel.

---

[3]Manifolds for which there exists such a *global frame* $\{E_i(x)\}_{i=1}^d$ are referred as *parallelizable*. $\mathbb{S}^2$ is a well-known example of *non-parallelizable* manifold as per the *Hairy ball theorem*.

Table 1: Differences between SGM on Euclidean spaces and RSGM on compact Riemannian manifolds.

Ingredient \ Space	Euclidean	Compact manifold
Forward process	Ornstein–Ulhenbeck	Brownian motion
Easy-to-sample distribution	Gaussian	Uniform
Time reversal	Cattiaux et al. (2021, Theorem 4.9)	Theorem 1
Sampling of the forward process	Direct	Geodesic Random Walk
Sampling of the backward process	Euler–Maruyama	Geodesic Random Walk

We choose the noising process to be the Brownian motion $(\mathbf{B}_t^{\mathcal{M}})_{t\geq 0}$ since it converges geometrically to the uniform distribution p_{ref} under compactness assumptions on $\mathcal{M}$, see Urakawa (2006, Proposition 2.6). For most manifolds of interest, one can either sample exactly from p_{ref} or approximately with high accuracy. With this choice in place, Table 1 highlights the main differences between RSGMs on compact manifolds and Euclidean score-based models.

Heat kernel on compact Riemannian manifolds. For any $x_0 \in \mathcal{M}$ and $t \geq s \geq 0$, the heat kernel $p_{t|s}(\cdot|x_s)$ is defined as the density of $\mathbf{B}_t^{\mathcal{M}}$ w.r.t.

Contrary to the Gaussian transition density of the OU process (or the Brownian motion) in the Euclidean setting, it is typically only available as an infinite series. In order to circumvent this issue we consider two techniques: i) a truncation approach, ii) a Taylor expansion around $t = 0$ called a Varadhan asymptotics. First, we recall that in the case of compact manifolds the heat kernel is given by the Sturm–Liouville decomposition (Chavel, 1984) given for any $t > 0$ and $x_0, x_t \in \mathcal{M}$ by

$$p_{t|0}(x_t|x_0) = \sum_{j\in\mathbb{N}} \mathrm{e}^{-\lambda_j t}\phi_j(x_0)\phi_j(x_t), \tag{5}$$

where the convergence occurs in $\mathrm{L}^2(p_{\mathrm{ref}} \otimes p_{\mathrm{ref}})$, $(\lambda_j)_{j\in\mathbb{N}}$ and $(\phi_j)_{j\in\mathbb{N}}$ are the eigenvalues, respectively the eigenvectors, of $-\Delta_{\mathcal{M}}$ in $\mathrm{L}^2(p_{\mathrm{ref}})$ (see Saloff-Coste, 1994, Section 2). When the eigenvalues and eigenvectors are known, we rely on an approximation of the logarithmic gradient of $p_{t|0}$ by truncating the sum in Equation (S8) with $J \in \mathbb{N}$ terms to obtain for any $t > 0$ and $x_0, x_t \in \mathcal{M}$

$$\nabla_{x_t} \log p_{t|0}(x_t|x_0) \approx S_{J,t}(x_0,x_t) \triangleq \sum_{j=0}^{J} \mathrm{e}^{-\lambda_j t}\phi_j(x_0)\nabla_{x_t}\phi_j(x_t) / \sum_{j=0}^{J} \mathrm{e}^{-\lambda_j t}\phi_j(x_0)\phi_j(x_t). \tag{6}$$

Under regularity conditions on $\mathcal{M}$ it can be shown that for any $x, y \in \mathcal{M}$ and $t \geq 0$, $\lim_{J\to+\infty} S_{J,t}(x_0,x_t) = \nabla_{x_t} \log p_{t|0}(x_t|x_0)$ (see Jones et al., 2008, Lemma 1). In the case of the d-dimensional torus or sphere the eigenvalues and eigenvectors are computable (see Saloff-Coste, 1994, Section 2) and we can apply this method to approximate $p_{t|0}$ for any $t > 0$, see Appendix E

When the eigenvalues and eigenvectors are unknown or not tractable, we can still derive an approximation of the heat kernel for small times t. Using Varadhan's asymptotics—see Bismut (1984, Theorem 3.8) or Chen et al. (2021, Theorem 2.1)—for any $x, y \in \mathcal{M}$ with $y \notin \mathrm{Cut}(x)$ (where $\mathrm{Cut}(x)$ is the cut-locus of x in $\mathcal{M}$ (see Lee, 2018, Chapter 10)) we have that

$$\lim_{t\to 0} t\nabla_{x_t} \log p_{t|0}(x_t|x_0) = \exp_{x_t}^{-1}(x_0). \tag{7}$$

Using the previously defined score-matching losses and the approximations to the heat kernel above, we highlight three methods to compute $\nabla \log p_t$ in Table 2.

3.5 Convergence results in the compact setting

We now provide a theoretical analysis of RSGM under the assumption that $\mathcal{M}$ is compact. The following result ensures that RSGM generates samples whose distribution is close to the data distribution p_0. Let us denote $\{Y_k\}_{n\in\{0,\dots,N\}}$ the sequence generated by Algorithm 2. This result relies on the following assumption, which is satisfied for a large class of manifolds $\mathcal{M}$ such as the d-dimensional sphere and torus, compact matrix groups and products of these manifolds.

A1. *There exist $C, \alpha > 0$ such that for any $t \in (0, 1]$ and $x \in \mathcal{M}$, $p_{t|0}(x|x) \leq Ct^{-\alpha/2}$, where $p_{t|0}(\cdot|x_0)$ is the density of the heat kernel, i.e. the density of $\mathbf{B}_t^{\mathcal{M}}$ with initial condition x_0 [4].*

---

[4] The diagonal upper-bound is implied by Sobolev inequalities which control of the growth of some functions by the growth of their gradient. **A1** is satisfied in our experiments, see Saloff-Coste (1994); Gross (1992).

Table 2: Computational complexity of score matching losses w.r.t. score network forward and backward passes. ε is a random variable on $\mathrm{T}_{\mathbf{X}_t}\mathcal{M}$ such that $\mathbb{E}[\varepsilon] = 0$ and $\mathbb{E}[\varepsilon\varepsilon^\top] = \mathrm{Id}$.

Loss	Approximation	Loss function	Requirements $p_{t\mid 0}$	Requirements $\log_{\mathbf{X}_t}$	Complexity
$\ell_{t\mid 0}$ (DSM)	Truncation (6)	$\frac{1}{2}\mathbb{E}\left[\|\mathbf{s}(\mathbf{X}_t) - S_{J,t}(\mathbf{X}_0, \mathbf{X}_t)\|^2\right]$	✓	✗	$\mathcal{O}(1)$
	Varhadan (7)	$\frac{1}{2}\mathbb{E}\left[\|\mathbf{s}(\mathbf{X}_t) - \exp_{\mathbf{X}_t}^{-1}(\mathbf{X}_0)/t\|^2\right]$	✗	✓	$\mathcal{O}(1)$
$\ell_{t\mid s}$ (DSM)	Varhadan (7)	$\frac{1}{2}\mathbb{E}\left[\|\mathbf{s}(\mathbf{X}_t) - \exp_{\mathbf{X}_t}^{-1}(\mathbf{X}_s)/(t-s)\|^2\right]$	✗	✓	$\mathcal{O}(1)$
ℓ_t^{im} (ISM)	Deterministic	$\mathbb{E}\left[\frac{1}{2}\|\mathbf{s}(\mathbf{X}_t)\|^2 + \mathrm{div}(\mathbf{s})(\mathbf{X}_t)\right]$	✗	✗	$\mathcal{O}(d)$
	Stochastic	$\mathbb{E}\left[\frac{1}{2}\|\mathbf{s}(\mathbf{X}_t)\|^2 + \varepsilon^\top \partial\mathbf{s}(\mathbf{X}_t)\varepsilon\right]$	✗	✗	$\mathcal{O}(1)$

Theorem 4. *Assume* **A**1, *that* p_0 *is smooth and positive and that there exists* $\mathtt{M} \geq 0$ *such that for any* $t \in [0, T]$ *and* $x \in \mathcal{M}$, $\|\mathbf{s}_{\theta^\star}(t, x) - \nabla\log p_t(x)\| \leq \mathtt{M}$, *with* $\mathbf{s}_{\theta^\star} \in \mathrm{C}([0, T], \mathcal{X}(\mathcal{M}))$. *Then if* $T > 1/2$, *there exists* $C \geq 0$ *independent on* T *such that*

$$\mathbf{W}_1(\mathcal{L}(Y_N), p_0) = C(\mathrm{e}^{-\lambda_1 T} + \sqrt{T/2}\mathtt{M} + \mathrm{e}^T\gamma^{1/2}),$$

where $\mathbf{W}_1$ *is the Wasserstein distance of order one on the probability measures on* $\mathcal{M}$.

The proof is postponed to Appendix H. In particular, for any $\varepsilon > 0$, choosing $T > 0$ large enough, $\mathtt{M}$ small enough (which can be achieved using the universal property of neural networks) and γ small enough, we get that $\mathbf{W}_1(\mathcal{L}(Y_N), p_0) \leq \varepsilon$. This result might seem weaker than the result obtained for Moser flows in (Rozen et al., 2021, Theorem 3), but we emphasize that our bound takes into account the time-discretization contrary to Rozen et al. (2021) which considers the continuous-time flow. If we consider the time-reversed continuous-time SDE then we recover a bound in total variation distance, see Appendix H. Note that the upper bound $\mathtt{M}$ encompasses both the bias introduced by the use of a neural network and the bias introduced by the use of an approximation of the score.

4   Related work

In this section we discuss previous work on parametrizing family of distributions for manifold-valued data. Here, the manifold structure is considered to be prescribed, in contrast with methods that jointly learn the manifold structure and density (e.g. Brehmer and Cranmer, 2020; Caterini et al., 2021).

Parametric family of distributions. The various parametric families of manifold-valued distributions that have been proposed can be categorized into three main approaches (Navarro et al., 2017): wrapping, projecting and conditioning. Wrapped distributions consider a parametric distribution on $\mathbb{R}^n$ that is pushed-forward along a surjective map $\psi : \mathbb{R}^n \to \mathcal{M}$. Projected distributions are defined by marginalizing out some distribution along the normal bundle of $\mathcal{M}$. Conditioning distributions encompass the von Mises-Fisher and Kent distributions (Fisher, 1953; Kent, 1982). Considering mixtures of these distributions is key to increase flexibility (Peel et al., 2001; Mardia et al., 2008).

Push-forward of Euclidean normalizing flows. More recently, approaches leveraging the flexibility of normalizing flows (Papamakarios et al., 2019) have been proposed. Following the wrapping method described above, these methods parametrize a normalizing flow in $\mathbb{R}^n$ before being pushed along an invertible map $\psi : \mathbb{R}^n \to \mathcal{M}$. However, to globally represent the manifold, the map ψ needs to be a homeomorphism, which can only happen if $\mathcal{M}$ is topologically equivalent to $\mathbb{R}^n$, hence limiting the scope of that approach. One natural choice for this map is the exponential map $\exp_x : \mathrm{T}_x\mathcal{M} \cong \mathbb{R}^d$. This approach has been taken, for instance, by Falorsi et al. (2019) and Bose et al. (2020), respectively parametrizing distributions on Lie groups and hyperbolic space.

Neural ODE on manifolds. To avoid artifacts or numerical instabilities due to the manifold embedding, another line of work uses tools from Riemannian geometry to define flows directly on the manifold of interest (Falorsi and Forré, 2020; Mathieu and Nickel, 2020; Falorsi, 2021). Since these methods do not require a specific embedding mapping, they are referred as *Riemannian*. They extend continuous normalizing flows (CNFs) (Grathwohl et al., 2019) to the manifold setting, by implicity parametrizing flows as solutions of Ordinary Differential Equations (ODEs). As such, the parametric flow is a *continuous* function of time. This approach has recently been extended by Rozen et al.

Table 3: Summary of computational complexity (w.r.t. neural network forward and backward passes) for different methods. d is the manifold dimension, k the number of Monte Carlo batches in Moser flow's regularizer, N is the number of steps in the (adaptive) ODE solver, whereas N^* is the number of steps in the SDE Euler-Maruyama solver–which can usually be lower than N. Moser flow and RSGM training complexity varies if the Hutchinson stochastic estimator is used. See Table 2 for score matching losses complexity.

Method	Training	Likelihood evaluation	Sampling
RCNF	Solving ODE $\mathcal{O}(dN)$	Solving augmented ODE $\mathcal{O}(dN)$	Solving ODE $\mathcal{O}(N)$
Moser flow	Computing div $\mathcal{O}(dk)$ or $\mathcal{O}(k)$	Solving augmented ODE $\mathcal{O}(dN)$	Solving ODE $\mathcal{O}(N)$
RSGM	Score matching $\mathcal{O}(d)$ or $\mathcal{O}(1)$	Solving augmented ODE $\mathcal{O}(dN)$	Solving SDE $\mathcal{O}(N^*)$

(2021) introducing Moser flows, whose main appeal being that it circumvents the need to solve an ODE in the training process. We refer to Appendix J for an in-depth discussion on the links between our work and Moser flows.

Optimal transport on manifolds. Another line of work has developed flows on manifolds using tools from optimal transport. Sei (2013) introduced a flow that is given by $f_\theta : x \mapsto \exp_x(\nabla \psi_\theta^c)$ with ψ_θ^c a c-convex function and $c = d_{\mathcal{M}}^2$ the squared geodesic distance. This approach is motivated by the fact that the optimal transport map takes such an expression (Ambrosio, 2003). These methods operate directly on the manifold, similarly to CNFs, yet in contrast they are *discrete* in time. The benefits of this approach depend on the specific choice of parametric family of c-convex functions (Rezende and Racanière, 2021; Cohen et al., 2021), trading-off expressivity with scalability. In the case of tori and spheres, Rezende et al. (2020) introduced *discrete* Riemannian flows based on Möbius transformations and spherical splines.

5 Experiments

In this section we benchmark the empirical performance of RSGMs along with other manifold-valued methods introduced in Section 4. We also compare to a 'Stereographic' score-based model, introduced in Appendix N. First, we assess their modelling capacity on earth and climate science spherical data. Then, we test the methods scalability with respect to manifold dimensions with a synthetic experiment on the torus $\mathbb{T}^d$. Eventually, we evaluate the models' regularity and time complexity with a synthetic $SO_3(\mathbb{R})$ target. Experimental details are provided in Appendix O.

5.1 Earth and climate science datasets on the sphere

We start by evaluating RSGMs on a collection of simple datasets, each containing an empirical distribution of occurrences of earth and climate science events on the surface of the earth. These events are: volcanic eruptions (NGDC/WDS), earthquakes (NGDC/WDS), floods (Brakenridge, 2017) and wild fires (EOSDIS, 2020). We compare to previous baseline methods: Riemannian Continuous Normalizing Flows (Mathieu and Nickel, 2020), Moser Flows (Rozen et al., 2021) and a mixture of Kent distributions (Peel et al., 2001). Additionally, we consider a standard SGM on the 2D plane followed by the inverse stereographic projection which induces a density on the sphere (Gemici et al., 2016). We evaluate the log-likelihood of each model, extending to the manifold setting the likelihood computation techniques of SGMs, see Appendix C. We observe from Table 4,

Table 4: Negative log-likelihood scores for each method on the earth and climate science datasets. Bold indicates best results (up to statistical significance). Means and confidence intervals are computed over 5 different runs. Novel methods are shown with blue shading.

Method	Volcano	Earthquake	Flood	Fire
Mixture of Kent	$-0.80_{\pm 0.47}$	$0.33_{\pm 0.05}$	$0.73_{\pm 0.07}$	$-1.18_{\pm 0.06}$
Riemannian CNF	$\mathbf{-6.05_{\pm 0.61}}$	$0.14_{\pm 0.23}$	$1.11_{\pm 0.19}$	$\mathbf{-0.80_{\pm 0.54}}$
Moser Flow	$-4.21_{\pm 0.17}$	$\mathbf{-0.16_{\pm 0.06}}$	$\mathbf{0.57_{\pm 0.10}}$	$-1.28_{\pm 0.05}$
Stereographic Score-Based	$-3.80_{\pm 0.27}$	$\mathbf{-0.19_{\pm 0.05}}$	$0.59_{\pm 0.07}$	$-1.28_{\pm 0.12}$
Riemannian Score-Based	$-4.92_{\pm 0.25}$	$\mathbf{-0.19_{\pm 0.07}}$	$0.45_{\pm 0.17}$	$-1.33_{\pm 0.06}$
Dataset size	827	6120	4875	12809

| (a) Volcano | (b) Earthquake | (c) Flood | (d) Fire |

Figure 1: Trained score-based generative models on earth sciences data. The learned density is colored green-blue. Blue and red dots represent training and testing datapoints, respectively.

that all benchmarked methods have comparable performance when evaluated on these simple tasks with RSGM performing marginally better on most datasets. However, we empirically notice that Moser flows are slow to train and additionally that both Moser flows and stereographic SGMs are computationally expensive to evaluate.

5.2   Synthetic data on tori

We now move to another manifold, that is the torus $\mathbb{T}^d = \mathbb{S}^1 \times \cdots \times \mathbb{S}^1$, so as to assess the scalability of the different methods with respect to the dimension d. We consider a wrapped Gaussian target distribution on $\mathbb{T}^d$ with a random mean and unit variance. Moser flows' (Rozen et al., 2021) loss involves a regularization term which involves an integral over the manifold, approximated by a Monte Carlo (MC) estimator with uniform proposal. This term regularizes Moser flows towards probability measures, i.e. with unit volume. We thus expect Moser flows to fail in high-dimension as the number of samples K required for the MC estimator to be accurate will grows as $\mathcal{O}(e^d)$, and the memory required to compute this estimator grows either in $\mathcal{O}(Kd)$ for exact divergences or $\mathcal{O}(K)$ for approximated divergences (see Table 3).

In Figure 2, we observe that RSGMs are able to fit well the target distribution even in high dimension, with a linear or constant computational cost—depending on the divergence estimator. In contrast, Moser flows scale poorly with the dimension, to the extent that we are unable to train them for $d \geq 10$. This is due to the combination of the complexity which grows linearly with both the dimension d and the number of MC samples K, which itself ought to grow exponentially with d—as discussed in the previous paragraph. This is illustrated by the gap between the 'Moser' and 'ODE' likelihoods which increases with the manifold dimension (see left Figure 2).

Figure 2: Comparison of Moser flows and RSGMs training speed and performance on the synthetic high-dimension torus task. Moser flows trained with $\lambda_{\min} = 1$. We report two likelihoods, the 'Moser' closed form density—not guaranteed to be normalized—and the 'ODE' likelihood given by solving an augmented ODE (as in CNFs) with the vector field induced by the Moser flow density—which is guaranteed to have unit volume.

(a) Histograms of $SO_3(\mathbb{R})$ samples from a target mixture distribution with $M = 4$ components, represented via their Euler angles.

(b) RSGMs are much more robust to hyperparameters than Exp-wrapped SGMs. The diffusion coefficient is given by $\sigma(t, \mathbf{X}_t) = \sqrt{\beta(t)}$, $\beta(t) = \beta_0 + (\beta_f - \beta_0)t$.

Figure 3: Trained score-based generative models on synthetic $SO_3(\mathbb{R})$ data.

5.3 Synthetic data on the Special Orthogonal group

In order to demonstrate the broad range of applicability of our model we now turn to the task of density estimation on the special orthogonal group $SO_d(\mathbb{R}) = \{Q \in M_d(\mathbb{R}) : QQ^\top = \mathrm{Id}, \det(Q) = 1\}$. We consider the synthetic dataset consisting of samples in $SO_3(\mathbb{R})$ from a mixture of wrapped normal distributions with M components.

We compare RSGMs against Moser flows and a wrapped-exponential baseline inspired by Falorsi et al. (2019)—where we parametrize a standard Euclidean SGM on $\mathfrak{so}(3)$ that is then pushed-forward on $SO_3(\mathbb{R})$. RSGMs are trained using the $\ell_{t|0}$ (DSM) loss with the Varadhan approximation (see Table 2). From Table 5 we observe that, RSGMs perform consistently, whether the target distribution has few or many mixture components M, as opposed to Exp-wrapped SGMs and Moser flows which only perform well in some range of M. Similarly to Section 5.2, we find Moser flows to be much slower to train due to the large number of Monte Carlo samples needed in the regularizer ($K = 10^4$). We also note from Table 5 that the number of score network evaluations (NFE) is significantly lower for RSGMs, and is particularly detrimental for Moser flows ($\gg 10^3$).

Table 5: Test log-likelihood and associated number of function evaluations (NFE) in 10^3 on the synthetic mixture distribution with M components on $SO_3(\mathbb{R})$. Bold indicates best results (up to statistical significance). Means and standard deviations are computed over 5 different runs. Novel methods are shown with blue shading.

Method	$M = 16$		$M = 32$		$M = 64$	
	log-likelihood	NFE	log-likelihood	NFE	log-likelihood	NFE
Moser Flow	$0.85_{\pm 0.03}$	$2.3_{\pm 0.5}$	$0.17_{\pm 0.03}$	$2.3_{\pm 0.9}$	$\mathbf{-0.49_{\pm 0.02}}$	$7.3_{\pm 1.4}$
Exp-wrapped SGM	$\mathbf{0.87_{\pm 0.04}}$	$0.5_{\pm 0.1}$	$0.16_{\pm 0.03}$	$0.5_{\pm 0.0}$	$-0.58_{\pm 0.04}$	$0.5_{\pm 0.0}$
RSGM	$\mathbf{0.89_{\pm 0.03}}$	$\mathbf{0.1_{\pm 0.0}}$	$\mathbf{0.20_{\pm 0.03}}$	$\mathbf{0.1_{\pm 0.0}}$	$\mathbf{-0.49_{\pm 0.02}}$	$\mathbf{0.1_{\pm 0.0}}$

6 Discussion and limitations

In this paper we introduced Riemannian Score-Based Generative Models (RSGMs), a class of deep generative models that represent target densities supported on compact manifolds, as the time-reversal of a Brownian motion. The main benefits of our method stems from its scalability to high dimensions, its applicability to a broad class of manifolds due to the diversity of available loss functions, its robustness and crucially its capacity to model complex datasets. We also provided theoretical guarantees on the convergence of RSGMs.

In future work, we would like to experimentally explore our generic approach to non-compact manifolds following Appendix M. Another promising extension concerns stochastic control on manifolds and more precisely, deriving efficient algorithms to solve Schrödinger bridges in the same spirit as De Bortoli et al. (2021) on Euclidean state spaces.

References

Allgower, E. L. and Georg, K. (2012). *Numerical Continuation Methods: An Introduction*, volume 13. Springer Science & Business Media.

Ambrosio, L. (2003). Optimal transport maps in Monge-Kantorovich problem. *arXiv preprint arXiv:0304389v1*.

Atkinson, K. and Han, W. (2012). *Spherical Harmonics and Approximations on the Unit Sphere: An Introduction*, volume 2044. Springer Science & Business Media.

Bakry, D., Gentil, I., and Ledoux, M. (2014). *Analysis and Geometry of Markov Diffusion Operators*, volume 348. Springer.

Bao, F., Li, C., Zhu, J., and Zhang, B. (2022). Analytic-dpm: an analytic estimate of the optimal reverse variance in diffusion probabilistic models. *arXiv preprint arXiv:2201.06503*.

Barfoot, T., Forbes, J. R., and Furgale, P. T. (2011). Pose estimation using linearized rotations and quaternion algebra. *Acta Astronautica*, 68(1):101–112.

Batzolis, G., Stanczuk, J., Schönlieb, C.-B., and Etmann, C. (2021). Conditional image generation with score-based diffusion models. *arXiv preprint arXiv:2111.13606*.

Bismut, J.-M. (1984). Large deviations and the Malliavin calculus. *Birkhauser Prog. Math.*, 45.

Bose, J., Smofsky, A., Liao, R., Panangaden, P., and Hamilton, W. (2020). Latent variable modelling with hyperbolic normalizing flows. In *International Conference on Machine Learning*.

Bradbury, J., Frostig, R., Hawkins, P., Johnson, M. J., Leary, C., Maclaurin, D., Necula, G., Paszke, A., VanderPlas, J., Wanderman-Milne, S., and Zhang, Q. (2018). JAX: composable transformations of Python+NumPy programs.

Brakenridge, G. (2017). Global active archive of large flood events. http://floodobservatory.colorado.edu/Archives/index.html.

Brehmer, J. and Cranmer, K. (2020). Flows for simultaneous manifold learning and density estimation. *arXiv preprint arXiv:2003.13913*.

Caterini, A. L., Loaiza-Ganem, G., Pleiss, G., and Cunningham, J. P. (2021). Rectangular flows for manifold learning. *arXiv preprint arXiv:2106.01413*.

Cattiaux, P., Conforti, G., Gentil, I., and Léonard, C. (2021). Time reversal of diffusion processes under a finite entropy condition. *arXiv preprint arXiv:2104.07708*.

Chavel, I. (1984). *Eigenvalues in Riemannian Geometry*. Academic press.

Chen, X., Li, X. M., and Wu, B. (2021). Logarithmic heat kernels: estimates without curvature restrictions. *arXiv preprint arXiv:2106.02746*.

Chen, Y., Georgiou, T., and Pavon, M. (2016). Entropic and displacement interpolation: a computational approach using the Hilbert metric. *SIAM Journal on Applied Mathematics*, 76(6):2375–2396.

Cheng, X., Zhang, J., and Sra, S. (2022). Theory and algorithms for diffusion processes on Riemannian manifolds. *arXiv preprint arXiv:2204.13665*.

Choi, K., Meng, C., Song, Y., and Ermon, S. (2021). Density ratio estimation via infinitesimal classification. *arXiv preprint arXiv:2111.11010*.

Chung, H., Sim, B., and Ye, J. C. (2021). Come-closer-diffuse-faster: Accelerating conditional diffusion models for inverse problems through stochastic contraction. *arXiv preprint arXiv:2112.05146*.

Cohen, S., Amos, B., and Lipman, Y. (2021). Riemannian convex potential maps. *arXiv preprint arXiv:2106.10272*.

Croke, C. B. (1980). Some isoperimetric inequalities and eigenvalue estimates. In *Annales scientifiques de l'École normale supérieure*, volume 13, pages 419–435.

De Bortoli, V., Thornton, J., Heng, J., and Doucet, A. (2021). Diffusion Schrödinger bridge with applications to score-based generative modeling. In *Advances in Neural Information Processing Systems*.

Dhariwal, P. and Nichol, A. (2021). Diffusion models beat GAN on image synthesis. *arXiv preprint arXiv:2105.05233*.

Dormand, R. J. and Prince, J. P. (1980). A family of embedded Runge-Kutta formulae. *Journal of Computational and Applied Mathematics*, pages 19–26.

EOSDIS (2020). Land, atmosphere near real-time capability for eos (lance) system operated by nasa's earth science data and information system (esdis). https://earthdata.nasa.gov/earth-observation-data/near-real-time/firms/active-fire-data.

Falorsi, L. (2021). Continuous normalizing flows on manifolds. *arXiv:2104.14959*.

Falorsi, L., de Haan, P., Davidson, T. R., and Forré, P. (2019). Reparameterizing distributions on lie groups. In *International Conference on Artificial Intelligence and Statistics*, pages 3244–3253.

Falorsi, L. and Forré, P. (2020). Neural ordinary differential equations on manifolds. *arXiv preprint arXiv:2006.06663*.

Federer, H. (2014). *Geometric Measure Theory*. Springer.

Feiten, W., Lang, M., and Hirche, S. (2013). Rigid motion estimation using mixtures of projected Gaussians. In *International Conference on Information Fusion*, pages 1465–1472. IEEE.

Fisher, R. A. (1953). Dispersion on a sphere. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 217(1130):295–305.

Ganea, O.-E., Huang, X., Bunne, C., Bian, Y., Barzilay, R., Jaakkola, T. S., and Krause, A. (2022). Independent SE(3)-Equivariant models for end-to-end rigid protein docking. In *International Conference on Learning Representations*.

García-Zelada, D. and Huguet, B. (2021). Brenier–Schrödinger problem on compact manifolds with boundary. *Stochastic Analysis and Applications*, pages 1–29.

Gemici, M. C., Rezende, D., and Mohamed, S. (2016). Normalizing flows on Riemannian manifolds. *arXiv preprint arXiv:1611.02304*.

Grathwohl, W., Chen, R. T. Q., Bettencourt, J., and Duvenaud, D. (2019). Scalable reversible generative models with free-form continuous dynamics. In *International Conference on Learning Representations*.

Gretton, A., Borgwardt, K. M., Rasch, M. J., Schölkopf, B., and Smola, A. (2012). A kernel two-sample test. *Journal of Machine Learning Research*, 13(null):723–773.

Grigor'yan, A. (1999). Estimates of heat kernels on Riemannian manifolds. *London Math. Soc. Lecture Note Ser*, 273:140–225.

Gross, L. (1992). Logarithmic sobolev inequalities on lie groups. *Illinois journal of mathematics*, 36(3):447–490.

Gunther, M. (1991). Isometric embeddings of Riemannian manifolds, Kyoto, 1990. In *Proc. Intern. Congr. Math.*, pages 1137–1143. Math. Soc. Japan.

Hauberg, S. (2018). Directional Statistics with the Spherical Normal Distribution. In *2018 21st International Conference on Information Fusion (FUSION)*, pages 704–711, Cambridge, United Kingdom. IEEE.

Haussmann, U. G. and Pardoux, E. (1986). Time reversal of diffusions. *The Annals of Probability*, 14(4):1188–1205.

He, Y. (2013). A lower bound for the first eigenvalue in the Laplacian operator on compact Riemannian manifolds. *Journal of Geometry and Physics*, 71:73–84.

Ho, J., Jain, A., and Abbeel, P. (2020). Denoising diffusion probabilistic models. *Advances in Neural Information Processing Systems*.

Hou, B., Miolane, N., Khanal, B., Lee, M. C. H., Alansary, A., McDonagh, S., Hajnal, J. V., Rueckert, D., Glocker, B., and Kainz, B. (2018). Computing CNN Loss and Gradients for Pose Estimation with Riemannian Geometry. In Frangi, A. F., Schnabel, J. A., Davatzikos, C., Alberola-López, C., and Fichtinger, G., editors, *Medical Image Computing and Computer Assisted Intervention – MICCAI 2018*, pages 756–764, Cham. Springer International Publishing.

Hsu, E. (1999). Estimates of derivatives of the heat kernel on a compact Riemannian manifold. *Proceedings of the american mathematical society*, 127(12):3739–3744.

Hsu, E. P. (2002). *Stochastic Analysis on Manifolds*. Number 38. American Mathematical Society.

Hutchinson, M. F. (1989). A stochastic estimator of the trace of the influence matrix for laplacian smoothing splines. *Communications in Statistics-Simulation and Computation*, 18(3):1059–1076.

Hyvärinen, A. (2005). Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6(4).

Ikeda, N. and Watanabe, S. (1989). *Stochastic Differential Equations and Diffusion Processes*, volume 24 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam; Kodansha, Ltd., Tokyo, second edition.

Jolicoeur-Martineau, A., Piché-Taillefer, R., Tachet des Combes, R., and Mitliagkas, I. (2021). Adversarial score matching and improved sampling for image generation. *International Conference on Learning Representations*.

Jones, P. W., Maggioni, M., and Schul, R. (2008). Manifold parametrizations by eigenfunctions of the Laplacian and heat kernels. *Proceedings of the National Academy of Sciences of the United States of America*, 105(6):1803–1808.

Jørgensen, E. (1975). The central limit problem for geodesic random walks. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 32(1-2):1–64.

Karpatne, A., Ebert-Uphoff, I., Ravela, S., Babaie, H. A., and Kumar, V. (2018). Machine learning for the geosciences: Challenges and opportunities. *IEEE Transactions on Knowledge and Data Engineering*, 31(8):1544–1554.

Kawar, B., Vaksman, G., and Elad, M. (2021a). Snips: Solving noisy inverse problems stochastically. *arXiv preprint arXiv:2105.14951*.

Kawar, B., Vaksman, G., and Elad, M. (2021b). Stochastic image denoising by sampling from the posterior distribution. *arXiv preprint arXiv:2101.09552*.

Kent, J. T. (1982). The Fisher-Bingham distribution on the sphere. *Journal of the Royal Statistical Society: Series B (Methodological)*, 44(1):71–80.

Kingma, D. P. and Ba, J. (2015). Adam: A Method for Stochastic Optimization. *arXiv:1412.6980 [cs]*.

Klimovskaia, A., Lopez-Paz, D., Bottou, L., and Nickel, M. (2020). Poincaré maps for analyzing complex hierarchies in single-cell data. *Nature communications*, 11(1):1–9.

Kloeden, P. and Platen, E. (2011). *Numerical Solution of Stochastic Differential Equations*. Stochastic Modelling and Applied Probability. Springer Berlin Heidelberg.

Köhler, J., Klein, L., and Noé, F. (2020). Equivariant Flows: Exact likelihood generative learning for symmetric densities. *arXiv:2006.02425 [physics, stat]*.

Kullback, S. (1997). *Information Theory and Statistics*. Dover Publications, Inc., Mineola, NY. Reprint of the second (1968) edition.

Kurtz, T. G., Pardoux, É., and Protter, P. (1995). Stratonovich stochastic differential equations driven by general semimartingales. In *Annales de l'IHP Probabilités et statistiques*, volume 31, pages 351–377.

Kuwada, K. (2012). Convergence of time-inhomogeneous geodesic random walks and its application to coupling methods. *The Annals of Probability*, 40(5):1945–1979.

Lee, J. (2010). *Introduction to Topological Manifolds*, volume 202. Springer Science & Business Media.

Lee, J. M. (2006). *Riemannian Manifolds: An Introduction to Curvature*, volume 176. Springer Science & Business Media.

Lee, J. M. (2013). Smooth manifolds. In *Introduction to Smooth Manifolds*, pages 1–31. Springer.

Lee, J. M. (2018). *Introduction to Riemannian manifolds*. Springer.

Lee, S.-g., Kim, H., Shin, C., Tan, X., Liu, C., Meng, Q., Qin, T., Chen, W., Yoon, S., and Liu, T.-Y. (2021). Priorgrad: Improving conditional denoising diffusion models with data-driven adaptive prior. *arXiv preprint arXiv:2106.06406*.

Leobacher, G. and Steinicke, A. (2021). Existence, uniqueness and regularity of the projection onto differentiable manifolds. *Annals of Global Analysis and Geometry*, 60(3):559–587.

Léonard, C. (2012a). From the Schrödinger problem to the Monge–Kantorovich problem. *Journal of Functional Analysis*, 262(4):1879–1920.

Léonard, C. (2012b). Girsanov theory under a finite entropy condition. In *Séminaire de Probabilités XLIV*, pages 429–465. Springer.

Léonard, C., Rœlly, S., Zambrini, J.-C., et al. (2014). Reciprocal processes: a measure-theoretical point of view. *Probability Surveys*, 11:237–269.

Li, P. (1986). Large time behavior of the heat equation on complete manifolds with non-negative ricci curvature. *Annals of Mathematics*, 124(1):1–21.

Liptser, R. S. and Shiryaev, A. N. (2001). *Statistics of Random Processes. I*, volume 5 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, expanded edition. General theory, Translated from the 1974 Russian original by A. B. Aries, Stochastic Modelling and Applied Probability.

Lui, Y. M. (2012). Advances in matrix manifolds for computer vision. *Image and Vision Computing*, 30(6-7):380–388.

Mardia, K. V., Hughes, G., Taylor, C. C., and Singh, H. (2008). A multivariate von Mises distribution with applications to bioinformatics. *Canadian Journal of Statistics*, 36(1):99–109.

Mathieu, E., Lan, C. L., Maddison, C. J., Tomioka, R., and Teh, Y. W. (2019). Continuous hierarchical representations with Poincaré variational auto-encoders. *arXiv preprint arXiv:1901.06033*.

Mathieu, E. and Nickel, M. (2020). Riemannian continuous normalizing flows. *arXiv preprint arXiv:2006.10605*.

Miolane, N., Guigui, N., Brigant, A. L., Mathe, J., Hou, B., Thanwerdas, Y., Heyder, S., Peltre, O., Koep, N., Zaatiti, H., Hajri, H., Cabanes, Y., Gerald, T., Chauchat, P., Shewmake, C., Brooks, D., Kainz, B., Donnat, C., Holmes, S., and Pennec, X. (2020). Geomstats: A python package for riemannian geometry in machine learning. *Journal of Machine Learning Research*, 21(223):1–9.

Nagano, Y., Yamaguchi, S., Fujita, Y., and Koyama, M. (2019). A wrapped normal distribution on hyperbolic space for gradient-based learning. In *International Conference on Machine Learning*.

Navarro, A. K., Frellsen, J., and Turner, R. E. (2017). The multivariate generalised von Mises distribution: inference and applications. In *AAAI Conference on Artificial Intelligence*.

(NGDC/WDS), N. G. D. C. . W. D. S. (2022a). Ncei/wds global significant earthquake database. https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ngdc.mgg.hazards:G012153.

(NGDC/WDS), N. G. D. C. . W. D. S. (2022b). Ncei/wds global significant volcanic eruptions database. https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ngdc.mgg.hazards:G10147.

Nichol, A. and Dhariwal, P. (2021). Improved denoising diffusion probabilistic models. *arXiv preprint arXiv:2102.09672*.

Papamakarios, G., Nalisnick, E., Rezende, D. J., Mohamed, S., and Lakshminarayanan, B. (2019). Normalizing flows for probabilistic modeling and inference. *arXiv preprint arXiv:1912.02762*.

Peel, D., Whiten, W. J., and McLachlan, G. J. (2001). Fitting mixtures of Kent distributions to aid in joint set identification. *Journal of the American Statistical Association*, 96(453):56–63.

Pennec, X. (2006). Intrinsic Statistics on Riemannian Manifolds: Basic Tools for Geometric Measurements. *Journal of Mathematical Imaging and Vision*, 25(1):127–154.

Prokudin, S., Gehler, P., and Nowozin, S. (2018). Deep directional statistics: Pose estimation with uncertainty quantification. In *European Conference on Computer Vision (ECCV)*.

Revuz, D. and Yor, M. (1999). *Continuous Martingales and Brownian Motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition.

Rezende, D. J., Papamakarios, G., Racanière, S., Albergo, M. S., Kanwar, G., Shanahan, P. E., and Cranmer, K. (2020). Normalizing flows on tori and spheres. *arXiv preprint arXiv:2002.02428*.

Rezende, D. J. and Racanière, S. (2021). Implicit Riemannian concave potential maps. *arXiv:2110.01288*.

Roy, D. M., Kemp, C., Mansinghka, V., and B Tenenbaum, J. (2007). Learning annotated hierarchies from relational data.

Rozen, N., Grover, A., Nickel, M., and Lipman, Y. (2021). Moser flow: Divergence-based generative modeling on manifolds. *Advances in Neural Information Processing Systems*.

Saloff-Coste, L. (1994). Precise estimates on the rate at which certain diffusions tend to equilibrium. *Mathematische Zeitschrift*, 217(1):641–677.

Santambrogio, F. (2017). {Euclidean, metric, and Wasserstein} gradient flows: an overview. *Bulletin of Mathematical Sciences*, 7(1):87–154.

Schrödinger, E. (1932). Sur la théorie relativiste de l'électron et l'interprétation de la mécanique quantique. *Annales de l'Institut Henri Poincaré*, 2(4):269–310.

Sei, T. (2013). A Jacobian inequality for gradient maps on the sphere and its application to directional statistics. *Communications in Statistics-Theory and Methods*, 42(14):2525–2542.

Senanayake, R. and Ramos, F. (2018). Directional grid maps: modeling multimodal angular uncertainty in dynamic environments. In *2018 IEEE/RSJ International Conference on Intelligent Robots and Systems (IROS)*, pages 3241–3248. IEEE.

Shapovalov, M. V. and Dunbrack Jr, R. L. (2011). A smoothed backbone-dependent rotamer library for proteins derived from adaptive kernel density estimates and regressions. *Structure*, 19(6):844–858.

Sinha, A., Song, J., Meng, C., and Ermon, S. (2021). D2c: Diffusion-denoising models for few-shot conditional generation. *arXiv preprint arXiv:2106.06819*.

Song, Y. and Ermon, S. (2019). Generative modeling by estimating gradients of the data distribution. In *Advances in Neural Information Processing Systems*.

Song, Y. and Ermon, S. (2020). Improved techniques for training score-based generative models. In *Advances in Neural Information Processing Systems*.

Song, Y., Sohl-Dickstein, J., Kingma, D. P., Kumar, A., Ermon, S., and Poole, B. (2021). Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*.

Steyvers, M. and Tenenbaum, J. B. (2005). The large-scale structure of semantic networks: Statistical analyses and a model of semantic growth. *Cognitive science*, 29(1):41–78.

Sun, Y., Flammarion, N., and Fazel, M. (2019). Escaping from saddle points on Riemannian manifolds. *Advances in Neural Information Processing Systems*, 32.

Urakawa, H. (2006). Convergence rates to equilibrium of the heat kernels on compact Riemannian manifolds. *Indiana University Mathematics Journal*, pages 259–288.

Vargas, F., Thodoroff, P., Lawrence, N. D., and Lamacraft, A. (2021). Solving Schrödinger bridges via maximum likelihood. *arXiv preprint arXiv:2106.02081*.

Vincent, P. (2011). A connection between score matching and denoising autoencoders. *Neural Computation*, 23(7):1661–1674.

Watson, D., Ho, J., Norouzi, M., and Chan, W. (2021). Learning to efficiently sample from diffusion probabilistic models. *arXiv preprint arXiv:2106.03802*.