Sparse Gaussian Process Hyperparameters: Optimize or Integrate?

Abstract

The kernel function and its hyperparameters are the central model selection choice in a Gaussian process [Rasmussen and Williams, 2006]. Typically, the hyperparameters of the kernel are chosen by maximising the marginal likelihood, an approach known as *Type-II maximum likelihood* (ML-II). However, ML-II does not account for hyperparameter uncertainty, and it is well-known that this can lead to severely biased estimates and an underestimation of predictive uncertainty. While there are several works which employ a fully Bayesian characterisation of GPs, relatively few propose such approaches for the sparse GPs paradigm. In this work we propose an algorithm for sparse Gaussian process regression which leverages MCMC to sample from the hyperparameter posterior within the variational inducing point framework of [Titsias, 2009]. This work is closely related to Hensman et al. [2015b], but side-steps the need to sample the inducing points, thereby significantly improving sampling efficiency in the Gaussian likelihood case. We compare this scheme against natural baselines in literature along with stochastic variational GPs (SVGPs) along with an extensive computational analysis.

1 Introduction

Gaussian processes (GPs) are a prominent class of models for supervised learning which can quantify uncertainty and incorporate inductive biases in function space via the kernel function. Hand-crafting a kernel function is a powerful way to incorporate prior knowledge. In many instances not all properties of a kernel function can be specified from prior knowledge alone, and parameters are chosen via ML-II. However, defining a complex kernel function with a large number of hyperparameters can make the marginal likelihood prone to multiple local optima and overfitting. Further, several local optima may correspond to priors that do not sensibly model the data. Weakly identified hyperparameters can manifest in flat ridges in the marginal likelihood surface[1] making gradient based optimisation extremely sensitive to starting values [Warnes and Ripley, 1987]. Overall, the ML-II point estimates for the hyperparameters are subject to high variability and underestimate prediction uncertainty.

The problem of ridges in the marginal likelihood surface also does not necessarily go away as more observations are collected. For example, if f_1 and f_2 are Brownian motions, $\sigma f_1(x/\ell)$ is equal in distribution to $\sqrt{\alpha}\sigma f_2(x/\alpha\ell)$, which means observations do not provide any information about the product $\sigma\ell$. More generally, for a greater class of kernels, including the Matérn–$1/2$ kernel, $\sigma f_1(x/\ell)$ is *equivalent* to $\sqrt{\alpha}\sigma f_2(x/\alpha\ell)$, which means that is not possible to consistently estimate the product $\sigma\ell$ from data, even if infinitely many observations are collected in a fixed domain [Chapter 6, Stein, 1999]. This implies that one cannot estimate the individual hyperparameters (σ, ℓ) consistently. It also motivates why there can be benefits to estimating the hyperparameter posterior even in large data regimes, and ML-II may be insufficient. A more satisfactory treatment of hyperparameters involves placing a prior over the hyperparameters and performing Bayesian inference to compute a

---

[1]where different combinations of hyperparameters give very similar marginal likelihood values

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

Figure 1: Negative log marginal likelihood surface as a function of two hyperparameters: σ_f^2 and l for a squared exponential kernel and 1d function. The red cross indicates the true hyperparameters. The hyperparameters selected via gradien- based optimisation are sensitive to the initialisation due to the long ridge of almost identical height at values of the hyperparameters not concordant with the ground truth.

(hyper)posterior. For large datasets, this motivates using scalable GP inference (e.g. sparse methods) in conjunction with Markov chain Monte Carlo (MCMC) for the hyperparametrs.

The Bayesian treatment of weakly identified hyperparameters may also be fraught with difficulties. Gradient-based samplers like Hamiltonian Monte Carlo [Neal et al., 2011] and its variants have difficulty navigating regions of high curvature and flat ridges where the gradient offers no information for transition [Betancourt, 2017]. This leads to over-concentration of samples from the flat region. (Usually, this can at least partially be rectified with informative priors.) These pathologies are also typical of other hierarchical models [Betancourt and Girolami, 2015]. Figure 1 shows that the GP marginal likelihood surface can manifest these pathologies. The evidence lower bound (ELBO) used in hyperparameter selection for variational sparse GPs relying on inducing points inherits similar, or even less favourable, characteristics to the exact marginal likelihood. As a result, for weakly informative priors, gradient based samplers are susceptible to getting *stuck* at the boundary of these pathological regions hence biasing the sample estimates. Fortunately, the effective sample size metric used for diagnoising mixing in MCMC is indicative of this behaviour when directly observing the phase space of the target distribution is infeasible in high-dimensions.

Historical justification for ML-II (also called the *evidence* framework) comes from [MacKay, 1994] which highlighted several conditions for ML-II to yield reasonable estimates for the hyperparameters. Crucially, the evidence is unlikely to manifest multiple local optima for a model well-matched to the data and with a high signal-to-noise ratio. Transferring this insight to the Gaussian process paradigm we show how the evidence can have a significant tail or no well-defined maximum in settings with a low signal-to-noise ratio which arises with high aleatoric uncertainty or sparse data, frequently both. In these settings, a point estimate may not adequately summarise the hyperparameter posterior and the benefits of marginalisation stand out. The situation is only exacerbated in high-dimensions (i.e. when there are many hyperparameter) where increasingly more volume of the posterior is captured in a thin-shell making the density peak extremely unrepresentative of the posterior.

Unlike the likelihood of parameters, the marginal likelihood inherently contains a trade-off between the data-fit and complexity penalty term. This is one of the main properties that makes the marginal likelihood objective a viable choice for model selection [Rasmussen and Williams, 2006]. For example, for the Gaussian process regression model,

$$y_n = f(x_n) + \epsilon_n, \ \epsilon_n \sim \mathcal{N}(0, \sigma^2) \ f \sim \mathcal{GP}(0, k_\theta) \tag{1}$$

the marginal likelihood takes the form,

$$\log p(\boldsymbol{y}|\boldsymbol{\theta}) = \log \int p(\boldsymbol{y}|f)p(f)df = c \underbrace{-\tfrac{1}{2}\boldsymbol{y}^T(K_\theta + \sigma^2 I)^{-1}\boldsymbol{y}}_{\text{data fit term}} \underbrace{-\tfrac{1}{2}|K_\theta + \sigma^2 I|}_{\text{complexity penalty}} \tag{2}$$

where c is a constant, $p(\boldsymbol{y}|f)$ denotes the data likelihood and $\boldsymbol{\theta}$ denotes kernel hyperparameters. This trade-off is a well-established idea which embodies the *automatic Occam's razor* effect [Rasmussen and Ghahramani, 2001] where models well-suited to the data are automatically selected just by using the marginal likelihood objective.

This may seem to contradict earlier claims regarding overfitting, but by shying away from dealing with the hyperparameter posterior we risk overfitting even with the marginal likelihood approach. In other words, the evidence framework subdues the overfitting effect induced by the canonical maximum likelihood approach which does not have any complexity penalty term. The parameters in this approach are free to fit the data as well as possible making them prone to overfitting and poor generalisation. Overparameterized kernels based on neural networks like deep kernel models [Wilson et al., 2016] are well known to exacerbate overfitting [Ober et al., 2021].

The main motivation for this work is to highlight that fully Bayesian schemes in sparse Gaussian process models are practically beneficial. While several works in the literature employ fully Bayesian scheme of integrating out the hyperparameters (see table 1), in this work we attempt to analyse them in an orthogonal direction, focusing on comparison with the evidence framework and other benchmarks by extending the main sparse variational formulation in the literature Titsias [2009]. We present a generalised inference scheme for fully Bayesian GP regression and counteract some of the computational cost of sampling both inducing variables and hyperparameters by deriving a *doubly collapsed* bound which selects the optimal distribution over the inducing points analytically and targets the kernel hyperparameters with HMC.

2   Related Work

Table 1: Existing literature on fully Bayesian inference in GPs, sparse GPs and generic likelihoods.

Index	Reference	Sparse	Posterior ($\boldsymbol{f}/\boldsymbol{u}$)	Posterior ($\boldsymbol{\theta}$)	Methods
1.	Murray and Adams [2010]	✗	sampling / NA	sampling	Slice sampling
2.	Filippone et al. [2013]	✗	sampling / NA	sampling	MH + HMC + MA-LA
3.	Filippone and Girolami [2014]	✗	Gaussian / NA	sampling	Deterministic + Pseudo-Marginal
4.	Hensman et al. [2015b]	✓	Gaussian / sampling	sampling	HMC
5.	Bui et al. [2018]	✓	Gaussian / (sampling & VFE)	sampling / VFE	MCMC
6.	Lalchand and Rasmussen [2020]	✗	Gaussian / NA	sampling / VFE	NUTS / VI
7.	Rossi et al. [2021]	✓	Gaussian / sampling	sampling	SG-HMC
8.	Simpson et al. [2021]	✗	Gaussian / NA	sampling	NUTS / Nested Sampling
9.	This work	✓	Gaussian / (sampling & VFE)	sampling / VFE	HMC / VI / Structured VI

Fully Bayesian Gaussian processes have been used by several authors spawning several variants. In early accounts, Neal [1998], Williams and Rasmussen [1996] explore the integration over covariance hyperparameters using HMC in the regression setting. Barber and Williams [1997] extend this to the classification setting using HMC for sampling in the hyperparameter space and Laplace approximation for the integrand over function values. Murray and Adams [2010] and Filippone et al. [2013] focused on MCMC schemes to sample covariance hyperparameters in conjunction with latent function values, mainly mitigating the coupling effect through reparameterisation. Hensman et al. [2015b] considered joint sampling of inducing variables and hyperparameters from the optimal variational posterior distribution while [Bui et al., 2018] consider inference schemes for fully Bayesian sparse GPs in a streaming setting. More recently, Rossi et al. [2021] studied fully Bayesian sparse GPs using SG-HMC. Rossi et al. [2021] modify the generative model by adding a prior over the inducing inputs, and peform inference using SG-HMC over both Z and θ. We list the most recent works in Table 1.

3   Background

Let $f \sim \mathcal{GP}(0, k_{\boldsymbol{\theta}})$ be a Gaussian process prior with covariance function $k_{\boldsymbol{\theta}}$ depending on hyperparameters $\boldsymbol{\theta}$. We are given noisy observations $\boldsymbol{y} = (y_n)_{n=1}^N \subseteq \mathbb{R}$ of $\boldsymbol{f} = (f(\boldsymbol{x}_n))_{n=1}^N$ at input data $X = (\boldsymbol{x}_n)_{n=1}^N \subseteq \mathbb{R}^D$. We consider likelihoods which factorise over the data, like $p(\boldsymbol{y}|f) = \prod_{n=1}^N \mathcal{N}(y_n|f_n, \sigma^2)$ for regression or $p(\boldsymbol{y}|f) = \prod_{n=1}^N \mathcal{B}(y_n|\phi(f_n))$ for binary classification where $\phi(\cdot) = \int_{-\infty}^x \mathcal{N}(a|0, 1) \, \mathrm{d}a$ denotes the probit link function and $\mathcal{B}(y_n|\phi(f_n)) = \phi(f_n)^{y_n}(1 - \phi(f_n))^{(1-y_n)}$ is the Bernoulli distribution. We wish to compute the posterior $p(f|\boldsymbol{y}, X, \boldsymbol{\theta})$. In this section, we recapitulate the canonical inducing variable approximation of $p(f|\boldsymbol{y}, X, \boldsymbol{\theta})$ by Titsias [2009] and its extension to a Bayesian treatment of the hyperparameters.

3.1 Sparse variational inference in Gaussian processes

Following Titsias [2009], we consider a set of inducing variables $\boldsymbol{u} = \{f(\boldsymbol{z}_m)\}_{m=1}^M \subseteq \mathbb{R}$ at inducing inputs $Z = \{\boldsymbol{z}_m\}_{m=1}^M, \boldsymbol{z}_m \in \mathbb{R}^d$. The complete generative model can then be factored as,

$$p(\boldsymbol{y}, f, \boldsymbol{u}|\boldsymbol{\theta}) = p(\boldsymbol{y}|f, \boldsymbol{\theta})p(f|\boldsymbol{u}, \boldsymbol{\theta})p(\boldsymbol{u}|\boldsymbol{\theta}). \tag{3}$$

We approximate the posterior $p(f, \boldsymbol{u}|\boldsymbol{y}, \boldsymbol{\theta})$ with a variational distribution:

$$p(f, \boldsymbol{u}|\boldsymbol{y}, \boldsymbol{\theta}) \approx q(f, \boldsymbol{u}|\boldsymbol{\theta}) = p(f|\boldsymbol{u}, \boldsymbol{\theta})q(\boldsymbol{u}) \tag{4}$$

where $q(\boldsymbol{u})$ is chosen to minimise the Kullback–Leibler divergence $\mathrm{KL}(q(f|\boldsymbol{\theta}) \| p(f|\boldsymbol{y}, X, \boldsymbol{\theta}))$. Minimising this KL divergence corresponds to maximising the evidence lower bound [Matthews et al., 2016], henceforth called the ELBO:

$$\mathrm{ELBO}(q(\boldsymbol{u}), \boldsymbol{\theta}) = \mathbb{E}_{q(f|\boldsymbol{\theta})}[\log p(\boldsymbol{y}|f, \boldsymbol{\theta})] - \mathrm{KL}(q(\boldsymbol{u}|\boldsymbol{\theta}) \| p(\boldsymbol{u}|\boldsymbol{\theta})).$$

Because the ELBO still depends on $q(\boldsymbol{u})$, this bound is called *uncollapsed*. Hensman et al. [2013, 2015a] let $q(\boldsymbol{u})$ be a Gaussian, which is optimal if the likelihood is Gaussian [Titsias, 2009], approximate the expectation using Monte Carlo, and maximise the ELBO using stochastic optimisation. On the other hand, if the likelihood is Gaussian, Titsias [2009] computes optimal form for $q(\boldsymbol{u})$ directly:

$$q^*(\boldsymbol{u}|\boldsymbol{\theta}) = \mathrm{argmax}_{q(\boldsymbol{u})} \, \mathrm{ELBO}(q(\boldsymbol{u}), \boldsymbol{\theta}) \propto p(\boldsymbol{u}|\boldsymbol{\theta}) \exp \mathbb{E}_{p(f|\boldsymbol{u}, \boldsymbol{\theta})}[\log p(\boldsymbol{y}|f, \boldsymbol{\theta})] \tag{5}$$

Plugging $q^*(\boldsymbol{u}|\boldsymbol{\theta})$ back into Equation (5), the resulting bound is called *collapsed*, because it now only depends on $\boldsymbol{\theta}$ and Z. The collapsed bound is the objective that Titsias [2009] proposes.

$$\log p(\boldsymbol{y}|\boldsymbol{\theta}) \geq \log \mathcal{N}(\boldsymbol{y}; \boldsymbol{0}, K_{nm}K_{mm}^{-1}K_{mn} + \sigma^2 I) - \frac{1}{2\sigma^2}\mathrm{Tr}(K_{nn} - K_{nm}K_{mm}^{-1}K_{mn}) =: \mathcal{L}_{\boldsymbol{\theta}} \tag{6}$$

Using the collapsed bound, approximate ML-II the consists of finding,

$$\theta^* \in \mathrm{argmax}_{\boldsymbol{\theta}, Z} \, \mathcal{L}_{\boldsymbol{\theta}}. \tag{7}$$

Predictions at new functions values can be made in $O(M^2)$ after an initial cost of $O(NM^2)$. Under certain assumptions on the data generating process, even when $M \ll N$, the approximate posterior closely resembles the posterior, and Equation (6) is a provably accurate approximation to Equation (2) [Burt et al., 2020].

3.2 Bayesian treatment of hyperparameters and sparse methods

The extension of the sparse variational framework to a Bayesian treatment of the hyperparameters has been previously considered by Hensman et al. [2015b]. Extend the generative model with a prior $p(\boldsymbol{\theta})$ over the hyperparameters $\boldsymbol{\theta}$, and let the variational approximation of the posterior $p(f, \boldsymbol{u}, \boldsymbol{\theta}|\boldsymbol{y}, X)$ be $q(f, \boldsymbol{u}, \boldsymbol{\theta}) = p(f|\boldsymbol{u}, \boldsymbol{\theta})q(\boldsymbol{u}, \boldsymbol{\theta})$. The analogue of Equation (5) is

$$\mathrm{ELBO}(q(\boldsymbol{u}, \boldsymbol{\theta})) = \mathbb{E}_{q(f, \boldsymbol{\theta})}[\log p(\boldsymbol{y}|f, \boldsymbol{\theta})] - \mathbb{E}_{q(\boldsymbol{\theta})}[\mathrm{KL}(q(\boldsymbol{u}) \| p(\boldsymbol{u}|\boldsymbol{\theta}))] - \mathrm{KL}(q(\boldsymbol{\theta}) \| p(\boldsymbol{\theta})) \tag{8}$$

and the optimal form for $q(\boldsymbol{u}, \boldsymbol{\theta})$ can again be determined:

$$q^*(\boldsymbol{u}, \boldsymbol{\theta}) \propto p(\boldsymbol{u}, \boldsymbol{\theta}) \exp \mathbb{E}_{p(f|\boldsymbol{u}, \boldsymbol{\theta})}[\log p(\boldsymbol{y}|f, \boldsymbol{\theta})]. \tag{9}$$

The distribution $q^*(\boldsymbol{u}, \boldsymbol{\theta})$ does not have a closed form, and for general likelihoods, Hensman et al. [2015b] propose to approximate the expectation in (9) with quadrature and to sample from $q^*(\boldsymbol{u}, \boldsymbol{\theta})$ using HMC. While this approach is quite general, in the case of Gaussian regression, it vastly increases the dimensionality of the state space over which HMC must be run relative to HMC in GPR, since the $\boldsymbol{u}$ are sampled in addition to the $\boldsymbol{\theta}$. This increases the cost of the procedure, and impact the success of the sampler.

An alternative approach to approximately inferring hyperparameters is to assume a parametric form for $q(\boldsymbol{u}, \boldsymbol{\theta})$ and maximise Equation (8) with respect to the variational parameters. Bui et al. [2018] took such an approach, assuming that $q(\boldsymbol{u}, \boldsymbol{\theta}) = q(\boldsymbol{u})q(\boldsymbol{\theta})$, with both distributions Gaussian. Similar approaches have been applied to variational inference in state-space modelling, sometimes leveraging the optimal form of $q(\boldsymbol{u}|\boldsymbol{\theta})$ discussed earlier.

Table 2: Comparison of a variety of approaches to approximating the posterior over hyperparameters in Gaussian process regression. Compares the quality of the posterior (QUALITY); the time complexity per iteration (TIME/IT.); the memory complexity per iteration (MEM./IT.); the number of parameters and/or variables (PARS/VARS); and whether the approach supports non-Gaussian likelihoods (LIK.).

APPROACH	QUALITY	TIME/IT.	MEM./IT.	PARS/VARS	LIK.
Maximum a posteriori	−	n^3	n^2	n_θ	✗
VI					
Inducing points; non-collapsed	±	nm^2	m^2	$n_\theta^2 + m^2$	✓
Inducing points; collapsed	±	nm^2	m^2	n_θ^2	✗
SAMPLING					
Exact with Gaussian lik.	+	n^3	n^2	n_θ	✗
Exact with non-Gaussian lik.	+	n^3	n^2	$n_\theta + n$	✓
Inducing points; non-collapsed	±	m^3	m^2	$n_\theta + m$	✓
Inducing points; collapsed	±	nm^2	m^2	n_θ	✗

3.3 Making predictions

The predictive posterior distribution for unknown test inputs X^* integrates over the joint posterior,

$$p(\boldsymbol{f}^*|\boldsymbol{y}) \approx \int p(\boldsymbol{f}^*|\boldsymbol{f},\boldsymbol{u},\boldsymbol{\theta})p(\boldsymbol{f}|\boldsymbol{u},\boldsymbol{\theta})q(\boldsymbol{u}|\boldsymbol{\theta})q(\boldsymbol{\theta})d\boldsymbol{f}d\boldsymbol{u}d\boldsymbol{\theta} \tag{10}$$

(where we have suppressed the conditioning over inputs X, X^* for brevity). The inner integral simplifies to $\int p(\boldsymbol{f}^*|\boldsymbol{f},\boldsymbol{u},\boldsymbol{\theta})p(\boldsymbol{f}|\boldsymbol{u},\boldsymbol{\theta})d\boldsymbol{f} = p(\boldsymbol{f}^*|\boldsymbol{u},\boldsymbol{\theta})$. We discuss the predictive posterior in such models in section 4.3.

4 Fully Bayesian SGPR with HMC: Doubly collapsed formulation

In the previous section, we observed that a major drawback of the approach taken in Hensman et al. [2015b] is the need to sample the $\boldsymbol{u}$, which for high-dimensional inputs or in cases where many inducing points are needed could introduce thousands of additional variables to sample. In this section, we leverage the optimal form of $q(\boldsymbol{u}|\boldsymbol{\theta})$ derived in Titsias [2009] to alleviate this sampling problem.

4.1 Collapsing the evidence lower bound (again)

We first derive the lower bound for this formulation and provide pseudo-code for the algorithm in Algorithm 1. As in the previous section, we wish to minimize the KL divergence between the true and approximate posterior, $\mathrm{KL}(q(\boldsymbol{f},\boldsymbol{u},\boldsymbol{\theta})||p(\boldsymbol{f},\boldsymbol{u},\boldsymbol{\theta}\,|\,\boldsymbol{y})$. Subtracting from the marginal likelihood and rearranging,

$$\log p(\boldsymbol{y}) \geq \int q(\boldsymbol{\theta}) \underbrace{\int q(\boldsymbol{u}|\boldsymbol{\theta}) \int p(\boldsymbol{f}|\boldsymbol{u},\boldsymbol{\theta}) \log \frac{p(\boldsymbol{y}|\boldsymbol{f},\boldsymbol{\theta})p(\boldsymbol{u}|\boldsymbol{\theta})}{q(\boldsymbol{u}|\boldsymbol{\theta})} d\boldsymbol{f}d\boldsymbol{u}}_{\mathcal{L}_\theta} d\boldsymbol{\theta} - \mathrm{KL}(q(\boldsymbol{\theta})\,\|\,p(\boldsymbol{\theta})) \tag{11}$$

$$\geq \int q(\boldsymbol{\theta})\mathcal{L}_\theta d\boldsymbol{\theta} - \mathrm{KL}(q(\boldsymbol{\theta})\,\|\,p(\boldsymbol{\theta})) = \int q(\boldsymbol{\theta}) \log \frac{M_\theta p(\boldsymbol{\theta})}{q(\boldsymbol{\theta})} d\boldsymbol{\theta}, \tag{12}$$

where we assign $\mathcal{L}_\theta = \log M_\theta$, which is defined in Equation (6). In Equation (12), we substitute in the optimal marginal distribution $q^*(\boldsymbol{u}|\boldsymbol{\theta})$ derived in Titsias [2009]. We can interpret the RHS of Equation (12) as a negative KL divergence as long as we account for a normalisation constant C for the un-normalised numerator term $M_\theta p(\boldsymbol{\theta})$. Hence, we can re-write the ELBO as,

$$\mathcal{L}^* = \log C - \mathrm{KL}(q(\boldsymbol{\theta})\,\|\,q^*(\boldsymbol{\theta})) \tag{13}$$

where, $q^*(\boldsymbol{\theta}) = M_\theta p(\boldsymbol{\theta})/C$. Crucially, we know the optimal variational distribution over $\boldsymbol{\theta}$ (up to the normalising constant C), so by sampling directly from $q^*(\boldsymbol{\theta})$ using MCMC we eliminate the need to sample the variables $\boldsymbol{u}$. By setting $q(\boldsymbol{\theta}) = q^*(\boldsymbol{\theta})$, we find the *doubly collapsed* ELBO $\mathcal{L}^{**} = \log C$. Although the value of $\mathcal{L}^{**}$ is computationally intractable, given samples $(\boldsymbol{\theta}_j)_{j=1}^J$ from

168    $q^*(\boldsymbol{\theta})$, gradients of $\mathcal{L}^{**}$ with respect to variational parameters ϕ can be estimated: using the chain
169   rule (showing explicit dependence of $\mathcal{L}^*$ and $\mathcal{L}^{**}$ on ϕ),

$$\frac{\mathrm{d}}{\mathrm{d}\phi}\mathcal{L}^{**}_\phi = \left.\frac{\partial}{\partial\phi}\mathcal{L}^*_\phi\right|_{q=q^*} + \left\langle \left.\frac{\delta}{\delta q}\mathcal{L}^*_\phi(q)\right|_{q=q^*}, \frac{\partial}{\partial\phi}q^*_\phi \right\rangle \approx \frac{1}{J}\sum_{j=1}^{J}\frac{\partial}{\partial\phi}\mathcal{L}_{\phi,\boldsymbol{\theta}_j} \qquad (14)$$

170   where $\frac{\delta}{\delta q}\mathcal{L}^*_\phi(q)$ is the functional derivative of $\mathcal{L}^*_\phi$ with respect to q, which is zero at $q=q^*$, because
171   q^* optimises $\mathcal{L}^*_\phi$ (it is a critical point, so the derivative is zero).

172   4.2   Performing approximate inference

173   We deploy HMC to sample from the optimal variational posterior $q^*(\boldsymbol{\theta})$ along with optimising the
174   inducing inputs Z in a hybrid scheme. We alternate between the two steps allocating longer intervals
175   for optimising Z for every HMC sampling run for the hyperparameters. We note that this hybrid
176   scheme is much more computationally efficient than sampling $\boldsymbol{u}$ and $\boldsymbol{\theta}$ jointly where one has to
177   tackle the coupling between inducing variables and hyperparameters in joint space. Further, joint
178   sampling is only feasible for moderate number of inducing variables while this scheme can scale to
179   much larger datasets as the efficiency of sampling in the hyperparameter space is only dependent on
180   the dimensionality of the hyperparameter space rather than the number of inducing variables. The
181   entire inference scheme is summarized in Algorithm 1.

Algorithm 1 Fully Bayesian Sparse GPR with HMC

1:   **Input:** ELBO objective $\mathcal{L}(\boldsymbol{\theta}, Z)$, gradient based optimiser `optim()`
2:   Initialise model params $\boldsymbol{\theta}$ (including data noise variance σ_y^2) and
3:   variational params $Z \in \mathbb{R}^M$ with a warm start strategy.
4:   **procedure** WARM-START
5:      **for** fixed number of steps **do**
6:        Gradient step: $Z, \boldsymbol{\theta} \longleftarrow$ `optim(`$\mathcal{L}(\boldsymbol{\theta}, Z)$`)`
7:   **return** initial values Z_{init} and $\boldsymbol{\theta}_{init}$
8:   **procedure** TRAIN
9:      **while** not converged **do**
10:       ▶ Initialise $\mathcal{L}$ at warm-start values by setting $\hat{\mathcal{L}} = \mathcal{L}(\boldsymbol{\theta}_{init}, Z_{init})$.
11:      ▶ Freeze kernel hyperparameters in the ELBO objective by setting
12: `requires_grad=False`.
13:      **for** t = 1 . . . T **do**
14:        ● Gradient step: $Z_{opt} \longleftarrow$ `optim(`$\hat{\mathcal{L}}$`)`
15:       **if** t mod L == 0 **then**
16:         (For every L gradient steps)
17:         ● Draw J samples from the optimal hyperparameter variational
18:         distribution $\log q^*(\boldsymbol{\theta}) \propto \mathcal{L}(\boldsymbol{\theta}, Z_{opt})p(\boldsymbol{\theta})$.
19:

$$(\boldsymbol{\theta}_j, p_j) \overset{\text{HMC}}{\longleftarrow} \mathcal{H}(\boldsymbol{\theta}, \boldsymbol{p}),$$

20:         where p denotes the zero-mean momentum variable in phase space with
21:         the same dimensionality as $\boldsymbol{\theta}$.
22:         ● Compute stochastic ELBO $\hat{\mathcal{L}} = \frac{1}{J}\sum_{j=1}^{J}\mathcal{L}(\boldsymbol{\theta}_j, Z_{opt})$
23:   **return** $Z_{opt}, \{\boldsymbol{\theta}\}_{j=1}^{J}$

---

182   4.3   Predictive posterior

183   It is ultimately the posterior predictive (PP) distribution that is of interest rather than point estimates
184   or the hyperparameter posterior. The Bayesian sparse GP predictive posterior entails integrating out
185   the posterior over inducing variables $\boldsymbol{u}$ and hyperparameters $\boldsymbol{\theta}$. Once we have performed inference,
186   we can approximate this directly,

$$p(\boldsymbol{f}^*|\boldsymbol{y}) = \int\int p(\boldsymbol{f}^*|\boldsymbol{u}, \boldsymbol{\theta})p(\boldsymbol{u}, \boldsymbol{\theta}|\boldsymbol{y})d\boldsymbol{u}d\boldsymbol{\theta} = \int\int p(\boldsymbol{f}^*|\boldsymbol{u}, \boldsymbol{\theta})p(\boldsymbol{u}|\boldsymbol{\theta}, \boldsymbol{y})p(\boldsymbol{\theta}|\boldsymbol{y})d\boldsymbol{u}d\boldsymbol{\theta} \qquad (15)$$

$$\approx \int\int p(\boldsymbol{f}^*|\boldsymbol{u}, \boldsymbol{\theta})q^*(\boldsymbol{u}|\boldsymbol{\theta}, \boldsymbol{y})p(\boldsymbol{\theta}|\boldsymbol{y})d\boldsymbol{u}d\boldsymbol{\theta} = \int \mathcal{N}(\boldsymbol{f}^*|A\boldsymbol{m}^*, K_{**} + A(S^* - K_{mm})A^T)p(\boldsymbol{\theta}|\boldsymbol{y})d\boldsymbol{\theta}$$

Figure 2: Top - 1d regression with Left: SGPR, Middle: SGPR + HMC, Right: Below - Left: Samples from the mixture posterior, Middle: Length-scale distribution under HMC and ML-II. Right: Noise std. deviation distribution from HMC and ML-II.

where $q^*(\boldsymbol{u}|\boldsymbol{\theta}, \boldsymbol{y}) = \mathcal{N}(\boldsymbol{m}^*, S^*)$ is the optimal Gaussian variational distribution and $A = K_{*m}K_{mm}^{-1}$ under the SGPR scheme (Section 4) is available in closed form with $\boldsymbol{m}^* = \sigma^{-2}(K_{mm} + \sigma^{-2}K_{mn}K_{nm})$ and $S^* = K_{mm}(K_{mm} + \sigma^{-2}K_{mn}K_{nm})^{-1}K_{mm}$. In either case, the internal integral with respect to the inducing variables is analytic the outer integral can be estimated using the samples collected from HMC to perform Monte Carlo estimation. This yields a mixture of Gaussians,

$$p(\boldsymbol{f}^*|\boldsymbol{y}) \approx \frac{1}{J}\sum_{j=1}^{J}\mathcal{N}(\boldsymbol{\mu}^{\theta_j}, \Sigma^{\theta_j}), \qquad \boldsymbol{\theta_j} \sim q^*(\boldsymbol{\theta}), \tag{16}$$

$$\boldsymbol{\mu}^{\theta_j} = A^{(\theta_j)}\boldsymbol{m}^{(\theta_j)}, \qquad \Sigma^{\theta_j} = K_{**}^{\theta_j} + A^{(\theta_j)}(S^{\theta_j} - K_{mm}^{(\theta_j)})A^{T^{(\theta_j)}} \tag{17}$$

where J samples are (approximately) drawn from $q^*(\boldsymbol{\theta})$ via HMC. The distribution inside the summation is the Gaussian posterior predictive distribution for fixed hyperparameters with identical mixing proportions. The compute cost for the predictive posterior scales the sparse GPR cost linearly in the number of samples. The Monte Carlo approximation costs $\mathcal{O}(JNM^2)$ for M inducing points and J hyperparameter samples.

5 Experiments

In the previous section, we discussed a hybrid scheme which leverages MCMC within the variational sparse inducing variable formulation leading to fully Bayesian sparse Gaussian processes. In the experiments we demonstrate the feasibility of this scheme relative to several benchmarks and assess regression performance on a 1-dimensional illustrative example and a range of other datasets. We also compare with exact GPs where the cost of using a gradient based sampler is prohibitive for even moderately large datasets requiring several inversions of the full covariance matrix. We show that using the doubly collapsed scheme proposed in this work is a much more attractive alternative for large datasets as compared to direct HMC in GPR, and sampling is more efficient than in the uncollapsed bound used in Hensman et al. [2015b].

We henceforth refer to benchmark methods as follows: **SGPR + HMC** refers to Bayesian GPs with doubly collapsed variational inference with NUTS, as described in section 3 (compatible with Gaussian likelihoods); we benchmark this model against sparse GPs (**SGPR**) [Titsias, 2009] and Stochastic Variational GPs (**SVGP**) both using approximate ML-II [Hensman et al., 2015a]. Additionally, we consider the sparse, joint sampling inference scheme proposed in [Hensman et al., 2015b] which gives a natural benchmark. We call this model **JointHMC**.

Dataset	N	d	GPR + HMC	SGPR	SGPR + HMC	SVGP	JointHMC
$|M|$	-	-	-	100	100	100	100
$q(\boldsymbol{\theta})$	-	-	free-form	δ	free-form	δ	free-form
Boston	506	13	3.049 (0.14)	3.291 (0.11)	3.286 (0.09)	3.619 (0.11)	3.28 (0.11)
Concrete	1030	8	4.864 (0.12)	5.459 (0.09)	5.402 (0.05)	5.617 (0.09)	5.612 (0.09)
Energy	768	8	0.441 (0.01)	0.477 (0.008)	0.469 (0.009)	0.500 (0.01)	0.755 (0.02)
WineRed	1599	11	0.640 (0.01)	0.636 (0.008)	0.635 (0.008)	0.641 (0.007)	0.641 (0.007)
Yacht	308	6	0.353 (0.03)	0.412 (0.03)	0.387 (0.03)	0.606 (0.04)	0.794 (0.07)

Table 4: A comparison of Sparse GP approaches for UCI benchmarks. RMSE ($\pm$ standard error of mean) evaluated on average of 10 splits with 80% of the data used for training.

We also present analysis where we fix inducing point locations at a random subset of the training data (as opposed to interleaving as per Algorithm 1) and only learn hyperparameters using NUTS. We provide several details about the experimental set-up in the supplementary.

Figure 3: Negative test log likelihoods with standard error of mean across 10 splits with 80% of the data reserved for training. Our method is SGPR + HMC.

5.1 One dimensional synthetic data

We sample noisy observations from $f(x) = \sin(3x) + 0.3\cos(\pi x)$ with the constraint $(x < -2)$ and $(x > 2)$. Figure 2 shows the results for SGPR along with the fully Bayesian schemes. We keep data split and noise identical across the three models to facilitate a comparison. While there is significant data to identify the hyperparameters we notice that the models mainly differ in their extrapolation abilities away from the training data. SGPR with ML-II overfits to the training data and recovers a low noise solution while the SGPR + HMC scheme recovers a more moderate fit and performs significantly better in terms of RMSE and NLPD on unseen data Table 3. We note that the JointHMC scheme which samples both $(\boldsymbol{u}, \boldsymbol{\theta})$ overfits in the central missing data region. We use $M = 25$ inducing locations across all methods which are optimised according to the protocol of each method and recover a very similar spatial distribution.

Table 3: Prediction performance in 1D synthetic regression across SGPR, SGPR + HMC and JointHMC methods with identical number of inducing points and train/test split.

Method	SGPR	SGPR + HMC	JointHMC
RMSE	0.580	**0.537**	0.682
NLPD	0.214	**0.065**	0.74

5.2 UCI regression benchmarks

We compare our approach across methods on 5 standard small to medium-sized UCI benchmark datasets. Following common practice we use a 20% randomly selected held out test set and scale the inputs and outputs to zero mean and unit standard deviation within the training set (we restore the output scaling for evaluation). While we could use any kernel, we choose the RBF-ARD kernel with a lengthscale for each dimension. For consistency we initialise all the inducing locations (Z) identically across the methods, i.e. by using the same random subset of training data split. We note that adapting the inducing locations brings serious gains in prediction performance versus keeping them fixed (Figure 3). Further, the JointHMC scheme underperforms SGPR (with ML-II) and SGPR + HMC. This is not surprising given that the JointHMC bound Equation (9) does not incorporate the optimal setting for $q(\boldsymbol{u})$ and was originally motivated by the need for a fully Bayesian scheme for generalised likelihoods. The method SGPR + HMC significantly improves upon JointHMC in the specific Gaussian likelihood case.

5.3 Ablation study

In order to understand the efficacy of Algorithm 1 we conduct an ablation study where we perform inference in the same manner, but keeping inducing locations fixed. Algorithmically, this implies that we don't need to compute the stochastic ELBO $\hat{\mathcal{L}}$ and just conduct a single sampling run for the hyperparameters. The results across 10 splits are summarised in Table 5.

Sensitivity to M: We benchmark SGPR+HMC and JointHMC on the Elevator dataset ($N = 16599, D = 18$) which demands a larger M. SGPR+HMC outperforms JointHMC for this dataset across different M but the advantage is more pronounced at smaller M. Further, our method took 1248 vs. 2109 wall clock sec. for the joint scheme for the same number of hyper-parameter samples and 500 inducing points.

Dataset	Metric	Boston	Concrete	Energy	WineRed	Yacht
Fixed Z	RMSE	3.624 (0.11)	6.021 (0.12)	0.499 (0.014)	0.640 (0.007)	0.533 (0.036)
Adapt Z		**3.286**(0.09)	**5.405** (0.07)	**0.469** (0.009)	0.635 (0.008)	**0.387** (0.03)
Fixed Z	NLPD	2.640 (0.02)	3.20 (0.06)	0.696 (0.014)	0.969 (0.012)	0.791 (0.13)
Adapt Z		**2.524** (0.022)	**3.065** (0.01)	**0.644** (0.013)	0.961 (0.011)	**0.391** (0.146)

Table 5: An ablation study for the doubly collapsed Sparse GPR scheme comparing performance with and without adapting the inducing locations during training. The table reports test NLPDs and test RMSEs over 10 splits.

5.4 Runtimes

While it is possible to train Exact GPR with HMC for datasets of this size it is important to look at the trade-off in terms of compute cost. In Figure 4 we record the average number of wall clock seconds to draw 500 samples under each method. The cost of sampling is $\mathcal{O}(N)$ for SGPR + HMC but $\mathcal{O}(N^3)$ for Exact GPR + HMC. JointHMC deals with a higher dimensional phase space $(\boldsymbol{u}, \boldsymbol{\theta})$ hence requires more tuning. We don't include tuning time for a fair comparison.

6 Discussion

The evidence framework continues to be the pre-dominant method for training Gaussian processes since their inception into modern machine learning [Rasmussen and Williams, 2006]. While the marginal likelihood is a compelling model selection objective as it offers an inherent trade-off between data-fit and complexity, it is susceptible to overfitting and other pathologies leading to biased inference. This work investigates sparse Gaussian processes relying on inducing variables in conjunction with Bayesian hyperparameter inference.

Bayesian hyperparameter inference in GPs is however intractable and one has to consider balancing between objectives of computational cost, prediction accuracy and robustness of uncertainty intervals. While in straightforward conditions the

Figure 4: Sampling performance measured in terms of the time it took to draw the combined set of samples during the training phase (excluding tuning) as defined by the python standard library *time.perf_counter (wall time)*. We use the `pymc3 pm.NUTS` sampler for GPR + HMC and SGPR + HMC, and `tfp.mcmc.HamiltonianMonteCarlo` for JointHMC. All experiments were conducted on an Intel Core i7-10700 CPU @ 2.90GHz x 16.

fully Bayesian approach might be counter-productive, most real-world applications of GPs rely on engineering sophisticated hand-crafted kernels involving many hyperparameters where there risk of overfitting is pronounced and further, harder to detect. A more robust solution is to incorporate prediction intervals that reflect these uncertainties in the model choice. Studying full Bayesian inference in more sophisticated GP models like deep [Damianou and Lawrence, 2013], warped [Snelson et al., 2004] and convolutional GPs [Van der Wilk et al., 2017] will offer greater insight to this question and is an imminent direction of future work.

References

David Barber and Christopher KI Williams. Gaussian processes for Bayesian classification via hybrid Monte Carlo. In *Advances in neural information processing systems*, pages 340–346, 1997.

Michael Betancourt. A conceptual introduction to Hamiltonian Monte Carlo. *arXiv preprint arXiv:1701.02434*, 2017.

Michael Betancourt and Mark Girolami. Hamiltonian monte carlo for hierarchical models. *Current trends in Bayesian methodology with applications*, 79(30):2–4, 2015.

Thang D Bui, Cuong V Nguyen, Siddharth Swaroop, and Richard E Turner. Partitioned variational inference: A unified framework encompassing federated and continual learning. *arXiv preprint arXiv:1811.11206*, 2018.

David R. Burt, Carl Edward Rasmussen, and Mark van der Wilk. Convergence of sparse variational inference in Gaussian processes regression. *Journal of Machine Learning Research*, 21(131):1–63, 2020.

Andreas Damianou and Neil D Lawrence. Deep Gaussian processes. In *Artificial intelligence and statistics*, pages 207–215. PMLR, 2013.

Maurizio Filippone and Mark Girolami. Pseudo-marginal Bayesian inference for Gaussian processes. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 36(11):2214–2226, 2014.

Maurizio Filippone, Mingjun Zhong, and Mark Girolami. A comparative evaluation of stochastic-based inference methods for gaussian process models. *Machine Learning*, 93(1):93–114, 2013.

James Hensman, Nicolo Fusi, and Neil D Lawrence. Gaussian processes for big data. *arXiv preprint arXiv:1309.6835*, 2013.

James Hensman, Alexander Matthews, and Zoubin Ghahramani. Scalable variational Gaussian process classification. *18th International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2015a.

James Hensman, Alexander G Matthews, Maurizio Filippone, and Zoubin Ghahramani. Mcmc for variationally sparse gaussian processes. In *Advances in Neural Information Processing Systems*, pages 1648–1656, 2015b.

Vidhi Lalchand and Carl Edward Rasmussen. Approximate inference for fully bayesian Gaussian process regression. In *Symposium on Advances in Approximate Bayesian Inference*, pages 1–12. PMLR, 2020.

David JC MacKay. Hyperparameters: optimise or integrate out. *Maximum entropy and Bayesian methods*, 1994.

A. G. d. G. Matthews, J. Hensman, R. E. Turner, and Z. Ghahramani. On sparse variational methods and the Kullback-Leibler divergence between stochastic processes. In *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics*, volume 54 of *Proceedings of Machine Learning Research*. PMLR, 2016.

Iain Murray and Ryan P Adams. Slice sampling covariance hyperparameters of latent gaussian models. In *Advances in neural information processing systems*, pages 1732–1740, 2010.

Radford Neal. Regression and classification using Gaussian process priors. *Bayesian statistics*, 6:475, 1998.

Radford M Neal et al. MCMC using Hamiltonian dynamics. *Handbook of markov chain monte carlo*, 2(11):2, 2011.

Sebastian W Ober, Carl E Rasmussen, and Mark van der Wilk. The promises and pitfalls of deep kernel learning. *arXiv preprint arXiv:2102.12108*, 2021.

Carl Edward Rasmussen and Zoubin Ghahramani. Occam's razor. *Advances in neural information processing systems*, pages 294–300, 2001.

Carl Edward Rasmussen and Christopher KI Williams. Gaussian processes for machine learning (adaptive computation and machine learning), 2006.

Simone Rossi, Markus Heinonen, Edwin Bonilla, Zheyang Shen, and Maurizio Filippone. Sparse Gaussian processes revisited: Bayesian approaches to inducing-variable approximations. In *International Conference on Artificial Intelligence and Statistics*, pages 1837–1845. PMLR, 2021.

Fergus Simpson, Vidhi Lalchand, and Carl Edward Rasmussen. Marginalised Gaussian processes with nested sampling. *Advances in neural information processing systems*, 2021.

Edward Snelson, Carl Edward Rasmussen, and Zoubin Ghahramani. Warped gaussian processes. *Advances in neural information processing systems*, 16:337–344, 2004.

Michael Stein. *Interpolation of Spatial Data*. Springer-Verlag New York, 1 edition, 1999.

Michalis Titsias. Variational learning of inducing variables in sparse Gaussian processes. In *Artificial intelligence and statistics*, pages 567–574. PMLR, 2009.

Mark Van der Wilk, Carl Edward Rasmussen, and James Hensman. Convolutional Gaussian processes. *arXiv preprint arXiv:1709.01894*, 2017.

JJ Warnes and BD Ripley. Problems with likelihood estimation of covariance functions of spatial Gaussian processes. *Biometrika*, 74(3):640–642, 1987.

Christopher KI Williams and Carl Edward Rasmussen. Gaussian processes for regression. In *Advances in neural information processing systems*, pages 514–520, 1996.

Andrew Gordon Wilson, Zhiting Hu, Ruslan Salakhutdinov, and Eric P Xing. Deep kernel learning. In *Artificial intelligence and statistics*, pages 370–378. PMLR, 2016.