History, Hypergraphs, and Memory: The Exact Complexity of Deviation-Rational Control

Keywords: reinforcement learning, big worlds, bounded-memory agents, history processes, deviation regret, hypergraph transversals, convex geometry, combinatorial optimization.

Summary

We study the minimum internal memory required for ε-deviation-rational control in arbitrary history processes. Our starting point is a past-compression view of bounded agency: after each attainable history, the agent compresses the entire past into one of finitely many memory labels, and that label must decode to a continuation controller whose deviation regret is at most ε. This formulation is entirely agent-side. It does not assume that the environment is Markov, fully observable, deterministic, finitely enumerable, or representable by a small latent model.

Our main result is an exact structural law: retentive complexity equals the transversal number of the family of history-indexed low-regret continuation sets. In the finite-controller mixed-continuation model, this yields a precise hypergraph law at exact rationality ($\varepsilon = 0$): the required memory is exactly the transversal number of the family of rational-support sets. We further show that every finite hypergraph can be realized as such a support family in a one-step fully observable instance, establishing combinatorial universality. Finally, we prove that pairwise compatibility is insufficient in general—even the complete compatibility graph can conceal large memory gaps—while convexity yields a Helly-type local certificate for one-state memory. Together, these results recast bounded-agent memory in big worlds as a problem in combinatorial optimization and convex geometry rather than environment-state abstraction.

Contribution(s)

1. We prove an exact structural law: for every deviation-stable continuation system and every $\varepsilon \geq 0$, retentive complexity equals the transversal number of the family of history-indexed ε-rational continuation sets.
   Context: This is an exact characterization of memory complexity, not an algorithmic tractability result.

2. In the finite-controller mixed-continuation model, we show that low-regret sets are convex polytopes and that, at exact rationality, retentive complexity is exactly the hypergraph transversal number of the family of rational-support sets; moreover, every finite hypergraph is realizable as such a support family in a one-step fully observable instance.
   Context: The hypergraph law and universality statement are proved for finite controller dictionaries with continuation values affine under mixing.

3. We prove a higher-order impossibility result: pairwise compatibility does not determine exact retentive complexity, and for every order q admitting a finite projective plane there exist two instances with identical complete compatibility graph K_{q^2+q+1} but exact retentive complexities 1 and $q + 1$.
   Context: This is a negative structural result; the projective-plane construction provides an explicit obstruction, not a classification of all obstructions.

4. We identify the convex regime in which local structure suffices: one-state memory admits a Helly-type certificate, and in the finite-controller simplex model any deviation class containing the external deviations yields exactly the same low-regret sets and retentive complexities as the external-deviation model for all $\varepsilon \geq 0$.
   Context: The Helly and robustness statements are finite-controller simplex results; they are not claimed for arbitrary continuation spaces.

History, Hypergraphs, and Memory: The Exact Complexity of Deviation-Rational Control

Anonymous authors

Abstract

We study the minimum internal memory required for ε-deviation-rational control in arbitrary history processes. We formalize this as the minimum number of labels in a past-compression architecture that maps each attainable history to a finite memory state and decodes that state to a continuation controller. Our first result is an exact law: this retentive complexity equals the transversal number of the family of history-indexed ε-rational continuation sets. We then specialize to finite controller dictionaries with mixed continuations on a simplex and show that the low-regret sets are convex polytopes; at exact rationality ($\varepsilon = 0$), the problem collapses to a hypergraph law, in which the required memory is exactly the transversal number of the family of rational-support sets. We further prove universality by showing that every finite hypergraph arises as such a support family in a one-step fully observable instance. Finally, we prove that pairwise compatibility is insufficient in general—even the complete compatibility graph can conceal large memory gaps—while convexity yields a Helly-type local certificate for one-state memory. Together, these results recast bounded-agent memory in big worlds as a problem in combinatorial optimization and convex geometry rather than environment-state abstraction.

1   Introduction

Reinforcement-learning agents act in histories, not in states supplied by the analyst. In large or partially observed worlds, the agent must compress the past into a bounded internal memory and act from that compressed state thereafter. This raises a basic question: *how much memory is required to summarize the past while preserving rational future behavior?* We study that question in an arbitrary history-based interface, under a deviation-regret criterion, and ask for the minimum memory cardinality sufficient to remain ε-deviation-rational from every attainable history.

This question is naturally history-first. Elelimy et al. (2025) advocate a continual-RL viewpoint based on histories, deviations, and behavioral consequences rather than on optimality in a fixed MDP, while hindsight-rationality provides rich deviation classes for sequential decision problems (Morrill et al., 2021). Related recent work studies what capable agents must internally represent, including predictive/world-model necessity results (Richens et al., 2025; Nayebi, 2026), bounded-interaction quotients (Nixon, 2026), and decision-centric memory compression (Zou et al., 2026). Those results are closely related in spirit, but they do not provide an exact cardinal law for the minimum memory required for ε-deviation-rational control in arbitrary history processes.

Our formulation separates the *past* from the *future*. For each attainable history h, let K_h^ε be the set of continuation controllers that are ε-deviation-rational from h. Memory then becomes a compression problem: histories are mapped to finitely many labels, and each label must decode to a continuation controller lying in the appropriate low-regret set for every history assigned to that label. This formulation is entirely agent-side: the relevant objects are the continuation-controller class, the de-

37 viation class, and the number of internal memory labels, not any latent state-space or observability
38 assumption.

39 Our contribution is an exact theoremic characterization of this memory problem. First, retentive
40 complexity equals the transversal number of the family of history-indexed low-regret continuation
41 sets. Second, in the finite-controller mixed-continuation model, exact rationality ($\varepsilon = 0$) collapses to
42 a hypergraph law: the required memory is exactly the transversal number of the family of rational-
43 support sets. Third, every finite hypergraph arises as such a support family in a one-step, fully
44 observable instance. Fourth, pairwise compatibility is insufficient in general: even the complete
45 compatibility graph can conceal large memory gaps. Finally, convexity yields a Helly-type local
46 certificate for one-state memory.

47 These results yield a different conceptual picture of bounded-agent memory than the one suggested
48 by quotient or metric viewpoints. The governing object is the higher-order intersection geometry
49 of the family of low-regret continuation sets. In the finite-controller model this geometry becomes
50 a hypergraph transversal problem; under convexity it admits local-to-global certificates; and, in
51 general, it is strictly richer than anything visible from pairwise compatibility alone. Finally, the
52 theory interfaces directly with ordinary finite-state agents: any deterministic recurrent agent whose
53 current internal state determines a continuation controller must have at least as many reachable
54 internal states as the retentive complexity defined here.

2   Model and Retentive Complexity

56 We work directly with histories and deviations, following the recent history-process and deviation-
57 regret perspective in continual reinforcement learning and hindsight-rationality (Elelimy et al., 2025;
58 Morrill et al., 2021). The formulation below is intentionally agent-side: the environment enters only
59 through the attainable-history set and the continuation values induced after each history, while the
60 complexity notion itself concerns only the family of continuation controllers available to the agent
61 and the deviations against which those controllers are evaluated.

62 **Definition 2.1** (Deviation-stable continuation system). A *deviation-stable continuation system* is a
63 quadruple

$$\mathfrak{M} = (\mathcal{H}, \mathcal{C}, \Phi, J),$$

64 where $\mathcal{H} \neq \varnothing$ is a set of attainable histories, $\mathcal{C} \neq \varnothing$ is a class of admissible continuation controllers,
65 $\Phi \neq \varnothing$ is a family of self-maps $\phi : \mathcal{C} \to \mathcal{C}$, and $J : \mathcal{H} \times \mathcal{C} \to \mathbb{R}$ is a bounded continuation-value
66 functional. We write ϕc for $\phi(c)$. The deviation regret of $c \in \mathcal{C}$ after $h \in \mathcal{H}$ is

$$\text{Reg}_\Phi(h, c) := \sup_{\phi \in \Phi} \Big(J(h, \phi c) - J(h, c) \Big). \tag{2.1}$$

67 **Definition 2.2** (ε-rational continuation set). For $\varepsilon \geq 0$ and $h \in \mathcal{H}$, the ε-rational continuation set is

$$K_h^\varepsilon := \{c \in \mathcal{C} : \text{Reg}_\Phi(h, c) \leq \varepsilon\}. \tag{2.2}$$

68 An element $c \in \mathcal{C}$ should be read as the entire closed-loop behavior the agent can execute *from*
69 *the current moment onward* once its internal memory label has been fixed. Thus K_h^ε is the set of
70 continuations that remain ε-deviation-rational from history h.

71 **Definition 2.3** (Past-compression architecture). For $m \in \mathbb{N}$, write $[m] := \{1, \ldots, m\}$. A *past-*
72 *compression architecture of size m* is a pair

$$(\sigma, d), \qquad \sigma : \mathcal{H} \to [m], \quad d : [m] \to \mathcal{C}.$$

73 It is *ε-rational* if

$$d(\sigma(h)) \in K_h^\varepsilon \qquad \forall h \in \mathcal{H}. \tag{2.3}$$

74 **Definition 2.4** (Retentive complexity)**.** The *retentive complexity* of $\mathfrak{M}$ at tolerance ε is

$$m_\Phi^{\mathrm{pc}}(\varepsilon) := \inf\{m \in \mathbb{N} : \exists\ \varepsilon\text{-rational past-compression architecture of size } m\}. \qquad (2.4)$$

75 If no finite ε-rational architecture exists, we interpret $m_\Phi^{\mathrm{pc}}(\varepsilon) = \infty$.

76 The quantity $m_\Phi^{\mathrm{pc}}(\varepsilon)$ is the minimum number of memory labels needed to compress the *entire past*
77 while still ensuring that, from every attainable history, at least one continuation chosen solely from
78 the compressed label is ε-deviation-rational.

79 **Proposition 2.5** (Finite-state compilation lower bound)**.** *Let Z be a finite internal state set, let*
80 $\zeta : \mathcal{H} \to Z$ *be a deterministic history-to-state map, and let $d : Z \to \mathcal{C}$ assign a continuation*
81 *controller to each internal state. If*

$$d(\zeta(h)) \in K_h^\varepsilon \qquad \forall h \in \mathcal{H}, \qquad (2.5)$$

82 *then*
$$|Z| \geq m_\Phi^{\mathrm{pc}}(\varepsilon).$$

83 *Proof.* Let $Z_{\mathrm{reach}} := \zeta(\mathcal{H}) \subseteq Z$, and choose a bijection $b : Z_{\mathrm{reach}} \to [|Z_{\mathrm{reach}}|]$. Define

$$\sigma := b \circ \zeta, \qquad \tilde{d}(i) := d\big(b^{-1}(i)\big) \quad \text{for } i \in [|Z_{\mathrm{reach}}|].$$

84 Then, for every $h \in \mathcal{H}$,
$$\tilde{d}(\sigma(h)) = d(\zeta(h)) \in K_h^\varepsilon$$

85 by (2.5). Hence $(\sigma, \tilde{d})$ is an ε-rational past-compression architecture of size $|Z_{\mathrm{reach}}|$. By Defini-
86 tion 2.4,
$$m_\Phi^{\mathrm{pc}}(\varepsilon) \leq |Z_{\mathrm{reach}}| \leq |Z|,$$

87 which proves the claim. $\qquad\qquad\square$

88 3 Exact Reduction to a Transversal Problem

89 The central structural fact of the paper is that retentive complexity is exactly a hitting-set problem
90 over the family of low-regret continuation sets. This reduction is purely set-theoretic: it uses only
91 the definitions of K_h^ε and $m_\Phi^{\mathrm{pc}}(\varepsilon)$, and therefore does not rely on any simplifying assumption about
92 the environment or the deviation class.

93 For $\varepsilon \geq 0$, write
$$\mathcal{K}_\varepsilon := \{K_h^\varepsilon : h \in \mathcal{H}\}.$$

94 A set $P \subseteq \mathcal{C}$ is a *transversal* for $\mathcal{K}_\varepsilon$ if

$$P \cap K_h^\varepsilon \neq \varnothing \qquad \forall h \in \mathcal{H}.$$

95 Its minimum possible cardinality is the transversal number

$$\tau(\mathcal{K}_\varepsilon) := \inf\{m \in \mathbb{N} : \exists P \subseteq \mathcal{C},\ |P| \leq m,\ P \cap K_h^\varepsilon \neq \varnothing\ \forall h \in \mathcal{H}\}, \qquad (3.1)$$

96 with the convention $\inf \varnothing = \infty$.

97 **Theorem 3.1** (Exact transversal law)**.** *For every deviation-stable continuation system and every*
98 $\varepsilon \geq 0$,
$$m_\Phi^{\mathrm{pc}}(\varepsilon) = \tau(\mathcal{K}_\varepsilon). \qquad (3.2)$$

99 *Proof.* We show that, for each memory budget $m \in \mathbb{N}$, there exists an ε-rational past-compression
100 architecture of size m if and only if there exists a transversal of $\mathcal{K}_\varepsilon$ of cardinality at most m.

101 First, suppose (σ, d) is an ε-rational past-compression architecture of size m, so

$$\sigma : \mathcal{H} \to [m], \qquad d : [m] \to \mathcal{C},$$

102 and

$$d(\sigma(h)) \in K_h^\varepsilon \qquad \forall h \in \mathcal{H}.$$

103 Let

$$P := d([m]) \subseteq \mathcal{C}.$$

104 Then $|P| \le m$. Moreover, for every $h \in \mathcal{H}$,

$$d(\sigma(h)) \in P \cap K_h^\varepsilon,$$

105 so $P \cap K_h^\varepsilon \neq \varnothing$. Hence P is a transversal for $\mathcal{K}_\varepsilon$ of size at most m.

106 Conversely, suppose $P \subseteq \mathcal{C}$ is a transversal for $\mathcal{K}_\varepsilon$ with $|P| \le m$. Enumerate its distinct elements
107 as

$$P = \{c_1, \ldots, c_r\}, \qquad r \le m.$$

108 Define $d : [m] \to \mathcal{C}$ by $d(i) = c_i$ for $1 \le i \le r$, and pad the remaining labels arbitrarily, say
109 $d(i) = c_1$ for $r < i \le m$. Since P intersects every K_h^ε, the set

$$I(h) := \{i \in [r] : c_i \in K_h^\varepsilon\}$$

110 is nonempty for every $h \in \mathcal{H}$. Define

$$\sigma(h) := \min I(h).$$

111 Then $\sigma : \mathcal{H} \to [m]$ is well defined, and by construction

$$d(\sigma(h)) = c_{\sigma(h)} \in K_h^\varepsilon \qquad \forall h \in \mathcal{H}.$$

112 Thus (σ, d) is an ε-rational past-compression architecture of size m.

113 We have proved the equivalence

$$\exists \, \varepsilon\text{-rational architecture of size } m \iff \exists \, \text{transversal of } \mathcal{K}_\varepsilon \text{ of size } \le m.$$

114 Taking the infimum over $m \in \mathbb{N}$ on both sides yields (3.2). $\qquad \square$

115 Theorem 3.1 identifies the exact combinatorial object governing memory: not a quotient of histories,
116 not a pairwise metric, and not a descriptive compression criterion, but the piercing number of the
117 family of history-indexed low-regret continuation sets.

4 Finite Controller Dictionaries: Hypergraph Law and Universality

119 To expose the combinatorial content of Theorem 3.1, we specialize to a finite controller dictionary
120 $\{e_1, \ldots, e_k\}$ and its simplex of mixed continuations

$$\Delta_k := \left\{ x \in \mathbb{R}_+^k : \sum_{i=1}^k x_i = 1 \right\}.$$

121 We assume that continuation values are affine under mixing, so that for every history $h \in \mathcal{H}$ and
122 $x \in \Delta_k$,

$$J(h, x) = \sum_{i=1}^k x_i J(h, e_i).$$

123   Let

$$v(h) := (v_1(h), \dots, v_k(h)) \in \mathbb{R}^k, \qquad v_i(h) := J(h, e_i),$$

124   and take the deviation class to be the external deviations

$$\Phi_{\text{ext}} := \{\phi_i : i \in [k]\}, \qquad \phi_i(x) := e_i.$$

125   Thus, after any history, a deviation may replace the mixed continuation by any pure library con-
126   troller.

127   **Theorem 4.1** (Polyhedral low-regret sets). *For every history $h \in \mathcal{H}$ and every $x \in \Delta_k$,*

$$\text{Reg}_{\Phi_{\text{ext}}}(h, x) = \max_{i \in [k]} v_i(h) - \langle v(h), x \rangle. \tag{4.1}$$

128   *Consequently, for every $\varepsilon \geq 0$,*

$$K_h^\varepsilon = \left\{ x \in \Delta_k : \langle v(h), x \rangle \geq \max_{i \in [k]} v_i(h) - \varepsilon \right\}. \tag{4.2}$$

129   *In particular, K_h^ε is the intersection of the simplex with a closed half-space, and hence a convex*
130   *polytope.*

131   *Proof.* By affineness of J under mixing,

$$J(h, x) = \sum_{i=1}^k x_i J(h, e_i) = \langle v(h), x \rangle.$$

132   Moreover, for each external deviation ϕ_i,

$$J(h, \phi_i x) = J(h, e_i) = v_i(h).$$

133   Substituting into Definition 2.1 gives

$$\text{Reg}_{\Phi_{\text{ext}}}(h, x) = \sup_{i \in [k]} (v_i(h) - \langle v(h), x \rangle) = \max_{i \in [k]} v_i(h) - \langle v(h), x \rangle,$$

134   which proves (4.1). The description (4.2) then follows by Definition 2.2.   □

135   At exact rationality, the geometry collapses to simplex faces. Define the maximizer set

$$S_h := \arg\max_{i \in [k]} v_i(h) \subseteq [k],$$

136   and for any nonempty $S \subseteq [k]$, write

$$\Delta(S) := \{x \in \Delta_k : \text{supp}(x) \subseteq S\}.$$

137   **Corollary 4.2** (Exact hypergraph law). *For every $h \in \mathcal{H}$,*

$$K_h^0 = \Delta(S_h).$$

138   *Hence, if*

$$\mathcal{S} := \{S_h : h \in \mathcal{H}\},$$

139   *then*

$$m_{\Phi_{\text{ext}}}^{\text{pc}}(0) = \tau(\mathcal{S}), \tag{4.3}$$

140   *where*

$$\tau(\mathcal{S}) := \inf\{|T| : T \subseteq [k], \ T \cap S \neq \varnothing \ \forall S \in \mathcal{S}\}.$$

Proof. By Theorem 4.1,

$$K_h^0 = \left\{ x \in \Delta_k : \langle v(h), x \rangle = \max_{i \in [k]} v_i(h) \right\}.$$

A convex combination attains the maximum of the coordinates of $v(h)$ iff it places mass only on maximizing indices. Therefore $K_h^0 = \Delta(S_h)$.

Now apply Theorem 3.1 to the family $\{\Delta(S_h) : h \in \mathcal{H}\}$. If $T \subseteq [k]$ hits every support set in $\mathcal{S}$, then the vertex set $\{e_i : i \in T\}$ pierces every face $\Delta(S_h)$. Conversely, if $P \subseteq \Delta_k$ pierces every $\Delta(S_h)$, choose for each $x \in P$ an arbitrary index $i(x) \in \mathrm{supp}(x)$, and set

$$T := \{i(x) : x \in P\} \subseteq [k].$$

Whenever $x \in P \cap \Delta(S_h)$, one has $\mathrm{supp}(x) \subseteq S_h$, hence $i(x) \in S_h$; thus T hits every set in $\mathcal{S}$. The minimum piercing number of the faces $\Delta(S_h)$ is therefore exactly $\tau(\mathcal{S})$, proving (4.3). □

Corollary 4.2 identifies exact retentive complexity with a classical hypergraph transversal number. The next theorem shows that this is not merely an analogy: every finite hypergraph can arise exactly from a history-dependent decision problem.

Theorem 4.3 (Universality)**.** *For every finite hypergraph*

$$\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\},$$

there exists a one-step, fully observable history process with controller dictionary $\{e_1, \ldots, e_k\}$ such that

$$\{S_h : h \in \mathcal{H}\} = \mathcal{S}.$$

Equivalently, every finite hypergraph can be realized as the family of exact rational-support sets of some mixed-controller instance.

Proof sketch. For each hyperedge $S \in \mathcal{S}$, create a distinct observable history h_S. After observing h_S, the agent chooses an action $A \in [k]$ and receives reward

$$R = \mathbf{1}\{A \in S\},$$

after which the episode terminates. Then

$$J(h_S, e_i) = \mathbf{1}\{i \in S\},$$

so the maximizing indices are exactly those in S, i.e.,

$$S_{h_S} = \arg\max_{i \in [k]} J(h_S, e_i) = S.$$

Hence the family $\{S_h : h \in \mathcal{H}\}$ is exactly $\mathcal{S}$. The full proof in Supplement E gives the stronger statement that, for every $\varepsilon \geq 0$,

$$K_{h_S}^\varepsilon = \left\{ x \in \Delta_k : \sum_{i \in S} x_i \geq 1 - \varepsilon \right\}.$$

□

Combining Theorem 4.3 with Corollary 4.2, every finite hitting-set instance already appears as an exact retentive-complexity problem in a one-step history process. Thus the memory geometry of exact deviation-rational control is combinatorially universal even in the simplest fully observable setting.

5 Pairwise Compatibility is Not Enough

The exact hypergraph law identifies retentive complexity with a transversal number of the family $\mathcal{S} = \{S_h : h \in \mathcal{H}\}$. A natural question is whether this family can be reduced to pairwise intersection data alone.

For the mixed-controller model at exact rationality, define the *pairwise compatibility graph*

$$G_{\text{pair}} = (\mathcal{H}, E_{\text{pair}})$$

by declaring distinct histories $h, h' \in \mathcal{H}$ adjacent iff

$$K_h^0 \cap K_{h'}^0 \neq \varnothing, \tag{5.1}$$

which, by Corollary 4.2, is equivalent to $S_h \cap S_{h'} \neq \varnothing$. Thus G_{pair} retains only pairwise intersections and forgets all higher-order intersection structure.

Theorem 5.1 (Projective-plane separation). *Let $\Pi_q = (P_q, \mathcal{L}_q, \in)$ be a finite projective plane of order q, and set $n := q^2 + q + 1$. Then there exist two one-step, fully observable mixed-controller instances, each with controller dictionary $\{e_1, \dots, e_n\}$ and external deviation class Φ_{ext}, such that both have pairwise compatibility graph K_n, but their exact retentive complexities are 1 and $q + 1$, respectively.*

The only numerical input is the following classical counting fact.

Lemma 5.2. *If $\mathscr{L}_q = \{L \subseteq P_q : L \in \mathcal{L}_q\}$ is the line hypergraph of a projective plane of order q, then*

$$\tau(\mathscr{L}_q) = q + 1.$$

Proof sketch of Theorem 5.1. Construct two instances on the same controller simplex Δ_n. In Instance A, reward is always $\mathbf{1}\{A = 1\}$, so $K_{h_j}^0 = \{e_1\}$ for all histories and the pairwise graph is K_n; hence $m_{\Phi_{\text{ext}}}^{\text{pc}}(0) = 1$. In Instance B, histories are indexed by lines $L \in \mathcal{L}_q$, reward at history h_L is $\mathbf{1}\{A \in L\}$, and Theorem 4.3 yields $K_{h_L}^0 = \Delta(L)$. Distinct lines intersect, so the pairwise graph is again K_n. But Corollary 4.2 gives $m_{\Phi_{\text{ext}}}^{\text{pc}}(0) = \tau(\mathscr{L}_q) = q + 1$. $\square$

Proof sketch of Lemma 5.2. Any line $L^\star \in \mathcal{L}_q$ contains $q + 1$ points and intersects every other line, so $\tau(\mathscr{L}_q) \leq q + 1$. Conversely, any set of at most q points lies on at most $q(q + 1) = q^2 + q$ lines, while the plane contains $q^2 + q + 1$ lines in total. Hence some line is missed, so no q-point set can be a transversal. $\square$

Corollary 5.3 (No pairwise characterization). *No function of the pairwise compatibility graph alone can determine $m_{\Phi_{\text{ext}}}^{\text{pc}}(0)$ for all finite-controller mixed-continuation instances.*

Proof. Theorem 5.1 gives two instances with the same graph $G_{\text{pair}} = K_n$ but different exact retentive complexities. $\square$

In particular, exact deviation-rational memory cannot in general be reduced to any pairwise graph-theoretic abstraction.

Thus even the complete compatibility graph can be maximally uninformative: what governs memory is the higher-order intersection geometry of the family $\{K_h^0 : h \in \mathcal{H}\}$, or equivalently of the support hypergraph $\mathcal{S}$. This is precisely the structural gap between pairwise compatibility and exact memory.

6 Local Convex Certificate and Discussion

The preceding sections show that exact retentive complexity is governed by a higher-order set system and, in general, cannot be read from pairwise compatibility alone. In the mixed-controller model, however, the geometry of the low-regret sets yields a sharp local certificate for the special case of one-state memory.

Corollary 6.1 (Helly certificate for one-state memory). *Consider the finite-controller mixed-continuation model of Section 4, with controller simplex Δ_k and external deviations Φ_{ext}. For every $\varepsilon \geq 0$, the following are equivalent:*

1. $m^{\text{pc}}_{\Phi_{\text{ext}}}(\varepsilon) = 1$;

2. $\bigcap_{h \in \mathcal{H}} K^{\varepsilon}_h \neq \varnothing$;

3. *every subfamily of $\{K^{\varepsilon}_h : h \in \mathcal{H}\}$ of cardinality at most k has nonempty intersection.*

Proof. The equivalence of (1) and (2) is immediate from Theorem 3.1: a one-state architecture exists if and only if the family $\{K^{\varepsilon}_h : h \in \mathcal{H}\}$ admits a transversal of size 1, and this is equivalent to the existence of a single point $x \in \Delta_k$ belonging to every K^{ε}_h.

The implication (2) $\Rightarrow$ (3) is trivial. It remains to prove (3) $\Rightarrow$ (2). By Theorem 4.1, each K^{ε}_h is a convex polytope in Δ_k. Since Δ_k lies in the affine hyperplane

$$\text{aff}(\Delta_k) = \left\{ x \in \mathbb{R}^k : \sum_{i=1}^{k} x_i = 1 \right\},$$

the relevant affine dimension is $k - 1$. Hence, by Helly's theorem in affine dimension $k - 1$ (Matoušek, 2002), every finite subfamily $\mathcal{F} \subseteq \{K^{\varepsilon}_h : h \in \mathcal{H}\}$ has nonempty intersection whenever every k or fewer members of $\mathcal{F}$ intersect. Assumption (3) therefore implies that *every finite* subfamily has nonempty intersection.

Finally, each K^{ε}_h is closed and contained in the compact set Δ_k. Since the family has the finite intersection property, compactness of Δ_k implies

$$\bigcap_{h \in \mathcal{H}} K^{\varepsilon}_h \neq \varnothing.$$

Thus (2) holds. $\qquad\square$

Corollary 6.1 reveals a striking contrast with Section 5. For general memory budgets, exact retentive complexity is governed by the full higher-order intersection geometry of the family $\{K^{\varepsilon}_h : h \in \mathcal{H}\}$, and pairwise data can be maximally uninformative. For the special case of one-state memory, by contrast, convexity collapses the problem to a local certificate of order k, where k is the size of the controller dictionary rather than any measure of environmental complexity. This threshold is exactly the Helly number of the affine hull of Δ_k, whose dimension is $k - 1$. In this sense, one-state certification depends only on controller complexity, whereas general retentive complexity remains a genuinely higher-order set-system invariant.

Taken together, Theorems 3.1 and 5.1, Corollary 4.2, and Corollary 6.1 yield a unified picture: deviation-rational memory is neither a quotient of histories nor a pairwise metric invariant. Its exact complexity is a piercing problem over history-indexed low-regret continuation sets; in the finite-controller model this becomes a hypergraph transversal law, while convexity supplies additional local-to-global structure only in special regimes. This places bounded-agent memory in arbitrary worlds squarely at the interface of reinforcement learning, combinatorial optimization, and convex geometry.

References

Charalambos D. Aliprantis and Kim C. Border. *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer, Berlin, Heidelberg, 3 edition, 2006.

Charles Castaing and Michel Valadier. *Convex Analysis and Measurable Multifunctions*, volume 580 of *Lecture Notes in Mathematics*. Springer, Berlin, Heidelberg, 1977.

Esraa Elelimy, David Szepesvari, Martha White, and Michael Bowling. Rethinking the foundations for continual reinforcement learning. *Reinforcement Learning Journal*, 6:2174–2194, 2025.

Jiří Matoušek. *Lectures on Discrete Geometry*, volume 212 of *Graduate Texts in Mathematics*. Springer, 2002.

Dustin Morrill, Ryan D'Orazio, Marc Lanctot, James R Wright, Michael Bowling, and Amy R Greenwald. Efficient deviation types and learning for hindsight rationality in extensive-form games. In Marina Meila and Tong Zhang (eds.), *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pp. 7818–7828. PMLR, 2021.

Aran Nayebi. What capable agents must know: Selection theorems for robust decision-making under uncertainty. *arXiv preprint arXiv:2603.02491*, 2026. URL https://arxiv.org/abs/2603.02491.

Anthony T. Nixon. The myhill–nerode theorem for bounded interaction: Canonical abstractions via agent-bounded indistinguishability. *arXiv preprint arXiv:2603.21399*, 2026. URL https://arxiv.org/abs/2603.21399.

Jonathan Richens, Tom Everitt, and David Abel. General agents need world models. In Aarti Singh, Maryam Fazel, Daniel Hsu, Simon Lacoste-Julien, Felix Berkenkamp, Tegan Maharaj, Kiri Wagstaff, and Jerry Zhu (eds.), *Proceedings of the 42nd International Conference on Machine Learning*, volume 267 of *Proceedings of Machine Learning Research*, pp. 51659–51687. PMLR, 2025. URL https://proceedings.mlr.press/v267/richens25a.html.

Mingxi Zou, Zhihan Guo, Langzhang Liang, Zhuo Wang, Qifan Wang, Qingsong Wen, Irwin King, Lizhen Qu, and Zenglin Xu. Remember the decision, not the description: A rate-distortion framework for agent memory. *arXiv preprint arXiv:2605.10870*, 2026. URL https://arxiv.org/abs/2605.10870.

Supplementary Materials

The following content was not necessarily subject to peer review.

A Formal Interface and Measurable Refinements

This section serves two purposes. First, it records a concrete interface from a standard history-based reinforcement-learning model to the abstract continuation-system formalism used in the main text. Second, it gives a measurable refinement of Theorem 3.1, showing that under standard regularity conditions the finite-memory compression map can be chosen measurably without changing the exact memory law.

A.1 From measurable history processes to continuation systems

We formalize a general history-based decision problem in the spirit of the history-process viewpoint used in recent continual RL work (Elelimy et al., 2025).

Definition A.1 (Discounted measurable history process)**.** A *discounted measurable history process* consists of the following data:

1. a standard Borel action space $\mathcal{A}$;

2. a standard Borel signal space X, whose elements may encode observations, rewards, terminal indicators, or any other percept available to the agent;

3. the set of pre-action histories
$$\mathcal{H} := \bigsqcup_{t \geq 0} (\mathsf{X} \times \mathcal{A})^t \times \mathsf{X};$$

4. a stochastic kernel
$$Q(\cdot \mid h, a)$$
from $\mathcal{H} \times \mathcal{A}$ to X, interpreted as the law of the next signal after history h and action a;

5. a bounded measurable reward extractor
$$\rho : \mathsf{X} \to [-R, R]$$
for some finite $R > 0$;

6. a discount factor $\gamma \in [0, 1)$;

7. a nonempty class Π of admissible continuation controllers, where each $\pi \in \Pi$ is a universally measurable stochastic kernel from $\mathcal{H}$ to $\mathcal{A}$;

8. a nonempty deviation class Φ, where each $\phi \in \Phi$ is a self-map of Π.

For $h \in \mathcal{H}$ and $\pi \in \Pi$, let
$$(H_t, A_t, X_{t+1})_{t \geq 0}$$
denote the controlled process started from initial history $H_0 = h$, with
$$A_t \sim \pi(\cdot \mid H_t), \qquad X_{t+1} \sim Q(\cdot \mid H_t, A_t),$$
and history recursion
$$H_{t+1} := H_t \oplus (A_t, X_{t+1}),$$
where $\oplus$ denotes concatenation of one action-signal pair. Standard kernel-composition arguments yield a unique law for this process.

300 **Proposition A.2** (Induced continuation system). *Let $(\mathcal{A}, \mathsf{X}, \mathcal{H}, Q, \rho, \gamma, \Pi, \Phi)$ be a discounted mea-*
301 *surable history process. Define*

$$J(h, \pi) := \mathbb{E}_h^\pi \left[\sum_{t=0}^\infty \gamma^t \rho(X_{t+1}) \right]. \tag{A.1}$$

302 *Then J is well defined, bounded by $R/(1 - \gamma)$ in absolute value, and*

$$(\mathcal{H}, \Pi, \Phi, J)$$

303 *is a deviation-stable continuation system in the sense of Definition 2.1.*

304 *Proof.* Fix $h \in \mathcal{H}$ and $\pi \in \Pi$. Since $|\rho(X_{t+1})| \leq R$ almost surely for every t,

$$\sum_{t=0}^\infty \gamma^t |\rho(X_{t+1})| \leq \sum_{t=0}^\infty \gamma^t R = \frac{R}{1 - \gamma} \qquad \text{a.s.}$$

305 Hence the discounted sum in (A.1) converges absolutely almost surely and is integrable. Therefore
306 $J(h, \pi)$ is well defined and satisfies

$$|J(h, \pi)| \leq \frac{R}{1 - \gamma}.$$

307 All ingredients of Definition 2.1 are thus satisfied, with continuation class $\mathcal{C} = \Pi$ and value func-
308 tional J given by (A.1). $\square$

309 *Remark* A.3 (Why the abstraction is lossless for our purposes). The memory theory developed in the
310 main text depends only on the attainable history set, the continuation-controller class, the deviation
311 class, and the continuation values $J(h, c)$. Once these objects are fixed, no additional representation
312 of the environment is needed. In particular, none of the main results requires that histories arise
313 from a Markov state process, from a finite latent model, or even from an explicitly parameterized
314 environment.

A.2   Measurable low-regret correspondences

316 The main text deliberately states the transversal law in a purely set-theoretic form. For measurable
317 decision problems, one can strengthen it by requiring the history-to-memory encoder to be measur-
318 able.

319 We isolate a convenient regularity hypothesis. It is stronger than necessary, but it cleanly covers all
320 finite-controller constructions used in the paper.

321 **Assumption A.4** (Regular measurable setting). Fix $\varepsilon \geq 0$. In addition to the hypotheses of Defini-
322 tion 2.1, assume:

323 1. $\mathcal{H}$ is a standard Borel space;

324 2. $\mathcal{C}$ is a compact metric space;

325 3. the deviation class is countable,

$$\Phi = \{\phi_1, \phi_2, \ldots\},$$

326   and each $\phi_n : \mathcal{C} \to \mathcal{C}$ is continuous;

327 4. the continuation-value functional $J : \mathcal{H} \times \mathcal{C} \to \mathbb{R}$ is bounded and Carathéodory, meaning:

328   (a) for each fixed $c \in \mathcal{C}$, the map $h \mapsto J(h, c)$ is Borel measurable;

329   (b) for each fixed $h \in \mathcal{H}$, the map $c \mapsto J(h, c)$ is continuous.

330 **Proposition A.5** (Borel graph and closed values). *Under Assumption A.4, the regret map*

$$(h, c) \longmapsto \operatorname{Reg}_\Phi(h, c)$$

331 *is Borel measurable on $\mathcal{H} \times \mathcal{C}$. Moreover, for each fixed $h \in \mathcal{H}$, the function $c \mapsto \operatorname{Reg}_\Phi(h, c)$ is*
332 *lower semicontinuous. Consequently, for every $h \in \mathcal{H}$, the set*

$$K_h^\varepsilon = \{c \in \mathcal{C} : \operatorname{Reg}_\Phi(h, c) \le \varepsilon\}$$

333 *is closed, and the graph*

$$\operatorname{Gr}(K^\varepsilon) := \{(h, c) \in \mathcal{H} \times \mathcal{C} : c \in K_h^\varepsilon\}$$

334 *is a Borel subset of $\mathcal{H} \times \mathcal{C}$.*

335 *Proof.* For each $n \in \mathbb{N}$, define

$$f_n(h, c) := J(h, \phi_n(c)) - J(h, c).$$

336 Because ϕ_n is continuous and J is Carathéodory, each $f_n : \mathcal{H} \times \mathcal{C} \to \mathbb{R}$ is Borel measurable. Indeed,
337 for fixed c,

$$h \longmapsto J(h, \phi_n(c)) - J(h, c)$$

338 is Borel measurable, and for fixed h,

$$c \longmapsto J(h, \phi_n(c)) - J(h, c)$$

339 is continuous. Therefore f_n is jointly Borel measurable.

340 By definition of deviation regret,

$$\operatorname{Reg}_\Phi(h, c) = \sup_{n \ge 1} f_n(h, c).$$

341 A countable supremum of Borel functions is Borel, so Reg_Φ is Borel measurable on $\mathcal{H} \times \mathcal{C}$. For
342 each fixed h, the map $c \mapsto f_n(h, c)$ is continuous, hence

$$c \longmapsto \sup_{n \ge 1} f_n(h, c)$$

343 is lower semicontinuous as a pointwise supremum of continuous functions. This proves the lower
344 semicontinuity of $c \mapsto \operatorname{Reg}_\Phi(h, c)$.

345 Now fix $h \in \mathcal{H}$. Since K_h^ε is the ε-sublevel set of the lower semicontinuous function $c \mapsto$
346 $\operatorname{Reg}_\Phi(h, c)$, it is closed in $\mathcal{C}$. Finally,

$$\operatorname{Gr}(K^\varepsilon) = \{(h, c) \in \mathcal{H} \times \mathcal{C} : \operatorname{Reg}_\Phi(h, c) \le \varepsilon\} = \bigcap_{n \ge 1} \{(h, c) : f_n(h, c) \le \varepsilon\},$$

347 which is Borel because each f_n is Borel. $\qquad\square$

348 **Corollary A.6** (Borel membership sections). *Under Assumption A.4, for every fixed $c \in \mathcal{C}$, the*
349 *membership set*

$$A_c^\varepsilon := \{h \in \mathcal{H} : c \in K_h^\varepsilon\}$$

350 *is Borel measurable.*

351 *Proof.* For fixed $c \in \mathcal{C}$,

$$A_c^\varepsilon = \{h \in \mathcal{H} : (h, c) \in \operatorname{Gr}(K^\varepsilon)\}.$$

352 Since $\operatorname{Gr}(K^\varepsilon)$ is Borel in $\mathcal{H} \times \mathcal{C}$, its section at the fixed point c is Borel in $\mathcal{H}$. $\qquad\square$

A.3 A measurable refinement of the transversal law

354 We now define the measurable analogue of retentive complexity.

355 **Definition A.7** (Measurable past-compression complexity). Assume $\mathcal{H}$ is equipped with a σ-
356 algebra. For $\varepsilon \geq 0$, the *measurable retentive complexity* is

$$m_\Phi^{\mathrm{pc,meas}}(\varepsilon) := \inf \left\{ m \in \mathbb{N} : \begin{array}{l} \exists\, \varepsilon\text{-rational past-compression architecture } (\sigma, d) \\ \text{of size } m \text{ with measurable } \sigma \end{array} \right\},$$

357 with the convention that the infimum of the empty set is ∞.

358 **Theorem A.8** (Measurable finite-transversal law). *Under Assumption A.4, for every $\varepsilon \geq 0$,*

$$m_\Phi^{\mathrm{pc,meas}}(\varepsilon) = m_\Phi^{\mathrm{pc}}(\varepsilon) = \tau(\mathcal{K}_\varepsilon). \tag{A.2}$$

359 *Proof.* Any measurable past-compression architecture is, in particular, a past-compression architec-
360 ture, so

$$m_\Phi^{\mathrm{pc}}(\varepsilon) \leq m_\Phi^{\mathrm{pc,meas}}(\varepsilon).$$

361 By Theorem 3.1, it therefore suffices to prove

$$m_\Phi^{\mathrm{pc,meas}}(\varepsilon) \leq \tau(\mathcal{K}_\varepsilon).$$

362 If $\tau(\mathcal{K}_\varepsilon) = \infty$, there is nothing to prove. Assume $\tau(\mathcal{K}_\varepsilon) = m < \infty$. Then there exists a transversal

$$P = \{c_1, \ldots, c_m\} \subseteq \mathcal{C}$$

363 for $\mathcal{K}_\varepsilon$. Define $d : [m] \to \mathcal{C}$ by $d(i) = c_i$. For each $i \in [m]$, let

$$A_i := \{h \in \mathcal{H} : c_i \in K_h^\varepsilon\}.$$

364 By Corollary A.6, each A_i is Borel. Since P is a transversal, for every $h \in \mathcal{H}$,

$$\{i \in [m] : h \in A_i\} \neq \varnothing.$$

365 Define

$$\sigma(h) := \min\{i \in [m] : h \in A_i\}. \tag{A.3}$$

366 This is well defined. Moreover, for each $i \in [m]$,

$$\{h \in \mathcal{H} : \sigma(h) = i\} = A_i \cap \bigcap_{j<i} A_j^c,$$

367 which is Borel. Hence $\sigma : \mathcal{H} \to [m]$ is measurable.

368 Finally, by construction of σ,

$$c_{\sigma(h)} \in K_h^\varepsilon \qquad \forall h \in \mathcal{H}.$$

369 Equivalently,

$$d(\sigma(h)) \in K_h^\varepsilon \qquad \forall h \in \mathcal{H}.$$

370 Thus (σ, d) is a measurable ε-rational past-compression architecture of size m. Therefore

$$m_\Phi^{\mathrm{pc,meas}}(\varepsilon) \leq m = \tau(\mathcal{K}_\varepsilon).$$

371 Combining this with Theorem 3.1 yields (A.2). $\qquad\qquad\square$

372 *Remark* A.9 (Why finite transversals avoid measurable-selection subtleties). The proof of The-
373 orem A.8 does not invoke any general measurable-selection theorem. Once a finite transversal
374 $P = \{c_1, \ldots, c_m\}$ is fixed, the encoder is obtained canonically by the minimum-index rule (A.3).
375 Measurability then reduces to the Borel measurability of the membership sections A_i.

376 **A.4 A selection corollary**

377 Although not needed for the finite-memory results in the main text, it is worth recording the corre-
378 sponding one-selector statement. Standard measurable-selection theory implies that, under suitable
379 hypotheses, one can choose a measurable continuation $s(h) \in K_h^\varepsilon$ whenever every low-regret set is
380 nonempty.

381 *Remark* A.10 (Classical measurable-selection consequence). Suppose, in addition to Assump-
382 tion A.4, that the correspondence $h \mapsto K_h^\varepsilon$ is weakly measurable and has nonempty closed values.
383 Then the Kuratowski–Ryll-Nardzewski theorem yields a Borel selector

$$s_\varepsilon : \mathcal{H} \to \mathcal{C} \qquad \text{such that} \qquad s_\varepsilon(h) \in K_h^\varepsilon \ \forall h \in \mathcal{H};$$

384 see, for example, Aliprantis & Border (2006, Theorem 18.13) or Castaing & Valadier (1977, Chap-
385 ter III). We do not use this result in the main text, because Theorem A.8 already gives the finite-
386 memory statement needed in the paper.

387 *Remark* A.11 (Coverage of the main constructions). All finite-controller constructions in Sections 4
388 and 5 satisfy Assumption A.4. Indeed, $\mathcal{C} = \Delta_k$ is compact metrizable, the external deviation class
389 is finite and continuous, and $J(h, \cdot)$ is affine (hence continuous) for each fixed history h. Thus no
390 additional measurability burden is hidden by the abstract formulation used in the main paper.

391 # B   Finite-State Realizations and Proof of Proposition 2.5

392 This section proves the lower bound that links our retentive-complexity notion to ordinary bounded-
393 memory agents. The argument is entirely internal to the framework: it uses only the definitions of
394 the low-regret continuation sets K_h^ε, the past-compression architectures, and the induced complexity
395 $m_\Phi^{\mathrm{pc}}(\varepsilon)$. In particular, the proof does *not* assume any Markov structure, latent-state enumerability,
396 observability condition, or simplicity property of the environment.

397 We begin by isolating the exact abstraction of a finite-memory agent that is needed for the proposi-
398 tion.

399 **Definition B.1** (Finite-state realization). Fix a deviation-stable continuation system

$$\mathfrak{M} = (\mathcal{H}, \mathcal{C}, \Phi, J)$$

400 in the sense of Definition 2.1. A *finite-state realization* of $\mathfrak{M}$ is a triple

$$(Z, \zeta, d),$$

401 where:

402 1. Z is a finite set of internal states;

403 2. $\zeta : \mathcal{H} \to Z$ is a deterministic history-to-state map;

404 3. $d : Z \to \mathcal{C}$ is a decoder assigning to each internal state an admissible continuation controller.

405 We interpret $\zeta(h)$ as the internal state reached by the agent after history h, and $d(z)$ as the contin-
406 uation controller induced by starting the agent from internal state z. The realization is said to be
407 *ε-deviation-rational* if

$$\mathrm{Reg}_\Phi(h, d(\zeta(h))) \leq \varepsilon \qquad \forall h \in \mathcal{H}. \tag{B.1}$$

408 The next proposition is stronger than Proposition 2.5 in the main text because it lower bounds the
409 number of *reachable* internal states rather than the ambient cardinality of the declared state space.

410 **Proposition B.2** (Reachable-state strengthening). *Let (Z, ζ, d) be a finite-state realization of $\mathfrak{M}$ that
411 is ε-deviation-rational in the sense of* (B.1). *Define the reachable set*

$$Z_{\mathrm{reach}} := \zeta(\mathcal{H}) \subseteq Z.$$

412   *Then*

$$|Z_{\text{reach}}| \geq m_{\Phi}^{\text{pc}}(\varepsilon). \tag{B.2}$$

413   *Consequently,*

$$|Z| \geq m_{\Phi}^{\text{pc}}(\varepsilon). \tag{B.3}$$

414   *Proof.* Fix an ε-deviation-rational finite-state realization (Z, ζ, d), and let

$$m := |Z_{\text{reach}}|.$$

415   Since Z_{reach} is finite, there exists a bijection

$$b : Z_{\text{reach}} \to [m].$$

416   Using this bijection, define

$$\sigma := b \circ \zeta : \mathcal{H} \to [m]$$

417   and

$$\tilde{d} : [m] \to \mathcal{C}, \qquad \tilde{d}(i) := d\big(b^{-1}(i)\big).$$

418   We claim that $(\sigma, \tilde{d})$ is an ε-rational past-compression architecture of size m in the sense of Defini-
419   tion 2.3.

420   To verify the claim, fix any history $h \in \mathcal{H}$. Because the realization is ε-deviation-rational, (B.1)
421   yields

$$\text{Reg}_{\Phi}(h, d(\zeta(h))) \leq \varepsilon.$$

422   By Definition 2.2, this is equivalent to the membership statement

$$d(\zeta(h)) \in K_h^{\varepsilon}.$$

423   On the other hand, since b is a bijection on Z_{reach} and $\zeta(h) \in Z_{\text{reach}}$, we have

$$\tilde{d}(\sigma(h)) = \tilde{d}(b(\zeta(h))) = d\big(b^{-1}(b(\zeta(h)))\big) = d(\zeta(h)).$$

424   Combining the two displays gives

$$\tilde{d}(\sigma(h)) \in K_h^{\varepsilon} \qquad \forall h \in \mathcal{H}. \tag{B.4}$$

425   Thus $(\sigma, \tilde{d})$ satisfies the defining feasibility condition for an ε-rational past-compression architec-
426   ture.

427   By Definition 2.4 of $m_{\Phi}^{\text{pc}}(\varepsilon)$, the existence of an ε-rational past-compression architecture of size m
428   implies

$$m_{\Phi}^{\text{pc}}(\varepsilon) \leq m = |Z_{\text{reach}}|.$$

429   This proves (B.2). Since $Z_{\text{reach}} \subseteq Z$, we immediately obtain

$$m_{\Phi}^{\text{pc}}(\varepsilon) \leq |Z_{\text{reach}}| \leq |Z|,$$

430   which is (B.3). $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\square$

431   *Proof of Proposition 2.5.* Proposition 2.5 is the weaker ambient-cardinality statement (B.3), which
432   follows immediately from Proposition B.2. $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\square$

433   *Remark* B.3 (Why this proposition is the correct bridge). The proof uses only the following two
434   facts: first, every history h is mapped to a current internal state $\zeta(h)$; second, every internal state z
435   determines an admissible continuation controller $d(z)$. No further structure is needed. In particular,
436   the proposition applies verbatim to any deterministic recurrent architecture whose hidden state is a
437   deterministic function of the realized history and whose future closed-loop behavior is completely
438   determined by the current hidden state.

439   The proposition is intentionally stated for deterministic history-to-state realizations. If the internal
440   state after a fixed history were itself randomized, then the natural comparison object would be a
441   *distributional* or *mixed* version of past-compression complexity. That richer formalism is orthogonal
442   to the present paper and is not needed for any of our main results.

C   Full Proof of Theorem 3.1

This section proves the exact reduction of retentive complexity to a transversal problem. The proof is purely set-theoretic and requires no topological, measurable, or compactness assumptions. In particular, it does not use any simplifying assumption on the environment, such as latent-state finiteness, Markovity, observability, or determinism.

Throughout this section, we write

$$\mathcal{K}_\varepsilon := \{K_h^\varepsilon : h \in \mathcal{H}\},$$

where K_h^ε is the ε-rational continuation set from Definition 2.2. Since $\mathcal{H}$ is the set of attainable histories, it is nonempty in every sequential decision problem of interest. If one wishes to allow the degenerate case $\mathcal{H} = \varnothing$, all statements below remain valid after the harmless convention that both sides of Theorem 3.1 are interpreted as 0.

We begin by making the hitting-set notion explicit.

Definition C.1 (Transversal number)**.** Let $\mathcal{F}$ be a family of subsets of $\mathcal{C}$. A set $P \subseteq \mathcal{C}$ is a *transversal* (or *piercing set*) for $\mathcal{F}$ if

$$P \cap F \neq \varnothing \qquad \forall F \in \mathcal{F}.$$

The *transversal number* of $\mathcal{F}$ is

$$\tau(\mathcal{F}) := \inf\{m \in \mathbb{N} : \exists P \subseteq \mathcal{C} \text{ with } |P| \leq m \text{ and } P \cap F \neq \varnothing \, \forall F \in \mathcal{F}\},$$

with the convention $\inf \varnothing = \infty$.

Remark C.2 (Budgeted versus exact size)*.* A past-compression architecture of size m is, by Definition 2.3, a pair (σ, d) with $\sigma : \mathcal{H} \to [m]$ and $d : [m] \to \mathcal{C}$. Unused labels are permitted. Consequently, for each fixed budget m, the existence of an ε-rational architecture of size m is equivalent to the existence of one using at most m labels. We use this monotonicity implicitly below.

The heart of the argument is the following budget-by-budget equivalence.

Proposition C.3 (Budget equivalence between memory and transversals)**.** *Fix $m \in \mathbb{N}$. The following are equivalent.*

1. There exists an ε-rational past-compression architecture of size m.

2. There exists a transversal $P \subseteq \mathcal{C}$ for $\mathcal{K}_\varepsilon$ with $|P| \leq m$.

Proof. We prove the two implications separately.

(1) $\Rightarrow$ (2).   Assume there exists an ε-rational past-compression architecture (σ, d) of size m, so

$$\sigma : \mathcal{H} \to [m], \qquad d : [m] \to \mathcal{C},$$

and

$$d(\sigma(h)) \in K_h^\varepsilon \qquad \forall h \in \mathcal{H}. \tag{C.1}$$

Define

$$P := d([m]) \subseteq \mathcal{C}.$$

Then $|P| \leq m$. Fix any history $h \in \mathcal{H}$. By (C.1),

$$d(\sigma(h)) \in K_h^\varepsilon.$$

Since $d(\sigma(h)) \in P$ by construction, we obtain

$$P \cap K_h^\varepsilon \neq \varnothing.$$

Because h was arbitrary, P is a transversal for $\mathcal{K}_\varepsilon$ of cardinality at most m.

474 **(2) ⇒ (1).** Assume there exists a transversal $P \subseteq \mathcal{C}$ for $\mathcal{K}_\varepsilon$ with $|P| \leq m$. Since $\mathcal{H} \neq \varnothing$ and P
475 intersects every K_h^ε, the set P is nonempty. Write

$$P = \{c_1, \ldots, c_r\} \qquad \text{with} \qquad 1 \leq r \leq m,$$

476 where the c_i are pairwise distinct.

477 Define a decoder $d : [m] \to \mathcal{C}$ by

$$d(i) := \begin{cases} c_i, & 1 \leq i \leq r, \\ c_1, & r < i \leq m. \end{cases} \tag{C.2}$$

478 Thus the first r labels encode the transversal points, and the remaining labels (if any) are padded
479 arbitrarily using c_1.

480 Now fix $h \in \mathcal{H}$. Since P is a transversal for $\mathcal{K}_\varepsilon$, we have

$$P \cap K_h^\varepsilon \neq \varnothing.$$

481 Equivalently,

$$I(h) := \{i \in [r] : c_i \in K_h^\varepsilon\}$$

482 is nonempty. Define

$$\sigma(h) := \min I(h). \tag{C.3}$$

483 This is well defined because $I(h) \neq \varnothing$ and $I(h) \subseteq [r] \subseteq [m]$. By the definition of $I(h)$,

$$c_{\sigma(h)} \in K_h^\varepsilon.$$

484 Using (C.2), we have $d(\sigma(h)) = c_{\sigma(h)}$, hence

$$d(\sigma(h)) \in K_h^\varepsilon \qquad \forall h \in \mathcal{H}.$$

485 Therefore (σ, d) is an ε-rational past-compression architecture of size m. □

486 We now derive the main theorem.

487 *Proof of Theorem 3.1.* Recall that

$$m_\Phi^{\mathrm{pc}}(\varepsilon) = \inf\{m \in \mathbb{N} : \exists \, \varepsilon\text{-rational past-compression architecture of size } m\},$$

488 with the convention that the infimum of the empty set is ∞; see Definition 2.4. By Proposition C.3,
489 for each $m \in \mathbb{N}$,

$$\exists \, \varepsilon\text{-rational architecture of size } m \quad \Longleftrightarrow \quad \exists \, P \subseteq \mathcal{C}, \ |P| \leq m, \ P \cap K_h^\varepsilon \neq \varnothing \ \forall h \in \mathcal{H}.$$

490 Therefore the set of feasible memory budgets coincides exactly with the set of feasible transversal
491 budgets:

$$\{m \in \mathbb{N} : \exists \, \varepsilon\text{-rational architecture of size } m\} = \left\{ m \in \mathbb{N} : \begin{array}{c} \exists \, P \subseteq \mathcal{C}, \ |P| \leq m, \\ P \cap K_h^\varepsilon \neq \varnothing \ \forall h \in \mathcal{H} \end{array} \right\}.$$

492 Taking infima of these two identical subsets of $\mathbb{N}$ yields

$$m_\Phi^{\mathrm{pc}}(\varepsilon) = \tau(\mathcal{K}_\varepsilon).$$

493 This is exactly the claimed identity

$$m_\Phi^{\mathrm{pc}}(\varepsilon) = \tau(\{K_h^\varepsilon : h \in \mathcal{H}\}).$$

494 □

495 *Remark* C.4 (No choice-theoretic subtlety). The reverse implication in Proposition C.3 does not
496 invoke any form of the axiom of choice. Once a finite transversal $P = \{c_1, \ldots, c_r\}$ is enumerated,
497 the encoder σ is defined canonically by the minimum-index rule (C.3). This finitary feature is
498 important for later measurable refinements: whenever the membership events $\{h : c_i \in K_h^\varepsilon\}$ carry
499 additional structure, the minimum-index construction inherits that structure automatically.

500 *Remark* C.5 (What the theorem does *not* use). Theorem 3.1 is purely structural. It does not rely on
501 convexity of the continuation class, finiteness of the history set, compactness of $\mathcal{C}$, or any algebraic
502 property of the deviation class Φ. All such structure becomes relevant only when we attempt to
503 compute, bound, or geometrically analyze the transversal number in subsequent sections.

D   Full Proof of Theorem 4.1 and Corollary 4.2

505 This section gives the full derivation of the simplex geometry used in Section 4. Throughout, we fix:

506 • a controller dictionary $\{e_1, \ldots, e_k\}$,

507 • the mixed-continuation class

$$\Delta_k := \left\{ x \in \mathbb{R}_+^k : \sum_{i=1}^k x_i = 1 \right\},$$

508 • the external deviation class

$$\Phi_{\text{ext}} := \{\phi_i : i \in [k]\}, \qquad \phi_i(x) := e_i,$$

509 • and a continuation-value functional that is affine under mixing:

$$J(h, x) = \sum_{i=1}^k x_i J(h, e_i) \qquad \forall h \in \mathcal{H}, \ \forall x \in \Delta_k.$$

510 For each history $h \in \mathcal{H}$, write

$$v(h) := (v_1(h), \ldots, v_k(h)) \in \mathbb{R}^k, \qquad v_i(h) := J(h, e_i).$$

511 We begin with a simple geometric lemma that will be used repeatedly.

512 **Lemma D.1** (Simplex maximization identity). *Let $v \in \mathbb{R}^k$, and let*

$$M(v) := \max_{i \in [k]} v_i, \qquad A(v) := \arg\max_{i \in [k]} v_i.$$

513 *Then, for every $x \in \Delta_k$,*

$$\langle v, x \rangle \leq M(v). \tag{D.1}$$

514 *Moreover, equality holds if and only if*

$$\text{supp}(x) \subseteq A(v). \tag{D.2}$$

515 *Proof.* Fix $x \in \Delta_k$. Since $x_i \geq 0$ for all i and $\sum_{i=1}^k x_i = 1$,

$$\langle v, x \rangle = \sum_{i=1}^k x_i v_i \leq \sum_{i=1}^k x_i M(v) = M(v),$$

516 proving (D.1).

517   For the equality characterization, observe that

$$M(v) - \langle v, x \rangle = \sum_{i=1}^{k} x_i \big(M(v) - v_i \big). \tag{D.3}$$

518   Each summand on the right-hand side is nonnegative, because $x_i \geq 0$ and $M(v) - v_i \geq 0$. Therefore
519   the sum in (D.3) vanishes if and only if every summand vanishes. Since $x_i \geq 0$, this is equivalent to

$$x_i > 0 \implies M(v) - v_i = 0,$$

520   or, equivalently,

$$x_i > 0 \implies i \in A(v).$$

521   This is exactly (D.2). $\qquad\square$

522   We now prove the polyhedral description of the low-regret continuation sets.

523   *Proof of Theorem 4.1.* Fix a history $h \in \mathcal{H}$ and a mixed continuation $x \in \Delta_k$. By affineness of J
524   under mixing,

$$J(h, x) = \sum_{i=1}^{k} x_i J(h, e_i) = \sum_{i=1}^{k} x_i v_i(h) = \langle v(h), x \rangle. \tag{D.4}$$

525   Moreover, for each $i \in [k]$,

$$J(h, \phi_i x) = J(h, e_i) = v_i(h).$$

526   Substituting these identities into the definition of deviation regret yields

$$
\begin{aligned}
\mathrm{Reg}_{\Phi_{\mathrm{ext}}}(h, x) &= \sup_{i \in [k]} \Big(J(h, \phi_i x) - J(h, x) \Big) \\
&= \sup_{i \in [k]} \Big(v_i(h) - \langle v(h), x \rangle \Big) \\
&= \max_{i \in [k]} v_i(h) - \langle v(h), x \rangle.
\end{aligned}
$$

527   This proves (4.1).

528   Now apply Definition 2.2. A point $x \in \Delta_k$ belongs to K_h^ε if and only if

$$\mathrm{Reg}_{\Phi_{\mathrm{ext}}}(h, x) \leq \varepsilon,$$

529   which, by the displayed identity above, is equivalent to

$$\max_{i \in [k]} v_i(h) - \langle v(h), x \rangle \leq \varepsilon.$$

530   Rearranging gives

$$\langle v(h), x \rangle \geq \max_{i \in [k]} v_i(h) - \varepsilon,$$

531   and therefore

$$K_h^\varepsilon = \left\{ x \in \Delta_k : \langle v(h), x \rangle \geq \max_{i \in [k]} v_i(h) - \varepsilon \right\}.$$

532   This is exactly (4.2).

533   Finally, Δ_k is a simplex, hence a compact polytope, and

$$\left\{ x \in \mathbb{R}^k : \langle v(h), x \rangle \geq \max_{i \in [k]} v_i(h) - \varepsilon \right\}$$

534   is a closed half-space. Their intersection is therefore a convex polytope. $\qquad\square$

535 To derive the exact hypergraph law, it is convenient to isolate the conversion between hitting subsets
536 of $[k]$ and piercing simplex faces.

537 **Proposition D.2** (Budgeted face-piercing equivalence). *Let $\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\}$ be any family of*
538 *nonempty subsets, and for each $S \in \mathcal{S}$ define the simplex face*

$$\Delta(S) := \{x \in \Delta_k : \operatorname{supp}(x) \subseteq S\}.$$

539 *Fix $m \in \mathbb{N}$. Then the following are equivalent:*

540 *1. there exists a set $P \subseteq \Delta_k$ with $|P| \le m$ such that*

$$P \cap \Delta(S) \ne \varnothing \qquad \forall S \in \mathcal{S};$$

541 *2. there exists a set $T \subseteq [k]$ with $|T| \le m$ such that*

$$T \cap S \ne \varnothing \qquad \forall S \in \mathcal{S}.$$

542 *Consequently,*
$$\tau(\{\Delta(S) : S \in \mathcal{S}\}) = \tau(\mathcal{S}), \tag{D.5}$$

543 *where the left-hand side is the transversal number in the ambient space Δ_k, and the right-hand side*
544 *is the usual hypergraph transversal number on the vertex set $[k]$.*

545 *Proof.* We prove the two implications separately.

546 **(2) $\Rightarrow$ (1).** Assume $T \subseteq [k]$ hits every set in $\mathcal{S}$ and $|T| \le m$. Define

$$P_T := \{e_i : i \in T\} \subseteq \Delta_k.$$

547 Then $|P_T| = |T| \le m$. Let $S \in \mathcal{S}$. Since $T \cap S \ne \varnothing$, choose $i \in T \cap S$. Then $\operatorname{supp}(e_i) = \{i\} \subseteq S$,
548 so $e_i \in \Delta(S)$. Because $e_i \in P_T$, we obtain

$$P_T \cap \Delta(S) \ne \varnothing.$$

549 Since $S \in \mathcal{S}$ was arbitrary, P_T pierces every face $\Delta(S)$.

550 **(1) $\Rightarrow$ (2).** Assume $P \subseteq \Delta_k$ pierces every face $\Delta(S)$ and $|P| \le m$. Because P is finite, we may
551 write
$$P = \{x^{(1)}, \ldots, x^{(r)}\} \qquad \text{with} \qquad r \le m.$$

552 Each $x^{(j)} \in \Delta_k$ has nonempty support, since its coordinates are nonnegative and sum to 1. For each
553 $j \in [r]$, choose an index

$$i_j \in \operatorname{supp}\left(x^{(j)}\right).$$

554 Define
$$T := \{i_1, \ldots, i_r\} \subseteq [k].$$

555 Then $|T| \le r \le m$. We claim that T hits every $S \in \mathcal{S}$.

556 Fix $S \in \mathcal{S}$. Since P pierces $\Delta(S)$, there exists some $x^{(j)} \in P \cap \Delta(S)$. By definition of $\Delta(S)$,

$$\operatorname{supp}\left(x^{(j)}\right) \subseteq S.$$

557 Because $i_j \in \operatorname{supp}(x^{(j)})$, it follows that $i_j \in S$. Since $i_j \in T$, we conclude that

$$T \cap S \ne \varnothing.$$

558 Thus T is a hitting set for $\mathcal{S}$.

559 The identity (D.5) follows immediately by taking the infimum over $m \in \mathbb{N}$. □

560   *Proof of Corollary 4.2.* Fix $h \in \mathcal{H}$, and recall the definition

$$S_h := \arg\max_{i \in [k]} v_i(h).$$

561   Since $[k]$ is finite, $S_h \neq \varnothing$.

562   We first identify K_h^0. By Theorem 4.1,

$$K_h^0 = \left\{ x \in \Delta_k : \langle v(h), x \rangle \geq \max_{i \in [k]} v_i(h) \right\}.$$

563   By Lemma D.1, every $x \in \Delta_k$ satisfies

$$\langle v(h), x \rangle \leq \max_{i \in [k]} v_i(h),$$

564   with equality if and only if

$$\operatorname{supp}(x) \subseteq \arg\max_{i \in [k]} v_i(h) = S_h.$$

565   Hence

$$K_h^0 = \{ x \in \Delta_k : \operatorname{supp}(x) \subseteq S_h \} = \Delta(S_h).$$

566   Now let

$$\mathcal{S} := \{ S_h : h \in \mathcal{H} \}.$$

567   By Theorem 3.1,

$$m_{\Phi_{\mathrm{ext}}}^{\mathrm{pc}}(0) = \tau\big(\{ K_h^0 : h \in \mathcal{H} \} \big).$$

568   Using the identity $K_h^0 = \Delta(S_h)$, this becomes

$$m_{\Phi_{\mathrm{ext}}}^{\mathrm{pc}}(0) = \tau(\{ \Delta(S_h) : h \in \mathcal{H} \}).$$

569   Finally, Proposition D.2 implies

$$\tau(\{ \Delta(S_h) : h \in \mathcal{H} \}) = \tau(\mathcal{S}).$$

570   Therefore

$$m_{\Phi_{\mathrm{ext}}}^{\mathrm{pc}}(0) = \tau(\mathcal{S}),$$

571   which is exactly (4.3). $\qquad\square$

572   *Remark* D.3 (Why exact rationality collapses to support geometry). At tolerance $\varepsilon = 0$, the only
573   mixed continuations that remain deviation-rational after history h are those that place mass exclu-
574   sively on maximizing pure controllers. Thus the exact low-regret set retains only the *support pattern*
575   S_h, and all numerical information in the values $v_i(h)$ beyond the maximizer set becomes irrelevant.
576   This is precisely what allows exact retentive complexity to collapse to a hypergraph transversal
577   problem.

E   Full Proof of Theorem 4.3

579   This section proves a stronger version of Theorem 4.3. Rather than realizing only the family of exact
580   rational-support sets, we will construct, for an arbitrary finite hypergraph, a one-step history process
581   whose entire family of ε-rational continuation sets is available in closed form for every $\varepsilon \geq 0$.

582   The proof is intentionally constructive and minimal. The realized instance is a fully observable
583   one-step decision problem with binary rewards. This is important conceptually: it shows that the
584   universality phenomenon established by Theorem 4.3 is not an artifact of a rich latent-state model,
585   long-horizon dependence, or partial observability. It already appears in the simplest possible se-
586   quential setting.

587 **E.1 Construction**

588 Fix $k \in \mathbb{N}$, and let

$$\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\}$$

589 be a finite hypergraph on vertex set $[k]$. We first dispose of the degenerate case $\mathcal{S} = \varnothing$: if the
590 hypergraph is empty, the theorem is trivial, since one may take a decision process with no attainable
591 pre-action histories, in which case the realized family of rational-support sets is empty. Accordingly,
592 throughout the remainder of this section we assume

$$\mathcal{S} \neq \varnothing.$$

593 For each hyperedge $S \in \mathcal{S}$, introduce a distinct observation symbol o_S. Consider the following
594 one-step history process:

595 1. Nature samples a hyperedge $S \in \mathcal{S}$ from any distribution μ on $\mathcal{S}$ with full support, e.g.

$$\mu(S) = \frac{1}{|\mathcal{S}|} \qquad \forall S \in \mathcal{S}.$$

596 2. The agent observes o_S.

597 3. The agent selects an action $A \in [k]$.

598 4. The reward is

$$R := \mathbf{1}\{A \in S\}.$$

599 5. The episode terminates immediately.

600 Let h_S denote the pre-action history consisting of the single observation o_S. Since μ has full support,
601 the attainable pre-action histories are exactly

$$\mathcal{H}_{\mathcal{S}} := \{h_S : S \in \mathcal{S}\}.$$

602 We now specify the continuation-controller class and the deviation class. Let

$$\mathcal{C}_{\mathcal{S}} := \Delta_k$$

603 be the simplex over the canonical basis $e_1, \dots, e_k$, where e_i is the pure controller that deterministi-
604 cally chooses action i. Thus the controller dictionary has size k, namely $\{e_1, \dots, e_k\}$, and $\mathcal{C}_{\mathcal{S}}$ is the
605 class of all mixed continuations over this dictionary.

606 Let the deviation class be the external-deviation family

$$\Phi_{\text{ext}} := \{\phi_i : i \in [k]\}, \qquad \phi_i(x) := e_i \quad \forall x \in \Delta_k.$$

607 Finally, define the continuation value functional

$$J_{\mathcal{S}}(h_S, x) := \mathbb{E}[R \mid h_S, \, x] \qquad \forall S \in \mathcal{S}, \, \forall x \in \Delta_k.$$

608 Since the process terminates after one action, this expectation is completely explicit.

609 **Lemma E.1** (Continuation values in the realization instance). *For every $S \in \mathcal{S}$ and every $x \in \Delta_k$,*

$$J_{\mathcal{S}}(h_S, x) = \sum_{i \in S} x_i. \tag{E.1}$$

610 *In particular, for each pure library controller e_i,*

$$J_{\mathcal{S}}(h_S, e_i) = \mathbf{1}\{i \in S\}. \tag{E.2}$$

611 *Proof.* Fix $S \in \mathcal{S}$ and $x \in \Delta_k$. Under controller x, the action A is sampled from $[k]$ according to
612 x, so

$$\mathbb{P}(A = i \mid h_S, x) = x_i \qquad \forall i \in [k].$$

613 By definition of the reward,

$$R = \mathbf{1}\{A \in S\}.$$

614 Therefore

$$
\begin{aligned}
J_{\mathcal{S}}(h_S, x) &= \mathbb{E}[\mathbf{1}\{A \in S\} \mid h_S, x] \\
&= \sum_{i=1}^{k} \mathbb{P}(A = i \mid h_S, x)\,\mathbf{1}\{i \in S\} \\
&= \sum_{i=1}^{k} x_i\,\mathbf{1}\{i \in S\} = \sum_{i \in S} x_i,
\end{aligned}
$$

615 which proves (E.1). Evaluating this identity at $x = e_i$ gives

$$J_{\mathcal{S}}(h_S, e_i) = \sum_{j \in S} (e_i)_j = \mathbf{1}\{i \in S\},$$

616 which is (E.2). □

617 The next proposition is the stronger realization statement alluded to above.

618 **Proposition E.2** (Strong realization theorem)**.** *For the one-step history process constructed above,*
619 *the following hold for every hyperedge $S \in \mathcal{S}$.*

620 *1. The pure-controller value vector satisfies*

$$v(h_S) = (\mathbf{1}\{1 \in S\}, \ldots, \mathbf{1}\{k \in S\}) \in \{0, 1\}^k.$$

621 *2. The exact rational-support set at history h_S is*

$$\arg\max_{i \in [k]} J_{\mathcal{S}}(h_S, e_i) = S. \tag{E.3}$$

622 *3. For every $\varepsilon \geq 0$, the ε-rational continuation set is*

$$K_{h_S}^{\varepsilon} = \left\{ x \in \Delta_k : \sum_{i \in S} x_i \geq 1 - \varepsilon \right\}. \tag{E.4}$$

623 *4. In particular, at exact rationality $\varepsilon = 0$,*

$$K_{h_S}^{0} = \Delta(S) := \{x \in \Delta_k : x_i = 0 \text{ for all } i \notin S\}. \tag{E.5}$$

624 *Proof.* Fix $S \in \mathcal{S}$.

625 **Part 1.** By Lemma E.1,

$$v_i(h_S) := J_{\mathcal{S}}(h_S, e_i) = \mathbf{1}\{i \in S\} \qquad \forall i \in [k].$$

626 This is exactly the stated formula for $v(h_S)$.

627 **Part 2.** Because $S \neq \varnothing$, there exists at least one $i \in S$, and hence

$$\max_{j \in [k]} J_{\mathcal{S}}(h_S, e_j) = 1.$$

628 By Part 1,

$$J_{\mathcal{S}}(h_S, e_i) = 1 \iff i \in S.$$

629 Therefore the set of maximizers is precisely S, proving (E.3).

630   **Part 3.**   Let $x \in \Delta_k$. By definition of external deviations,

$$\phi_i(x) = e_i \qquad \forall i \in [k].$$

631   Hence the deviation regret at history h_S is

$$\begin{aligned}
\mathrm{Reg}_{\Phi_{\mathrm{ext}}}(h_S, x) &= \sup_{i \in [k]} \left(J_{\mathcal{S}}(h_S, \phi_i(x)) - J_{\mathcal{S}}(h_S, x) \right) \\
&= \sup_{i \in [k]} \left(J_{\mathcal{S}}(h_S, e_i) - J_{\mathcal{S}}(h_S, x) \right).
\end{aligned}$$

632   By Part 1 and Lemma E.1,

$$\sup_{i \in [k]} J_{\mathcal{S}}(h_S, e_i) = 1, \qquad J_{\mathcal{S}}(h_S, x) = \sum_{i \in S} x_i.$$

633   Therefore

$$\mathrm{Reg}_{\Phi_{\mathrm{ext}}}(h_S, x) = 1 - \sum_{i \in S} x_i. \tag{E.6}$$

634   By Definition 2.2,

$$x \in K_{h_S}^{\varepsilon} \iff \mathrm{Reg}_{\Phi_{\mathrm{ext}}}(h_S, x) \leq \varepsilon.$$

635   Using (E.6), this is equivalent to

$$1 - \sum_{i \in S} x_i \leq \varepsilon,$$

636   or, equivalently,

$$\sum_{i \in S} x_i \geq 1 - \varepsilon.$$

637   This proves (E.4).

638   **Part 4.**   Setting $\varepsilon = 0$ in (E.4) yields

$$K_{h_S}^0 = \left\{ x \in \Delta_k : \sum_{i \in S} x_i \geq 1 \right\}.$$

639   Since $x \in \Delta_k$ implies $x_i \geq 0$ for all i and $\sum_{i=1}^k x_i = 1$, the inequality

$$\sum_{i \in S} x_i \geq 1$$

640   is equivalent to the equality

$$\sum_{i \in S} x_i = 1,$$

641   which in turn is equivalent to

$$\sum_{i \notin S} x_i = 0.$$

642   Because all coordinates of x are nonnegative, this happens if and only if

$$x_i = 0 \qquad \forall i \notin S.$$

643   Thus

$$K_{h_S}^0 = \{ x \in \Delta_k : x_i = 0 \; \forall i \notin S \} = \Delta(S),$$

644   proving (E.5). $\square$

645   We can now derive the main-text realization theorem immediately.

646 *Proof of Theorem 4.3.* If $\mathcal{S} = \varnothing$, the result is trivial, as noted above. Assume henceforth that
647 $\mathcal{S} \neq \varnothing$, and consider the one-step history process constructed in this section.

648 By Proposition E.2, for every $S \in \mathcal{S}$,

$$\arg\max_{i \in [k]} J_{\mathcal{S}}(h_S, e_i) = S.$$

649 Therefore the family of exact rational-support sets of the constructed instance is

$$\left\{ \arg\max_{i \in [k]} J_{\mathcal{S}}(h, e_i) : h \in \mathcal{H}_{\mathcal{S}} \right\} = \left\{ \arg\max_{i \in [k]} J_{\mathcal{S}}(h_S, e_i) : S \in \mathcal{S} \right\} = \mathcal{S}.$$

650 Equivalently, if the main text denotes

$$S_h := \arg\max_{i \in [k]} v_i(h), \qquad v_i(h) = J(h, e_i),$$

651 then the realized family satisfies

$$\{ S_h : h \in \mathcal{H}_{\mathcal{S}} \} = \mathcal{S}.$$

652 This is exactly the claim of Theorem 4.3. $\square$

653 **Corollary E.3** (Memory universality at exact rationality). *For every finite hypergraph $\mathcal{S} \subseteq 2^{[k]} \setminus$*
654 *$\{\varnothing\}$, there exists a one-step history process such that*

$$m^{\mathrm{pc}}_{\Phi_{\mathrm{ext}}}(0) = \tau(\mathcal{S}).$$

655 *Proof.* Apply Theorem 4.3 to realize $\mathcal{S}$ as the family of exact rational-support sets. Then Corol-
656 lary 4.2 implies that the exact retentive complexity of the realized instance is the transversal number
657 of $\mathcal{S}$. $\square$

658 *Remark* E.4 (Why nonempty hyperedges are necessary). The restriction $\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\}$ is not
659 cosmetic. If $S = \varnothing$, then in the one-step binary-reward model one has

$$J_{\mathcal{S}}(h_S, e_i) = 0 \qquad \forall i \in [k],$$

660 so the maximizer set would be all of $[k]$, not $\varnothing$. Thus empty hyperedges cannot be realized as exact
661 rational-support sets in this model. This is precisely why the theorem is formulated for hypergraphs
662 with nonempty edges.

663 *Remark* E.5 (Minimality of the construction). The realizing instance is a contextual bandit with a
664 single decision, binary rewards, and full observability. In particular, the universality statement does
665 not rely on recurrence, hidden state, long temporal dependencies, or inaccurate models. These richer
666 phenomena are central elsewhere in the paper, but they are not needed to show that exact deviation-
667 rational memory already has the full combinatorial expressivity of arbitrary finite hypergraphs.

F Proof of Theorem 5.1 and Lemma 5.2

669 This section proves the quantitative separation showing that pairwise compatibility information
670 can be maximally uninformative about exact retentive complexity. The proof combines a finite-
671 geometric construction with the exact hypergraph law from Section 4.

F.1 Projective planes and their line hypergraphs

673 We first record the finite-geometric object that drives the separation.

674 **Definition F.1** (Finite projective plane of order q)**.** Let $q \in \mathbb{N}$. A *finite projective plane of order q* is
675 an incidence structure

$$\Pi_q = (P_q, \mathcal{L}_q, \in),$$

676 where P_q is a finite set of points and $\mathcal{L}_q$ is a finite set of lines, satisfying:

677 1. every line $L \in \mathcal{L}_q$ contains exactly $q + 1$ points;

678 2. every point $p \in P_q$ lies on exactly $q + 1$ lines;

679 3. any two distinct points lie on a unique common line;

680 4. any two distinct lines intersect in a unique common point;

5.

$$|P_q| = |\mathcal{L}_q| = q^2 + q + 1.$$

681 Associated to Π_q is the *line hypergraph*

$$\mathscr{L}_q := \{L \subseteq P_q : L \in \mathcal{L}_q\},$$

682 viewed as a hypergraph on the vertex set P_q.

683 *Remark* F.2. Theorem 5.1 is conditional on the existence of a projective plane of order q. We do
684 not need any classification result about which orders admit such planes. Once Π_q is given, the proof
685 below is entirely elementary.

686 The next lemma is the key finite-geometric counting fact.

687 **Lemma F.3.** *Let $\Pi_q = (P_q, \mathcal{L}_q, \in)$ be a finite projective plane of order q, and let $\mathscr{L}_q$ be its line*
688 *hypergraph. Then*

$$\tau(\mathscr{L}_q) = q + 1.$$

689 *Proof.* We prove the upper and lower bounds separately.

690 **Upper bound.** Fix any line $L^\star \in \mathcal{L}_q$. Since $L^\star$ contains exactly $q + 1$ points, it is a subset of P_q
691 of cardinality $q + 1$. We claim that $L^\star$ is a transversal for $\mathscr{L}_q$. Indeed, let $L \in \mathcal{L}_q$ be any line. If
692 $L = L^\star$, then trivially $L^\star \cap L = L^\star \neq \varnothing$. If $L \neq L^\star$, then by the projective-plane axiom that any
693 two distinct lines intersect in a unique point,

$$L^\star \cap L \neq \varnothing.$$

694 Thus $L^\star$ intersects every hyperedge of $\mathscr{L}_q$, so

$$\tau(\mathscr{L}_q) \leq |L^\star| = q + 1.$$

695 **Lower bound.** Let $T \subseteq P_q$ be any set with $|T| \leq q$. We will show that T is *not* a transversal for
696 $\mathscr{L}_q$.

697 For each point $p \in T$, let

$$\mathcal{L}(p) := \{L \in \mathcal{L}_q : p \in L\}$$

698 denote the set of lines through p. By Definition F.1,

$$|\mathcal{L}(p)| = q + 1 \qquad \forall p \in T.$$

699 Hence the set of lines incident to at least one point of T satisfies

$$\left| \bigcup_{p \in T} \mathcal{L}(p) \right| \leq \sum_{p \in T} |\mathcal{L}(p)|$$
$$= |T|(q + 1)$$
$$\leq q(q + 1)$$
$$= q^2 + q.$$

700   On the other hand, the total number of lines in Π_q is

$$|\mathcal{L}_q| = q^2 + q + 1.$$

701   Therefore

$$\left| \bigcup_{p \in T} \mathcal{L}(p) \right| < |\mathcal{L}_q|,$$

702   so there exists at least one line $L \in \mathcal{L}_q$ such that

$$L \notin \bigcup_{p \in T} \mathcal{L}(p).$$

703   By definition of the union, this means $p \notin L$ for every $p \in T$, i.e.

$$T \cap L = \varnothing.$$

704   Thus T fails to intersect the hyperedge L, and hence T is not a transversal.

705   We have shown that no subset of P_q of cardinality at most q can hit all lines. Therefore

$$\tau(\mathscr{L}_q) \geq q + 1.$$

706   Combining the upper and lower bounds yields

$$\tau(\mathscr{L}_q) = q + 1.$$

707 $\square$

F.2   Pairwise compatibility graph

709   For clarity, we restate the graph notion used in the main theorem.

710   **Definition F.4** (Pairwise compatibility graph at exact rationality). Let $\mathfrak{M}$ be an instance of the
711   mixed-controller model with attainable history set $\mathcal{H}$, controller class Δ_n, and exact low-regret sets
712    $\{K_h^0 : h \in \mathcal{H}\}$. The *pairwise compatibility graph* of $\mathfrak{M}$ is the simple graph

$$G_{\mathrm{pair}}(\mathfrak{M}) = (\mathcal{H}, E),$$

713   where two distinct histories $h, h' \in \mathcal{H}$ are adjacent iff

$$K_h^0 \cap K_{h'}^0 \neq \varnothing.$$

714   *Remark* F.5. The graph $G_{\mathrm{pair}}(\mathfrak{M})$ records only whether pairs of exact low-regret sets intersect. It
715   forgets all higher-order intersection geometry. Theorem 5.1 shows that this loss of information can
716   be severe: even the complete graph does not determine exact retentive complexity.

717   We now prove the main separation theorem.

718   *Proof of Theorem 5.1.* Fix a projective plane $\Pi_q = (P_q, \mathcal{L}_q, \in)$ of order q, and let

$$n := q^2 + q + 1 = |P_q| = |\mathcal{L}_q|.$$

719   Enumerate the points of P_q as $[n]$. We construct two one-step instances, denoted $\mathfrak{M}_q^{(A)}$ and $\mathfrak{M}_q^{(B)}$,
720   each with:

721   • controller dictionary $\{e_1, \ldots, e_n\}$,

722   • mixed continuation class Δ_n,

723 • external deviation class

$$\Phi_{\text{ext}} = \{\phi_i : i \in [n]\}, \qquad \phi_i(x) = e_i.$$

724 Both instances have exactly n attainable pre-action histories.

725 We will show:

726 1. $G_{\text{pair}}(\mathfrak{M}_q^{(A)})$ is the complete graph on n vertices;

727 2. $G_{\text{pair}}(\mathfrak{M}_q^{(B)})$ is also the complete graph on n vertices;

3.

$$m_{\Phi_{\text{ext}}}^{\text{pc}}(0; \mathfrak{M}_q^{(A)}) = 1, \qquad m_{\Phi_{\text{ext}}}^{\text{pc}}(0; \mathfrak{M}_q^{(B)}) = q + 1.$$

728 This will prove the theorem.

729 **Instance A: complete pairwise graph with memory** 1. Let the attainable histories of $\mathfrak{M}_q^{(A)}$ be

$$\mathcal{H}^{(A)} := \{h_1, \ldots, h_n\},$$

730 represented concretely by n distinct observations. The agent then chooses an action $A \in [n]$, and
731 the reward is

$$R^{(A)} := \mathbf{1}\{A = 1\}.$$

732 The episode terminates immediately.

733 For any history h_j and any mixed controller $x \in \Delta_n$,

$$J^{(A)}(h_j, x) = x_1.$$

734 Consequently, for each pure controller e_i,

$$J^{(A)}(h_j, e_i) = \mathbf{1}\{i = 1\}.$$

735 Hence the unique maximizing controller is e_1, and therefore

$$K_{h_j}^{0,(A)} = \Delta(\{1\}) = \{e_1\} \qquad \forall j \in [n].$$

736 It follows immediately that, for any two distinct histories $h_j, h_{j'}$,

$$K_{h_j}^{0,(A)} \cap K_{h_{j'}}^{0,(A)} = \{e_1\} \neq \varnothing.$$

737 Thus

$$G_{\text{pair}}(\mathfrak{M}_q^{(A)})$$

738 is the complete graph on n vertices.

739 Moreover, since the singleton set $\{e_1\}$ intersects every exact low-regret set, Theorem 3.1 implies

$$m_{\Phi_{\text{ext}}}^{\text{pc}}(0; \mathfrak{M}_q^{(A)}) = 1.$$

740 **Instance B: projective-plane geometry.** Let the attainable histories of $\mathfrak{M}_q^{(B)}$ be indexed by the
741 lines of Π_q:

$$\mathcal{H}^{(B)} := \{h_L : L \in \mathcal{L}_q\}.$$

742 Since $|\mathcal{L}_q| = n$, this instance also has n histories.

743 At history h_L, the agent chooses an action $A \in [n]$, interpreted as a point of P_q, and receives reward

$$R^{(B)} := \mathbf{1}\{A \in L\}.$$

744 The episode terminates immediately.

745 By Proposition E.2 applied to the hypergraph

$$\mathscr{L}_q = \{L \subseteq P_q : L \in \mathcal{L}_q\},$$

746 the exact low-regret set at history h_L is

$$K_{h_L}^{0,(B)} = \Delta(L) \qquad \forall L \in \mathcal{L}_q.$$

747 Now let $L, L' \in \mathcal{L}_q$ be distinct. Since Π_q is a projective plane, the two lines intersect in a unique
748 point $p \in P_q$. Therefore

$$e_p \in \Delta(L) \cap \Delta(L') = K_{h_L}^{0,(B)} \cap K_{h_{L'}}^{0,(B)}.$$

749 Hence every pair of distinct histories in $\mathfrak{M}_q^{(B)}$ is adjacent, so

$$G_{\mathrm{pair}}(\mathfrak{M}_q^{(B)})$$

750 is also the complete graph on n vertices.

751 Finally, by Corollary 4.2, the exact retentive complexity of $\mathfrak{M}_q^{(B)}$ is the transversal number of the
752 family of support sets $\mathscr{L}_q$. Lemma 5.2 therefore gives

$$m_{\Phi_{\mathrm{ext}}}^{\mathrm{pc}}(0; \mathfrak{M}_q^{(B)}) = \tau(\mathscr{L}_q) = q + 1.$$

753 **Conclusion.** The two instances $\mathfrak{M}_q^{(A)}$ and $\mathfrak{M}_q^{(B)}$ have the same pairwise compatibility graph—
754 namely the complete graph on n vertices, where $n = q^2 + q + 1$—but their exact retentive complex-
755 ities are 1 and $q + 1$, respectively. This proves Theorem 5.1. □

756 **Corollary F.6** (Quantitative failure of pairwise determination). *For every projective plane of or-*
757 *der q, there exist two exact-rationality instances with identical pairwise compatibility graph whose*
758 *retentive complexities differ by a factor of $q + 1$.*

759 *Proof.* This is immediate from Theorem 5.1, since the two realized complexities are 1 and $q+1$. □

760 *Remark* F.7 (Why the complete graph is the strongest possible host graph). Theorem 5.1 uses the
761 complete compatibility graph, which is the graph containing the *most* pairwise information one
762 could hope for at the graph level. Thus the obstruction is not that the graph is too sparse to reveal
763 the right answer. Rather, the obstruction is genuinely higher-order: pairwise intersections do not
764 determine the transversal number of the family of exact low-regret sets.

765 *Remark* F.8 (Same controller class, same deviation class). The two instances in the proof are
766 matched in a stronger sense than claimed in the theorem statement. They use the same controller
767 simplex Δ_n, the same external deviation family, and the same number n of attainable histories. The
768 only difference lies in the geometry of the exact low-regret continuation sets. This isolates the source
769 of the separation precisely.

G Full Proof of Corollary 6.1

771 This section gives a complete proof of Corollary 6.1. The argument has two ingredients. The first is a
772 finite-dimensional Helly reduction inside the affine hull of the simplex. The second is a compactness
773 argument upgrading finite intersection to intersection of the entire history-indexed family.

774 ### G.1 Finite-family Helly reduction inside the simplex

775 We begin with the finite-family statement.

776 **Lemma G.1** (Helly reduction on the simplex). *Let $\mathcal{F}$ be a finite family of convex subsets of Δ_k. If*
777 *every subfamily of $\mathcal{F}$ of cardinality at most k has nonempty intersection, then*

$$\bigcap_{F \in \mathcal{F}} F \neq \varnothing.$$

778 *Proof.* Let

$$H := \left\{ x \in \mathbb{R}^k : \sum_{i=1}^k x_i = 1 \right\}.$$

779 Then $\Delta_k \subseteq H$, and H is an affine hyperplane of dimension $k - 1$. Consider the affine bijection

$$T : \mathbb{R}^{k-1} \to H, \qquad T(z_1, \ldots, z_{k-1}) = \left(z_1, \ldots, z_{k-1}, 1 - \sum_{i=1}^{k-1} z_i \right).$$

780 For each $F \in \mathcal{F}$, define

$$\widetilde{F} := T^{-1}(F) \subseteq \mathbb{R}^{k-1}.$$

781 Since T is affine and F is convex, each $\widetilde{F}$ is convex in $\mathbb{R}^{k-1}$. Moreover, because T is bijective,

$$\bigcap_{j=1}^m F_j \neq \varnothing \quad \Longleftrightarrow \quad \bigcap_{j=1}^m \widetilde{F}_j \neq \varnothing$$

782 for every finite subfamily $\{F_1, \ldots, F_m\} \subseteq \mathcal{F}$.

783 Hence every subfamily of $\left\{ \widetilde{F} : F \in \mathcal{F} \right\}$ of cardinality at most $k = (k - 1) + 1$ has nonempty
784 intersection. By Helly's theorem in $\mathbb{R}^{k-1}$ (Matoušek, 2002, Chapter 2),

$$\bigcap_{F \in \mathcal{F}} \widetilde{F} \neq \varnothing.$$

785 Applying T gives

$$\bigcap_{F \in \mathcal{F}} F \neq \varnothing,$$

786 as claimed. $\square$

787 ### G.2 Closedness and compactness

788 The next lemma upgrades the finite-family conclusion to the full history-indexed family.

789 **Lemma G.2** (Finite intersection property on the simplex). *Let $\{F_\alpha : \alpha \in I\}$ be a family of closed*
790 *subsets of the simplex Δ_k. If every finite subfamily has nonempty intersection, then*

$$\bigcap_{\alpha \in I} F_\alpha \neq \varnothing.$$

791 *Proof.* The simplex Δ_k is compact as a closed and bounded subset of $\mathbb{R}^k$. Suppose, for contradic-
792 tion, that

$$\bigcap_{\alpha \in I} F_\alpha = \varnothing.$$

793 Then

$$\Delta_k \subseteq \bigcup_{\alpha \in I} (\Delta_k \setminus F_\alpha).$$

794 Because each F_α is closed in Δ_k, the relative complement $\Delta_k \setminus F_\alpha$ is open in Δ_k. Thus
795 $\{\Delta_k \setminus F_\alpha : \alpha \in I\}$ is an open cover of the compact space Δ_k, and so admits a finite subcover:

$$\Delta_k \subseteq \bigcup_{j=1}^m (\Delta_k \setminus F_{\alpha_j})$$

796 for some $\alpha_1, \ldots, \alpha_m \in I$. Equivalently,

$$\bigcap_{j=1}^m F_{\alpha_j} = \varnothing,$$

797 contradicting the finite intersection hypothesis. □

798 We now verify the closedness property needed to apply Lemma G.2 to the low-regret continuation
799 sets.

800 **Lemma G.3** (Closedness of low-regret sets in the mixed-controller model)**.** *In the finite-controller*
801 *mixed-continuation model of Section 4, for every history $h \in \mathcal{H}$ and every $\varepsilon \geq 0$, the set K_h^ε is a*
802 *closed convex subset of Δ_k.*

803 *Proof.* By Theorem 4.1,

$$K_h^\varepsilon = \left\{ x \in \Delta_k : \langle v(h), x \rangle \geq \max_{i \in [k]} v_i(h) - \varepsilon \right\}.$$

804 The map $x \mapsto \langle v(h), x \rangle$ is affine, hence continuous, on $\mathbb{R}^k$. It follows that

$$\left\{ x \in \mathbb{R}^k : \langle v(h), x \rangle \geq \max_{i \in [k]} v_i(h) - \varepsilon \right\}$$

805 is a closed half-space. Therefore K_h^ε, being the intersection of this closed half-space with the sim-
806 plex Δ_k, is closed in Δ_k. It is also convex because both sets in the intersection are convex. □

G.3 Proof of the corollary

808 *Full proof of Corollary 6.1.* We prove the equivalence of the three statements in the corollary.

809 **(1) $\Longleftrightarrow$ (2).** By Theorem 3.1,

$$m_{\Phi_{\mathrm{ext}}}^{\mathrm{pc}}(\varepsilon) = 1 \quad \Longleftrightarrow \quad \tau(\{K_h^\varepsilon : h \in \mathcal{H}\}) = 1.$$

810 Since $\mathcal{H} \neq \varnothing$ by Definition 2.1, a transversal of cardinality at most 1 must be a singleton $\{x\}$ for
811 some $x \in \Delta_k$. Such a singleton pierces every K_h^ε if and only if

$$x \in K_h^\varepsilon \qquad \forall h \in \mathcal{H},$$

812 which is equivalent to

$$\bigcap_{h \in \mathcal{H}} K_h^\varepsilon \neq \varnothing.$$

813 This proves (1) $\Longleftrightarrow$ (2).

814 **(2) $\Longrightarrow$ (3).** This implication is immediate: if the total intersection is nonempty, then every sub-
815 family, in particular every subfamily of cardinality at most k, also has nonempty intersection.

816 **(3) $\implies$ (2).** Assume that every subfamily of

$$\mathcal{K}_\varepsilon := \{K_h^\varepsilon : h \in \mathcal{H}\}$$

817 of cardinality at most k has nonempty intersection.

818 We first show that every *finite* subfamily of $\mathcal{K}_\varepsilon$ has nonempty intersection. Let

$$\mathcal{F} \subseteq \mathcal{K}_\varepsilon$$

819 be finite. By Lemma G.3, every member of $\mathcal{F}$ is convex and contained in Δ_k. Moreover, every sub-
820 family of $\mathcal{F}$ of cardinality at most k has nonempty intersection by the assumption of (3). Therefore
821 Lemma G.1 applies and yields

$$\bigcap_{F \in \mathcal{F}} F \neq \varnothing.$$

822 Since $\mathcal{F}$ was arbitrary, the family $\mathcal{K}_\varepsilon$ has the finite intersection property.

823 Next, Lemma G.3 shows that every K_h^ε is closed in the compact simplex Δ_k. Hence Lemma G.2
824 implies

$$\bigcap_{h \in \mathcal{H}} K_h^\varepsilon \neq \varnothing.$$

825 This is exactly statement (2).

826 Combining the three implications proves the equivalence of (1), (2), and (3). $\qquad\square$

827 *Remark* G.4 (Why the threshold is k). The threshold k is not arbitrary. The family $\{K_h^\varepsilon : h \in \mathcal{H}\}$
828 lives inside the affine hull of the simplex,

$$\mathrm{aff}(\Delta_k) = \left\{ x \in \mathbb{R}^k : \sum_{i=1}^k x_i = 1 \right\},$$

829 which has dimension $k-1$. Helly's theorem therefore yields a local intersection certificate of order
830 $(k-1)+1 = k$. The certificate depends only on the size of the controller dictionary and not on any
831 measure of environmental complexity.

832 *Remark* G.5 (Finite-history special case). If $\mathcal{H}$ is finite, the compactness step is unnecessary. In that
833 case statement (3) implies statement (2) directly from Lemma G.1 by applying it to the finite family
834 $\{K_h^\varepsilon : h \in \mathcal{H}\}$.

835 # H   Richer Deviation Classes

836 The main text focuses on external deviations because they already yield a sharp, exact memory law.
837 This section shows that the geometric picture is robust to substantially richer deviation families. In
838 particular:

839 1. finite affine deviation classes still induce polyhedral low-regret sets;

840 2. the one-state Helly certificate extends unchanged to such classes;

841 3. enriching the deviation class can only increase memory complexity in general;

842 4. once the deviation class contains the external deviations, the entire regret geometry is already
843   saturated: for every $\varepsilon \geq 0$, the low-regret sets and retentive complexities coincide exactly with
844   those of the external-deviation model.

845 This places the external-deviation analysis in the main text on a firmer foundation: the support ge-
846 ometry is not an artifact of using only the simplest deviation family, but rather a stable consequence
847 of allowing at least the pure-action replacement deviations emphasized in the hindsight-rationality
848 literature (Morrill et al., 2021).

H.1 Finite affine deviation classes

Fix the finite-controller mixed-continuation model of Section 4. Thus the continuation class is the simplex

$$\Delta_k := \left\{ x \in \mathbb{R}^k_+ : \sum_{i=1}^k x_i = 1 \right\},$$

and continuation values are affine under mixing:

$$J(h, x) = \langle v(h), x \rangle, \qquad v(h) := (v_1(h), \dots, v_k(h)) \in \mathbb{R}^k.$$

Definition H.1 (Finite affine deviation class)**.** A *finite affine deviation class* on Δ_k is a finite family

$$\Phi = \{\phi^1, \dots, \phi^m\}$$

of affine self-maps $\phi^\ell : \Delta_k \to \Delta_k$, each of the form

$$\phi^\ell(x) = M_\ell x + b_\ell,$$

where $M_\ell \in \mathbb{R}^{k \times k}$ and $b_\ell \in \mathbb{R}^k$ satisfy $\phi^\ell(\Delta_k) \subseteq \Delta_k$.

Remark H.2 (Examples)*.* This class includes:

1. *external deviations*: $\phi_i(x) = e_i$, obtained by taking $M_i = 0$ and $b_i = e_i$;

2. *permutation deviations*: $\phi(x) = Px$, where P is a permutation matrix;

3. *stochastic mixing deviations*: $\phi(x) = Mx$, where M is a column-stochastic matrix;

4. *reset-and-mix deviations*: $\phi(x) = \alpha e_i + (1 - \alpha)Mx$ with $\alpha \in [0, 1]$, $i \in [k]$, and M column-stochastic.

Thus the affine formalism captures a broad family of structured deviations while preserving exact tractability.

Proposition H.3 (Polyhedral low-regret sets for finite affine deviations)**.** *Let* $\Phi = \{\phi^1, \dots, \phi^m\}$ *be a finite affine deviation class on* Δ_k, *with*

$$\phi^\ell(x) = M_\ell x + b_\ell.$$

Then, for every history $h \in \mathcal{H}$ *and every* $x \in \Delta_k$,

$$\mathrm{Reg}_\Phi(h, x) = \max_{1 \le \ell \le m} \left(\langle (M_\ell^\top - I)v(h), x \rangle + \langle v(h), b_\ell \rangle \right). \tag{H.1}$$

Consequently, for every $\varepsilon \ge 0$,

$$K^\varepsilon_{h,\Phi} = \Delta_k \cap \bigcap_{\ell=1}^m \left\{ x \in \mathbb{R}^k : \langle (M_\ell^\top - I)v(h), x \rangle \le \varepsilon - \langle v(h), b_\ell \rangle \right\}. \tag{H.2}$$

In particular, $K^\varepsilon_{h,\Phi}$ *is a closed convex polytope.*

Proof. Fix $h \in \mathcal{H}$, $x \in \Delta_k$, and $\ell \in [m]$. Since $J(h, \cdot)$ is affine under mixing and $\phi^\ell(x) = M_\ell x + b_\ell$,

$$J(h, \phi^\ell(x)) = \langle v(h), M_\ell x + b_\ell \rangle = \langle M_\ell^\top v(h), x \rangle + \langle v(h), b_\ell \rangle.$$

Therefore

$$J(h, \phi^\ell(x)) - J(h, x) = \langle (M_\ell^\top - I)v(h), x \rangle + \langle v(h), b_\ell \rangle.$$

Taking the maximum over $\ell \in [m]$ gives (H.1).

Now $x \in K^\varepsilon_{h,\Phi}$ iff $\mathrm{Reg}_\Phi(h, x) \le \varepsilon$. By (H.1), this is equivalent to the simultaneous inequalities

$$\langle (M_\ell^\top - I)v(h), x \rangle + \langle v(h), b_\ell \rangle \le \varepsilon \qquad \forall \ell \in [m],$$

873   or equivalently

$$\left\langle (M_\ell^\top - I)v(h), x \right\rangle \leq \varepsilon - \langle v(h), b_\ell \rangle \qquad \forall \ell \in [m].$$

874   This proves (H.2). Since $K_{h,\Phi}^\varepsilon$ is the intersection of the simplex Δ_k with finitely many closed
875   half-spaces, it is a closed convex polytope.   □

Corollary H.4 (Helly certificate for finite affine deviations). *Let Φ be any finite affine deviation*
877   *class on Δ_k. Then, for every $\varepsilon \geq 0$, the following are equivalent:*

878   *1. $m_\Phi^{\mathrm{pc}}(\varepsilon) = 1$;*

879   *2. $\bigcap_{h \in \mathcal{H}} K_{h,\Phi}^\varepsilon \neq \varnothing$;*

880   *3. every subfamily of $\left\{ K_{h,\Phi}^\varepsilon : h \in \mathcal{H} \right\}$ of cardinality at most k has nonempty intersection.*

881   *Proof.* By Proposition H.3, every $K_{h,\Phi}^\varepsilon$ is a closed convex subset of Δ_k. The proof of Corollary 6.1
882   in Supplement G applies verbatim with K_h^ε replaced by $K_{h,\Phi}^\varepsilon$.   □

H.2   Monotonicity under richer deviation classes

884   We next record a monotonicity principle that will be useful for interpreting the main results as lower
885   bounds for more demanding deviation notions.

Proposition H.5 (Monotonicity under deviation enrichment). *Let Φ_1 and Φ_2 be two deviation*
887   *classes on the same continuation system, with*

$$\Phi_1 \subseteq \Phi_2.$$

888   *Then, for every history $h \in \mathcal{H}$ and every $\varepsilon \geq 0$,*

$$K_{h,\Phi_2}^\varepsilon \subseteq K_{h,\Phi_1}^\varepsilon. \tag{H.3}$$

889   *Consequently,*

$$m_{\Phi_2}^{\mathrm{pc}}(\varepsilon) \geq m_{\Phi_1}^{\mathrm{pc}}(\varepsilon). \tag{H.4}$$

890   *Proof.* Fix $h \in \mathcal{H}$ and $c \in \mathcal{C}$. Since $\Phi_1 \subseteq \Phi_2$,

$$\mathrm{Reg}_{\Phi_1}(h, c) = \sup_{\phi \in \Phi_1} \left(J(h, \phi c) - J(h, c) \right) \leq \sup_{\phi \in \Phi_2} \left(J(h, \phi c) - J(h, c) \right) = \mathrm{Reg}_{\Phi_2}(h, c).$$

891   Hence $\mathrm{Reg}_{\Phi_2}(h, c) \leq \varepsilon$ implies $\mathrm{Reg}_{\Phi_1}(h, c) \leq \varepsilon$, which is exactly (H.3).

892   Now let

$$\mathcal{K}_{\Phi_r,\varepsilon} := \left\{ K_{h,\Phi_r}^\varepsilon : h \in \mathcal{H} \right\}, \qquad r \in \{1, 2\}.$$

893   By (H.3), every transversal of $\mathcal{K}_{\Phi_2,\varepsilon}$ is also a transversal of $\mathcal{K}_{\Phi_1,\varepsilon}$. Therefore

$$\tau(\mathcal{K}_{\Phi_2,\varepsilon}) \geq \tau(\mathcal{K}_{\Phi_1,\varepsilon}).$$

894   Applying Theorem 3.1 on both sides yields (H.4).   □

H.3   Saturation by external deviations

896   The next theorem is the corrected robustness statement. In the finite-controller simplex model, once
897   external deviations are available, no additional simplex-valued deviations can produce larger regret.
898   Thus the external deviations already determine the *entire* low-regret geometry, not merely its exact-
899   rationality slice.

900 **Theorem H.6** (Saturation by external deviations). *Consider the finite-controller mixed-continuation*
901 *model on Δ_k, and let Φ be any family of self-maps $\phi : \Delta_k \to \Delta_k$ such that*

$$\Phi_{\text{ext}} \subseteq \Phi.$$

902 *Then, for every history $h \in \mathcal{H}$, every $x \in \Delta_k$, and every $\varepsilon \geq 0$,*

$$\text{Reg}_{\Phi}(h, x) = \text{Reg}_{\Phi_{\text{ext}}}(h, x) = \max_{i \in [k]} v_i(h) - \langle v(h), x \rangle. \tag{H.5}$$

903 *Consequently,*

$$K_{h,\Phi}^{\varepsilon} = K_{h,\Phi_{\text{ext}}}^{\varepsilon} = \left\{ x \in \Delta_k : \langle v(h), x \rangle \geq \max_{i \in [k]} v_i(h) - \varepsilon \right\}, \tag{H.6}$$

904 *and*

$$m_{\Phi}^{\text{pc}}(\varepsilon) = m_{\Phi_{\text{ext}}}^{\text{pc}}(\varepsilon) \qquad \forall \varepsilon \geq 0. \tag{H.7}$$

905 *Proof.* Fix $h \in \mathcal{H}$ and $x \in \Delta_k$, and write

$$M_h := \max_{i \in [k]} v_i(h).$$

906 Because $\Phi_{\text{ext}} \subseteq \Phi$, we immediately have

$$\text{Reg}_{\Phi}(h, x) \geq \text{Reg}_{\Phi_{\text{ext}}}(h, x).$$

907 By Theorem 4.1,

$$\text{Reg}_{\Phi_{\text{ext}}}(h, x) = M_h - \langle v(h), x \rangle.$$

908 Hence

$$\text{Reg}_{\Phi}(h, x) \geq M_h - \langle v(h), x \rangle. \tag{H.8}$$

909 For the reverse inequality, let $\phi \in \Phi$. Since $\phi(x) \in \Delta_k$, Lemma D.1 implies

$$J(h, \phi(x)) = \langle v(h), \phi(x) \rangle \leq M_h.$$

910 Therefore

$$J(h, \phi(x)) - J(h, x) \leq M_h - \langle v(h), x \rangle.$$

911 Taking the supremum over $\phi \in \Phi$ yields

$$\text{Reg}_{\Phi}(h, x) \leq M_h - \langle v(h), x \rangle.$$

912 Combining this with (H.8) proves (H.5).

913 The set identity (H.6) follows immediately from the definition of low-regret sets:

$$x \in K_{h,\Phi}^{\varepsilon} \iff \text{Reg}_{\Phi}(h, x) \leq \varepsilon \iff M_h - \langle v(h), x \rangle \leq \varepsilon.$$

914 Finally, applying Theorem 3.1 to the two equal families $\left\{ K_{h,\Phi}^{\varepsilon} : h \in \mathcal{H} \right\}$ and $\left\{ K_{h,\Phi_{\text{ext}}}^{\varepsilon} : h \in \mathcal{H} \right\}$
915 gives (H.7). □

916 **Corollary H.7** (Exact hypergraph law under arbitrary external enrichment). *In the finite-controller*
917 *mixed-continuation model, for every deviation class Φ satisfying*

$$\Phi_{\text{ext}} \subseteq \Phi \subseteq \{\phi : \Delta_k \to \Delta_k\},$$

918 *one has*

$$K_{h,\Phi}^0 = \Delta(S_h) \qquad \forall h \in \mathcal{H},$$

919 *and therefore*

$$m_{\Phi}^{\text{pc}}(0) = \tau(\{S_h : h \in \mathcal{H}\}).$$

920 *Proof.* Set $\varepsilon = 0$ in (H.6), then invoke Lemma D.1 exactly as in the proof of Corollary 4.2. $\qquad\square$

921 **Corollary H.8** (Helly certificate under arbitrary external enrichment)**.** *Under the same hypotheses*
922 *as Corollary H.7, for every $\varepsilon \geq 0$, the following are equivalent:*

923 *1.* $m_{\Phi}^{\text{pc}}(\varepsilon) = 1$;

924 *2.* $\bigcap_{h \in \mathcal{H}} K_{h,\Phi}^{\varepsilon} \neq \varnothing$;

925 *3. every subfamily of $\left\{ K_{h,\Phi}^{\varepsilon} : h \in \mathcal{H} \right\}$ of cardinality at most k has nonempty intersection.*

926 *Proof.* By Theorem H.6,

$$K_{h,\Phi}^{\varepsilon} = K_{h,\Phi_{\text{ext}}}^{\varepsilon} \qquad \forall h \in \mathcal{H}, \ \forall \varepsilon \geq 0.$$

927 Thus the statement is exactly Corollary 6.1. $\qquad\square$

928 *Remark* H.9 (Where richer deviations still matter). If Φ is an arbitrary finite affine deviation class,
929 then Proposition H.3 shows that the geometry of $K_{h,\Phi}^{\varepsilon}$ generally depends on the chosen deviations.
930 The saturation theorem identifies the precise threshold at which that dependence disappears: once
931 the external deviations are included, the finite-controller simplex model is already maximally adver-
932 sarial, and no further simplex-valued deviation can enlarge regret or increase memory complexity at
933 any tolerance $\varepsilon \geq 0$.

I Fractional Memory Relaxation and LP Duality

935 This section develops a linear-programming relaxation of exact retentive complexity in the finite-
936 controller model. The resulting object is a fractional hitting-set problem, or equivalently a fractional
937 face-piercing problem on the simplex. This viewpoint is not needed for any claim in the main paper,
938 but it sharpens the optimization structure underlying the exact hypergraph law.

939 Throughout this section, fix a finite support family

$$\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\},$$

940 typically $\mathcal{S} = \{S_h : h \in \mathcal{H}\}$ from Corollary 4.2.

I.1 Fractional exact retentive complexity

942 **Definition I.1** (Fractional transversal number)**.** The *fractional transversal number* of $\mathcal{S}$ is

$$\tau^*(\mathcal{S}) := \min\left\{ \sum_{i=1}^{k} w_i : w \in \mathbb{R}_+^k, \ \sum_{i \in S} w_i \geq 1 \ \forall S \in \mathcal{S} \right\}. \tag{I.1}$$

943 The vector w can be interpreted as assigning nonnegative *memory mass* to the pure library controllers
944 $e_1, \ldots, e_k$, with the requirement that each support set $S \in \mathcal{S}$ receive total mass at least 1. This is
945 the natural LP relaxation of the integral hitting-set problem.

946 **Proposition I.2** (Fractional relaxation of exact retentive complexity)**.** *For every finite support family*
947 $\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\}$,

$$\tau^*(\mathcal{S}) \leq \tau(\mathcal{S}). \tag{I.2}$$

948 *Consequently, in the exact-rationality mixed-controller model,*

$$\tau^*(\{S_h : h \in \mathcal{H}\}) \leq m_{\Phi_{\text{ext}}}^{\text{pc}}(0). \tag{I.3}$$

949 *Proof.* Let $T \subseteq [k]$ be any transversal for $\mathcal{S}$. Define

$$w_i := \mathbf{1}\{i \in T\} \qquad \forall i \in [k].$$

950 Then $w \in \mathbb{R}_+^k$, $\sum_{i=1}^k w_i = |T|$, and for every $S \in \mathcal{S}$,

$$\sum_{i \in S} w_i = |S \cap T| \geq 1$$

951 because T hits S. Thus w is feasible for (I.1), so

$$\tau^*(\mathcal{S}) \leq |T|.$$

952 Taking the minimum over all integral transversals T proves (I.2). The inequality (I.3) now follows
953 from Corollary 4.2. $\qquad\square$

I.2 Fractional face-piercing on the simplex

955 The main text identifies exact memory with the transversal number of simplex faces $\Delta(S)$. The next
956 proposition shows that the *fractional* relaxation is equally insensitive to mixing: allowing weighted
957 mixed continuations yields exactly the same optimum as weighting only the vertices $e_1, \ldots, e_k$.

958 **Definition I.3** (Fractional face-piercing number). For a family $\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\}$, define

$$\Delta(S) := \{x \in \Delta_k : \operatorname{supp}(x) \subseteq S\}.$$

959 Its *fractional face-piercing number* is

$$\tau_{\text{face}}^*(\mathcal{S}) := \inf \left\{ \sum_{x \in P} \lambda_x : \begin{array}{l} P \subseteq \Delta_k \text{ finite, } \lambda_x \geq 0 \, \forall x \in P, \\ \sum_{x \in P \cap \Delta(S)} \lambda_x \geq 1 \ \forall S \in \mathcal{S} \end{array} \right\}. \tag{I.4}$$

960 **Proposition I.4** (Fractional face-piercing equals fractional hitting). *For every finite support family*
961 $\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\}$,
$$\tau_{\text{face}}^*(\mathcal{S}) = \tau^*(\mathcal{S}). \tag{I.5}$$

962 *Proof.* We prove the two inequalities separately.

963 **Step 1:** $\tau_{\text{face}}^*(\mathcal{S}) \leq \tau^*(\mathcal{S})$. Let $w \in \mathbb{R}_+^k$ be feasible for (I.1). Set

$$P := \{e_i : w_i > 0\} \subseteq \Delta_k, \qquad \lambda_{e_i} := w_i.$$

964 Fix $S \in \mathcal{S}$. Since $e_i \in \Delta(S)$ whenever $i \in S$,

$$\sum_{x \in P \cap \Delta(S)} \lambda_x = \sum_{i \in S} w_i \geq 1.$$

965 Hence (P, λ) is feasible for (I.4) with objective value $\sum_{i=1}^k w_i$. Taking the infimum over feasible w
966 gives
$$\tau_{\text{face}}^*(\mathcal{S}) \leq \tau^*(\mathcal{S}).$$

967 **Step 2:** $\tau^*(\mathcal{S}) \leq \tau_{\text{face}}^*(\mathcal{S})$. Let $P \subseteq \Delta_k$ be finite, and let $\lambda_x \geq 0$ for $x \in P$, be feasible for (I.4).
968 Since every $x \in \Delta_k$ has nonempty support, choose once and for all an index

$$i(x) \in \operatorname{supp}(x) \qquad \forall x \in P.$$

969 Define $w \in \mathbb{R}_+^k$ by

$$w_i := \sum_{x \in P : i(x) = i} \lambda_x \qquad \forall i \in [k].$$

970 Then

$$\sum_{i=1}^k w_i = \sum_{x \in P} \lambda_x.$$

971 Now fix $S \in \mathcal{S}$. If $x \in P \cap \Delta(S)$, then $\mathrm{supp}(x) \subseteq S$, hence $i(x) \in S$. Therefore

$$\sum_{i \in S} w_i = \sum_{x \in P : i(x) \in S} \lambda_x \geq \sum_{x \in P \cap \Delta(S)} \lambda_x \geq 1.$$

972 Thus w is feasible for (I.1) with the same objective value as (P, λ). Taking the infimum over feasible
973 (P, λ) yields

$$\tau^*(\mathcal{S}) \leq \tau^*_{\mathrm{face}}(\mathcal{S}).$$

974 Combining the two inequalities proves (I.5). □

975 *Remark* I.5 (No mixed-controller advantage in the fractional relaxation). Proposition I.4 shows that
976 the LP relaxation already lives on the vertices of the controller simplex. Allowing weighted mixed
977 continuations does not decrease the optimum beyond what can already be achieved by weighting
978 pure controllers.

I.3   LP duality and fractional packings

980 Let $A \in \{0,1\}^{\mathcal{S} \times [k]}$ denote the incidence matrix of the hypergraph $\mathcal{S}$:

$$A_{S,i} := \mathbf{1}\{i \in S\} \qquad \forall S \in \mathcal{S}, \ \forall i \in [k].$$

981 Then (I.1) can be written compactly as

$$\tau^*(\mathcal{S}) = \min\{\mathbf{1}^\top w : Aw \geq \mathbf{1}, \ w \geq 0\}. \tag{I.6}$$

982 Its dual is the fractional packing problem.

983 **Definition I.6** (Fractional packing number). The *fractional packing number* of $\mathcal{S}$ is

$$\nu^*(\mathcal{S}) := \max\left\{ \sum_{S \in \mathcal{S}} y_S : y \in \mathbb{R}_+^{\mathcal{S}}, \ \sum_{S \ni i} y_S \leq 1 \ \forall i \in [k] \right\}. \tag{I.7}$$

984 **Theorem I.7** (Strong duality for fractional memory). *For every finite support family $\mathcal{S} \subseteq 2^{[k]} \setminus \{\varnothing\}$,*

$$\tau^*(\mathcal{S}) = \nu^*(\mathcal{S}). \tag{I.8}$$

985 *Moreover, both optima are attained.*

986 *Proof.* The primal problem (I.6) is feasible, since the vector

$$w = \mathbf{1} \in \mathbb{R}^k$$

987 satisfies $Aw \geq \mathbf{1}$ because every hyperedge $S \in \mathcal{S}$ is nonempty. It is also bounded below by 0.

988 We next show that an optimal primal solution exists. Let w be any primal-feasible vector. Define
989 $\widetilde{w} \in \mathbb{R}_+^k$ coordinatewise by

$$\widetilde{w}_i := \min\{w_i, 1\}.$$

990 Then $\widetilde{w}_i \leq w_i$ for all i, so

$$\mathbf{1}^\top \widetilde{w} \leq \mathbf{1}^\top w.$$

991 Moreover, for every $S \in \mathcal{S}$,

$$\sum_{i \in S} \widetilde{w}_i \geq 1.$$

992 Indeed, if some $w_i \geq 1$ for $i \in S$, then $\widetilde{w}_i = 1$, and the sum is already at least 1; otherwise all
993 coordinates on S satisfy $w_i < 1$, so $\widetilde{w}_i = w_i$ on S, and feasibility is preserved. Thus the primal
994 optimum is unchanged if one restricts to the compact set

$$\{w \in [0,1]^k : Aw \geq \mathbf{1}\}.$$

995   By continuity of the objective, the minimum is attained.

996   The dual problem (I.7) is feasible, since $y = 0$ is feasible. It is bounded above because, for any
997   dual-feasible y,

$$\sum_{S \in \mathcal{S}} y_S \le \sum_{i=1}^{k} \sum_{S \ni i} y_S \le k.$$

998   Thus the dual optimum is finite; by standard finite-dimensional LP theory it is also attained.

999   Since (I.6) and (I.7) are a primal-dual pair of feasible finite linear programs with finite optimal
1000 values, strong duality applies, yielding

$$\tau^*(\mathcal{S}) = \nu^*(\mathcal{S}).$$

1001 $\square$

1002 The dual variables y_S assign load to histories/support sets, subject to the constraint that the total
1003 load incident to any pure controller i is at most 1. In this sense, $\nu^*(\mathcal{S})$ measures the maximum
1004 normalized amount of *history obstruction* that can be packed into the support family.

1005 **Corollary I.8** (Complementary slackness interpretation). *Let $w^\star$ and $y^\star$ be primal and dual optimal*
1006 *solutions. Then:*

1007   *1. if $w_i^\star > 0$, then*

$$\sum_{S \ni i} y_S^\star = 1;$$

1008   *2. if $y_S^\star > 0$, then*

$$\sum_{i \in S} w_i^\star = 1.$$

1009 *Proof.* This is the standard complementary-slackness condition for the primal-dual pair (I.6)–(I.7).
1010 $\square$

I.4   Projective-plane example

1012 The projective-plane family used in Section 5 also admits an exact fractional analysis.

1013 **Proposition I.9** (Fractional value for projective-plane line hypergraphs). *Let $\Pi_q = (P_q, \mathcal{L}_q, \in)$ be*
1014 *a projective plane of order q, and let*

$$\mathscr{L}_q = \{L \subseteq P_q : L \in \mathcal{L}_q\}$$

1015 *be its line hypergraph. Then*

$$\tau^*(\mathscr{L}_q) = \nu^*(\mathscr{L}_q) = \frac{q^2 + q + 1}{q + 1}. \tag{I.9}$$

1016 *Proof.* For the primal problem, assign uniform weight

$$w_p := \frac{1}{q + 1} \qquad \forall p \in P_q.$$

1017 Since each line $L \in \mathscr{L}_q$ contains exactly $q + 1$ points,

$$\sum_{p \in L} w_p = 1,$$

1018 so w is primal-feasible. Its objective value is

$$\sum_{p \in P_q} w_p = \frac{|P_q|}{q+1} = \frac{q^2 + q + 1}{q+1}.$$

1019 Hence

$$\tau^*(\mathscr{L}_q) \le \frac{q^2 + q + 1}{q+1}.$$

1020 For the dual problem, assign uniform weight

$$y_L := \frac{1}{q+1} \qquad \forall L \in \mathcal{L}_q.$$

1021 Each point $p \in P_q$ lies on exactly $q + 1$ lines, so

$$\sum_{L \ni p} y_L = 1.$$

1022 Thus y is dual-feasible. Its objective value is

$$\sum_{L \in \mathcal{L}_q} y_L = \frac{|\mathcal{L}_q|}{q+1} = \frac{q^2 + q + 1}{q+1}.$$

1023 Hence

$$\nu^*(\mathscr{L}_q) \ge \frac{q^2 + q + 1}{q+1}.$$

1024 Now apply Theorem I.7:

$$\tau^*(\mathscr{L}_q) = \nu^*(\mathscr{L}_q).$$

1025 The upper and lower bounds above therefore coincide, proving (I.9). $\qquad\square$

1026 **Corollary I.10** (Fractional value of the projective-plane realization)**.** *In the exact-rationality in-*
1027 *stance of Section 5 generated from the line hypergraph of a projective plane of order q,*

$$\tau^*(\{S_h : h \in \mathcal{H}\}) = \frac{q^2 + q + 1}{q+1}, \qquad m_{\Phi_{\mathrm{ext}}}^{\mathrm{pc}}(0) = q + 1.$$

1028 *Proof.* The first identity is Proposition I.9; the second is Lemma 5.2 together with Corollary 4.2. $\quad\square$

1029 *Remark* I.11 (Integrality gap). For the projective-plane family,

$$\frac{m_{\Phi_{\mathrm{ext}}}^{\mathrm{pc}}(0)}{\tau^*(\mathscr{L}_q)} = \frac{q+1}{(q^2 + q + 1)/(q+1)} = \frac{(q+1)^2}{q^2 + q + 1} = 1 + \frac{q}{q^2 + q + 1}.$$

1030 Thus the projective-plane obstruction separating exact memory from pairwise compatibility has only
1031 a modest fractional gap: the LP relaxation is strictly smaller than the exact complexity, but asymp-
1032 totically close to it.

1033 *Remark* I.12 (Total unimodularity and exactness). The gap between $\tau^*(\mathcal{S})$ and $\tau(\mathcal{S})$ is precisely an
1034 integrality phenomenon of the incidence matrix A. In particular, if the system

$$Aw \ge \mathbf{1}, \qquad w \ge 0$$

1035 defines an integral polyhedron—for example, if the relevant incidence matrix is totally unimodular
1036 after sign normalization—then the fractional and exact retentive complexities coincide. We do not
1037 pursue such structural conditions here, but they suggest a natural bridge from bounded-memory RL
1038 to polyhedral combinatorics.

J Small Examples and Counterexamples

1040 This section collects small finite examples that isolate the structural phenomena proved in the main
1041 text. The goal is not to add new theory, but to make the geometry of exact and approximate retentive
1042 complexity completely explicit in low-dimensional instances.

J.1 A minimal pairwise obstruction

1044 We begin with the smallest nontrivial support family exhibiting failure of pairwise determination.

Definition J.1 (Triangle support family). Let

$$\mathcal{S}_\triangle := \{\{1,2\},\{1,3\},\{2,3\}\} \subseteq 2^{[3]} \setminus \{\varnothing\}.$$

Proposition J.2 (Minimal complete-graph obstruction). *There exists a one-step, fully observable mixed-controller instance with controller dictionary $\{e_1, e_2, e_3\}$ such that:*

1. the family of exact rational-support sets is

$$\mathcal{S}_\triangle = \{\{1,2\},\{1,3\},\{2,3\}\};$$

2. the pairwise compatibility graph is the complete graph K_3;

3. the exact retentive complexity is

$$m^{\mathrm{pc}}_{\Phi_{\mathrm{ext}}}(0) = 2.$$

Proof. By Theorem 4.3, there exists a one-step instance realizing the support family $\mathcal{S}_\triangle$. Let the attainable histories be h_{12}, h_{13}, h_{23}, with

$$S_{h_{12}} = \{1,2\}, \qquad S_{h_{13}} = \{1,3\}, \qquad S_{h_{23}} = \{2,3\}.$$

Every two sets in $\mathcal{S}_\triangle$ intersect, so

$$S_{h_{12}} \cap S_{h_{13}} = \{1\} \neq \varnothing, \quad S_{h_{12}} \cap S_{h_{23}} = \{2\} \neq \varnothing, \quad S_{h_{13}} \cap S_{h_{23}} = \{3\} \neq \varnothing.$$

Hence the pairwise compatibility graph is K_3.

To compute the exact retentive complexity, apply Corollary 4.2. Thus it suffices to compute $\tau(\mathcal{S}_\triangle)$. No singleton hits all three sets, because

$$\{1\} \cap \{2,3\} = \varnothing, \qquad \{2\} \cap \{1,3\} = \varnothing, \qquad \{3\} \cap \{1,2\} = \varnothing.$$

Hence $\tau(\mathcal{S}_\triangle) \geq 2$. On the other hand, $\{1,2\}$ hits all three sets, so $\tau(\mathcal{S}_\triangle) \leq 2$. Therefore

$$\tau(\mathcal{S}_\triangle) = 2,$$

and so

$$m^{\mathrm{pc}}_{\Phi_{\mathrm{ext}}}(0) = 2.$$

$\square$

Remark J.3. Proposition J.2 is the smallest complete-graph obstruction in the sense that it uses only three histories and three pure controllers. It is the finite toy version of the projective-plane separation in Section 5.

1063 ## J.2 A tiny integrality gap

1064 The same triangle family also exhibits the smallest nontrivial fractional gap.

1065 **Proposition J.4** (Fractional value of the triangle family). *For the support family $\mathcal{S}_\triangle$,*

$$\tau(\mathcal{S}_\triangle) = 2, \qquad \tau^*(\mathcal{S}_\triangle) = \frac{3}{2}.$$

1066 *Proof.* The exact value $\tau(\mathcal{S}_\triangle) = 2$ was proved in Proposition J.2.

1067 For the fractional value, consider the primal LP

$$\tau^*(\mathcal{S}_\triangle) = \min\left\{ w_1 + w_2 + w_3 : \begin{array}{c} w_1 + w_2 \geq 1, \\ w_1 + w_3 \geq 1, \\ w_2 + w_3 \geq 1, \\ w_1, w_2, w_3 \geq 0 \end{array} \right\}.$$

1068 The uniform vector

$$w_1 = w_2 = w_3 = \frac{1}{2}$$

1069 is feasible, so

$$\tau^*(\mathcal{S}_\triangle) \leq \frac{3}{2}.$$

1070 Conversely, summing the three covering constraints gives

$$2(w_1 + w_2 + w_3) \geq 3,$$

1071 and therefore

$$w_1 + w_2 + w_3 \geq \frac{3}{2}$$

1072 for every feasible w. Hence

$$\tau^*(\mathcal{S}_\triangle) = \frac{3}{2}.$$

1073 $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\quad\square$

1074 *Remark* J.5. The triangle family therefore has integrality gap

$$\frac{\tau(\mathcal{S}_\triangle)}{\tau^*(\mathcal{S}_\triangle)} = \frac{2}{3/2} = \frac{4}{3}.$$

1075 This is the smallest explicit example in the paper where exact memory exceeds its fractional relax-
1076 ation.

1077 ## J.3 A sharp threshold for one-state memory

1078 We now turn to positive tolerance $\varepsilon > 0$. The next two-history example shows that approximate
1079 retentive complexity can exhibit a sharp phase transition.

1080 **Proposition J.6** (Sharp threshold for one-state approximate memory). *Consider the mixed-*
1081 *controller model with $k = 2$, controller simplex Δ_2, external deviations Φ_{ext}, and two histories*
1082 *h_1, h_2 with value vectors*

$$v(h_1) = (1,0), \qquad v(h_2) = (0,1).$$

1083 *Then, for every $\varepsilon \geq 0$,*

$$K_{h_1}^\varepsilon = \{x \in \Delta_2 : x_1 \geq 1 - \varepsilon\}, \qquad K_{h_2}^\varepsilon = \{x \in \Delta_2 : x_1 \leq \varepsilon\}.$$

1084 *Consequently,*

$$m_{\Phi_{\text{ext}}}^{\text{pc}}(\varepsilon) = \begin{cases} 2, & 0 \leq \varepsilon < \frac{1}{2}, \\ 1, & \varepsilon \geq \frac{1}{2}. \end{cases} \tag{J.1}$$

1085 *Proof.* By Theorem 4.1,

$$K_{h_1}^\varepsilon = \{x \in \Delta_2 : \langle (1,0), x \rangle \geq 1 - \varepsilon\} = \{x \in \Delta_2 : x_1 \geq 1 - \varepsilon\},$$

1086 and similarly

$$K_{h_2}^\varepsilon = \{x \in \Delta_2 : \langle (0,1), x \rangle \geq 1 - \varepsilon\} = \{x \in \Delta_2 : x_2 \geq 1 - \varepsilon\} = \{x \in \Delta_2 : x_1 \leq \varepsilon\}.$$

1087 Their intersection is nonempty iff there exists $x_1 \in [0,1]$ satisfying

$$1 - \varepsilon \leq x_1 \leq \varepsilon,$$

1088 which is equivalent to $\varepsilon \geq \frac{1}{2}$. By Corollary 6.1, this is equivalent to

$$m_{\Phi_{\text{ext}}}^{\text{pc}}(\varepsilon) = 1 \iff \varepsilon \geq \frac{1}{2}.$$

1089 It remains to show that, when $\varepsilon < \frac{1}{2}$, the exact value is 2. Since the total intersection is empty,
1090 one-state memory is impossible. On the other hand,

$$e_1 \in K_{h_1}^\varepsilon, \qquad e_2 \in K_{h_2}^\varepsilon \qquad \forall \varepsilon \geq 0,$$

1091 so the set $\{e_1, e_2\}$ is a transversal for $\{K_{h_1}^\varepsilon, K_{h_2}^\varepsilon\}$. By Theorem 3.1,

$$m_{\Phi_{\text{ext}}}^{\text{pc}}(\varepsilon) \leq 2 \qquad \forall \varepsilon \geq 0.$$

1092 Hence $m_{\Phi_{\text{ext}}}^{\text{pc}}(\varepsilon) = 2$ for $0 \leq \varepsilon < \frac{1}{2}$, proving (J.1). □

1093 *Remark* J.7. At the threshold $\varepsilon = \frac{1}{2}$, the total intersection is the singleton

$$K_{h_1}^{1/2} \cap K_{h_2}^{1/2} = \left\{ \left(\frac{1}{2}, \frac{1}{2} \right) \right\}.$$

1094 Thus the one-state feasible continuation appears through a single barycentric point of the simplex.

J.4 Mixed controllers can strictly help at positive tolerance

1096 The previous example also shows that mixed continuations are not merely a technical convenience:
1097 they can strictly reduce approximate memory complexity.

1098 **Definition J.8** (Pure-restricted approximate retentive complexity). Fix a finite-controller mixed-
1099 continuation instance with dictionary $\{e_1, \ldots, e_k\}$. Its *pure-restricted approximate retentive com-*
1100 *plexity* is

$$m_{\text{pure}}^{\text{pc}}(\varepsilon) := \inf \left\{ m \in \mathbb{N} : \begin{array}{l} \exists\, (\sigma, d) \text{ of size } m \text{ with } d : [m] \to \{e_1, \ldots, e_k\}, \\ d(\sigma(h)) \in K_h^\varepsilon \; \forall h \in \mathcal{H} \end{array} \right\}.$$

1101 **Proposition J.9** (Strict mixed-controller advantage). *In the two-history instance of Proposition J.6,*

$$m_{\Phi_{\text{ext}}}^{\text{pc}} \left(\frac{1}{2} \right) = 1, \qquad m_{\text{pure}}^{\text{pc}} \left(\frac{1}{2} \right) = 2.$$

1102 *Proof.* The first identity is Proposition J.6. Indeed, the mixed controller

$$x^\star = \left(\frac{1}{2}, \frac{1}{2} \right)$$

1103 belongs to both $K_{h_1}^{1/2}$ and $K_{h_2}^{1/2}$, so one-state memory is feasible.

1104   For the pure-restricted problem, note that

$$e_1 \notin K_{h_2}^{1/2}, \qquad e_2 \notin K_{h_1}^{1/2},$$

1105   because

$$K_{h_1}^{1/2} = \left\{ x \in \Delta_2 : x_1 \geq \frac{1}{2} \right\}, \qquad K_{h_2}^{1/2} = \left\{ x \in \Delta_2 : x_1 \leq \frac{1}{2} \right\}.$$

1106   Hence no single pure controller belongs to both low-regret sets, so $m_{\text{pure}}^{\text{pc}}(1/2) \neq 1$.

1107   On the other hand, the pure controllers e_1 and e_2 satisfy

$$e_1 \in K_{h_1}^{1/2}, \qquad e_2 \in K_{h_2}^{1/2},$$

1108   so a two-label pure architecture exists. Therefore

$$m_{\text{pure}}^{\text{pc}} \left(\frac{1}{2} \right) = 2.$$

1109 $\square$

1110   *Remark* J.10. At exact rationality $\varepsilon = 0$, the face-piercing reduction in Section 4 shows that mixed
1111   continuations do not reduce memory beyond what can already be achieved by selecting vertices
1112   of the simplex. The strict separation in Proposition J.9 is therefore a genuinely positive-tolerance
1113   phenomenon.

1114   **J.5   A small complete-graph example below the Helly threshold**

1115   The final example ties together the pairwise and Helly phenomena.

1116   **Proposition J.11** (Complete pairwise graph does not imply one-state memory). *For the triangle*
1117   *family $\mathcal{S}_\triangle$, every pair of exact low-regret sets intersects, but the total intersection is empty:*

$$\Delta(\{1,2\}) \cap \Delta(\{1,3\}) \neq \varnothing, \qquad \Delta(\{1,2\}) \cap \Delta(\{2,3\}) \neq \varnothing, \qquad \Delta(\{1,3\}) \cap \Delta(\{2,3\}) \neq \varnothing,$$

1118   *whereas*

$$\Delta(\{1,2\}) \cap \Delta(\{1,3\}) \cap \Delta(\{2,3\}) = \varnothing.$$

1119   *Proof.* The pairwise intersections are

$$\Delta(\{1,2\}) \cap \Delta(\{1,3\}) = \{e_1\},$$
$$\Delta(\{1,2\}) \cap \Delta(\{2,3\}) = \{e_2\},$$
$$\Delta(\{1,3\}) \cap \Delta(\{2,3\}) = \{e_3\}.$$

1120   all of which are nonempty.

1121   Now let $x \in \Delta(\{1,2\}) \cap \Delta(\{1,3\}) \cap \Delta(\{2,3\})$. Then simultaneously

$$\text{supp}(x) \subseteq \{1,2\}, \qquad \text{supp}(x) \subseteq \{1,3\}, \qquad \text{supp}(x) \subseteq \{2,3\}.$$

1122   Hence

$$\text{supp}(x) \subseteq \{1,2\} \cap \{1,3\} \cap \{2,3\} = \varnothing,$$

1123   which is impossible for a point $x \in \Delta_3$. Therefore the triple intersection is empty. $\square$

1124   *Remark* J.12. Proposition J.11 is consistent with Corollary 6.1: the simplex Δ_3 has affine dimension
1125   2, so one-state memory is controlled by intersections of subfamilies of size at most 3, not merely by
1126   pairwise intersections.