Score-based generative modeling secretly minimizes the Wasserstein distance

Abstract

Score-based generative models are shown to achieve remarkable empirical per-
formances in various applications such as image generation and audio synthesis.
However, a theoretical understanding of score-based diffusion models is still in-
complete. Recently, Song et al. showed that the training objective of score-based
generative models is equivalent to minimizing the Kullback-Leibler divergence of
the generated distribution from the data distribution. In this work, we show that
score-based models also minimize the Wasserstein distance between them under
suitable assumptions on the model. Specifically, we prove that the Wasserstein
distance is upper bounded by the square root of the objective function up to mul-
tiplicative constants and a fixed constant offset. Our proof is based on a novel
application of the theory of optimal transport, which can be of independent interest
to the society. Our numerical experiments support our findings. By analyzing our
upper bounds, we provide a few techniques to obtain tighter upper bounds.

1 Introduction

The idea of gradually perturbing the data distribution to define the forward process and learning
the reverse process has recently demonstrated remarkable performances in generative modeling.
Score-based generative models [33, 34, 35] and diffusion probabilistic models [30, 17] have been
used in numerous applications, including image generation [26, 12], and audio synthesis [9, 21].

Both model families have the following two processes. The given data distribution is incrementally
perturbed by adding small noise in the forward process. This forward process can be understood as
the standard diffusion process [35]. The reverse process requires the so-called *score function* [2], the
gradient of the log probability density function. To approximate the score function, it is common to
minimize a weighted combination of score matching losses. Once the score function is approximated,
one can generate samples from randomly generated noise by applying the learned reverse process.

Note that score-based models do not explicitly minimize any sort of similarity between *the model
distribution* (the distribution of the generated samples) and the data distribution. Therefore, it
is unclear whether or not scored-based generative models can minimize divergence (or distance)
between the model distribution and the data distribution. In very recent work [32], the authors show
that optimizing the score matching loss is indeed equivalent to minimizing an upper bound on the
Kullback–Leibler divergence (KL divergence). Therefore, it is expected that the model distribution is
similar (in the sense of KL divergence) to the data distribution if the score matching loss is small
enough.

A natural question follows – *do score-based models minimize any other divergence or distance
than the KL divergence?* At first glance, this seems unlikely. For instance, Generative Adversarial
Networks (GANs) [16] were shown to minimize different divergences depending on the choice
of the loss function. The original GAN loss function turns out to capture the Jensen–Shannon

divergence [16], while the W-GAN loss function captures the Wasserstein distance [3]. We refer the readers to [10] for more details.

On the other hand, a unique aspect of score-based models is that the score function is defined in the same space where data lies in. In stark contrast, all the other popular generative models such as GANs [16], VAEs [19], and Normalizing Flows [20] use loss functions that only care about likelihoods and do not directly operate in the knowledge of the underlying data space.

In this work, we answer this question by revealing the following property of score-based modeling:

> Score-based modeling minimizes the Wasserstein distance.

We note that this is quite surprising as there is no direct relationship between the KL divergence and the Wasserstein distance, in general. Therefore, this finding is not implied by the aforementioned finding.

(a) 1 cluster in 1D   (b) 2 clusters in 1D   (c) 1 cluster in 2D   (d) 4 clusters in 2D

Figure 1: Training curves of loss function J_{DSM} (eq. (14)), Wasserstein distance and JS divergence during training. The training data is sampled from mixture of 1D and 2D Gaussian clusters. (a): samples from one cluster $\mathcal{N}(0, 0.1)$. (b): samples from two clusters $\mathcal{N}(0, 0.1)$ and $\mathcal{N}(0.5, 0.1)$ with equal weights. (c): samples from one cluster $\mathcal{N}(\mathbf{0}, 0.01\mathbf{I})$. (d): samples from four clusters $\mathcal{N}((\pm 0.5, \pm 0.5)^\top, 0.01\mathbf{I})$ with equal weights.

Using the theory of optimal transport, we investigate the evolution of data distribution based on Partial differential equations (PDEs). This allows us to estimate the Wasserstein distance between the given data distribution and the distribution of the generated samples. Our work theoretically guarantees that the score-based generative models approximate data distributions in the space of probability measures equipped with the Wasserstein distance. Our main contributions and findings are summarized as follows:

- (Theorem 1 & Corollary 1) Using the theory of optimal transport, we prove that the Wasserstein distance between the data distribution and the model distribution is bounded from above. The upper bound is given as the square root of the score-matching loss J_{SM} up to multiplicative constants and a fixed constant offset. The main idea is to estimate the time-derivative of the Wasserstein metric by using the form of the continuity equations.

- (Theorem 2) We prove $J_{SM} \leq J_{DSM}$ (as defined in eq. (5) and eq. (14), respectively) under suitable assumptions, which gives us a more practical upper bound since J_{DSM} is much easier to estimate in practice and is widely used as a loss function to train score-based models.

- (Theorem 3) We analyze how the model distribution changes if the data distribution is perturbed initially. This guarantees that the score based-model is robust under noise.

- (Section 4) Our numerical experiments support the upper bound of our analysis. We also explore the effect on the weight decay coefficient in training and different choices of the number of total timesteps T.

Notations: $x = (x_1, x_2, \cdots, x_d)^\top \in \mathbb{R}^d$ and $t \in [0, T]$ denote the spatial variable and the time variable, respectively. Here, $\mathbb{R}^d$ is the d-dimensional Euclidean space and $T > 0$ is the time horizon. The gradient of a real-valued function ρ with respect to the spatial variable and the time-derivative of ρ are denoted by $\nabla \rho = \left(\frac{\partial \rho}{\partial x_1}, \frac{\partial \rho}{\partial x_2}, \cdots, \frac{\partial \rho}{\partial x_d} \right)^\top$, and $\partial_t \rho$, respectively. The Laplacian of ρ is given as $\Delta \rho = \nabla \cdot (\nabla \rho)$. Here, $\nabla \cdot F = \frac{\partial F_1}{\partial x_1} + \frac{\partial F_2}{\partial x_2} + \cdots + \frac{\partial F_d}{\partial x_d}$ denotes the divergence of $F = (F_1, F_2, \cdots, F_d)$ with respect to the spatial variable x. The Hessian of ρ is a $d \times d$ matrix

given by $(D^2\rho)_{ij} = \frac{\partial^2 \rho}{\partial x_i \partial x_j}$ for $1 \leq i, j \leq d$. We indicate Gaussian distribution with mean μ and covariance Σ by $\mathcal{N}(\mu, \Sigma)$, and L_2 norm by $\|\cdot\|$. We denote the data distribution and the distribution of the generated samples by p_0 and q_0.

2 Preliminaries and related works

In this section, we analyze the forward process and the reverse process in the score-based model through Partial differential equations (PDEs). In particular, we investigate the time evolution of marginal distributions through the two processes.

2.1 The forward process

As in [35], the forward process can be modeled by the standard diffusion process $\{x(t)\}_{t=0}^T$ for $t \in [0, T]$ such that $x(0) \sim p_0$ and

$$dx = f(x, t)dt + g(t)d\mathbf{w}. \tag{1}$$

Here, $\mathbf{w}$ is the standard Wiener process, $f : \mathbb{R}^d \times [0, T] \to \mathbb{R}^d$ is the drift coefficient of $x(t)$, and $g : [0, T] \to \mathbb{R}$ is the diffusion coefficient of $x(t)$.

Let $p_t = p(\cdot, t)$ be a probability density function of the random variable $x(t)$ on $\mathbb{R}^d$ at time $t \in [0, T]$. Then, the function $p : \mathbb{R}^d \times [0, T] \to [0, +\infty)$ satisfies the following partial differential equations (PDEs), namely the Fokker–Planck equation or the forward Kolmogorov equation as in [5, Theorem 2.2]:

$$\partial_t p + \nabla \cdot (pf) - \frac{g^2}{2}\Delta p = 0, \;\; p(\cdot, 0) = p_0. \tag{2}$$

where the pdf p_0 represents the data distribution. Under suitable assumptions on f and g, a solution to the above initial value problem uniquely exists. We refer the readers to [6] for more details.

2.2 Understanding the reverse process through PDEs

To generate samples from $t = T$ to 0, the score-based model uses the reverse-time SDE [2]:

$$dx = (f(x, t) - g(t)^2 \nabla \log p)dt + g(t)d\bar{\mathbf{w}} \tag{3}$$

Here, $\bar{\mathbf{w}}$ is a standard Wiener process when time flows backward from $t = T$ to $t = 0$.

As in the previous section, for given $x(T) \sim p_T$, a probability density function of the random variable $x(t)$ from the above process (3) satisfies the final value problem:

$$\partial_t p + \nabla \cdot \left(p\left(f - g^2 \nabla \log p\right)\right) + \frac{g^2}{2}\Delta p = 0, \;\; p(\cdot, T) = p_T. \tag{4}$$

Using this simple identity, $\Delta p = \nabla \cdot (\nabla p) = \nabla \cdot \left(p\frac{\nabla p}{p}\right) = \nabla \cdot (p\nabla(\log p))$, it can be seen that the differential equations in (2) and (4) are indeed identical.

2.3 The objective function

The score of the probability distribution, $\nabla \log p_t$, is approximated by a function $s_\theta : \mathbb{R}^d \times [0, T] \to \mathbb{R}^d$. For instance, in [32], the authors consider the weighted mean squared error (MSE):

$$J_{SM}(\theta; \lambda) := \frac{1}{2} \int_0^T \lambda(t)\mathbb{E}_{p_t}\left[\|\nabla \log p_t(x) - s_\theta(x, t)\|^2\right] dt \tag{5}$$

where $\lambda : [0, T] \to (0, \infty)$ is a positive weighting function.

Replacing $\nabla \log p$ in (4) with s_θ, we deduce a new reverse process. For given noise distribution q_T, a probability density function $q_t = q(\cdot, t)$ at time $t \in [0, T]$ solves

$$\partial_t q + \nabla \cdot \left(q\left(f - g^2 s_\theta\right)\right) + \frac{g^2}{2}\Delta q = 0, \;\; q(T, \cdot) = q_T. \tag{6}$$

Here, we introduced a new function $q : \mathbb{R}^d \times [0, T] \to [0, +\infty)$ to illustrate the reverse process.

2.4 Related works

Optimal transport The Wasserstein distance has a solid connection to the Fokker–Planck equation. In particular, the Fokker–Planck equation can be understood as the gradient flow in the space of probability measures equipped with the 2-Wasserstein distance [1, 18, 27]. For two probability measures μ and ν on $\mathbb{R}^d$, we define the 2-Wasserstein or Monge-Kantorovich distance as

$$W_2(\mu, \nu) := \inf \left\{ \int_{\mathbb{R}^d \times \mathbb{R}^d} \|x - y\|^2 d\gamma : \gamma \in \Pi(\mu, \nu) \right\}^{\frac{1}{2}}. \tag{7}$$

Here, $\Pi(\mu, \nu)$ is the set of all couplings π of μ and ν where a Borel probability measure π on $\mathbb{R}^d \times \mathbb{R}^d$ satisfies $\pi(A \times \mathbb{R}^d) = \mu(A)$ and $\pi(\mathbb{R}^d \times B) = \nu(B)$ for all measurable subsets $A, B \subset \mathbb{R}^d$.

Under suitable assumptions on the drift coefficient and the diffusion coefficient, a solution to the Fokker–Planck equation satisfies the so-called Wasserstein contraction property [7, 8, 29], which plays an important role in our analysis. More discussions can be found in Section 3.3 and Section 5.

Score matching and diffusion probabilistic models Score matching models [33, 35] use the score function to reverse the noise corruption in a continuous time range by solving stochastic differential equations (SDEs). Diffusion probabilistic models [30, 17] adopt a similar idea of modeling tractable denoising process to reverse each discrete step of noise corruption with probabilistic models. In fact, [35] show that diffusion models are score matching models with certain forms of SDE, which unifies the whole framework. Diffusion models are gaining increasing attention: recent advances [31, 26, 12] show that diffusion models can generate even better images than GANs while enjoying a more stable training process, which encourages it to be widely applied in various tasks, such as text-to-image and image editing [25, 24], audio generation [9, 21], etc.

3 Approximation in the Wasserstein space

In this section, we estimate the Wasserstein distance between a given data distribution p_0 and the marginal of the generated samples q_0 obtained by the score-based model. The proofs of the main results are deferred to Appendix A and Appendix D. The key ideas are illustrated in Section 5.

3.1 The main results

We present our main results under the following assumptions. **(A1)** The drift coefficient $f : \mathbb{R}^d \times [0, T] \to \mathbb{R}^d$ is Lipschitz continuous in the space variable x: there exists a positive constant $L_f(t) \in (0, \infty)$, depending on $t \in [0, T]$, such that for all $x, y \in \mathbb{R}^d$

$$\|f(x, t) - f(y, t)\| \leq L_f(t)\|x - y\|. \tag{8}$$

(A2) $s_\theta : \mathbb{R}^d \times [0, T] \to \mathbb{R}^d$ satisfies *the one-sided Lipschitz condition* [13, Definition 2.1]: there exists a constant $L_s(t) \in \mathbb{R}$, depending on $t \in [0, T]$, satisfying

$$(s_\theta(x, t) - s_\theta(y, t)) \cdot (x - y) \leq L_s(t)\|x - y\|^2 \tag{9}$$

for all $x, y \in \mathbb{R}^d$. The usual Lipschitz constant of s_θ satisfies the above inequality (9). Unlike the usual Lipschitz constant, *the one-sided Lipschitz constant $L_s(t)$ is not necessarily nonnegative* (see Appendix B). More technical assumptions can be found in Appendix A. We now present a general result.

Theorem 1. *Let p_0 and q_0 be the data distribution and a pdf obtained by the score-based model as in Section 2. Then the Wasserstein distance between p_0 and q_0 is bounded as follows:*

$$W_2(p_0, q_0) \leq \int_0^T g^2(t) I(t) \mathbb{E}_{p_t} \left[\|\nabla \log p_t(x) - s_\theta(x, t)\|^2 \right]^{\frac{1}{2}} dt + I(T) W_2(p_T, q_T). \tag{10}$$

Here, a function $I : [0, T] \to (0, +\infty)$ defined by

$$I(t) := \exp \left(\int_0^t L_f(r) + L_s(r) g^2(r) dr \right) \tag{11}$$

142 is an integrating factor of the differential inequality (22) given later in Proposition 1. The main idea
143 of the proofs is to estimate the derivatives of the Wasserstein distance along with the time evolution,
144 which is illustrated in Section 5.

145 The first term on the right-hand side of (10) naturally arises from the estimation of $W_2(p_0, q_0)$.
146 It is similar to a loss function J_{SM} commonly used in the score-based models. Applying the
147 Cauchy-Schwarz inequality, we obtain the upper bound in terms of the weighted MSE J_{SM}.

148 **Corollary 1.** *Let p_0 and q_0 be given in Theorem 1. If $\lambda(t) = g(t)^2$ for all $t \in [0, T]$, then the*
149 *following inequality holds:*

$$W_2(p_0, q_0) \le \sqrt{2 \left(\int_0^T g^2(t) I(t)^2 dt \right) J_{SM}} + I(T) W_2(p_T, q_T). \tag{12}$$

150 We defer the proofs of Theorem 1, and Corollary 1 to Appendix A.

151 **Remark 1.** *From the above result, the Wasserstein distance converges to zero as the loss function*
152 *J_{SM} goes to zero if $p_T = q_T$. However, due to the tension between $I(T)$ and $W_2(p_T, q_T)$, a general*
153 *convergence result remains open (see Section 6 and Appendix H).*

154 **Remark 2.** *If $g^4 I^2 \lambda^{-1}$ is integrable in $[0, T]$, the similar inequality as in (32) holds for other choices*
155 *of λ. See Corollary 4 in Appendix A.*

156 **Remark 3.** *Note that the above upper bound depends on the time horizon T. Numerical experiments*
157 *for different choices of T are provided in Section 4.3.2.*

158 We further investigate the offset $I(T) W_2(p_T, q_T)$ for Denoising Diffusion Probabilistic Models [17].

159 **Corollary 2.** *Under the same setting as in Corollary 1, let $f(t) = -\frac{\beta(t)}{2} x$, $g(t) = \sigma \sqrt{\beta(t)}$, and*
160 *$q_T = \phi \sim \mathcal{N}(0, \sigma^2 I)$. Then, we have*

$$W_2(p_0, q_0) \le \sqrt{2 \left(\int_0^T g^2(t) I(t)^2 dt \right) J_{SM}} + \exp \left(\int_0^T \beta(r) L_s(r) dr \right) W_2(p_0, \phi). \tag{13}$$

161 *Here, $\sigma > 0$, and $\beta : [0, T] \to (0, +\infty)$ is a positive function.*

162 The proof of Corollary 2 and the analysis of L_s can be found in Appendix H. As the arguments are
163 technical, we provide a formal analysis below.

164 **Remark 4.** *If s_θ is close enough to $\nabla \log p_t(x)$, then we have $L_s(t) \sim (-\sigma^{-2} + \|D^2(\log h_t)\|_\infty)$.*
165 *Here, $h_t := p_t/\phi$ exponentially converges to 1 as $t \to \infty$ (see Lemma 4). Thus, $L_s(t)$ converges to*
166 *$-\sigma^{-2}$, as observed in our experiments with $\sigma = 1$ in Fig. 4a. Consequently, the second term in (13)*
167 *decays as T increases (see Fig. 4b).*

3.2 The relation between J_{SM} and J_{DSM}

169 The above results give the guarantee that J_{SM} can upper-bound Wasserstein distance. However, J_{SM}
170 is generally intractable due to the unknown quantity $\nabla_x \log p_t(x)$ [32], and is rarely used in practice.
171 Alternatively, there is a conditional version of the score-matching loss J_{DSM}, which is equivalent
172 to J_{SM} up to a constant, as shown in [36]. J_{DSM} is also widely used as a loss function to train
173 score-based models:

$$J_{DSM}(\theta, \lambda) := \frac{1}{2} \int_0^T \lambda(t) \, \mathbb{E}_{p_0(x(0))} \, \mathbb{E}_{p_{0t}(x|x(0))} [\|s_\theta(x, t) - \nabla_x \log p_{0t}(x|x(0))\|^2] dt, \tag{14}$$

174 where p_{0t} indicates the conditional probability of x at step t given $x(0)$. Now we present our new
175 result, which shows that J_{DSM} is also an upper bound of J_{SM} under the certain assumption.

176 **Theorem 2.** *If p_{0t} satisfies*

$$\text{Var}[\mathbb{E}[(\nabla_x \log p_{0t}(x|x(0)))^\top | x(0)]] = 0, \tag{15}$$

177 *then we have*

$$J_{SM} \le J_{DSM} \quad and \quad W_2(p_0, q_0) \le \sqrt{2 \left(\int_0^T g^2(t) I(t)^2 dt \right) J_{DSM}} + I(T) W_2(p_T, q_T). \tag{16}$$

178 It is worth noting that Denoising Diffusion Probabilistic Models (DDPM) [17] satisfy eq. (15).

179 **Corollary 3.** *Let $p(x(t)|x(t-1)) = \mathcal{N}(\sqrt{1-\beta_t}x(t-1), \beta_t I)$ as in Denoising Diffusion Prob-*
180 *abilistic Models (DDPM) [17], which corresponds to $f(x,t) = -\frac{1}{2}\beta_t x$, and $g(t) = \sqrt{\beta_t}$. Here,*
181 *$\beta_t = \beta(t) > 0$ is the noise schedule for all t, and let $\lambda(t) = g^2(t)$. Then eq. (16) holds.*

182 The proofs of Theorem 2 and Corollary 3 are in Appendix D. We also provide numerical results
183 which show $J_{SM} \leq J_{DSM}$ for DDPM in Appendix E.

184 **Remark 5.** *Using Theorem 2 and Corollary 3, we can present an upper bound of the Wasserstein*
185 *distance in terms of J_{DSM}, where both $W_2(p_0, q_0)$ and J_{DSM} are tractable. This enables us to*
186 *empirically verify this upper bound via DDPM in Section 4.*

3.3 Perturbation and sensitivity analysis

188 Let $\tilde{p}_0$ be a small perturbation of the original data distribution p_0. For a given corrupted distribution
189 $\tilde{p}_0$, denote the probability density functions in the forward process and the reverse process by $\tilde{p}_t$ and
190 $\tilde{q}_t$, respectively. We investigate how this perturbation changes the score-based model.

191 It is well known from the Wasserstein contraction property [8] that the Wasserstein distance between
192 two probability distributions in the forward process can be estimated as follows: for all $t \in [0, T]$

$$W_2(p_t, \tilde{p}_t) \leq \exp\left(\int_0^t L_f(r)dr\right) W_2(p_0, \tilde{p}_0) \tag{17}$$

193 In the reverse process, the score function in the drift coefficients may vary, and thus the above
194 inequality does not directly give us the upper bound on $W_2(q_0, \tilde{q}_0)$. Instead, our analysis in the
195 previous sections allows us to estimate $W_2(q_0, \tilde{q}_0)$.

196 It is worth noting that the Wasserstein distance satisfies the triangle inequality, unlike other statistical
197 measures, including the Kullback–Leibler (KL) divergence.

198 **Theorem 3.** *Let $\tilde{p}_0$ be the corrupted data distribution and $\tilde{q}_0$ be a probability density function*
199 *obtained by the score-based model starting from $\tilde{p}_0$. It holds that*

$$W_2(q_0, \tilde{q}_0) \leq W_2(p_0, \tilde{p}_0) + \sqrt{2\left(\int_0^T g^2(t)I(t)^2 dt\right)J_{SM}} + \sqrt{2\left(\int_0^T g^2(t)\tilde{I}(t)^2 dt\right)\tilde{J}_{SM}}$$
$$+ \tilde{I}(T)W_2(p_T, q_T) + \tilde{I}(T)W_2(\tilde{p}_T, \tilde{q}_T). \tag{18}$$

200 Here, $\tilde{J}_{SM}$ and $\tilde{I}$ denote the weighted MSE and the integrating factor from the perturbed model.

201 **Remark 6.** *The Wasserstein distance between the original data distribution p_0 and the model*
202 *distribution $\tilde{q}_0$ starting from the corrupted data $\tilde{p}_0$ can also be estimated similarly.*

4 Experiments

204 Here we present experiments on simulation datasets to verify the upper bound in eq. (16).

4.1 Experiment settings

206 **Datasets** Here we adopt three 2D datasets for simulation: One cluster Gaussian $\mathcal{N}(\mathbf{0}, 0.1\mathbf{I})$, two
207 moons in [28], and four clusters Gaussian mixture $\mathcal{N}((\pm0.5, \pm0.5)^\top, 0.01\mathbf{I})$ with equal weights for
208 each cluster. Visualization for these datasets is in Fig 2.

209 For forward dynamics, we use discrete timesteps with $T = 10$. For $t \in [1, 10]$, $p(x(t)|x(t-1)) =$
210 $\mathcal{N}(\sqrt{1-\beta_t}x(t-1), \beta_t I)$ as in DDPM[17], i.e., $f(x,t) = -\frac{1}{2}\beta_t x$, and $g(t) = \sqrt{\beta_t}$. For β_t, we
211 adopt a sigmoid schedule from $\beta_1 = 10^{-5}$ to $\beta_T = 10^{-2}$.

212 **Training and evaluation** We use a 3-layer neural network as the score matching model, with ReLU
213 nonlinearity and skip-connection at the final output. Each layer is composed of a linear layer with 128
214 hidden neurons, and an embedding layer for 10 timesteps. For optimizer, we use AdamW [22] with

215 learning rate = 0.001 and weight decay coefficient 0.01. To evaluate Wasserstein distance, we use
216 POT [15] to compute $W_2(p_0, q_0)$ at the end of each training epoch. To avoid the term of $W_2(p_T, q_T)$
217 in eq. (16) for verification, we first perform forward pass on sampled $x(0)$ to get $x(T)$, and perform
218 data generation on the same $x(T)$ to fix $q_T = p_T$ and $W_2(p_T, q_T) = 0$. For loss function, we use
219 J_{DSM} with $\lambda(t) = g^2(t)$ and batch size = 512. To get both lower and higher losses to verify our
220 bound, we first maximize the loss for 15 epochs, and then minimize the loss till convergence.

4.2 Results

222 To verify the upper bound in eq. (16), we take log on each side of the equation with $W_2(p_T, q_T) = 0$
223 and get:

$$\log W_2(p_0, q_0) \leq \underbrace{\frac{1}{2} \log J_{DSM}}_{\text{slope}} + \underbrace{\frac{1}{2} \log \left(\int_0^T g^2(t) I(t)^2 dt \right)}_{\text{intercept}} \tag{19}$$

224 We present a log-log plot to verify the slope and the intercept in the upper bound, respectively: see
225 Fig 2. The log-log plots are obtained by evaluating Wasserstein distance and the loss function during
226 training. The red dots are the sampled data from each dataset, the green dots are generated data after
227 loss maximization, and the purple dots are the generated data sampled at convergence. From the
228 log-log plots, we can observe that the theoretical line (the black line) is a valid upper bound of the
229 empirical one (the blue dots).

Figure 2: Log-log plots of the loss function and Wasserstein distance, with corresponding generated data at the start of the loss minimization and at convergence for each dataset. In the log-log plots, the darkness of the datapoints corresponds to training epochs: the darkest ones are at convergence.

4.3 Exploring the upper bound

231 Here we explore different factors that can affect the tightness of the theoretical upper bound. As
232 shown in Fig 2, the gap between the theoretical bound and the empirical one mainly results from the
233 intercept in eq. (19). Recall our intercept is $\frac{1}{2} \log \left(\int_0^T g^2(t) I(t)^2 dt \right)$ for $I(t)$ given in (11).

4.3.1 Effect of weight regularization

235 Since $g(t)$ and L_f are fixed, only the one-sided Lipchitz constant $L_s(r)$ of the model s_θ is subject to
236 the training process, which is estimated via grid search in our experiments.

237 Notice that the loss function J_{DSM} aims to minimize the distance between $s_\theta(x, t)$ and
238 $\nabla_x \log p_{0t}(x|x(0))$. Thus, L_s at convergence is determined by dynamics $\nabla_x \log p_{0t}(x|x(0))$. If
239 the one-sided Lipschitz constant for $\nabla_x \log p_{0t}(x|x(0))$ is small, then the intercept would be ex-
240 pected to be small (see example in Fig 2a of Gaussian cluster with moderately high variance). If
241 the one-sided Lipschitz constant for $\nabla_x \log p_{0t}(x|x(0))$ is large, to minimize the loss, the intercept
242 would also be large (see example in Fig 2c of Gaussian clusters with lower variance). However, it can
243 be possible to make the upper bound tighter while sacrificing the training loss by constraining L_s.

244 To constrain L_s, we adopt spectral normalization (as used in [4] to regularize Lipschitz constant),
245 weight clipping (used in [3] to regularize Lipschitz constant), and simple weight decay. As we can
246 observe in Fig. 3, loss at convergence will increase when we impose very strong regularization on
247 the weights, while the intercept decreases. Importantly, the intercept approaches the empirical one
248 without not exceeding it. See more examples of weight decay in Appendix G, where we also present
249 the log-log training plots for weight decay $= 0.01, 0.1, 0.5, 1, 5$ respectively.

Figure 3: Effect of weight clipping and spectral normalization.

4.3.2 Effect of the number of total timesteps

251 In our Remark 4, we argue that the offset in our upper bound would decay to 0 when choosing
252 appropriate SDE and sufficiently large T. Here we provide empirical evidence for our claim without
253 fixing $p_T = q_T$. We use the four clusters Gaussian dataset and fix β to be sigmoid schedule between
254 $[10^{-5}, 10^{-2}]$ for any T. In Fig. 4, we plot $L_s(t)$ for $T = 100$ to show that $L_s(t)$ would converge
255 to -1 when $t \to T$. As a result, since $W_2(p_T, q_T)$ decays exponentially (see eq. 48), we also show
256 that as T increases, our offset $I(T)W_2(p_T, q_T)$ would also decay in a rate that is approximately
257 exponential when T is sufficiently large, as the model converges to the real data distribution since
258 $W_2(p_0, q_0) \approx 0$.

5 The main idea of the proofs

260 We investigate the time evolution of probability measures arising in the score-based models. Using
261 the optimal transport theory, we differentiate the Wasserstein distances, $W_2(p_t, q_t)$ with respect to
262 time. The main idea is to express these equations (2) and (6) in the form of the continuity equation:

$$\partial_t \rho + \nabla \cdot (\rho v) = 0 \tag{20}$$

263 where v is a so-called velocity field. Let us denote the velocity fields of p, and q by

$$v[p] = f - \frac{1}{2}g^2 \nabla \log p, \text{ and } v[q] = \left(f - g^2 s_\theta\right) + \frac{1}{2}g^2 \nabla \log q.$$

264 Then, the equations (2), and (6) can be written in the form of (20) with $v[p]$ and $v[q]$, respectively.

265 Using these vector fields, we compute the time derivative of the Wasserstein distance between two
266 evolving probability measures. For velocity fields $v[\rho^{(i)}]$ depending on $\rho^{(i)}$, let $\rho^{(i)}$ for $i = 1, 2$ be
267 two solutions of the continuity equations $\partial_t \rho^{(i)} + \nabla \cdot (\rho^{(i)} v[\rho^{(i)}]) = 0$. Then, the relation between
268 the Wasserstein distance and velocity fields arises from the theory of optimal transport:

$$\frac{1}{2}\frac{d}{dt}W_2^2(\rho_t^{(1)}, \rho_t^{(2)}) = \mathbb{E}_{\pi_t}\left[(x - y) \cdot (v[\rho_t^{(1)}](x) - v[\rho_t^{(2)}](y))\right] \tag{21}$$

(a) $L_s(t)$ for $T = 100$. We can verify that when T is sufficiently large, $L_s(T) \to -1$ as suggested in Remark 4. $L_s(t)$ when $t \leq 50$ is cut due to visibility of the threshold 0 and -1. The full figure is in Appendix F.

(b) $I_T W_2(p_T, q_T)$ and $W_2(p_0, q_0)$ for $T = 10, 50, 100, 150, 200$ respectively at the convergence of training, while β to be sigmoid schedule from $\beta_1 = 10^{-5}$ to $\beta_T = 10^{-2}$ for all choices of T. We can observe that $I(T)W_2(p_T, q_T)$ decays almost exponentially as T increases, as suggested in Remark 4.

Figure 4: Effect of T. We can observe that $L_s(T)$ would converge to -1, and $I(T)W_2(p_T, q_T)$ would decay almost exponentially when T is sufficiently large.

where π_t is an optimal transport plan between $\rho_t^{(1)}$ and $\rho_t^{(2)}$. See [1, Theorem 8.4.7] and [29, Corollary 5.25]. This equality plays an important role when we estimate the Wasserstein distance. The following proposition is a consequence of (21) and Lemma 1 in Appendix C.

Proposition 1. *Let p_t and q_t be solutions to* (2) *and* (6)*, respectively, in $[0, T]$. Then,*

$$-\frac{d}{dt} W_2(p_t, q_t) \leq (L_f + L_s g^2) W_2(p_t, q_t) + g^2 b^{\frac{1}{2}} \text{ where } b(t) := \mathbb{E}_{p_t} \left[\|\nabla \log p_t(x) - s_\theta(x, t)\|^2 \right]. \tag{22}$$

6 Discussions

Convergence of the Wasserstein distance If $p_T = q_T$, then our result yields that the Wasserstein distance $W_2(p_0, q_0)$ converges to zero as J_{SM} goes to zero. However, if p_T is not equal to q_T and f and g are not explicitly given, then we find that it is not easy to obtain such a convergence result. This is due to the nature of the Langevin dynamics in the reverse process. Typically, additional assumptions such as the log-concavity are required to guarantee the Langevin dynamics convergence, for instance, [11].

In addition, the dependence of the neural network s_θ makes it hard to evaluate the precise upper bound. We expect that appropriate assumptions on s_θ and the initial distribution p_0 may resolve this issue, but we do not pursue them in this paper.

Estimating L_s Estimating our theoretical bound requires estimating the one-sided Lipschitz constant L_s, which is NP-hard [37], making it difficult to estimate in high dimensional data. Note that the two-sided Lipschitz constant is an upper bound of the one-sided one, which can be more efficiently estimated as shown in [14]. However, it would result in a looser theoretical upper bound if we adopt the two-sided Lipschitz constant, and estimating the one-sided Lipschitz constant does not enjoy the semi-definiteness of estimating the two-sided one in [14].

7 Conclusion

In this paper, we mainly discuss the relationship between the loss function of score-based models and the Wasserstein distance. With the help of the theory in optimal transport, we present a novel upper bound on Wasserstein distance in terms of the loss function used in practice. We also provide numerical experiments to verify our theoretical bound with various simulation datasets. Moreover, we explore factors that affect the estimation of the upper bound and show the corresponding influence on the training process.

References

[1] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows in metric spaces and in the space of probability measures.* Second edition. Lecture notes in Mathematics ETH Zürich. Birkhäuser Verlag, Bassel, (2008).

[2] B. D. Anderson. Reverse-time diffusion equation models. *Stochastic Processes and their Applications*, 12(3):313–326, 1982.

[3] M. Arjovsky, S. Chintala, and L. Bottou. Wasserstein generative adversarial networks. In *International conference on machine learning*, pages 214–223. PMLR, 2017.

[4] J. Behrmann, W. Grathwohl, R. T. Chen, D. Duvenaud, and J.-H. Jacobsen. Invertible residual networks. In *International Conference on Machine Learning*, pages 573–582. PMLR, 2019.

[5] R. N. Bhattacharya and E. C. Waymire. *Stochastic processes with applications*. SIAM, 2009.

[6] V. I. Bogachev, N. V. Krylov, M. Röckner, and S. V. Shaposhnikov. *Fokker-Planck-Kolmogorov Equations*, volume 207. American Mathematical Soc., 2015.

[7] F. Bolley and J. A. Carrillo. Nonlinear diffusion: geodesic convexity is equivalent to wasserstein contraction. *Communications in Partial Differential Equations*, 39(10):1860–1869, 2014.

[8] J. A. Carrillo, R. J. McCann, and C. Villani. Contractions in the 2-wasserstein length space and thermalization of granular media. *Archive for Rational Mechanics and Analysis*, 179(2):217–263, 2006.

[9] N. Chen, Y. Zhang, H. Zen, R. J. Weiss, M. Norouzi, and W. Chan. Wavegrad: Estimating gradients for waveform generation. *arXiv preprint arXiv:2009.00713*, 2020.

[10] I. Csiszár and P. C. Shields. Information theory and statistics: A tutorial. 2004.

[11] A. S. Dalalyan. Theoretical guarantees for approximate sampling from smooth and log-concave densities. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(3):651–676, 2017.

[12] P. Dhariwal and A. Nichol. Diffusion models beat gans on image synthesis. *Advances in Neural Information Processing Systems*, 34, 2021.

[13] T. Donchev and E. Farkhi. Stability and euler approximation of one-sided lipschitz differential inclusions. *SIAM journal on control and optimization*, 36(2):780–796, 1998.

[14] M. Fazlyab, A. Robey, H. Hassani, M. Morari, and G. Pappas. Efficient and accurate estimation of lipschitz constants for deep neural networks. *Advances in Neural Information Processing Systems*, 32, 2019.

[15] R. Flamary, N. Courty, A. Gramfort, M. Z. Alaya, A. Boisbunon, S. Chambon, L. Chapel, A. Corenflos, K. Fatras, N. Fournier, L. Gautheron, N. T. Gayraud, H. Janati, A. Rakotomamonjy, I. Redko, A. Rolet, A. Schutz, V. Seguy, D. J. Sutherland, R. Tavenard, A. Tong, and T. Vayer. Pot: Python optimal transport. *Journal of Machine Learning Research*, 22(78):1–8, 2021.

[16] I. Goodfellow, J. Pouget-Abadie, M. Mirza, B. Xu, D. Warde-Farley, S. Ozair, A. Courville, and Y. Bengio. Generative adversarial nets. *Advances in neural information processing systems*, 27, 2014.

[17] J. Ho, A. Jain, and P. Abbeel. Denoising diffusion probabilistic models. *Advances in Neural Information Processing Systems*, 33:6840–6851, 2020.

[18] R. Jordan, D. Kinderlehrer, and F. Otto. The variational formulation of the Fokker–Planck equation. *SIAM journal on Mathematical Analysis*, 29(1):1–17, 1998.

[19] D. P. Kingma and M. Welling. Auto-encoding variational bayes. *arXiv preprint arXiv:1312.6114*, 2013.

[20] I. Kobyzev, S. J. Prince, and M. A. Brubaker. Normalizing flows: An introduction and review of current methods. *IEEE transactions on pattern analysis and machine intelligence*, 43(11):3964–3979, 2020.

[21] M. W. Lam, J. Wang, D. Su, and D. Yu. Bddm: Bilateral denoising diffusion models for fast and high-quality speech synthesis. *arXiv preprint arXiv:2203.13508*, 2022.

[22] I. Loshchilov and F. Hutter. Fixing weight decay regularization in adam. 2018.

[23] P. A. Markowich and C. Villani. On the trend to equilibrium for the fokker-planck equation: an interplay between physics and functional analysis. *Mat. Contemp*, 19:1–29, 2000.

[24] C. Meng, Y. He, Y. Song, J. Song, J. Wu, J.-Y. Zhu, and S. Ermon. Sdedit: Guided image synthesis and editing with stochastic differential equations. In *International Conference on Learning Representations*, 2021.

[25] A. Nichol, P. Dhariwal, A. Ramesh, P. Shyam, P. Mishkin, B. McGrew, I. Sutskever, and M. Chen. Glide: Towards photorealistic image generation and editing with text-guided diffusion models. *arXiv preprint arXiv:2112.10741*, 2021.

[26] A. Q. Nichol and P. Dhariwal. Improved denoising diffusion probabilistic models. In *International Conference on Machine Learning*, pages 8162–8171. PMLR, 2021.

[27] F. Otto. The geometry of dissipative evolution equations: the porous medium equation. 2001.

[28] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, and E. Duchesnay. Scikit-learn: Machine learning in Python. *Journal of Machine Learning Research*, 12:2825–2830, 2011.

[29] F. Santambrogio. *Optimal Transport for Applied Mathematicians*. Calculus of Variations, PDEs, and Modeling. Birkhäuser, Oct. 2015.

[30] J. Sohl-Dickstein, E. Weiss, N. Maheswaranathan, and S. Ganguli. Deep unsupervised learning using nonequilibrium thermodynamics. In *International Conference on Machine Learning*, pages 2256–2265. PMLR, 2015.

[31] J. Song, C. Meng, and S. Ermon. Denoising diffusion implicit models. *arXiv preprint arXiv:2010.02502*, 2020.

[32] Y. Song, C. Durkan, I. Murray, and S. Ermon. Maximum likelihood training of score-based diffusion models. *Advances in Neural Information Processing Systems*, 34:1415–1428, 2021.

[33] Y. Song and S. Ermon. Generative modeling by estimating gradients of the data distribution. *Advances in Neural Information Processing Systems*, 32, 2019.

[34] Y. Song and S. Ermon. Improved techniques for training score-based generative models. *Advances in neural information processing systems*, 33:12438–12448, 2020.

[35] Y. Song, J. Sohl-Dickstein, D. P. Kingma, A. Kumar, S. Ermon, and B. Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*, 2020.

[36] P. Vincent. A connection between score matching and denoising autoencoders. *Neural computation*, 23(7):1661–1674, 2011.

[37] A. Virmaux and K. Scaman. Lipschitz regularity of deep neural networks: analysis and efficient estimation. *Advances in Neural Information Processing Systems*, 31, 2018.

A Assumptions and Proofs

In addition to **(A1)** and **(A2)** in Section 3.1, we assume the following:

(A3) For the initial distribution p_0 of (2) and the final distribution q_T of (6), the following quantities are finite:

$$\mathbb{E}_{p_0}[\|\log p_0\|], \quad \mathbb{E}_{p_0}[\Lambda(x)] \quad \mathbb{E}_{q_T}[\|\log q_T\|], \quad \text{and} \quad \mathbb{E}_{q_T}[\Lambda(x)] \tag{23}$$

where $\Lambda(x) := \log \max(\|x\|, 1)$.

(A4) There exist constants C_1 and C_2 such that

$$f(x, t) \cdot x \le C_1 \|x\|^2 + C_2 \text{ for all } x \in \mathbb{R}^d \text{ and } t \in [0, T]. \tag{24}$$

(A5) There exists a positive constant C_3 such that

$$\frac{1}{C_3} < g(t) < C_3 \text{ for all } t \in [0, T]. \tag{25}$$

(A6) $\int_0^T \mathbb{E}_{p_t}[f^2]dt$ and $\int_0^T \mathbb{E}_{q_t}[(f - g^2 s_\theta)^2]dt$ are finite.

The above assumptions guarantee the integrability of the score function $\nabla \log p$.

Proposition 2. *[6, Theorem 4.2.1] A solution p_t to (2) satisfies $\nabla p_t \in L^1(\mathbb{R}^d)$ and*

$$\int_0^T \mathbb{E}_{p_t}\left[\|\nabla \log p_t(x)\|^2\right] dt < +\infty. \tag{26}$$

We give proofs in smooth settings.

(A7) $p_0, q_T \in C^2(\mathbb{R}^d)$, and $g \in C^2([0, T])$. f and s_θ are C^2 in both $x \in \mathbb{R}^d$ and $t \in [0, T]$.

(A8) A solution p_t to (2) and a solution q_t to (6) are positive C^2 probability densities decaying rapidly in $\mathbb{R}^d$: there exists $k > 0$ such that $p_t(x) = O(\exp(-\|x\|^k))$ and $q_t(x) = O(\exp(-\|x\|^k))$ for all $t \in [0, T]$. Furthermore, the vector fields $v[p_t] = f - \frac{1}{2}g^2 \nabla \log p_t$, and $v[q_t] = \left(f - g^2 s_\theta\right) + \frac{1}{2}g^2 \nabla \log q_t$ grow at most linearly as $\|x\| \to \infty$.

Proof of Theorem 1. Recall the differential inequality (22) below from Proposition 1:

$$-\frac{d}{dt}W_2(p_t, q_t) \le (L_f + L_s g^2)W_2(p_t, q_t) + g^2 b^{\frac{1}{2}}.$$

It can be solved by introducing the integrating factor

$$I(t) := \exp\left(\int_0^t L_f(r) + L_s(r)g^2(r)dr\right) \text{ where } b(t) := \mathbb{E}_{p_t}\left[\|\nabla \log p_t(x) - s_\theta(x, t)\|^2\right]. \tag{27}$$

As $\frac{d}{dt}I(t) = (L_f(t) + L_s(t)g^2(t))I(t)$, the above inequality (22) can be written as

$$-\frac{d}{dt}\{I(t)W_2(p_t, q_t)\} \le g^2(t)I(t)b^{\frac{1}{2}}(t).$$

Integrating both sides from 0 to T, we obtain that

$$I(0)W_2(p_0, q_0) - I(T)W_2(p_T, q_T) \le \int_0^T g^2(t)I(t)b^{\frac{1}{2}}(t)dt.$$

As $I(0) = 1$, we conclude that

$$W_2(p_0, q_0) \le \int_0^T g^2(t)I(t)b^{\frac{1}{2}}(t)dt + I(T)W_2(p_T, q_T). \tag{28}$$

$\square$

Proof of Corollary 1. Let

$$J_I(\theta) = \int_0^T g^2(t)I(t)b^{\frac{1}{2}}(t)dt. \tag{29}$$

Here, $I(t)$ and $b(t) = \mathbb{E}_{p_t}\left[\|\nabla \log p_t(x) - s_\theta(x,t)\|^2\right]$ are given in (27). The Cauchy-Schwarz inequality yields

$$J_I(\theta) \leq \left(\int_0^T 2g^4(t)I^2(t)\lambda^{-1}(t)dt\right)^{\frac{1}{2}} \left(\frac{1}{2}\int_0^T \lambda(t)b(t)dt\right)^{\frac{1}{2}}. \tag{30}$$

Since $\lambda = g^2$ and J_{SM} given in (5) satisfies

$$J_{SM}(\theta;\lambda) = \frac{1}{2}\int_0^T \lambda(t)\mathbb{E}_{p_t}\left[\|\nabla \log p_t(x) - s_\theta(x,t)\|_2^2\right]dt = \frac{1}{2}\int_0^T \lambda(t)b(t)dt, \tag{31}$$

we conclude (32):

$$W_2(p_0, q_0) \leq \sqrt{2\left(\int_0^T g^2(t)I(t)^2dt\right)J_{SM} + I(T)W_2(p_T, q_T)}.$$

$\square$

Remark 7. *It is expected that the parallel results hold for weak solutions to* (2) *and* (4) *by using suitable approximations. This approach has been studied for the Wasserstein contraction property in [8, Section 6.2].*

From (30) and (31) in the proof of Corollary 1, we obtain the following result for other choices of λ.

Corollary 4. *Let p_0 and q_0 be given in Theorem 1. Suppose that $g^4 I^2 \lambda^{-1}$ is integrable in $[0, T]$. Then the following inequality holds:*

$$W_2(p_0, q_0) \leq \sqrt{2\left(\int_0^T g^4(t)I(t)^2\lambda^{-1}(t)dt\right)J_{SM} + I(T)W_2(p_T, q_T)}. \tag{32}$$

B One-sided Lipschitzness

For an arbitrary Lipschitz function $F : \mathbb{R}^d \to \mathbb{R}^d$, the Cauchy-Schwarz inequality yields that

$$(F(x) - F(y)) \cdot (x - y) \leq \|F(x) - F(y)\|\|x - y\| \leq L\|x - y\|^2 \tag{33}$$

where L is the Lipschitz constant of F. Therefore, all Lipschitz function satisfies the one-sided Lipschitz condition:

$$(F(x) - F(y)) \cdot (x - y) \leq L\|x - y\|^2. \tag{34}$$

As pointed out earlier in Section 3.1, the one-sided Lipschitz constant is not necessarily to be positive. For instance, if $F(x) = -ax + b$ for $a > 0$ and $b \in \mathbb{R}^d$, then $-a < 0$ can be the one-sided Lipschitz constant of F while its Lipschitz constant is $a > 0$. Figure 5 visualizes this.

C Technical lemmas

Lemma 1. *Let π_t be an optimal transport plan between p_t and q_t. Then, we have*

$$\mathbb{E}_{\pi_t}\left[(x - y) \cdot (v[q_t](y) - v[p_t](x))\right] \leq W_2(p_t, q_t)\left\{(L_f + L_s g^2)W_2(p_t, q_t) + g^2 b^{\frac{1}{2}}\right\} \tag{35}$$

where $b(t) := \mathbb{E}_{p_t}\left[\|\nabla \log p_t(x) - s_\theta(x,t)\|^2\right]$.

(a) Two-sided Lipschitz   (b) One-sided Lipschitz; $L > 0$ (c) One-sided Lipschitz; $L < 0$

Figure 5: Two-sided and one-sided Lipschitzness.

Proof. The left-hand side of (35) is given by

$$\mathbb{E}_{\pi_t}\left[(x - y) \cdot (v[q_t](y) - v[p_t](x))\right] = \mathbb{E}_{\pi_t}\left[(x - y) \cdot (f(y, t) - f(x, t))\right]$$
$$+ g^2 \mathbb{E}_{\pi_t}\left[(x - y) \cdot (\nabla \log p_t(x) - s_\theta(y, t))\right]$$
$$+ \frac{g^2}{2}\mathbb{E}_{\pi_t}\left[(x - y) \cdot (\nabla \log q_t(y) - \nabla \log p_t(x))\right].$$

In Lemma 2 below, we prove that the last term is less than or equal to zero.

In what follows, we estimate the first two terms. First, using the Lipschitzness of f in space, we get

$$\mathbb{E}_{\pi_t}\left[(x - y) \cdot (f(y, t) - f(x, t))\right] \le L_f \mathbb{E}_{\pi_t}\left[\|x - y\|^2\right] = L_f W_2^2(p_t, q_t).$$

The last equality follows from the fact that π_t is an optimal plan between p_t and q_t.

Next, the second term $g^2 \mathbb{E}_{\pi_t}\left[(x - y) \cdot (\nabla \log p_t(x) - s_\theta(y, t))\right]$ is the sum of the following two terms:

$$I_1 := g^2 \mathbb{E}_{\pi_t}\left[(x - y) \cdot (s_\theta(x, t) - s_\theta(y, t))\right]$$

and

$$I_2 := g^2 \mathbb{E}_{\pi_t}\left[(x - y) \cdot (\nabla \log p_t(x) - s_\theta(x, t))\right].$$

As shown above, the former one I_1 is bounded from above by $g^2 L_s W_2^2(p_t, q_t)$.

It suffices to find an upper bound on the latter one I_2. By the Cauchy-Schwarz inequality, we have

$$I_2 \le g^2 \mathbb{E}_{\pi_t}\left[\|x - y\|^2\right]^{\frac{1}{2}} \mathbb{E}_{\pi_t}\left[|\nabla \log p_t(x) - s_\theta(x, t)|^2\right]^{\frac{1}{2}}.$$

As the marginals of π_t are p_t and q_t, it holds that

$$\mathbb{E}_{\pi_t}\left[\|\nabla \log p_t(x) - s_\theta(x, t)\|^2\right] = \mathbb{E}_{p_t}\left[\|\nabla \log p_t(x) - s_\theta(x, t)\|^2\right].$$

As a consequence, we conclude that

$$I_1 + I_2 \le g^2(t) W_2(p_t, q_t) \left\{L_s W_2(p_t, q_t) + b(t)^{\frac{1}{2}}\right\}$$

where $b(t) = \mathbb{E}_{p_t}\left[\|\nabla \log p_t(x) - s_\theta(x, t)\|^2\right]$. $\qquad\qquad\square$

Before proving the lemma below, let us recall some basic definitions from the theory of optimal transport. The Wasserstein distance defined in (7) has an equivalent formulation:

$$W_2(\mu, \nu) = \inf \left\{\int_{\mathbb{R}^d} \|x - T(x)\|^2 d\mu : T_\# \mu = \nu\right\}^{\frac{1}{2}}. \tag{36}$$

The optimizer of the above problem is called the optimal map from μ to ν. It is well known that there exists a convex function ϕ such that $T = \nabla \phi$.

Lemma 2. $\mathbb{E}_{\pi_t}\left[(x - y) \cdot (\nabla \log q_t(y) - \nabla \log p_t(x))\right]$ *is nonpositive.*

Proof. Let T_t be an optimal transport map from p_t to q_t and a convex function ϕ_t satisfy $\nabla \phi_t = T_t$ for all $t \in [0, T]$. As in the proof of [7, Theorem 1], we have

$$\mathbb{E}_{\pi_t}\left[(x - y) \cdot (\nabla \log q_t(y) - \nabla \log p_t(x))\right] = -\mathbb{E}_{p_t}[\Delta \phi_t + \Delta \phi_t^*(\nabla \phi_t) - 2d] \tag{37}$$

where ϕ_t^* is a convex conjugate of ϕ_t. The convexity of ϕ_t yields that $\Delta \phi_t + \Delta \phi_t^*(\nabla \phi_t) - 2d$ and we conclude. $\qquad\qquad\square$

 # D   Proofs of Theorem 2 and Corollary 3

479   **Definitions**   For a given t, define

$$J_{SM}(\theta, t) = \frac{1}{2} \mathbb{E}_{p_t(x)}[\|s_\theta(x, t) - \nabla_x \log p_t(x)\|^2], \tag{38}$$

480   and

$$J_{DSM}(\theta, t) = \frac{1}{2} \mathbb{E}_{p_0(x(0))} \mathbb{E}_{p_{0t}(x|x(0))}[\|s_\theta(x, t) - \nabla_x \log p_{0t}(x|x(0))\|^2]. \tag{39}$$

481   **Lemma 3.** *(Appendix in [36])*

$$\mathbb{E}_x[\nabla_x \log p_t(x)] = \mathbb{E}_{x(0)}\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))]. \tag{40}$$

482   *Proof of Theorem 2.*   From [36], we have

$$\begin{aligned}
J_{SM}(\theta, t) &= J_{DSM}(\theta, t) \\
&+ \frac{1}{2}(\mathbb{E}_{p_t(x)}\left[\|\nabla_x \log p_t(x)\|^2\right] - \mathbb{E}_{p_0(x(0))}\mathbb{E}_{p_{0t}(x|x(0))}[\|\nabla_x \log p_{0t}(x|x(0))\|^2])
\end{aligned} \tag{41}$$

483   Note that

$$\begin{aligned}
&\mathbb{E}_{p_t(x)}\left[\|\nabla_x \log p_t(x)\|^2\right] - \mathbb{E}_{p_0(x(0))}\mathbb{E}_{p_{0t}(x|x(0))}[\|\nabla_x \log p_{0t}(x|x(0))\|^2] \\
&= \mathbb{E}_x[(\nabla_x \log p_t(x))^\top (\nabla_x \log p_t(x))] \\
&- \mathbb{E}_{x(0)}[\mathbb{E}_{x|x(0)}[(\nabla_x \log p_{0t}(x|x(0)))^\top (\nabla_x \log p_{0t}(x|x(0)))|x(0)]] \\
&= \mathrm{Var}[(\nabla_x \log p_t(x))^\top] - \mathbb{E}[\mathrm{Var}[(\nabla_x \log p_{0t}(x|x(0)))^\top]|x(0)] \\
&+ (\mathbb{E}_x[\nabla_x \log p_t(x)])^\top (\mathbb{E}_x[\nabla_x \log p_t(x)]) \\
&- \mathbb{E}_{x(0)}[(\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)])^\top (\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)])] \\
&\leq (\mathbb{E}_x[\nabla_x \log p_t(x)])^\top (\mathbb{E}_x[\nabla_x \log p_t(x)]) \\
&- \mathbb{E}_{x(0)}[(\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)])^\top (\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)])]
\end{aligned} \tag{42}$$

484   where the inequality comes from the law of total variance and our condition:

$$\begin{aligned}
&\mathrm{Var}[(\nabla_x \log p_t(x))^\top] - \mathbb{E}[\mathrm{Var}[(\nabla_x \log p_{0t}(x|x(0)))^\top|x(0)]] \\
&= \mathrm{Var}[\mathbb{E}[(\nabla_x \log p_{0t}(x|x(0)))^\top|x(0)]] = 0.
\end{aligned} \tag{43}$$

485   Then we have

$$\begin{aligned}
&(\mathbb{E}_x[\nabla_x \log p_t(x)])^\top (\mathbb{E}_x[\nabla_x \log p_t(x)]) \\
&- \mathbb{E}_{x(0)}[(\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)])^\top (\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)])] \\
&\leq (\mathbb{E}_x[\nabla_x \log p_t(x)])^\top (\mathbb{E}_x[\nabla_x \log p_t(x)]) \\
&- (\mathbb{E}_{x(0)}\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)])^\top (\mathbb{E}_{x(0)}\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)]) \\
&= (\mathbb{E}_x[\nabla_x \log p_t(x)] - \mathbb{E}_{x(0)}\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)])^\top \\
&(\mathbb{E}_x[\nabla_x \log p_t(x)] + \mathbb{E}_{x(0)}\mathbb{E}_{x|x(0)}[\nabla_x \log p_{0t}(x|x(0))|x(0)]) \\
&= 0,
\end{aligned} \tag{44}$$

486   where first inequality comes from Jensen's inequality, and the last equality comes from eq. (40).

487   Recall that $J_{SM}(\theta, \lambda) = \int_0^T \lambda(t) J_{SM}(\theta, t)dt$ and $J_{DSM}(\theta, \lambda) = \int_0^T \lambda(t) J_{DSM}(\theta, t)dt$, $\lambda(t) > 0$,
488   we have that $J_{DSM} \geq J_{SM}$. Plugging it in eq. (32), we can get eq. (16). $\square$

489   *Proof of Corollary 3.*   We have $p_{0t}(x|x(0)) = \mathcal{N}(\sqrt{\bar{\alpha}_t}x(0), (1 - \bar{\alpha}_t)I)$ where $\bar{\alpha}_t = \prod_{r=1}^t (1 - \beta_t)$,
490   which can be inferred from $p(x(t)|x(t-1)) = \mathcal{N}(\sqrt{1 - \beta_t}x(t-1), \beta_t I)$.

491   Thus we can show that $\mathbb{E}[\nabla_x \log p_{0t}(x|x(0))^\top|x(0)]$ is constant with respect to $x(0)$: recall
492   $p_{0t}(x|x(0)) = \mathcal{N}(\sqrt{\bar{\alpha}_t}x(0), (1 - \bar{\alpha}_t)I)$, we have $\nabla_x \log p_{0t}(x|x(0)) = -((1 - \bar{\alpha}_t)I)^{-1}(x(t) -$

$\sqrt{\bar{\alpha}_t}x(0))$, which is a linear function of $x(t) - \sqrt{\bar{\alpha}_t}x(0)$. Using the Gaussian density function, we have:

$$\int p_{0t}(x|x(0))\nabla_x \log p_{0t}(x|x(0))dx = 0, \tag{45}$$

As a result, $\mathrm{Var}[\mathbb{E}[(\nabla_x \log p_{0t}(x|x(0)))^\top|x(0)]] = 0$, which satisfies the condition of eq. (15) in Theorem 2.

Note that here the assumption of f and g is only a sufficient condition for $\mathrm{Var}[\mathbb{E}[(\nabla_x \log p_{0t}(x|x(0)))^\top|x(0)]] = 0$. In fact, any conditional distribution p_{0t} that satisfies $\mathrm{Var}[\mathbb{E}[(\nabla_x \log p_{0t}(x|x(0)))^\top|x(0)]] = 0$ can lead to the same conclusion. $\square$

E   Numerical results on J_{DSM} upper-bounding J_{SM} in DDPM

To verify $J_{SM} \leq J_{DSM}$ in (16) for DDPM, we adopt the same datasets as in Fig 1, and the same training and evaluation settings in Section 4.1.

Moreover, to estimate J_{SM} numerically, we estimate $p_t(x)$ by performing Gaussian kernel density estimation with bandwidth = 0.05 on sampled data. $\nabla_x p_t(x)$ is estimated by central difference approximation with interval = 0.01. The resulting plots of J_{DSM} and J_{SM} are shown in Fig 6, which shows that J_{DSM} is an upper bound of J_{SM} in DDPM during training, where p_0 is the dataset and q_0 is the generated data distribution at the convergence of training.

(a) 1 cluster in 2D   (b) 4 clusters in 2D

Figure 6: J_{SM} and J_{DSM} during training. The datasets are the same as in Fig 1 in 2D. The training curves are obtained via training DDPM with modification of J_{DSM} loss.

F   Full plot of $L_s(t)$ when $T = 100$

See Fig. 7.

G   Log-log plots with weight decay

See Fig. 8.

H   Further analysis of the upper bound

H.1   Proof of Corollary 2

We further investigate the upper bound that appeared in the previous sections. To illustrate this, let us consider the following case:

$$f(t) = -\frac{\beta(t)}{2}x \text{ and } g(t) = \sigma\sqrt{\beta(t)} \tag{46}$$

for a positive function $\beta : [0, T] \rightarrow (0, +\infty)$.

Figure 7: Full plot of $L_s(t)$ when $T = 100$.

(a) Coefficient=0.01   (b) Coefficient=0.1   (c) Coefficient=0.5   (d) Coefficient=1   (e) Coefficient=5

Figure 8: Log-log plots for different weight decay coefficients. As the weight decay coefficient increases, the theoretical upper bound is approaching the empirical one.

Proof of Corollary 2. In the case of (46), the stationary solution to (2) is given as the probability distribution of $\mathcal{N}(0, \sigma^2 I)$:

$$\phi(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{\|x\|^2}{2\sigma^2}\right). \tag{47}$$

Based on the contraction property [8], we quantify the Wasserstein distance between p_T and ϕ.

$$W_2(p_T, \phi) \leq \exp\left(-\int_0^T \frac{\beta(t)}{2} dt\right) W_2(p_0, \phi) \tag{48}$$

Using the above, the definition of $I(t)$, and $q_T = \phi$, we conclude (13). $\qquad\square$

It worth noting that as a consequence of (48), $W_2(p_T, \phi)$ is small for an appropriate choice of T and $\beta(t)$.

H.2 Exponential convergence of h_t

For simplicity of notations, we define the norm in $L^2(\phi)$ as follows,

$$\|f\|_{L^2(\phi)} := \left(\int_{\mathbb{R}^d} f^2 d\phi\right)^{\frac{1}{2}}, \tag{49}$$

where ϕ is given in (47). In addition, assume that

$$\beta(t) > c > 0 \text{ for all } t \geq 0. \tag{50}$$

Lemma 4. *Under the same setting as in Corollary 2, we have*

$$\|h_t - 1\|_{L^2(\phi)} \leq \exp\left(-\frac{\sigma^2\lambda}{2}\int_0^T \beta(t)dt\right)\|h_0 - 1\|_{L^2(\phi)}. \tag{51}$$

where $h_t = p_t/\phi$ for some constant $\lambda > 0$. In particular, h_t exponentially converges to 1 in $L^2(\phi)$ for β satisfying (50).

Remark 8. *The convergence of h_t to 1 can be shown under more general assumption: $\beta > 0$ satisfying*

$$\lim_{T\to\infty}\int_0^T \beta(t)dt = \infty.$$

For a constant function β, Lemma 4 is proven in [23]. For the sake of completeness, we provide the proof, which is a small modification of [23, Section 2].

Proof of Lemma 4. In our case, the equation (2) of p_t is given by

$$\partial_t p - \frac{\beta}{2}\left(\nabla \cdot (px) + \sigma^2\Delta p\right) = 0, \quad p(\cdot, 0) = p_0. \tag{52}$$

By direct computations, we obtain the equation of $h(x, t) = h_t(x)$ as follows:

$$\partial_t h - \frac{\beta}{2}\left(-\nabla h \cdot x + \sigma^2\Delta h\right) = 0, \quad h(\cdot, 0) = h_0. \tag{53}$$

We estimate $\|h - 1\|_{L^2(\phi)}^2$ by differentiating it with respect to time. Using the integration by parts and Poincaré inequality, we obtain that

$$\frac{d}{dt}\|h - 1\|_{L^2(\phi)}^2 = -\sigma^2\beta(t)\|\nabla h\|_{L^2(\phi)}^2 \leq -\sigma^2\lambda\beta(t)\|h - 1\|_{L^2(\phi)}^2 \tag{54}$$

This yields (51). Lastly, for β satisfying (50), $\|h_t - 1\|_{L^2(\phi)}^2 \leq \exp(-\sigma^2\lambda ct)\|h_0 - 1\|_{L^2(\phi)}^2$. Thus, we conclude the exponential convergence of h to 1. $\qquad\square$

Further analysis is plausible based on the techniques in the study of partial differential equations.

Remark 9. *As p_t is given as the convolution between p_0 and the Gaussian distribution, it is smooth for $t > 0$. Therefore, h_t is also smooth, and the higher order derivatives of h_t are all bounded. As a consequence, the above result combining with Gagliardo–Nirenberg interpolation inequality yield that the gradient of h, Dh, and the Hessian of h, D^2h, also converge to zero in $L^2(\phi)$.*

Remark 10. *Proving the uniform convergence of h, Dh or D^2h requires an additional technical assumption that the support of h is bounded. Under the assumption, another interpolation inequality, Agmon's inequality yields the desired uniform convergence result.*

H.3   Estimation of L_s

In this subsection, we investigate the estimation of L_s. If J_{SM} is sufficiently small, then s_θ is close to $\nabla \log p_t$. We first investigate the one-sided Lipschitz constant of $\nabla \log p_t$.

Lemma 5. *Under the same setting as in Corollary 2, $\nabla \log p_t$ satisfies the one-sided Lipschitz condition with a constant $(-\sigma^{-2} + \|D^2(\log h)\|_\infty)$ i.e.,*

$$(\nabla \log p_t(x) - \nabla \log p_t(y)) \cdot (x - y) \leq (-\sigma^{-2} + \|D^2(\log h)\|_\infty) \|x - y\|^2. \tag{55}$$

where $\| \cdot \|_\infty$ denotes the supremum of the matrix norm,

$$\|D^2(\log h)\|_\infty := \sup_{x \in \mathbb{R}^d} \|D^2(\log h(x))\| = \sup_{x,y \in \mathbb{R}^d, \|y\| \leq 1} \|D^2(\log h(x))y\|. \tag{56}$$

Proof. From the definition of h_t, we have

$$\log p_t(x) = \log h_t(x) + \log \phi(x) = \log h_t(x) - \frac{x^2}{2\sigma^2} - c \tag{57}$$

for some constant c. As a consequence,

$$(\nabla \log p_t(x) - \nabla \log p_t(y)) \cdot (x - y) = (\nabla \log h_t(x) - \nabla \log h_t(y)) \cdot (x - y) - \sigma^{-2} \|x - y\|^2. \tag{58}$$

To prove (55), it suffices to estimate $(\nabla \log h_t(x) - \nabla \log h_t(y)) \cdot (x - y)$. From the fundamental theorem of calculus, we have

$$(\nabla \log h_t(x) - \nabla \log h_t(y)) = \int_0^1 D^2(\log h_t)(sx + (1 - s)y)ds \cdot (x - y). \tag{59}$$

Using $|z^\top D^2(\log h_t(w))z| \leq \|D^2(\log h)\|_\infty \|z\|^2$ for all $w, z \in \mathbb{R}^d$, we have

$$(\nabla \log h_t(x) - \nabla \log h_t(y)) \cdot (x - y) \leq \|D^2(\log h_t)\|_\infty \|x - y\|^2 \tag{60}$$

and conclude (55). $\square$

Remark 11. *Based on the similar relation as in (59), the difference between L_s and $-\sigma^{-2} + \|D^2(\log h)\|_\infty$ can be estimated. More precisely, we have $L_s(t) = (-\sigma^{-2} + \|D^2(\log h)\|_\infty) + \epsilon(t)$. Here, $\epsilon(t)$ depends on the difference between s_θ and $\nabla \log p_t$. Therefore, it is expected that the upper bound of $\int_0^T \epsilon(t)dt$ is given by J_{SM} under suitable regularity assumptions.*

I   Verifying the upper bound in high-dimensional data

Here we present numerical results on some moderately high dimensional data, where estimating both $W_2(p_0, q_0)$ and L_s would become more computationally expensive.

I.1   Gaussian clusters in higher dimension

Here we present the log-log plots for Gaussian cluster $\mathcal{N}(\mathbf{0}, 0.1\mathbf{I})$ with $d = 10, 20$ and 100 respectively in Fig 9, where the training configuration is the same as Section 4. As we can observe, our upper bound is still a valid one of the empirical Wasserstein distance, though the gap between the empirical one and the theoretical one becomes larger when the dimensionality is higher.

(a) 10D   (b) 20D   (c) 100D

Figure 9: Log-log plots for Gaussian cluster in different dimensionalities.

I.2 MNIST data

We also trained the diffusion model on MNIST dataset for 100 epochs with $T = 100$, while we modify the neural network architecture to UNet, omit the loss maximization training, and following other experimental configurations as in [17]. As in Section 4, we still sample from $p_T = q_T$ when calculating $W_2(p_0, q_0)$, in order to satisfy $W_2(p_T, q_T) \approx 0$ for the log-log plot. The generated images and the corresponding log-log plot obtained during training are shown in Fig 10, where we also show the validity of our upper bound on image data.

(a) Generated images for MNIST after 100 epochs.

(b) Log-log plot for MNIST training.

Figure 10: Results on MNIST data.

J Computing resources

We use Google Colab (`https://research.google.com/colaboratory`) for our training.