Revisiting the stability of stochastic gradient descent: a tightness analysis

Abstract

The technique of algorithmic stability has been used to capture the generalization power of several learning models, especially those trained with stochastic gradient descent (SGD). This paper investigates the tightness of the algorithmic stability bounds for SGD given by Hardt et al. (2016). We show that the analysis of Hardt et al. (2016) is tight for convex objective functions, but loose for non-convex objective functions. In the non-convex case we provide a tighter upper bound on the stability (and hence generalization error), and provide evidence that it is asymptotically tight up to a constant factor.

However, deep neural networks trained with SGD exhibit much better stability and generalization in practice than what is suggested by these (tight) bounds, namely, linear or exponential degradation with time for SGD with constant step size. We aim towards characterizing deep learning loss functions with good generalization guarantees, despite being trained using SGD with constant step size.

We propose the Hessian Contractive (HC) condition, which specifies the contractivity of regions containing local minima in the neural network loss landscape. We provide empirical evidence that this condition holds for several loss functions, and provide theoretical evidence that the known tight SGD stability bounds for convex and non-convex loss functions can be circumvented by HC loss functions, thus partially explaining the generalization of deep neural networks.

1 Introduction

Stochastic gradient descent (SGD) has gained great popularity in solving machine learning optimization problems (Kingma & Ba, 2014; Johnson & Zhang, 2013). SGD leverages the finite-sum structure of the objective function, avoids the expensive computation of exact gradients, and thus provides a feasible and efficient optimization solution in large-scale settings (Bottou, 2012). The convergence and the optimality of SGD have been thoroughly studied (Ge et al., 2015; Rakhlin et al., 2012; Reddi et al., 2018; Zhou & Gu, 2019; Carmon et al., 2019a;b; Shamir & Zhang, 2013) .

In recent years, new research questions have been raised regarding SGD's impact on a model's generalization power. The seminal work (Hardt et al., 2016) tackled the problem using the *algorithmic stability* of SGD, i.e., the progressive sensitivity of the trained model w.r.t. the replacement of a single (test) datum in the training set. The stability-based analysis of the generalization gap allows one to bypass classical model capacity theorems (Vapnik, 1998; Koltchinskii & Panchenko, 2000) or weight-based complexity theorems (Neyshabur et al., 2017; Bartlett et al., 2017; Arora et al., 2018). This framework also provides theoretical insights into many phenomena observed in practice, e.g., the "train faster, generalize better" phenomenon, the power of regularization techniques such as weight decay (Krogh & Hertz, 1992), Dropout (Srivastava et al., 2014), and gradient clipping. Other works have applied the stability analysis to more sophisticated settings such as Stochastic Gradient Langevin Dynamics and momentum SGD (Mou et al., 2018; Chaudhari et al., 2019; Chen et al., 2018).

Despite the promises of this stability-based analysis, it remains open whether this framework can explain the strong generalization performance of deep neural networks in practice. Existing theoretical upper bounds of the stability (and thus, generalization) (Hardt et al., 2016) are ideal for strongly convex loss functions: the upper bound remains constant even as the number of training iterations increases. However, the same bound deteriorates significantly when we relax to more general and realistic settings. In particular, for convex (but not strongly convex) and non-convex loss functions, if

SGD has constant step size, then the upper bound grows linearly and exponentially with the number of training iterations. This bound fails to match the superior generalization performance of deep neural networks, and leads to the following question:

Question 1: *Can we find a better stability upper bound for convex or non-convex loss functions?*

In this paper, we first address the question above and investigate the tightness of the algorithmic stability analysis for stochastic gradient methods (SGM) proposed by (Hardt et al., 2016).

R1. We show in Theorem 1 that the analysis in (Hardt et al., 2016) is tight for convex and smooth objective functions; in other words, there is a convex loss function whose stability grows linearly with the number of training iterations, with constant step size ($\alpha_t = \alpha$) in SGD.

R2. We show that in Theorem 2 that for linear models, the analysis in the convex case can be tightened to show that ϵ_{stab} does not increase with t.

R3. In Theorem 3 we show that the analysis in (Hardt et al., 2016) for decreasing step size ($\alpha_t = O(1/t)$) is loose for non-convex objective functions by providing a tighter upper bound on the stability (and hence generalization error).

R4. The bound on the stability of SGD by (Hardt et al., 2016) is achieved by bounding the divergence at time t, defined as $\delta_t := \mathbb{E}\|w_t - w_t'\|$, where w_t is the model trained on data set S and w_t' is the model trained on a data set S' that differs from S in exactly one sample. In Theorem 4 we provide evidence that our new upper bound in the non-convex case is tight, by showing a non-convex loss function whose divergence matches the upper bound for our divergence.

R5. Although it is not derived formally, the techniques in (Hardt et al., 2016) can be employed to show an exponential upper bound for non-convex loss functions minimized using SGD with constant-size step. In Theorem 5, we give evidence that this abysmal upper bound is likely tight for non-convex loss functions, by exhibiting a non-convex loss function for which the divergence δ_t increases exponentially.

Thus the only functions whose stability provably does not increase with the number of iterations when a constant step-size during SGD is employed, are strongly convex functions. However, a) it has been empirically observed that for deep neural network loss, near the local minima, the Hessians are usually low rank (Chaudhari et al., 2017; Yao et al., 2019), and b) neural networks trained with constant step-size SGD do generalize well in practice (Lin & Jegelka, 2018; Huang et al., 2017; Smith et al., 2017). Combined with our lower bounds on convex and non-convex functions, we seem to hit an obstacle on the way to explaining generalization using the stability framework.

Question 2: *What is it that makes constant-step SGD on deep learning loss function generalize well?*

Realizing the limitation of the current state of stability analysis, we investigate whether a stronger-than-convex, but weaker-than-strongly-convex assumption of the loss function can be made, at least near local minima. If we can show algorithmic stability near local minima, we can still show the stability using similar argument as (Du et al., 2019; Allen-Zhu et al., 2019).

Aiming towards a characterization of loss functions exhibiting good stability, we propose a new condition for loss near local minima. This condition, called *Hessian contractive*, is slightly stronger than a general convex condition, but considerably weaker than strongly convex. Formally, the Hessian contractive condition stipulates that near any local minima, (1) the function is convex; and (2) a data dependent Hessian is positive definite in the gradient direction. Theoretically, we show that such a condition is sufficient to guarantee a constant stability bound for SGD (constant step size) near the local minima, while allowing the Hessian to be low rank. We also provide examples showing Hessian Contractive is a reasonable condition for several loss functions. Empirically, we verify the Hessian Contractive condition near a local minima of the loss while training deep neural networks. We sample points from a neighborhood of current iterates by adding Gaussian noise and verify the HC condition locally by Hessian product approximation. Summarizing our second set of contributions:

R6. In Observation 1 we show that the family of widely used (convex) linear model loss functions will satisfy the Hessian Contractive condition. One typical example of such linear model loss is the regression loss function. These observation suggests that Hessian Contractive is a condition satisfied by (potentially many) machine learning loss functions.

R7. In Theorem 6 we show that the Hessian Contractive condition will localize SGD iterates in a neighborhood of minima, which implies a constant stability bound for SGD near the local minima.

Table 1: Current landscape of stability bounds. [H] indicates results in (Hardt et al., 2016), and * indicates results in this paper. Bounds without [H] or * are trivial. β is the smoothness parameter.

SGD Step Size	Constant $\alpha_t = a/\beta$			$\alpha_t = a/(\beta t)$
Loss function	Strongly Convex	Hessian Contractive	Convex	Non-Convex
Upper Bound	$O(1)$ [H]	$O(1)$*	$O(aT/n)$ [H]	$O\left(T^{\frac{a}{1+a}}/n\right)$ [H] $O\left(T^a/n^{1+a}\right)$*
Lower Bound	$\Omega(1)$	$\Omega(1)$	$\Omega(aT/n)$*	Open, evidence*

1.1 RELATED WORKS

Stability and generalization. The stability framework suggests that a stable machine learning algorithm results in models with good generalization performance (Kearns & Ron, 1999; Bousquet & Elisseeff, 2002; Elisseeff et al., 2005; Shalev-Shwartz et al., 2010; Devroye & Wagner, 1979a;b; Rogers & Wagner, 1978). It serves as a mechanism for provable learnability when uniform convergence fails (Shalev-Shwartz et al., 2010; Nagarajan & Kolter, 2019). The concept of uniform stability was introduced in order to derive high probability bounds on the generalization error (Bousquet & Elisseeff, 2002). Uniform stability describes the worst case change in the loss of a model trained on an algorithm when a single data point in the dataset is replaced. In (Hardt et al., 2016), a uniform stability analysis for *iterative algorithms* is proposed to analyze SGD, generalizing the one-shot version in (Bousquet & Elisseeff, 2002). Algorithmic uniform stability is widely used in analyzing the generalization performance of SGD (Mou et al., 2018; Feldman & Vondrak, 2019; Chen et al., 2018). The worst case leave-one-out type bounds also closely connect uniform stability with *differential private learning* (Feldman et al., 2018; 2020; Dwork et al., 2006; Wu et al., 2017b), where the uniform stability can lead to provable privacy guarantee. While the upper bounds of algorithmic stability of SGD have been extensively studied, the tightness of those bounds remains open. In this work, we report for the first time lower bounds on the uniform stability of SGD. Our tightness analysis suggests necessity of additional assumptions for analyzing the generalization of SGD on deep learning.

Geometry of local minima. The geometry of local minima plays an important role in the generalization performance of deep neural networks (Hochreiter & Schmidhuber, 1995; Wu et al., 2017a). The flat minima, i.e., minima whose Hessians have a large portion of zero-valued eigenvalues, are believed to attain better generalization (Keskar et al., 2016; Li et al., 2018). In (Chaudhari et al., 2019), the authors construct a local entropy-based objective function which converges to a solution with good generalization in a flat region, where "flatness" means that the Hessian matrix has a large portion of nearly-zero eigenvalues. However, these observations have not been supported theoretically. In this paper, we propose the Hessian contractive condition that is slight stronger than flat minima. Such condition suggests that the minima is sharp only in the gradient direction while remains flat in other directions, which unifies the geometrical interpretation of flat minima and uniform stability analysis.

2 PRELIMINARIES

In this section we introduce the notion of uniform stability and establish our notations. We first introduce the quantities *Empirical and Population Risk* and *Generalization Gap*. Given an unknown distribution $\mathcal{D}$ on labeled sample space $Z = X \times \{-1, +1\}$, let $S = \{z_1, ..., z_n\}$ denote a set of n samples $z_i = (x_i, y_i)$ drawn i.i.d. from $\mathcal{D}$. Let $w \in \mathbb{R}^d$ be the parameter(s) of a model that tries to predict y given x, and let f be a loss function where $f(w; z)$ denotes the loss of the model with parameter w on sample z. Let $f(w; S)$ denote the *empirical risk* $f(w; S) = E_{z \sim S}[f(w; z)] = \frac{1}{n}\sum_{i=1}^n f(w; z_i)$ with corresponding *population risk* $E_{z \sim \mathcal{D}}[f(w; z)]$. The *generalization error* of the model with parameter w is defined as the difference between the empirical and population risks:

$$|E_{z \sim \mathcal{D}}[f(w; z)] - E_{z \sim S}[f(w; z)]|.$$

Next we introduce the Stochastic Gradient Descent (SGD) method. We follow the setting of (Hardt et al., 2016), and starting with some initialization $w_0 \in \mathbb{R}^d$, consider the following SGD update step:

$$w_{t+1} = w_t - \alpha_t \nabla_w f(w; z_{i_t}),$$

where i_t is drawn from $[n] := \{1, 2, \cdots, n\}$ uniformly and independently in each round. The analysis of SGD requires the following crucial properties of the loss function $f(., z)$ at any fixed point z, viewed solely as a function of the parameter w:

Definition 1 (L-Lipschitz)**.** *A function $f(w)$ is L-Lipschitz if $\forall u, v \in \mathbb{R}^d$: $|f(u) - f(v)| \leq L\|u - v\|$.*

Definition 2 (β-smooth)**.** *A function $f(w)$ is β-smooth if $\forall u, v \in \mathbb{R}^d$: $|\nabla f(u) - \nabla f(v)| \leq \beta\|u - v\|$.*

Definition 3 (γ-strongly convex)**.** *A function $f(w)$ is γ strongly convex if $\forall u, v \in \mathbb{R}^d$:*

$$f(u) > f(v) + \nabla f(v)^\top [u - v] + \frac{\gamma}{2}\|u - v\|^2.$$

Algorithmic Stability   Next we define the key concept of *algorithmic stability*, which was introduced by (Bousquet & Elisseeff, 2002) and adopted by (Hardt et al., 2016). Informally, an algorithm is *stable* if its output only varies slightly when we change a single sample in the input dataset. When this stability is *uniform* over all datasets differing at a single point, this leads to an upper bound on the generalization gap. More formally:

Definition 4. *Two sets of samples S, S' are twin datasets if they differ at a single entry, i.e., $S = \{z_1, ...z_i, ..., z_n\}$ and $S' = \{z_1, ..., z_i', ..., z_n\}$.*

Consider a possibly randomized algorithm A that given a sample S of size n outputs a parameter $A(S)$. Define the algorithmic stability parameter of A by:

$$\varepsilon_{\text{stab}}(A, n) := \inf\{\varepsilon : \sup_{z \in Z, S, S'} \mathbb{E}_A |f(A(S); z) - f(A(S'); z)| \leq \varepsilon\}.$$

Here $\mathbb{E}_A$ denote expectation over the random coins of A. Also, for such an algorithm, one can define its expected generalization error as:

$$GE(A, n) := \mathbb{E}_{S,A}[E_{z \sim \mathcal{D}}[f(A(S); z)] - E_{z \sim S}[f(A(S); z)]].$$

Stability and generalization: It was proved in (Hardt et al., 2016) that $GE(A, n) \leq \varepsilon_{\text{stab}}(A, n)$. Furthermore, the authors observed that an L-Lipschitz condition on the loss function f enforces a uniform upper bound: $\sup_z |f(w; z) - f(w'; z)| \leq L\|w - w'\|$. This implies that for Lipschitz loss, the algorithmic stability $\varepsilon_{\text{stab}}(A, n)$ (and hence the generalization error $GE(A, n)$) can be bounded by obtaining bounds on $\|w - w'\|$.

Let w_T and w_T' be the parameters obtained by running SGD starting on twin datasets S and S', respectively. Throughout this paper we will focus on the *divergence quantity* $\delta_T := \mathbb{E}_A\|w_T - w_T'\|$. While (Hardt et al., 2016) reports upper bounds on δ_T with different types of loss functions, e.g., convex and non-convex loss functions, we investigate the tightness of those bounds.

3   TIGHTNESS OF EXISTING BOUNDS

In this section we report our main results. We first consider the convex case with constant step size, where we prove 1) that the existing bounds stating $\epsilon_{\text{stab}} \propto t$ are tight, and 2) for linear models, the analysis can be tightened to show that ϵ_{stab} does not increase with t. Then we move on to the non-convex case, where a) for decreasing step size we improve the existing upper bound, and give evidence that our new upper bound is tight, and b) for constant step size we give loss functions whose divergence δ_t increases exponentially with t.

3.1   CONVEX CASE

In this section we analyze the stability of SGD when loss function is convex and smooth. We begin with a construction which shows that Theorem 3.8 in (Hardt et al., 2016) is tight. Our lower bound analysis will require the following quadratic function:

$$f(w; z) = \frac{1}{2}w^\top A w - yx^\top w \tag{1}$$

where A is a $d \times d$ matrix. In the construction of lower bounds, we carefully choose A and S so that the single data point replaced in twin data set will cause the instability of SGD.

Theorem 1. *Let w_t, w_t' be the outputs of SGD on twin datasets S, S' respectively. Let $\Delta_t = w_t - w_t'$ and α_t be the step size of SGD. There exists a function f which is convex, β-smooth, and L-Lipschitz on the domain of w, and twin datasets S, S' such that:*

$$\mathbb{E}\|\Delta_T\| \geq \frac{1}{n}\sum_{t=1}^{T}\alpha_t, \;\; and \;\; \varepsilon_{stab} \geq \frac{1}{n}\sum_{t=1}^{T}\alpha_t. \tag{2}$$

The convex upper bound in Theorem 3.8 in (Hardt et al., 2016) states that $\|\Delta_T\| \leq \frac{L\sum_{i=1}^{T}\alpha_t}{n}$, which implies that the divergence increases throughout training. The lower bound in Theorem 1 suggests the tightness of the upper bound. However, in practice, such phenomenon is not commonly observed, i.e., for a family of convex but not-strongly-convex loss functions, the generalization performance does not deteriorate as the number of training iterations increases. This motivates us to investigate a weaker condition which still can enforce an $O(1)$ stability, without strong convexity. In the next theorem, we restrict ourselves to a family of linear model loss functions and show that the divergence will not increase unboundedly during training.

We shall need the following definition of a ξ-self correlated data set. Essentially, a self-correlated dataset requires an average linear dependence of each x. Recall that the i'th sample $z_i = (x_i, y_i)$.

Definition 5. *A set $S = \{z_1, ..., z_n\}$ is ξ-self correlated if $\forall j \in [n], \frac{1}{n}\sum_{i=1}^{n}(x_j^\top x_i)^2 \geq \xi > 0$.*

Assuming that $\forall j \in [n], \|x_j\| \geq r$ for some $r > 0$, definition 5 implies that S is at least $\frac{r^2}{n}$-self correlated. Thus the above condition holds for all datasets S not containing the zero-feature vector. In our next theorem, we leverage on the ξ-self correlated condition to prove a non-accumulate uniform stability bound for SGD on a loss function of *Linear Models*. We characterize a linear model by rewriting the loss function $f(w; z)$ in terms of $f_y(w^\top x)$ where $f_y(\cdot)$ is a scalar function depending only on the inner product of the model parameter w and the input feature x.

Theorem 2. *Suppose a loss function $f(w, z)$ is of the form $f(w, S) = \frac{1}{n}\sum_{j=1}^{n}f_{y_j}(w^\top x_j)$, where $f_y(w^\top x)$ satisfies (1) $|f_y'(\cdot)| \leq L$, (2) $0 < \gamma \leq f_y''(\cdot) \leq \beta$, (3) $\|x\| \leq R$ and (4) S, S' are ξ-self correlated, twin datasets. Let w_t and w_t' be the outputs of SGD on S and S' after t steps, respectively. Let the divergence $\Delta_t := w_t - w_t'$ and $\alpha \leq \frac{1}{\beta}$ be the step size of SGD. Then,*

$$\mathbb{E}\|\Delta_T\| \leq \frac{4LR}{\xi\gamma n}, \;\; and \;\; \varepsilon_{stab} \leq \frac{4L^2R^2}{\xi\gamma n}.$$

Remark 1. *In (Hardt et al., 2016), an $O\left(\frac{L^2}{\gamma n}\right)$ stability bound is derived on a loss function $f(w, z_i)$ which is strongly convex, i.e., $\nabla^2 f(w) \succ \gamma I$. In practice one can incorporate a strongly convex regularizer to impose strong convexity, often resulting in improved generalization performance in practice (Shalev-Shwartz et al., 2010; Bousquet & Elisseeff, 2002). The ξ-self correlated condition allows SGD to maintain a uniformly upper-bounded stability guarantee for a family of widely used models for arbitrary long training without using strongly convex regularizer. The theorem suggests that if the dataset S is reasonably simple, e.g., every x_i lies in a low dimensional subspace, the algorithmic stability of SGD is comparable with a strongly convex loss function. This analysis suggests an alternative condition other can strong convexity can empower SGD an $O(1)$ stability which is data-dependent. This motivates us to go beyond linear model and seek for a generalized condition in Theorem 2. In section 4.1, we propose the Hessian Contractive condition for more general loss function driven by the observation on the linear model.*

Example: Linear and logistic regression. Linear regression minimizes the quadratic loss on w: $f(w, S) = \frac{1}{2n}\sum_{x_j \in S}(x_j^\top w - y_j)^2$. Note that the Hessian of an individual linear regression loss term is $x_j x_j^\top$ which is *not strongly convex* since it has rank 1. One cannot apply the strongly convex bound, and the bound for convex suggests stability will increase linearly. However, one can rewrite the loss function as $f_y(w^\top x)$ where $f_y''(\cdot) = 1$. Hence Thm. 2 can be applied to give a non-accumulative bound on SGD's stability. A similar result can be derived for the *logistic regression* loss.

3.2 Non-Convex Case

In this section, we construct a non-convex loss function to analyze the tightness of the divergence bound in (Hardt et al., 2016). We first focus on the case where SGD applies a step size that *decreases*

with t. We first fix a hitting time t_0 and prove Lemma 4 in the Appendix. The following theorem follows from Lemma 4 by optimizing over t_0. The choice of hitting time t_0 plays an important role in the analysis, which is also illustrated in the "burn-in Lemma" 3.11 in (Hardt et al., 2016).

Theorem 3. *Let w_t, w_t' be the SGD outputs on twin datasets S, S', $\Delta_t := w_t - w_t'$. There exists f that is non-convex and β-smooth, twin datasets S, S' and a constant $a > 0$ such that the SGD divergence after T rounds ($T > n$) using decreasing step size $\alpha_t = \frac{a}{0.99\beta t}$ satisfies: $\mathbb{E}\|\Delta_T\| \geq \frac{T^a}{3n^{1+a}}$.*

In above theorem, the lower bound is derived by choosing $t_0 = n$ in Lemma 4. The bound in (Hardt et al., 2016) is of the form $O\left(\frac{T^{\frac{a}{1+a}}}{n}\right)$ which does not match the above lower bound. According to the lower bound provided in Theorem 3, the bound in Hardt et al. (2016) may not be tight in the region $T^{\frac{a}{1+a}} \leq n$. We investigate this gap and derive a tighter bound in the next theorem which improves the Theorem 3.12 in Hardt et al. (2016).

Theorem 4. *Assume f is β-smooth and L-lipschitz. Running T ($T > n$) iterations of SGD on $f(w; S)$ with step size $\alpha_t = \frac{a}{\beta n}$, the stability of SGD satisfies:*

$$\mathbb{E}\|\Delta_T\| \leq \frac{2LT^a}{n^{1+a}}, \; \varepsilon_{stab} \leq \frac{2L^2T^a}{n^{1+a}}. \tag{3}$$

Dividing our bound with the one in Theorem 3.12 in Hardt et al. (2016), we obtain the ratio $\Omega\left(\frac{T^{\frac{a^2}{1+a}}}{n^a}\right)$. This factor is less than 1 (and so we improve the upper bound) exactly when $T^{\frac{a}{1+a}} \leq n$. Note that this is potentially a large range as a is a small and positive constant. We remark that our tight bound is for *permutation SGD*. The bound for SGD (see Appendix Theorem 4b) using sampling without replacement has an additional $\log(n)$ factor (which we conjecture can be removed), which nevertheless is also a polynomial improvement over the known bounds.

4 DEEP LEARNING AND LIMITATIONS OF BOUNDS

While Theorem 4 improves existing upper bounds on stability by an polynomial factor, it still can not explain the generalization performance of deep learning model. In particular, the analysis relies on a decreasing step size while in training deep learning model, constant instead of a decreasing step size is a common choice. In the next result, we adopt the same construction as in Theorem 3 but with a constant learning rate to show that, unfortunately, SGD may have an exponential divergence rate for general β smooth function.

Theorem 5. *Let w_t, w_t' be the outputs of SGD on twin datasets S, S', and let $\Delta_t := w_t - w_t'$. There exists a function f which is non-convex and β-smooth, twin sets S, S' and constants a, γ such that the divergence of SGD after T rounds ($T > n$) using constant step size $\alpha = \frac{a}{0.99\beta}$ satisfies: $\mathbb{E}\|\Delta_T\| \geq \frac{1}{n^2}e^{aT/2}$.*

The above theorem implies that the generalization gap may deteriorate at an exponential rate if SGD is run with a constant step size for a non-convex function. However, in Fig. 1 we observe that the distance between the weights of two deep models trained on twin datasets (in other words, the divergence δ_T *stabilizes after* ≈ 20

Figure 1: Following the experiment performed by (Hardt et al., 2016), we measure the normalized Euclidean distance between the parameters of two identical deep models–here we used ResNet18–trained on twin datasets: i.e., CIFAR10, with a different random image removed. We repeated this experiment for 35 random pairs of images removed: the variance is shown around the mean curve. The figure shows that the distance initially increases but then stabilizes after ≈ 20 epochs of training. The curve labeled "all" is the distance between the full parameter vectors of the respective models, whereas conv$i$ is the distance between the weights of the first convolutional layer in the i-th "block" of the ResNet18 model.

epochs of training. Fig. 1 suggests that the generalization gap for deep models may remain constant

after sufficiently many training iterations. However, the analysis of (Hardt et al., 2016) suggests that local minima of the loss landscape must be in strongly convex regions in order to achieve this sort of stability. Indeed, strongly convex Hessians have been proven to be contractive, leading to the stability of SGD (Ge et al., 2015). On the other hand, it is known that the Hessians of the loss functions of common deep learning architectures have a large fraction of eigenvalues which are close to zero (Yao et al., 2019; Chaudhari et al., 2017). This phenomenon requires further investigation to close the gap between theory and practice. Next, we formalize a condition requiring that local minima of deep learning loss landscapes lie in contractive regions, which aids in explaining why the generalization gap stabilizes. We then provide empirical evidence that this property does indeed hold.

4.1 GEOMETRY OF LOCAL MINIMA

In Theorems 1 and 5, we show that in general, divergence (and hence stability) may deteriorate at a linear or exponential rate with the number of training iterations when the local minima are not in strongly convex regions. We introduce the following property to aid in understanding the stability of the training curve of SGD.

Definition 6. *[Hessian Contractive] For a given set $S = \{z_1, ...z_n\}$, a local minimum w^* is in a (σ, γ)-Hessian contractive region if $\forall w, w'$ with $\max(\|w' - w^*\|, \|w - w^*\|) \leq \sigma$ and $z \in S$,*

$$\nabla_w f(w, z)^\top \mathcal{H}_{w'} \nabla_w f(w, z) \geq \gamma \|\nabla_w f(w, z)\|^2 \tag{4}$$

where $\mathcal{H}_{w'} = \nabla_w^2 f(w'; S) = \frac{1}{n} \sum_{j=1}^n \nabla_w^2 f(w', z_j)$. When (σ, γ)-Hessian contractive holds for all $\sigma > 0$ and local minima w^*, we say $f(w; S)$ is globally γ-Hessian contractive.

Hessian contractivity describes a region where the stochastic gradient will lie in the range of a positive semi-definite Hessian matrix. Such a condition prevents the iterates of SGD from escaping from a local minima in the sense that all gradient descent steps will be pulled back to the minima by the power of Hessian, which mimics the structure of a strongly convex loss function but with a weaker assumption. To show that this class is nonempty, we observe the (not strongly convex) linear models loss functions in Theorem 2 satisfy Hessian contractive condition globally.

Observation 1. *Suppose loss function $f(w; S)$ conditions in Theorem 2, i.e., $f(w, S) = \frac{1}{n} \sum_{j=1}^n f_{y_j}(w^\top x_j)$ satisfies (1) $|f_y'(\cdot)| \leq L$, (2) $\gamma \leq f_y''(\cdot) \leq \beta$, (3) $\|x\| \leq R$ and (4) S is ξ-self correlated, we have $f(w; S)$ is globally $\frac{\xi\gamma}{R^2}$ Hessian Contractive.*

Dependence on dataset The Hessian Contractive condition states that the gradient of loss function evaluated on z can be regulated by Hessian evaluated on the whole dataset, which is a generalization of ξ-self correlated set condition. In the linear model case, the power of Hessian Contractive condition depends on the the average correlation between data points which indicates the effect of data complexity in generalization. We postulate that the value of γ in Hessian Contractive condition can be one complexity measurement of the data, which is believed to affect generalization power (Kawaguchi et al., 2017).

Next, we leverage on the Hessian contractive condition to show that once SGD hits a stationary point around w^* in a (σ, γ)-Hessian contractive region, it will be localized near the minima. The localization of SGD implies that the divergence curve will be stable eventually.

Theorem 6. *Given a set S and a β-smooth and L-Lipschitz loss funtion f, suppose SGD with a fixed step size $\alpha \leq \frac{\sigma}{2L}$ reaches a stationary point w_0 with $\|\nabla_w f(w_0, S)\| \leq \frac{\gamma\sigma}{2}$, which is close to a local minimum w^* in a (σ, γ)-Hessian contractive region, i.e., $\|w_0 - w^*\| \leq \frac{\sigma}{2}$. Then, $\forall T \geq 1$, we have $\|w_T - w^*\| \leq \sigma$.*

Theorem 6 states that once the iterate w of SGD encounters a stationary point in a (σ, γ)-Hessian contractive region, w will remain in this region. In terms of the stability of SGD, assuming SGD will bring w_t into one of the local minima quickly (Ge et al., 2015; Allen-Zhu et al., 2019), the divergence $\|w_t - w_t'\|$ will be uniformly bounded by the difference between two local minima $w^*, w^{*\prime}$ of loss functions $f(w; S)$ and $f(w'; S')$ plus the σ-radius of the Hessian contractive region. The $O(1)$ uniform stability of SGD follows from the fact that the loss f is L-lipschitz. The above suggests that a weaker condition than strong convexity is sufficient to explain the observation in Fig. 1. In the next section, we provide empirical evidence that the local minima of deep learning loss landscapes lie in Hessian contractive regions.

Figure 2: Approximate normalized Hessian contractivity. For both neural networks, the "mean" contractivity value at each epoch is averaged over 100 perturbations and 5 different copies of the network trained on CIFAR10. Similarly for "mean(min)", except the min of the 100 approximations after perturbing is taken. Both quantities for both networks provide evidence that SGD converges to an increasingly contractive solution.

4.2 EMPIRICAL SUPPORT FOR HESSIAN CONTRACTIVITY

In order to explain the stabilization of the generalization gap observed when training deep learning models via SGD, we hypothesize that local minima of deep loss landscapes found by SGD lie in Hessian contractive regions. We propose to empirically support this hypothesis by locally approximating the LHS of Eq. (6)–normalized by $\|\nabla_w f(\tilde{w}, z)\|^2$–during the training of a model, and showing that this quantity steadily increases as training progresses. This indicates that SGD leads the training into regions of increasing contractivity.

We now explain the nature of our approximation: For a model w^*, we generate many random samples $\tilde{w}$ from the unit sphere around w^*. For each such $\tilde{w}$, we locally estimate a stochastic version of Hessian-gradient product to get the approximation

$$\nabla_w f(\tilde{w}, z)^\top \mathcal{H}(z) \nabla_w f(\tilde{w}, z) \approx \frac{1}{\eta} [\nabla_w f(\tilde{w}, z) - \nabla_w f(\tilde{w} - \eta \nabla_w f(\tilde{w}, z), z)]^\top \nabla_w f(\tilde{w}, z) \quad (5)$$

where the gradients on the RHS approximation are taken with respect to a random minibatch of samples. For each perturbation $\tilde{w}$ we normalize the corresponding approximation by $\|\nabla_w f(\tilde{w}, z)\|^2$, and then take the mean and minimum over all of these quantities.

For our experiments, we used the well-known CIFAR10 dataset (Krizhevsky et al., 2009) and two popular deep architectures: ResNet18, and DenseNet121 (Huang et al., 2017; Lin & Jegelka, 2018). To match the settings in the theorems of this paper as much as possible and avoid any confounding factors, we avoided standard regularization (e.g. weight decay), data augmentation, and adaptive learning rates. We used SGD with a constant learning rate of 0.01 and momentum of 0.9. The models were trained for 200 epochs and batch size of 128.

We trained five copies of each network and checkpointed the models every 8 epochs. For each checkpoint, we generated 100 random perturbations from the unit sphere around the checkpoint weights. We used $\eta = 0.001$ and batch size 128 for the approximation in Eq. (5). We averaged over these 100 approximations and then further took the average over the five networks. The results of these experiments are shown in Fig. 2, and indicate a clear increase in Hessian contractivity as training progresses.

5 CONCLUSION

In this paper, we studied the stability bound of SGD with regard to different types of loss functions. We proved a better upper bound and proved the tightness of various bounds. These tightness results suggest that existing stability bounds may not suffice in answering the generalization problem of deep neural nets. We propose a new Hessian contractive condition that is slighly stronger than convex, but with $O(1)$ stability bound. We provide empirical evidence to support the hypothesis that deep learning loss function is Hessian contractive near local minima.

REFERENCES

Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. A convergence theory for deep learning via over-parameterization. In *International Conference on Machine Learning*, pp. 242–252. PMLR, 2019.

Sanjeev Arora, Rong Ge, Behnam Neyshabur, and Yi Zhang. Stronger generalization bounds for deep nets via a compression approach. In *Proceedings of the 35th International Conference on Machine Learning*, ICML, 2018.

Peter L Bartlett, Dylan J Foster, and Matus J Telgarsky. Spectrally-normalized margin bounds for neural networks. In *Advances in Neural Information Processing Systems*, pp. 6240–6249, 2017.

Léon Bottou. Stochastic gradient descent tricks. In *Neural networks: Tricks of the trade*, pp. 421–436. Springer, 2012.

Olivier Bousquet and André Elisseeff. Stability and generalization. *Journal of machine learning research*, 2(Mar):499–526, 2002.

Yair Carmon, John C Duchi, Oliver Hinder, and Aaron Sidford. Lower bounds for finding stationary points i. *Mathematical Programming*, pp. 1–50, 2019a.

Yair Carmon, John C Duchi, Oliver Hinder, and Aaron Sidford. Lower bounds for finding stationary points ii: First-order methods. *Mathematical Programming*, pp. 1–50, 2019b.

Pratik Chaudhari, Anna Choromanska, Stefano Soatto, Yann LeCun, Carlo Baldassi, Christian Borgs, Jennifer Chayes, Levent Sagun, and Riccardo Zecchina. Entropy-sgd: Biasing gradient descent into wide valleys. In *ICLR*, 2017.

Pratik Chaudhari, Anna Choromanska, Stefano Soatto, Yann LeCun, Carlo Baldassi, Christian Borgs, Jennifer Chayes, Levent Sagun, and Riccardo Zecchina. Entropy-sgd: Biasing gradient descent into wide valleys. *Journal of Statistical Mechanics: Theory and Experiment*, 2019(12):124018, 2019.

Yuansi Chen, Chi Jin, and Bin Yu. Stability and convergence trade-off of iterative optimization algorithms. *arXiv preprint arXiv:1804.01619*, 2018.

Luc Devroye and T Wagner. Distribution-free performance bounds with the resubstitution error estimate (corresp.). *IEEE Transactions on Information Theory*, 25(2):208–210, 1979a.

Luc Devroye and Terry Wagner. Distribution-free inequalities for the deleted and holdout error estimates. *IEEE Transactions on Information Theory*, 25(2):202–207, 1979b.

Simon Du, Jason Lee, Haochuan Li, Liwei Wang, and Xiyu Zhai. Gradient descent finds global minima of deep neural networks. In *International Conference on Machine Learning*, pp. 1675–1685, 2019.

Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam Smith. Calibrating noise to sensitivity in private data analysis. In *Theory of cryptography conference*, pp. 265–284. Springer, 2006.

Andre Elisseeff, Theodoros Evgeniou, and Massimiliano Pontil. Stability of randomized learning algorithms. *Journal of Machine Learning Research*, 6(Jan):55–79, 2005.

Vitaly Feldman and Jan Vondrak. High probability generalization bounds for uniformly stable algorithms with nearly optimal rate. *arXiv preprint arXiv:1902.10710*, 2019.

Vitaly Feldman, Ilya Mironov, Kunal Talwar, and Abhradeep Thakurta. Privacy amplification by iteration. In *2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS)*, pp. 521–532. IEEE, 2018.

Vitaly Feldman, Tomer Koren, and Kunal Talwar. Private stochastic convex optimization: optimal rates in linear time. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, pp. 439–449, 2020.

Rong Ge, Furong Huang, Chi Jin, and Yang Yuan. Escaping from saddle points—online stochastic gradient for tensor decomposition. In *Conference on Learning Theory*, pp. 797–842, 2015.

Moritz Hardt, Ben Recht, and Yoram Singer. Train faster, generalize better: Stability of stochastic gradient descent. In *International Conference on Machine Learning*, pp. 1225–1234, 2016.

Sepp Hochreiter and Jürgen Schmidhuber. Simplifying neural nets by discovering flat minima. In *Advances in neural information processing systems*, pp. 529–536, 1995.

Gao Huang, Zhuang Liu, Laurens Van Der Maaten, and Kilian Q Weinberger. Densely connected convolutional networks. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pp. 4700–4708, 2017.

Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. In *Advances in neural information processing systems*, pp. 315–323, 2013.

Kenji Kawaguchi, Leslie Pack Kaelbling, and Yoshua Bengio. Generalization in deep learning. *arXiv preprint arXiv:1710.05468*, 2017.

Michael Kearns and Dana Ron. Algorithmic stability and sanity-check bounds for leave-one-out cross-validation. *Neural computation*, 11(6):1427–1453, 1999.

Nitish Shirish Keskar, Dheevatsa Mudigere, Jorge Nocedal, Mikhail Smelyanskiy, and Ping Tak Peter Tang. On large-batch training for deep learning: Generalization gap and sharp minima. *arXiv preprint arXiv:1609.04836*, 2016.

Diederik P Kingma and Jimmy Ba. Adam: A method for stochastic optimization. In *ICLR*, 2014.

Vladimir Koltchinskii and Dmitriy Panchenko. Rademacher processes and bounding the risk of function learning. In *High dimensional probability II*, pp. 443–457. Springer, 2000.

Alex Krizhevsky, Geoffrey Hinton, et al. Learning multiple layers of features from tiny images. 2009.

Anders Krogh and John A Hertz. A simple weight decay can improve generalization. In *Advances in neural information processing systems*, pp. 950–957, 1992.

Hao Li, Zheng Xu, Gavin Taylor, Christoph Studer, and Tom Goldstein. Visualizing the loss landscape of neural nets. In *Advances in Neural Information Processing Systems*, pp. 6389–6399, 2018.

Hongzhou Lin and Stefanie Jegelka. Resnet with one-neuron hidden layers is a universal approximator. In *Advances in Neural Information Processing Systems*, pp. 6172–6181, 2018.

Wenlong Mou, Liwei Wang, Xiyu Zhai, and Kai Zheng. Generalization bounds of sgld for non-convex learning: Two theoretical viewpoints. volume 75 of *Proceedings of Machine Learning Research*, pp. 605–638. PMLR, 06–09 Jul 2018. URL `http://proceedings.mlr.press/v75/mou18a.html`.

Vaishnavh Nagarajan and J Zico Kolter. Uniform convergence may be unable to explain generalization in deep learning. In *Advances in Neural Information Processing Systems*, pp. 11615–11626, 2019.

Behnam Neyshabur, Srinadh Bhojanapalli, David McAllester, and Nati Srebro. Exploring generalization in deep learning. In *Advances in Neural Information Processing Systems*, pp. 5947–5956, 2017.

Alexander Rakhlin, Ohad Shamir, and Karthik Sridharan. Making gradient descent optimal for strongly convex stochastic optimization. In *Proceedings of the 29th International Coference on International Conference on Machine Learning*, pp. 1571–1578. Omnipress, 2012.

Sashank J Reddi, Satyen Kale, and Sanjiv Kumar. On the convergence of adam and beyond. In *ICLR*, 2018.

William H Rogers and Terry J Wagner. A finite sample distribution-free performance bound for local discrimination rules. *The Annals of Statistics*, pp. 506–514, 1978.

Shai Shalev-Shwartz, Ohad Shamir, Nathan Srebro, and Karthik Sridharan. Learnability, stability and uniform convergence. *The Journal of Machine Learning Research*, 11:2635–2670, 2010.

Ohad Shamir and Tong Zhang. Stochastic gradient descent for non-smooth optimization: Convergence results and optimal averaging schemes. In *International Conference on Machine Learning*, pp. 71–79, 2013.

Samuel L Smith, Pieter-Jan Kindermans, Chris Ying, and Quoc V Le. Don't decay the learning rate, increase the batch size. *arXiv preprint arXiv:1711.00489*, 2017.

Nitish Srivastava, Geoffrey Hinton, Alex Krizhevsky, Ilya Sutskever, and Ruslan Salakhutdinov. Dropout: a simple way to prevent neural networks from overfitting. *The journal of machine learning research*, 15(1):1929–1958, 2014.

Vladimir N Vapnik. Statistical learning theory. *A Wiley-Interscience Publication*, 1998.

Lei Wu, Zhanxing Zhu, et al. Towards understanding generalization of deep learning: Perspective of loss landscapes. *arXiv preprint arXiv:1706.10239*, 2017a.

Xi Wu, Fengan Li, Arun Kumar, Kamalika Chaudhuri, Somesh Jha, and Jeffrey Naughton. Bolt-on differential privacy for scalable stochastic gradient descent-based analytics. In *Proceedings of the 2017 ACM International Conference on Management of Data*, pp. 1307–1322, 2017b.

Zhewei Yao, Amir Gholami, Kurt Keutzer, and Michael Mahoney. Pyhessian: Neural networks through the lens of the hessian. *arXiv preprint arXiv:1912.07145*, 2019.

Dongruo Zhou and Quanquan Gu. Lower bounds for smooth nonconvex finite-sum optimization. *arXiv preprint arXiv:1901.11224*, 2019.

A PROOFS

Lemma 1 (Dynamics of divergence). *Let $f(w; x) = \frac{1}{2}w^\top A w - yx$. Let Δ_t be $w_t - w_t'$ and α_t be the step size of SGD. Suppose $[x_i - x_i']/\|x_i - x_i'\|$ is an eigenvector of A where $A[x_i - x_i'] = \lambda_{xx'}[x_i - x_i']$. Suppose one run SGD on $f(w, S)$, the dynamics of Δ_t are given by:*

$$\mathbb{E}\|\Delta_{t+1}\| = (1 - \alpha_t \lambda_{xx'})\mathbb{E}\|\Delta_t\| + \frac{\alpha_t}{n}\|x_i - x_i'\| \tag{6}$$

Proof: In case the different entry z_i, z_i is not picked, the gradient difference of $f(w; z)$ and $f(w; z')$ is
$$\nabla f(w, z) - \nabla f(w', z') = A[w - w']$$
and in case different entry z_i, z_i is picked,
$$\nabla f(w, z) - \nabla f(w', z') = A[w - w'] + [x_i - x_i']$$
SGD selects $z_t = z_t'$ with probability $1 - \frac{1}{n}$ and a different entry with probability $\frac{1}{n}$. The lemma follows by combining the above two facts and $A[x_i - x_i'] = \lambda_{xx'}[x_i - x_i']$. $\qquad \square$

Lemma 2 (Lower bound on divergence). *Let Δ_t be $w_t - w_t'$, α_t be the step size of SGD and $\Delta_0 = 0$. Suppose $[x_i - x_i']/\|x_i - x_i'\|$ is an eigenvector of A where $A[x_i - x_i'] = \lambda_{xx'}[x_i - x_i']$. Running SGD on $f(w, S)$, we have:*

$$\mathbb{E}\|\Delta_T\| \geq \frac{\|x_i - x_i'\|}{n} \sum_{t=1}^{T-1} \prod_{\tau=t+1}^{T-1} \alpha_t (1 - \alpha_\tau \lambda_{xx'})$$

Proof: By Lemma 1 we have

$$\mathbb{E}\|\Delta_T\| = \mathbb{E}\|(I - \alpha_{T-1}A)\Delta_{T-1} + \frac{\alpha_{T-1}}{n}[x_i - x_i']\|$$

$$= (1 - \alpha_{T-1}\lambda_{xx'})\mathbb{E}\|\Delta_{T-1}\| + \frac{\alpha_{T-1}}{n}\|x_i - x_i'\|$$

$$= \|[x_i - x_i']\|\frac{1}{n} \sum_{t=1}^{T-1} \alpha_t \prod_{\tau=t+1}^{T-1} (1 - \alpha_\tau \lambda_{xx'})$$

$$\square$$

Theorem 1. *Let w_t, w'_t be the outputs of SGD on twin datasets S, S' respectively, Δ_t be $w_t - w'_t$ and α_t be the step size of SGD. There exists a function f which is convex and β-smooth, L-Lipschitz on domain of w_t, w'_t and twin datasets S, S' such that the divergence of the two SGD outputs satisfies:*

$$\mathbb{E}\|\Delta_T\| \geq \frac{1}{n}\sum_{t=1}^{T}\alpha_t; \quad \varepsilon_{stab} \geq \frac{1}{n}\sum_{t=1}^{T}\alpha_t \tag{7}$$

Proof We set $x_i = v, y_i = 0.5, x'_i = -v, y'_i = 0.5$. The proof is constructive. Let $f(w, z) = \frac{1}{2}w^\top A w - yx$. The function is β smooth and convex by setting $\beta = \|A\|$ and $u^\top A u \geq 0$ for all u. We set $S \setminus \{z_i\} = S' \setminus \{z'_i\}$ to lie in the range of A, where A is a PSD matrix which is not full-rank. We further set x_i, x'_i to lie in the null space of A so that $Av = 0$. The lower bound $\|\Delta\|_T$ follows from Lemma 2 from the fact that $\Delta_0 = 0$. Since $w^\top A w = w'^\top A w'$, $\varepsilon_{stab} = \sup_z \mathbb{E}|f(w_T, z) - f(w'_T, z)| \geq \mathbb{E}|v^\top[w_T - w'_T]| \geq \frac{1}{n}\sum_{t=1}^{T}\alpha_t$.

The last part we show for any sequence w_t generated from stochastic gradient descent step using data set S will have a constant gradient thus the Lipshitzness will follow. Let U be the range of A (so any $u \in U$ satisfy $u^A u > 0$. Pick γ so that $u^\top A u \geq \gamma > 0$ holds for all $u \in U$. For all u, $\|w_{t+1}^\top u\| \leq \|(I - \alpha_t A)w_t^\top u + \alpha z^\top u\| \leq (1 - \alpha_t\gamma)\|w_t^\top u\| + \alpha$. This implies $\|w_t^\top u\| \leq \frac{1}{\gamma}$. Now we bound gradient $\|\nabla f_w(w_t, z)\| = \|Aw_t - x\| \leq |v^\top A w| + |v^\top yx| + |u^\top A w| + |u^\top yx| \leq 1 + \frac{\beta}{\gamma}$. This implies that the function is L-Lipschitz with $L = 1 + \frac{\beta}{\gamma}$. $\qquad\square$

Theorem 2. *Suppose loss function $f(w, z)$ of the form $f(w, S) = \frac{1}{n}\sum_{j=1}^{n} f_{y_j}(w^\top x_j)$ and $f_y(w^\top x)$ satisfies (1) $|f'_y(\cdot)| \leq L$, (2) $0 < \gamma \leq f''_y(\cdot) \leq \beta$, (3) $\|x\| \leq R$ and (4), S' are ξ-self correlated. Let w_t, w'_t be the outputs of SGD on twin datasets S, S', $\Delta_t := w_t - w'_t$ and $\alpha \leq \frac{1}{\beta}$ is the step size of SGD. Then,*

$$\mathbb{E}\|\Delta_T\| \leq \frac{4LR}{\xi\gamma n} \quad ; \varepsilon_{stab} \leq \frac{4L^2R^2}{\xi\gamma n}$$

Proof: For simplicity we omit the dependence of f on y_j so that $f_{y_j}(w^\top x_j) = f(w, z_j)$. Note that the gradient of the loss function is $\nabla f_{y_j}(w_t^\top x_j) = f'_{y_j}(w_t^\top x_j)x_j$ and the Hessian is $\nabla^2 f_{y_j}(w_t^\top x_j) = f''_{y_j}(w_t^\top x_j)x_j x_j^\top$. The stochastic gradient step of $f_{y_j}(w_t^\top x_j)$ is

$$w_{t+1} = w_t - \alpha_t f'_{y_j}(w_t^\top x_j)x_j.$$

The dynamics of the divergence can be described as:

$$\mathbb{E}_{1:t+1}\|\Delta_{t+1}\| = \mathbb{E}_{1:t}\|\frac{1}{n}\sum_{j\neq i}\Delta_t - \alpha_t[f'_{y_j}y_j(w_t^\top x_j) - f'_{y_j}(w_t'^\top x_j)]x_j$$
$$\frac{1}{n}(\Delta_t - \alpha_t[f'_{y_i}(w_t^\top x_i)x_i - f'_{y'_i}(w_t'^\top x_i)x'_i])\| \tag{8}$$

Note that $[f'_{y_j} y_j(w_t^\top x_j) - f'_{y_j}(w_t'^\top x_j)]x_j$ can be rewritten as $f''_{y_j}(w_t^{\theta_j \top} x_j)x_j x_j^\top \Delta_t$ where $w^{\theta_j} = (1-\theta_j)w_t + \theta_j w_t', 0 < \theta_j < 1$. Similarly we can also rewrite $f'_{y_i}(w_t^\top x_i)x_i - f'_{y_i'}(w_t'^\top x_i)x_i'$ as

$$
\begin{aligned}
&f'_{y_i}(w_t^\top x_i)x_i - f'_{y_i'}(w_t'^\top x_i)x_i' \\
&= \frac{1}{2}\{f'_{y_i}(w_t^\top x_i)x_i - f'_{y_i}(w_t'^\top x_i)x_i\} \\
&\quad + \frac{1}{2}\{f'_{y_i'}(w_t^\top x_i')x_i' - f_{y_i'}(w_t'^\top x_i')x_i'\} \\
&\quad + \frac{1}{2}\{f'_{y_i}(w_t'^\top x_i) + f'_{y_i}(w_t^\top x_i)\}x_i \\
&\quad - \frac{1}{2}\{f'_{y_i'}(w_t^\top x_i') + f_{y_i'}(w_t'^\top x_i')\}x_i' \\
&= \frac{1}{2}f''_{y_i}(w_t^{\theta_i \top} x_i)x_i x_i^\top \Delta_t + \frac{1}{2}f''_{y_i'}(w_t^{\theta'_i \top} x_i')x_i' x_i'^\top \Delta_t \\
&\quad + \frac{1}{2}\{f'_{y_i}(w_t'^\top x_i) + f'_{y_i}(w_t^\top x_i)\}x_i \\
&\quad - \frac{1}{2}\{f'_{y_i'}(w_t^\top x_i') + f_{y_i'}(w_t'^\top x_i')\}x_i'
\end{aligned}
\tag{9}
$$

Thus equation equation 8 can be written as

$$
\begin{aligned}
\mathbb{E}_{1:t+1}\|\Delta_{t+1}\| = \mathbb{E}_{1:t}\|(I - \frac{\alpha_t}{2n}\sum_{j=1}^n f''_{y_j}(w_t^{\theta_i \top} x_i)x_i x_i^\top)\Delta_t \\
+ \frac{\alpha_t}{2n}\{f'_{y_i}(w_t'^\top x_i) + f'_{y_i}(w_t^\top x_i)\}x_i \\
- \frac{\alpha_t}{2n}\{f'_{y_i'}(w_t^\top x_i') + f_{y_i'}(w_t'^\top x_i')\}x_i'\|
\end{aligned}
$$

By the ξ-self correlated assumption and letting $\mathcal{H} = \frac{1}{n}\sum_{x_j \in S} x_j x_j^\top$ and $\mathcal{H}' = \frac{1}{n}\sum_{x_j \in S'} x_j x_j^\top$, we have:

$$
\begin{aligned}
\mathbb{E}\|\Delta_{t+1}\| &\le \mathbb{E}\|(I - \frac{\alpha_t \gamma}{2}\{\mathcal{H}+\mathcal{H}'\})\Delta_t\| + \frac{\alpha_t L}{n}\|x_i\| + \frac{\alpha_t L}{n}\|x_i'\| \\
&\le (1 - \frac{\alpha_t \xi \gamma}{2})\mathbb{E}\|\Delta_t\| + \frac{2\alpha_t L R}{n}.
\end{aligned}
\tag{10}
$$

The first inequality follows from the fact that $f''(\cdot) \ge \gamma$ and $x_j x_j^\top$s are all PSD. The second inequality follows from the fact that $\Delta_t \in Span\{x_1, .., x_n\}$. Fix $\alpha_t = \alpha$ and the theorem follows. $\qquad\square$

Lemma 3. *Suppose $x_{t_0} = 0$, $x_{t+1} = (1 + \frac{a}{0.99t})x_t + \frac{y}{t}$, we have $x_T \ge y(\frac{T}{t_0})^a$ if $a > 0$ is a sufficiently small constant.*

Proof: In the proof we use following inequality:

$$
e^a \le 1 + \frac{a}{0.99} \le e^{\frac{a}{0.99}}
$$

where $a > 0$ is a sufficiently small constant.

$$
\begin{aligned}
x_T &= \sum_{t=t_0+1}^{T} \frac{y}{t} \prod_{s=t+1}^{T} \left(1 + \frac{a}{0.99s}\right) \\
&\geq \sum_{t=t_0+1}^{T} \frac{y}{t} \exp\left(a \sum_{s=t+1}^{T} \frac{1}{s}\right) \\
&\geq \sum_{t=t_0+1}^{T} \frac{y}{t} \exp\left(a \log(T/t)\right) \\
&\geq y T^a \sum_{t=t_0+1}^{T} \frac{1}{t^{1+a}} \\
&\geq y \left(\frac{T}{t_0}\right)^a
\end{aligned}
\tag{11}
$$

$\square$

Lemma 4. *Let w_t, w_t' be the outputs of SGD on twin datasets S, S', and $\Delta_t := w_t - w_t'$. There exists a function f which is non-convex and β-smooth, twin datasets S, S' and constants a such that the divergence of SGD using step size $\alpha_t = \frac{a}{0.99\beta t}$ satisfies:*

$$
\mathbb{E}\|\Delta_T\| \geq \frac{1}{2n}\left(\frac{T}{t_0}\right)^a
\tag{12}
$$

Proof: For simplicity we set $x_i = v, y_i = 0.5, x_i' = -v, y_i' = 0.5$. We set S/x_i and S/x_i' to lie in the range of A. $x_j^\top A x_j > 0$. And we set $v^\top A v = -\beta$. We set $\alpha = a/(0.99\beta)$ where a is a sufficiently small constant. This construction yields:

$$
\mathbb{E}\|\Delta_{t+1}\| = (1 + \frac{1}{t})\mathbb{E}\|\Delta_t\| + \frac{\alpha_t}{n}\|v\|.
\tag{13}
$$

The theorem follows from Lemma 3. $\square$

Theorem 3. *Let w_t, w_t' be the outputs of SGD on twin datasets S, S', and $\Delta_t := w_t - w_t'$. There exists a function f which is non-convex and β-smooth, twin datasets S, S' and constants a such that the divergence of SGD after T rounds $(T > n)$ using constant step size $\alpha_t = \frac{a}{0.99\beta t}$ satisfies:*

$$
\mathbb{E}\|\Delta_T\| \geq \frac{T^a}{3n^{1+a}}
\tag{14}
$$

Proof: The proof is based on Theorem 4 plus the idea of a "burn-in" period. We have:

$$
\begin{aligned}
\mathbb{E}\|\Delta_T\| &= \mathbb{E}[\|w_t - w_t'\| | \Delta_n = 0]\mathbb{P}[\Delta_n = 0] + \mathbb{E}[\|w_t - w_t'\| | \Delta_n \neq 0]\mathbb{P}[\Delta_n \neq 0] \\
&\geq \mathbb{E}[\|w_t - w_t'\| | \Delta_n \neq 0]\mathbb{P}[\Delta_n \neq 0] \\
&= \left(1 - \left(1 - \frac{1}{n}\right)^n\right) \frac{T^a}{n^{1+a}}\|x_i - x_i'\| \\
&\geq \frac{T^a}{3n^{1+a}}\|x_i - x_i'\|
\end{aligned}
\tag{15}
$$

$\square$

Lemma 5 ((Hardt et al., 2016)). *Assume f is β-smooth and L-lipschitz. Let w_t, w_t' be outputs of SGD on twin datasets S, S' respectively after t iterations and let $\Delta_t := [w_t - w_t']$ and $\delta_t = \mathbb{E}\|\Delta_t\|$. Running SGD on $f(w; S)$ with step size $\alpha_t = \frac{a}{\beta t}$ satisfies the following conditions:*

- *The SGD update rule is a $(1 + \alpha_t\beta)$-expander and $2\alpha_t L$-bounded.*

- $\mathbb{E}[\|\Delta_t\| | \Delta_{t-1}] \leq (1 + \alpha_t\delta)\|\Delta_{t-1}\| + \frac{2\alpha_t L}{n}$.

- $\mathbb{E}[\|\Delta_T\| | \Delta_{t_k} = 0] \leq \left(\frac{T}{t_{k-1}}\right)^a \frac{2L}{n}$.

Lemma 6. *Let w_t, w'_t be outputs of SGD on twin datasets S, S' respectively after t iterations and let $\Delta_t := w_t - w'_t$. Suppose that $t_k = ct_{k-1}$. Then the following conditions hold:*

- $\mathbb{P}[\Delta_{t_{k-1}} = 0 | \Delta_{t_k} \neq 0] \leq \frac{n}{n + t_{k-1}}$.

- $\mathbb{P}[\Delta_{t_{k-1}} \neq 0 | \Delta_{t_k} \neq 0] \leq \frac{t_{k-1}}{t_k}\left(1 + \frac{t_k}{n}\right)$.

- $\mathbb{E}[\|\Delta_T\| | \Delta_{t_k} \neq 0] \leq \mathbb{E}[\|\Delta_T\| | \Delta_{t_{k-1}} \neq 0]\frac{t_{k-1}}{t_k}\left(1 + \frac{t_k}{n}\right) + \left(\frac{T}{t_{k-1}}\right)^a \frac{2L}{n}$.

Proof. In the proof we will use the following inequality with $r \geq 1$:

$$\frac{n-r}{n} \leq (1 - \frac{1}{n})^r \leq \frac{n}{n-r}$$

i):

$$\begin{aligned}
&\mathbb{P}[\Delta_{t_{k-1}} = 0 | \Delta_{t_k} \neq 0] \\
&= \frac{\mathbb{P}[\Delta_{t_{k-1}} = 0, \Delta_{t_k} \neq 0]}{\mathbb{P}[\Delta_{t_k} \neq 0]} \\
&= \frac{(1 - 1/n)^{t_{k-1}}[1 - (1 - 1/n)^{t_k - t_{k-1}}]}{1 - (1 - 1/n)^{t_k}} \\
&= 1 - \frac{1 - (1 - 1/n)^{t_{k-1}}}{1 - (1 - 1/n)^{t_k}} \\
&\leq (1 - 1/n)^{t_{k-1}} \leq \frac{n}{n + t_{k-1}}
\end{aligned} \tag{16}$$

ii):

$$\begin{aligned}
&\mathbb{P}[\Delta_{t_{k-1}} \neq 0 | \Delta_{t_k} \neq 0] \\
&= \frac{\mathbb{P}[\Delta_{t_k} \neq 0, \Delta_{t_{k-1}} \neq 0]}{\mathbb{P}[\Delta_{t_k} \neq 0]} \\
&= \frac{\mathbb{P}[\Delta_{t_{k-1}} \neq 0]}{\mathbb{P}[\Delta_{t_k} \neq 0]} \\
&= \frac{1 - (1 - 1/n)^{t_{k-1}}}{1 - (1 - 1/n)^{t_k}} \\
&\leq \frac{t_{k-1}}{t_k}(1 + \frac{t_k}{n})
\end{aligned} \tag{17}$$

iii):
By applying i) and ii) in the decomposition of $\mathbb{E}[\Delta_T | \Delta_{t_k} \neq 0]$ we have

$$\begin{aligned}
\mathbb{E}[\|\Delta_T\| | \Delta_{t_k} \neq 0] &\leq \mathbb{E}[\|\Delta_T\| | \Delta_{t_{k-1}} \neq 0]\mathbb{P}[\Delta_{t_{k-1}} \neq 0 | \Delta_{t_k} \neq 0] \\
&\quad + \mathbb{E}[\|\Delta_T\| | \Delta_{t_{k-1}} = 0]\mathbb{P}[\Delta_{t_{k-1}} = 0 | \Delta_{t_k} \neq 0] \\
&\leq \mathbb{E}[\|\Delta_T\| | \Delta_{t_{k-1}} \neq 0]\frac{t_{k-1}}{t_k}(1 + \frac{t_k}{n}) \\
&\quad + \left(\frac{T}{t_{k-1}}\right)^a \frac{2L}{n + t_{k-1}}
\end{aligned} \tag{18}$$

where the last inequality uses the fact that $\mathbb{E}[\|\Delta_T\| | \Delta_{t_k} = 0] \leq \left(\frac{T}{t_{k-1}}\right)^a \frac{2L}{n}$. $\qquad\square$

Theorem 4 ((Permutation)). *Assume f is β-smooth and L-lipschitz. Running T ($T > n$) iterations of SGD on $f(w; S)$ with step size $\alpha_t = \frac{a}{\beta n}$, the stability of SGD satisfies:*

$$\mathbb{E}\|\Delta_T\| \leq \frac{2LT^a}{n^{1+a}}, \varepsilon_{stab} \leq \frac{2L^2 T^a}{n^{1+a}} \tag{19}$$

Proof. Let $H = t$ represents the event that the first time the SGD pick the different entry is at time t:

$$
\begin{aligned}
\mathbb{E}\|\Delta_T\| &\leq (\mathbb{E}\|\Delta_n\|) \left(\frac{T}{n}\right)^a + \frac{LT^a}{n^{1+a}} \\
&\leq \frac{1}{n} \sum_{t=1}^{n} \mathbb{E}[\|\Delta_n\| | H = t] \left(\frac{n}{t}\right)^a + \frac{LT^a}{n^{1+a}} \\
&\leq \frac{1}{n} \int_{t=1}^{n} \frac{n^a}{t^{1+a}} \left(\frac{T}{n}\right)^a + \frac{LT^a}{n^{1+a}} \\
&\leq \frac{2LT^a}{n^{1+a}}
\end{aligned}
\tag{20}
$$

$\square$

Theorem4b (Uniformly Independent) Assume f is β-smooth and L-lipschitz. Running T $(T > n)$ iterations of SGD on $f(w; S)$ with step size $\alpha_t = \frac{a}{\beta n}$, the stability of SGD satisfies:

$$
\mathbb{E}\|\Delta_T\| \leq 16 \log(n) L \frac{T^a}{n^{1+a}}; \quad \varepsilon_{stab} \leq 16 \log(n) L^2 \frac{T^a}{n^{1+a}}
\tag{21}
$$

Proof. We first decompose Δ_T as follows by selecting $t_k = n$:

$$
\mathbb{E}\|\Delta_T\| = \underbrace{\mathbb{E}[\|\Delta_T\| | \Delta_{t_k} = 0] \mathbb{P}[\Delta_{t_k} = 0]}_{\text{Term 1} \leq \frac{2LT^a}{2n^{1+a}} \text{ (Lemma 5)}} + \underbrace{\mathbb{E}[\|\Delta_T\| | \Delta_{t_k} \neq 0] \mathbb{P}[\Delta_{t_k} \neq 0]}_{\text{Term 2} \leq \frac{30L \log(n) T^a}{n^{1+a}}}
\tag{22}
$$

Term 1 is easily bounded by applying Lemma 5 with $\alpha_t = \frac{a}{t\beta}$. To bound Term 2, plug in $\mathbb{P}[\Delta_{t_k} \neq 0] = 1 - (1 - 1/n)^{t_k} \leq \frac{t_k}{n}$ and recursively apply point (iii) from Lemma 6 by setting $t_{i+1} = ct_i$. We get:

$$
\begin{aligned}
&\mathbb{E}[\|\Delta_T\| | \Delta_{t_k} \neq 0] \mathbb{P}[\Delta_{t_k} \neq 0] \\
&\leq \frac{2L}{n} \frac{t_k}{n} \sum_{i=1}^{k-1} \left(\frac{T}{t_i}\right)^a \frac{n}{n + t_i} \prod_{\tau=i+1}^{k-1} \left(1 + \frac{t_{\tau+1}}{n}\right) \frac{t_\tau}{t_{\tau+1}} \\
&\leq \frac{2L}{n} \sum_{i=1}^{k-1} \left(\frac{T}{t_i}\right)^a \frac{t_{i+1}}{n + t_i} exp\left(\sum_{\tau=i+1}^{k-1} \frac{t_{\tau+1}}{n}\right) \\
&\leq \frac{16L}{n} \sum_{i=1}^{k-1} \left(\frac{T}{t_i}\right)^a \frac{t_i}{n + t_i} \\
&\leq \frac{8kLT^a}{n^{1+a}}
\end{aligned}
\tag{23}
$$

In the second and third inequality we use the fact that $1 + x \leq \exp(x)$ and $t_{i+1} = ct_i$ to get $\prod_{\tau=i+1}^{k-1} (1 + \frac{t_{\tau+1}}{n}) \leq exp(\sum_{\tau=i+1}^{k-1} \frac{t_{\tau+1}}{n})$ and $\exp(\frac{t_k}{n}) \leq 3$. In the last inequality, we use the fact that $t_i \leq t_k = n$. Indeed, we can conclude that $k = O(\log(n))$ as $t_{i+1} = ct_i, t_k = n$.

$\square$

Theorem 5. *Let w_t, w_t' be the outputs of SGD on twin datasets S, S', and let $\Delta_t := w_t - w_t'$. There exists a function f which is non-convex and β-smooth, twin sets S, S' and constants a, γ such that the divergence of SGD after T rounds $(T > n)$ using constant step size $\alpha = \frac{a}{0.99\gamma}$ satisfies:*

$$
\mathbb{E}\|\Delta_T\| \geq \frac{1}{n^2} e^{aT/2}
\tag{24}
$$

Proof: The proof is similar to Theorem 3. Since $\Delta_t \in Span\{x_i - x_i'\}$, we have:

$$
\mathbb{E}\|\Delta_{t+1}\| \geq (1 - \frac{1}{n})(1 + \alpha_t \beta)\mathbb{E}\|\Delta_t\| + \frac{\alpha_t}{n}\|x_i - x_i\|
$$

Suppose t_0 is the hitting time when $\|\Delta_{t_0}\| > 0$ and $\|\Delta_{t_0-1}\| = 0$, $\|\Delta_T\| \geq \frac{\|x_i - x_i'\|}{3n} e^{a(T-t_0)/2}$.

$$
\begin{aligned}
\mathbb{E}\|\Delta_T\| &= \mathbb{E}[\|w_t - w_t'\| \,|\, \Delta_1 = 0]\mathbb{P}[\Delta_1 = 0] + \mathbb{E}[\|w_t - w_t'\| \,|\, \Delta_1 \neq 0]\mathbb{P}[\Delta_1 \neq 0] \\
&\geq \mathbb{E}[\|w_t - w_t'\| \,|\, \Delta_1 \neq 0]\mathbb{P}[\Delta_1 \neq 0] \\
&= \frac{1}{n}\left(\frac{\|x_i - x_i'\|}{n} e^{aT/2}\right) \\
&= \frac{\|x_i - x_i'\|}{n^2} e^{aT/2}.
\end{aligned}
\tag{25}
$$

$\square$

Theorem 6. *Given a set S and a β-smooth and L-Lipschitz loss funtion f, suppose SGD with a fixed step size $\alpha \leq \frac{\delta}{2L}$ reaches a stationary point w_0 with $\|\nabla_w f(w_0, S)\| \leq \frac{\gamma\delta}{2}$, which is close to a local minimum w^* in a (δ, γ)-Hessian contractive region: $\|w_0 - w^*\| \leq \frac{\delta}{2}$, we have $\forall T \geq 1$, $\|w_T - w^*\| \leq \delta$.*

Proof: Let $\mathcal{H}_{w_t, w^*, \theta} = \frac{1}{n}\sum_{z_j \in S}\nabla_w^2 f(\theta w_t + (1-\theta)w_0, z_j)$ We derive that:

$$
\begin{aligned}
&\mathbb{E}[\|w_{t+1} - w_0\|^2] \\
&= \mathbb{E}[\|w_t - w_0 - \alpha\nabla_w f(w, z)\|^2] \\
&= \mathbb{E}[\|w_t - w_0\|^2 + \alpha^2\|\nabla_w f(w_s, z)\|^2 - 2\alpha \underbrace{\mathbb{E}\{\nabla_w f(w, z)^\top[w_t - w_0]]\}}_{\mathbb{E}_z \nabla_w f(w,z)^\top[w_t - w_0] = \nabla_w f(w,S)^\top[w_t - w_0]} \\
&= \mathbb{E}[\|w_t - w_0\|^2 + \alpha^2\|\nabla_w f(w_s, z)\|^2 - 2\alpha\nabla_w f(w_0, S)^\top[w_t - w_0] \\
&\quad - 2\alpha\nabla_w[f(w_t, S) - \nabla_w f(w_0, S)]^\top[w_t - w_0] \\
&\leq \mathbb{E}[\|w_t - w_0\|^2 + \alpha^2 L^2 - 2\alpha[w_t - w_0]^\top\mathcal{H}_{w_t, w^*, \theta}[w_t - w_0]] + \alpha\gamma\delta^2
\end{aligned}
\tag{26}
$$

Due to the fact that $w_t - w_0 \in Span\{\nabla_w f(w_0, z), ..., \nabla_w f(w_{t-1}, z)\}$, we have

$$
\begin{aligned}
&\mathbb{E}[\|w_t - w_0\|^2 + \alpha^2 L^2 - 2\alpha[w_t - w_0]^\top\mathcal{H}_{w_t, w^*, \theta}[w_t - w_0]] + \alpha\gamma\delta^2 \\
&\leq (1 - 2\alpha\gamma)\|w_t - w_0\|^2 + \alpha^2 L^2 + \alpha\gamma\delta^2
\end{aligned}
\tag{27}
$$

This implies $\|w_t - w_0\| \leq \delta$ for all t. $\qquad\square$

Observation 1. *Suppose loss function $f(w; S)$ satisfies condition in Theorem 2 so that $f(w, S) = \frac{1}{n}\sum_{j=1}^n f_{y_j}(w^\top x_j)$ satisfies (1) $|f_y'(\cdot)| \leq L$, (2) $\gamma \leq f_y''(\cdot) \leq \beta$, (3) $r_i\|x\| \leq R$ and (4) S is ξ-self correlated, we have $f(w; S)$ is globally $\frac{\xi\gamma}{R^2}$ Hessian Contractive.*

Proof. The gradient of $f(w, z_j)$ is

$$
\nabla f(w, z_j) = f'(w^\top x_j)x_j
$$

and Hessian is

$$
\nabla^2 f(w, z_j) = f''(w^\top x_j)x_j x_j^\top
$$

. For all w, z_j,

$$
\begin{aligned}
&\nabla f(w, z_j)^\top \mathcal{H}_{w'}\nabla f(w, z_j) \\
&= (f'(w^\top x_j))^2 x_j^\top\left[\frac{1}{n}\sum_{j=1}^n f''(w'^\top x_j)x_j x_j^\top\right]x_j \\
&\geq (f'(w^\top x_j))^2\xi\gamma \geq \frac{\xi\gamma}{R^2}\|\nabla f(w, z_j)\|^2
\end{aligned}
\tag{28}
$$

. $\qquad\square$

B   EMPIRICAL SUPPORT FOR THEOREM 6

In order to empirically support Theorem 6, we ran the follow experiment: We started with a fully-trained model w^*, and generated random perturbations $\tilde{w} = w^* + \nu$ where $\nu \sim \mathcal{N}(0, \sigma^2)$. We then trained each $\tilde{w}$ until convergence $\tilde{w}^*$, and computed the normalized distance $\|\tilde{w}^* - w^*\|/\|w^*\|$. To match Theorem 6, this distance should be no more than $2\|\tilde{w}^* - w^*\|/\|w^*\|$, i.e., the trained weights $\tilde{w}^*$ after perturbing stay close to w^*, as in the conclusion of Theorem 6.

For the above experiment, we first trained both ResNet18 and Densenet121 for 200 epochs, with constant learning rate of 0.01, a momentum of 0.9, and batch size of 128. We generated 20 Gaussian perturbations $\tilde{w}$ of the trained model weights w^* and trained each for 50 more epochs, resulting in weights $\tilde{w}^*$. We used standard deviation $\sigma \in \{0.01, 0.005\}$ for the Gaussian perturbations. We then computed the normalized differences as explained in the previous paragraph. The results are shown in Table 2. Inspecting the table shows that the distances between the trained weights after perturbing are quite close to the distances before further training, with low variance. This indicates that the weights after training the perturbed model are well within the desired range as specified in Theorem 6.

		$\|\tilde{w} - w^*\|/\|w^*\|$	$\|\tilde{w}^* - w^*\|/\|w^*\|$
ResNet	$\sigma = 0.01$	0.549±0.00009	0.557±0.0003
	$\sigma = 0.005$	0.274±0.00005	0.277±0.007
DenseNet	$\sigma = 0.01$	0.123±0.00003	0.126±0.0004
	$\sigma = 0.005$	0.062±0.00001	0.065±0.003

Table 2: Pertubed SGD Results