On the symmetries of the synchronization problem in Cryo-EM: Multi-Frequency Vector Diffusion Maps on the Projective Plane

Abstract

Cryo-Electron Microscopy (Cryo-EM) is an important imaging method which allows high-resolution reconstruction of the 3D structures of biomolecules. It produces highly noisy 2D images by projecting a molecule's 3D density from random viewing directions. Because the projection directions are *unknown*, estimating the images' poses is necessary to perform the reconstruction. We focus on this task and study it under the *group synchronization* framework: if the relative poses of pairs of images can be approximated from the data, an estimation of the images' poses is given by the assignment which is most consistent with the relative ones. In particular, by studying the symmetries of cryo-EM, we show that relative poses in the group $O(2)$ provide sufficient constraints to identify the images' poses, up to the molecule's chirality. With this in mind, we improve the existing multi-frequency vector diffusion maps (MFVDM) method: by using $O(2)$ relative poses, our method not only predicts the similarity between the images' viewing directions but also recovers their poses. Hence, we can leverage all input images in a 3D reconstruction algorithm by initializing the poses with our estimation rather than just clustering and averaging the input images. We validate the recovery capabilities and robustness of our method on randomly generated synchronization graphs and a synthetic cryo-EM dataset.

1   Introduction

Cryo-Electron Microscopy (Cryo-EM) revolutionized the field of structural biology, enabling the reconstruction of protein structures at unprecedented resolutions. Indeed, the 2017 Nobel Prize in Chemistry was awarded to three scientists for their pioneering works on it [1]. In single-particle cryo-EM, a purified solution containing the molecule of interest is frozen on a thin film and then bombarded with electrons to obtain a 2D tomographic (integral) projection of it. The resulting image contains the projection of each copy of the molecule in the solution; in a *particle picking* phase, these projections are cropped to obtain a dataset of 2D images. Since each copy in the solution is randomly rotated in 3D, each image is a projection of the molecule's density in a random *unknown* pose. The objective is reconstructing the molecule's 3D structure from these observations. However, the produced images are characterized by very low signal-to-noise ratios (SNR) [2], e.g. see Fig. 1; the high noise and the unknown poses make this inverse problem particularly challenging.

Common methods attempt to estimate the images' poses while performing the reconstruction. This is typically done via iterative refinement with an Expectation-Maximization (EM) algorithm [3, 4]. Unfortunately, EM-like methods are known to suffer from convergence issues due to local minima and generally require a sufficiently good initialization and restarts

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

[5]. The 3D density of known molecules can be used as initial guess but this often causes *model bias* [6]. *Ab-initio* methods, which don't rely on an initial guess, are particularly useful to initialize the iterative methods above and, therefore, are a fundamental element of the reconstruction pipeline. Moreover, a good initialization of the poses can speed up iterative refinement by providing a non-flat prior over poses, and so, reducing the search space.

Figure 1: Example of Cryo-EM images at different SNR values.

Due to the high level of noise, a first phase of *2D classification and averaging* is performed, where a smaller denoised set of images is generated by clustering images with a similar viewing direction and averaging images within each cluster. Because this blurs out the fine details, it can reduce the final reconstruction's resolution and, therefore, it is desirable to minimize the averaging performed [7, 5]. Vector Diffusion Maps (VDM), and their variants, [8, 9, 10] are powerful methods to estimate the similarity between images' viewing directions and, therefore, provide a robust metric for clustering. In short, these methods leverage the *equivariance* of Cryo-EM to planar rotations (SO(2)) and interpret the planar rotations aligning pairs of images as an approximation of the local parallel transport over a sphere of viewing directions. The closer the viewing directions of two images are, the more consistent is the transport along different paths between them. This notion of consistency is encoded in the top eigenvectors of an Hermitian matrix representing the discretized parallel transport operator.

Contributions and Outline   In this work, we consider the problem of estimating Cryo-EM images' poses under the *group synchronization* framework, i.e. using the relative poses of pairs of images to recover an estimation of the absolute poses. First, in Sec. 2, we relate the symmetries of Cryo-EM with those of the synchronization task: we show that SO(2) *relative poses are not sufficient* to solve the multi-view synchronization problem, *but* O(2) *relative poses are*. Second, in Sec. 3, we use this insight to construct an improved VDM method, which is able to recover not only a similarity metric between the images, but also an estimation of each image's pose. In particular, we extend VDM by *i)* using vector diffusion over a *projective plane* (rather than a sphere) to reconstruct denoised similarity and parallel transport and by *ii)* including a *final synchronization* step to recover the absolute poses. This minor modification of VDM allows one to replace the 2D classification phase with a pose estimation phase and, therefore, leverage all the raw cryo-EM images in the final 3D reconstruction without the loss of resolution caused by averaging. Additionally, our improved VDM method shares the same spectral properties of the original VDM method and, hence, the same noise stability and recovery guarantees. Finally, we validate experimentally our theoretical results in Sec. 5.

1.1   The mathematics of Cryo-EM

Following [11], we consider the problem of single-particle homogeneous reconstruction and assume no image translations for simplicity. Then, a molecule's 3D density is a function $\Psi : \mathbb{R}^3 \to \mathbb{R} \in L^2(\mathbb{R}^3)$ with compact support around the origin of $\mathbb{R}^3$. An observation is a gray-scale image generated by the *integral projection* along the Z axis $\Pi : L^2(\mathbb{R}^3) \to L^2(\mathbb{R}^2)$, which is defined as:

$$\Pi(\Psi)(x, y) = \int_z \Psi(x, y, z)\mathrm{d}z \quad x, y, z \in \mathbb{R} .$$

Let $\{o_i\}_{i=1}^N$ be the collection of images observed. The image $o_i := \Pi(g_i^{-1}.\Psi) \in L^2(\mathbb{R}^2)$ is the projection of a copy of the molecule Ψ rotated by a random $g_i^{-1} \in \mathrm{SO}(3)$, where the action of $\mathrm{SO}(3)$ on $L^2(\mathbb{R}^3)$ is the standard action:

$$[g.\Psi](\boldsymbol{x}) := \Psi(g^{-1}.\boldsymbol{x}) \quad \forall g \in \mathrm{SO}(3) .$$

An element $g_i \in \mathrm{SO}(3)$ is identified with a real orthonormal matrix $(\boldsymbol{x}_i, \boldsymbol{y}_i, \boldsymbol{z}_i) \in \mathbb{R}^{3 \times 3}$ with positive determinant. The three columns $\boldsymbol{x}_i, \boldsymbol{y}_i, \boldsymbol{z}_i \in \mathbb{R}^3$ form an orthonormal basis for $\mathbb{R}^3$. We can write:

$$o_i(x, y) = \left[\Pi(g_i^{-1}\Psi)\right](x, y) = \int_z \Psi(x\boldsymbol{x}_i + y\boldsymbol{y}_i + z\boldsymbol{z}_i)\mathrm{d}z . \tag{1}$$

In particular, if $g_i = (\boldsymbol{x}_i, \boldsymbol{y}_i, \boldsymbol{z}_i)$, then $\boldsymbol{z}_i \in \mathbb{R}^3$ is the viewing direction along which Ψ is projected, while $(\boldsymbol{x}_i, \boldsymbol{y}_i)$ defines the camera rotation around $\boldsymbol{z}_i$. We define the *projection map* $\pi(g_i) = \boldsymbol{z}_i \in \mathbb{R}^3$.

Figure 2: Generation process of Cryo-EM images. Projections from similar or opposite directions are related by elements of O(2), see also Fig. 3.

Figure 3: *First row*: images sharing a similar viewing direction with the top-left image. *Second row*: images with opposite viewing direction with respect to the top-left image. The colored group elements are the relative transformation with respect the top-left image; e is the identity, r is a rotation by $\frac{2\pi}{5}$ around the viewing axis Z and f mirrors along the X axis.

2 On the symmetries of the Cryo-EM synchronization problem

One of the main challenges in cryo-EM is the fact that the pose $g_i \in \mathrm{SO}(3)$ of each observation o_i is unknown. This prevents one to glue the observations together to reconstruct the density Ψ. However, the set of observations $\{o_i\}_i$ can be used to estimate the relative poses $\{\widehat{g}_{ij} \approx g_{ij} = g_j^{-1} g_i\}_{ij}$. A synchronization problem is defined by a "synchronization graph" $\mathcal{G} = (\mathcal{V}, \mathcal{E})$, i.e. a graph with a node in $\mathcal{V}$ per image and an edge $(i, j) \in \mathcal{E}$ associated with $\widehat{g}_{ij}$ if its estimation is available. Given a synchronization graph $\mathcal{G}$, the goal is now to "synchronize" them, i.e. find a global assignment of the poses $\{\widehat{g}_i\}_i$ consistent with the estimated relative poses:

$$\widehat{g}_j \approx \widehat{g}_i \widehat{g}_{ij}^{-1} \quad \forall (i, j) \in \mathcal{G} . \tag{2}$$

Global $\mathrm{SO}(3)$ **ambiguity** If $\{\widehat{g}_i\}_i$ is a solution, so is $\{g\widehat{g}_i\}_i$ for any $g \in \mathrm{SO}(3)$. This ambiguity is not unexpected and corresponds to the fact that any rotated molecule $g^{-1}.\Psi$ is an equally valid reconstruction, i.e. cryo-EM can not recover the orientation of a molecule.

2.1 Relative poses in $\mathrm{SO}(2)$: the impossibility of synchronization

Before solving the synchronization problem, it is necessary to estimate the relative poses $\{g_{ij} = g_j^{-1} g_i\}_{ij}$. Unfortunately, since the projection Π loses information about the 3D structure, it is not possible to estimate the relative pose of any pair of images. However, when the viewing directions $\pi(g_i) = \boldsymbol{z}_i$ and $\pi(g_j) = \boldsymbol{z}_j$ of two images are sufficiently close, they generally differ only by a planar rotation $r_{ij} \in \mathrm{SO}(2)$ which can be directly estimated as

$$r_{ij} = \underset{r \in \mathrm{SO}(2)}{\arg\min} \|o_j - r.o_i\|_2^2 \text{ if } \pi(g_i) \approx \pi(g_j) . \tag{3}$$

where r acts on image o_i by rotating it. The closer $\boldsymbol{z}_i$ and $\boldsymbol{z}_j$ are, the better r_{ij} approximates g_{ij}. Indeed, if $\boldsymbol{z}_i = \boldsymbol{z}_j$, then $g_{ij} = g_j^{-1} g_i$ has form[1] $\begin{bmatrix} * & * & 0 \\ * & * & 0 \\ 0 & 0 & 1 \end{bmatrix}$, where the top-left 2×2 block is an $\mathrm{SO}(2)$ matrix. We identify all the rotations of this form as the subgroup[2] $\mathrm{SO}(2) \cong R_z < \mathrm{SO}(3)$ of planar rotations around the Z axis. The projection operator Π is $\mathrm{SO}(2)$-**equivariant**[3], i.e.:

$$\Pi(r.\Psi) = r.\Pi(\Psi) \quad \forall r \in R_z \cong \mathrm{SO}(2) . \tag{4}$$

Hence, if $g_j^{-1} = r g_i^{-1}$, it follows that $o_j = r.o_i$. For example, all projections generated by the top view in Fig 2 are related by a rotations r as shown in the top row of Fig. 3.

---

[1]This is shown using the orthogonality of g_i and g_j's columns.

[2]$\mathrm{SO}(3)$ has many subgroups isomorphic to $\mathrm{SO}(2)$ so we use R_z to indicate this particular one.

[3]For simplicity, the action of R_z on 3D densities or 2D images is not distinguished.

Global $\mathrm{SO}(2)$ **ambiguity** While this is a common approach in the literature, e.g. in VDM [8, 12, 10, 13], it does not provide sufficient information to estimate the poses $\{g_i\}_i$. Indeed, the estimated relative rotations $\{r_{ij} \in \mathrm{SO}(2)\}_{(ij) \in \mathcal{G}}$ yield the following constraints:

$$g_j \approx g_i r_{ij}^{-1} \quad \forall (i,j) \in \mathcal{G} \ .$$

Since $\mathrm{SO}(2)$ is *abelian*, if $\{\widehat{g}_i\}_i$ is a solution to this set of constraints, then $\{\widehat{g}_i r\}_i$ is an equally valid solution for any $r \in R_z \cong \mathrm{SO}(2)$:

$$\widehat{g}_j \approx \widehat{g}_i r_{ij}^{-1} \implies \widehat{g}_j r \approx \widehat{g}_i r_{ij}^{-1} r \implies \widehat{g}_j r \approx (\widehat{g}_i r) \, r_{ij}^{-1}$$

This implies that any method which estimates the poses $\{g_i\}_i$ from the estimated relative ones $\{r_{ij} \in \mathrm{SO}(2)\}_{ij}$ will only be able to recover them up to a global rotation $r \in R_z$. This global ambiguity prevents using these estimations to directly invert the linear projection Π and recover the original molecule Ψ, since it would result in averaging randomly rotated versions of the images.

Nevertheless, the ambiguous poses $\{g_i r\}_i$ still contain information about the viewing direction of the projections. Indeed, note that $\pi(g_i r) = \pi(g_i)$, i.e. the viewing directions are invariant to this global ambiguity and, therefore, can be recovered from this method. Previous works such as VDM [8, 12, 10] exploit this invariance to estimate the relative distance between different images. This distance is generally used for *class averaging* i.e. to cluster images and average images within each cluster to generate a smaller, de-noised, images dataset.

2.2 Relative poses in $\mathrm{O}(2)$: a sufficient condition for synchronization

In this work, we solve this global $\mathrm{SO}(2)$ ambiguity and directly estimate the absolute poses $\{g_i\}_i$ of the images. To do so, we will exploit another symmetry of Cryo-EM. Indeed, the tomographic projection Π is also invariant to *mirroring* along the Z axis $\mathrm{m}_z = \left[\begin{smallmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{smallmatrix}\right] \in \mathrm{O}(3)$ i.e.:

$$\Pi(\mathrm{m}_z \,.\, \Psi) = \int_z \Psi(x, y, -z) dz = \int_z \Psi(x, y, z) dz = \Pi(\Psi)$$

This additional symmetry turns out useful to break the global $\mathrm{SO}(2)$ symmetry explained earlier.

First, let $r_y = \left[\begin{smallmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{smallmatrix}\right] \in \mathrm{SO}(3)$ be a π rotation along the Y axis; note that $r_y = \mathrm{m}_z \, \mathrm{m}_x$, where m_x is a mirroring along the X axis. Then, let $\mathrm{f} = \left[\begin{smallmatrix} -1 & 0 \\ 0 & 1 \end{smallmatrix}\right] \in \mathrm{O}(2)$ be the flip of a planar image along the X axis. Then:

$$\Pi(r_y . \Psi) = \Pi(\mathrm{m}_z \, \mathrm{m}_x \,.\, \Psi) = \Pi(\mathrm{m}_x \,.\, \Psi) = \mathrm{f} \,.\, \Pi(\Psi) \ , \tag{5}$$

i.e. the projection operator Π is **flip equivariant**: projections along a direction ($\Pi(g_i^{-1}.\Psi)$) are related to projections from the opposite direction ($\mathrm{f} \,.\, \Pi(r_y g_i^{-1}.\Psi)$) by a planar reflection f. For example, all projections generated by the bottom view in Fig. 2 are related by a rotation r followed by the reflection f with the images generated from the top view, as shown in the bottom row of Fig. 3. To simplify the notation, we will also let r_y act on a image; this action should be intended as that of f.

That means that we can not only estimate the relative pose $g_{ij} \approx h_{ij} \in R_z$ when the viewing directions $\pi(g_i)$ and $\pi(g_j)$ are sufficiently close, but also $h_{ij} \in H \cong \mathrm{O}(2)$ when $\pi(g_i) \approx -\pi(g_j)$, i.e.:

$$h_{ij} = \underset{h \in H \cong \mathrm{O}(2)}{\arg\min} \|o_j - h.o_i\|_2^2 \quad \text{if } \pi(g_i) \approx \pm\pi(g_j) \ . \tag{6}$$

where $H \cong \mathrm{O}(2)$ is the subgroup[4] of $\mathrm{SO}(3)$ containing R_z (planar rotations along the Z axis) and r_y.

Global chirality ambiguity The relative poses $\{h_{ij} \in \mathrm{O}(2)\}_{ij}$ still don't fully constrain the absolute ones; if $\{\widehat{g}_i\}_i$ is a solution, so is $\{\widehat{g}_i r_z\}_i$, with $r_z \in R_z$ being a π-rotation around the Z axis[5]:

$$\widehat{g}_j \approx \widehat{g}_i h_{ij}^{-1} \implies \widehat{g}_j r_z \approx \widehat{g}_i h_{ij}^{-1} r_z \implies \widehat{g}_j r_z \approx (\widehat{g}_i r_z) \, h_{ij}^{-1}$$

---

[4] More precisely, $H := R_z \rtimes \{e, r_y\} \cong \mathrm{O}(2)$.

[5] One can verify that r_z commutes with any $r \in R_z$ and r_x.

Table 1: Summary of the Cryo-EM symmetries.

	description of the transformation	symmetry name
$o_i = \Pi(g_i^{-1}\Psi)$	$g_i \in \mathrm{SO}(3)$	generative process
$r.o_i = \Pi(rg_i^{-1}\Psi)$	$r \in \mathrm{SO}(2)$ any rotation around Z axis	$\mathrm{SO}(2)$ equivariance
$o_i = \Pi(\mathrm{m}_z\, g_i^{-1}\Psi)$	$\mathrm{m}_z \in \mathrm{O}(3)$ mirroring along Z axis	Mirroring (Z) invariance
$\mathrm{f}.o_i = \Pi(m_x g_i^{-1}\Psi)$	$\mathrm{m}_x \in \mathrm{O}(3), \mathrm{f} \in \mathrm{O}(2)$ mirroring along X axis	Mirroring (XY) equivariance
$\mathrm{f}.o_i = \Pi(r_y g_i^{-1}\Psi)$	$r_y = \mathrm{m}_x\, \mathrm{m}_z \in \mathrm{SO}(3)$ rotation by π around Y axis	Flip equivariance

This ambiguity is related to the well known problem that Cryo-EM can not recover the **chirality** of a molecule. To see why, let $\mathrm{i} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$ be the *inversion*, i.e. a mirroring along all axes; note that $\mathrm{m}_z = r_z\, \mathrm{i}$ and that i commutes with any element $g \in \mathrm{SO}(3)$. Then:

$$o_i = \Pi(g_i^{-1}.\Psi) = \Pi(\mathrm{m}_z\, g_i^{-1}.\Psi) = \Pi(r_z\, \mathrm{i}\, g_i^{-1}.\Psi) = \Pi(r_z g_i^{-1}.(\mathrm{i}\,.\Psi)) = \Pi((g_i r_z)^{-1}.(\mathrm{i}\,.\Psi))$$

i.e., o_i is equally likely to be generated by the molecule Ψ with pose g_i or its mirrored version $\mathrm{i}\,.\Psi$ with pose $g_i r_z$. Together with the global $\mathrm{SO}(3)$ symmetry introduced earlier, this means Cryo-EM intrinsically suffers from a **global** $\mathrm{O}(3)$ **ambiguity** which we can not hope to resolve. Nevertheless, since r_z is the *only* element of $\mathrm{SO}(3)$ commuting with all elements in $H \cong \mathrm{O}(2)$, we have that relative poses in H constrain the synchronization problem precisely up to this $\mathrm{O}(3)$ symmetry and, thus provide *sufficient* constraints to identify the images' poses. We provide a theoretical argument based on Vector Diffusion Maps (VDM) [9] supporting this informal claim in Apx. C.

3 Multi-Frequency Vector Diffusion Maps on the Projective Plane

In this section, we use the new insights on the symmetries of cryo-EM to enable the Vector Diffusion Map (VDM) method to recover the poses $\{g_i\}_i$. The key component is the Graph Connection Laplacian (GCL) of the synchronization graph $\mathcal{G}$ with the estimated relative poses $\{h_{ij} \in \mathrm{O}(2)\}_{ij}$ on its edges. Under certain conditions, its eigenvectors converge to the eigen-vector-fields of the connection Laplacian operator on the projective plane, which contain information about the absolute poses. Our method involves *two steps*: a *denoising step* (like the original VDM) and a *synchronization step*; Alg. 1 provides an overview of the proposed method. We study its computational complexity in Apx. F.

3.1 Quotient Spaces and Projective Plane

Each image o_i is uniquely identified by an element $g_i \in \mathrm{SO}(3)$ via Eq. 1, assuming the molecule has no symmetries. Two images o_i and o_j related by a rotation $r_{ij} \in R_z$ contain the same information and are considered equivalent. We can define a similar equivalence relation on their respective poses, i.e. $g_i \sim g_j = g_i r_{ij}$. This defines the quotient space $\mathrm{SO}(3)/R_z$ which is isomorphic to the *sphere* $\mathcal{S}^2$. An element $g_i \in \mathrm{SO}(3)$ is interpreted as a point $\pi(g_i) = z_i$ in $\mathcal{S}^2$ and a choice of *frame* on it, i.e. the choice of basis[6] (x_i, y_i) for the tangent space at z_i or, equivalently, the rotation of the camera around the axis z_i. Then, the relative rotation $r_{ij} \in R_z$ estimated in Eq. 3 is interpreted as the unique rotation such that $g_j r_{ij}^{-1}$ is the *parallel transport* of g_i along the shortest geodesic connecting $\pi(g_i)$ to $\pi(g_j)$ (this is unique unless $\pi(g_i) = -\pi(g_j)$). Here, we want to leverage the larger symmetry H. We define the new equivalence relation $g \sim gh$ for all $h \in H$ on $\mathrm{SO}(3)$, which determines the quotient space $\mathrm{SO}(3)/H$, isomorphic to the *projective plane*[7] $\mathrm{P}\mathbb{R}^2$. While it is harder to visualize, elements of $\mathrm{SO}(3)$ are now interpreted as points in $\mathrm{P}\mathbb{R}^2$ together with a choice of frame for their tangent spaces[8]. Finally, the relative pose $h_{ij} \in H$ from Eq. 6 is interpreted as the element such that $g_j h_{ij}^{-1}$ is the *parallel transport* of g_i along the shortest geodesic between the projections of g_i and g_j on $\mathrm{P}\mathbb{R}^2$.

3.2 Vector Diffusion Maps on $\mathrm{P}\mathbb{R}^2$

An initial synchronization graph $\mathcal{G}$ can be generated by comparing all pairs of images using Eq. 6 and preserving only the edges with smallest distances. Ideally, this graph is a discretization of

---

[6]Indeed, (x_i, y_i) span a 2D plane through the origin parallel to the tangent plane at z_i.

[7]The projective plane can be thought as a sphere where antipodal points are identified together.

[8]By moving from $\mathcal{S}^2$ to $\mathrm{P}\mathbb{R}^2$, we change the *structure group* and consider frames in $\mathrm{O}(2)$ rather than $\mathrm{SO}(2)$.

$\text{P}\mathbb{R}^2 \cong \text{SO}(3)/H$ with local connectivity, but the noise on the images negatively affects the estimated distances and relative poses, introducing "shortcut" edges in this graph. Therefore, the first step of our method is re-estimating the geodesic distances and the relative poses of the points in the graph.

Like previous works, we leverage the consistency of the local alignments to do this. Indeed, since a manifold is locally Euclidean, parallel transport along sufficiently short paths is approximately path-independent in the local neighborhood of a point. On the other hand, the further away two points are, the more inconsistent the cycles through them will be. When summing the transport over multiple paths between two points, inconsistent paths tend to average out.

We use the synchronization graph to build a discretization of this local parallel transport operator. Let the graph connection Laplacian W be the $2N \times 2N$ matrix with its i,j-th 2×2 block defined as:

$$W(i,j) = \begin{cases} \rho(h_{ij}) & \text{if } (i,j) \in \mathcal{G} \\ 0 & \text{otherwise} \end{cases} \in \mathbb{R}^{2 \times 2} \tag{7}$$

where $\rho(h_{ij})$ is the standard representation of $h_{ij} \in \text{O}(2)$ as a 2×2 orthogonal matrix. Let D be the $2N \times 2N$ diagonal matrix containing the *degree* of each node in $\mathcal{G}$ (repeated twice) in its diagonal. The matrix W acts by parallel-transporting tangent vectors along each edge of the graph while $A := D^{-1}W$ averages all vectors transported to a node. Then, W^t transports vectors along all length-t paths in $\mathcal{G}$ and the block $A^t(i,j)$ is the average of the transport over all length-t paths from i to j. If the paths are mostly inconsistent, $A^t(i,j)$ tends to 0, being the average of uncorrelated orthogonal matrices. Hence, $A^t(i,j)$ provides information about the geodesic distance between i and j.

Like [12], we take a **multi-frequency** approach to improve robustness to noise. In practice, we build a different graph connection Laplacian A_k for different choices[9] of *irreducible representation* (irrep) ρ_k of $\text{O}(2)$. For $k > 0$, ρ_k is a 2×2 orthogonal matrix with rotational frequency k; see Apx. A

In the limit $t \mapsto \infty$, the power iteration of A_k^t converges to its top eigenspace and $A_k^t(i,j)$ will be the product of the top eigenvectors of A_k. It is therefore important to discuss the spectral properties of A_k. [14] proved that the normalized graph connection Laplacian is a discretization of the connection Laplacian operator and that, under certain conditions, its eigenvectors and eigenvalues converge to those of the connection Laplacian operator. In Apx. B, we study the spectral properties of the connection Laplacian operator over $\text{P}\mathbb{R}^2$ and relate it with the operator over $\mathcal{S}^2$ studied in [13]. In particular, we show that the eigenspaces of our Laplacian operator over $\text{P}\mathbb{R}^2$ are subspaces of the eigenspaces of the spherical Laplacian and they share the same eigenvalues. This also implies our graph connection Laplacians enjoy the same spectral properties studied in [13], including the spectral gap and the robustness to noise.

For frequency $k > 0$, the top eigen-space of A_k is $2k + 1$ dimensional. Additionally, denote $\varphi_k : \mathcal{V} \to \mathbb{R}^{(2k+1) \times 2}$ the stack of the top $2k + 1$ eigen-vectors. Let $\underline{\varphi_k}(i) = \frac{\sqrt{2}}{\|\varphi_k(i)\|_F} \varphi_k(i)$ and define $\widehat{A_k}(i,j) = \underline{\varphi_k}(j)^T \underline{\varphi_k}(i) \in \mathbb{R}^{2 \times 2}$. Then:

Theorem 3.1. *Let* $s_{ij} = \langle \mathbf{z}_j, \mathbf{z}_i \rangle = \langle \pi(g_i), \pi(g_j) \rangle \in [-1, 1]$ *be the* cosine similarity *of the viewing directions* $\pi(g_i), \pi(g_j) \in \mathcal{S}^2$. *The two following identities hold:*

$$\frac{1}{2} \left\| (\widehat{A_k})(i,j) \right\|_F^2 \approx \left(\frac{1 + s_{ij}}{2} \right)^{2k} + \left(\frac{1 - s_{ij}}{2} \right)^{2k} \quad , \quad \det(\widehat{A_k}(i,j)) \approx \left(\frac{1 + s_{ij}}{2} \right)^{2k} - \left(\frac{1 - s_{ij}}{2} \right)^{2k}$$

The proof is in Apx. D. Since we need a similarity over $\text{P}\mathbb{R}^2$ rather than $\mathcal{S}^2$, we use $\widehat{w}_{ij} := |s_{ij}|$. In other words, the top eigenspace of A_k allows the estimation of the geodesic distance between points.

Denoised similarity and parallel transport The information obtained from the different A_k matrices can be combined in multiple ways to obtain an estimation of s_{ij}. Define

$$S_{ij}^{k\pm} := \frac{1}{4} \left\| (\widehat{A_k})(i,j) \right\|_F^2 \pm \frac{1}{2} \det(\widehat{A_k}(i,j)) \approx \left(\frac{1 \pm s_{ij}}{2} \right)^{2k} \tag{8}$$

For numerical stability, we estimate s_{ij} as

$$\widehat{s_{ij}} = \exp\left(\frac{1}{L} \sum_{k=1}^{L} \frac{\log S_{ij}^{k+}}{2k} \right) - \exp\left(\frac{1}{L} \sum_{k=1}^{L} \frac{\log S_{ij}^{k-}}{2k} \right) \tag{9}$$

---

[9]The standard representation of $\text{O}(2)$ as rotation matrices correspond to $k = 1$.

Algorithm 1 MFVDM based synchronization over $\mathbb{PR}^2$

Require: $\{o_i \mid i = 1, \ldots, N \in \mathcal{V}\}$ the input raw images, max frequency $L > 0$

1: $\mathcal{G}(\mathcal{V}, \mathcal{E}) \leftarrow$ synchronization-graph $(\{h_{ij} = \arg\min_h \|o_j - h.o_i\|_2^2\}_{ij}, \{d_{ij} = \min_h \|o_j - h.o_i\|_2^2\}_{ij})$

2: **for** $0 < k \leq L$ **do**

3:     $A_k(i,j) \leftarrow \rho_k(h_{ij})$ if $(i,j) \in \mathcal{E}$ else 0 $\forall i, j \in \mathcal{V}$ ▷ Frequency-k graph connection Laplacian; Eq. 7

4:     $\varphi_k \leftarrow$ top-eigenvectors$(A_k, 2k+1)$

5:     $\widehat{A_k}(i,j) \leftarrow \left(\frac{\sqrt{2}}{\|\varphi_k(j)\|_F} \varphi_k(j) \right)^T \left(\frac{\sqrt{2}}{\|\varphi_k(i)\|_F} \varphi_k(i) \right)$ $\forall i, j \in \mathcal{V}$

6:     $S_{ij}^{k\pm} \leftarrow \frac{1}{4} \left\| (\widehat{A_k})(i,j) \right\|_F^2 \pm \frac{1}{2} \det(\widehat{A_k}(i,j))$ $\forall i, j \in \mathcal{V}$ ▷ Eq. 8

7: $\widehat{s}_{ij} \leftarrow \exp\left(\frac{1}{L} \sum_{k=1}^{L} \frac{\log S_{ij}^{k+}}{2k} \right) - \exp\left(\frac{1}{L} \sum_{k=1}^{L} \frac{\log S_{ij}^{k-}}{2k} \right)$ $\forall i, j \in \mathcal{V}$ ▷ **Denoised similarity** over $\mathbb{PR}^2$; Eq. 9

8: $\widehat{h}_{ij} \leftarrow \arg\max_h \sum_{0 < k \leq L} \sqrt{\dim_{\rho_k}} \operatorname{Tr}\left(\widehat{A_k}(i,j) \rho_k(h)^T \right)$ $\forall i, j \in \mathcal{V}$ ▷ **Denoised parallel transport**; Eq. 10

9: $\mathcal{G}'(\mathcal{V}, \mathcal{E}') \leftarrow$ synchronization-graph $\left(\{\widehat{h}_{ij}\}_{ij}, \{|\widehat{s}_{ij}|\}_{ij} \right)$ ▷ Build denoised Synchronization Graph

10: $A' \leftarrow \left[\rho_1(\widehat{h}_{ij}) \text{ if } (i,j) \in \mathcal{E}' \text{ else } 0 \right]_{ij}$ ▷ Build denoised graph connection Laplacian; Eq. 7

11: $x, y \leftarrow \varphi \leftarrow$ top-eigenvectors$(A', 3)$ ▷ **Final Synchronization**

12: $\widehat{g}_i \leftarrow$ projectSVD$(x(i), y(i), x(i) \times y(i))$

---

where L is the largest frequency considered. Additionally, note that, due to the decaying spectrum of the Laplacian operators, the noise has a stronger effect on the lowest eigenvalues; discarding the lower eigenvectors also helps denoising the GCL matrices. That means that the top eigenvectors of A_k can be used to partially denoise the frequency-k parallel transport. Next, we combine all denoised blocks $\{\widehat{A_k}(i,j)\}_k$ by interpreting them as the Fourier coefficients of a function on H; we estimate the relative pose with the element $h \in H$ maximizing this function:

$$\widehat{h}_{ij} = \arg\max_{h \in H} \sum_{\rho_k \in \widehat{H}} \sqrt{\dim_{\rho_k}} \operatorname{Tr}\left(\widehat{A_k}(i,j) \rho_k(h)^T \right) \tag{10}$$

Final Synchronization So far, we have used the MFVDM method on $\mathbb{PR}^2$ to estimate the similarity $\widehat{w}_{ij} = |\widehat{s}_{ij}|$ and parallel transport $\widehat{h}_{ij}$ over $\mathbb{PR}^2$. With respect to [12], our method includes a final synchronization step, where these quantities are used to recover $\{\widehat{g}_i\}_i$. We construct a new graph connection Laplacian matrix A' as in Eq. 7 using the new estimations and the standard representation ($k = 1$) of O(2). The top eigen-space of A' is 3 dimensional and its 3 top eigenvectors define a tangent frame at each node in the following way. Let $\varphi : \mathcal{V} \to \mathbb{R}^{3 \times 2}$ be the stack of the top 3 eigen-vectors, $\underline{\varphi}(i) = \frac{\sqrt{2}}{\|\varphi(i)\|_F} \varphi(i)$ and $x : \mathcal{V} \to \mathbb{R}^3$ and $y : \mathcal{V} \to \mathbb{R}^3$ be its two columns. Then, $x(i)$ and $y(i) \in \mathbb{R}^3$ define a basis for the tangent space at $\pi(g_i) \in \mathcal{S}^2$, see Apx. C. Recall that $g_i = (\boldsymbol{x}_i, \boldsymbol{y}_i, \boldsymbol{z}_i) \in$ SO(3), then $x(i)$ and $y(i)$ approximates respectively $\boldsymbol{x}_i$ and $\boldsymbol{y}_i$, while $\boldsymbol{z}_i$ can be recovered[10] as $z(i) = x(i) \times y(i)$. Note that recovered poses $\{\widehat{g}_i = (x(i), y(i), z(i))\}_i$ present the global O(3) symmetry discussed in Sec. 2. Indeed, eigenvalue decomposition is unique up to an orthogonal change of basis in each eigenspace. Since the top eigenspace is 3 dimensional, the solution is unique up to a global $g \in$ O(3) transformation, i.e. if φ is a set of orthogonal eigenvectors, so is $g\varphi$, defined as $[g\varphi](i) = g\varphi(i)$. Hence, all tangent frames $\{(x(i), y(i))\}_i$ can be simultaneously rotated by any $g \in$ O(3). Finally, because $x(i)$ and $y(i)$ are not perfectly unitary and orthogonal to each other, the matrix $(x(i), y(i), z(i))$ will not be orthogonal; therefore, we project it to the closest SO(3) element via SVD.

Limitations VDM assumes uniformly distributed poses, which might not always be true; normalization methods such as [15, 16] can help relaxing this requirement. Our analysis of the synchronization symmetries, however, is not limited to VDM but applies to any cryo-EM synchronization method. 2D classification or pose estimation methods which do not account for the image formation model might fail at the lowest SNR regimes [5]. Indeed, if the noise is too high, the relative poses estimation itself can be impossible and increasing the number of observations can not solve this problem [17, 18], although methods like steerable PCA [19] can alleviate that. Nevertheless, in practice, 2D classification with VDM is leveraged as a denoising step in reconstruction pipelines (e.g. in ASPIRE); our method can replace this step, providing pose estimation and reducing the need for averaging. Finally, while we assumed non-symmetric molecules, symmetries directly affect the spectral properties of the GCL matrices and can theoretically be accounted for. We leave this as future work.

---

[10] $\langle z(j), z(i) \rangle = \langle x(j) \times y(j), x(i) \times y(i) \rangle = \det\left(\underline{\varphi}(j)^T \underline{\varphi}(i) \right) \approx s_{ij}$ by using Theorem 3.1 with $k = 1$.

4 Related Works

Vector-Diffusion Maps (VDM) were initially proposed in [9, 8] for solving the cryo-EM synchronization problem with $SO(2)$ constraints. The $O(2)$ symmetry was noted also in [8] which suggests augmenting a dataset with mirrored images; still, it only uses $SO(2)$ relative alignments and, therefore, recovers only a similarity metric over the sphere. VDMs were later extended to a multi-frequency setting in [12, 10, 13]. [20] describes a strategy based on the classical *method of moments* to solve the generic problem of recovering orbits from invariants. Additionally, the authors provide theoretical analysis of sample complexity of such problems; in particular, they show that in the cryo-EM synchronization problem with uniform viewing distribution, the sample complexity scales as $\frac{1}{SNR^3}$. Some works exploit the property that the Fourier transforms of any pair of cryo-EM images must agree on a line passing through the origin (*common lines approach*) to define a set of constraints on each pair of absolute poses. [21] use a Semi-Definite Programming (SDP) relaxation to find the solution; however, SDP is known for its high computational cost. Instead, [7, 22] rely on a spectral relaxation of the problem; still, common lines estimation is very sensitive to noise and is generally performed only after a first 2D classification and averaging phase [23]. Popular reconstruction methods include RELION [24] and cryoSPARC [25], which are based on an EM-like algorithm. [26] suggested using stochastic rather than batch optimization for iterative refinement, thereby improving the computational cost. Recent advances employ neural networks optimized by SGD to represent the 3D density [27, 28]. Note that these methods still need to periodically perform an expensive search over the images' poses and [28] found resetting the model's weights throughout training essential for convergence.

5 Experiments

We first evaluate our method on a synchronization task over some synthetic datasets of vectors which share the same symmetries of cryo-EM. Then, we include our method in a simple 3D reconstruction pipeline which we test on synthetic cryo-EM data. We include additional experiments in Appendix E.

Figure 4: Correlation with the $SO(3)$ poses in different variations of the vector dataset.

5.1 Synthetic Vector dataset

We generate a number of synthetic datasets which imitate the symmetry of the cryo-EM generative process. To do so, we first construct a matrix[11] $M \in \mathbb{R}^{d \times b}$ which has the symmetries of Π described in Table 1, where H acts on the left with a d-dimensional representation ρ of the group $O(2)$ while $O(3)$ acts on its right with a b-dimensional representation ψ. Both representation contain multiple irreps of different frequencies of the two groups. A dataset is generated by sampling a random vector $\Psi \in \mathbb{R}^b$ and a set of random poses $\{g_i \in SO(3)\}_i^n$ and then evaluating $\{o_i = M\psi(g_i^{-1})\Psi + \frac{1}{\lambda}\epsilon_i \in \mathbb{R}^d\}_i$, where ϵ_i's entries are normally distributed and λ^2 is the SNR.

We evaluate variants of our method on this data and compare it with MFVDM with $SO(2)$ poses. We use the $SO(3)$ correlation[12] $\frac{1}{n\sqrt{3}}\left\|\sum_i g_i\widehat{g}_i^{-1}\right\|_F$ and the correlation between the real and estimated cosine similarities $\{s_{ij}\}_{ij}$ as metrics. Fig. 4 shows the $SO(3)$

Figure 5: Estimated vs real cosine similarities $\{s_{ij}\}_{ij}$ in the synthetic vector dataset. We compare our method with $O(2)$ relative poses with the original MFVDM with $SO(2)$ poses.

correlation achieved by our method in different settings; in the low-noise regime, perfect reconstruction is achieved. As expected, in the highest noise regime, reconstruction is almost impossible as increasing the number of samples helps only marginally (SNR=0.16). Fig. 5 compares our method

---

[11]M contains a subset of the matrix decomposing $\mathrm{Ind}_{H \times \{e, m_z\}}^{O(3)} \rho$ into $O(3)$ irreps.

[12]Note that this metric is invariant to the global $O(3)$ symmetry.

with the original MFVDM-SO(2), for two choices of maximum frequency L. While the original estimator from MFVDM is stable when $s_{ij} \gg 0$, our estimator from Eq. 9 is stable when $|s_{ij}| \gg 0$. We also observe the effect of the multi-frequency approach: leveraging higher frequencies provide a more robust estimation.

5.2 3D reconstruction of synthetic cryo-EM data

For a more realistic evaluation, we use the 70S Ribosome [29] structure from the Protein Data Bank database (structure 5O60). We adopt a simple 3D reconstruction pipeline from [30] which optimizes via SGD a Gaussian posterior over the 3D density in Fourier space leveraging the Fourier Slice Theorem. Like [30], we ignore the contrast transfer function (CTF) in the rendering step for simplicity. The differentiability of the render allows us to also optimize the poses in the quaternion space. We compare three different initializations of the poses - with the ground truth, with the estimation produced by our method and with estimations produced via SO(2) alignment - and a version where we learn full posteriors over the poses using a discretization of SO(3) and the Gumbel-Softmax trick [31]. In the SO(2)

Figure 6: Random projections of reconstructions. *Rows*: 3000, 8000 and 8000 training samples with SNR $0.3, 0.09$ and 0.02. *Columns*: Gumbel-Softmax, MFVDM-SO(2), MFVDM-O(2) and GT methods.

alignment case, we use the similarity metric recovered by MFVDM to build a diffusion operator on $\mathcal{S}^2$, whose top eigenvectors estimate z_i; we initialize (x_i, y_i) to random orthogonal frames.

To evaluate the reconstruction, we compute the likelihood of a number of new *clean* images. We estimate the test images' poses by, first, estimating the optimal alignment $g \in O(3)$ between the training images' ground truth and estimated poses and, then, correcting the test images' poses by g. We also fine-tune these estimate poses before computing their final likelihood. Before synchronization, we perform a first dimensionality reduction and denoising step of the images using steerable PCA [32, 19] with a Fourier-Bessel basis. Since this basis is O(2)-steerable, we use PCA's embeddings to estimate the relative poses rather than the raw images. We estimate the initial synchronization graph by picking for each node its nearest neighbors according to the H-invariant distance.

Table 2 reports the correlation of the training images' poses and the negative log-likelihood (NLL) of 500 clean test images in different training settings. Our MFVDM-O(2) method can accurately recover the poses, yielding reconstruction quality close to the one obtained using ground truth poses (GT). Conversely, we do not find initializing the poses with the spherical embeddings recovered by MFVDM-SO(2) useful. Unfortunately, we find that the NLL - essentially, a mean-squared error - does not reflect the reconstruction quality in the lowest SNR regimes (e.g., the baselines have losses lower than the GT model). Still, the sample projections of the produced densities in Fig. 6 demonstrate that our method (MFVDM-O(2)) achieves similar visual quality to the ground-truth. Instead, the baseline models did not converge to meaningful structures, emphasizing the importance of pose initialization.

6 Conclusions and Discussion

In this work, we studied the symmetries of cryo-EM and found that the O(2) relative poses estimated from the images are sufficient to solve the SO(3) poses synchronization task. With this insight, we proposed a new method which can replace current VDM-based 2D classification methods with limited changes to directly estimate poses rather than clustering and averaging images, while enjoying the same noise robustness properties.

Table 2: Pose Correlation (and Negative Log-Likelihood) on test images with different poses initialization.

| n. samples | 1000 | 1000 | 1000 | 3000 | 3000 | 8000 | 8000 |
SNR	clean	2.7	0.67	0.67	0.3	0.09	0.02
Gumbel-Softmax						0.01 (35.3)	0.01 (50.9)
MFVDM-SO(2)	34.4 (23.6)	33.9 (33.8)	33.1 (50.3)	33.1 (46.4)	25.2 (27.8)	18.4 (44.0)	18.0 (45.0)
MFVDM-O(2)	100.0 (2.7)	99.6 (10.3)	99.3 (33.7)	99.6 (11.0)	99.0 (27.8)	88.8 (52.6)	57.7 (74.4)
Ground Truth	— (2.0)	— (10.4)	— (28.7)	— (11.7)	— (20.9)	— (30.2)	— (55.8)

References

[1] Daniel Cressey and Ewen Callaway. Cryo-electron microscopy wins chemistry nobel. *Nature*, 550(7675):167–167, October 2017. (Cited on 1)

[2] William T. Baxter, Robert A. Grassucci, Haixiao Gao, and Joachim Frank. Determination of signal-to-noise ratios and spectral snrs in cryo-em low-dose imaging of molecules. *Journal of Structural Biology*, 166(2):126–132, 2009. (Cited on 1)

[3] A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM Algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1):1–22, September 1977. (Cited on 1)

[4] F.J. Sigworth. A maximum-likelihood approach to single-particle image refinement. *Journal of Structural Biology*, 122(3):328–339, 1998. (Cited on 1)

[5] Amit Singer and Fred J Sigworth. Computational methods for single-particle cryo-em. 2020. (Cited on 2, 7)

[6] Jon Cohen. Is high-tech view of hiv too good to be true? *Science*, 341(6145):443–444, 2013. (Cited on 2)

[7] A. Singer and Y. Shkolnisky. Three-dimensional structure determination from common lines in cryo-EM by eigenvectors and semidefinite programming. *SIAM Journal on Imaging Sciences*, 4(2):543–572, January 2011. (Cited on 2, 8, 23)

[8] A. Singer, Z. Zhao, Y. Shkolnisky, and R. Hadani. Viewing angle classification of cryo-electron microscopy images using eigenvectors. *SIAM J. Img. Sci.*, 4(2):723–759, June 2011. (Cited on 2, 4, 8)

[9] Amit Singer and H-T Wu. Vector diffusion maps and the connection laplacian. *Communications on pure and applied mathematics*, 65(8):1067–1144, 2012. (Cited on 2, 5, 8)

[10] Yifeng Fan and Zhizhen Zhao. Cryo-electron microscopy image analysis using multi-frequency vector diffusion maps. *arXiv preprint arXiv:1904.07772*, 2019. (Cited on 2, 4, 8)

[11] Ronny Hadani and Amit Singer. Representation theoretic patterns in three-dimensional cryo-electron microscopy ii—the class averaging problem. *Foundations of computational mathematics*, 11(5):589–616, 2011. (Cited on 2)

[12] Yifeng Fan and Zhizhen Zhao. Multi-frequency vector diffusion maps. In *International Conference on Machine Learning*, pages 1843–1852. PMLR, 2019. (Cited on 4, 6, 7, 8, 24)

[13] Yifeng Fan, Tingran Gao, and Zhizhen Zhao. Representation theoretic patterns in multi-frequency class averaging for three-dimensional cryo-electron microscopy. *Information and Inference: A Journal of the IMA*, (iaab012), May 2021. (Cited on 4, 6, 8, 15, 16, 19)

[14] Amit Singer and Hau-Tieng Wu. Spectral convergence of the connection laplacian from random samples. *Information and Inference: A Journal of the IMA*, 6(1):58–123, 2017. (Cited on 6)

[15] Lihi Zelnik-manor and Pietro Perona. Self-tuning spectral clustering. In L. Saul, Y. Weiss, and L. Bottou, editors, *Advances in Neural Information Processing Systems*, volume 17. MIT Press, 2004. (Cited on 7)

[16] Ronald R. Coifman and Stéphane Lafon. Diffusion maps. *Applied and Computational Harmonic Analysis*, 21(1):5–30, 2006. Special Issue: Diffusion Maps and Wavelets. (Cited on 7)

[17] Cecilia Aguerrebere, Mauricio Delbracio, Alberto Bartesaghi, and Guillermo Sapiro. Fundamental limits in multi-image alignment. *IEEE Transactions on Signal Processing*, 64(21):5707–5722, 2016. (Cited on 7)

[18] Tamir Bendory, Alberto Bartesaghi, and Amit Singer. Single-particle cryo-electron microscopy: Mathematical theory, computational challenges, and opportunities. *IEEE Signal Processing Magazine*, 37(2):58–76, 2020. (Cited on 7)

[19] Zhizhen Zhao, Yoel Shkolnisky, and Amit Singer. Fast steerable principal component analysis. *IEEE transactions on computational imaging*, 2(1):1–12, 2016. (Cited on 7, 9, 24)

[20] Afonso S Bandeira, Ben Blum-Smith, Joe Kileel, Amelia Perry, Jonathan Weed, and Alexander S Wein. Estimation under group actions: recovering orbits from invariants. *arXiv preprint arXiv:1712.10163*, 2017. (Cited on 8)

[21] Afonso S Bandeira, Yutong Chen, Roy R Lederman, and Amit Singer. Non-unique games over compact groups and orientation estimation in cryo-em. *Inverse Problems*, 36(6):064002, 2020. (Cited on 8)

[22] Yoel Shkolnisky and Amit Singer. Viewing direction estimation in cryo-em using synchronization. *SIAM journal on imaging sciences*, 5(3):1088–1110, 2012. (Cited on 8, 23)

[23] Amit Singer, Ronald R. Coifman, Fred J. Sigworth, David W. Chester, and Yoel Shkolnisky. Detecting consistent common lines in cryo-em by voting. *Journal of Structural Biology*, 169(3):312–322, 2010. (Cited on 8, 23)

[24] Sjors H.W. Scheres. Relion: Implementation of a bayesian approach to cryo-em structure determination. *Journal of Structural Biology*, 180(3):519–530, 2012. (Cited on 8)

[25] Ali Punjani, John L Rubinstein, David J Fleet, and Marcus A Brubaker. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nature Methods*, 14(3):290–296, February 2017. (Cited on 8)

[26] Marcus A Brubaker, Ali Punjani, and David J Fleet. Building proteins in a day: Efficient 3d molecular reconstruction. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, pages 3099–3108, 2015. (Cited on 8)

[27] Ellen D Zhong, Tristan Bepler, Joseph H Davis, and Bonnie Berger. Reconstructing continuous distributions of 3d protein structure from cryo-em images. *arXiv preprint arXiv:1909.05215*, 2019. (Cited on 8)

[28] Ellen D. Zhong, Adam Lerer, Joseph H. Davis, and Bonnie Berger. Cryodrgn2: Ab initio neural reconstruction of 3d protein structures from real cryo-em images. In *Proceedings of the IEEE/CVF International Conference on Computer Vision (ICCV)*, pages 4066–4075, October 2021. (Cited on 8)

[29] Jendrik Hentschel, Chloe Burnside, Ingrid Mignot, Marc Leibundgut, Daniel Boehringer, and Nenad Ban. The complete structure of the mycobacterium smegmatis 70s ribosome. *Cell Reports*, 20(1):149–160, 2017. (Cited on 9, 22)

[30] Karen Ullrich, Rianne van den Berg, Marcus Brubaker, David Fleet, and Max Welling. Differentiable probabilistic models of scientific imaging with the fourier slice theorem. In *proceedings of the Conference on Uncertainty in Artificial Intelligence (UAI)*, 2019. (Cited on 9, 20, 21)

[31] Eric Jang, Shixiang Gu, and Ben Poole. Categorical reparameterization with gumbel-softmax. 2016. (Cited on 9, 22)

[32] Zhizhen Zhao and Amit Singer. Fourier–bessel rotational invariant eigenimages. *JOSA A*, 30(5):871–877, 2013. (Cited on 9)

[33] Gabriele Cesa, Leon Lang, and Maurice Weiler. A program to build E(N)-equivariant steerable CNNs. In *International Conference on Learning Representations*, 2022. (Cited on 14)

[34] Zhizhen Zhao and Amit Singer. Rotationally invariant image representation for viewing direction classification in cryo-em. *Journal of structural biology*, 186(1):153–166, 2014. (Cited on 22, 23, 24)