A Topological Perspective on Causal Inference

Abstract

This paper presents a topological learning-theoretic perspective on causal inference by introducing a series of topologies defined on general spaces of structural causal models (SCMs). As an illustration of the framework we prove a topological causal hierarchy theorem, showing that substantive assumption-free causal inference is possible only in a meager set of SCMs. Thanks to a known correspondence between open sets in the weak topology and statistically verifiable hypotheses, our results show that inductive assumptions sufficient to license valid causal inferences are statistically unverifiable in principle. Similar to no-free-lunch theorems for statistical inference, the present results clarify the inevitability of substantial assumptions for causal inference. An additional benefit of our topological approach is that it easily accommodates SCMs with infinitely many variables. We finally suggest that the framework may be helpful for the positive project of exploring and assessing alternative causal-inductive assumptions.

1 Introduction and Motivation

In the background of any investigation into learning algorithms are no-free-lunch phenomena: roughly, the observation that assumption-free statistical learning is infeasible in general (see, e.g., [32, Ch. 5] for a formal statement). Common wisdom is that learning algorithms and architectures must adequately reflect non-trivial features of the data-generating distribution to gain inductive purchase.

For many purposes we need to move beyond passive observation, focusing instead on what *would* happen were we to *act* upon a given system. Even further, we sometimes desire to *explain* the behavior of a system, raising questions about what *would have* occurred had some aspects of a situation been different. Such questions depend not just on the data distribution; they depend on deeper features of underlying data-generating *processes* or *mechanisms*. It is thus generally acknowledged that stronger assumptions are required if we want to draw *causal* conclusions from data [34, 27, 19, 29, 31].

Whether implicit or explicit, any approach to causal inference involves a space of candidate causal models, viz. data-generating processes. Indeed, a blunt way of incorporating inductive bias is simply to *omit* some class of possible causal hypotheses from consideration. Many (im)possibility results in the literature can accordingly be understood as pertaining to all models within a class. For instance, if we can restrict attention to *Markovian* models that satisfy *faithfulness*, then we can *always* identify the structure of a model from experimental data (e.g., [11, 34]). If we can restrict attention to Markovian (continuous) models with linear functions and *non*-Gaussian noise, then *every* model can be learned even from purely observational data [33]. As a negative example, in the larger class of (not necessarily Markovian) models, *no* model can *ever* be determined from observational data alone [34, 3].

At the same time, in many settings it is sensible to aim for results with "nearly universal" force. It is natural to ask, e.g., within the class of all Markovian models, how "typical" are those in which the faithfulness condition is violated? This might tell us, for instance, how typically we could expect failure of a method that depended on these assumptions. A well-known result shows that, fixing

any particular causal dependence graph, such violations have *measure zero* for any smooth (e.g., Lebesgue) measure on the parameter space of distributions consistent with that graph [23]. In fact, the very notion of statistical consistency itself, which underlies many possibility results in causal inference, must generally omit some purportedly "negligible" set of possible data streams [9, 34].

There are two standard mathematical approaches to making concepts like "typical" and "negligible" rigorous: *measure-theoretic* and *topological*. While the two approaches often agree, they capture slightly different intuitions [24]. One virtue of the measure-theoretic approach is its natural probabilistic interpretation: intuitively, we are exceedingly *unlikely* to hit upon a set with measure zero. At the same time, the measure-theoretic approach is sometimes criticized in statistical settings for its alleged dependence on a measure, and this has been argued to favor topological approaches (see, e.g., [4] on no-free-lunch theorems). The latter of course in turn demands an appropriate topology.

In the present work we show how to define a sequence of meaningful topologies on the space of causal models, each corresponding to a progressively coarser level of the so called *causal hierarchy* ([28, 3]; see Fig. 1 for an abbreviated pictorial summary). We aim to demonstrate that topologizing causal models in this way helps clarify the scope and limits of causal inference under different assumptions, as well as the potential empirical status of those very assumptions, in a highly general setting.

Our starting point is a canonical topology on the space of Borel probability distributions called the *weak topology*. The weak topology is grounded in the fundamental notion of *weak convergence* of probability distributions [5] and is thereby closely related to problems of statistical inference (see, e.g., [8]). Recent work has sharpened this correspondence, showing that open sets in the weak topology correspond exactly to the statistical hypotheses that can be naturally deemed (almost-surely) *verifiable* [14, 15]. We extend the correspondence to higher levels of the causal hierarchy, including the most refined and expansive "top" level consisting of all (well-founded) causal models. Lower levels and natural subspaces (e.g., corresponding to prominent causal assumption classes) emerge as coarsenings and continuous projections of this largest space. As an illustration of the general approach, we prove a topological version of the causal hierarchy theorem from [3]. Rather than showing that collapse happens only in a measure zero set as in [3], our Theorem 3 show that collapse is *topologically meager*. Conceptually, this highlights a different (but complementary) intuition: not only is collapse exceedingly unlikely in the sense of measure, meagerness implies that collapse could never be *statistically verified*. Correlatively, this implies that any causal assumption that would generally allow us to infer counterfactual probabilities from experimental (or "interventional") probabilities must itself be statistically unverifiable (Corollary 1).

To derive such a result we actually show something slightly stronger (see Lem. 2): even with respect to the subspace of models consistent with a fixed *temporal order* on variables, the causal hierarchy theorem holds. Merely knowing the temporal order of the variables is not enough to render collapse of the hierarchy a statistically verifiable proposition.

In §2 we give background on causal models, and in §3 we present a model-theoretic characterization of the causal hierarchy as a sequence of spaces. Topology is introduced in §4, and the main results about collapse appear in §5. For the technical results, we include proof sketches in the main text to provide the core intuitions, relegating some of the details to an exhaustive technical appendix, which also includes additional supplementary material.

2 Structural Causal Models

A fundamental building block in the theory of causality is the *structural causal model* [25, 34, 27] or SCM, which formalizes the notion of a data-generating process. In addition to specifying data-generating distributions, these models also specify the generative mechanisms that produce them. For the purpose of causal inference and learning, SCMs provide a broad, fine-grained hypothesis space.

The notions in this section have their usual definition following, e.g., [27], but we have recast them in the standard language of Borel probability spaces so as to handle the case of infinitely many variables rigorously. We start with notation, basic assumptions, and some probability theory.

Notation. The signature (or range) of a variable V is denoted χ_V. Where $\mathbf{S}$ is a set of variables, let $\chi_{\mathbf{S}} = \bigtimes_{S \in \mathbf{S}} \chi_S$. Given an indexed family of sets $\{S_\beta\}_{\beta \in B}$ and elements $s_\beta \in S_\beta$, let $(s_\beta)_\beta$ denote the tuple whose element at index β is s_β, for all β. For $B' \subset B$ write $\pi_{B'} : \bigtimes_{\beta \in B} S_\beta \to \bigtimes_{\beta \in B'} S_\beta$ for the *projection map* sending each $(s_\beta)_{\beta \in B} \mapsto (s_{\beta'})_{\beta' \in B'}$; abbreviate $\pi_{\beta'} = \pi_{\{\beta'\}}$, where $\beta' \in B$.

91 The reader is referred to standard texts [20, 6] for elaboration on the concepts used below.

92 **Definition 1** (Topology). For discrete spaces (like χ_S, for a single categorical variable S) we use
93 the discrete topology and for product spaces (like $\chi_{\mathbf{S}}$ for a set of variables $\mathbf{S}$) we use the product
94 topology. Note that *cylinder sets* of the form $\pi_{\mathbf{Y}}^{-1}(\{\mathbf{y}\})$ for finite subsets $\mathbf{Y} \subset \mathbf{S}$ and $\mathbf{y} \in \chi_{\mathbf{Y}}$ form a
95 basis on $\chi_{\mathbf{S}}$. Following standard statistical notation this cylinder is abbreviated as simply $\mathbf{y}$.

96 **Definition 2** (Probability). Where ϑ is a topological space write $\mathcal{B}(\vartheta)$ for its Borel σ-algebra of
97 measurable subsets. Let $\mathfrak{P}(\vartheta)$ be the set of probability measures on $\mathcal{B}(\vartheta)$. Specifically, elements
98 of $\mathfrak{P}(\vartheta)$ are functions $\mu : \mathcal{B}(\vartheta) \to [0,1]$ assigning a probability to each measurable set such that
99 $\mu(\vartheta) = 1$ and $\mu\big(\bigcup_{i=1}^{\infty}(S_i)\big) = \sum_{i=1}^{\infty} \mu(S_i)$ for each sequence $S_1, S_2, \ldots$ of pairwise disjoint sets
100 from $\mathcal{B}(\vartheta)$. A map $f : \vartheta_1 \to \vartheta_2$ is said to be *measurable* if $f^{-1}(S_2) \in \mathcal{B}(\vartheta_1)$ for every $S_2 \in \mathcal{B}(\vartheta_2)$.

101 **Fact 1** (Lemma 1.9.4 [6]). A Borel probability measure is determined by its values on a basis.

2.1 SCMs, Observational Distributions

103 Let $\mathbf{V}$ be a set of *endogenous variables*. We assume for simplicity every variable $V \in \mathbf{V}$ is
104 dichotomous with $\chi_V = \{0,1\}$, although the results here generalize to any larger countable range.
105 Influences among endogenous variables are the main phenomena our formalism aims to capture. A
106 well-founded *direct influence* relation $\to$ on $\mathbf{V}$ encapsulates the notion of one endogenous variable
107 possibly influencing another. For each $V \in \mathbf{V}$, we call $\{V' \in \mathbf{V} : V' \to V\} = \mathbf{Pa}(V)$ the *parents*
108 of $\mathbf{V}$. We assume every set $\mathbf{Pa}(V)$ is finite; this is called *local finiteness*. These two assumptions
109 (well-foundedness and local finiteness) generalize the well known *recursiveness* assumption to the
110 infinitary setting, and have an alternative characterization in terms of "temporal" orderings:

111 **Fact 2.** Say that a total order $\prec$ on $\mathbf{V}$ is ω-*like* if for each $V \in \mathbf{V}$, the set $\{V' : V' \prec V\}$ is finite.
112 Then the influence relation $\to$ is extendible to an ω-like order iff $\to$ is well-founded and locally finite.

113 In addition to endogenous variables, causal models have *exogenous variables* $\mathbf{U}$. Each endogenous V
114 depends on a subset $\mathbf{U}(V) \subset \mathbf{U}$ of "exogenous parents" and uncertainty enters via exogenous noise,
115 that is, a distribution from $\mathfrak{P}(\chi_{\mathbf{U}})$. A *structural function* (or *mechanism*) for $V \in \mathbf{V}$ is a measurable
116 $f_V : \chi_{\mathbf{Pa}(V)} \times \chi_{\mathbf{U}(V)} \to \chi_V$ mapping parental endogenous and exogenous valuations to values.

117 **Definition 3.** A *structural causal model* is a tuple $\mathcal{M} = \langle \mathbf{U}, \mathbf{V}, \{f_V\}_{V \in \mathbf{V}}, P \rangle$ where $\mathbf{U}$ is a
118 collection of exogenous variables, $\mathbf{V}$ is a collection of endogenous variables, f_V is a structural
119 function for each $V \in \mathbf{V}$, and $P \in \mathfrak{P}(\chi_{\mathbf{U}})$ is a probability measure on (the Borel σ-algebra of) $\chi_{\mathbf{U}}$.

120 Recursiveness implies that each $\mathbf{u} \in \chi_{\mathbf{U}}$ induces a unique $\mathbf{v} \in \chi_{\mathbf{V}}$ that solves $\{V = f_V\}_V$:

121 **Proposition 1.** Any SCM $\mathcal{M}$ with well-founded, locally finite parent relation $\to$ induces a unique
122 measurable $m^{\mathcal{M}} : \chi_{\mathbf{U}} \to \chi_{\mathbf{V}}$ such that $f_V\big(\pi_{\mathbf{Pa}(V)}(m^{\mathcal{M}}(\mathbf{u})), \pi_{\mathbf{U}(V)}(\mathbf{u})\big) = \pi_V\big(m^{\mathcal{M}}(\mathbf{u})\big)$ for all
123 $\mathbf{u} \in \chi_{\mathbf{U}}$ and $V \in \mathbf{V}$.

124 Measurability then entails that the exogenous noise P induces a distribution on joint valuations of $\mathbf{V}$,
125 called the *observational distribution*, which characterizes passive observations of the system.

126 **Definition 4.** The observational distribution $p^{\mathcal{M}} \in \mathfrak{P}(\chi_{\mathbf{V}})$ is defined on open sets by $p^{\mathcal{M}}(\mathbf{y}) =$
127 $P\big((m^{\mathcal{M}})^{-1}(\mathbf{y})\big)$. Here recall that $\mathbf{y}$ represents a cylinder subset of $\chi_{\mathbf{V}}$.

2.2 Interventions

129 What makes SCMs distinctively causal is the way they accommodate statements about possible
130 manipulations of a causal setup capturing, e.g., observations resulting from a controlled experimental
131 trial. This is formalized in the following definition.

132 **Definition 5.** An *intervention* is a choice of a finite subset of variables $\mathbf{W} \subset \mathbf{V}$ and $\mathbf{w} \in \chi_{\mathbf{W}}$. This
133 intervention is written $\mathbf{W} := \mathbf{w}$, and we let A be the set of all interventions. Under this intervention,
134 each $W \in \mathbf{W}$ is held fixed to its value $\pi_W(\mathbf{w}) \in \chi_W$ in $\mathbf{w}$ while the mechanism for any $V \in \mathbf{V} \setminus \mathbf{W}$
135 is left unchanged. Specifically, where $\mathcal{M}$ is as in Definition 3, the manipulated model for $\mathbf{W} := \mathbf{w}$ is
136 the model $\mathcal{M}_{\mathbf{W}:=\mathbf{w}} = \langle \mathbf{U}, \mathbf{V}, \{f_V^{\mathbf{W}:=\mathbf{w}}\}_{V \in \mathbf{V}}, P \rangle$ where

$$f_V^{\mathbf{W}:=\mathbf{w}} = \begin{cases} f_V, & V \notin \mathbf{W} \\ \text{constant func. mapping to } \pi_V(\mathbf{w}), & V \in \mathbf{W}. \end{cases}$$

137 The *interventional* or *experimental distribution* $p^{\mathcal{M}_{\mathbf{W}:=\mathbf{w}}} \in \mathfrak{P}(\chi_{\mathbf{V}})$ is just the observational distribu-
138 tion for the manipulated model $\mathcal{M}_{\mathbf{W}:=\mathbf{w}}$, and it encodes the probabilities for an experiment in which
139 the variables $\mathbf{W}$ are fixed to the values $\mathbf{w}$.

140 **Remark 1.** Empty interventions $\varnothing := ()$ are just passive observations, i.e., $p^{\mathcal{M}_{\varnothing:=()}} = p^{\mathcal{M}}$.

2.3 Counterfactuals

142 By permitting multiple manipulated settings to share exogenous noise, not only the distribution
143 arising from a single manipulation, but also joint distributions over multiple can be considered. These
144 are often called *counterfactuals*. The set $\mathfrak{P}(\chi_{A \times \mathbf{V}})$ encompasses the combined joint distributions
145 over $\mathbf{V}$ for any combination of interventions from A. A basis for the space $\chi_{A \times \mathbf{V}}$ are the cylinder
146 sets of the following form, for some sequence $(\mathbf{X} := \mathbf{x}, \mathbf{Y}), \ldots, (\mathbf{W} := \mathbf{w}, \mathbf{Z})$ of pairs, where
147 $\mathbf{Y}, \ldots, \mathbf{Z} \subset \mathbf{V}$ are finite, and $\mathbf{X} := \mathbf{x}, \ldots, \mathbf{W} := \mathbf{w} \in A$ are interventions:

$$\pi_{\{\mathbf{X}:=\mathbf{x}\} \times \mathbf{Y}}^{-1}(\{\mathbf{y}\}) \cap \cdots \cap \pi_{\{\mathbf{W}:=\mathbf{w}\} \times \mathbf{Z}}^{-1}(\{\mathbf{z}\}).$$

148 We will abbreviate this open set as $\mathbf{y}_{\mathbf{x}}, \ldots, \mathbf{z}_{\mathbf{w}}$, writing, e.g. simply $\mathbf{x}$ for the intervention $\mathbf{X} = \mathbf{x}$.

149 **Definition 6.** Given $\mathcal{M}$, define a counterfactual distribution $p_{\text{cf}}^{\mathcal{M}} \in \mathfrak{P}(\chi_{A \times \mathbf{V}})$ on a basis as follows:

$$p_{\text{cf}}^{\mathcal{M}}(\mathbf{y}_{\mathbf{x}}, \ldots, \mathbf{z}_{\mathbf{w}}) = P\big((m^{\mathcal{M}_{\mathbf{X}:=\mathbf{x}}})^{-1}(\mathbf{y}) \cap \cdots \cap (m^{\mathcal{M}_{\mathbf{W}:=\mathbf{w}}})^{-1}(\mathbf{z})\big).$$

150 **Remark 2.** Marginalizing $p_{\text{cf}}^{\mathcal{M}}$ to any single intervention $\mathbf{W} := \mathbf{w}$ yields $p^{\mathcal{M}_{\mathbf{W}:=\mathbf{w}}}$. If $\chi_{\mathbf{U}}$ is finite, we
151 obtain a familiar [13] sum formula $p_{\text{cf}}^{\mathcal{M}}(\mathbf{y}_{\mathbf{x}}, \ldots, \mathbf{z}_{\mathbf{w}}) = \sum_{\{\mathbf{u}|m^{\mathcal{M}_{\mathbf{X}:=\mathbf{x}}}(\mathbf{u}) \in \mathbf{y}, \ldots, m^{\mathcal{M}_{\mathbf{W}:=\mathbf{w}}}(\mathbf{u}) \in \mathbf{z}\}} P(\mathbf{u})$.

2.4 SCM classes

153 We now define several subclasses of SCMs that we will use throughout the paper. Notably, we do
154 not require their endogenous variable sets $\mathbf{V}$ to be finite. It is infinite in many applications, for
155 instance, in time series models, or generative models defined by probabilistic programs (see, e.g.,
156 [17, 35]). Because the proofs call for slightly different methods, we deal with the infinite and finite
157 cases separately. We make one additional assumption in the infinite case.

158 **Definition 7.** $\mu \in \mathfrak{P}(\vartheta)$ is *atomless* if $\mu(\{t\}) = 0$ for each $t \in \vartheta$; $\mathcal{M}$ is atomless if $p_{\text{cf}}^{\mathcal{M}}$ is atomless.

159 Intuitively, an atomless distribution is one in which weight is always "smeared" out continuously
160 and there are no point masses; infinitely many fair coin flips, for example, generate an atomless
161 distribution as the probability of obtaining any given infinite sequence is zero.

162 **Definition 8.** For the remainder of the paper, fix a countable endogenous variable set $\mathbf{V}$. Define the
163 following classes of SCMs:

$$\mathfrak{M}_{\prec} = \text{SCMs over } \mathbf{V} \text{ whose influence relation is extendible to the } \omega\text{-like order } \prec;$$
$$\mathfrak{M}_X = \text{SCMs over } \mathbf{V} \text{ in which the variable } X \text{ has no parents: } \mathbf{Pa}(X) = \varnothing;$$
$$\mathfrak{M} = \text{all SCMs over } \mathbf{V} = \bigcup_{\prec} \mathfrak{M}_{\prec} = \bigcup_X \mathfrak{M}_X.$$

164 If $\mathbf{V}$ is infinite then all SCMs in the classes above are assumed to be atomless.

3 The Causal Hierarchy

166 Implicit in §2, and indeed in much of the literature on causal inference, is a hierarchy of causal
167 expressivity. Following the metaphor offered in [28], it is natural to characterize three levels of
168 the hierarchy as the *observational*, *interventional* (experimental), and *counterfactual* (explanatory).
169 Drawing on recent work [3, 18] we make this characterization explicit. The levels will be defined in
170 descending order of causal expressivity (the reverse of §2). Fig. 1(a) summarizes our definitions.

171 Higher levels determine lower levels—counterfactuals determine interventionals, and the obser-
172 vational is just an (empty) interventional. Thus movement "downward" in the causal hierarchy
173 corresponds to a kind of projection. For indexed $\{S_\beta\}_{\beta \in B}$ and $B' \subset B$ let $\varsigma_{B'} : \mathfrak{P}(\bigtimes_{\beta \in B} S_\beta) \to$
174 $\mathfrak{P}(\bigtimes_{\beta \in B'} S_\beta)$ be the *marginalization* map taking a joint distribution to its marginal on B'.

Definition 9. Define three composable *causal projections* $\{\varpi_i\}_{1 \leq i \leq 3}$ with signatures and definitions

$$\varpi_3 : \mathfrak{M} \to \mathfrak{P}(\chi_{A \times \mathbf{V}}), \quad \varpi_2 : \mathfrak{P}(\chi_{A \times \mathbf{V}}) \to \underset{\alpha \in A}{\bigtimes} \mathfrak{P}(\chi_{\mathbf{V}}), \quad \varpi_1 : \underset{\alpha \in A}{\bigtimes} \mathfrak{P}(\chi_{\mathbf{V}}) \to \mathfrak{P}(\chi_{\mathbf{V}});$$

$$\varpi_3(\mathcal{M}) = p^{\mathcal{M}}, \quad \varpi_2(\mu) = \left(\varsigma_{\{\alpha\} \times \mathbf{V}}(\mu) \right)_{\alpha \in A}, \quad \varpi_1(\mu) = \pi_{\varnothing}(\mu).$$

The *causal hierarchy* consists of three sets $\{\mathfrak{S}_i\}_{1 \leq i \leq 3}$ defined as images or projections of $\mathfrak{M}$:

$$\mathfrak{S}_3 = \varpi_3(\mathfrak{M}), \quad \mathfrak{S}_2 = \varpi_2(\mathfrak{S}_3), \quad \mathfrak{S}_1 = \varpi_1(\mathfrak{S}_2).$$

These are the three *Levels* of the hierarchy. The definitions cohere with those of §2:

Fact 3. Let $\mathcal{M} \in \mathfrak{M}$. Then $\mu_3 = \varpi_3(\mathcal{M}) \in \mathfrak{S}_3$ trivially coincides with its counterfactual distribution as defined in §2.3, while $\mu_2 = \varpi_2(\mu_3) \in \mathfrak{S}_2$ coincides with the indexed family of all its interventional distributions (§2.2), i.e., $\pi_{\mathbf{W} := \mathbf{w}}(\mu_2) = p^{\mathcal{M}_{\mathbf{W} := \mathbf{w}}}$ for each $\mathbf{W} := \mathbf{w} \in A$. Finally $\mu_1 = \varpi_1(\mu_2) \in \mathfrak{S}_1$ coincides with its observational distribution (§2.1).

Thus, e.g., $\mathfrak{S}_3$ is the set of counterfactual distributions that are consistent with at least some SCM from $\mathfrak{M}$. It is a fact that $\mathfrak{S}_3 \subsetneq \mathfrak{P}(\chi_{A \times \mathbf{V}})$ and similarly not every interventional family belongs to $\mathfrak{S}_2$; see Appendix B for explicit characterizations. At the observational level, this is simple:

Fact 4. $\mathfrak{S}_1 = \mathfrak{P}(\chi_{\mathbf{V}})$ in the finite case. In the infinite case, $\mathfrak{S}_1 = \{\mu \in \mathfrak{P}(\chi_{\mathbf{V}}) : \mu \text{ is atomless}\}$.

We will also use the subsets $\{\mathfrak{S}_i^{\prec}\}_i$ and $\{\mathfrak{S}_i^X\}_i$, which are defined analogously but via projection from $\mathfrak{M}_{\prec}$ and $\mathfrak{M}_X$ respectively.

3.1 Problems of Causal Inference

As elucidated in [28, 3], the causal hierarchy helps characterize many standard problems of causal inference, in as far as these problems typically involve ascending levels of the hierarchy. Some examples include:

1. Classical identifiability: given observational data about some variables in $\mathbf{V}$, estimate a *causal effect* of setting variables $\mathbf{X}$ to values $\mathbf{x}$ [25, 34]. In the notation here, given information about $p^{\mathcal{M}}(\mathbf{V})$, can we determine $p^{\mathcal{M}_{\mathbf{X} := \mathbf{x}}}(\mathbf{Y})$?

2. General identifiability: given a mix of observational data and limited experimental data—that is, information about $p^{\mathcal{M}}(\mathbf{V})$ as well as some experimental distributions of the form $p^{\mathcal{M}_{\mathbf{W} := \mathbf{w}}}(\mathbf{V})$—determine $p^{\mathcal{M}_{\mathbf{X} := \mathbf{x}}}(\mathbf{Y})$ [37, 21].

3. Structure learning: given observational data, and perhaps experimental data, infer properties of the underlying causal influence relation $\to$ [34, 29].

4. Counterfactual estimation: given a combination of observational and experimental data, infer a counterfactual quantity, such as probability of necessity [30], or probability of necessity and sufficiency [26, 36] (see also §3.3 below).

5. Global identifiability: given observational data drawn from $p^{\mathcal{M}}(\mathbf{V})$ infer the full counterfactual distribution $p_{\text{cf}}^{\mathcal{M}}(A \times \mathbf{V})$ [33, 10].

This is not an exhaustive list, and these problems are not all independent of one another. They are also all unsolvable in general. Problems 1, 2, and 3 involve ascending to Level 2 given information at Level 1 (and perhaps partial information at Level 2); problems 4 and 5 ask us to ascend to Level 3 given only Level 1 (and perhaps also Level 2) information. The upshot of the causal hierarchy theorem from [3] is that these steps are impossible without assumptions, formalizing the common wisdom, "no causes in, no causes out" [7]. To understand the statement of the causal hierarchy theorem—and our topological version of it—we explain what it means for the hierarchy to collapse.

3.2 Collapse of the Hierarchy

In the present setting a *collapse* of the hierarchy can be understood in terms of injectivity of the functions ϖ_i. For $i = 1, 2$ let $\mathfrak{C}_i \subset \mathfrak{S}_i$ be the injective fibers of ϖ_i, i.e., $\mathfrak{C}_i = \{\mu_i \in \mathfrak{S}_i : \mu_{i+1} = \mu'_{i+1} \text{ whenever } \varpi_i(\mu_{i+1}) = \varpi_i(\mu'_{i+1}) = \mu_i\}$. Every element $\mu \in \mathfrak{C}_i$ is a witness to (global) collapse of the hierarchy: knowing μ would be sufficient to determine the Level $i + 1$ facts completely.

| (a) Causal Hierarchy | (b) Collapse Set $\mathfrak{C}_2$ |

Figure 1: (a) $\mathfrak{S}_3$ can be seen as a coarsening of $\mathfrak{M}$, abstracting from irrelevant "intensional" details. $\mathfrak{S}_2$ is obtained from $\mathfrak{S}_3$ by marginalization (also a coarsening), while $\mathfrak{S}_1$ is a projection of $\mathfrak{S}_2$ via the "empty" intervention. Each map ϖ_i, $i = 1, 2, 3$, is continuous for the respective weak topologies (Prop. 4). The projections $\varpi_{\text{NS}(Y)}^X$ from $\mathfrak{S}_2$ to the PNS-spaces are likewise continuous (Prop. 4).
(b) The shaded region, $\mathfrak{C}_2 \subset \mathfrak{S}_2$, is the collapse set in which Level 2 facts determine all Level 3 facts: those points in $\mathfrak{S}_2$ whose ϖ_2-preimage in $\mathfrak{S}_3$ is a singleton set. The main result of this paper is that $\mathfrak{C}_2$ is *meager* in weak topology on $\mathfrak{S}_2$ (Thm. 3). This means $\mathfrak{C}_2$ contains no open subset, which by Thm. 2 implies no part of $\mathfrak{C}_2$ is statistically verifiable, even with infinitely many ideal experiments.

A first observation is that ϖ_1 is *never* injective. In other words, the distribution $p^{\mathcal{M}}(\mathbf{V})$ never determines all the interventional distributions $p^{\mathcal{M}_{\mathbf{x}:=\mathbf{x}}}(\mathbf{Y})$. This is essentially a way of stating that correlation never implies causation absent assumptions. (See also [3, Thm. 1].)

Proposition 2. $\mathfrak{C}_1 = \varnothing$. That is, Level 2 *never* collapses to Level 1 without assumptions.

To overcome this formidable inferential barrier, researchers often assume we are not working in the "full" space $\mathfrak{M}$ of all causal models, but rather some proper subset embodying a range of causal assumptions. This may effectively eliminate counterexamples to collapse (cf. Fig. 1(b)). For problems of type 1 or 2 (from the list above in §3.1) it is common to assume we are only dealing with models whose graph (direct influence relation) $\to$ satisfies a fixed set of properties. For problems of type 3 it is common to assume that $p^{\mathcal{M}}$ and $\to$ relate in some way (for instance, through an assumption like *faithfulness* or *minimality* [34]). All of these problems become solvable with sufficiently strong assumptions about the form of the functions $\{f_V\}_V$ or the probability space P.

In some cases, the relevant causal assumptions are justified by appeal to background or expert knowledge. In other cases, however, an assumption will be justified by the fact that it rules out only a "small" or "negligible" or "measure zero" part of the full set $\mathfrak{M}$ of possibilities. As emphasized recently [22], not all "small" subsets are the same, and it seems reasonable to demand further justification for eliminating one over another (see [12] for related, if more pessimistic, arguments). We believe that the framework presented here can contribute to this positive project, but our immediate interest is in solidifying and clarifying limitative results about what cannot be done.

The issue of collapse becomes especially delicate when we turn to $\mathfrak{C}_2$. When do interventional distributions fully determine counterfactual distributions? In contrast to Prop. 2 we have:

Proposition 3. $\mathfrak{C}_2 \neq \varnothing$. That is, there exists an SCM in which Level 3 collapses to Level 2.

Proof sketch. As a very simple example in the finite case, any fully deterministic SCM will result in collapse. This is because, if $(\mu_\alpha)_{\alpha \in A}$ are all binary-valued then the measure $\mu \in \mathfrak{P}\big(\bigtimes_{\alpha \in A} \chi_{\mathbf{v}}\big)$ that produces the marginals μ_α is completely determined: each μ_α specifies an element of $\chi_{\mathbf{V}}$, so μ must assign unit probability to the tuple that matches μ_α at the α projection. In the infinite case, any example must be non-deterministic by atomlessness, but collapse is still possible; see Example 1 in Appendix B. $\qquad\square$

3.3 Probability of Necessity and Sufficiency

There is a particular counterfactual quantity that will play a starring role in the proof of our main result, thanks to its robustness against collapse. This quantity is known as the *probability of necessity and sufficiency* (PNS), commonly written $p(y_x, y'_{x'})$: the joint probability that Y would take on value y if X is set by intervention to x, and y' if X is set to x'. PNS has been thoroughly studied [26, 36, 2], in part due to its widespread relevance: from medical treatment to online advertising, we would like to

251 assess which interventions are likely to be both *necessary* and *sufficient* for a given outcome. Using the
252 notation from §2.3, PNS concerns the measure of sets $y_x, y'_{x'} = \pi^{-1}_{(X:=x,Y)}(\{y\}) \cap \pi^{-1}_{(X:=x',Y)}(\{y'\})$.

253 PNS is a paradigmatically Level 3 notion, and we will be interested in its manifestations at Level 2.
254 In that direction we introduce a small part of $\mathfrak{S}_2$, just enough to witness the behavior of Y (and X)
255 under the empty intervention and the two possible interventions on X:

256 **Definition 10.** Let $A^X_{\mathrm{NS}} = \{\varnothing := (), X := 0, X := 1\}$. We define a small subspace $\mathfrak{S}^X_{\mathrm{NS}(Y)} \subset$
257 $\bigtimes_{\alpha \in A^X_{\mathrm{NS}}} \mathfrak{P}(\chi_{\{X,Y\}})$, as the image of the map $\varpi^X_{\mathrm{NS}(Y)} = \left(\varsigma_{\{X,Y\}} \times \varsigma_{\{X,Y\}} \times \varsigma_{\{X,Y\}}\right) \circ \pi_{A^X_{\mathrm{NS}}}$ (see
258 Fig. 1(a)). Call $\mathfrak{S}^X_{\mathrm{NS}(Y)}$ a *PNS-space*; fixing X, we have a PNS-space for every Y.

259 It is known in the literature that PNS is not identifiable from the data $p(X, Y)$, $p(Y_x)$, and $p(Y_{x'})$
260 (see, e.g., [2]). As part of our proof of Theorem 3 below, we will strengthen this considerably to show
261 that PNS is *generically* unidentifiable, in a topological sense to be made precise.

4 The Weak Topology

263 We now demonstrate how $\mathfrak{S}_1, \mathfrak{S}_2$ and $\mathfrak{S}_3$ can be topologized. In general, given a space ϑ and the set
264 $\mathfrak{S} = \mathfrak{P}(\vartheta)$ of Borel probability measures on ϑ, a natural topology on $\mathfrak{S}$ can be defined as follows:

265 **Definition 11.** For a sequence $(\mu_n)_n$ of measures in $\mathfrak{S}$, write $(\mu_n)_n \Rightarrow \mu$ and say it *converges*
266 *weakly* [5, p. 7] to μ if $\int_\vartheta f \, d\mu_n \to \int_\vartheta f \, d\mu$ for all bounded, continuous $f : \vartheta \to \mathbb{R}$. Then the *weak*
267 *topology* τ^{w} on $\mathfrak{S}$ is that with the following closed sets: $E \subset \mathfrak{S}$ is closed in τ^{w} iff for any weakly
268 convergent sequence $(\mu_n)_n \Rightarrow \mu$ in which every $\mu_n \in E$, the limit point μ is in E.

269 There are several alternative characterizations of τ^{w}, which hold under very general conditions. For
270 instance, it coincides with the topology induced by the so called Lévy-Prohorov metric [5]. The most
271 useful characterization for our purposes is that it can be generated by subbasic open sets of the form

$$\{\mu : \mu(X) > r\} \tag{1}$$

272 with X ranging over basic clopens in ϑ and r over rationals (see [15, Lemma A.5] or [1, Eq. 28]).

273 Conceptually, the explication of τ^{w} in terms of weak convergence strongly suggests a connection
274 with statistical learning. We now make this connection precise, building on existing work [8, 14, 15].

4.1 Connection to Learning Theory

276 Roughly speaking, we will say a hypothesis $H \subseteq \mathfrak{S}$ is *statistically verifiable* if there is some error
277 bound ϵ and a sequence of statistical tests that converge on H with error at most ϵ, when data are
278 generated from H. More formally, a *test* is a function $\lambda : \vartheta^n \to \{\mathsf{accept}, \mathsf{reject}\}$, where ϑ^n is the
279 n-fold product of ϑ, viz. finite data streams from ϑ. The interest is in whether a "null" hypothesis
280 can be rejected given data observed thus far. The *boundary* of a set $A \subseteq \vartheta$, written $\mathrm{bd}(A)$, is the
281 difference of its closure and its interior. Intuitively, a learner will not be able to decide whether to
282 accept or reject on the boundary. Consequently it is assumed that λ is *feasible* in the sense that
283 the boundary of its acceptance zone (in the product topology on ϑ^n) always has measure 0, i.e.,
284 $\mu^n[\mathrm{bd}(\lambda^{-1}(\mathsf{accept}))] = 0$ for every $\mu \in \mathfrak{S}$, where μ^n is the n-fold product measure of μ.

285 Say a hypothesis $H \subseteq \mathfrak{S}$ is *(almost-surely) verifiable* [14] if there is $\epsilon > 0$ and a sequence $(\lambda_n)_{n \in \mathbb{N}}$
286 of feasible tests (of the complement of H in $\mathfrak{S}$, i.e., the "null hypothesis") such that

287   1. $\mu^n[\lambda_n^{-1}(\mathsf{reject})] \leq \epsilon$ for all n, whenever $\mu \notin H$;

288   2. $\lim_{n \to \infty} \mu^n[\lambda_n^{-1}(\mathsf{reject})] = 1$, whenever $\mu \in H$.

289 That is, to be verifiable we only require a sequence of tests that (almost-surely) converge on the true
290 hypothesis in the limit of infinite data (requirement 2), while incurring (type 1) error only up to a
291 given bound at finite stages (requirement 1). As an illustrative example, *conditional dependence* is
292 verifiable [14]. In general, this is a relatively lax notion of verifiability. For instance, a verifiable
293 hypothesis need not even be *decidable*, in that the learner may not know at any finite stage that they
294 are on track to convergence (cf. [8]). For our purposes this generality is a virtue: we want to show
295 that certain hypotheses are not statistically verifiable by any method, even in this wide sense. The
296 fundamental link between verifiability and the weak topology is the following, due to [14, 15]:

Theorem 1. A set $H \subseteq \mathfrak{S}$ is almost-surely verifiable if and only if it is open in the weak topology.

4.2 Topologizing Causal Models

We now reinterpret τ^{w} at each level of the causal hierarchy:

Definition 12. The *weak causal topology* τ_i^{w}, $1 \leq i \leq 3$, is the subspace topology on $\mathfrak{S}_i$, induced by

$$\text{if } i = 3 : \tau^{\mathrm{w}} \text{ on } \mathfrak{P}(\chi_{A \times \mathbf{V}}); \quad \text{if } i = 2 : \text{product of } \tau^{\mathrm{w}} \text{ on } \bigtimes_{\alpha \in A} \mathfrak{P}(\chi_{\mathbf{V}}); \quad \text{if } i = 1 : \tau^{\mathrm{w}} \text{ on } \mathfrak{P}(\chi_{\mathbf{V}}).$$

Proposition 4. All projections $\{\varpi_i\}_i, \varpi_{\mathrm{NS}(Y)}^X$ are continuous in the weak causal topologies.

A significant observation is that the learning theoretic interpretation, originally intended for τ_1^{w}, naturally extends to τ_2^{w}. While data streams at Level 1 amount to passive observations of $\mathbf{V}$, data streams at Level 2 can be seen as sequences of experimental results, i.e., observations of "potential outcomes" $\mathbf{Y_x}$. To make verifiability as easy as possible we assume a learner can observe a sample from all conceivable experiments at each step. A learner is thus a function $\lambda : \mathcal{E}^n \to \{\mathsf{accept, reject}\}$, where $\mathcal{E}^n = ((\chi_{\mathbf{V}})^n)_\alpha$ is the set of potential experimental observations over n trials (with α indexing the experiments). Construing $\mathcal{E}^n$ as a product space we can again speak of *feasibility* of λ.

Recall that elements of $\mathfrak{S}_2$ are tuples $(\mu_\alpha)_{\alpha \in A}$ of measures. Say a hypothesis $H \subseteq \mathfrak{S}_2$ is *experimentally verifiable* if there is $\epsilon > 0$ and a sequence $(\lambda_n)_{n \in \mathbb{N}}$ of feasible tests such that 1 and 2 above hold, replacing $\mu^n[\lambda_n^{-1}(\mathsf{reject})]$ with $\prod_\alpha \mu_\alpha^n[(\lambda_n^{-1}(\mathsf{reject}))_\alpha]$. That is, when experimental data are drawn from the interventional distributions $(\mu_\alpha)_{\alpha \in A} \in H$, we require that the learner eventually converge on H with bounded error at finite stages. We can then show (see Appendix C):

Theorem 2. A set $H \subseteq \mathfrak{S}_2$ is experimentally verifiable if and only if it is open in τ_2^{w}.

A similar result can be given for $(\mathfrak{S}_3, \tau_3^{\mathrm{w}})$, although it is less clear what the empirical content of this result would be. Note also that $\tau_1^{\mathrm{w}}, \tau_2^{\mathrm{w}}, \tau_3^{\mathrm{w}}$ give a sequence of increasingly fine topologies on the set of actual SCMs $\mathfrak{M}$ by simply pulling back the projections. The point is that τ_2^{w} is the finest that has clear empirical significance, while τ_3^{w} is the finest in terms of relevance to the causal hierarchy.

5 Collapse is Meager

Recall that a set $X \subseteq \vartheta$ is *nowhere dense* if every open set contains an open Y with $X \cap Y = \varnothing$. A countable union of nowhere dense sets is said to be *meager* (or *of first category*). The complement of a meager set is *comeager*. Intuitively, a meager set is one that can be "approximated" by sets "perforated with holes" [24]. Meagerness is notably preserved under continuous preimages.

As discussed above, one intuition highlighted by the weak topology τ^{w} is that open sets are the kinds of probabilistic propositions that could, in the limit of infinite data, be corroborated (Thms. 1, 2). Correlatively, meager sets in τ^{w} are so negligible as to be non-empirical: as a meager set contains no non-empty open subsets (by the Baire Category Theorem [24]), it is statistically unverifiable. We will now show that the injective collapse set $\mathfrak{C}_2$ from §3.2 is topologically meager.

The crux is to identify a "good" comeager PNS-subspace where collapse *never* occurs, with PNS as witness. In this subspace, the constraints circumscribing Level 3 have sufficient slack to allow tweaking PNS without thereby disturbing Level 2 (cf. Figure 2). Below, fix some $Y \neq X \in \mathbf{V}$, $x \neq x' \in \chi_X$, and $y \neq y' \in \chi_Y$. We define the good set as the locus of a set of strict inequalities:

Definition 13. A family $(\mu_\alpha)_{\alpha \in A_{\mathrm{NS}}^X} \in \mathfrak{S}_{\mathrm{NS}(Y)}^X$ is *Y-good* if we have the following, abbreviating the members of A_{NS}^X as $(), x, x'$:

$$0 < \mu_x(y') - \mu_{()}(x, y') < \mu_{()}(x'), \tag{2}$$

$$0 < \mu_{()}(x', y') < \mu_{()}(x'). \tag{3}$$

Lemma 1. The subspace of Y-good families is comeager in $\mathfrak{S}_{\mathrm{NS}(Y)}^X$.

Proof sketch. The non-strict versions of (2), (3) hold universally, so the complement of the good set is defined by equalities. This is closed and contains no nonempty open by the weak subbasis (1). $\square$

	$\mathcal{M}$			
u	$P(u)$	X_u	$Y_{x,u}$	$Y_{x',u}$
u_0	$^{1}/_{2}$	x'	y	y
u_1	$^{1}/_{2}$	x'	y'	y'

(a) Y-good Model

	$\mathcal{M}'$			
u	$P(u)$	X_u	$Y_{x,u}$	$Y_{x',u}$
u_0	$^{1}/_{2} - \varepsilon$	x'	y	y
u_1	ε	x'	y	y'
u_2	$^{1}/_{2} - \varepsilon$	x'	y'	y'
u_3	ε	x'	y'	y

(b) Example Separating Levels 2 and 3

Figure 2: (a): the structural functions and exogenous noise for a model $\mathcal{M}$ with direct influence $X \to Y$. This $\mathcal{M}$ meets (2) and (3), so we may apply Lemma 2, constructing the model $\mathcal{M}'$ in (b), where $0 < \varepsilon < ^{1}/_{2}$. Note that $p_{\mathrm{cf}}^{\mathcal{M}}(y_x, y'_{x'}) = 0$ while $p_{\mathrm{cf}}^{\mathcal{M}'}(y_x, y'_{x'}) = \varepsilon$, so that the two models disagree on a Level 3 PNS quantity; on the other hand, it is easy to check agreement on all of Level 2.

Figure 2 presents the construction in a small, two-variable case, and Lemma 2 below is proven by generalizing it to arbitrary $\mathbf{V}$. Guaranteeing agreement on every interventional distribution in the general case is subtle (Appendix D): it has been observed that enlarging $\mathbf{V}$ can enable additional inferences (e.g., [31]), though the next result reflects a dependence on further assumptions.

Lemma 2. Suppose $\prec$ is an order in which X comes first and $\mu_2 \in \mathfrak{S}_2^{\prec}$ is such that $\varpi_{\mathrm{NS}(Y)}^X(\mu_2)$ is Y-good. Then for any $\mu_3 \in \mathfrak{S}_3^{\prec}$ such that $\varpi_2(\mu_3) = \mu_2$, there exists a $\mu'_3 \in \mathfrak{S}_3^{\prec}$ such that $\varpi_2(\mu'_3) = \mu_2$ but $\mu'_3(y_x, y'_{x'}) \neq \mu_3(y_x, y'_{x'})$.

The main theorem and its important learning-theoretic corollary are now straightforward.

Theorem 3 (Topological Hierarchy). The set $\mathfrak{C}_2$ of points where all Level 3 facts are identifiable from Level 2 is meager in $(\mathfrak{S}_2, \tau_2^{\mathrm{w}})$. The preimage $\varpi_2^{-1}(\mathfrak{C}_2) = \mathfrak{C}_3$ is likewise meager in $(\mathfrak{S}_3, \tau_3^{\mathrm{w}})$.

Proof. Let $\mathfrak{D}_2^{X,Y} \subset \mathfrak{S}_2^X$ be the preimage under $\varpi_{\mathrm{NS}(Y)}^X$ of the set of Y-good tuples in $\mathfrak{S}_{\mathrm{NS}(Y)}^X$. Lemma 2 implies that $\mathfrak{C}_2 \cap \mathfrak{S}_2^X$ is contained in $\mathfrak{S}_2^X \setminus \mathfrak{D}_2^{X,Y}$, for *any* $Y \neq X$. Meanwhile, since $\varpi_{\mathrm{NS}(Y)}^X$ is continuous, Lemma 1 implies that $\mathfrak{S}_2^X \setminus \mathfrak{D}_2^{X,Y}$ is meager in $\mathfrak{S}_2^X$, and thereby also in $\mathfrak{S}_2$. Thus $\mathfrak{C}_2 = \bigcup_{X \in \mathbf{V}} \mathfrak{C}_2 \cap \mathfrak{S}_2^X$ is a countable union of meager sets, and hence meager. $\square$

Corollary 1. No causal hypothesis licensing arbitrary counterfactual (and specifically PNS) inferences from observational and experimental data is itself statistically (even experimentally) verifiable.

6   Conclusion

We introduced a general framework for topologizing spaces of causal models, including the space of all (discrete, well-founded) causal models. As an illustration of the framework we characterized levels of the causal hierarchy topologically, and proved a topological version of the causal hierarchy theorem from [3]. While the latter shows that collapse of the hierarchy (specifically of Level 3 to Level 2) is *exceedingly unlikely* in the sense of (Lebesgue) measure, we offer a complementary result: any condition guaranteeing that we could infer arbitrary Level 3 information from purely Level 2 information must be *statistically unverifiable*, even by experimental means. Both results capture an important sense in which collapse is "negligible" in the space of all possible models. As an added benefit, the topological approach extends seamlessly to the setting of infinitely many variables.

There are many natural extensions of these results. For instance, we have begun work on a version for continuous endogenous variables. Also of interest are subspaces embodying familiar causal assumptions or other well-studied coarsenings of SCMs (see, e.g., [22] on Bayesian networks, or [16] on linear non-Gaussian models), which often render important inference problems solvable, though sometimes only "generically" so. As emphasized by [3], a causal hierarchy theorem should not be construed as a purely limitative result, but rather as further motivation for understanding the whole range of causal-inductive assumptions, how they relate, and what they afford. We submit that the topological constructions presented here can help clarify and systematize this broader lanscape.

References

[1] N. L. Ackerman, C. E. Freer, and D. M. Roy. On the computability of conditional probability. *Journal of the ACM*, 66(3), 2019.

[2] C. Avin, I. Shpitser, and J. Pearl. Identifiability of path-specific effects. In *Proceedings of IJCAI*, 2005.

[3] E. Bareinboim, J. D. Correa, D. Ibeling, and T. Icard. On Pearl's hierarchy and the foundations of causal inference. Technical Report R-60, Causal AI Lab, Columbia University, 2020.

[4] G. Belot. Absolutely no free lunches! *Theoretical Computer Science*, 845:159–180, 2020.

[5] P. Billingsley. *Convergence of Probability Measures*. Wiley, 2nd edition, 1999.

[6] V. I. Bogachev. *Measure Theory*. Springer Berlin Heidelberg, 2007.

[7] N. Cartwright. *Nature's Capacities and their Measurements*. Clarendon Press, 1989.

[8] A. Dembo and Y. Peres. A topological criterion for hypothesis testing. *Annals of Statistics*, 22(1):106–117, 1994.

[9] P. Diaconis and D. Freedman. On the consistency of Bayes estimates. *Annals of Statistics*, 14(1):1–26, 1986.

[10] M. Drton, R. Foygel, and S. Sullivant. Global identifiability of linear structural equation models. *Annals of Statistics*, 39(2):865–886, 2011.

[11] F. Eberhardt, C. Glymour, and R. Scheines. On the number of experiments sufficient and in the worst case necessary to identify all causal relations among n variables. In *Proceedings of UAI*, page 178–184, 2005.

[12] D. Freedman. From association to causation via regression. *Advances in Applied Mathematics*, 18:59–110, 1997.

[13] D. Galles and J. Pearl. An axiomatic characterization of causal counterfactuals. *Foundations of Science*, 3 (1):151–182, Jan 1998.

[14] K. Genin. *The Topology of Statistical Inquiry*. PhD thesis, Carnegie Mellon University, 2018.

[15] K. Genin and K. Kelly. The topology of statistical verifiability. In *Proceedings of TARK*, pages 236–250, 2017.

[16] K. Genin and C. Mayo-Wilson. Statistical decidability in linear, non-Gaussian models. In *Causal Discovery & Causality-Inspired Machine Learning*, NeurIPS, 2020.

[17] D. Ibeling and T. Icard. On open-universe causal reasoning. In *Proceedings of UAI*, 2019.

[18] D. Ibeling and T. Icard. Probabilistic reasoning across the causal hierarchy. In *Proceedings of AAAI*, 2020.

[19] G. W. Imbens and D. B. Rubin. *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press, 2015.

[20] J. L. Kelley. *General Topology*. Springer, 1975.

[21] S. Lee, J. Correa, and E. Bareinboim. General identifiability with arbitrary surrogate experiments. In *Proceedings of UAI*, 2019.

[22] H. Lin and J. Zhang. On learning causal structures from non-experimental data without any faithfulness assumption. In *Proceedings of ALT*, pages 554–582, 2020.

[23] C. Meek. Strong completeness and faithfulness in Bayesian networks. In *Proceedings of UAI*, page 411–418, 1995.

[24] J. C. Oxtoby. *Measure and Category*. Springer, 1971.

[25] J. Pearl. Causal diagrams for empirical research. *Biometrika*, 82(4):669–710, 1995.

[26] J. Pearl. Probabilities of causation: Three counterfactual interpretations and their identification. *Synthese*, 121(1):93–149, 1999.

[27] J. Pearl. *Causality*. Cambridge University Press, 2009.

[28] J. Pearl and D. Mackenzie. *The book of why: The new science of cause and effect*. Basic Books, 2018.

[29] J. Peters, D. Janzing, and B. Schölkopf. *Elements of Causal Inference: Foundations and Learning Algorithms*. MIT Press, 2017.

[30] J. Robins and S. Greenland. The probability of causation under a stochastic model for individual risk. *Biometrics*, 45(4):1125–1138, 1989.

[31] B. Schölkopf, F. Locatello, S. Bauer, N. R. Ke, N. Kalchbrenner, A. Goyal, and Y. Bengio. Toward causal representation learning. *Proceedings of the IEEE*, 109(5):612–634, 2021.

[32] S. Shalev-Shwartz and S. Ben-David. *Understanding Machine Learning: From Theory to Algorithms*. Cambridge University Press, 2014.

[33] S. Shimizu, P. O. Hoyer, A. Hyvärinen, and A. Kerminen. A linear non-Gaussian acyclic model for causal discovery. *Journal of Machine Learning Research*, 7(72):2003–2030, 2006.

[34] P. Spirtes, C. Glymour, and R. Scheines. *Causation, Prediction, and Search*. MIT Press, 2001.

[35] Z. Tavares, J. Koppel, X. Zhang, R. Das, and A. S. Lezama. A language for counterfactual generative models. In *Proceedings of ICML*, 2021.

[36] J. Tian and J. Pearl. Probabilities of causation: Bounds and identification. *Annals of Mathematics and Artificial Intelligence*, 28(1):287–313, 2000.

[37] J. Tian and J. Pearl. A general identification condition for causal effects. In *Proceedings of AAAI*. 2002.