Sampling in Constrained Domains with Orthogonal-Space Variational Gradient Descent

Abstract

Sampling methods, as important inference and learning techniques, are typically designed for unconstrained domains. However, constraints are ubiquitous in machine learning problems, such as those on safety, fairness, robustness, and many other properties that must be satisfied to apply sampling results in real-life applications. Enforcing these constraints often leads to implicitly-defined manifolds, making efficient sampling with constraints very challenging. In this paper, we propose a new variational framework with a designed orthogonal-space gradient flow (O-Gradient) for sampling on a manifold $\mathcal{G}_0$ defined by general equality constraints. O-Gradient decomposes the gradient into two parts: one decreases the distance to $\mathcal{G}_0$ and the other decreases the KL divergence in the orthogonal space. While most existing manifold sampling methods require initialization on $\mathcal{G}_0$, O-Gradient does not require such prior knowledge. We prove that O-Gradient converges to the target constrained distribution with rate $\widetilde{O}(1/\text{the number of iterations})$ under mild conditions. Our proof relies on a new Stein characterization of conditional measure which could be of independent interest. We implement O-Gradient through both Langevin dynamics and Stein variational gradient descent and demonstrate its effectiveness in various experiments, including Bayesian deep neural networks.

1 Introduction

Sampling methods, such as Markov chain Monte Carlo (MCMC) [1] and Stein variational gradient descent (SVGD) [22, 21], have been widely used for getting samples from or approximating intractable distributions in machine learning (ML) problems, such as estimating Bayesian neural network posteriors [35], generating new images [30], and training energy-based models [14]. While being powerful, most sampling methods usually can only be used in unconstrained domains or some special geometric spaces. This greatly limits the application of sampling to many real-life tasks.

We consider sampling from a distribution π with an equality constraint $g(x) = 0$ where $g : \mathbb{R}^d \to \mathbb{R}$ is a general differentiable function. The domain in this case is the level set $\mathcal{G}_0 = \{x \in \mathbb{R}^d : g(x) = 0\}$ which is a submanifold in $\mathbb{R}^d$. We do not require additional information about $\mathcal{G}_0$, such as explicit parameterization or known in-domain points, which is in contrast, often demanded by previous methods [3, 32, 2, 18]. The problem defined above includes many ML applications, such as disease diagnosis with logic rules constraint, policymaking with fairness constraint for different demographic subgroups, and autonomous driving with robustness constraint to unseen scenarios.

In this paper, we propose a new variational framework which transforms the above constrained sampling problem into a constrained functional minimization problem. A special gradient flow, denoted *orthogonal-space gradient flow* (O-Gradient), is developed to minimize the objective. As illustrated in Figure 1a, the direction of O-Gradient v can be decomposed into two parts: the first

| (a) O-Gradient | (b) O-Langevin | (c) O-SVGD |

Figure 1: Visualization of our methods. (a) O-Gradient v is formed by $v_\sharp$ which follows ∇g, and $v_\perp$ which is perpendicular to ∇g. (b)-(c) Applying O-Gradient to Langevin dynamics and SVGD. Both methods can approach the manifold and sample on it.

part $v_\sharp$ drives the sampler towards the manifold $\mathcal{G}_0$ following ∇g and keeps it on $\mathcal{G}_0$ once arrived; the second part $v_\perp$ makes the sampler explore $\mathcal{G}_0$ following the density $\pi(x)$. We prove the convergence of O-Gradient in the continuous-time mean-field limit. O-Gradient can be applied to both Langevin dynamics and SVGD, resulting in O-Langevin and O-SVGD respectively. As shown in Figure 1b&c, both methods can converge to the target distribution on the manifold. In particular, O-Langevin converges following a noisy trajectory while O-SVGD converges smoothly, similar to their standard unconstrained counterparts. We empirically demonstrate the sampling performance of O-Langevin and O-SVGD across different constrained ML problems. We summarize our contributions as follows:

- We reformulate the hard-constrained sampling problem into a functional optimization problem and derive a special gradient flow, O-Gradient, to obtain the solution.
- We prove that O-Gradient converges to the target constrained distribution with rate $\widetilde{O}(1/\text{the number of iterations})$ under mild conditions. Our proof technique includes a new Stein characterization of conditional measure which could be of independent interest.
- We implement O-Gradient through both Langevin dynamics and SVGD and demonstrate its effectiveness in various experiments, including a constrained synthetic distribution, income classification with fairness constraint, loan classification with logic rules and image classification with robust Bayesian deep neural networks.

2 Related Work

Sampling on Explicitly Defined Manifolds Manifolds with special shapes, such as geometric or physics structures, can sometimes be explicitly parameterized in lower dimension spaces. For example, a torus embedded in $\mathbb{R}^3$ can be explicitly defined in two dimensions using polar coordinates. Variants of classical methods have been developed to sample on such manifolds, including rejection sampling [6], Langevin dynamics [32], Hamiltonian Monte Carlo (HMC) [3] and Riemannian manifold HMC [15, 27]. However, explicit parameterization is only applicable to a few special cases and cannot be used for general machine learning problems. In contrast to this line of work, our method is able to work with more general manifolds defined in the original domain $\mathbb{R}^d$.

Sampling on Implicitly Defined Manifolds Many common applications are not endowed with simple manifolds, such as molecular dynamics [16], matrix factorization [26] and free energy calculations [31]. Motivated by these applications, sampling methods on implicitly defined manifolds have been developed. Brubaker et al. [2] has proposed a family of constrained MCMC methods by applying Lagrangian mechanics to Hamiltonian dynamics. Zappa et al. [34] has introduced a constrained Metropolis-Hastings (MH) with a reverse projection check to ensure the reversibility. Later, this method has been extended to HMC [18] and multiple projections [19]. However, the implementation and analysis of these methods often assume the algorithm starts on the manifold and never leaves it, requiring prior known points on the manifold and expensive projection subroutines, such as Newton's method [2, 34, 17–19] or a long time ordinary differential equation (ODE) [36, 28, 13]. In contrast, our method works with distributions supported on the ambient space and thus gets rid of the above strong assumptions, leading to a much faster update per iteration. This makes our method especially suitable for complex ML models such as deep neural networks.

75 **Sampling with a Moment Constraint** Recently, sampling with a general moment constraint, such
76 as $\mathbb{E}_q[g] \leq 0$ where q is the approximated distribution, has been studied [24]. However, this type
77 of constraint can not guarantee every sample to satisfy $g(x) = 0$. From a technical view, the target
78 distribution with a moment constraint is usually not singular w.r.t. π, so the problem is conceptually
79 less challenging compared to the problem considered in this work.

3 Preliminaries

81 **Variational Framework** We review the derivation of Langevin dynamics and SVGD from a
82 unified variational framework. The variational approach frames the problem of sampling into a KL
83 divergence minimization problem: $\min_{q \in \mathcal{P}} \mathrm{KL}(q \,||\, \pi)$ where $\mathcal{P}$ is the space of probability measures.
84 Starting from an initial q_0 and $x_0 \sim q_0$, assume we update x_t following $dx_t = v_t(x_t)dt$, where
85 $v_t \colon \mathbb{R}^d \to \mathbb{R}^d$ is a velocity field at time t. The density q_t of x_t follows Fokker-Planck equation:
86 $dq_t/dt = -\nabla \cdot (v_t q_t)$, and the KL divergence decreases with the following rate [22]:

$$-\frac{d}{dt}\mathrm{KL}(q_t \,||\, \pi) = \mathbb{E}_{q_t}[\mathcal{A}_\pi v_t] = \mathbb{E}_{q_t}[(s_\pi - s_{q_t})^\top v_t], \tag{1}$$

87 where $\mathcal{A}_\pi v(x) = s_\pi(x)^\top v(x) + \nabla \cdot v(x)$ is the Stein operator, and $s_p = \nabla \log p$ is the score function
88 of distribution p. The optimal v_t is obtained by solving an optimization in a Hilbert space $\mathcal{H}$,

$$\max_{v \in \mathcal{H}} \mathbb{E}_{q_t}[(s_\pi - s_{q_t})^\top v] - \frac{1}{2} \|v\|_\mathcal{H}^2. \tag{2}$$

89 The above objective makes sure that v_t decreases the KL divergence as fast as possible.

90 **Langevin Dynamics and SVGD Algorithms** Both Langevin dynamics and SVGD can be derived
91 from this variational framework by taking $\mathcal{H}$ to be different spaces. Taking $\mathcal{H}$ to be $\mathcal{L}_q^2$, the velocity
92 field becomes $v_t(\cdot) = \nabla s_\pi(\cdot) - \nabla q_t(\cdot)$ which can be simulated by Langevin dynamics $dx_t =$
93 $s_\pi(x_t)dt + dW_t$ with W_t being a standard Browninan motion. After discretization with a step size
94 $\eta > 0$, the update step of Langevin dynamics is $x_{t+1} = x_t + \mathrm{Langevin}(x_t)$, where

$$\mathrm{Langevin}(x_t) = \eta \nabla \log \pi(x_t) + \sqrt{2\eta}\xi_t, \ \ \xi_t \sim \mathcal{N}(0, I). \tag{3}$$

95 Taking $\mathcal{H}$ to be the reproducing kernel Hilbert space (RKHS) of a continuously differentiable kernel
96 $k \colon \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$, the velocity field becomes $v_t(\cdot) = \mathbb{E}_{x \sim q_t}[k_t(\cdot, x)s_\pi(x) + \nabla_x k_t(\cdot, x)]$. After
97 discretization, the update step of SVGD for particles $\{x_i\}_{i=1}^n$ is $x_{i,t+1} = x_{i,t} + \eta \cdot \mathrm{SVGD}_k(x_{i,t})$,
98 for $i = 1, \ldots, n$, where η is a step size and

$$\mathrm{SVGD}_k(x_{i,t}) = \frac{1}{n} \sum_{j=1}^n k(x_{i,t}, x_{j,t}) \nabla_{x_{j,t}} \log \pi(x_{j,t}) + \nabla_{x_{j,t}} k(x_{i,t}, x_{j,t}). \tag{4}$$

4 Main Method

100 In this section, we formulate the constrained sampling problem into a constrained optimization
101 through the variational lens in Section 4.1, and introduce a new gradient flow to solve the problem
102 in Section 4.2. We apply this general framework to Langevin dynamics and SVGD, leading to two
103 practical algorithms in Section 4.3.

4.1 Constrained Variational Optimization

105 Recall that our goal is to draw samples according to the probability of π, but restricted to a low
106 dimensional manifold specified by an equality: $\mathcal{G}_0 := \{x \in \mathbb{R}^d \colon g(x) = 0\}$. Similar to the standard
107 variational framework in Section 3, we can formulate the problem into a constrained optimization in
108 the space of probability measures: $\min_{q \in \mathcal{P}} \mathrm{KL}(q \,||\, \pi)$, s.t. $q(g(x) = 0) = 1$. However, this problem
109 is in general ill-posed, because when q satisfies the constraint, q is singular w.r.t. π, so both $\frac{dq}{d\pi}$ and
110 $\mathrm{KL}(q \,||\, \pi)$ are not defined. Although the problem is ill-posed, we are actually still able to derive
111 a KL-gradient flow to solve the problem by considering q supported on $\mathbb{R}^d$. The intuition of the
112 derivation is that, in addition to minimizing the objective as in Eq. (1), the velocity filed v_t should also
113 push q towards $\mathcal{G}_0$ to satisfy the constraint. Surprisingly, the distribution q_t following such a gradient
114 flow indeed converges to the target distribution on the manifold. We will focus on the derivation of
115 the gradient flow here and leave its rigorous justification in Section 5.

4.2 Orthogonal-Space Gradient Flow (O-Gradient)

As mentioned above, besides maximizing the decay of $\mathrm{KL}(q \,\|\, \pi)$, the velocity field v_t also needs to drive q towards the manifold satisfying $g(x) = 0$. In particular, we add to (2) a requirement that value of $g(x)$ is driven towards 0 with a given rate:

$$v_t = \arg\max_{v \in \mathcal{H}} \mathbb{E}_{q_t}[(s_\pi - s_{q_t})^\top v] - \frac{1}{2} \|v\|_{\mathcal{H}}^2, \quad s.t \ \ v_t(x)^\top \nabla g(x) = -\psi(g(x)) \tag{5}$$

where $\psi(x)$ is an increasing odd function. We use $\psi(x) = \alpha \mathrm{sign}(x)|x|^{1+\beta}$ with $\alpha > 0$ and $\beta \in (0, 1]$ in this paper. The objective is the same as Eq. (2) in the standard variational framework while the constraint ensures that v_t pushes q_t towards the manifold and keeps it stay. It is easy to see that the solution of the above problem can be decomposed as $v_t = v_\sharp + v_\perp$ where

$$v_\sharp(x) = \frac{-\psi(g(x))\nabla g(x)}{\|\nabla g(x)\|^2}, \quad v_\perp \perp \nabla g, \tag{6}$$

where we use $f \perp g$ to denote (pointwise) orthogonality: $f(x)^\top g(x) = 0, \forall x \in \mathbb{R}^d$. Note that $v_\sharp$ is parallel to ∇g and the remaining is to determine $v_\perp$. Note that $v_\perp$ can be represented as a projection of an arbitrary function u to the orthogonal space of ∇g:

$$v_\perp = D(x)u(x), \quad \text{where } D(x) := I - \frac{\nabla g(x) \nabla g(x)^\top}{\|\nabla g(x)\|^2}. \tag{7}$$

The projection operator D makes sure that $v_\perp \perp \nabla g$ holds for any u, of which the optimal value we can get by maximizing the unconstrained objective in Eq. (2),

$$\max_{v_\perp} E_{q_t}[(s_\pi - s_{q_t})^\top (v_\sharp + v_\perp)] - \frac{1}{2}\|v_\perp\|_{\mathcal{H}}^2 \Rightarrow \max_u \mathbb{E}_{q_t}[(D(s_\pi - s_{q_t}))^\top u] - \frac{1}{2}\|Du\|_{\mathcal{H}}^2. \tag{8}$$

The optimal solution of u depends on the choice of space $\mathcal{H}$, which we discuss in Section 4.3.

Overall, we obtain the velocity field v_t by first formulating a constrained optimization and then transforming it into an unconstrained optimization via orthogonal decomposition. We call v_t *Orthogonal-Space Gradient Flow* (O-Gradient) and it drives q_t to the target distribution when only knowing ∇g, requiring no explicit representation of the manifold $\mathcal{G}_0$.

4.3 Practical Algorithms

After deriving O-Gradient for general Hilbert spaces $\mathcal{H}$, we explain how to implement it using SVGD and Langevin dynamics. The resulting O-SVGD and O-Langevin are outlined in Algorithm 1. At a high level, our algorithms keep the original SVGD or Langevin dynamics movement in the directions perpendicular to ∇g, while pushing the density towards $\mathcal{G}_0$ along the ∇g direction.

O-SVGD We apply O-Gradient to SVGD first since it is fairly straightforward. Recall that $v_\sharp$ can be obtained using Eq. (6). We solve Eq. (8) to get $v_\perp$ through the following lemma.

Lemma 4.1. *When $\mathcal{H}$ is an RKHS with kernel $k \colon \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$, a solution to Eq. (8) is $v_\perp = Du(x) = \mathbb{E}_{y \sim q_t}(k_\perp(x, y)s_\pi(y) + \nabla_y \cdot k_\perp(x, y))$ with the orthogonal-space kernel $k_\perp(x, y) = k(x, y)D(x)D(y)$. Here $k_\perp \colon \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}^{d \times d}$ is matrix valued, and $\nabla_y \cdot k_\perp = \sum_j \partial_{y_j} k_\perp^{ij}(x, y)$.*

Then the combined velocity is obtained using the original SVGD with the kernel $k_\perp$,

$$v_t(x) = v_\sharp(x) + \int k_\perp(x, y)s_\pi(y)q_t(y)dy + \int \nabla_y \cdot (k_\perp(x, y))q_t(y)dy.$$

Numerically, we iteratively update a set of n particles $\{x_{i,t}\}_{i=1}^n \subset \mathbb{R}^d$, such that its empirical distribution $\sum_{i=1}^n \delta_{\theta_{i,t}}/n$ is an approximation of q_t in a proper sense when step size $\eta \to 0$ and particle size $n \to +\infty$. Similar to the update of standard SVGD in Eq. (4), the update of O-SVGD is $x_{i,t+1} = x_{i,t} + \eta \cdot (v_\sharp(x_{i,t}) + \mathrm{SVGD}_{k_\perp}(x_{i,t}))$ where

$$\mathrm{SVGD}_{k_\perp}(x_{i,t}) = \frac{1}{n}\sum_{j=1}^n k_\perp(x_{i,t}, x_{j,t})\nabla_{x_{j,t}} \log \pi(x_{j,t}) + \nabla_{x_{j,t}} k_\perp(x_{i,t}, x_{j,t}). \tag{9}$$

It is worth noting that $\mathrm{SVGD}_{k_\perp}$ is identical to Eq. 4 but with kernel $k_\perp$ rather than k.

Algorithm 1 O-SVGD and O-Langevin.

---

given: Initialization $\{x_{i,0}\}_{i=1}^n$ for O-SVGD and x_0 for O-Langevin, step size η.
loop

    compute $v_\sharp(x) = \frac{-\psi(g(x))\nabla g(x)}{\|\nabla g(x)\|^2}$ and the projection operator $D(x) = I - \frac{\nabla g(x)\nabla g(x)^\top}{\|\nabla g(x)\|^2}$.
    if `O-SVGD`, update $\{x_{i,t}\}_{i=1}^n$ by Eq. (9).
    if `O-Langevin`, update x_t by Eq. (12).
end loop

O-Langevin The Langevin implementation requires some additional derivation. First of all, with $\mathcal{H} = L_q^2$, we can show that the optimal velocity field is $v_t(x) = \phi(x) - D(x)s_{q_t}(x)$ where $\phi(x) = v_\sharp(x) + D(x)s_\pi(x)$. This leads to a density flow

$$\frac{d}{dt}q_t(x) = -\nabla \cdot (\phi(x)q_t(x)) + \nabla \cdot (D(x)\nabla q_t(x)). \tag{10}$$

Next, we try to design a stochastic differential equation (SDE) of which the Fokker–Plank equation (FPE) is identical to Eq. (10). The result is given by the following:

Theorem 4.2. *Consider a vector field $r(x) = \nabla \cdot D(x)$, or its component-wise formulation $r_i(x) = \sum_{j=1}^d \partial_{x_j} D_{i,j}(x)$, where x_j denotes the jth dimension of x. Consider the SDE*

$$dx_t = (\phi(x_t) + r(x_t))dt + \sqrt{2}D(x_t)dW_t \tag{11}$$

with $\phi(x) = v_\sharp(x) + D(x)s_\pi(x)$, then its FPE is identical to Eq. (10). Moreover, i) the value $g(x_t)$ has deterministic decay $\frac{d}{dt}g(x_t) = -\psi(x_t)$; ii) for any f with $\nabla f \perp \nabla g = 0$, the generator of x_t satisfies the Langevin equation: $\frac{d}{dt}\mathbb{E}[f(x_t)|x_0 = x]\big|_{t=0} := \mathcal{L}f(x) = \nabla f^\top(x)s_\pi(x) + \Delta f(x)$.

It is worth pointing out that if $g(x_0) = 0$, then $g(x_t) \equiv 0$, that is, x_t always stays on $\mathcal{G}_0$. In this case, Eq. (11) degenerates to manifold Langevin dynamics studied in previous work [9, 32]. However, our SDE does not have this requirement since *it is still well-defined off* $\mathcal{G}_0$. This is especially useful for numerical implementations, leading to a fast algorithm without expensive projection steps.

Similar to the standard Langevin dynamics update in Eq. (3), the update rule of O-Langevin is $x_{t+1} = x_t + \eta \cdot v_\sharp(x_t) + \text{Langevin}_\perp(x_t)$ where

$$\text{Langevin}_\perp(x_t) = \eta D(x_t)s_\pi(x_t) + \eta r(x_t) + \sqrt{2\eta}D(x_t)\xi_t, \ \ \xi_t \sim \mathcal{N}(0, I). \tag{12}$$

5 Theoretical Analysis

We theoretically justify the convergence of O-Gradient in this section. To do so, we first describe the target measure as a conditioned measure Π_0, then derive its associated orthogonal-space Fisher divergence, and finally prove that O-Gradient converges to Π_0.

5.1 Conditioned measure and its Stein characterization

First of all, conditioning on a zero-measure set is a challenging concept. Assume we have a distribution Π with density π. Let A be a set with $\Pi(A) \neq 0$, then $\Pi(B|A) = \frac{\Pi(B \cap A)}{\Pi(A)}$. However, if $\Pi(A) = 0$, this definition is ill-posed. Instead, a standard way to define a conditioned measure on a zero measure set is using disintegration theorem [5]. Let $\Pi_z(\cdot) = \Pi(\cdot|g(x) = z)$ be the measure such that

$$\mathbb{E}_\Pi[f(x)] = \mathbb{E}_{z \sim \Pi^g}\mathbb{E}_{x \sim \Pi_z}[f(x)], \ \ \forall f, \tag{13}$$

where Π^g is the pushforward measure of Π under g. It is worth noting that Π_0 *is not* the measure with Hausdorff density π on $\mathcal{G}_0$. Instead, under regularity conditions, Π_0 can also be defined through Hausdorff density $\pi(x)/|\nabla g(x)|$ on $\mathcal{G}_0$, see Lemma 6.4.1 of [29].

One can also avoid this rather abstract definition, and consider an alternative definition which is more intuitive and Bayesian. Suppose we observe $z = g(x) + \sigma\xi$, where $\xi \sim \mathcal{N}(0, 1)$. Under prior $x \sim \pi$, the posterior $\pi_{\sigma^2,z}(x) \propto \pi(x)\exp(-|z - g(x)|^2/2\sigma^2)$. Then we can show that Π_z is the (weak) limit of $\pi_{\sigma^2,z}$ as $\sigma \to 0$, moreover, it satisfies a *Stein characterization of conditional measure*. We need some regularity assumptions to state our results.

Definition 5.1. *A density q is g-regular if there is a L so that for small enough $\eta > 0$*

$$|\mathbb{E}_{\xi \sim \mathcal{N}(0,1)}[q^g(z + \eta\xi)/q^g(z)] - 1| \le L\eta, \quad \forall z. \tag{14}$$

Note that Eq. (14) holds if q^g is a Lipschitz function. Moreover, while Definition 5.1 requires Eq. (14) to hold for all z, in practice we only concern Π_0, thus only need the equation to hold for z near 0.

Proposition 5.2. *Suppose π is g-regular. Then the weak limit of $\pi_{\eta,z}(x) \propto \pi(x)\exp(-\frac{1}{2\eta}(g(x)-z)^2)$ as $\eta \to 0$ concentrates on $\mathcal{G}_z$ and satisfies Eq. (13). Moreover, for each z,*

$$\mathbb{E}_{\Pi_z}[\mathcal{A}_\pi \phi] = 0, \quad \forall \phi \perp \nabla g. \tag{15}$$

The first part of Proposition 5.2 also provides us with a way to sample Π_z, as we can sample a sequence of distribution $\pi_{\eta_k,z}$ with $\eta_k \to 0$. However, this method involves double loops and is usually much more expensive than single-loop algorithms.

The second part of Proposition 5.2 can be used to formulate as an orthogonal-space Stein equation. Specifically, given a q concentrated on $\mathcal{G}_z$, we consider checking whether $\mathbb{E}_q[\mathcal{A}_\pi\phi]$ is zero for $\phi \in \mathcal{H}_\perp = \{\phi : \phi \perp \nabla g\}$, which is a necessary condition for $q = \Pi_z$. Notably, this equation is well defined in $\mathbb{R}^d$ without using any parameterization of $\mathcal{G}_z$.

Compared with the standard Stein equation, we add an additional restriction that $\phi \in \mathcal{H}_\perp$. We argue such restriction is actually very natural. To see that, we note Π_z is concentrated on $\mathcal{G}_z$, so we should check the Stein equation with vector field that "live" only on $\mathcal{G}_z$, namely the tangent bundle $\mathcal{T}\mathcal{G}_z$. Notably, if $\phi(x) \in \mathcal{T}\mathcal{G}_z$, then $\phi(x) \perp \nabla g(x)$, therefore it is natural to consider $\mathcal{H}_\perp$.

5.2 Orthogonal-Space Fisher divergence

Next we consider how to generalize Fisher divergence for constrained sampling. Since the orthogonal-space Stein equation can only discern discrepancy in $\mathcal{H}_\perp$, it is natural to consider an orthogonal-space Fisher divergence

$$F_\perp(q, \pi) := \|D(s_q - s_\pi)\|_{\mathcal{H}}^2, \tag{16}$$

where D is the $\mathcal{H}_\perp$-projection operator defined pointwisely using Eq. (7). Its easy to see $F_\perp$ is well defined for a density q on $\mathbb{R}^d$. But it remains unclear whether it can be used to measure the distance between q and Π_z.

To answer this question, we need some form of strong convexity conditions. Recall that in the unconstrained case, such conditions can be characterized by Poincaré inequalities (PI) when $\mathcal{H}$ is L_q^2, so we will focus on this setting. PI counterparts of SVGD remain an open question [10, 7], which we hope to be addressed by future works. When constrained on $\mathcal{G}_z$, PI's formulation involves gradients so it depends on the Riemannian metric used in $\mathcal{G}_z$. One natural choice of Riemannian metric is the metric inherited from $\mathbb{R}^d$ which equals $\|D(x)\nabla f(x)\|$, since $D(x)$ is also the projection onto the tangent bundle of $\mathcal{G}_z$. This leads to the following definition.

Definition 5.3. *Π_z follows a κ-PI, if for any smooth f on $\mathbb{R}^d$, $\mathrm{var}_{\Pi_z}[f] \le \kappa \mathbb{E}_{\Pi_z}[\|D\nabla f\|^2]$.*

PI on Riemannian manifold has been studied in previous work [11, 20]. In particular, Theorem 2.10 of [11] shows that PI holds for the Riemannian measure if the manifold $\mathcal{G}_z$ is compact, and hence it also holds for any equivalent distributions.

Proposition 5.4. *Suppose that Π_z satisfies κ-PI for $|z| \le \delta$, and q is supported on $\{x : |g(x)| \le \delta\}$. Suppose also that Π_z, q^g, π^g admit C^1 density functions. Then for any function f such that $|f| \le 1$, the following holds*

$$|\mathbb{E}_q[f] - \mathbb{E}_{\Pi_0}[f]| \le \sqrt{8\kappa \mathbb{E}_q[\|D(s_q - s_\pi)\|^2]} + \max_{|z| \le \delta} |\mathbb{E}_{\Pi_z}[f] - \mathbb{E}_{\Pi_0}[f]|.$$

In particular, Proposition 5.4 shows that if we have a q such that it is 1) supported close to $\mathcal{G}_0$ and 2) $F_\perp(q, \pi)$ is small, then q is close to Π_0 in total variation. We can also interpret the bound alternatively as a decomposition of the difference between q and Π_0: the first part measures the difference along with $\mathcal{H}_\perp$ directions and it is controlled by $F_\perp$; the second part measures the difference along ∇g direction, and it is controlled by the distance between q's support to $\mathcal{G}_0$.

As a final remark, the orthogonal-space Stein equation (15) and orthogonal-space Fisher divergence (16) can be applied to densities supported on $\mathbb{R}^d$, and their formulations do not require information of $\mathcal{G}_0$ such as parameterization and geodesic. So they can be used as very computationally friendly statistical divergence in practice.

Proposition 5.5. *Suppose we apply $v_t = v_\sharp + Du$ with (6) and (8) to the density field q_t.*

1. *Let $M_t = \max\{|g(x)|, x \in supp(q_t)\}$. Let S_t be the solution of an ordinary differential equation $\dot{S}_t = -\psi(S_t)$. Suppose $S_0 = M_0 < \infty$, then $M_t < S_t$ a.s..*

2. *Assume ψ is differentiable with derivative $\dot{\psi}$. Suppose g is also smooth enough so that*

$$\left| \nabla g^T s_\pi + \Delta g - \frac{2\nabla g^T \nabla^2 g \nabla g}{\|\nabla g\|^2} \right| \leq C_0 \|\nabla g\|^2. \tag{17}$$

Then the KL-divergence follows $\frac{d}{dt}\mathrm{KL}(q_t\|\pi) \leq -F_\perp(q_t, \pi) + \mathbb{E}_{q_t}[|\dot{\psi}(g)|] + C_0\mathbb{E}_{q_t}[|\psi(g)|\|\nabla g\|^2]$.

The first result shows that the support of q_t will shrink towards $\mathcal{G}_0$. But in order to avoid possible stagnation case, e.g. $\|g\|$ is too small, it is necessary to impose (17). The second result provides us some hint on the choice of ψ. Note that under q_t for large enough t, $g(x)$ will take values close to 0, so we need $\psi(0) = \dot{\psi}(0) = 0$. Since $\psi(x)$ should have the same sign with x, one natural choice would be $\psi(y) = \alpha\mathrm{sign}(y)|y|^{1+\beta}$.

Theorem 5.6. *Suppose we choose $\psi(y) = \alpha sign(y)|y|^{1+\beta}$ for an $\alpha > 0, \beta \in (0,1]$ and the conditions of Proposition 5.5 hold. Suppose also that $\mathrm{KL}(q_0, \pi) < \infty$ and $\|\nabla g(x)\| \leq D_0$ if $|\psi(x)| < M_0$, then $M_T = O(T^{-\frac{1}{\beta}})$, $\min_{t \leq T} F_\perp(q_t, \pi) = O(\log T/T)$. The orthogonal-space Stein equality holds approximately, in the sense that for any $\phi(x) \perp \nabla g(x)$ and $\|\phi\|_{\mathcal{H}} = 1$, $\min_{t \leq T}|\mathbb{E}_{q_t}\mathcal{A}_\pi\phi| \leq \sqrt{F_\perp(q_t, \pi)} = O(\sqrt{\log T/T})$.*

Theorem 5.6 shows that we can obtain a q_t that is supported on $\mathcal{G}_{[-\delta,\delta]} = \{x : |g(x)| \leq \delta\}$, while it has close to zero orthogonal-space Fisher divergence and the Stein identity holds approximately on $\mathcal{H}_\perp$. Combining it with Proposition 5.4, we have

Corollary 5.7. *Under the conditions of Theorem 5.6, for a bounded f with $\mathcal{H} = L_q^2$, suppose $h_f(z) = \mathbb{E}_\pi[f|g(X) = z] = \mathbb{E}_{\Pi_z}[f]$ is Lipschitz with near $z = 0$, then $\min_{t \leq T}|\mathbb{E}_{q_t}[f] - \mathbb{E}_{\Pi_0}[f]| \leq O(\sqrt{\log T/T})$.*

6 Experiments

We demonstrate the effectiveness of our methods on various tasks, including a constrained synthetic distribution, an income classification with fairness constraint, a loan application with logic constraint and Bayesian deep neural networks with robustness constraint. For all MCMC methods, we run n parallel chains and collect the final samples. For all SVGD, we use n particles with an RBF kernel.

Synthetic Distribution   We first demonstrate our methods on a two-dimensional synthetic distribution where the ground truth samples are available. Let $y \sim \mathcal{N}(0, I)$, and we transform y to $x = [x_1, x_2]$ by $x = \phi^{-1}(y)$ where $\phi(x) = [x_1 + x_2^3, x_2]^\top$. We aim to sample from $\pi(x)$ with constraint $g(x) = x_1 + x_2^3 = 0$. We are able to generate ground truth samples by first generating $y_0 \sim \mathcal{N}(0, 1)$ and then setting $x_{gt} = [-y_0^3, y_0]$. We report the energy distance between the collected samples and the ground truth samples (measuring how well samples approximate the target distribution) and the mean absolute error (MAE) $|g(x)|$ (measuring how well the samples satisfy the constraint). We set $n = 50$ for this task.

In order to compare with previous manifold sampling methods, which all require initializations on the manifold, we first consider the case when the sampler starts on the manifold. We compare with constrained Langevin dynamics (CLangevin) and constrained Hamiltonian Monte Carlo (CHMC), which are two advanced manifold samplers [18]. From Figure 2a, we observe that O-SVGD and O-Langevin converge after 2000 and 4000 steps respectively whereas CLangevin and CHMC have not fully converged even after 5000 steps. These results indicate that projection-based methods could

(a)   (b)   (c)   (d)

Figure 2: Energy distance and MAE w.r.t. iterations when starting: (a)&(b) on the manifold and (c)&(d) outside the manifold.

(a) Training Curve   (b) Testing Curve

Figure 3: Income Classification: O-Langevin and O-SVGD have almost zero fairness loss and disparate impact while maintaining great prediction performance. Standard Langevin and SVGD fail to make fair predictions.

(a) Training Curve   (b) Testing Curve

Figure 4: Loan Classification: O-Langevin and O-SVGD satisfy the constraint well whereas standard Langevin and SVGD violate the constraint severely.

be inefficient due to finding sub-optimal solutions. In terms of runtime, we observe that O-Langevin converges the fastest (see Appendix for comparisons). Figure 2b shows that previous methods have almost zero MAE all the time since they are required to stay on the manifold. Our methods have reasonably small MAE (but not zero due to operating in the ambient space) even without expensive projection steps. O-Langevin has more wiggly trajectories than O-SVGD due to Gaussian noise.

Next, we test our methods starting outside the manifold, where previous manifold sampling methods are not applicable. From Figure 2d&e, we observe again that O-SVGD converges the fastest in terms of iterations and both methods converge very quickly to the manifold and are able to stay close to it.

Income Classification with Fairness Constraint   ML Fairness has gained increasing attention in recent years, which aims to guarantee unbiased treatments for individuals w.r.t. gender, race, disabilities, etc. Following previous work [25, 23], we predict whether an individual's annual income is $\geq 50,000$ unfavorably in terms of the gender. We use *Adult Income* dataset [12] and train a Bayesian neural network with two-layer multilayer perceptron (MLP). During testing, we use CV score [4] as the measure of disparate impact. Training and testing performance of our methods and standard algorithms are shown in Figure 3. We find that during training and testing, both O-Langevin and O-SVGD satisfy the constraint very well while maintaining high training LL and test accuracy. In contrast, standard Langevin and SVGD violate the constraint significantly, indicating that they fail to make fair predictions. These results demonstrate the power of O-Gradient which keeps the sampler on the manifold while still correctly sampling from $\pi(x)$.

Table 1: Results on Image Classification (%).

	Test Error ($\downarrow$)	ECE ($\downarrow$)	AUROC ($\uparrow$)
SGLD	15.00	2.21	89.41
Tempered SGLD	4.73	0.83	97.63
O-Langevin	**4.46**	0.87	**98.68**
SVGD	6.11	0.93	93.55
O-SVGD	4.92	**0.77**	94.69

Loan Classification with Logic Rules Following previous work [24], we predict whether to lend loans to applicants and encode two logic rules: (1) an applicant must be rejected if not employed and having the lowest credit rank; (2) an applicant must be approved if employed over 15 years and having the highest credit rank. We use a Bayesian logistic regression model and set $g(x) = l_{\text{logic}}(x) = \mathbb{E}_{(\mu,\nu)\sim\mathcal{D}_{\text{logic}}}[\text{Loss}(\nu, \hat{\nu}(\mu; x))]$ where $\mathcal{D}_{\text{logic}}$ is a uniform distribution over all datapoints (μ, ν) that satisfy the constraint, Loss refers to the classification loss and $\hat{\nu}$ is the prediction made by the model. In Figure 4, we see that both of our methods satisfy the logic rules well whereas standard algorithms violate the constraint severely. A drop in the accuracy of our methods might be caused by conflicts between the data and the logic rules, as indicated in Liu et al. [24]. Nevertheless, our methods can obtain models satisfying the constraint, thus guarantee their safety and interpretability.

Prior-Agnostic Bayesian Neural Networks Bayesian neural network (BNN) has been widely used in deep learning (DL) for quantifying uncertainty. The posterior of BNNs is determined by the prior which reflects our prior knowledge, and the likelihood which quantifies the data fitness. Due to complex NN architectures, specifying an appropriate prior for BNNs has been shown to be difficult [8]. A poor prior often leads to a bad posterior and thus bad inference results. To automatically control the influence of the prior, one approach is to sample from the posterior with the constraint of a reasonably high data fitness. We apply our methods to image classification on CIFAR10 with ResNet-18. Since the training loss is typically very small in DL, we set the constraint to be $g(x) = \text{Loss}(x)$. Similar to stochastic Langevin dynamics (SGLD) [33] and SVGD, our methods are easy to be combined with stochastic gradients which are used on this task. We also ignore the second-order derivative terms in our methods for speedup. Besides standard algorithms, we also compare with a tempered SGLD which has been used in the literature for good performance [35]. We report test error, expected calibration error (ECE) and AUROC on the SVHN dataset (measuring out-of-distribution detection).

From Table 1, we see that O-Langevin and O-SVGD improve significantly over their unconstrained counterparts on all three metrics. O-Langevin has the lowest test error and the highest AUROC while O-SVGD has the lowest ECE. These results indicate that the constraint can automatically limit the effect of the prior, leading to a better posterior and thus better generalization and calibration.

7 Conclusion and Limitations

We propose a new variational framework with a designed gradient flow (O-Gradient) for sampling in general, implicitly defined constrained domains. O-Gradient is formed by two orthogonal directions where one drives the sampler towards the domain and the other explores the domain by decreasing a KL divergence. We prove the convergence of O-Gradient and apply the framework to both Langevin dynamics and SVGD. We empirically demonstrate the power of our methods on a variety of ML tasks. While ML achieves impressive performance, we must take realistic constraints into consideration when deploying it in daily life. This work provides a principled framework for solving constrained sampling problem with theoretical guarantees and practical algorithms.

One possible limitation of our methods is that the obtained samples are not exactly on the manifold due to working in an ambient space, though can be made arbitrarily close to the manifold by setting hyperapameters. This might be solved by including a projection step. Another limitation is that we only consider an equality constraint. It will be interesting to incorporating inequality constraints into the framework. Besides, our theory only considers the convergence in the continuous-time limit. Future work could provide convergence analysis in discrete time, which may characterize the properties of practical algorithms more accurately.

References

[1] Steve Brooks, Andrew Gelman, Galin Jones, and Xiao-Li Meng. *Handbook of markov chain monte carlo*. CRC press, 2011.

[2] Marcus Brubaker, Mathieu Salzmann, and Raquel Urtasun. A family of mcmc methods on implicitly defined manifolds. In *Artificial intelligence and statistics*, pages 161–172. PMLR, 2012.

[3] Simon Byrne and Mark Girolami. Geodesic monte carlo on embedded manifolds. *Scandinavian Journal of Statistics*, 40(4):825–845, 2013.

[4] Toon Calders and Sicco Verwer. Three naive bayes approaches for discrimination-free classification. *Data Mining and Knowledge Discovery*, 21(2):277–292, 2010.

[5] Joseph T Chang and David Pollard. Conditioning as disintegration. *Statistica Neerlandica*, 51 (3):287–317, 1997.

[6] Persi Diaconis, Susan Holmes, and Mehrdad Shahshahani. Sampling from a manifold. In *Advances in modern statistical theory and applications: a Festschrift in honor of Morris L. Eaton*, pages 102–125. Institute of Mathematical Statistics, 2013.

[7] Andrew Duncan, Nikolas Nüsken, and Lukasz Szpruch. On the geometry of stein variational gradient descent. *arXiv preprint arXiv:1912.00894*, 2019.

[8] Vincent Fortuin, Adrià Garriga-Alonso, Florian Wenzel, Gunnar Rätsch, Richard Turner, Mark van der Wilk, and Laurence Aitchison. Bayesian neural network priors revisited. *arXiv preprint arXiv:2102.06571*, 2021.

[9] Mark Girolami and Ben Calderhead. Riemann manifold langevin and hamiltonian monte carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(2): 123–214, 2011.

[10] Jackson Gorham and Lester Mackey. Measuring sample quality with kernels. In *International Conference on Machine Learning*, pages 1292–1301. PMLR, 2017.

[11] Emmanuel Hebey. *Nonlinear Analysis on Manifolds: Sobolev Spaces and Inequalities: Sobolev Spaces and Inequalities*, volume 5. American Mathematical Soc., 2000.

[12] Ron Kohavi. Scaling up the accuracy of naive-bayes classifiers: A decision-tree hybrid. In *Kdd*, volume 96, pages 202–207, 1996.

[13] Yunbum Kook, Yin Tat Lee, Ruoqi Shen, and Santosh S Vempala. Sampling with riemannian hamiltonian monte carlo in a constrained space. *arXiv preprint arXiv:2202.01908*, 2022.

[14] Yann LeCun, Sumit Chopra, Raia Hadsell, M Ranzato, and F Huang. A tutorial on energy-based learning. *Predicting structured data*, 1, 2006.

[15] Yin Tat Lee and Santosh S Vempala. Convergence rate of riemannian hamiltonian monte carlo and faster polytope volume computation. In *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing*, pages 1115–1121, 2018.

[16] Tony Lelievre and Gabriel Stoltz. Partial differential equations and stochastic methods in molecular dynamics. *Acta Numerica*, 25:681–880, 2016.

[17] Tony Lelievre, Mathias Rousset, and Gabriel Stoltz. Langevin dynamics with constraints and computation of free energy differences. *Mathematics of computation*, 81(280):2071–2125, 2012.

[18] Tony Lelièvre, Mathias Rousset, and Gabriel Stoltz. Hybrid monte carlo methods for sampling probability measures on submanifolds. *Numerische Mathematik*, 143(2):379–421, 2019.

[19] Tony Lelièvre, Gabriel Stoltz, and Wei Zhang. Multiple projection markov chain monte carlo algorithms on submanifolds. *IMA Journal of Numerical Analysis*, 2022.

[20] Peter Li and Jiaping Wang. Weighted poincaré inequality and rigidity of complete manifolds. In *Annales scientifiques de l'Ecole normale supérieure*, volume 39, pages 921–982, 2006.

[21] Qiang Liu. Stein variational gradient descent as gradient flow. *Advances in Neural Information Processing Systems (NeurIPS)*, 2017.

[22] Qiang Liu and Dilin Wang. Stein variational gradient descent: A general purpose Bayesian inference algorithm. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2016.

[23] Suyun Liu and Luis Nunes Vicente. Accuracy and fairness trade-offs in machine learning: A stochastic multi-objective approach. *arXiv preprint arXiv:2008.01132*, 2020.

[24] Xingchao Liu, Xin Tong, and Qiang Liu. Sampling with trusthworthy constraints: A variational gradient framework. *Advances in Neural Information Processing Systems*, 34, 2021.

[25] Natalia Martinez, Martin Bertran, and Guillermo Sapiro. Minimax pareto fairness: A multi objective perspective. In *International Conference on Machine Learning*, pages 6755–6764. PMLR, 2020.

[26] Gilles Meyer, Silvere Bonnabel, and Rodolphe Sepulchre. Linear regression under fixed-rank constraints: a riemannian approach. In *Proceedings of the 28th international conference on machine learning*, 2011.

[27] Sam Patterson and Yee Whye Teh. Stochastic gradient riemannian langevin dynamics on the probability simplex. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2013.

[28] Upanshu Sharma and Wei Zhang. Nonreversible sampling schemes on submanifolds. *SIAM Journal on Numerical Analysis*, 59(6):2989–3031, 2021.

[29] Leon Simon. Introduction to geometric measure theory. *Tsinghua Lectures*, 2014.

[30] Yang Song and Stefano Ermon. Generative modeling by estimating gradients of the data distribution. In *Advances in Neural Information Processing Systems*, pages 11895–11907, 2019.

[31] Gabriel Stoltz, Mathias Rousset, et al. *Free energy computations: A mathematical perspective*. World Scientific, 2010.

[32] Xiao Wang, Qi Lei, and Ioannis Panageas. Fast convergence of langevin dynamics on manifold: Geodesics meet log-sobolev. *Advances in Neural Information Processing Systems*, 33:18894–18904, 2020.

[33] Max Welling and Yee W Teh. Bayesian learning via stochastic gradient Langevin dynamics. In *International Conference on Machine Learning*, 2011.

[34] Emilio Zappa, Miranda Holmes-Cerfon, and Jonathan Goodman. Monte carlo on manifolds: sampling densities and integrating functions. *Communications on Pure and Applied Mathematics*, 71(12):2609–2647, 2018.

[35] Ruqi Zhang, Chunyuan Li, Jianyi Zhang, Changyou Chen, and Andrew Gordon Wilson. Cyclical stochastic gradient mcmc for bayesian deep learning. *International Conference on Learning Representations*, 2020.

[36] Wei Zhang. Ergodic sdes on submanifolds and related numerical sampling schemes. *ESAIM: Mathematical Modelling and Numerical Analysis*, 54(2):391–430, 2020.

A proofs

Proof of Lemma 4.1. When $\mathcal{H}$ is the RKHS with kernel k

$$\mathbb{E}_{q_t}[(D(s_\pi - s_{q_t}))^\top u] - \frac{1}{2}\|Du\|_{\mathcal{H}}^2$$

$$= \mathbb{E}_{q_t}[(D(s_\pi - s_{q_t}))^\top Du] - \frac{1}{2}\|Du\|_{\mathcal{H}}^2$$

$$= \langle R_{D(s_\pi - s_{q_t})}, Du \rangle_{\mathcal{H}} - \frac{1}{2}\|Du\|_{\mathcal{H}}^2$$

where R_v is the Riesz representation of the linear function $u \to \mathbb{E}_{q_t} v^\top u$. Therefore a solution to (8) is

$$Du = DR_{D(s_\pi - s_{q_t})}$$

which can be written as

$$Du(x) = \int D(x)k(x,y)D(y)(s_\pi(y) - s_q(y))q_t(y)dy$$

$$= \int k_\perp(x,y)(s_\pi(y) - s_q(y))q_t(y)dy$$

$$= \int (k_\perp(x,y)s_\pi(y) + \nabla_y k_\perp(x,y))q_t(y)dy$$

$$= \mathbb{E}_{y \sim q_t}(k_\perp(x,y)s_\pi(y) + \nabla_y k_\perp(x,y))$$

$\square$

Proof of Theorem 4.2. First, we show the components of r can be written as

$$r_i(x) = \sum_j \partial_j D_{i,j}(x) = -\frac{\sum_j \partial_{i,j}^2 g \partial_j g + \partial_i g \partial_{j,j} g}{\|\nabla g\|^2} + \frac{2(\nabla g^T \nabla^2 g \nabla g)\partial_i g}{\|\nabla g\|^4}.$$

To this end, note that

$$D_{i,j}(x) = 1 - \frac{\partial_i g \partial_j g}{\|\nabla g\|^2}$$

Then by product rule, and the fact that $\partial_j \|\nabla g\|^2 = 2\sum_k \partial_{k,j}^2 g \partial_k g = 2[\nabla^2 g \nabla g]_j$

$$\partial_j D_{i,j}(x) = \frac{-\partial_{i,j}^2 g \partial_j g - \partial_i g \partial_{jj}^2 g}{\|\nabla g\|^2} + \frac{2\partial_i g \partial_j g[\nabla^2 g \nabla g]_j}{\|\nabla g\|^4}.$$

So

$$\sum_j \partial_j D_{i,j}(x) = \frac{-\sum_j \partial_{i,j}^2 g \partial_j g - \sum_j \partial_i g \partial_{jj}^2 g}{\|\nabla g\|^2} + \frac{2\partial_i g (\nabla g^\top \nabla^2 g \nabla g)}{\|\nabla g\|^4}.$$

Comparing with the each component of

$$r = -\frac{\nabla^2 g \nabla g}{\|\nabla g\|^2} - \frac{\text{tr}(\nabla^2 g)}{\|\nabla g\|^2}\nabla g + \frac{2\nabla g^T \nabla^2 g \nabla g}{\|\nabla g\|^4}\nabla g,$$

it is easy to see they are the same.

Next, recall the O-gradient density flow (10)

$$\frac{d}{dt}q = -\nabla \cdot (\phi(x)q(x)) + \nabla \cdot (D(x)\nabla q(x))$$

$$= -\nabla \cdot (\phi(x)q(x)) + \sum_{i,j} \partial_i(D_{i,j}(x)\partial_j q(x))$$

$$= -\nabla \cdot (\phi(x)q(x)) + \sum_{i,j} \partial_i D_{i,j}(x)\partial_j q(x) + \sum_{i,j} D_{i,j}(x)\partial_{i,j}^2 q(x)$$

$$=: -\nabla \cdot (\phi(x)q(x)) + (I).$$

Meanwhile, the FPE of the SDE follows

$$\frac{d}{dt}q = -\nabla \cdot (\phi(x)q(x)) - \nabla \cdot (r(x)q(x)) + \sum_{i,j} \partial_{i,j}^2 (D_{i,j}(x)q(x)) \qquad (18)$$

Then note that

$$(II) := -\nabla \cdot (r(x)q(x)) = -\sum_i \partial_i \left(\sum_j \partial_j D_{i,j}(x)q(x) \right)$$

$$= -\sum_{i,j} \partial_{i,j}^2 D_{i,j}(x)q(x) - \sum_{i,j} \partial_j D_{i,j}(x)\partial_i q(x).$$

Also note that

$$\sum_{i,j} \partial_{i,j}^2 (D_{i,j}(x)q(x)) = \sum_{i,j} \partial_{i,j}^2 D_{i,j}(x)q(x) + \sum_{i,j} \partial_j D_{i,j}(x)\partial_i q(x)$$

$$+ \sum_{i,j} \partial_i D_{i,j}(x)\partial_j q(x) + \sum_{i,j} D_{i,j}(x)\partial_{i,j}^2 q(x)$$

$$= (I) - (II).$$

So we arrive at our first claim.

For the second claim, using $\nabla g(x)^T D(x) = 0$, the Ito formula indicates that

$$dg(X_t) = \nabla g(X_t)^T [D(X_t)\nabla \log \pi(X_t) - \frac{cg(X_t)}{\|\nabla g(X_t)\|^2}\nabla g(X_t) + r(X_t) + D(X_t)\sqrt{2}dW_t]$$

$$+ \operatorname{tr}(D(X_t)\nabla^2 g(X_t)D(X_t))dt$$

$$= -cg(X_t)dt + \nabla g(X_t)^T r(X_t) + \operatorname{tr}(D(X_t)\nabla^2 g(X_t)D(X_t))dt.$$

So it suffices to show that

$$\nabla g(x)^T r(x) + \operatorname{tr}(D(x)\nabla^2 g(x)D(x)) = 0.$$

To continue, we suppress the expression of x in below to keep formulas short. We first note that $D_{i,j} = 1_{i=j} - \partial_i g\partial_j g/\|\nabla g\|^2$,

$$\partial_j D_{i,j} = \frac{-\partial_{i,j}g\partial_j g - \partial_i g\partial_{j,j}g}{\|\nabla g\|^2} + \frac{2\sum_k \partial_i g\partial_j g\partial_{j,k}^2 g\partial_k g}{\|\nabla g\|^4}$$

We plug this into the computation of $\nabla g^T r = \sum_i \partial_i g\partial_j D_{i,j}$. We note that

$$\sum_{i,j} -\frac{\partial_i g\partial_{i,j}g\partial_j g}{\|\nabla g\|^2} = \frac{-(\nabla g)^T \nabla^2 g\nabla g}{\|\nabla g\|^2}.$$

$$\sum_{i,j} -\frac{(\partial_i g)^2 \partial_{j,j}g\partial_j g}{\|\nabla g\|^2} = -\operatorname{tr}(\nabla^2 g).$$

$$\frac{2\sum_{i,j}\sum_k \partial_i g\partial_i g\partial_j g\partial_{j,k}^2 g\partial_k g}{\|\nabla g\|^4} = \frac{\sum_{j,k}\|\nabla g\|^2 \partial_j g\partial_{j,k}^2 g\partial_k g}{\|\nabla g\|^4} = \frac{2(\nabla g)^T \nabla^2 g\nabla g}{\|\nabla g\|^2}$$

Therefore

$$\sum_i \partial_i g\partial_j D_{i,j} = \frac{(\nabla g)^T \nabla^2 g\nabla g}{\|\nabla g\|^2} - \operatorname{tr}(\nabla^2 g)$$

$$= \operatorname{tr}\left(-\nabla^2 g + \nabla^2 g\frac{\nabla g(\nabla g)^T}{\|\nabla g\|^2} \right)$$

$$= \operatorname{tr}\left(-\nabla^2 gD \right) = \operatorname{tr}\left(-D\nabla^2 gD \right), \quad \text{since } D^2 = D.$$

This completes our proof.

For the last claim, we note that

$$\mathcal{L}f = \nabla f^T[D\nabla \log \pi - \frac{\psi(g)}{\|\nabla g\|^2}\nabla g + r] + \text{tr}(D\nabla^2 f D)$$

$$= \nabla f^T[\nabla \log \pi - \frac{\nabla^2 g}{\|\nabla g\|^2}\nabla g] + \text{tr}(D\nabla^2 f)$$

$$= \nabla f^T \nabla \log \pi + \text{tr}(\nabla^2 f) - \frac{\nabla f^T \nabla^2 g \nabla g}{\|\nabla g\|^2} + \frac{\nabla g^T \nabla^2 f \nabla g}{\|\nabla g\|^2}$$

Finally, we note that $\nabla f^T \nabla g = 0$. Take derivative of this identity we find

$$\nabla^2 f \nabla g + \nabla^2 g \nabla f = 0$$

Therefore

$$\nabla f^T \nabla^2 g \nabla g = -\nabla g^T \nabla^2 f \nabla g$$

and

$$\mathcal{L}f = \nabla f^T \nabla \log \pi + \text{tr}(\nabla^2 f)$$

which is the same as the generator of the Langevin diffusion. $\qquad\square$

Remark If we drop r and implementing a naive SDE:

$$dx_t = \phi(x) + \sqrt{2}D(x)dw_t,$$

the FPE of this SDE will be

$$\frac{d}{dt}q = -\nabla \cdot (\phi(x)q(x)) + \sum_{i,j} \partial_{i,j}^2 (D_{i,j}(x)q(x)).$$

It is identical to (18) but without the term $-\nabla \cdot (r(x)q(x)) = (II)$. In other words, it will not match (10) unless $(II) = 0$, which happens when $\sum_{i,j} \partial_{i,j}^2 D_{i,j} = \sum_j \partial_j D_{i,j} \equiv 0$. But it should be pointed out that if g is an affine function, $D(x)$ is a constant matrix, then $(II) \equiv 0$, and r is safe to be dropped out.

Proof of Proposition 5.2. We will first show that for any function f, the following holds

$$\int \pi^g(z)f(x)\pi_{\eta,z}(x)dxdz \to \int f(x)\pi(x)dx,$$

when $\eta \to 0$. So (13) holds for Π_z as the weak limit of $\pi_{\eta,z}$.

We note that

$$\pi_{\eta,z}(x) = \frac{\pi(x)\exp(-\frac{1}{2\eta}(g(x) - z)^2)}{\sqrt{2\eta\pi}Z_{\eta,z}}$$

where the normalizing constant is given by

$$Z_{\eta,z} = \frac{1}{\sqrt{2\pi\eta}}\int \pi(x)\exp(-\frac{1}{2\eta}(g(x) - z)^2)dx.$$

Because π^g is the density of $g(X)$, so for any function h:

$$\int h(g(x))\pi(x)dx = \mathbb{E}_{X\sim\pi}[h(g(X))] = \int h(y)\pi^g(y)dy.$$

We pick $h(y) = \exp(-\frac{1}{2\eta}(y - z)^2)$, we obtain that

$$Z_{\eta,z} = \int \frac{1}{\sqrt{2\pi\eta}}\exp(-\frac{1}{2\eta}(y - z)^2)\pi^g(y)dy$$

$$= \mathbb{E}\pi^g(z + \sqrt{\eta}\xi) = \pi^g(z)(1 + R(z))^{-1}.$$

494   where the $|R| \leq 2L\sqrt{\eta}$ and L is the regularity constant of π^g. Therefore

$$\int \pi^g(z)f(x)\pi_{\eta,z}(x)dxdz$$

$$= \int \pi(x)f(x)\left(\int \frac{1}{\sqrt{2\pi\eta}}\exp(-\frac{1}{2\eta}(g(x)-z)^2)\frac{\pi^g(z)}{Z_{\eta,z}}dz\right)dx$$

$$= \int \pi(x)f(x)\left(\int \frac{1}{\sqrt{2\pi\eta}}\exp(-\frac{1}{2\eta}(g(x)-z)^2)dz\right)dx + \mathbb{E}R(g(X))$$

$$= \int \pi(x)f(x)dx + \mathbb{E}R(g(X)).$$

495   Since $\mathbb{E}R(g(X)) \leq 2L\sqrt{\eta}$, we find our first claim when $\eta \to 0$.

496   Next note that if we pick $f(x) = 1_{|g(x)-z|\geq\epsilon}$

$$\int f(x)\pi_{\eta,z}(x)dx$$

$$= \int 1_{|g(x)-z|\geq\epsilon}\frac{1}{\sqrt{2\pi\eta}}\exp(-\frac{1}{2\eta}(g(x)-z)^2)\frac{\pi(x)}{Z_{\eta,z}}dx$$

$$\leq \int \frac{1}{\sqrt{2\pi\eta}}\exp(-\frac{\epsilon^2}{2\eta})\frac{\pi(x)}{Z_{\eta,z}}dx = \frac{1}{\sqrt{2\pi\eta}Z_{\eta,z}}\exp(-\frac{\epsilon^2}{2\eta}).$$

497   When $\eta \to 0$, since $Z_{\eta,z} \to \pi^g(z)$, and $\frac{1}{\sqrt{2\eta}}\exp(-\frac{\epsilon^2}{2\eta}) \to 0$, we find that

$$\Pi_z(|g(X)-z| \geq \epsilon) = \lim_{\eta\to0}\int f(x)\pi_{\eta,z}(x)dx \to 0$$

For the Stein equation part, note that for each $\pi_{\eta,z}$, we have the following by Stein's identity:

$$\mathbb{E}_{\pi_{\eta,z}}[(\nabla\log\pi(x) - \frac{1}{2\eta}(g(x)-z)\nabla g(x))^\top\phi(x) + \nabla^\top\phi(x)] = 0.$$

But $\nabla g(x)^\top\phi(x) = 0$ for $\phi \in \mathcal{H}_\perp$. This gives

$$\mathbb{E}_{\pi_{\eta,z}}[\mathcal{A}_\pi\phi] = 0, \quad \forall\eta.$$

498   Taking $\eta \to 0$ yields that $\mathbb{E}_{\pi_z}[\mathcal{A}_\pi\phi] = 0$. $\square$

499

500   **Lemma A.1.** *Suppose $q(x), q^g(z)$ and $\pi(x), \pi^g(z), \pi_z(x)$ are all C^1 functions, then*

$$q_z(x) = \frac{\pi^g(z)q(x)}{q^g(z)\pi(x)}\pi_z(x), \quad z = g(x).$$

501   *In particular,*

$$D(s_{q_z}(x) - s_{\pi_z}(x)) = D(s_q(x) - s_\pi(x)).$$

502   *Proof of Lemma A.1.* We will show (13) holds with this choice of $q_z(x)$. Suppose $x \in \mathcal{G}_z$, let
503   y be the its coordinate on the manifold $\mathcal{G}_z$. Let t be the map such that $t(z,y) = x$. We write
504    $f_z(y) = f(t(z,y))$ Then we have

$$\mathbb{E}_{z\sim q^g}\mathbb{E}_{x\sim q_z}[f(x)] = \int_R dzq^g(z)\int_{\mathcal{G}_z}q_z(y)f_z(y)dy$$

$$= \int_R dzq^g(z)\int_{\mathcal{G}_z}\frac{\pi^g(z)q(t(y,z))}{q^g(z)\pi(t(y,z))}\pi_z(y)f_z(y)dy$$

$$= \int_R \pi^g(z)dz\int_{\mathcal{G}_z}\frac{q(t(y,z))}{\pi(t(y,z))}\pi_z(y)f_z(y)dy$$

$$= \mathbb{E}_{z\sim\pi^g}\mathbb{E}_{y\sim\pi_z}[f_z(y)q(t(z,y))/\pi(t(z,y))]$$

$$= \mathbb{E}_{x\sim\pi}[f(x)q(x)/\pi(x)] = \mathbb{E}_{x\sim q}[f(x)].$$

505 Moreover

$$D(s_{q_z}(x)-s_{\pi_z}(x)) = D(s_q(x)-s_\pi(x))+D(\nabla g(x)s_{q^g}(g(x))-\nabla g(x)s_{q^g}(g(x))) = D(s_q(x)-s_\pi(x)).$$

506 $\square$

507 *Proof of Proposition 5.4.* Note that

$$\left| \mathbb{E}_{\Pi_0}[f] - \int_{-\delta}^{\delta} q_g(z)\mathbb{E}_{\Pi_z}[f]dz \right| \leq \int_{-\delta}^{\delta} q_g(z)\left| \mathbb{E}_{\Pi_0}[f] - \mathbb{E}_{\Pi_z}[f] \right| dz$$
$$\leq \max_{|z|\leq\delta} |\mathbb{E}_{\Pi_z}[f] - \mathbb{E}_{\Pi_0}[f]|.$$

508 Meanwhile

$$\mathbb{E}_q[f] - \int_{-\delta}^{\delta} q_g(z)\mathbb{E}_{\Pi_z}[f]dz = \int_{-\delta}^{\delta} q_g(z)\left(\mathbb{E}_{q_z}[f] - \mathbb{E}_{\Pi_z}[f] \right) dz.$$

509 Then we note the following holds when we restrict f on $\mathcal{G}_z$. We use a paramaterization of $\mathcal{G}_z$ with
510 dummy variable y and the inherited metric form $\mathbb{R}^d$,

$$(\mathbb{E}_{q_z}[f] - \mathbb{E}_{\Pi_z}[f])^2 = \left(\int_{\mathcal{G}_z} \Pi_z(y)(\frac{q_z(y)}{\Pi_z(y)} - 1)f(y)dy \right)^2$$
$$\leq \int_{\mathcal{G}_z} \Pi_z(y)(\sqrt{\frac{q_z(y)}{\Pi_z(y)}} - 1)^2 dy \cdot \int_{\mathcal{G}_z} \Pi_z(y)(\sqrt{\frac{q_z(y)}{\Pi_z(y)}} + 1)^2 f(y)^2 dy$$
$$\leq 2\int_{\mathcal{G}_z} \Pi_z(y)(\sqrt{\frac{q_z(y)}{\Pi_z(y)}} - 1)^2 dy \cdot \int_{\mathcal{G}_z} \Pi_z(y)(\frac{q_z(y)}{\Pi_z(y)} + 1)dy$$
$$= 4\int_{\mathcal{G}_z} \Pi_z(y)(\sqrt{\frac{q_z(y)}{\Pi_z(y)}} - 1)^2 dy$$
$$= 8(1 - \mathbb{E}_{\Pi_z}\sqrt{\frac{q_z(y)}{\Pi_z(y)}}) = 8(1 - b) \text{ with } b = \mathbb{E}_{\Pi_z}\sqrt{\frac{q_z(y)}{\Pi_z(y)}} \leq 1.$$

511 Then note that

$$\mathrm{var}_{\Pi_z}\sqrt{\frac{q_z(y)}{\Pi_z(y)}} = \mathbb{E}_{\Pi_z}[\frac{q_z(y)}{\Pi_z(y)}] - b^2 = 1 - b^2.$$

512 Therefore by κ-PI, we have

$$1 - b \leq (1 - b^2) \leq \kappa \int_{\mathcal{G}_z} \Pi_z(y) \left\| \nabla\sqrt{\frac{q_z(y)}{\Pi_z(y)}} \right\|_{\mathcal{G}_z}^2 dy$$
$$= \kappa \int_{\mathcal{G}_z} q_z(y)\|(s_{q_z}(y) - s_{\Pi_z}(y))\|_{\mathcal{G}_z}^2 dy$$
$$= \kappa \int_{\mathcal{G}_z} q_z(y)\|D(y)(s_{q_z}(y) - s_{\Pi_z}(y))\|^2 dy$$
$$= \kappa\mathbb{E}_{q_z}\|D(s_{q_z} - s_{\Pi_z})\|^2.$$

513 So in combination

$$(\mathbb{E}_{q_z}[f] - \mathbb{E}_{\Pi_z}[f])^2 \leq 8\kappa\mathbb{E}_{q_z}\|D(s_{q_z} - s_{\Pi_z})\|^2.$$

514 Then by Lemma A.1 we have

$$D\nabla(\log q_z - \log \Pi_z) = D(s_q - s_\pi).$$

515 So we find that

$$(\mathbb{E}_{q_z}[f] - \mathbb{E}_{\Pi_z}[f])^2 \leq 4\kappa\mathbb{E}_{q_z}\|D(s_q - s_\pi)\|^2.$$

516 Integrating both sides with $q_g(z)$ we find our final claim.

517 $\square$

Proof of Proposition 5.5. Fix a particle z_t in the density of q_t, we track its g-value trajectory:

$$\frac{d}{dt}g(z_t) = \nabla g(z_t)^\top v_t(z_t) = -\psi(g(z_t)),$$

we will show that $g(z_t) \leq M_t$ for all t (The proof for $g(z_t) \geq -M_t$ is identifical is omitted). Suppose $g(z_t) > M_t + \epsilon$ for some t and $\epsilon > 0$. Let $t_0 = \inf\{t > 0, g(z_t) > M_t + \epsilon\}$. Then $\frac{d}{dt}g(z_{t_0}) = -\psi(M_{t_0} + \epsilon) < -\psi(M_{t_0})$, so for a sufficiently small $\delta > 0$, $g(z_{t_0-\delta}) > M_{t_0-\delta}$, this contradicts the definition of t_0.

For the second claim, note that

$$\frac{d}{dt}\mathrm{KL}(q_t\|\pi) = -\int q_t(x)v_t(x)^\top(s_\pi(x) - s_{q_t}(x))dx$$

$$= -\int q_t(x)(v_\perp(x) + v_\sharp(x))^\top(s_\pi(x) - s_{q_t}(x))dx$$

$$= -F_\perp(q_t, \pi) + \int q_t(x)\frac{\psi(g(x))\nabla g(x)^\top(s_\pi(x) - s_{q_t}(x))}{\|\nabla g(x)\|^2}dx$$

$$= -F_\perp(\pi, q_t) + \int \frac{\psi(g(x))\nabla g(x)^\top s_\pi(x)}{\|\nabla g(x)\|^2}dx$$

$$+ \int \frac{\|\nabla g(x)\|^2\dot\psi(g(x)) + \psi(g(x))\Delta g(x)}{\|\nabla g(x)\|^2}q_t(x)dx$$

$$- \int \frac{2\psi(g(x))\nabla g(x)^\top\nabla^2 g(x)\nabla g(x)}{\|\nabla g(x)\|^4}q_t(x)dx$$

$$\leq -F_\perp(\pi, q_t) + \mathbb{E}_{q_t}[|\dot\psi(g)|] + C_0\mathbb{E}_{q_t}[|\psi(g)|\|\nabla g\|^2].$$

$\square$

Proof of Theorem 5.6. We use the notation M_t from Proposition 5.5. When we take $\psi(z) = -\alpha\mathrm{sgn}(z)z^{1+\beta}$, we find that for some c_0 that depends on M_0

$$\frac{d}{dt}M_t = -\alpha|M_t|^{1+\beta} \Rightarrow M_t = (\alpha\beta)^{-\frac{1}{\beta}}(t + c_0)^{-\frac{1}{\beta}}.$$

Moreover, we have that for some constant C_1

$$\frac{d}{dt}\mathrm{KL}(q_t\|\pi)$$

$$\leq -F_\perp(q_t, \pi) + \alpha\mathbb{E}_{q_t}(1 + \beta)[|g|^\beta] + \alpha C_0\mathbb{E}_{q_t}[|g|^{1+\beta}\|\nabla g\|^2]$$

$$\leq -F_\perp(q_t, \pi) + \frac{C_1}{t + c_0} + \frac{C_1}{(t + c_0)^{1+1/\beta}}.$$

Integrating both sides yields the following for some constant M_α

$$\int_{T/2}^T F_\perp(q_t, \pi)dt \leq \int_0^T F_\perp(q_t, \pi)dt$$

$$\leq \mathrm{KL}(q_0, \pi) + C_1\log\frac{T + c_0}{c_0} + C_1 M_\alpha.$$

So

$$\min_{T/2 \leq t_0 \leq T} F_\perp(q_t, \pi) \leq \frac{2}{T}\mathrm{KL}(q_0, \pi) + \frac{2C_1}{T}\log\frac{T + c_0}{c_0} + \frac{2}{T}C_1 M_\alpha.$$

Finally, we note that if $r_{q,\pi}^D$ is the Riez representation of $D(s_\pi - s_q)^T$,

$$\mathbb{E}_q\mathcal{A}_\pi\phi = \mathbb{E}_q(D(s_\pi - s_q))^\top\phi = \langle r_{q,\pi}^D, \phi\rangle_\mathcal{H} \leq \frac{1}{2}\|r_{q,\pi}^D\|_\mathcal{H} = \frac{1}{2}\sqrt{F_\perp(q, \pi)}.$$

$\square$

B Additional Experiments Results and Setting Details

We use NVIDIA GeForce RTX 2080 Ti for neural network experiments.

B.1 Synthetic Distribution

For both O-Langevin and O-SVGD, we use $\alpha = 100$ and $\beta = 0$. We set $\eta = 0.01$ and 0.5 for O-Langevin and O-SVGD respectively. For CLangevin and CHMC, we use a python implementation [1] and tune the step size and the number of leapfrog steps. We report the best results which are achieved at step size $= 0.3$ for CLangevin, and step size $= 1$ and number of leapfrog steps $= 2$ for CHMC.

Runtime Comparison   We report runtime comparison in Figure 5. We did not include O-SVGD since one iteration of it already > 5s (SVGD is known to take more time per iteration than MCMC due to computing particle interaction). When starting on the manifold, we observe that O-Langevin converges much faster than previous manifold sampling methods. It takes about 4s to fully converge whereas previous methods have not fully converged after 5s. Previous methods cannot work with initializations outside the manifold, thus we only report the runtime of O-Langevin in Figure 5b.

Effect of Hyperparameters   Besides the hyperparameter step size η as in standard Langevin and SVGD, our methods have hyperparameters $\alpha > 0$ and $\beta \in (0, 1]$ in $\psi(x) = \alpha\text{sign}(x)|x|^{1+\beta}$ to control the speed of the sampler to approach the manifold and the closeness of the sampler to the manifold after converging. In theory, as α increases, the sampler approaches the manifold faster and stays closer. As β increases, the sampler first converges faster to the manifold but stays relatively far away after converging. We report the results of O-Langevin with varying α (with fixed $\beta = 0$) and β (with fixed $\alpha = 1$) when starting outside the manifold in Figure 6. We can see that the results of MAE, which measures the closeness to the manifold, align with the theoretical analysis. The energy distance with varying α and β are similar while $\alpha = 10$ and $\beta = 0$ perform slightly better. The theoretical analysis could not tell which values of hyperparameters give the fastest convergence to the target distribution thus we still need to tune α and β to achieve the optimal performance. In practice, we recommend to set $\beta = 0$ (though our theoretical results apply to $\beta \in (0, 1]$, we find $\beta = 0$ generally works well in practice.) and tune α to achieve a desirable MAE and energy distance.

Density Estimation   To compare the estimated density with the ground truth, we plot the collected samples after 5000 epochs when starting on the manifold and 8000 epochs when starting outside the manifold in Figure 7. The density estimation from our methods is closer to the ground truth than previous methods, aligning with the results of the energy distance in Figure 2.

(a)   (b)

Figure 5: Runtime comparison when (a) starting on the manifold and (b) starting outside the manifold.

---

[1] https://matt-graham.github.io/mici/

Figure 6: Effect of hyperparameters α and β

Figure 7: Density estimation when (a)-(d) starting on the manifold and (e)-(f) starting outside the manifold.

B.2 Income Classification with Fairness Constraint

The Adult Income dataset contains 30,162 training samples and 15,060 test samples. The feature dimension is 86. Following previous work [25, 23], we obtain the training set by randomly subsampling 20,000 data points from the training samples. The model is a two-layer multilayer perceptron

(MLP), which has 50 hidden units and RELU nonlinearities. The metric values are the mean over all particles. For both methods, we use $n = 10$ particles and $\beta = 0$. For O-Langevin, $\alpha = 100$ and $\eta = 10^{-5}$. For O-SVGD, $\alpha = 130$ and $\eta = 10^{-4}$. The results are averaged over 3 runs with the standard error as the error bar. CV score is a standard fairness measure of disparate impact Calders and Verwer [4]. It is defined as |p(income$\geq$ 50,000|female) - p(income$\geq$ 50,000|male)|

B.3 Loan Classification with Logic Rules

The dataset[2] contains loans issued through 2007-2015 of several banks. Each data point contains 28 features such as the current loan status and latest payment information. We define the logic loss to be the binary cross-entropy loss. The metric values are the mean over all particles. For both methods, we use $n = 10$ and $\beta = 0$. For O-Langevin, $\alpha = 80$ and $\eta = 10^{-4}$. For O-SVGD, $\alpha = 100$ and $\eta = 10^{-3}$. The results are averaged over 3 runs with the standard error as the error bar. The test violation refers to the percentage of predictions that do not follow the logic rules.

B.4 Prior-Agnostic Bayesian Neural Networks

For large models, such as ResNet-18 on this task, computing second-order derivatives is slow. To speed up our methods on large models, we ignore the second-order terms in O-Langevin and O-SVGD and empirically find that they still perform well. We leave the theoretical analysis for future work.

Specifically, we ignore the r term in the update of O-Langevin and obtain

$$x_{t+1} = x_t + \eta \cdot v_\sharp(x_t) + \text{Langevin}_\perp(x_t),$$
$$\text{where } \text{Langevin}_\perp(x_t) = \eta D(x_t)\nabla \log \pi(x_t) + \sqrt{2\eta}D(x_t)\xi_t, \ \ \xi_t \sim \mathcal{N}(0, I).$$

For O-SVGD, the update becomes

$$x_{i,t+1} = x_{i,t} + \eta \cdot \left(v_\sharp(x_{i,t}) + \text{SVGD}_{K_\perp}(x_{i,t}) \right),$$
$$\text{where } \text{SVGD}_{K_\perp}(x_{i,t}) = \frac{1}{n}\sum_{j=1}^{n} k_\perp(x_{i,t}, x_{j,t})\nabla_{x_{j,t}} \log \pi(x_{j,t}) + \tilde{\nabla}_{x_{j,t}} k_\perp(x_{i,t}, x_{j,t})$$
$$\text{and } \tilde{\nabla}_{x_{j,t}} k_\perp(x_{i,t}, x_{j,t}) = D(x_{i,t})(D(x_{j,t})\nabla_{x_{j,t}} k(x_{i,t}, x_{j,t}))$$

For all results, we use 200 epochs, 64 batchsize, $n = 4$, $\beta = 0$, $\alpha = 1000$ and $\eta = 10^{-4}$. During testing, we do Bayesian model averaging to obtain test error, ECE and AUROC. The test error refers to the percentage of the wrong predictions on the test set.

B.5 Further Comparison to Previous Methods

Manifold Sampling Methods   Previous manifold sampling methods assume that the initialization is on the manifold. One may wonder if we can obtain such an initialization by optimization algorithms so that we can still use previous methods when the sampler starts outside the manifold. This will not work because the initialization must be exactly on the manifold whereas the solutions found by optimization always have some intolerable error. Finding a point that is exactly on the manifold without any prior knowledge is by itself a hard problem. Therefore, we are not able to compare our methods with previous methods when there are no known in-domain points, such as the income, loan and image classification tasks in Section 6.

Moment Constraints   As mentioned in the related work, sampling with moment constraints $\mathbb{E}_q[g]$ cannot guarantee every sample to satisfy the constraint. To empirically show the difference between our methods and this type of methods, we compare O-Langevin to Control+ Langevin, which is a recently proposed moment constraint sampling method [24], on the income classification task. We report the mean and the maximum value of the fairness loss in Figure 8. While both methods have small mean fairness loss, the maximum value of O-Langevin is always much smaller than that of Control+Langevin. This suggests that every sample of our method satisfies the constraint well whereas some samples of Control+Langevin violate the constraint significantly, since the moment constraint can only guarantee the mean value instead of the value of each sample.

---

[2]https://www.kaggle.com/wendykan/lending-club-loan-data

Figure 8: Every sample of O-Langevin satisfies the constraint well whereas some samples of Control+ Langevin violate the constraint significantly.