GENERATIVE PRINCIPAL COMPONENT ANALYSIS

ABSTRACT

In this paper, we study the problem of principal component analysis with generative modeling assumptions, adopting a general model for the observed matrix that encompasses notable special cases, including spiked matrix recovery and phase retrieval. The key assumption is that the underlying signal lies near the range of an L-Lipschitz continuous generative model with bounded k-dimensional inputs. We propose a quadratic estimator, and show that it enjoys a statistical rate of order $\sqrt{\frac{k \log L}{m}}$ on the squared error, where m is the number of samples. Moreover, we provide a variant of the classic power method, which projects the calculated data onto the range of the generative model during each iteration. We show that under suitable conditions, this method converges linearly to a point achieving the above-mentioned statistical rate. This rate is conjectured in (Aubin et al., 2019; Cocola et al., 2020) to be the best possible even when we only restrict to the special case of spiked matrix models. We perform experiments on various image datasets for spiked matrix and phase retrieval models, and illustrate performance gains of our method to the classic power method and the truncated power method devised for sparse principal component analysis.

1 INTRODUCTION

Principal component analysis (PCA) is one of the most popular techniques for data processing and dimensionality reduction (Jolliffe, 1986), with an abundance of applications such as image recognition (Hancock et al., 1996), gene expression data analysis (Alter et al., 2000), and clustering (Ding & He, 2004; Liu & Tan, 2019). PCA seeks to find the directions that capture maximal variances in vector-valued data. In detail, letting $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_m$ be m realizations of a random vector $\mathbf{x} \in \mathbb{R}^n$ with a population covariance matrix $\bar{\mathbf{\Sigma}} \in \mathbb{R}^{n \times n}$, PCA aims to reconstruct the top r principal eigenvectors of $\bar{\mathbf{\Sigma}}$. The first principal eigenvector can then be computed as follows:

$$\mathbf{u}_1 = \arg \max_{\mathbf{w} \in \mathbb{R}^n} \mathbf{w}^T \mathbf{\Sigma} \mathbf{w} \quad \text{s.t.} \quad \|\mathbf{w}\|_2 = 1, \tag{1}$$

where the empirical covariance matrix is defined as $\mathbf{\Sigma} := \frac{1}{m} \sum_{i=1}^{m} (\mathbf{x}_i - \mathbf{c})(\mathbf{x}_i - \mathbf{c})^T$, with $\mathbf{c} := \frac{1}{m} \sum_{i=1}^{m} \mathbf{x}_i$. In addition, subsequent principal eigenvectors can be estimated by similar optimization problems subject to being orthogonal to the previous vectors.

PCA is consistent in the conventional setting where the dimension of the data n is relatively small compared to the sample size m (Anderson, 1962), but leads to rather poor estimates in the high-dimensional setting where $m \ll n$. In particular, it has been shown in various papers that the empirical principal eigenvectors are no longer consistent estimates of their population counterparts (Nadler, 2008; Johnstone & Lu, 2009; Jung & Marron, 2009; Birnbaum et al., 2013). In order to tackle the curse of dimensionality, a natural approach is to impose certain structural constraints on the principal eigenvectors, e.g., chosen based on prior knowledge. One of the most widely used assumptions is that the principal eigenvectors are sparse, and this gives rise to the problem of sparse principal component analysis (SPCA) (Zou et al., 2006). In particular, for recovering the top principal eigenvector, the optimization problem of SPCA is given by

$$\mathbf{u}_1 = \arg \max_{\mathbf{w} \in \mathbb{R}^n} \mathbf{w}^T \mathbf{\Sigma} \mathbf{w} \quad \text{s.t.} \quad \|\mathbf{w}\|_2 = 1, \|\mathbf{w}\|_0 \leq s, \tag{2}$$

where $\|\mathbf{w}\|_0 = |\{i : w_i \neq 0\}|$ denotes the number of non-zero entries of $\mathbf{w}$, and $s \in \mathbb{N}$ represents the sparsity level. In addition to reducing the effective number of parameters, the sparsity assumption also enhances the interpretability (Zou et al., 2006).

Departing momentarily from the PCA problem, recent years have seen tremendous advances in deep generative models in a wide variety of real-world applications (Foster, 2019). This has motivated a new perspective of the related problem of compressed sensing (CS), in which the standard sparsity assumption is replaced by a generative modeling assumption. That is, the underlying signal is assumed to lie near the range of a (deep) generative model (Bora et al., 2017). The authors of (Bora et al., 2017) characterized the number of samples required to attain an accurate recovery, and also presented numerical results on image datasets showing that compared to sparsity-based methods, generative priors can lead to large reductions (e.g., a factor of 5 to 10) in the number of measurements needed to recover the signal up to a given accuracy. Additional numerical and theoretical results concerning inverse problems using generative models have been provided in (Van Veen et al., 2018; Heckel & Hand, 2019; Jalal et al., 2020; Liu & Scarlett, 2020a; Ongie et al., 2020; Whang et al., 2020; Jalal et al., 2021; Nguyen et al., 2021), among others.

In this paper, following the developments in both PCA/SPCA and inverse problems with generative priors, we study the use of generative priors in principal component analysis (GPCA), which gives a generative counterpart of SPCA in (2), formulated as follows:

$$\mathbf{u}_1 = \arg \max_{\mathbf{w} \in \mathbb{R}^n} \mathbf{w}^T \mathbf{\Sigma} \mathbf{w} \quad \text{s.t.} \quad \mathbf{w} \in \text{Range}(G), \tag{3}$$

where G is a (pre-trained) generative model, which we assume has a range contained in the unit sphere of $\mathbb{R}^n$.[1] Similarly to SPCA, the motivation for this problem is to incorporate prior knowledge on the vector being recovered (or alternatively, a prior preference), and to permit meaningful recovery and theoretical bounds even in the high-dimensional regime $m \ll n$.

1.1 RELATED WORK

In this subsection, we summarize some relevant works, which can roughly be divided into (i) the SPCA problem, and (ii) signal recovery with generative models.

SPCA: It has been proved that the SPCA problem in (2) attains the optimal statistical rate $\sqrt{s \log n/m}$ (Vu & Lei, 2012), where m is the number of samples, n is the ambient dimension, and s is the sparsity level of the first principal eigenvector. However, due to the combinatorial constraint, the computation of (2) is intractable. To address this computational issue, in recent years, a huge body of practical approaches for estimating sparse principal eigenvectors have been proposed in the literature, including (d'Aspremont et al., 2007; Vu et al., 2013; Chang et al., 2016; Moghaddam et al., 2006; d'Aspremont et al., 2008; Jolliffe et al., 2003; Zou et al., 2006; Shen & Huang, 2008; Journée et al., 2010; Hein & Bühler, 2010; Kuleshov, 2013; Yuan & Zhang, 2013; Asteris et al., 2011; Papailiopoulos et al., 2013), just to name a few.

Notably, statistical guarantees for several approaches have been provided. The authors of (Yuan & Zhang, 2013) propose the truncated power method (TPower), which adds a truncation operation to the power method to ensure the desired level of sparsity. It is shown that this approach attains the optimal statistical rate under appropriate initialization. While most approaches for SPCA only focus on estimating the first principal eigenvector, with a certain deflation method (Mackey, 2008) being leveraged to reconstruct the rest, an iterative thresholding approach is proposed in (Ma, 2013), and is shown to attain a near-optimal statistical rate when estimating multiple individual principal eigenvectors. In addition, the authors of (Cai et al., 2013) propose a regression-type method that gives an optimal principal subspace estimator. Both works (Ma, 2013; Cai et al., 2013) rely on the assumption of a spiked covariance model to ensure a good initial vector. To avoid the spiked covariance model assumption, the work (Wang et al., 2014) proposes a two-stage procedure that attains the optimal subspace estimator in polynomial time.

Signal recovery with generative models: Since the seminal work (Bora et al., 2017), there has been a substantial volume of papers studying various inverse problems with generative priors. One of the problems more closely related to PCA is spectral initialization in phase retrieval, which amounts to solving an eigenvalue problem. Phase retrieval with generative priors has been studied in (Hand et al., 2018; Hyder et al., 2019; Jagatap & Hegde, 2019; Wei et al., 2019; Shamshad & Ahmed,

---

[1]Similarly to (Liu et al., 2020; 2021a), we assume that the range of G is contained in the unit sphere for convenience. Our results transfer to any general (unnormalized) generative model readily by considering its normalized version. See Remark 1 for a detailed discussion.

2020; Aubin et al., 2020; Liu et al., 2021b). In particular, the work (Hand et al., 2018) models the underlying signal as being in the range of a fully-connected ReLU neural network with no offsets, and all the weight matrices of the ReLU neural network are assumed to have i.i.d. zero-mean Gaussian entries. In addition, the neural network needs to be sufficiently expansive in the sense that $n_i \geq \Omega(n_{i-1} \log n_{i-1})$, where n_i is the width of the i-th layer. Under these assumptions, the authors establish favorable global optimization landscapes for the corresponding objective, and derive near-optimal sample complexity upper bound. They minimize the objective function directly over the latent space in $\mathbb{R}^k$ using gradient descent, which may suffer from local minima in general optimization landscapes (Hyder et al., 2019; Shah & Hegde, 2018).

In (Aubin et al., 2020), the assumptions on the neural network are similar to those in (Hand et al., 2018), relaxing to general activation functions (beyond ReLU) and $n_i \geq \Omega(n_{i-1})$. The authors focus on the high dimensional regime where $n, m, k \to \infty$ with the ratio m/n being fixed, and assume that the input vector in $\mathbb{R}^k$ is drawn from a separable distribution. They derive sharp asymptotics for the information-theoretically optimal performance and for the associated approximate message passing (AMP) algorithm. Both works (Hand et al., 2018; Aubin et al., 2020) focus on noiseless phase retrieval. When only making the much milder assumption that the generative model is Lipschitz continuous, with no assumption on expansiveness, Gaussianity, and offsets, similarly to sparse phase retrieval, a spectral initialization step is typically required in order to accurately reconstruct the signal (Netrapalli et al., 2015; Candès et al., 2015). The authors of (Liu et al., 2021b) propose an optimization problem similar to (3) for the spectral initialization for phase retrieval with generative models. It was left open in (Liu et al., 2021b) how to solve (or approximate sufficiently accurately) the optimization problem in practice.

Understanding the eigenvalues of spiked random matrix models has been a central problem of random matrix theory, and spiked matrices have been widely used in the statistical analysis of SPCA. Recently, theoretical guarantees concerning spiked matrix models with generative priors have been provided in (Aubin et al., 2019; Cocola et al., 2020). In particular, in (Aubin et al., 2019), the assumptions are similar to those in (Aubin et al., 2020), except that the neural network is assumed to have exactly one hidden layer. The Bayes-optimal performance is analyzed, and it is shown that the AMP algorithm can attain this optimal performance. In addition, the authors of (Aubin et al., 2019) also propose the linearized approximate message passing (LAMP) algorithm, which is a spectral algorithm specifically designed for a single-layer feedforward neural network with no bias term. The authors show its superiority to classical PCA via numerical results on the Fashion-MNIST dataset. In (Cocola et al., 2020), the same assumptions are made as those in (Hand et al., 2018) on the neural network, and the authors demonstrate the benign global geometry for a nonlinear least squares objective. Similarly to (Hand et al., 2018), the objective is minimized over $\mathbb{R}^k$ using a gradient descent algorithm, which can get stuck in local minima for general global geometries.

Compared to the above discussed works that use generative models, we make no assumption on expansiveness, Gaussianity, and offsets for the generative model, and we consider a data model that simultaneously encompasses both spiked matrix and phase retrieval models, among others.

1.2 CONTRIBUTIONS

The main contributions of this paper are threefold:

- We study eigenvalue problems with generative priors (including GPCA), and characterize the statistical rate of a quadratic estimator similar to (3) under suitable assumptions.

- We propose a variant of the classic power method, which uses an additional projection operation to ensure that the output of each iteration lies in the range of a generative model. We refer to our method as projected power method (PPower). We further show that under appropriate conditions (most notably, assuming exact projections are possible), PPower obtains a solution achieving a statistical rate that is conjectured to be optimal in (Aubin et al., 2019; Cocola et al., 2020) for spiked matrix models.

- For spiked matrix and phase retrieval models, we perform numerical experiments on image datasets, and demonstrate that when the number of samples is relatively small compared to the ambient dimension, PPower leads to significantly better performance compared to the classic power method and TPower.

1.3 Notation

We use upper and lower case boldface letters to denote matrices and vectors respectively. We write $[N] = \{1, 2, \cdots, N\}$ for a positive integer N, and we use $\mathbf{I}_N$ to denote the identity matrix in $\mathbb{R}^{N \times N}$. A *generative model* is a function $G : \mathcal{D} \to \mathbb{R}^n$, with latent dimension k, ambient dimension n, and input domain $\mathcal{D} \subseteq \mathbb{R}^k$. We focus on the setting where $k \ll n$. For a set $S \subseteq \mathbb{R}^k$ and a generative model $G : \mathbb{R}^k \to \mathbb{R}^n$, we write $G(S) = \{G(\mathbf{z}) : \mathbf{z} \in S\}$. We use $\|\mathbf{X}\|_{2 \to 2}$ to denote the spectral norm of a matrix $\mathbf{X}$. We define the ℓ_q-ball $B_q^k(r) := \{\mathbf{z} \in \mathbb{R}^k : \|\mathbf{z}\|_q \leq r\}$ for $q \in [0, +\infty]$. $\mathcal{S}^{n-1} := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\|_2 = 1\}$ represents the unit sphere in $\mathbb{R}^n$. The symbols C, C', C'' are absolute constants whose values may differ from line to line.

2 Problem Setup

In this section, we formally introduce the problem, and overview some important assumptions that we adopt. Except where stated otherwise, we will focus on the following setting:

- We have a matrix $\mathbf{V} \in \mathbb{R}^{n \times n}$ satisfying

$$\mathbf{V} = \bar{\mathbf{V}} + \mathbf{E}, \tag{4}$$

  where $\mathbf{E}$ is a perturbation matrix and $\bar{\mathbf{V}}$ is assumed to be positive semidefinite (PSD). For PCA and its constrained variants, $\mathbf{V}$ and $\bar{\mathbf{V}}$ can be thought of as the empirical and population covariance matrices respectively.

- We have an L-Lipschitz continuous generative model $G : B_2^k(r) \to \mathbb{R}^n$. For convenience, similarly to that in (Liu et al., 2020), we assume that $\mathrm{Range}(G) \subseteq \mathcal{S}^{n-1}$.
  Remark 1. *For a general (unnormalized) L-Lipschitz continuous generative model G, we can instead consider a corresponding normalized generative model $\tilde{G} : \mathcal{D} \to \mathcal{S}^{n-1}$ as in (Liu et al., 2021b), where $\mathcal{D} := \{\mathbf{z} \in B_2^k(r) : \|\mathbf{z}\|_2 > R_{\min}\}$ for some $R_{\min} > 0$. Then, the Lipschitz constant of $\tilde{G}$ becomes $L/R_{\min}$. For a d-layer neural network, we typically have $L = n^{\Theta(d)}$ (Bora et al., 2017). Thus, we can set $R_{\min}$ to be as small as $1/n^{\Theta(d)}$ without changing the scaling laws, which makes the dependence on $R_{\min}$ very mild.*

- We aim to solve the following eigenvalue problem with a generative prior:

$$\hat{\mathbf{v}} := \max_{\mathbf{w} \in \mathbb{R}^n} \mathbf{w}^T \mathbf{V} \mathbf{w} \quad \text{s.t.} \quad \mathbf{w} \in \mathrm{Range}(G). \tag{5}$$

  Note that since $\mathrm{Range}(G) \subseteq \mathcal{S}^{n-1}$, we do not need to impose the constraint $\|\mathbf{w}\|_2 = 1$.[2] Since $\mathbf{V}$ is not restricted to being an empirical covariance matrix, (5) is more general than GPCA in (3). However, we slightly abuse terminology and also refer to (5) as GPCA.

- To approximately solve (5), we perform a projected power method (PPower), which gives the following iterative procedure:[3]

$$\mathbf{w}^{(t+1)} = \mathcal{P}_G\left(\mathbf{V}\mathbf{w}^{(t)}\right), \tag{6}$$

  where $\mathcal{P}_G(\cdot)$ is the projection function onto $G(B_2^k(r))$,[4] and the initialization vector $\mathbf{w}^{(0)}$ may be chosen either manually or randomly, e.g., uniform over $\mathcal{S}^{n-1}$. Often the initialization vector $\mathbf{w}^{(0)}$ plays an important role and we may need a careful design of it. For example, for phase retrieval with generative models, as mentioned in (Liu et al., 2021b, Section V), we may choose the column corresponding to the largest diagonal entry of $\mathbf{V}$ as the starting point. See also (Yuan & Zhang, 2013, Section 3) for a discussion on the initialization strategy for TPower devised for SPCA. We present the algorithm corresponding to (6) in Algorithm 1.

---

[2]To find the top r rather than top one principal eigenvectors that are in the range of a generative model, we may follow the common approach to use the iterative deflation method for PCA/SPCA: Subsequent principal eigenvectors are derived by recursively removing the contribution of the principal eigenvectors that are calculated already under the generative model constraint. See for example (Mackey, 2008).

[3]In the iterative procedure, we may replace $\mathbf{V}$ by $\mathbf{V} + \rho \mathbf{I}_n$ for some $\rho \in \mathbb{R}$ to improve convergence.

[4]That is, for any $\mathbf{x} \in \mathbb{R}^n$, $\mathcal{P}_G(\mathbf{x}) := \arg\min_{\mathbf{w} \in \mathrm{Range}(G)} \|\mathbf{w} - \mathbf{x}\|_2$. We will implicitly assume that the projection step can be performed accurately, e.g., (Deshpande et al., 2014; Shah & Hegde, 2018; Peng et al., 2020), though in practice approximate methods might be needed, e.g., via gradient descent (Shah & Hegde, 2018) or GAN-based projection methods (Raj et al., 2019).

Algorithm 1 A projected power method for GPCA (`PPower`)

Input: $\mathbf{V}$, number of iterations T, pre-trained generative model G, initial vector $\mathbf{w}^{(0)}$
Output: $\mathbf{w}^{(T)}$
1: **for** $t = 0, 1, \ldots, T-1$ **do**
2:     $\mathbf{w}^{(t+1)} = \mathcal{P}_G\big(\mathbf{V}\mathbf{w}^{(t)}\big)$
3: **end for**

Remark 2. *To tackle generalized eigenvalue problems encountered in some specific applications, there are variants of the projected power method, which combine a certain power iteration with additional operations to ensure sparsity or enforce other constraints. These applications include but not limited to sparse PCA (Journée et al., 2010; Yuan & Zhang, 2013), phase synchronization (Boumal, 2016; Liu et al., 2017), the hidden clique problem (Deshpande & Montanari, 2015), the joint alignment problem (Chen & Candès, 2018), and cone-constrained PCA (Deshpande et al., 2014; Yi & Neykov, 2020). For example, under the simple spiked Wigner model (Perry et al., 2018) for the observed data matrix $\mathbf{V}$ with the underlying signal being assumed to lie in a convex cone, the authors of (Deshpande et al., 2014) show that cone-constrained PCA can be computed efficiently via a generalized projected power method. In general, the range of a Lipschitz-continuous generative model is non-convex and not a cone. In addition, we consider a matrix model that is more general than the spiked Wigner model.*

Under the assumption that $\mathbf{V}$ is PSD, we have the following useful lemma. The PSD assumption will be satisfied, for example, when $\mathbf{E} = \mathbf{0}$, or when $\mathbf{V}$ is a sample covariance matrix.

Lemma 1. *For any $\mathbf{x} \in \mathbb{R}^n$, let $Q(\mathbf{x}) = \mathbf{x}^T \mathbf{V} \mathbf{x}$. Then, if $\mathbf{V}$ is PSD, the sequence $\{Q(\mathbf{w}^{(t)})\}_{t>0}$ for $\mathbf{w}^{(t)}$ in (6) is monotonically non-decreasing.*

Proof. Since $\mathbf{w}^{(t+1)} = \mathcal{P}_G\big(\mathbf{V}\mathbf{w}^{(t)}\big)$ and $\mathbf{w}^{(t)} \in \mathrm{Range}(G)$, we have

$$\|\mathbf{V}\mathbf{w}^{(t)} - \mathbf{w}^{(t+1)}\|_2 \leq \|\mathbf{V}\mathbf{w}^{(t)} - \mathbf{w}^{(t)}\|_2, \tag{7}$$

and since $\|\mathbf{w}^{(t+1)}\|_2 = \|\mathbf{w}^{(t)}\|_2 = 1$, expanding the square gives

$$\left\langle \mathbf{V}\mathbf{w}^{(t)}, \mathbf{w}^{(t+1)} \right\rangle \geq Q(\mathbf{w}^{(t)}). \tag{8}$$

Then, we obtain

$$Q(\mathbf{w}^{(t+1)}) = \left\langle \mathbf{V}\mathbf{w}^{(t+1)}, \mathbf{w}^{(t+1)} \right\rangle \tag{9}$$

$$= Q(\mathbf{w}^{(t+1)} - \mathbf{w}^{(t)}) + 2\left\langle \mathbf{V}(\mathbf{w}^{(t+1)} - \mathbf{w}^{(t)}), \mathbf{w}^{(t)} \right\rangle + Q(\mathbf{w}^{(t)}) \tag{10}$$

$$\geq 2\left\langle \mathbf{V}(\mathbf{w}^{(t+1)} - \mathbf{w}^{(t)}), \mathbf{w}^{(t)} \right\rangle + Q(\mathbf{w}^{(t)}) \tag{11}$$

$$\geq Q(\mathbf{w}^{(t)}), \tag{12}$$

where (11) follows from the assumption that $\mathbf{V}$ is PSD, and (12) follows from (8). $\square$

3   SPECIALIZED DATA MODELS AND EXAMPLES

In this section, we make more specific assumptions on $\mathbf{V} = \bar{\mathbf{V}} + \mathbf{E}$, starting with the following.

Assumption 1 (Assumption on $\bar{\mathbf{V}}$). *Assume that $\bar{\mathbf{V}}$ is PSD with eigenvalues $\bar{\lambda}_1 > \bar{\lambda}_2 \geq \ldots \geq \bar{\lambda}_n \geq 0$. We use $\bar{\mathbf{x}}$ (a unit vector) to represent the eigenvector of $\bar{\mathbf{V}}$ that corresponds to $\bar{\lambda}_1$.*

In the following, it is useful to think of $\bar{\mathbf{x}}$ is being close to the range of the generative model G. In the special case of (3), letting m be the number of samples, it is natural to derive that the upper bound of $\|\mathbf{E}\|_{2\to2}$ grows linearly in $(n/m)^b$ for some positive constant b such as $\frac{1}{2}$ or 1 (with high probability; see, e.g., (Vershynin, 2010, Corollary 5.35)). In the following, we consider general scenarios with $\mathbf{V}$ depending on m samples (see below for specific examples). Similarly to (Yuan & Zhang, 2013), we may consider a restricted version of $\|\mathbf{E}\|_{2\to2}$, leading to the following.

Assumption 2 (Assumption on $\mathbf{E}$). *Let S_1, S_2 be two (arbitrary) finite sets in $\mathbb{R}^n$ satisfying $m = \Omega(\log(|S_1| \cdot |S_2|))$. Then, we have for all $\mathbf{s}_1 \in S_1$ and $\mathbf{s}_2 \in S_2$ that*

$$\left| \mathbf{s}_1^T \mathbf{E} \mathbf{s}_2 \right| \leq C \sqrt{\frac{\log(|S_1| \cdot |S_2|)}{m}} \cdot \|\mathbf{s}_1\|_2 \cdot \|\mathbf{s}_2\|_2, \tag{13}$$

where C is an absolute constant. In addition, we have $\|\mathbf{E}\|_{2 \to 2} = O(n/m)$.[5]

The following examples show that when the number of measurements is sufficiently large, the data matrices corresponding to certain spiked matrix and phase retrieval models satisfy the above assumptions with high probability. Short proofs are given in Appendix A for completeness.

Example 1 (Spiked covariance model). *In the spiked covariance model (Johnstone & Lu, 2009; Deshpande & Montanari, 2016), the observed vectors $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_m \in \mathbb{R}^n$ are of the form*

$$\mathbf{x}_i = \sum_{q=1}^{r} \sqrt{\beta_q} u_{q,i} \mathbf{s}_q + \mathbf{z}_i, \tag{14}$$

where $\mathbf{s}_1, \ldots, \mathbf{s}_r \in \mathbb{R}^n$ are orthonormal vectors that we want to estimate, while $\mathbf{z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ and $u_{q,i} \sim \mathcal{N}(0, 1)$ are independent and identically distributed. In addition, $\beta_1, \ldots, \beta_r$ are positive constants that dictate the signal-to-noise ratio (SNR). To simplify the exposition, we focus on the rank-one case and drop the subscript $q \in [r]$. Let

$$\mathbf{V} = \frac{1}{m} \sum_{i=1}^{m} (\mathbf{x}_i \mathbf{x}_i^T - \mathbf{I}_n), \tag{15}$$

and $\bar{\mathbf{V}} = \mathbb{E}[\mathbf{V}] = \beta \mathbf{s} \mathbf{s}^T$.[6] Then, $\bar{\mathbf{V}}$ satisfies Assumption 1 with $\bar{\lambda}_1 = \beta > 0$, $\bar{\lambda}_2 = \ldots = \bar{\lambda}_n = 0$, and $\bar{\mathbf{x}} = \mathbf{s}$. In addition, let $\mathbf{E} = \mathbf{V} - \bar{\mathbf{V}}$. From the Bernstein-type inequality (Vershynin, 2010, Proposition 5.10) for the sum of sub-exponential random variables, we observe that for any finite sets $S_1, S_2 \subset \mathbb{R}^n$, when $m = \Omega(\log(|S_1| \cdot |S_2|))$, with probability $1 - e^{-\Omega(\log(|S_1| \cdot |S_2|))}$, $\mathbf{E}$ satisfies (13) in Assumption 2. Moreover, standard concentration arguments give $\|\mathbf{E}\|_{2 \to 2} = O(n/m)$ with probability $1 - e^{-\Omega(n)}$.

Remark 3. *We can also consider the simpler spiked Wigner model (Perry et al., 2018; Chung & Lee, 2019) where $\mathbf{V} = \beta \mathbf{s} \mathbf{s}^T + \frac{1}{\sqrt{n}} \mathbf{H}$, with the signal $\mathbf{s}$ being a unit vector, $\beta > 0$ being an SNR parameter, and $\mathbf{H} \in \mathbb{R}^{n \times n}$ being a symmetric matrix with entries drawn i.i.d. (up to symmetry) from $\mathcal{N}(0, 1)$. In this case, when $m = n$ is sufficiently large, with high probability, $\bar{\mathbf{V}} := \mathbb{E}[\mathbf{V}] = \beta \mathbf{s} \mathbf{s}^T$ and $\mathbf{E} := \mathbf{V} - \bar{\mathbf{V}}$ similarly satisfy Assumptions 1 and 2 respectively.*

Example 2 (Phase retrieval). *Let $\mathbf{A} \in \mathbb{R}^{m \times n}$ be a matrix having i.i.d. $\mathcal{N}(0, 1)$ entries, and let $\mathbf{a}_i^T$ be the i-th row of $\mathbf{A}$. For some unit vector $\mathbf{s}$, suppose that the observed vector is $\mathbf{y} = |\mathbf{A}\mathbf{s}|$, where the absolute value is applied element-wise.[7] We construct the weighted empirical covariance matrix (Zhang et al., 2017; Liu et al., 2021b)*

$$\mathbf{V} = \frac{1}{m} \sum_{i=1}^{m} \left(y_i \mathbf{a}_i \mathbf{a}_i^T \mathbf{1}_{\{l < y_i < u\}} - \gamma \mathbf{I}_n \right), \tag{16}$$

where $u > l > 1$ are positive constants, and for $g \sim \mathcal{N}(0, 1)$, $\gamma := \mathbb{E}\big[|g| \mathbf{1}_{\{l < |g| < u\}}\big]$. Let $\bar{\mathbf{V}} = \mathbb{E}[\mathbf{V}] = \beta \mathbf{s} \mathbf{s}^T$, where $\beta := \mathbb{E}\big[(|g|^3 - |g|) \mathbf{1}_{\{l < |g| < u\}}\big]$. Then, $\bar{\mathbf{V}}$ satisfies Assumption 1 with $\bar{\lambda}_1 = \beta > 0$, $\bar{\lambda}_2 = \ldots = \bar{\lambda}_n = 0$, and $\bar{\mathbf{x}} = \mathbf{s}$. In addition, letting $\mathbf{E} = \mathbf{V} - \bar{\mathbf{V}}$, we have similarly to Example 1 that $\mathbf{E}$ satisfies Assumption 2 with high probability.

---

[5]For the spectral norm of $\mathbf{E}$, one often expects an even tighter bound $O(\sqrt{n/m})$, but we use $O(n/m)$ to simplify the analysis of our examples. Moreover, at least under the typical scaling where L is polynomial in n (Bora et al., 2017), the upper bound for $\|\mathbf{E}\|_{2 \to 2}$ can be easily relaxed to $O\big((n/m)^b\big)$ for any positive constant b, without affecting the scaling of our derived statistical rate.

[6]To avoid non-essential complications, β is typically assumed to be known (Johnstone & Lu, 2009).

[7]Without loss of generality, we assume that $\mathbf{s}$ is a unit vector. For a general signal $\mathbf{s}$, we may instead focus on estimating $\bar{\mathbf{s}} = \mathbf{s}/\|\mathbf{s}\|_2$, and simply use $\frac{1}{m} \sum_{i=1}^{m} y_i$ to approximates $\|\mathbf{s}\|_2$.

4 MAIN RESULTS

The following theorem concerns globally optimal solutions of (5). The proof is given in Appendix B.

Theorem 1. *Let* $\mathbf{V} = \bar{\mathbf{V}} + \mathbf{E}$ *with Assumptions 1 and 2 being satisfied by* $\bar{\mathbf{V}}$ *and* $\mathbf{E}$ *respectively, and let* $\mathbf{x}_G := \mathcal{P}_G(\bar{\mathbf{x}}) = \arg\min_{\mathbf{w} \in \text{Range}(G)} \|\mathbf{w} - \bar{\mathbf{x}}\|_2$. *Suppose that* $\hat{\mathbf{v}}$ *is a globally optimal solution to* (5). *Then, for any* $\delta \in (0, 1)$, *we have*

$$\min\left\{\|\bar{\mathbf{x}} - \hat{\mathbf{v}}\|_2^2, \|\bar{\mathbf{x}} + \hat{\mathbf{v}}\|_2^2\right\} \le O\left(\frac{\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}}}{\bar{\lambda}_1 - \bar{\lambda}_2}\right) + \frac{4\bar{\lambda}_1}{\bar{\lambda}_1 - \bar{\lambda}_2} \cdot \|\mathbf{x}_G - \bar{\mathbf{x}}\|_2 + O\left(\frac{\delta n}{m(\bar{\lambda}_1 - \bar{\lambda}_2)}\right). \tag{17}$$

We note that since both $\bar{\mathbf{x}}$ and $-\bar{\mathbf{x}}$ are leading eigenvectors of $\bar{\mathbf{V}}$, without further assumptions, a quadratic estimator of the form (5) can only recover the signal up to a global sign flip. From Theorem 1, we observe that when there is no representation error, i.e., $\bar{\mathbf{x}} \in \text{Range}(G)$, under the scaling $\bar{\lambda}_1 - \bar{\lambda}_2 = \Theta(1)$, $L = n^{\Omega(1)}$, and $\delta = O(\sqrt{m}/n)$, the solution $\hat{\mathbf{v}}$ to the quadratic estimator in (5) satisfies $\min\left\{\|\bar{\mathbf{x}} - \hat{\mathbf{v}}\|_2^2, \|\bar{\mathbf{x}} + \hat{\mathbf{v}}\|_2^2\right\} = O(\sqrt{\frac{k \log L}{m}})$.

Before providing our main theorem for PPower described in (6), we present the following important lemma, whose proof is presented in Appendix C. To simplify the statement of the lemma, we fix δ to be $O(1/n)$, though a more general form analogous to Theorem 1 is also possible.

Lemma 2. *Let* $\mathbf{V} = \bar{\mathbf{V}} + \mathbf{E}$ *with Assumptions 1 and 2 being satisfied by* $\bar{\mathbf{V}}$ *and* $\mathbf{E}$ *respectively, and further assume that* $\bar{\mathbf{x}} \in \text{Range}(G)$. *Let* $\bar{\gamma} = \bar{\lambda}_2/\bar{\lambda}_1$ *with* $\bar{\lambda}_1 = \Theta(1)$. *Then, for all* $\mathbf{s} \in \text{Range}(G)$ *satisfying* $\mathbf{s}^T\bar{\mathbf{x}} > 0$, *let* $\hat{\mathbf{s}} = \mathcal{P}_G(\mathbf{V}\mathbf{s})$, *and we have*

$$\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2 \le \frac{2\bar{\gamma}\|\mathbf{s} - \bar{\mathbf{x}}\|_2}{\mathbf{s}^T\bar{\mathbf{x}}} + O\left(\frac{1}{\mathbf{s}^T\bar{\mathbf{x}}}\sqrt{\frac{k \log(nLr)}{m}}\right). \tag{18}$$

Remark 4. *The assumption* $\mathbf{s}^T\bar{\mathbf{x}} > 0$ *will be generally satisfied when the range of* G *only contains nonnegative vectors. Such a condition holds for a neural network with the popular ReLU or Sigmoid activation functions. In addition, as mentioned in some works studying nonnegative SPCA (Zass & Shashua, 2007; Sigg & Buhmann, 2008; Asteris et al., 2014), for several practical fields such as economics, bioinformatics, and computer vision, it is natural to assume that the underlying signal has no negative entries. More generally, the assumption* $\mathbf{s}^T\bar{\mathbf{x}} > 0$ *can be removed if we additionally have that* $-\bar{\mathbf{x}}$ *is also contained in the range of* G. *For this case, when* $\mathbf{s}^T\bar{\mathbf{x}} < 0$, *we can instead derive an upper bound for* $\|\hat{\mathbf{s}} + \bar{\mathbf{x}}\|_2$.

Based on the above lemma, we have the following theorem, whose proof is given in Appendix D.

Theorem 2. *Suppose that the assumptions on the data model* $\mathbf{V} = \bar{\mathbf{V}} + \mathbf{E}$ *are the same as those in Lemma 2, and assume that there exists a* $t_0 \in \mathbb{N}$ *such that* $\bar{\mathbf{x}}^T\mathbf{w}^{(t_0)} \ge 2\bar{\gamma} + \nu$, *where* $\bar{\gamma} = \bar{\lambda}_2/\bar{\lambda}_1$ *and* ν *is a sufficiently small positive constant. Let* $\mu_0 = 2\bar{\gamma}/(\bar{\mathbf{x}}^T\mathbf{w}^{(t_0)}) < 1$. *Finally, suppose that* $m = \Omega(k \log(nLr))$ *with a large enough implied constant. Then, for any* $t \ge t_0$, *we have either*

$$\|\mathbf{w}^{(t)} - \bar{\mathbf{x}}\|_2 \le \frac{C}{(1 - \mu_0)\nu}\sqrt{\frac{k \log(nLr)}{m}}, \tag{19}$$

or

$$\|\mathbf{w}^{(t+1)} - \bar{\mathbf{x}}\|_2 \le \mu_0\|\mathbf{w}^{(t)} - \bar{\mathbf{x}}\|_2 + \frac{C}{\nu}\sqrt{\frac{k \log(nLr)}{m}} \le (1 - \xi)\|\mathbf{w}^{(t)} - \bar{\mathbf{x}}\|_2, \tag{20}$$

where C *is an absolute constant, and* ξ *is strictly positive.*

If $L = n^{\Omega(1)}$, then (20) reveals that we have established conditions under which PPower in (6) converges linearly to a point achieving the statistical rate of order $\sqrt{\frac{k \log L}{m}}$. Based on the minimax rates for SPCA (Vu & Lei, 2012; Birnbaum et al., 2013) and the information-theoretic lower bounds for CS with generative models (Liu & Scarlett, 2020b; Kamath et al., 2020), the optimal rate for GPCA is naturally conjectured to be of the same order $\sqrt{\frac{k \log L}{m}}$.

Regarding the assumption $\bar{\mathbf{x}}^T \mathbf{w}^{(t_0)} \geq 2\bar{\gamma} + \nu$, we note that if $t_0 = 1$, this condition can be viewed as having a good initialization. For both Examples 1 and 2, we have $\bar{\gamma} = 0$. Thus, for the spiked covariance and phase retrieval models corresponding to these examples, the assumption $\bar{\mathbf{x}}^T \mathbf{w}^{(t_0)} \geq 2\bar{\gamma} + \nu$ reduces to $\bar{\mathbf{x}}^T \mathbf{w}^{(t_0)} \geq \nu$ for a sufficiently small positive constant ν, which results in a mild assumption. Such an assumption is also required for the projected power method devised for cone-constrained PCA under the simple spiked Wigner model, with the underlying signal being assumed to lie in a convex cone; see (Deshpande et al., 2014, Corollary 4.1).

Perhaps the main caveat to Theorem 2 is that it assumes the projection step can be performed exactly. However, this is a standard assumption in analyses of projected gradient methods, e.g., see (Shah & Hegde, 2018), and both gradient-based projection and GAN-based projection have been shown to be highly effective in practice (Shah & Hegde, 2018; Raj et al., 2019).

5 EXPERIMENTS

In this section, we experimentally study the performance of Algorithm 1 (`PPower`). We note that these experiments are intended as a simple proof of concept rather than seeking to be comprehensive, as our contributions are primarily theoretical. We compare with the truncated power method (`TPower`) devised for SPCA proposed in (Yuan & Zhang, 2013, Algorithm 1) and the vanilla power method (`Power`) that performs the iterative procedure $\mathbf{w}^{(t+1)} = (\mathbf{V}\mathbf{w}^{(t)})/\|\mathbf{V}\mathbf{w}^{(t)}\|_2$. For a fair comparison, for `PPower`, `TPower`, and `Power`, we use the same initial vector, i.e., as mentioned in (Liu et al., 2021b, Section V), we choose the initialization vector $\mathbf{w}^{(0)}$ as the column of $\mathbf{V}$ that corresponds to its largest diagonal entry. For all these three algorithms, the total number of iterations T is set to be 30. To compare the performance across algorithms, we use the scale-invariant Cosine Similarity metric defined as $\text{Cossim}\left(\bar{\mathbf{x}}, \mathbf{w}^{(T)}\right) := \frac{\langle \bar{\mathbf{x}}, \mathbf{w}^{(T)} \rangle}{\|\bar{\mathbf{x}}\|_2 \|\mathbf{w}^{(T)}\|_2}$, where $\bar{\mathbf{x}}$ is the ground-truth signal to estimate, and $\mathbf{w}^{(T)}$ denotes the output vector of the algorithm.

The experiments are performed on the MNIST (LeCun et al., 1998) and Fashion-MNIST (Xiao et al., 2017) datasets, with the numerical results for the Fashion-MNIST dataset being presented in Appendix E. The MNIST dataset consists of $60,000$ images of handwritten digits. The size of each image in the MNIST dataset is 28×28, and thus $n = 784$. To reduce the impact of local minima, we perform 10 random restarts, and choose the best among these. The cosine similarity is averaged over the test images, and also over these 10 random restarts. The generative model G is set to be a pre-trained variational autoencoder (VAE) model with latent dimension $k = 20$. The encoder and decoder are both fully connected neural networks with two hidden layers, with the architecture being $20 - 500 - 500 - 784$. The VAE is trained by the Adam optimizer with a mini-batch size of 100 and a learning rate of 0.001. The projection step $\mathcal{P}_G(\cdot)$ is solved by the Adam optimizer with a learning rate of 0.03 and 200 steps. In each iteration of `TPower`, the calculated entries are truncate to zero except for the largest q entries, where $q \in \mathbb{N}$ is a tuning parameter. Since for `TPower`, q is usually selected as an integer larger than the true sparsity level, and it is unlikely that the image of the MNIST dataset can be well approximated by a k-sparse vector with $k = 20$, we choose a relatively large q, namely $q = 150$.

We perform two sets of experiments; one for the spiked covariance model, and another for phase retrieval. All experiments are run using Python 3.6 and Tensorflow 1.5.0, with a NVIDIA GeForce GTX 1080 Ti 11GB GPU.

1. **Spiked covariance model (Example 1)**: The numerical results are shown in Figures 1 and 2. We observe from Figure 1 that `Power` and `TPower` attain poor reconstructions, and the generative prior based method `PPower` attains significantly better reconstructions. To illustrate the effect of the sample size m, we fix the SNR parameter $\beta = 1$ and vary m in $\{100, 200, 300, 400, 500\}$. In addition, to illustrate the effect of the SNR parameter β, we fix $m = 300$, and vary β in $\{0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4\}$. From Figure 2, we observe that for these settings of m and β, `PPower` always leads to a much higher cosine similarity compared to `Power` and `TPower`, which is natural given the more precise modeling assumptions used.

2. **Phase retrieval (Example 2)**: The results are shown in Figure 2 (Right) and Figure 3. Again, we can observe that `PPower` significantly outperforms `Power` and `TPower`. In particular, for sparse phase retrieval, when performing experiments on image datasets, even for the noiseless

(a) $\beta = 1$ and $m = 200$ (b) $\beta = 2$ and $m = 100$

Figure 1: Examples of reconstructed images of the MNIST dataset for the spiked covariance model.

(a) Fixing $\beta = 1$ and varying m (b) Fixing $m = 300$ and varying β (c) Analog of (a) for phase ret.

Figure 2: Quantitative comparisons about the performance of `Power`, `TPower` and `PPower` according to Cosine Similarity over the MNIST dataset for the spiked covariance model (Left/Middle) and for phase retrieval (Right).

setting, solving an eigenvalue problem similar to (5) can typically only serve as a spectral initialization step, with a subsequent iterative algorithm being required to refine the initial guess. In view of this, it is notable that for phase retrieval with generative priors, `PPower` can return meaningful reconstructed images for $m = 200$, which is small compared to $n = 784$.

6 CONCLUSION AND FUTURE WORK

We have proposed a quadratic estimator for eigenvalue problems with generative models, and we showed that this estimator attains a statistical rate of order $\sqrt{\frac{k \log L}{m}}$ on the squared error. We provided a projected power method to efficiently solve (modulo the complexity of the projection step) the corresponding optimization problem, and showed that our method converges linearly to a point achieving a statistical rate of order $\sqrt{\frac{k \log L}{m}}$ under suitable conditions. An interesting direction for further research is to provide guarantees for estimating multiple individual signal vectors (or multiple spikes), or the subspace spanned by these signal vectors.

(a) $m = 200$ (b) $m = 400$

Figure 3: Examples of reconstructed images of the MNIST dataset for phase retrieval.

REFERENCES

Orly Alter, Patrick O Brown, and David Botstein. Singular value decomposition for genome-wide expression data processing and modeling. *PNAS*, 97(18):10101–10106, 2000.

Theodore Wilbur Anderson. An introduction to multivariate statistical analysis. Technical report, Wiley New York, 1962.

Megasthenis Asteris, Dimitris S Papailiopoulos, and George N Karystinos. Sparse principal component of a rank-deficient matrix. In *Int. Symp. Inf. Theory (ISIT)*, pp. 673–677. IEEE, 2011.

Megasthenis Asteris, Dimitris Papailiopoulos, and Alexandros Dimakis. Nonnegative sparse PCA with provable guarantees. In *Int. Conf. Mach. Learn. (ICML)*, pp. 1728–1736. PMLR, 2014.

Benjamin Aubin, Bruno Loureiro, Antoine Maillard, et al. The spiked matrix model with generative priors. *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, 32:8366–8377, 2019.

Benjamin Aubin, Bruno Loureiro, Antoine Baker, et al. Exact asymptotics for phase retrieval and compressed sensing with random generative priors. In *Math. Sci. Mach. Learn. (MSML)*, pp. 55–73. PMLR, 2020.

Aharon Birnbaum, Iain M Johnstone, Boaz Nadler, and Debashis Paul. Minimax bounds for sparse PCA with noisy high-dimensional data. *Ann. Stat.*, 41(3):1055, 2013.

Ashish Bora, Ajil Jalal, Eric Price, and Alexandros G Dimakis. Compressed sensing using generative models. In *Int. Conf. Mach. Learn. (ICML)*, pp. 537–546, 2017.

Nicolas Boumal. Nonconvex phase synchronization. *SIAM J. Optim.*, 26(4):2355–2377, 2016.

T Tony Cai, Zongming Ma, and Yihong Wu. Sparse PCA: Optimal rates and adaptive estimation. *Ann. Stat.*, 41(6):3074–3110, 2013.

Emmanuel J Candès, Xiaodong Li, and Mahdi Soltanolkotabi. Phase retrieval via Wirtinger flow: Theory and algorithms. *IEEE Trans. Inf. Theory*, 61(4):1985–2007, 2015.

Xiaojun Chang, Feiping Nie, Yi Yang, et al. Convex sparse PCA for unsupervised feature learning. *ACM T. Knowl. Discov. D.*, 11(1):1–16, 2016.

Yuxin Chen and Emmanuel J Candès. The projected power method: An efficient algorithm for joint alignment from pairwise differences. *Comm. Pure Appl. Math.*, 71(8):1648–1714, 2018.

Hye Won Chung and Ji Oon Lee. Weak detection of signal in the spiked Wigner model. In *Int. Conf. Mach. Learn. (ICML)*, pp. 1233–1241. PMLR, 2019.

Jorio Cocola, Paul Hand, and Vlad Voroninski. Nonasymptotic guarantees for spiked matrix recovery with generative priors. In *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, volume 33, 2020.

Alexandre d'Aspremont, Laurent El Ghaoui, Michael I Jordan, and Gert RG Lanckriet. A direct formulation for sparse PCA using semidefinite programming. *SIAM Rev.*, 49(3):434–448, 2007.

Alexandre d'Aspremont, Francis Bach, and Laurent El Ghaoui. Optimal solutions for sparse principal component analysis. *J. Mach. Learn. Res.*, 9(7), 2008.

Yash Deshpande and Andrea Montanari. Finding hidden cliques of size $\sqrt{N/e}$ in nearly linear time. *Found. Comput. Math.*, 15(4):1069–1128, 2015.

Yash Deshpande and Andrea Montanari. Sparse PCA via covariance thresholding. *J. Mach. Learn. Res.*, 17(1):4913–4953, 2016.

Yash Deshpande, Andrea Montanari, and Emile Richard. Cone-constrained principal component analysis. In *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, pp. 2717–2725, 2014.

Chris Ding and Xiaofeng He. K-means clustering via principal component analysis. In *Int. Conf. Mach. Learn. (ICML)*, pp. 29, 2004.

David Foster. *Generative Deep Learning : Teaching Machines to Paint, Write, Compose and Play*. O'Reilly Media, Inc, USA, 2019.

Peter JB Hancock, A Mike Burton, and Vicki Bruce. Face processing: Human perception and principal components analysis. *Mem. Cogn.*, 24(1):26–40, 1996.

Paul Hand, Oscar Leong, and Vladislav Voroninski. Phase retrieval under a generative prior. In *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, pp. 9154–9164, 2018.

Reinhard Heckel and Paul Hand. Deep decoder: Concise image representations from untrained non-convolutional networks. In *Int. Conf. Learn. Repr. (ICLR)*, 2019.

Matthias Hein and Thomas Bühler. An inverse power method for nonlinear eigenproblems with applications in 1-spectral clustering and sparse PCA. In *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, pp. 847–855, 2010.

Rakib Hyder, Viraj Shah, Chinmay Hegde, and M Salman Asif. Alternating phase projected gradient descent with generative priors for solving compressive phase retrieval. In *Int. Conf. Acoust. Sp. Sig. Proc. (ICASSP)*, pp. 7705–7709. IEEE, 2019.

Gauri Jagatap and Chinmay Hegde. Algorithmic guarantees for inverse imaging with untrained network priors. In *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, volume 32, 2019.

A Jalal, S Karmalkar, A Dimakis, and E Price. Instance-optimal compressed sensing via posterior sampling. In *Int. Conf. Mach. Learn. (ICML)*, 2021.

Ajil Jalal, Liu Liu, Alexandros G Dimakis, and Constantine Caramanis. Robust compressed sensing using generative models. *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, 2020.

Iain M Johnstone and Arthur Yu Lu. On consistency and sparsity for principal components analysis in high dimensions. *J. Am. Stat. Assoc.*, 104(486):682–693, 2009.

Ian T. Jolliffe. *Principal component analysis*. Springer-Verlag, 1986.

Ian T Jolliffe, Nickolay T Trendafilov, and Mudassir Uddin. A modified principal component technique based on the Lasso. *J. Comput. Graph. Stat.*, 12(3):531–547, 2003.

Michel Journée, Yurii Nesterov, Peter Richtárik, and Rodolphe Sepulchre. Generalized power method for sparse principal component analysis. *J. Mach. Learn. Res.*, 11(2), 2010.

Sungkyu Jung and J Stephen Marron. PCA consistency in high dimension, low sample size context. *Ann. Stat.*, 37(6B):4104–4130, 2009.

Akshay Kamath, Sushrut Karmalkar, and Eric Price. On the power of compressed sensing with generative models. In *Int. Conf. Mach. Learn. (ICML)*, pp. 5101–5109, 2020.

Volodymyr Kuleshov. Fast algorithms for sparse principal component analysis based on Rayleigh quotient iteration. In *Int. Conf. Mach. Learn. (ICML)*, pp. 1418–1425. PMLR, 2013.

Yann LeCun, Léon Bottou, Yoshua Bengio, and Patrick Haffner. Gradient-based learning applied to document recognition. *Proc. IEEE*, 86(11):2278–2324, 1998.

Huikang Liu, Man-Chung Yue, and Anthony Man-Cho So. On the estimation performance and convergence rate of the generalized power method for phase synchronization. *SIAM J. Optim.*, 27 (4):2426–2446, 2017.

Zhaoqiang Liu and Jonathan Scarlett. The generalized Lasso with nonlinear observations and generative priors. In *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, volume 33, 2020a.

Zhaoqiang Liu and Jonathan Scarlett. Information-theoretic lower bounds for compressive sensing with generative models. *IEEE J. Sel. Areas Inf. Theory*, 1(1):292–303, 2020b.

Zhaoqiang Liu and Vincent YF Tan. The informativeness of k-means for learning mixture models. *IEEE Trans. Inf. Theory*, 65(11):7460–7479, 2019.

Zhaoqiang Liu, Selwyn Gomes, Avtansh Tiwari, and Jonathan Scarlett. Sample complexity bounds for 1-bit compressive sensing and binary stable embeddings with generative priors. In *Int. Conf. Mach. Learn. (ICML)*, 2020.

Zhaoqiang Liu, Subhroshekhar Ghosh, Jun Han, and Jonathan Scarlett. Robust 1-bit compressive sensing with partial Gaussian circulant matrices and generative priors. *https://arxiv.org/2108.03570*, 2021a.

Zhaoqiang Liu, Subhroshekhar Ghosh, and Jonathan Scarlett. Towards sample-optimal compressive phase retrieval with sparse and generative priors. *https://arxiv.org/2106.15358*, 2021b.

Zongming Ma. Sparse principal component analysis and iterative thresholding. *Ann. Stat.*, 41(2): 772–801, 2013.

Lester W Mackey. Deflation methods for sparse PCA. In *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, volume 21, pp. 1017–1024, 2008.

Baback Moghaddam, Yair Weiss, and Shai Avidan. Spectral bounds for sparse PCA: Exact and greedy algorithms. *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, 18:915, 2006.

Boaz Nadler. Finite sample approximation results for principal component analysis: A matrix perturbation approach. *Ann. Stat.*, 36(6):2791–2817, 2008.

Praneeth Netrapalli, Prateek Jain, and Sujay Sanghavi. Phase retrieval using alternating minimization. *IEEE Trans. Sig. Proc.*, 63(18):4814–4826, 2015.

Thanh V Nguyen, Gauri Jagatap, and Chinmay Hegde. Provable compressed sensing with generative priors via Langevin dynamics. *https://arxiv.org/2102.12643*, 2021.

Gregory Ongie, Ajil Jalal, Christopher A Metzler, et al. Deep learning techniques for inverse problems in imaging. *IEEE J. Sel. Areas Inf. Theory*, 1(1):39–56, 2020.

Dimitris Papailiopoulos, Alexandros Dimakis, and Stavros Korokythakis. Sparse PCA through low-rank approximations. In *Int. Conf. Mach. Learn. (ICML)*, pp. 747–755. PMLR, 2013.

Pei Peng, Shirin Jalali, and Xin Yuan. Solving inverse problems via auto-encoders. *IEEE J. Sel. Areas Inf. Theory*, 1(1):312–323, 2020.

Amelia Perry, Alexander S Wein, Afonso S Bandeira, and Ankur Moitra. Optimality and sub-optimality of PCA I: Spiked random matrix models. *Ann. Stat.*, 46(5):2416–2451, 2018.

Ankit Raj, Yuqi Li, and Yoram Bresler. GAN-based projector for faster recovery with convergence guarantees in linear inverse problems. In *IEEE/CVF Int. Conf. Comp. Vision (ICCV)*, 2019.

Viraj Shah and Chinmay Hegde. Solving linear inverse problems using GAN priors: An algorithm with provable guarantees. In *Int. Conf. Acoust. Sp. Sig. Proc. (ICASSP)*, pp. 4609–4613. IEEE, 2018.

Fahad Shamshad and Ali Ahmed. Compressed sensing-based robust phase retrieval via deep generative priors. *IEEE Sens. J.*, 21(2):2286–2298, 2020.

Haipeng Shen and Jianhua Z Huang. Sparse principal component analysis via regularized low rank matrix approximation. *J. Multivar. Anal.*, 99(6):1015–1034, 2008.

Christian D Sigg and Joachim M Buhmann. Expectation-maximization for sparse and non-negative PCA. In *Int. Conf. Mach. Learn. (ICML)*, pp. 960–967, 2008.

Dave Van Veen, Ajil Jalal, Mahdi Soltanolkotabi, et al. Compressed sensing with deep image prior and learned regularization. *https://arxiv.org/1806.06438*, 2018.

Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. *https://arxiv.org/abs/1011.3027*, 2010.

Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.

Vincent Vu and Jing Lei. Minimax rates of estimation for sparse PCA in high dimensions. In *Int. Conf. Artif. Intell. Stat. (AISTATS)*, pp. 1278–1286. PMLR, 2012.

Vincent Q Vu, Juhee Cho, Jing Lei, and Karl Rohe. Fantope projection and selection: A near-optimal convex relaxation of sparse PCA. *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, 26, 2013.

Zhaoran Wang, Huanran Lu, and Han Liu. Tighten after relax: Minimax-optimal sparse PCA in polynomial time. *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, 2014:3383, 2014.

Xiaohan Wei, Zhuoran Yang, and Zhaoran Wang. On the statistical rate of nonlinear recovery in generative models with heavy-tailed data. In *Int. Conf. Mach. Learn. (ICML)*, pp. 6697–6706, 2019.

Jay Whang, Qi Lei, and Alexandros G Dimakis. Compressed sensing with invertible generative models and dependent noise. *https://arxiv.org/2003.08089*, 2020.

Han Xiao, Kashif Rasul, and Roland Vollgraf. Fashion-MNIST: A novel image dataset for benchmarking machine learning algorithms. *https://arxiv.org/1708.07747*, 2017.

Yufei Yi and Matey Neykov. Non-sparse PCA in high dimensions via cone projected power iteration. *https://arxiv.org/2005.07587*, 2020.

Xiao-Tong Yuan and Tong Zhang. Truncated power method for sparse eigenvalue problems. *J. Mach. Learn. Res.*, 14(4), 2013.

Ron Zass and Amnon Shashua. Nonnegative sparse PCA. In *Conf. Neur. Inf. Proc. Sys. (NeurIPS)*, pp. 1561–1568. Citeseer, 2007.

Huishuai Zhang, Yi Zhou, Yingbin Liang, and Yuejie Chi. A nonconvex approach for phase retrieval: Reshaped Wirtinger flow and incremental algorithms. *J. Mach. Learn. Res.*, 18, 2017.

Hui Zou, Trevor Hastie, and Robert Tibshirani. Sparse principal component analysis. *J. Comput. Graph. Stat.*, 15(2):265–286, 2006.

APPENDIX FOR GENERATIVE PRINCIPAL COMPONENT ANALYSIS

A PROOFS FOR SPIKED MATRIX AND PHASE RETRIEVAL EXAMPLES

Before proceeding, we present the following standard definitions.

Definition 1. *A random variable X is said to be sub-Gaussian if there exists a positive constant C such that $(\mathbb{E}[|X|^p])^{1/p} \leq C\sqrt{p}$ for all $p \geq 1$. The sub-Gaussian norm of a sub-Gaussian random variable X is defined as $\|X\|_{\psi_2} := \sup_{p \geq 1} p^{-1/2} (\mathbb{E}[|X|^p])^{1/p}$.*

Definition 2. *A random variable X is said to be sub-exponential if there exists a positive constant C such that $(\mathbb{E}[|X|^p])^{\frac{1}{p}} \leq Cp$ for all $p \geq 1$. The sub-exponential norm of X is defined as $\|X\|_{\psi_1} := \sup_{p \geq 1} p^{-1} (\mathbb{E}[|X|^p])^{\frac{1}{p}}$.*

The following lemma states that the product of two sub-Gaussian random variables is sub-exponential, regardless of the dependence between them.

Lemma 3. *(Vershynin, 2018, Lemma 2.7.7) Let X and Y be sub-Gaussian random variables (not necessarily independent). Then XY is sub-exponential, and satisfies*

$$\|XY\|_{\psi_1} \leq \|X\|_{\psi_2}\|Y\|_{\psi_2}. \tag{21}$$

The following lemma provides a useful concentration inequality for the sum of independent sub-exponential random variables.

Lemma 4. *(Vershynin, 2010, Proposition 5.16) Let $X_1, \ldots, X_N$ be independent zero-mean sub-exponential random variables, and $K = \max_i \|X_i\|_{\psi_1}$. Then for every $\boldsymbol{\alpha} = [\alpha_1, \ldots, \alpha_N]^T \in \mathbb{R}^N$ and $\epsilon \geq 0$, it holds that*

$$\mathbb{P}\left(\Big|\sum_{i=1}^N \alpha_i X_i\Big| \geq \epsilon\right) \leq 2\exp\left(-c \cdot \min\Big(\frac{\epsilon^2}{K^2\|\boldsymbol{\alpha}\|_2^2}, \frac{\epsilon}{K\|\boldsymbol{\alpha}\|_\infty}\Big)\right). \tag{22}$$

The widely-used notion of an ϵ-net is introduced as follows.

Definition 3. *Let $(\mathcal{X}, d)$ be a metric space, and fix $\epsilon > 0$. A subset $S \subseteq \mathcal{X}$ is said be an ϵ-net of $\mathcal{X}$ if, for all $x \in \mathcal{X}$, there exists some $s \in S$ such that $d(s, x) \leq \epsilon$. The minimal cardinality of an ϵ-net of $\mathcal{X}$, if finite, is denoted $\mathcal{C}(\mathcal{X}, \epsilon)$ and is called the covering number of $\mathcal{X}$ (at scale ϵ).*

The following lemma provides a useful upper bound for the covering number of a unit sphere.

Lemma 5. *(Vershynin, 2010, Lemma 5.2) The unit Euclidean sphere $\mathcal{S}^{N-1}$ equipped with the Euclidean metric satisfies for every $\epsilon > 0$ that*

$$\mathcal{C}(\mathcal{S}^{N-1}, \epsilon) \leq \left(1 + \frac{2}{\epsilon}\right)^N. \tag{23}$$

The following lemma provides an upper bound for the spectral norm of a symmetric matrix.

Lemma 6. *(Vershynin, 2010, Lemma 5.4) Let $\mathbf{X}$ be a symmetric $N \times N$ matrix, and let $\mathcal{C}_\epsilon$ be an ϵ-net of $\mathcal{S}^{N-1}$ for some $\epsilon \in [0, 1/2)$. Then,*

$$\|\mathbf{X}\|_{2 \to 2} = \sup_{\mathbf{r} \in \mathcal{S}^{N-1}} |\langle \mathbf{X}\mathbf{r}, \mathbf{r}\rangle| \leq (1 - 2\epsilon)^{-1} \sup_{\mathbf{r} \in \mathcal{C}_\epsilon} |\langle \mathbf{X}\mathbf{r}, \mathbf{r}\rangle|. \tag{24}$$

With the above auxiliary results in place, we provide the relevant proofs for our two example introduced in Section 3.

A.1 PROOF FOR THE SPIKED COVARIANCE MODEL (EXAMPLE 1)

For $r = 1$, we have $\mathbf{x}_i = \sqrt{\beta}u_i\mathbf{s} + \mathbf{z}_i$ and a direct calculation gives $\mathbb{E}[\mathbf{V}] = \bar{\mathbf{V}} = \beta\mathbf{s}\mathbf{s}^T$. For any $\mathbf{s}_1 \in S_1$, we have that $\mathbf{s}_1^T\mathbf{x}_i = \sqrt{\beta}u_i\mathbf{s}^T\mathbf{s}_1 + \mathbf{z}_i^T\mathbf{s}_1$ is sub-Gaussian, with the sub-Gaussian norm being upper bounded by $C(\sqrt{\beta} + 1)\|\mathbf{s}_1\|_2$. Similarly, we have that for any $\mathbf{s}_2 \in S_2$, $\|\mathbf{s}_2^T\mathbf{x}_i\|_{\psi_2} \leq$

$C(\sqrt{\beta}+1)\|\mathbf{s}_2\|_2$. Applying Lemma 3, we deduce that $(\mathbf{s}_1^T\mathbf{x}_i)(\mathbf{s}_2^T\mathbf{x}_i)$ is sub-exponential, with the sub-exponential norm being upper bounded by $C^2(\sqrt{\beta}+1)^2\|\mathbf{s}_1\|_2\|\mathbf{s}_2\|_2$. In addition, we have

$$\mathbf{s}_1^T\mathbf{E}\mathbf{s}_2 = \mathbf{s}_1^T(\mathbf{V}-\bar{\mathbf{V}})\mathbf{s}_2 \tag{25}$$

$$= \frac{1}{m}\sum_{i=1}^{m}\left((\mathbf{x}_i^T\mathbf{s}_1)(\mathbf{x}_i^T\mathbf{s}_2) - \left((\mathbf{s}_1^T\mathbf{s}_2) + \beta(\mathbf{s}^T\mathbf{s}_1)(\mathbf{s}^T\mathbf{s}_2)\right)\right), \tag{26}$$

where $\mathbb{E}[(\mathbf{x}_i^T\mathbf{s}_1)(\mathbf{x}_i^T\mathbf{s}_2)] = (\mathbf{s}_1^T\mathbf{s}_2) + \beta(\mathbf{s}^T\mathbf{s}_1)(\mathbf{s}^T\mathbf{s}_2)$. Then, from Lemma 4, we obtain that for any $t > 0$ satisfying $m = \Omega(t)$, with probability $1 - e^{-\Omega(t)}$,

$$\left|\frac{1}{m}\sum_{i=1}^{m}\left((\mathbf{x}_i^T\mathbf{s}_1)(\mathbf{x}_i^T\mathbf{s}_2) - \left((\mathbf{s}_1^T\mathbf{s}_2) + \beta(\mathbf{s}^T\mathbf{s}_1)(\mathbf{s}^T\mathbf{s}_2)\right)\right)\right| \le C(\sqrt{\beta}+1)^2\|\mathbf{s}_1\|_2\|\mathbf{s}_2\|_2 \cdot \frac{\sqrt{t}}{\sqrt{m}}. \tag{27}$$

Taking a union bound over all $\mathbf{s}_1 \in S_1$ and $\mathbf{s}_2 \in S_2$, and setting $t = \log(|S_1| \cdot |S_2|)$, we obtain with probability $1 - e^{-\Omega(\log(|S_1| \cdot |S_2|))}$ that (13) holds (with β being a fixed positive constant).

Moreover, from Lemma 4, we also obtain that for any $t > 0$ satisfying $t = \Omega(m)$, and any $\mathbf{r} \in \mathcal{S}^{n-1}$, with probability $1 - e^{-\Omega(t)}$,

$$|\mathbf{r}^T\mathbf{E}\mathbf{r}| = \left|\frac{1}{m}\sum_{i=1}^{m}\left((\mathbf{x}_i^T\mathbf{r})^2 - \left(1 + \beta(\mathbf{s}^T\mathbf{r})^2\right)\right)\right| \tag{28}$$

$$\le C(\sqrt{\beta}+1)^2 \cdot \frac{t}{m}. \tag{29}$$

From Lemma 5, there exists an $(1/4)$-net $\mathcal{C}_{\frac{1}{4}}$ of $\mathcal{S}^{n-1}$ satisfying $\log\left|\mathcal{C}_{\frac{1}{4}}\right| \le n\log 9$. Taking a union bound over all $\mathbf{r} \in \mathcal{C}_{\frac{1}{4}}$, and setting $t = Cn$, we obtain with probability $1 - e^{-\Omega(n)}$ that

$$\sup_{\mathbf{r}\in\mathcal{C}_{\frac{1}{4}}}\left|\mathbf{r}^T\mathbf{E}\mathbf{r}\right| = O\left(\frac{n}{m}\right). \tag{30}$$

Then, from Lemma 6, we have

$$\|\mathbf{E}\|_{2\to2} \le 2\sup_{\mathbf{r}\in\mathcal{C}_{\frac{1}{4}}}\left|\mathbf{r}^T\mathbf{E}\mathbf{r}\right| = O\left(\frac{n}{m}\right). \tag{31}$$

A.2 PROOF FOR PHASE RETRIEVAL (EXAMPLE 2)

Let $\mathbf{W} = \frac{1}{m}\sum_{i=1}^{m}y_i\mathbf{a}_i\mathbf{a}_i^T\mathbf{1}_{\{l<y_i<u\}}$ and $\bar{\mathbf{W}} = \beta\mathbf{s}\mathbf{s}^T + \gamma\mathbf{I}_n$. It is shown in (Liu et al., 2021b, Lemma 8) that

$$\mathbb{E}\left[\mathbf{W}\right] = \bar{\mathbf{W}}, \tag{32}$$

which implies

$$\mathbb{E}[\mathbf{V}] = \beta\mathbf{s}\mathbf{s}^T = \bar{\mathbf{V}}. \tag{33}$$

Then, for any $\mathbf{s}_1 \in S_1$ and $\mathbf{s}_2 \in S_2$, we have

$$\mathbf{s}_1^T\mathbf{E}\mathbf{s}_2 = \mathbf{s}_1^T(\mathbf{V}-\bar{\mathbf{V}})\mathbf{s}_2 = \mathbf{s}_1^T(\mathbf{W}-\bar{\mathbf{W}})\mathbf{s}_2 \tag{34}$$

$$= \frac{1}{m}\sum_{i=1}^{m}\left(y_i(\mathbf{a}_i^T\mathbf{s}_1)(\mathbf{a}_i^T\mathbf{s}_2)\mathbf{1}_{\{l<y_i<u\}} - \left(\beta(\mathbf{s}^T\mathbf{s}_1)(\mathbf{s}^T\mathbf{s}_2) + \gamma(\mathbf{s}_1^T\mathbf{s}_2)\right)\right). \tag{35}$$

We observe that $y_i(\mathbf{a}_i^T\mathbf{s}_1)(\mathbf{a}_i^T\mathbf{s}_2)\mathbf{1}_{\{l<y_i<u\}}$ is sub-exponential with the sub-exponential norm being upper bounded by $Cu\|\mathbf{s}_1\|_2\|\mathbf{s}_2\|_2$. In addition, from (32), we have $\mathbb{E}[y_i(\mathbf{a}_i^T\mathbf{s}_1)(\mathbf{a}_i^T\mathbf{s}_2)\mathbf{1}_{\{l<y_i<u\}}] = \beta(\mathbf{s}^T\mathbf{s}_1)(\mathbf{s}^T\mathbf{s}_2) + \gamma(\mathbf{s}_1^T\mathbf{s}_2)$. Then, from Lemma 4, we obtain that for any $t > 0$ satisfying $m = \Omega(t)$, with probability $1 - e^{-\Omega(t)}$,

$$\left|\frac{1}{m}\sum_{i=1}^{m}\left(y_i(\mathbf{a}_i^T\mathbf{s}_1)(\mathbf{a}_i^T\mathbf{s}_2)\mathbf{1}_{\{l<y_i<u\}} - (\beta(\mathbf{s}^T\mathbf{s}_1)(\mathbf{s}^T\mathbf{s}_2) + \gamma(\mathbf{s}_1^T\mathbf{s}_2))\right)\right| \le Cu\|\mathbf{s}_1\|_2\|\mathbf{s}_2\|_2 \cdot \frac{\sqrt{t}}{\sqrt{m}}. \tag{36}$$

Taking a union bound over all $\mathbf{s}_1 \in S_1$ and $\mathbf{s}_2 \in S_2$, and setting $t = \log(|S_1| \cdot |S_2|)$, we obtain that with probability $1 - e^{-\Omega(\log(|S_1| \cdot |S_2|))}$, (13) holds as desired (with u being a fixed positive constant). In addition, similarly to (31), we have with probability $1 - e^{-\Omega(n)}$ that

$$\|\mathbf{E}\|_{2\to2} = O\left(\frac{n}{m}\right). \tag{37}$$

B   PROOF OF THEOREM 1 (GUARANTEE ON THE GLOBAL OPTIMUM)

Let the singular value decomposition (SVD) of $\bar{\mathbf{V}}$ be

$$\bar{\mathbf{V}} = \bar{\mathbf{U}}\bar{\mathbf{D}}\bar{\mathbf{U}}^T, \tag{38}$$

where $\bar{\mathbf{D}} = \mathrm{Diag}([\bar{\lambda}_1, \ldots, \bar{\lambda}_n])$, and $\bar{\mathbf{U}} \in \mathbb{R}^n$ is an orthogonal matrix with the first column being $\bar{\mathbf{x}}$. For $i > 1$, let the i-th column of $\bar{\mathbf{U}}$ be $\bar{\mathbf{u}}_i$. Let M be a (δ/L)-net of $B_2^k(r)$; from Lemma 5, we know that there exists such a net with

$$\log|M| \leq k \log \frac{4Lr}{\delta}. \tag{39}$$

By the L-Lipschitz continuity of G, we have that $G(M)$ is a δ-net of $\mathrm{Range}(G) = G(B_2^k(r))$. We write

$$\hat{\mathbf{v}} = (\hat{\mathbf{v}} - \tilde{\mathbf{x}}) + \tilde{\mathbf{x}}, \tag{40}$$

where $\tilde{\mathbf{x}} \in G(M)$ satisfies $\|\hat{\mathbf{v}} - \tilde{\mathbf{x}}\|_2 \leq \delta$. Then, we have

$$\hat{\mathbf{v}}^T \mathbf{V} \hat{\mathbf{v}} = \hat{\mathbf{v}}^T (\bar{\mathbf{V}} + \mathbf{E}) \hat{\mathbf{v}} \tag{41}$$

$$= \hat{\mathbf{v}}^T \bar{\mathbf{V}} \hat{\mathbf{v}} + ((\hat{\mathbf{v}} - \tilde{\mathbf{x}}) + \tilde{\mathbf{x}})^T \mathbf{E}((\hat{\mathbf{v}} - \tilde{\mathbf{x}}) + \tilde{\mathbf{x}}) \tag{42}$$

$$= \hat{\mathbf{v}}^T \bar{\mathbf{V}} \hat{\mathbf{v}} + \tilde{\mathbf{x}}^T \mathbf{E} \tilde{\mathbf{x}} + (\hat{\mathbf{v}} - \tilde{\mathbf{x}})^T \mathbf{E} (\hat{\mathbf{v}} - \tilde{\mathbf{x}}) + 2(\hat{\mathbf{v}} - \tilde{\mathbf{x}})^T \mathbf{E} \tilde{\mathbf{x}} \tag{43}$$

$$\leq \hat{\mathbf{v}}^T \bar{\mathbf{V}} \hat{\mathbf{v}} + C\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}} + (2\delta + \delta^2)\|\mathbf{E}\|_{2\to 2} \tag{44}$$

$$\leq \hat{\mathbf{v}}^T \bar{\mathbf{V}} \hat{\mathbf{v}} + O\left(\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}} + \delta\|\mathbf{E}\|_{2\to 2}\right) \tag{45}$$

$$= \hat{\mathbf{v}}^T \left(\bar{\lambda}_1 \bar{\mathbf{x}}\bar{\mathbf{x}}^T + \sum_{i>1} \bar{\lambda}_i \bar{\mathbf{u}}_i \bar{\mathbf{u}}_i^T\right) \hat{\mathbf{v}} + O\left(\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}} + \delta\|\mathbf{E}\|_{2\to 2}\right) \tag{46}$$

$$\leq \bar{\lambda}_1 \left(\bar{\mathbf{x}}^T \hat{\mathbf{v}}\right)^2 + \bar{\lambda}_2 \sum_{i>1} \left(\bar{\mathbf{u}}_i^T \hat{\mathbf{v}}\right)^2 + O\left(\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}} + \delta\|\mathbf{E}\|_{2\to 2}\right) \tag{47}$$

$$= \bar{\lambda}_1 \left(\bar{\mathbf{x}}^T \hat{\mathbf{v}}\right)^2 + \bar{\lambda}_2 \left(1 - \left(\bar{\mathbf{x}}^T \hat{\mathbf{v}}\right)^2\right) + O\left(\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}} + \delta\|\mathbf{E}\|_{2\to 2}\right), \tag{48}$$

where (44) follows from Assumption 2 and (39), and we use $\left(\bar{\mathbf{x}}^T \hat{\mathbf{v}}\right)^2 + \sum_{i>1} \left(\bar{\mathbf{u}}_i^T \hat{\mathbf{v}}\right)^2 = 1$ in (48). In addition, from (5) and $\mathbf{x}_G \in \mathrm{Range}(G)$, we have

$$\hat{\mathbf{v}}^T \mathbf{V} \hat{\mathbf{v}} \geq \mathbf{x}_G^T \mathbf{V} \mathbf{x}_G \tag{49}$$

$$= \mathbf{x}_G^T \bar{\mathbf{V}} \mathbf{x}_G + \mathbf{x}_G^T \mathbf{E} \mathbf{x}_G \tag{50}$$

$$= ((\mathbf{x}_G - \bar{\mathbf{x}}) + \bar{\mathbf{x}})^T \bar{\mathbf{V}}((\mathbf{x}_G - \bar{\mathbf{x}}) + \bar{\mathbf{x}}) + \mathbf{x}_G^T \mathbf{E} \mathbf{x}_G \tag{51}$$

$$= \bar{\lambda}_1 + 2\bar{\lambda}_1 \bar{\mathbf{x}}^T (\mathbf{x}_G - \bar{\mathbf{x}}) + (\mathbf{x}_G - \bar{\mathbf{x}})^T \bar{\mathbf{V}}(\mathbf{x}_G - \bar{\mathbf{x}}) + \mathbf{x}_G^T \mathbf{E} \mathbf{x}_G \tag{52}$$

$$\geq \bar{\lambda}_1 - 2\bar{\lambda}_1 \|\mathbf{x}_G - \bar{\mathbf{x}}\|_2 - \left|\mathbf{x}_G^T \mathbf{E} \mathbf{x}_G\right|, \tag{53}$$

where (52) expands the brackets and uses the fact that $(\bar{\lambda}_1, \bar{\mathbf{x}})$ is an eigenvalue-eigenvector pair, and (53) uses $\|\bar{\mathbf{x}}\|_2 = 1$ and the PSD property of $\bar{\mathbf{V}}$. Since $\{\mathbf{x}_G, G(M)\}$ is also a δ-net of $\mathrm{Range}(G)$, in (53), we also have $\left|\mathbf{x}_G^T \mathbf{E} \mathbf{x}_G\right| \leq O\left(\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}}\right)$. Combining this with (48) and (53), we obtain for some absolute constants C, C' that

$$\bar{\lambda}_1 - 2\bar{\lambda}_1 \|\mathbf{x}_G - \bar{\mathbf{x}}\|_2 - C\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}} \leq \hat{\mathbf{v}}^T \mathbf{V} \hat{\mathbf{v}} \tag{54}$$

$$\leq \bar{\lambda}_1 \left(\bar{\mathbf{x}}^T \hat{\mathbf{v}}\right)^2 + \bar{\lambda}_2 \left(1 - \left(\bar{\mathbf{x}}^T \hat{\mathbf{v}}\right)^2\right) + C'\sqrt{\frac{k \log \frac{Lr}{\delta}}{m}} + C''\delta\|\mathbf{E}\|_{2\to 2}, \tag{55}$$

which simplifies to

$$\left(\bar{\lambda}_1 - \bar{\lambda}_2\right)\left(\bar{\mathbf{x}}^T\hat{\mathbf{v}}\right)^2 \geq \left(\bar{\lambda}_1 - \bar{\lambda}_2\right) - 2\bar{\lambda}_1\|\mathbf{x}_G - \bar{\mathbf{x}}\|_2 - (C + C')\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}} - C''\delta\|\mathbf{E}\|_{2\to 2}. \quad (56)$$

Hence, we obtain

$$\left(\bar{\mathbf{x}}^T\hat{\mathbf{v}}\right)^2 \geq 1 - \frac{2\bar{\lambda}_1}{\bar{\lambda}_1 - \bar{\lambda}_2} \cdot \|\mathbf{x}_G - \bar{\mathbf{x}}\|_2 - \frac{(C + C')\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}}{\bar{\lambda}_1 - \bar{\lambda}_2} - \frac{C''\delta\|\mathbf{E}\|_{2\to 2}}{\bar{\lambda}_1 - \bar{\lambda}_2}. \quad (57)$$

Without loss of generality, we assume $\bar{\mathbf{x}}^T\hat{\mathbf{v}} \geq 0$ (otherwise, we can similarly provide an upper bound for $\|\bar{\mathbf{x}} + \hat{\mathbf{v}}\|_2^2$), which yields

$$\|\bar{\mathbf{x}} - \hat{\mathbf{v}}\|_2^2 = 2(1 - \bar{\mathbf{x}}^T\hat{\mathbf{v}}) \quad (58)$$

$$\leq 2\left(1 - \left(\bar{\mathbf{x}}^T\hat{\mathbf{v}}\right)^2\right) \quad (59)$$

$$\leq \frac{4\bar{\lambda}_1}{\bar{\lambda}_1 - \bar{\lambda}_2} \cdot \|\mathbf{x}_G - \bar{\mathbf{x}}\|_2 + \frac{2(C + C')\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}}{\bar{\lambda}_1 - \bar{\lambda}_2} + \frac{2C''\delta\|\mathbf{E}\|_{2\to 2}}{\bar{\lambda}_1 - \bar{\lambda}_2} \quad (60)$$

$$\leq \frac{4\bar{\lambda}_1}{\bar{\lambda}_1 - \bar{\lambda}_2} \cdot \|\mathbf{x}_G - \bar{\mathbf{x}}\|_2 + O\left(\frac{\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}}{\bar{\lambda}_1 - \bar{\lambda}_2} + \frac{\delta n}{m(\bar{\lambda}_1 - \bar{\lambda}_2)}\right), \quad (61)$$

where (59) uses $\bar{\mathbf{x}}^T\hat{\mathbf{v}} \in [0, 1]$, (60) follows from (57), and (61) follows from the assumption $\|\mathbf{E}\|_{2\to 2} = O(n/m)$.

C PROOF OF LEMMA 2 (AUXILIARY RESULT FOR PPOWER ANALYSIS)

By the assumption $\text{Range}(G) \subseteq \mathcal{S}^{n-1}$, for any $\mathbf{x} \in \mathbb{R}^n$ and $a > 0$, we have

$$\mathcal{P}_G(\mathbf{x}) = \mathcal{P}_G(a\mathbf{x}). \quad (62)$$

Let $\bar{\eta} = 1/\bar{\lambda}_1 > 0$. Then, we have $\hat{\mathbf{s}} = \mathcal{P}_G(\mathbf{V}\mathbf{s}) = \mathcal{P}_G(\bar{\eta}\mathbf{V}\mathbf{s})$. Since $\bar{\mathbf{x}} \in \text{Range}(G)$, we have

$$\|\bar{\eta}\mathbf{V}\mathbf{s} - \hat{\mathbf{s}}\|_2 \leq \|\bar{\eta}\mathbf{V}\mathbf{s} - \bar{\mathbf{x}}\|_2. \quad (63)$$

This is equivalent to

$$\|(\bar{\eta}\mathbf{V}\mathbf{s} - \bar{\mathbf{x}}) + (\bar{\mathbf{x}} - \hat{\mathbf{s}})\|_2^2 \leq \|\bar{\eta}\mathbf{V}\mathbf{s} - \bar{\mathbf{x}}\|_2^2, \quad (64)$$

which gives

$$\|\bar{\mathbf{x}} - \hat{\mathbf{s}}\|_2^2 \leq 2\langle\bar{\eta}\mathbf{V}\mathbf{s} - \bar{\mathbf{x}}, \hat{\mathbf{s}} - \bar{\mathbf{x}}\rangle. \quad (65)$$

Note also that from $\bar{\mathbf{V}}\bar{\mathbf{x}} = \bar{\lambda}_1\bar{\mathbf{x}}$, we obtain $\bar{\mathbf{x}} = \bar{\eta}\bar{\mathbf{V}}\bar{\mathbf{x}}$, which we will use throughout the proof.

For $\delta > 0$, let M be a (δ/L)-net of $B_2^k(r)$; from Lemma 5, there exists such a net with

$$\log|M| \leq k\log\frac{4Lr}{\delta}. \quad (66)$$

By the L-Lipschitz continuity of G, we have that $G(M)$ is a δ-net of $\text{Range}(G) = G(B_2^k(r))$. We write

$$\mathbf{s} = (\mathbf{s} - \mathbf{s}_0) + \mathbf{s}_0, \quad \hat{\mathbf{s}} = (\hat{\mathbf{s}} - \tilde{\mathbf{s}}) + \tilde{\mathbf{s}}, \quad (67)$$

where $\tilde{\mathbf{s}} \in G(M)$ satisfies $\|\hat{\mathbf{v}} - \tilde{\mathbf{x}}\|_2 \leq \delta$ and $\mathbf{s}_0 \in G(M)$ satisfies $\|\mathbf{s} - \mathbf{s}_0\|_2 \leq \delta$. Then, we have

$$\langle\bar{\eta}\mathbf{V}\mathbf{s} - \bar{\mathbf{x}}, \hat{\mathbf{s}} - \bar{\mathbf{x}}\rangle = \langle\bar{\eta}\bar{\mathbf{V}}(\mathbf{s} - \bar{\mathbf{x}}), \hat{\mathbf{s}} - \bar{\mathbf{x}}\rangle + \langle\bar{\eta}\mathbf{E}\mathbf{s}, \hat{\mathbf{s}} - \bar{\mathbf{x}}\rangle, \quad (68)$$

where (68) follows from $\mathbf{V} = \bar{\mathbf{V}} + \mathbf{E}$ and $\bar{\mathbf{x}} = \bar{\eta}\bar{\mathbf{V}}\bar{\mathbf{x}}$. In the following, we control the two terms in (68) separately.

1. The term $\langle\bar{\eta}\bar{\mathbf{V}}(\mathbf{s} - \bar{\mathbf{x}}), \hat{\mathbf{s}} - \bar{\mathbf{x}}\rangle$: Let $\mathbf{s} = \alpha\bar{\mathbf{x}} + \beta\mathbf{t}$ and $\hat{\mathbf{s}} = \hat{\alpha}\bar{\mathbf{x}} + \hat{\beta}\hat{\mathbf{t}}$, where $\|\mathbf{t}\|_2 = \|\hat{\mathbf{t}}\|_2 = 1$ and $\mathbf{t}^T\bar{\mathbf{x}} = \hat{\mathbf{t}}^T\bar{\mathbf{x}} = 0$. Since $\|\mathbf{s}\|_2 = \|\hat{\mathbf{s}}\|_2 = 1$, we have $\alpha^2 + \beta^2 = \hat{\alpha}^2 + \hat{\beta}^2 = 1$. In addition, we have $\alpha = \mathbf{s}^T\bar{\mathbf{x}}$ and $\hat{\alpha} = \hat{\mathbf{s}}^T\bar{\mathbf{x}}$. Recall that in (38), we write the SVD of $\bar{\mathbf{V}}$

as $\bar{\mathbf{V}} = \bar{\mathbf{U}}\bar{\mathbf{D}}\bar{\mathbf{U}}^T$. Since $\mathbf{t}^T\bar{\mathbf{x}} = 0$, we can write $\mathbf{t}$ as $\mathbf{t} = \sum_{i>1} h_i \bar{\mathbf{u}}_i$. In addition, since $\|\mathbf{t}\|_2 = 1$, we have $\sum_{i>1} h_i^2 = 1$. Then, we obtain

$$|\langle \bar{\mathbf{V}}\mathbf{t}, \hat{\mathbf{t}} \rangle| \leq \|\bar{\mathbf{V}}\mathbf{t}\|_2 \tag{69}$$

$$= \left\| \sum_{i>1} \bar{\lambda}_i h_i \bar{\mathbf{u}}_i \right\|_2 \tag{70}$$

$$= \sqrt{\sum_{i>1} \bar{\lambda}_i^2 h_i^2} \tag{71}$$

$$\leq \sqrt{\bar{\lambda}_2^2 \sum_{i>1} h_i^2} \tag{72}$$

$$= \bar{\lambda}_2. \tag{73}$$

Therefore, we obtain

$$|\langle \bar{\eta}\bar{\mathbf{V}}(\mathbf{s} - \bar{\mathbf{x}}), \hat{\mathbf{s}} - \bar{\mathbf{x}} \rangle| = |\langle (\alpha - 1)\bar{\mathbf{x}} + \bar{\eta}\beta\bar{\mathbf{V}}\mathbf{t}, (\hat{\alpha} - 1)\bar{\mathbf{x}} + \hat{\beta}\hat{\mathbf{t}} \rangle| \tag{74}$$

$$= |(\alpha - 1)(\hat{\alpha} - 1) + \bar{\eta}\beta\hat{\beta}\langle \bar{\mathbf{V}}\mathbf{t}, \hat{\mathbf{t}} \rangle| \tag{75}$$

$$\leq (1 - \alpha)(1 - \hat{\alpha}) + \bar{\eta}|\beta\hat{\beta}|\bar{\lambda}_2 \tag{76}$$

$$= (1 - \alpha)(1 - \hat{\alpha}) + \bar{\gamma}\sqrt{1 - \alpha^2}\sqrt{1 - \hat{\alpha}^2}. \tag{77}$$

2. The term $\langle \bar{\eta}\mathbf{E}\mathbf{s}, \hat{\mathbf{s}} - \bar{\mathbf{x}} \rangle$: We have

$$|\langle \bar{\eta}\mathbf{E}\mathbf{s}, \hat{\mathbf{s}} - \bar{\mathbf{x}} \rangle| = \langle \bar{\eta}\mathbf{E}((\mathbf{s} - \mathbf{s}_0) + \mathbf{s}_0), \hat{\mathbf{s}} - \bar{\mathbf{x}} \rangle \tag{78}$$

$$= \langle \bar{\eta}\mathbf{E}(\mathbf{s} - \mathbf{s}_0), \hat{\mathbf{s}} - \bar{\mathbf{x}} \rangle + \langle \bar{\eta}\mathbf{E}\mathbf{s}_0, (\hat{\mathbf{s}} - \tilde{\mathbf{s}}) + (\tilde{\mathbf{s}} - \bar{\mathbf{x}}) \rangle \tag{79}$$

$$\leq \bar{\eta}\|\mathbf{E}\|_{2\to2}\delta\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2 + \bar{\eta}\|\mathbf{E}\|_{2\to2}\delta + O\left(\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}\right) \cdot \|\tilde{\mathbf{s}} - \bar{\mathbf{x}}\|_2 \tag{80}$$

$$\leq O\left(\delta\|\mathbf{E}\|_{2\to2}\right) + O\left(\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}\right) \cdot (\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2 + \delta), \tag{81}$$

where (80) follows from Assumption 2 and (66), and (81) follows from $\bar{\eta} = 1/\bar{\lambda}_1$, along with the fact that we assumed $\bar{\lambda}_1 = \Theta(1)$.

Note that $\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2^2 = 2(1 - \hat{\mathbf{s}}^T\bar{\mathbf{x}}) = 2(1 - \hat{\alpha})$. Combining (65), (68), (77), and (81), we obtain

$$2(1 - \hat{\alpha}) \leq 2\left((1 - \alpha)(1 - \hat{\alpha}) + \bar{\gamma}\sqrt{1 - \alpha^2}\sqrt{1 - \hat{\alpha}^2}\right)$$

$$+ O\left(\delta\|\mathbf{E}\|_{2\to2}\right) + O\left(\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}\right) \cdot (\sqrt{2(1 - \hat{\alpha})} + \delta). \tag{82}$$

Using $2(1 - \hat{\alpha}) - 2(1 - \alpha)(1 - \hat{\alpha}) = 2\alpha(1 - \hat{\alpha})$, $\sqrt{1 - \alpha^2} = \sqrt{1 - \alpha}\sqrt{1 + \alpha} \leq \sqrt{2(1 - \alpha)}$, and $\sqrt{1 - \hat{\alpha}^2} \leq \sqrt{2(1 - \hat{\alpha})}$, from (82), we obtain

$$2\alpha(1 - \hat{\alpha}) \leq 2\bar{\gamma}\sqrt{2(1 - \alpha)}\sqrt{2(1 - \hat{\alpha})} + O\left(\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}\right) \cdot (\sqrt{2(1 - \hat{\alpha})} + \delta) + O\left(\delta\|\mathbf{E}\|_{2\to2}\right). \tag{83}$$

Since $\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2^2 = 2(1 - \hat{\alpha})$ and $\|\mathbf{s} - \bar{\mathbf{x}}\|_2^2 = 2(1 - \alpha)$, this is equivalent to

$$\alpha\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2^2 \leq \left(2\bar{\gamma}\|\mathbf{s} - \bar{\mathbf{x}}\|_2 + O\left(\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}\right)\right)\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2 + O\left(\delta\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}\right) + O\left(\delta\|\mathbf{E}\|_{2\to 2}\right) \tag{84}$$

$$= \left(2\bar{\gamma}\|\mathbf{s} - \bar{\mathbf{x}}\|_2 + O\left(\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}}\right)\right)\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2 + O\left(\delta\|\mathbf{E}\|_{2\to 2}\right). \tag{85}$$

Therefore, we obtain

$$\|\hat{\mathbf{s}} - \bar{\mathbf{x}}\|_2 \leq \frac{2\bar{\gamma}\|\mathbf{s} - \bar{\mathbf{x}}\|_2}{\mathbf{s}^T\bar{\mathbf{x}}} + O\left(\frac{1}{\mathbf{s}^T\bar{\mathbf{x}}}\left(\sqrt{\frac{k\log\frac{Lr}{\delta}}{m}} + \sqrt{\mathbf{s}^T\bar{\mathbf{x}}\cdot\delta\|\mathbf{E}\|_{2\to 2}}\right)\right) \tag{86}$$

$$\leq \frac{2\bar{\gamma}\|\mathbf{s} - \bar{\mathbf{x}}\|_2}{\mathbf{s}^T\bar{\mathbf{x}}} + O\left(\frac{1}{\mathbf{s}^T\bar{\mathbf{x}}}\sqrt{\frac{k\log(nLr)}{m}}\right), \tag{87}$$

where we use $\alpha = \mathbf{s}^T\bar{\mathbf{x}}$ in (86), and we use the assumption $\|\mathbf{E}\|_{2\to 2} = O(n/m)$ and set $\delta = 1/n$ in (87).

D PROOF OF THEOREM 2 (MAIN THEOREM FOR PPOWER ANALYSIS)

From Lemma 2, we have

$$\|\mathbf{w}^{(t_0+1)} - \bar{\mathbf{x}}\|_2 \leq \mu_0\|\mathbf{w}^{(t_0)} - \bar{\mathbf{x}}\|_2 + \frac{C}{\nu}\cdot\sqrt{\frac{k\log(nLr)}{m}}. \tag{88}$$

In addition, if

$$\|\mathbf{w}^{(t_0)} - \bar{\mathbf{x}}\|_2 > \frac{C}{(1-\mu_0)\nu}\sqrt{\frac{k\log(nLr)}{m}}, \tag{89}$$

then we obtain from (88) that

$$\|\mathbf{w}^{(t_0+1)} - \bar{\mathbf{x}}\|_2 \leq \mu_0\|\mathbf{w}^{(t_0)} - \bar{\mathbf{x}}\|_2 + \frac{C}{\nu}\cdot\sqrt{\frac{k\log(nLr)}{m}}, \tag{90}$$

from which the assumed scaling $m = \Omega(k\log(nLr))$ (with a large enough implied constant) yields

$$\|\mathbf{w}^{(t_0+1)} - \bar{\mathbf{x}}\|_2 < \|\mathbf{w}^{(t_0)} - \bar{\mathbf{x}}\|_2. \tag{91}$$

Squaring both sides, expanding, and canceling equal terms, we obtain

$$\bar{\mathbf{x}}^T\mathbf{w}^{(t_0+1)} > \bar{\mathbf{x}}^T\mathbf{w}^{(t_0)}, \tag{92}$$

and by induction, we obtain the desired result.

E NUMERICAL RESULTS FOR THE FASHION-MNIST DATASET

The Fashion-MNIST dataset consists of Zalando's article images with a training set of $60,000$ examples and a test set of $10,000$ examples. The size of each image in the Fashion-MNIST dataset is also 28×28, and thus $n = 784$.

The generative model G is set to be a boundary-seeking generative adversarial network (BEGAN). The BEGAN architecture is summarized as follows.[8] The generator has latent dimension $k = 62$ and four layers. The first two are fully connected layers with the architecture $62 - 1024 - 6272$, and with ReLU activation functions. The output of the second layer, reshaped to $128 \times 7 \times 7$,

---

[8]Further details of the architecture can be found at `https://github.com/hwalsuklee/tensorflow-generative-model-collections`.

(a) $\beta = 1$ and $m = 200$ (b) $\beta = 2$ and $m = 100$

Figure 4: Examples of reconstructed Fashion-MNIST images for the spiked covariance model.

(a) Fixing $\beta = 1$ and varying m (b) Fixing $m = 300$ and varying β

Figure 5: Quantitative comparisons about the performance of Power, TPower and PPower according to Cosine Similarity over the Fashion-MNIST dataset for the spiked covariance model.

is forwarded to a deconvolution layer with kernel size 4 and stride 2. The third layer uses ReLU activation and has output size $64 \times 14 \times 14$, where 64 is the number of channels. The fourth layer is also a deconvolution layer with kernel size 4 and strides 2, and it uses ReLU activation and has output size $1 \times 28 \times 28$, where the number of channels is 1.

The BEGAN is trained with a mini-batch size of 256, a learning rate of 0.0002, and 100 epochs. Other settings are the same as those for the MNIST dataset. We perform two sets of experiments, considering the spiked covariance and phase retrieval models separately. The corresponding results are reported in Figures 4, 5, 6, and 7. From these figures, again, we can observe clear superiority of PPower to Power and TPower.

(a) $m = 200$ (b) $m = 400$

Figure 6: Examples of reconstructed images of the Fashion-MNIST dataset for phase retrieval.

Figure 7: Quantitative comparisons about the performance of `Power`, `TPower` and `PPower` according to Cosine Similarity on the Fashion-MNIST dataset for phase retrieval.