Proceedings of the 90th Annual International Meeting of the Psychometric Society, Minneapolis, United States (2025).

A Comparison of Joint and Stepwise Dynamic Cognitive Diagnostic Models

Y. Ma ⊙,*† G. Wallin ⊙,† A. Ushakova ⊙,‡ and K. Cain ⊙¶

†School of Mathematical Sciences, Lancaster, LA1 4YF, United Kingdom
‡Centre for Health Informatics, Computing, and Statistics, Lancaster Medical School, Lancaster, LA1 4YW, United Kingdom
¶Department of Psychology, Lancaster, LA1 4YF, United Kingdom
*Corresponding author. Email: y.ma24@lancaster.ac.uk

Abstract

To extend cognitive diagnostic models (CDMs) to longitudinal settings, stepwise approaches that integrate a CDM model with a latent transition model and covariates are widely used due to their flexibility. Previous research has shown that stepwise estimation can yield biased results, motivating classification-error correction as a means of improving inference over uncorrected stepwise procedures. In this study, we evaluate a unified Bayesian dynamic cognitive diagnostic model that jointly estimates measurement (item parameters, latent attribute profiles) and transition components (transition parameters) in longitudinal settings with covariates. We compare this joint approach with the bias-corrected stepwise latent transition CDM through a Monte Carlo study. Results demonstrate that joint modeling provides more accurate recovery of transition parameters, particularly under limited test length and sample size, underscoring its advantages for longitudinal diagnostic analysis and offering practical guidance for applied researchers.

Keywords: longitudinal cognitive diagnostic models, latent transition models with covariates, joint modeling, bias-corrected three-step estimation

1. Introduction

Longitudinal cognitive diagnostic models (CDMs) provide a principled framework for examining changes in individuals' latent attribute profiles over time, focusing on discrete mastery transitions de la Torre (2011), rather than continuous latent growth trajectories (e.g., second–order growth model; Zheng et al. (2021)). By modeling change in dynamic latent variables rather than relying on cross-sectional representations of ability, longitudinal CDMs offer interpretable insights into learning processes, intervention effects, and changes in attribute profiles over time. Consequently, these models have been increasingly applied in educational assessment (Chen et al., 2017; Li, 2015; S. Wang et al., 2018), and psychological measurement (Liang et al., 2024).

The evolution of longitudinal latent modeling has been strongly influenced by the development of stepwise estimation procedures, most notably the three-step approach. This framework typically involves: (1) estimating the latent measurement model (e.g., latent class analysis (Goodman, 1974), latent transition analysis (Collins & Lanza, 2010), latent profile analysis (Muthén, 2004), or Gaussian Mixture Models (Li & Harring, 2017; Nylund et al., 2007) using only observed responses; (2) assigning individuals to latent classes while accounting for classification uncertainty; and (3) examining the associations between latent classes and external variables, such as covariates or distal outcomes (Bakk et al., 2013; Clark & Muthén, 2009; Li, 2015). Naive implementations of the three-step method treat assigned class memberships as error-free in step 2, leading to biased structural parameter estimates

due to misclassification (Croon, 2002). A substantial body of methodological work addressed this limitation through bias-adjusted procedures designed to correct classification error using misclassification matrix or posterior probability (Asparouhov & Muthén, 2014; Bakk & Kuha, 2018; Bakk et al., 2013; Bolck et al., 2004; Vermunt, 2010). Because the classification error of the three-step method is introduced in its second step, alternative stepwise strategies have been proposed by eliminating this step; two-step approaches (Bakk & Kuha, 2018) retain separate estimation of the measurement model while avoiding explicit class assignment. Related developments trace back to earlier contributions (Bandeen-Roche et al., 1997; Bartolucci et al., 2014; Xue & Bandeen-Roche, 2002). In contrast, the one-step approach estimates the measurement and structural components simultaneously using full information maximum likelihood (Clogg, 1981; Dayton & Macready, 1988; Lanza et al., 2013) or by incorporating logistic regression within DINA and higher-order DINA models to examine how covariates affect attribute mastery probabilities (Ayers et al., 2013; de la Torre & Douglas, 2004a; Park & Lee, 2014). While statistically efficient, this one-step estimation can be computationally demanding, requires re-estimation of the entire model whenever structural components are modified, and blurs the distinction between measurement and structural parameters. These methodological advances highlight the importance of accounting for measurement uncertainty when examining relationships between latent classes and covariates. By estimating the measurement model independently, the latent class structure is protected from being altered by the inclusion of external variables. This separation further enables researchers to interpret the latent classes, particularly in educational contexts, to use socioeconomic and demographic variables, or other external covariates, to explain latent class membership.

In many applications, researchers seek not only to describe change, but also to explain why learners differ in their initial mastery states and subsequent transitions. Individual characteristics, social demographics, and instructional interventions are therefore commonly incorporated as covariates to examine their effects on both initial attribute mastery and learning trajectories. This motivation has led to the development of longitudinal CDMs that integrate covariates into transition and structural components, most commonly through stepwise modeling strategies based on latent transition models (Di Mari et al., 2016; Iaconangelo & de la Torre, 2016). The integration of CDMs and latent transition models is attractive in practice because of its flexibility and ease of implementation. By contrast, one-step approaches estimate measurement and regression components jointly, but prior work has largely focused on cross-sectional settings (Ayers et al., 2013; Park & Lee, 2014), leaving limited evidence for their performance in longitudinal CDMs with covariates.

To address this gap, the present study compared a joint Bayesian dynamic CDM and a bias-corrected three-step latent transition CDM with covariates, under identical measurement specifications. Simulation studies examined differences in parameter recovery and attribute classification accuracy, providing practical guidance for applied longitudinal CDMs.

2.   Methodology Framework

This study evaluates the joint dynamic CDM framework recently introduced in Y. Ma et al. (2025) for modeling transitions between attribute profiles, and compares it with the bias-corrected three-step approach proposed by Liang et al. (2023). Specifically, two estimation strategies were included in this study: 1) a joint Bayesian dynamic CDM that simultaneously estimates measurement and transition components, and 2) a bias-corrected three-step latent transition CDM with covariates. Both frameworks were examined using simulations under identical measurement specifications, including a common set of items administered at each time point, and time-invariant covariates. In the simulation study, the Q-matrix was assumed to be known and time-invariant for both estimation frameworks. To ensure interpretability and to focus the comparison on estimation strategy rather than measurement model choice, the deterministic-input, noisy-and-gate (DINA) model was adopted as the measurement model. In the following sections, we first briefly review DINA models and

latent transition models, then introduce the bias-corrected stepwise modeling framework, and finally present the joint Bayesian dynamic formulation.

2.1 A Brief Review of DINA Models

CDMs belong to the family of latent class models (LCMs) and impose a specific structure on the latent space and on the relationship between latent classes and item responses. CDMs aim to characterize the relationship between learners' latent attribute mastery profiles and their observed item responses. Let $\boldsymbol{\alpha}_{it} = (\alpha_{i1t}, \ldots, \alpha_{iKt})^{\top}$ denote the $K \times 1$ vector of binary attribute mastery profiles for learner i at time t ($t = 1, \ldots, T$), where $\alpha_{ikt} = 1$ indicates mastery of attribute k ($k = 1, \ldots, K$) and $\alpha_{ikt} = 0$ otherwise. The relationship between items and attributes is specified by the Q-matrix, where the element of Q-matrix $q_{jk} = 1$ indicates that the item j requires attribute k.

Let $\mathbf{Y}_{it} = (Y_{i1t}, \ldots, Y_{iJt})^{\top}$ denote the $J \times 1$ vector of binary item responses for learner i ($i = 1, \ldots, N$) at time t ($t = 1, \ldots, T$), where $Y_{ijt} \in \{0, 1\}$ indicates the response to item j ($j = 1, \ldots, J$), with 1 for a correct response and 0 otherwise. In CDMs, item response probabilities are expressed through a latent indicator variable η_{ijt}, which encodes whether learner i meets the attribute requirements of item j at time t. Conditional on η_{ijt}, the probability of a correct response is given by

$$P(Y_{ijt} = 1 \mid \boldsymbol{\alpha}_{it}) = (1 - s_j)^{\eta_{ijt}} g_j^{1 - \eta_{ijt}}, \tag{1}$$

where $s_j = P(Y_{ijt} = 0 \mid \eta_{ijt} = 1)$ and $g_j = P(Y_{ijt} = 1 \mid \eta_{ijt} = 0)$ are the slipping and guessing parameters for item j, respectively.

Under the deterministic-input, noisy-and-gate (DINA) model, η_{ijt} follows a conjunctive rule, reflecting the non-compensatory assumption that all required attributes must be mastered for a high probability of a correct response.

$$\eta_{ijt} = \prod_{k=1}^{K} \alpha_{ikt}^{q_{jk}},$$

so that $\eta_{ijt} = 1$ when learner i has mastered all attributes required by item j at time t.

2.2 Latent Transition Models with Covariates

To model longitudinal change in attribute mastery, we adopt a latent transition framework. The latent attribute process is assumed to follow a first-order Markov structure, such that the attribute profile α_{ikt} at time t depends on the profile at the previous time point $\alpha_{ik(t-1)}$. Covariates enter the model through the structural component, which affect both the initial attribute distribution and the transition probabilities. Let $\mathbf{Z}_{it} = (Z_{it,1}, \ldots, Z_{it,C})^{\top}$ denote the $C \times 1$ vector of covariates for learner i at time t. The initial mastery probability of attribute k ($t = 1$) is modelled as

$$\text{logit}\big(P(\alpha_{ik1} = 1 \mid \mathbf{Z}_{i0})\big) = \beta_{0k} + \sum_{c=1}^{C} \beta_{kc} Z_{i0,c}, \tag{2}$$

where β_{0k} is the intercept and β_{kc} represents the effect of covariate $\mathbf{Z}_{i0}$ on initial mastery of attribute k.

Attribute transitions from $t - 1$ to t are modelled conditionally on the previous attribute profiles and covariates. Specifically, the probabilities of acquiring and losing mastery for attribute k are

parameterized as

$$\text{logit}\big(P(\alpha_{ikt} = 1 \mid \alpha_{ik,t-1} = 0, \mathbf{Z}_{i,t-1})\big) = \gamma_{01,k,0} + \sum_{c=1}^{C} \gamma_{01,k,c} Z_{i,t-1,c}, \tag{3}$$

$$\text{logit}\big(P(\alpha_{ikt} = 0 \mid \alpha_{ik,t-1} = 1, \mathbf{Z}_{i,t-1})\big) = \gamma_{10,k,0} + \sum_{c=1}^{C} \gamma_{10,k,c} Z_{i,t-1,c}, \tag{4}$$

where $\gamma_{01,k,c}$ denotes the effects of covariates $\mathbf{Z}_{i,t-1}$ on the probability of acquiring attribute k, and $\gamma_{10,k,c}$ denotes the effects of covariates on the probability of losing mastery of attribute k.

In the present study, we assume that once an attribute is mastered, it remains mastered over time, so mastery is treated as an absorbing state. This assumption is common in educational applications of longitudinal cognitive diagnosis, where learning is often viewed as cumulative and mastered attributes are unlikely to be unlearned within a short period (Chen et al., 2017; Yigit & Douglas, 2021). It is particularly reasonable in domains such as early reading development, where foundational skills (e.g., word decoding and vocabulary) serve as prerequisites for advanced skills such as comprehension (Hoover & Gough, 1990), and in spatial rotation tasks, where skill acquisition is typically progressive and stable over repeated practice (Chen et al., 2017). It is also a practically useful structural restriction because it reduces the number of possible learning trajectories and simplifies the transition states (Chen & Culpepper, 2020). Importantly, both the stepwise and joint models can still be estimated without imposing the monotonicity assumption. Therefore, this assumption should not be interpreted as a requirement for identifiability; rather, it is a substantively motivated modeling choice that has been widely adopted in the literature (Liu et al., 2023). This assumption reflects the cumulative nature of skill acquisition and helps distinguish learning progression from measurement error. Accordingly, attribute transitions are restricted to be monotonic in this study, such that, for each attribute k,

$$P(\alpha_{ikt} = 0 \mid \alpha_{ik,t-1} = 1, \mathbf{Z}_{i,t-1}) = 0.$$

2.3   Bias-Corrected Stepwise Latent Transition CDMs

The bias–corrected stepwise approach separates the measurement and structural components and estimates them sequentially, while explicitly correcting for classification error induced by hard assignment of latent states. Following Di Mari et al. (2016) and Liang et al. (2023), the procedure consists of three steps, with measurement and transition specifications held identical to the joint framework presented in Section 2.4.

2.3.1   Step 1: Measurement Model Estimation at Each Time Point

At each time point, a DINA model is independently fitted to the response vector $\mathbf{Y}_{it}$ for learner i at time t under a pre-specified and time-invariant Q-matrix. Fitting the DINA model provides estimates of the item parameters and the posterior distribution over attribute profiles for each learner,

$$\widehat{p}_{it}(\boldsymbol{\alpha}) = P(\boldsymbol{\alpha}_{it} = \boldsymbol{\alpha} \mid \mathbf{Y}_{it}, \widehat{s}, \widehat{g}, \mathbf{Q}), \qquad \boldsymbol{\alpha} \in \{0, 1\}^K.$$

2.3.2   Step 2: Latent State Assignment and Classification Error Estimation

To facilitate subsequent modeling of transitions and covariate effects, learners are assigned to discrete attribute profiles based on the posterior distributions obtained in Step 1. Specifically, a maximum a posteriori (MAP) assignment rule is applied such that the estimated attribute profile for learner i at time t is defined as

$$\widehat{\boldsymbol{\alpha}}_{it} = \arg \max_{\boldsymbol{\alpha} \in \{0,1\}^K} \widehat{p}_{it}(\boldsymbol{\alpha}).$$

The resulting assigned $\widehat{\boldsymbol{\alpha}}_{it}$ are treated as observed indicators of attribute profiles and are used as inputs to the structural model. However, this hard assignment ignores posterior uncertainty and introduces misclassification error. To account for this error, classification error probabilities (CEPs) are introduced. Let $\mathbf{W}_i = (W_{i1}, \ldots, W_{iT})^\top$ denote the assigned attribute profiles across time. The CEP matrix ($\mathbf{M}_t$) summarizes the misclassification induced by the stepwise assignment, linking the unobserved true attribute profiles L_{it} to the observed assigned attribute profiles W_{it} at each time point t. The $\mathbf{M}_t$ is defined as

$$\mathbf{M}_t(r, s) = P(W_{it} = r \mid L_{it} = s), \qquad r, s \in \mathcal{S}, \mathcal{S} = \{0, 1\}^K,$$

where $\mathcal{S}$ denotes the set of all possible attribute profiles, with $|\mathcal{S}| = 2^K$. For example, when $K = 2$, $\mathcal{S} = \{(0, 0), (0, 1), (1, 0), (1, 1)\}$. Here, $L_{it} = s$ indicates that the value of the true attribute vector $\boldsymbol{\alpha}_{it}$ for learner i at time t is s, while $W_{it} = r$ denotes the attribute profile r assigned by Step 1. Hence, the entry $M_t(r, s)$ represents the probability that a learner with true attribute profile s is assigned to profile r at Step 1. In practice, $\mathbf{M}_t$ is estimated by averaging posterior classification probabilities obtained in Step 1 over individuals,

$$\widehat{\mathbf{M}}_t(r, s) = \frac{\sum_{i=1}^{N} P(L_{it} = s \mid \mathbf{Y}_{it}) \, \mathbf{I}(W_{it} = r)}{\sum_{i=1}^{N} P(L_{it} = s \mid \mathbf{Y}_{it})}.$$

2.3.3   Step 3: Structural Model Estimation with CEP Correction

In the final step, the latent transition model with covariates is estimated while correcting for misclassification in attribute profiles. The structural model specifies the initial attribute distribution $P(L_{i1} \mid \mathbf{Z}_{i0})$ and the transition probabilities $P(L_{it} \mid L_{i,t-1}, \mathbf{Z}_{i,t-1})$, allowing covariates to influence both initial mastery and transitions. Because the true latent states L_{it} are unobserved, estimation is carried out by marginalizing over all possible latent state sequences while conditioning on the observed assignments W_{it}. The likelihood contribution for individual i is given by

$$P(\mathbf{W}_i \mid \mathbf{Z}_i) = \sum_{L_{i1}} \cdots \sum_{L_{iT}} P(L_{i1} \mid \mathbf{Z}_{i0}) \prod_{t=2}^{T} P(L_{it} \mid L_{i,t-1}, \mathbf{Z}_{i,t-1}) \prod_{t=1}^{T} \mathbf{M}_t(W_{it}, L_{it}). \tag{5}$$

This formulation explicitly incorporates classification error by weighting each possible attribute profile according to the CEP matrix estimated in Step 2. The structural parameters (initial mastery probability and transition probabilities) are estimated by maximizing the likelihood in Equation (5).

2.4   A Joint Bayesian Dynamic CDM

In contrast, the joint approach retains the same separation between measurement and structural components at the model specification level, but jointly estimates all components within a joint Bayesian estimation framework. Specifically, latent attribute profiles ($\boldsymbol{\alpha}_{it}$), item parameters (g_j, s_j), initial attribute mastery parameters (β_{0k}, β_{kc}), and transition parameters ($\gamma_{01,k,0}, \gamma_{01,k,c}$) are estimated simultaneously using Markov chain Monte Carlo (MCMC) sampling.

2.4.1   Prior Specifications

Weakly informative priors are assigned to all model parameters to ensure that posterior inference is primarily driven by the likelihood while allowing sufficient flexibility. For the guessing and slipping parameters, we used non-informative priors (flat priors) following S. Wang et al. (2018). Specifically, for item parameters, independent priors were assigned to the guessing and slipping parameters,

$$g_j \sim \text{Beta}(1, 1), \qquad s_j \sim \text{Beta}(1, 1).$$

Regression coefficients for initial attribute mastery and transition processes were assigned normal priors $N(0, 1)$:

$$\beta_{0k}, \ \beta_{kc} \sim \mathcal{N}(0, 1), \qquad \gamma_{01,k,0}, \gamma_{01,k,c} \sim \mathcal{N}(0, 1).$$

All continuous covariates were standardized to have mean zero and unit variance; under this scaling, these priors are weakly informative. For the transition parameters $\gamma_{10,k,0}$ and $\gamma_{10,k,c}$, we specified more informative priors for the intercepts centered at -2 on the logit scale to reflect the absorbing learning assumption, while retaining weakly informative $\mathcal{N}(0, 1)$ priors for the covariate effects following (Y. Ma et al., 2025). Latent attribute profiles are treated as random variables and inferred jointly with all other parameters.

2.4.2 Inference

Inference is conducted by sampling from the joint posterior distribution of all unknown quantities given the observed data. Specifically, the joint posterior distribution is given by

$$
\begin{aligned}
P(\mathbf{g}, \mathbf{s}, \boldsymbol{\beta}, \gamma_{01}, \gamma_{10}, \boldsymbol{\alpha}_1, \ldots, \boldsymbol{\alpha}_T \mid \mathbf{Y}_1, \ldots, \mathbf{Y}_T, \mathbf{Z}, \mathbf{Q}) \propto & \prod_{t=1}^{T} P(\mathbf{Y}_t \mid \mathbf{Q}, \mathbf{g}, \mathbf{s}, \boldsymbol{\alpha}_t) \cdot \\
& \prod_{t=2}^{T} P(\boldsymbol{\alpha}_t \mid \boldsymbol{\alpha}_{t-1}, \gamma_{01}, \gamma_{10}, \mathbf{Z}_{t-1}) \cdot \\
& P(\boldsymbol{\alpha}_1 \mid \boldsymbol{\beta}, \mathbf{Z}_0) \cdot P(\mathbf{g}, \mathbf{s}) \cdot P(\boldsymbol{\beta}, \gamma_{01}, \gamma_{10}),
\end{aligned}
$$

where $\mathbf{Y}_t = \{Y_{ijt}\}_{i=1,\ldots,N; j=1,\ldots,J}$, and $\boldsymbol{\alpha}_t = (\boldsymbol{\alpha}_{1t}, \ldots, \boldsymbol{\alpha}_{Nt})$. The first term, $P(\mathbf{Y}_t \mid \mathbf{Q}, \mathbf{g}, \mathbf{s}, \boldsymbol{\alpha}_t)$, represents the likelihood of observed responses at time t based on DINA model defined in Equation (1). The second term, $P(\boldsymbol{\alpha}_t \mid \boldsymbol{\alpha}_{t-1}, \gamma_{01}, \gamma_{10}, \mathbf{Z}_{t-1})$, specifies the transition probabilities between attribute profiles across time , characterized by the acquisition and loss mastery models as defined in (3) and (4). While the third term, $P(\boldsymbol{\alpha}_1 \mid \boldsymbol{\beta}, \mathbf{Z}_0)$, models the initial attribute mastery distribution in Equation (2). The remaining terms $P(\mathbf{g}, \mathbf{s})$ and $P(\boldsymbol{\beta}, \gamma_{01}, \gamma_{10})$, represent prior distributions on item parameters and regression coefficients, respectively.

2.4.3 Computation

For the stepwise framework, we followed a three-step latent transition cognitive diagnostic modeling strategy (Liang et al., 2023). Specifically, DINA measurement models were fitted separately at each time point using the R package GDINA (W.-C. Ma & de la Torre, 2020). The CEPs were then estimated and incorporated to correct for misclassification. Finally, covariate effects on initial attribute mastery and attribute transitions were estimated via logistic regression, corresponding to Equations (2), (3), and (4).

For joint modeling, posterior inference was conducted using MCMC sampling implemented in JAGS (Just Another Gibbs Sampler) via the R package RJags. Full conditional distributions were derived for all parameters, and samples were drawn iteratively to approximate the joint posterior distribution. Convergence was assessed using the potential scale reduction factor ($\widehat{R}$) and visual inspection of trace plots. The DINA implementation follows standard cognitive diagnostic modeling practice, see Zhan et al. (2019) for further examples of RJAGS-based CDMs. The code used for the simulation study is publicly available, see Appendix 1 for details.

3.   Simulation

3.1   Simulation Design

We conducted a simulation study to compare joint and stepwise estimation frameworks for longitudinal CDMs with time-invariant covariates. The design included two time points and two latent attributes, with item parameters held constant across time to ensure longitudinal measurement invariance. The Q-matrix was assumed to be known and time-invariant for both estimation frameworks. Three combinations of sample size and test length were considered, (N, J) = (200, 6), (400, 18), (600, 30), and the joint and stepwise estimation approaches were compared across these conditions. Regarding the sample size and test length conditions, these values were selected to represent small to moderate testing scenarios commonly encountered in cognitive diagnostic assessments (Culpepper, 2015; de la Torre, 2009). Moreover, the same design settings were adopted as in our previous study under the Q-matrix unknown framework (Y. Ma et al., 2025), allowing for a direct and meaningful comparison between the current results and prior findings.

For each learner i, a vector of three time-constant covariates $\mathbf{Z}_i$ was generated and incorporated into both the initial attribute mastery and transition models. To better reflect realistic dependence among predictors, these covariates were generated from a correlated distribution, such that the covariate vector $\mathbf{Z}_i = (Z_{i1}, Z_{i2}, Z_{i3})^\top$ follows a three-dimensional multivariate normal (MVN) distribution,

$$\mathbf{Z}_i \sim \mathrm{MVN}(\mathbf{0}, \Sigma),$$

where

$$\Sigma = \begin{pmatrix} 1 & \rho & \rho \\ \rho & 1 & \rho \\ \rho & \rho & 1 \end{pmatrix}.$$

In the main simulation results, we set $\rho = 0.4$ to represent a moderate level of correlation among covariates. Sensitivity analyses for the most challenging simulation condition $(N, J) = (200, 6)$ under varying correlation levels $\rho \in \{0, 0.2, 0.4, 0.6, 0.8\}$ are provided in Appendix 5 (Tables 6 and 7). Item responses at each time point were generated under the DINA model using a fixed Q-matrix that satisfies the necessary and sufficient conditions for model identifiability (Gu & Xu, 2021). The Q-matrices used in simulations are provided in the Appendix 4. Guessing and slipping parameters were held constant across time. According to W. Ma et al. (2016), item quality was set at a moderate level by independently drawing guessing and slipping parameters from a Uniform$(0.15, 0.25)$ distribution.

Model performance was evaluated in terms of parameter recovery and classification accuracy. For regression and transition parameters, mean absolute error (MAE) and root mean square error (RMSE) were computed across replications, while classification accuracy was assessed using the attribute-wise agreement rate (AAR) at each time point, a standard metric in cognitive diagnostic modeling (e.g., Chiu and Douglas, 2013; Zhan et al., 2018). Definitions of the MAE and RMSE metrics are provided in the Appendix 2. Specifically, AAR for attribute k at time point t was defined as

$$\mathrm{AAR}_{k,t} = \frac{1}{NR} \sum_{r=1}^{R} \sum_{i=1}^{N} \mathbb{I}\left(\hat{\alpha}_{ikt}^{(r)} = \alpha_{ikt}\right),$$

where $\alpha_{ikt} \in \{0, 1\}$ is the true mastery state and $\hat{\alpha}_{ikt}^{(r)}$ is its estimate in replication r.

3.2   Simulation Results

We conducted 100 replications per simulation condition. We fitted the joint Bayesian model in JAGS using two parallel chains. After 1,000 burn-in iterations, we drew 2,000 MCMC iterations per chain. Convergence was assessed using the potential scale reduction factor ($\hat{R}$; Gelman and Rubin (1992)), and the maximum $\hat{R}$ across key parameters was below 1.2, with approximately 98%

of monitored parameters having PSRF values less than 1.02. The Monte Carlo error (MCE) for coverage probability is 0.02 across all conditions following (Koehler et al., 2009), confirming 100 replications are sufficient. Using a single CPU core, runtimes across conditions for each iteration ranged from approximately 0.8 to 6 minutes. Parameter estimates were obtained by averaging the estimates from two parallel chains with randomly chosen starting values for each replication, and summarizing over 100 replications.

Table 1 reports attribute recovery accuracy (AAR) for the joint and stepwise approaches across simulation conditions. Both methods achieved high AARs, with recovery accuracy increasing as sample size and test length increased. The joint approach showed slightly higher AARs under the smallest sample condition ($N = 200, J = 6$). For the large sample condition, both approaches achieved high recovery accuracy, with AARs exceeding 0.99 across attributes and time points. Table 4 in Appendix 3 summarises the estimation accuracy of guessing and slipping parameters (g_{jt} and s_{jt}) under the joint and stepwise frameworks. Estimation accuracy improved with increasing sample size and test length, with RMSEs for g_{jt} and s_{jt} substantially lower under moderate and large conditions. Across all conditions, MAE values remained small, generally below 0.06 for the joint approach and below 0.07 for the stepwise approach, indicating accurate recovery of item parameters for both methods.

Table 1. Attribute recovery (AAR) under Joint and Stepwise.

			Joint		Stepwise	
N	J	T	AAR1	AAR2	AAR1	AAR2
200	6	1	0.899	0.869	0.868	0.846
		2	0.891	0.858	0.861	0.843
400	18	1	0.978	0.972	0.971	0.973
		2	0.974	0.968	0.970	0.969
600	30	1	0.993	0.992	0.993	0.992
		2	0.994	0.993	0.992	0.992

Note. AAR1 and AAR2 denote attribute–wise recovery accuracy for Attribute 1 and Attribute 2, respectively.

Table 2 summarizes the estimation accuracy of initial covariate effects (β) under the joint and stepwise frameworks. Across all conditions, both MAE and RMSE decreased as sample size and test length increased. For the smallest condition ($N = 200, J = 6$), notable estimation error was observed particularly for intercept terms, whereas slope estimates were more stable. Both approaches exhibited similar accuracy, although the joint framework tended to yield slightly smaller errors for parameters under smaller sample sizes.

Table 3 summarises the estimation accuracy of transition–related covariate effects on attribute acquisition ($\gamma_{01,Z,k}$) under the joint and stepwise frameworks. Across all conditions, the joint model consistently achieved lower MAE and RMSE than the stepwise approach, with differences being observed under small sample sizes ($N = 200$) and short tests ($J = 6$). Estimation accuracy improved as N and J increased for both methods; however stepwise estimates of γ_{01} remained considerably more variable even in larger samples. Sensitivity analyses reported in Appendix 5 further suggest that the joint approach shows limited sensitivity to varying covariate correlation levels ρ, with MAE and RMSE for item parameters and regression parameters typically changing by less than 0.1 relative to the independent covariate setting (ρ = 0). Because mastery is treated as an absorbing state through an informative prior on γ_{10} that makes attribute loss effectively impossible, these parameters are not of substantive interest and are not reported.

Table 2. Estimation accuracy of initial covariate effects (β) evaluated by MAE and RMSE across conditions.

N	J	Metric	Joint				Stepwise			
			$\beta_{0,1}$	$\beta_{0,2}$	$\beta_{Z,1}$	$\beta_{Z,2}$	$\beta_{0,1}$	$\beta_{0,2}$	$\beta_{Z,1}$	$\beta_{Z,2}$
200	6	MAE	0.182	0.366	0.055	0.095	0.229	0.279	0.060	0.105
		RMSE	0.197	0.225	0.067	0.105	0.275	0.317	0.067	0.119
400	18	MAE	0.094	0.085	0.055	0.042	0.099	0.093	0.071	0.055
		RMSE	0.108	0.106	0.065	0.051	0.120	0.126	0.085	0.068
600	30	MAE	0.060	0.079	0.031	0.023	0.057	0.073	0.024	0.031
		RMSE	0.071	0.092	0.039	0.028	0.082	0.094	0.034	0.040

Note. $\beta_{0,k}$ denotes intercepts and $\beta_{Z,k}$ denotes covariate effects for Attribute k.

Table 3. Estimation accuracy of transition covariate effects (γ_{01} slopes) evaluated by MAE and RMSE across conditions.

N	J	Metric	Joint		Stepwise	
			$\gamma_{01,Z,1}$	$\gamma_{01,Z,2}$	$\gamma_{01,Z,1}$	$\gamma_{01,Z,2}$
200	6	MAE	0.100	0.140	1.008	0.805
		RMSE	0.125	0.164	1.208	0.911
400	18	MAE	0.060	0.064	0.533	0.489
		RMSE	0.085	0.077	0.629	0.570
600	30	MAE	0.063	0.058	0.489	0.381
		RMSE	0.073	0.070	0.674	0.457

Note. $\gamma_{01,Z,k}$ denotes the covariate effect Z on the transition probability from non-mastery to mastery for Attribute k.

4. Discussion & Future Directions

Across simulation conditions, the joint framework demonstrated better overall estimation performance relative to the stepwise framework, particularly in the recovery of covariate effects on attribute acquisition. This advantage was most evident under small sample and short test conditions, where joint modeling provided more stable and accurate estimates than stepwise procedures. Previous research has noted that transition parameters are generally more difficult to estimate accurately than initial state parameters under the stepwise framework (Liang et al., 2023). This pattern likely reflects the cumulative impact of classification uncertainty in stepwise procedures, where transition models are built on latent state assignments obtained from earlier stages. Findings of this study indicated that the proposed joint model improved the accuracy of transition parameter recovery. Notably, the joint framework achieved these gains while remaining computationally efficient, supporting its practical applicability in longitudinal diagnostic settings.

As noted by Liang et al. (2023), a key advantage of stepwise approaches is their flexibility at the measurement stage. Response data are modelled separately at each time point, allowing measurement model specification and model selection to be conducted independently. Future work should investigate extensions of the joint framework proposed in this paper to alternative measurement models, such as deterministic input noisy "or" gate (Templin & Henson, 2006), Generalized DINA (de la Torre & Chiu, 2016), and higher-order DINA (de la Torre & Douglas, 2004b). Additionally, future work could consider the inclusion of time–varying covariates.

From a broader methodological perspective, most existing longitudinal CDMs assume a known or fixed Q-matrix. In practice, however, the Q-matrix is often unknown, and researchers have developed various methods to estimate the Q-matrix from response data (see Chen and Culpepper

(2017), Chung (2019), and D. Wang et al. (2020)). For example, Y. Ma et al. (2025) investigate how data–driven Q–matrix estimation can be integrated within unified longitudinal CDM frameworks. Extending such joint approaches to real–world educational data represents an important direction for applied research, especially for large–scale assessments such as the Programme for International Student Assessment (PISA), the Programme for the International Assessment of Adult Competencies (PIAAC), and the Trends in International Mathematics and Science Study (TIMSS).

Funding Statement   This research was supported by the Engineering and Physical Research Council, which funded the first author through a PhD studentship.

Competing Interests   None.

References

Asparouhov, T., & Muthén, B. (2014). Auxiliary Variables in Latent Class Analysis: Three-Step Approaches. *Structural Equation Modeling: A Multidisciplinary Journal*, *21*(3), 329–341. https://doi.org/10.1080/10705511.2014.915181

Ayers, E., Nugent, R., & Junker, B. P. (2013). A Comparison of Covariate Models for Cognitive Diagnosis. *Applied Psychological Measurement*, *37*(8), 617–635. https://doi.org/10.1177/0146621613488257

Bakk, Z., & Kuha, J. (2018). Two-Step Estimation of Models Between Latent Classes and External Variables. *Psychometrika*, *83*(4), 871–892. https://doi.org/10.1007/s11336-017-9592-2

Bakk, Z., Tekle, F. B., & Vermunt, J. K. (2013). Estimating the Association between Latent Class Membership and External Variables Using Bias-Adjusted Three-Step Approaches. *Sociological Methodology*, *43*(1), 272–311. https://doi.org/10.1177/0081175012470644

Bandeen-Roche, K., Miglioretti, D. L., Zeger, S. L., & Rathouz, P. J. (1997). Latent Variable Regression for Multiple Discrete Outcomes. *Journal of the American Statistical Association*, *92*(440), 1375–1386. https://doi.org/10.1080/01621459.1997.10473658

Bartolucci, F., Montanari, G. E., & Pandolfi, S. (2014). Latent Markov Models: A Review of a Class of Models for Longitudinal Data with Discrete Latent Variables. *African Stat. J*, *18*, 43–82.

Bolck, A., Croon, M., & Hagenaars, J. (2004). Estimating Relationships Between Latent Class Membership and External Variables Using Stepwise Latent Class Analysis. *Political Analysis*, *12*(1), 3–20. https://doi.org/10.1093/pan/mgh008

Chen, Y., & Culpepper, S. A. (2017). Bayesian estimation of the DINA Q-matrix. *British Journal of Mathematical and Statistical Psychology*, *70*(3), 357–380. https://doi.org/10.1111/bmsp.12093

Chen, Y., Culpepper, S. A., & Douglas, J. A. (2017). A Hidden Markov Model for Learning Trajectories in Cognitive Diagnosis With Application to Spatial Rotation Skills. *Applied Psychological Measurement*, *42*(1), 3–19. https://doi.org/10.1177/0146621617721250

Chen, Y., & Culpepper, S. A. (2020). A multivariate probit model for learning trajectories: A fine-grained evaluation of an educational intervention. *Applied Psychological Measurement*, *44*(7-8), 515–530. https://doi.org/10.1177/0146621620920928

Chiu, C.-Y., & Douglas, J. A. (2013). A nonparametric approach to cognitive diagnosis by proximity to ideal response patterns. *Journal of Classification*, *30*(2), 225–250. https://doi.org/10.1007/s00357-013-9132-9

Chung, M. (2019). A Gibbs sampling algorithm that estimates the Q-matrix for the DINA model. *Journal of Mathematical Psychology*, *93*, 102275. https://doi.org/10.1016/j.jmp.2019.07.002

Clark, S., & Muthén, B. (2009). *Relating latent class analysis results to variables not included in the analysis* (tech. rep.) (Retrieved from http://www.statmodel.com/download/relatinglca.pdf). Muthén & Muthén.

Clogg, C. C. (1981). New Developments in Latent Structure Analysis. *Evaluation Review*, *5*(3), 445–459. https://doi.org/10.1177/0193841X8100500302

Collins, L. M., & Lanza, S. T. (2010). *Latent Class and Latent Transition Analysis: With Applications in the Behavioral, Social, and Health Sciences*. John Wiley & Sons. https://doi.org/10.1002/9780470567333

Croon, M. A. (2002). Using Predicted Latent Class Membership in Subsequent Analyses. In J. A. Hagenaars & A. L. McCutcheon (Eds.), *Applied latent class analysis* (pp. 194–223). Cambridge University Press. https://doi.org/10.1017/CBO9780511499531.008

Culpepper, S. A. (2015). Bayesian estimation of the dina model. *Journal of Educational and Behavioral Statistics*. https://doi.org/10.3102/1076998615595403

Dayton, C. M., & Macready, G. B. (1988). A Latent Class Covariate Model with Applications to Criterion-Referenced Testing. *Biometrika*, *75*(1), 173–178. https://doi.org/10.1093/biomet/75.1.173

de la Torre, J. (2011). The generalized DINA model framework. *Psychometrika*, *76*(2), 179–199. https://doi.org/10.1007/s11336-011-9207-7

de la Torre, J., & Chiu, C.-Y. (2016). A General Method of Empirical Q-matrix Validation. *Psychometrika*, *81*(2), 253–273. https://doi.org/10.1007/s11336-015-9467-8

de la Torre, J., & Douglas, J. A. (2004a). Higher-order Latent Trait Models for Cognitive Diagnosis. *Psychometrika*, *69*(3), 333–353. https://doi.org/10.1007/BF02295640

de la Torre, J., & Douglas, J. A. (2004b). Higher-Order Latent Trait Models for Cognitive Diagnosis. *Psychometrika*, *69*(3), 333–353. https://doi.org/10.1007/BF02295640

de la Torre, J. (2009). Dina model and parameter estimation: A didactic. *Journal of Educational and Behavioral Statistics*, *34*(1), 115–130. https://doi.org/10.3102/1076998607309474

Di Mari, R., Oberski, D. L., & Vermunt, J. K. (2016). Bias-Adjusted Three-Step Latent Markov Modeling With Covariates. *Structural Equation Modeling: A Multidisciplinary Journal*, *23*(5), 649–666. https://doi.org/10.1080/10705511.2016.1191015

Gelman, A., & Rubin, D. B. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7(4), 457–472. https://doi.org/10.1214/ss/1177011136

Goodman, L. A. (1974). Exploratory Latent Structure Analysis Using Both Identifiable and Unidentifiable Models. *Biometrika*, *61*(2), 215–231. https://doi.org/10.1093/biomet/61.2.215

Gu, Y., & Xu, G. (2021). Sufficient and necessary conditions for the identifiability of the q-matrix. *Statistica Sinica*, *31*(1), 449–472.

Hoover, W., & Gough, P. (1990). The simple view of reading. *Reading and Writing*, *2*, 127–160. https://doi.org/10.1007/BF00401799

Iaconangelo, C., & de la Torre, J. (2016). Three-step estimation of cognitive diagnosis models with covariates [Paper presentation]. *Proceedings of the 81st International Meeting of the Psychometric Society*, 32. http://hdl.handle.net/10722/247992

Koehler, E., Brown, E., & Haneuse, S. J.-P. A. (2009). On the assessment of monte carlo error in simulation-based statistical analyses. *The American Statistician*, *63*(2), 155–162. https://doi.org/10.1198/tast.2009.0030

Lanza, S. T., Tan, X., & Bray, B. C. (2013). Latent Class Analysis with Antenatal Predictors: A One-Step Approach Using Full Information Maximum Likelihood. *Structural Equation Modeling: A Multidisciplinary Journal*, *20*(4), 692–711. https://doi.org/10.1080/10705511.2013.824781

Li, M. (2015). *Investigating Methods of Incorporating Covariates in Growth Mixture Modeling: A Simulation Study* [Doctoral dissertation]. University of Maryland, College Park.

Li, M., & Harring, J. R. (2017). Investigating Approaches to Estimating Covariate Effects in Growth Mixture Modeling: A Simulation Study. *Educational and Psychological Measurement*, *77*(5), 766–791. https://doi.org/10.1177/0013164416653789

Liang, Q., de la Torre, J., Larimer, M. E., & Mun, E.-Y. (2024). Mental Health Symptom Profiles Over Time: A Three-Step Latent Transition Cognitive Diagnosis Modeling Analysis with Covariates. In M. Stemmler, W. Wiedermann, & F. Huang (Eds.), *Dependent Data in Social Sciences Research: Forms, Issues, and Methods of Analysis* (2nd, pp. XXX–XXX). Springer. https://doi.org/10.1007/978-3-031-56318-8_22

Liang, Q., de la Torre, J., & Law, N. (2023). Latent Transition Cognitive Diagnosis Model With Covariates: A Three-Step Approach. *Journal of Educational and Behavioral Statistics*, *48*(6), 690–718. https://doi.org/10.3102/10769986231163320

Liu, Y., Culpepper, S. A., & Chen, Y. (2023). Identifiability of hidden markov models for learning trajectories in cognitive diagnosis. *Psychometrika*, *88*(2), 361–386. https://doi.org/10.1007/s11336-023-09904-x

Ma, W.-C., & de la Torre, J. (2020). *Gdina: An r package for cognitive diagnosis modeling* [R package version 2.9.3]. https://CRAN.R-project.org/package=GDINA

Ma, W., Iaconangelo, C., & de la Torre, J. (2016). Model Similarity, Model Selection, and Attribute Classification. *Applied Psychological Measurement*, *40*(3), 401–416. https://doi.org/10.1177/0146621615621717

Ma, Y., Ushakova, A., Cain, K., & Wallin, G. (2025). A statistical framework for dynamic cognitive diagnosis in digital learning environments.

Muthén, B. (2004). Latent Variable Analysis: Growth Mixture Modeling and Related Techniques for Longitudinal Data. *Handbook of Quantitative Methodology for the Social Sciences*, 345–368.

Nylund, K. L., Asparouhov, T., & Muthén, B. (2007). Deciding on the Number of Classes in Latent Class Analysis and Growth Mixture Modeling: A Monte Carlo Simulation Study. *Structural Equation Modeling: A Multidisciplinary Journal*, *14*(4), 535–569. https://doi.org/10.1080/10705510701575396

Park, J., & Lee, Y.-S. (2014). The Effect of Covariates on Attribute Mastery: A Comparison of the One-Step and Two-Step Approaches in Cognitive Diagnosis Models. *Journal of Educational and Behavioral Statistics*, *39*(4), 257–285. https://doi.org/10.3102/1076998614541701

R Core Team. (2023). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. Vienna, Austria. https://www.R-project.org/

Templin, J. L., & Henson, R. A. (2006). Measurement of Psychological Disorders Using Cognitive Diagnosis Models. *Psychological Methods*, *11*(3), 287–305. https://doi.org/10.1037/1082-989X.11.3.287

Vermunt, J. K. (2010). Latent Class Modeling with Covariates: Two-Step Estimation Revisited. *Political Analysis*, *18*(4), 450–469. https://doi.org/10.1093/pan/mpq014

Wang, D., Cai, Y., & Tu, D. (2020). Q-Matrix Estimation Methods for Cognitive Diagnosis Models: Based on Partial Known Q-Matrix. *Multivariate Behavioral Research*, *55*(1), 1–13. https://doi.org/10.1080/00273171.2020.1746901

Wang, S., Zhang, S., Douglas, J., & Culpepper, S. (2018). Using response times to assess learning progress: A joint model for responses and response times. *Measurement: Interdisciplinary Research and Perspectives*, *16*(1), 45–58. https://doi.org/10.1080/15366367.2018.1435105

Xue, Q.-L., & Bandeen-Roche, K. (2002). Combining Multiple Outcomes to Derive Score-Based Stages of Physical Frailty: The Women's Health and Aging Study. *Epidemiology*, *13*(6), 661–670. https://doi.org/10.1097/00001648-200211000-00009

Yigit, H. D., & Douglas, J. A. (2021). First-Order Learning Models With the G-DINA: Estimation With the EM Algorithm and Applications. *Journal of Educational and Behavioral Statistics*, *45*(3), 264–302. https://doi.org/10.1177/0146621621990746

Zhan, P., Jiao, H., & Man, K. (2019). Using JAGS for Bayesian Cognitive Diagnosis Modeling: A Tutorial. *Journal of Educational and Behavioral Statistics*, *44*(4), 473–503. https://doi.org/10.3102/1076998619826040

Zhan, P., Jiao, H., & Liao, D. (2018). Cognitive diagnosis modelling incorporating item response times. *British Journal of Mathematical and Statistical Psychology*, *71*(2), 262–286. https://doi.org/10.1111/bmsp.12114

Zheng, X., Yang, J. S., & Harring, J. R. (2021). Latent growth curve analysis with item response data: A methodological investigation of model parameterization, estimation, and attrition. *Structural Equation Modeling: A Multidisciplinary Journal*, *29*(2), 182–206. https://doi.org/10.1080/10705511.2021.1930543

Appendix 1.   Code Availability

The implementation of the joint estimation framework using R (version 4.5.2; R Core Team, 2023) is available at https://github.com/Yawen-Ma/IMPS_proceeding2025. The stepwise procedure builds on the R implementation of Liang et al. (2023). A minor modification was made to ensure proper normalisation of the transition probability matrix. The corrected code is also available at the author's GitHub.

Appendix 2.   Simulation Evaluation Metrics

Model performance was evaluated with respect to both parameter estimation accuracy and classification accuracy. For item parameters and regression coefficients in the initial attribute and transition models, estimation accuracy was quantified using the mean absolute error (MAE) and root mean square error (RMSE) across replications, defined as

$$\text{MAE} = \frac{1}{R} \sum_{r=1}^{R} \left| \hat{p}^{(r)} - p \right|, \quad \text{RMSE} = \sqrt{\frac{1}{R} \sum_{r=1}^{R} \left(\hat{p}^{(r)} - p \right)^2},$$

where p denotes the true parameter value, $\hat{p}^{(r)}$ is the estimate obtained in replication r, and R is the total number of replications.

Classification accuracy was evaluated at the attribute level using the attribute accuracy rate (AAR). For each attribute k at a given time point, AAR was computed as

$$\text{AAR}_{k,t} = \frac{1}{NR} \sum_{r=1}^{R} \sum_{i=1}^{N} \mathbb{I}\left(\hat{\alpha}_{ikt}^{(r)} = \alpha_{ikt} \right),$$

where N is the number of examinees, α_{ik} and $\hat{\alpha}_{ik}$ denote the true and estimated mastery status of attribute k for examinee i, respectively, and $\mathbb{I}(\cdot)$ is an indicator function equal to 1 when the condition holds and 0 otherwise.

Appendix 3.   Item Parameter Recovery
Appendix 4.   Simulation Q-Matrices

The true Q-matrices used in the simulation study for $(N, J) = (200, 6)$, $(400, 18)$, and $(600, 30)$ are provided here.

Table 4. Estimation accuracy of item parameters (g, s) evaluated by MAE and RMSE across sample size (N) and test length (J).

			Joint				Stepwise			
N	J	Metric	g_{t1}	g_{t2}	s_{t1}	s_{t2}	g_{t1}	g_{t2}	s_{t1}	s_{t2}
200	6	MAE	0.060	0.048	0.042	0.058	0.055	0.062	0.049	0.059
		RMSE	0.073	0.061	0.053	0.069	0.069	0.077	0.061	0.072
400	18	MAE	0.026	0.022	0.022	0.027	0.023	0.023	0.023	0.024
		RMSE	0.032	0.028	0.027	0.034	0.028	0.030	0.028	0.030
600	30	MAE	0.019	0.022	0.018	0.019	0.020	0.022	0.018	0.018
		RMSE	0.024	0.027	0.022	0.023	0.024	0.027	0.022	0.023

Appendix 5.   Sensitivity Analysis with Varying Levels of Covariate Correlation

Sensitivity analyses for the most challenging simulation condition, $(N, J) = (200, 6)$, under varying correlation levels $\rho \in \{0, 0.2, 0.4, 0.6, 0.8\}$ are reported in Tables 6 and 7. Panels A–D summarise the results for attribute profiles, item parameters, initial attribute parameters, and transition parameters, respectively. The case $\rho = 0$ corresponds to the independent covariate setting, where the covariate vector $\mathbf{Z}_i$ follows a multivariate normal distribution, $\mathbf{Z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_3)$. Across $\rho \in 0, 0.2, 0.4, 0.6, 0.8$, the differences in AAR were generally small, typically within about 0.01–0.02 (Table 6, Panel A). A similar pattern was observed for item parameter recovery (Table 6, Panel B), where the estimation accuracy of the guessing and slipping parameters remained stable across correlation levels, with differences in MAE and RMSE typically within approximately 0.01–0.02.

As shown in Panels C and D of Table 7, the estimation of covariate effects exhibits a similar pattern across correlation levels. For the initial covariate effects (Panel C), under the joint approach, the variation in MAE and RMSE across ρ values is relatively small, typically within about 0.03–0.10 for intercept parameters and less than 0.03 for covariate effects. A similar pattern is observed for the stepwise method, where the variation across ρ is also small, generally within about 0.05–0.10 for intercept parameters and less than 0.05 for covariate effects. For the transition covariate effects (Panel D), the stepwise method exhibits substantially larger estimation bias overall. In addition, as ρ varies, the corresponding changes in MAE and RMSE are approximately 0.2–0.6.

Table 5. True Q-matrices with $K = 2$ attributes under different test lengths: (a) $J = 6$, (b) $J = 18$, and (c) $J = 30$.

(a) $J = 6$

Item	A_1	A_2
1	1	0
2	0	1
3	1	0
4	0	1
5	1	0
6	1	1

(b) $J = 18$

Item	A_1	A_2
1	1	0
2	0	1
3	1	0
4	0	1
5	1	0
6	0	1
7	1	0
8	0	1
9	1	0
10	0	1
11	1	0
12	0	1
13	1	0
14	0	1
15	1	1
16	1	1
17	1	1
18	1	1

(c) $J = 30$

Item	A_1	A_2
1	1	0
2	0	1
3	1	0
4	0	1
5	1	0
6	0	1
7	1	0
8	0	1
9	1	0
10	0	1
11	1	0
12	0	1
13	1	0
14	0	1
15	1	0
16	0	1
17	1	0
18	0	1
19	1	0
20	0	1
21	1	0
22	0	1
23	1	0
24	0	1
25	1	1
26	1	1
27	1	1
28	1	1
29	1	1
30	1	1

(a) $J = 6$ (b) $J = 18$ (c) $J = 30$

Table 6. Sensitivity analysis results under varying covariate correlation levels.

Panel A. Attribute recovery (AAR)

ρ	N	J	T	Joint AAR$_1$	AAR$_2$	Stepwise AAR$_1$	AAR$_2$
0	200	6	1	0.890	0.860	0.863	0.842
			2	0.898	0.854	0.859	0.847
0.2	200	6	1	0.901	0.863	0.873	0.843
			2	0.894	0.849	0.879	0.853
0.4	200	6	1	0.899	0.869	0.868	0.846
			2	0.891	0.858	0.861	0.843
0.6	200	6	1	0.904	0.866	0.870	0.835
			2	0.890	0.843	0.863	0.849
0.8	200	6	1	0.898	0.877	0.874	0.847
			2	0.885	0.855	0.853	0.845

Panel B. Item parameter recovery

ρ	N	J	Metric	Joint g_{t1}	g_{t2}	s_{t1}	s_{t2}	Stepwise g_{t1}	g_{t2}	s_{t1}	s_{t2}
0	200	6	MAE	0.064	0.047	0.043	0.059	0.066	0.055	0.058	0.059
			RMSE	0.073	0.058	0.051	0.070	0.080	0.067	0.068	0.068
0.2	200	6	MAE	0.049	0.047	0.041	0.068	0.062	0.053	0.055	0.045
			RMSE	0.059	0.056	0.053	0.081	0.076	0.064	0.066	0.052
0.4	200	6	MAE	0.060	0.048	0.042	0.058	0.055	0.062	0.049	0.059
			RMSE	0.073	0.061	0.053	0.069	0.069	0.077	0.061	0.072
0.6	200	6	MAE	0.048	0.044	0.039	0.071	0.067	0.050	0.053	0.048
			RMSE	0.057	0.050	0.047	0.084	0.081	0.063	0.062	0.057
0.8	200	6	MAE	0.065	0.050	0.041	0.061	0.062	0.058	0.058	0.056
			RMSE	0.077	0.062	0.050	0.070	0.077	0.073	0.067	0.069

Table 7. Sensitivity analysis results under varying covariate correlation levels for regression parameters.

Panel C. Initial covariate effects

				Joint				Stepwise			
ρ	N	J	Metric	$\beta_{0,1}$	$\beta_{0,2}$	$\beta_{Z,1}$	$\beta_{Z,2}$	$\beta_{0,1}$	$\beta_{0,2}$	$\beta_{Z,1}$	$\beta_{Z,2}$
0	200	6	MAE	0.206	0.304	0.097	0.071	0.300	0.279	0.115	0.082
			RMSE	0.231	0.371	0.113	0.087	0.383	0.364	0.140	0.109
0.2	200	6	MAE	0.080	0.362	0.057	0.073	0.226	0.251	0.054	0.065
			RMSE	0.115	0.430	0.073	0.083	0.256	0.380	0.073	0.081
0.4	200	6	MAE	0.182	0.366	0.055	0.095	0.229	0.279	0.060	0.105
			RMSE	0.197	0.425	0.067	0.105	0.275	0.317	0.067	0.119
0.6	200	6	MAE	0.097	0.301	0.059	0.073	0.214	0.303	0.080	0.080
			RMSE	0.127	0.359	0.069	0.082	0.262	0.442	0.092	0.091
0.8	200	6	MAE	0.160	0.301	0.057	0.053	0.247	0.225	0.051	0.050
			RMSE	0.184	0.347	0.064	0.067	0.255	0.277	0.055	0.064

Panel D. Transition covariate effects

				Joint				Stepwise			
ρ	N	J	Metric	$\gamma_{01,0,1}$	$\gamma_{01,0,2}$	$\gamma_{01,Z,1}$	$\gamma_{01,Z,2}$	$\gamma_{01,0,1}$	$\gamma_{01,0,2}$	$\gamma_{01,Z,1}$	$\gamma_{01,Z,2}$
0	200	6	MAE	0.085	0.175	0.105	0.111	1.254	1.548	0.835	0.746
			RMSE	0.102	0.211	0.114	0.127	1.387	1.793	1.147	0.817
0.2	200	6	MAE	0.120	0.177	0.101	0.086	1.716	1.598	0.944	0.732
			RMSE	0.143	0.199	0.125	0.110	2.032	1.655	1.100	0.968
0.4	200	6	MAE	0.122	0.210	0.100	0.140	1.263	1.468	1.008	0.805
			RMSE	0.146	0.246	0.125	0.164	1.352	1.542	1.208	0.911
0.6	200	6	MAE	0.103	0.192	0.114	0.079	1.639	1.427	0.672	0.535
			RMSE	0.122	0.204	0.126	0.104	1.931	1.542	0.791	0.745
0.8	200	6	MAE	0.109	0.228	0.092	0.135	1.209	1.523	0.704	0.623
			RMSE	0.137	0.262	0.106	0.152	1.280	1.890	0.831	0.776