Graph Scattering beyond Wavelet Shackles

Abstract

This work develops a flexible and mathematically sound framework for the design and analysis of graph scattering networks with variable branching ratios and generic functional calculus filters. Spectrally-agnostic stability guarantees for node- and graph-level perturbations are derived; the vertex-set non-preserving case is treated by utilizing recently developed mathematical-physics based tools. Additionally, conditions for energy decay are presented and subsequently utilized to prove asymptotic expressivity of corresponding scattering transforms. New methods of graph-level feature aggregation are introduced and stability of the resulting composite scattering architectures is established. Finally, scattering transforms are extended to edge- and higher order tensorial input. Theoretical results are complemented by numerical investigations: Suitably chosen scattering networks conforming to the developed theory perform better than traditional graph-wavelet based scattering approaches in social network graph classification tasks and significantly outperform other graph-based learning approaches to regression of quantum-chemical energies on QM7.

1 Introduction

Euclidean wavelet scattering networks [23, 4] are deep convolutional architectures where output-features are generated in each layer. Employed filters are designed rather than learned and derive from a fixed (tight) wavelet frame, resulting in a tree structured network with constant branching ratio. Such networks provide state of the art methods in settings with limited data availability and serve as a mathematically tractable model of standard convolutional neural networks (CNNs). Rigorous investigations — establishing remarkable invariance- and stability properties of wavelet scattering networks — were initially carried out in [23]. The extensive mathematical analysis [40] generalized the term 'scattering network' to include tree structured networks with varying branching rations and frames of convolutional filters, thus significantly narrowing the conceptual gap to general CNNs.

With increasing interest in data on graph-structured domains, well performing networks generalizing Euclidean CNNs to this geometric setting emerged [18, 5, 9]. If efficiently implemented, such graph convolutional networks (GCNs) replace Euclidean convolutional filters by functional calculus filters; i.e. scalar functions applied to a suitably chosen graph-shift-oprator capturing the geometry of the underlying graph [18, 14, 9]. Almost immediately, proposals aimed at extending the success story of Euclidean scattering networks to the graph convolutional setting began appearing: In [50], the authors utilize dyadic graph wavelets (see e.g. [14]) based on the non-normalized graph Laplacian resulting in a norm preserving graph wavelet scattering transform. In [10], diffusion wavelets (see e.g. [8]) are used to construct a graph scattering transform enjoying spectrum-dependent stability guarantees to graph level perturbations. For scattering transforms with N layers and K distinct functional calculus

filters, the work [11] derives node-level stability bounds of $\mathcal{O}(K^{N/2})$ and conducts corresponding numerical experiments choosing diffusion wavelets, monic cubic wavelets [14] and tight Hann wavelets [36] as filters. In [12] the authors, following [8], construct so called geometric wavelets and establish the expressivity of a scattering transform based on such a frame through extensive numerical experiments. A theoretical analysis of this and a closely related wavelet based scattering transform is the main focus of [29]. Additionally, graph-wavelet based scattering transforms have been extended to the spatio-temporal domain [28], utilized to overcome the problem of oversmoothing in GCNs [26] and pruned to deal with their exponential (in network depth) increase in needed computing resources [15].

Common among all these contributions is the focus on graph wavelets, which are generically understood to derive in a scale-sampling procedure from a common wavelet generating kernel function $g : \mathbb{R} \to \mathbb{R}$ satisfying various properties [14]. Established stability or expressivity properties — especially to structural perturbations — are then generally linked to the specific choice of the wavelet kernel g and utilized graph shift operator [10, 29]. This severely limits the diversity of available filter banks in the design of scattering networks and draws into question their validity as models for more general GCNs whose filters generically do not derive from a wavelet kernel.

A primary focus of this work is to provide alleviation in this situation: After reviewing the graph signal processing setting in Section 2, we introduce a general framework for the construction of (generalized) graph scattering transforms beyond the wavelet setting in Section 3. Section 4 establishes spectrum-agnostic stability guarantees on the node signal level and for the first time also for graph-level perturbations. To handle the vertex-set non-preserving case, a new 'distance measure' for operators capturing the geometry of varying graphs is utilized. After providing conditions for energy decay (with the layers) and deriving a related asymptotic expressivity result in Section 5, we consider graph level feature aggregation and higher order inputs in Sections 6 and 7 respectively. In Section 8 we then provide numerical results indicating that general functional calculus filter based scattering is at least as expressive as standard wavelet based scattering in graph classification tasks and outperforms leading graph neural network approaches to regression of quantum chemical energies on QM7.

2   Graph Signal Processing

Taking a signal processing approach, we consider signals on graphs as opposed to graph embeddings:

Node-Signals: Given a graph (G, E), we are primarily interested in node-signals, which are functions from the node-set G to the complex numbers, modelled as elements of $\mathbb{C}^{|G|}$. We equip this space with an inner product according to $\langle f, g \rangle = \sum_{i=1}^{|G|} \overline{f_i} g_i \mu_i$ (with all vertex weights $\mu_i \geqslant 1$) and denote the resulting inner product space by $\ell^2(G)$. We could consider arbitrary inner products on $\mathbb{C}^{|G|}$ without fundamentally changing any of the results below, but will forego this in the interest of better readability.

Functional Calculus Filters: Our fundamental objects in investigating node-signals will be functional calculus filters based on a normal operator $\Delta : \ell^2(G) \to \ell^2(G)$. Prominent examples include the adjacency matrix W, the degree matrix D, un-normalized $(D - W)$ or normalized $(\mathbb{1} - D^{-\frac{1}{2}} W D^{-\frac{1}{2}})$ graph Laplacians and the lazy random walk matrix $(\mathbb{1} + W D^{-1})$ (normal for node-weights $\propto$ inverse degree [29]). Writing normalized eigenvalue-eigenvector pairs of Δ as $(\lambda_i, \phi_i)_{i=1}^{|G|}$, the filter obtained from applying $g : \mathbb{C} \to \mathbb{C}$ is given by $g(\Delta)f = \sum_{i=1}^{|G|} g(\lambda_i)\langle \phi_i, f \rangle_{\ell^2(V)} \phi_i$.

Generalized Frames: We are most interested in filters that arise from a collection of functions adequately covering the spectrum of the operator to which they are applied. To this end we call a collection $\{g_i(\cdot)\}_{i \in I}$ of functions a **generalized frame** if it satisfies the **generalized frame condition** $A \leqslant \sum_{i \in I} |g_i(c)|^2 \leqslant B$ for any c in $\mathbb{C}$ for constants $A; B > 0$. As proved in Appendix B, this condition is sufficient to guarantee that the associated operators form a frame:

Theorem 2.1. Let $\Delta : \ell^2(G) \to \ell^2(G)$ be normal. If the family $\{g_i(\cdot)\}_{i \in I}$ of bounded continuous functions satisfies $A \leqslant \sum_{i \in I} |g_i(c)|^2 \leqslant B$ for all c in the spectrum $\sigma(\Delta)$, we have $(\forall f \in \ell^2(G))$

$$A\|f\|_{\ell^2(G)}^2 \leqslant \sum_{i \in I} \|g_i(\Delta)f\|_{\ell^2(G)}^2 \leqslant B\|f\|_{\ell^2(G)}^2.$$

3 The Generalized Graph Scattering Transform

A generalized graph scattering transform is a non-linear map Φ based on a tree structured multilayer graph convolutional network with constant branching factor in each layer. For an input signal $f \in \ell^2(G)$, outputs are generated in each layer of such a scattering network, and then concatenated to form a feature vector in a feature space $\mathscr{F}$. The network is built up from three ingredients:

Connecting Operators: To allow intermediate signal representations in the 'hidden' network layers to be further processed with functional calculus filters based on varying operators, which might not all be normal for the same choice of node-weights, we allow these intermediate representations to live in varying graph signal spaces. In fact, we do not even assume that these signal spaces are based on a common vertex set. This is done to allow for modelling of recently proposed networks where input- and 'processing' graphs are decoupled (see e.g. [1, 37]), as well as architectures incorporating graph pooling [21]. Instead, we associate one signal space $\ell^2(G_n)$ to each layer n. Connecting operators are then (not necessarily linear) operators $P_n : \ell^2(G_{n-1}) \rightarrow \ell^2(G_n)$ connecting the signal spaces of subsequent layers. We assume them to be Lipschitz continuous $(\|P(f) - P(g)\|_{\ell^2(G_{n-1})} \leqslant R^+ \|f - g\|_{\ell^2(G_n)})$ and triviality preserving $(P(0) = 0)$. For our original node-signal space we also write $\ell^2(G) \equiv \ell^2(G_0)$.

Non-Linearities: To each layer, we also associate a (possibly) non-linear function $\rho_n : \mathbb{C} \rightarrow \mathbb{C}$ acting poinwise on signals in $\ell^2(G_n)$. Similar to connecting operators, we assume ρ_n preserves zero and is Lipschitz-continuous with Lipschitz constant denoted by L_n^+. This definition allows for the absolute value non-linearity, but also ReLu or, trivially, the identity function.

Operator Frames: Beyond these ingredients, the central building block of our scattering architecture is comprised of a family of functional calculus filters in each layer. That is, we assume that in each layer, the node signal space $\ell^2(G_n)$ carries a normal operator Δ_n and an associated collection of continuous functions comprised of an **output generating function** $\chi_n(\cdot)$ as well as a **filter bank** $\{g_{\gamma_n}(\cdot)\}_{\gamma_n \in \Gamma_n}$ indexed by an index set Γ_n. As the network layer n varies (and in contrast to wavelet-scattering networks) we allow the index set Γ_n as well as the collection $\{\chi_n(\cdot)\} \bigcup \{g_{\gamma_n}(\cdot)\}_{\gamma_n \in \Gamma_n}$ of functions to vary. We only demand that in each layer the functions in the filter bank together with the output generating function constitute a generalized frame with frame constants $A_n, B_n > 0$.

We refer to the collection of functions $\Omega_N := (\rho_n, \{\chi_n(\cdot)\} \bigcup \{g_{\gamma_n}(\cdot)\}_{\gamma_n \in \Gamma_n})_{n=1}^N$ as a **module sequence** and call $\mathscr{D}_N := (P_n, \Delta_n)_{n=1}^N$ our **operator collection**. The generalized scattering transform is then constructed iteratively:

Figure 1: Schematic Scattering Architecture

To our initial signal $f \in \ell^2(G)$ we first apply the connecting operator P_1, yielding a signal representation in $\ell^2(G_1)$. Subsequently, we apply the pointwise non-linearity ρ_1. Then we apply our graph filters $\{\chi_1(\Delta_1)\} \bigcup \{g_{\gamma_1}(\Delta_1)\}_{\gamma_1 \in \Gamma_1}$ to $\rho_1(P_1(f))$ yielding the output $V_1(f) := \chi_1(\Delta_1)\rho_1(P_1(f))$ as well as the intermediate hidden representations $\{U_1[\gamma_1](f) := g_{\gamma_1}(\Delta_1)\rho_1(P_1(f))\}_{\gamma_1 \in \Gamma_1}$ obtained in the first layer. Here we have introduced the **one-step scattering propagator** $U_n[\gamma_n] : \ell^2(G_{n-1}) \rightarrow \ell^2(G_n)$ mapping $f \mapsto g_{\gamma_n}(\Delta_n)\rho_n(P_n(f))$ as well as the **output generating operator** $V_n : \ell^2(G_{n-1}) \rightarrow \ell^2(G_n)$ mapping f to $\chi_n(\Delta_n)\rho_n(P_n(f))$. Upon defining the **set** $\Gamma^{N-1} := \Gamma_{N-1} \times ... \times \Gamma_1$ **of paths** of length $(N-1)$ terminating in layer $N-1$ (with Γ^0 taken to be the one-element set) and iterating the above procedure, we see that the outputs generated in the N^{th}-layer are indexed by paths Γ^{N-1} terminating in the previous layer.

120 Outputs generated in the N^{th} layer are thus given by $\{V_N \circ U[\gamma_{N-1}] \circ ... \circ U[\gamma_1](f)\}_{(\gamma_{N-1},...,\gamma_1) \in \Gamma^{N-1}}$.
121 Concatenating the features obtained in the various layers of a network with depth N, our full feature
122 vectors thus live in the feature space

$$\mathscr{F}_N = \oplus_{n=1}^N \left(\ell^2(G_n)\right)^{|\Gamma^{n-1}|}. \tag{1}$$

123 The associated canonical norm is denoted $\|\cdot\|_{\mathscr{F}_N}$. For convenience, a brief review of direct sums of
124 spaces, their associated norms and a discussion of corresponding direct sums of maps is provided in
125 Appendix A. We denote the hence constructed generalized scattering transform of length N, based
126 on a module sequence Ω_N and operator collection $\mathscr{D}_N$ by Φ_N.

4 Stability Guarantees

128 In order to produce meaningful signal representations, a small change in input signal should produce
129 a small change in the output of our generalized scattering transforms. This property is captured in the
130 result below, which is proved in Appendix C.

131 **Theorem 4.1.** With the notation of Section 3, we have for all $f, h \in \ell^2(G)$:

$$\|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N}^2 \leqslant \left(1 + \sum_{n=1}^N \max\{[B_n - 1], [B_n(L_n^+ R_n^+)^2 - 1], 0\} \prod_{k=1}^{n-1} B_k\right) \|f - h\|_{\ell^2(G)}^2$$

132 We call a scattering transform Φ_N based on a module sequence Ω_N and operator collection $\mathscr{D}_N$
133 **admissible**, if the upper frame bounds B_n and Lipschitz constants L_n^+ and R_n^+ are all smaller than or
134 equal one in each layer $n \leqslant N$. Admissible scattering transforms are 1-Lipschitz,

$$\|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N} \leqslant \|f - h\|_{\ell^2(G)},$$

135 which greatly simplifies the presentation. We shall henceforth assume admissibility, as is not a severe
136 limitation since it can always be achieved through a rescaling operation.

137 Operators capturing graph geometries might only be known approximately in real world tasks; e.g. if
138 edge weights are only known to a certain level of precision. Hence it is important that our scattering
139 representation be insensitive to small perturbations in the underlying normal operators in each layer,
140 which is captured by our next result, proved in Appendix D. Smallness here is measured in Frobenius
141 norm $\|\cdot\|_F$, which for convenience is briefly reviewed in Appendix A).

142 **Theorem 4.2.** Let Φ_N and $\widetilde{\Phi}_N$ be two admissible scattering transforms based on the same module
143 sequence Ω_N and operator sequences $\mathscr{D}_N, \widetilde{\mathscr{D}}_N$ with the same connecting operators $(P_n = \widetilde{P}_n)$ in
144 each layer. Assume that the respective normal operators satisfy $\|\Delta_n - \widetilde{\Delta}_n\|_F \leqslant \delta$ for some $\delta > 0$.
145 Further assume that the functions $\{g_{\gamma_n}\}_{\gamma_n \in \Gamma_n}$ and χ_n in each layer are Lipschitz continuous with
146 associated Lipschitz constants satisfying $L_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_n} L_{g_{\gamma_n}}^2 \leqslant D^2$ for all $n \leqslant N$ and some $D > 0$.
147 Then we have for all $f \in \ell^2(G)$

$$\|\widetilde{\Phi}_N(f) - \Phi_N(f)\|_{\mathscr{F}_N} \leqslant \sqrt{2(2^N - 1)} \cdot D \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

148 We are also interested in perturbations that change the vertex set of the graphs in our architecture.
149 This is important for example in the context of social networks, when passing from nodes representing
150 individuals to nodes representing (close knit) groups of individuals. To investigate this setting, we
151 utilize tools originally developed within the mathematical physics community [30]:

152 **Definition 4.3.** Let $\mathcal{H}$ and $\widetilde{\mathcal{H}}$ be two finite dimensional Hilbert spaces. Let Δ and $\widetilde{\Delta}$ be normal
153 operators on these spaces. Let $J : \mathcal{H} \to \widetilde{\mathcal{H}}$ and $\widetilde{J} : \widetilde{\mathcal{H}} \to \mathcal{H}$ be linear maps — called **identification**
154 **operators**. We call the two spaces δ-**quasi-unitarily-equivalent** (with $\delta \geqslant 0$) if for any $f \in \mathcal{H}$ and
155 $u \in \widetilde{\mathcal{H}}$ we have

$$\|Jf\|_{\widetilde{\mathcal{H}}} \leqslant 2\|f\|_{\mathcal{H}}, \quad \|(J - \widetilde{J}^*)f\|_{\widetilde{\mathcal{H}}} \leqslant \delta\|f\|_{\mathcal{H}},$$

$$\|f - \widetilde{J}Jf\|_{\mathcal{H}} \leqslant \delta\sqrt{\|f\|_{\mathcal{H}}^2 + \langle f, |\Delta| \, f\rangle_{\mathcal{H}}}, \quad \|u - J\widetilde{J}u\|_{\widetilde{\mathcal{H}}} \leqslant \delta\sqrt{\|u\|_{\widetilde{\mathcal{H}}}^2 + \langle u, |\widetilde{\Delta}| \, u\rangle_{\widetilde{\mathcal{H}}}}.$$

156 If, for some $w \in \mathbb{C}$ the resolvent $R := (\Delta - \omega)^{-1}$ satisfies $\|(\widetilde{R}J - JR)f\|_{\widetilde{\mathcal{H}}} \leqslant \delta\|f\|_{\mathcal{H}}$ for all $f \in \mathcal{H}$,
157 we say that Δ and $\widetilde{\Delta}$ are ω-δ-**close** with identification operator J.

Absolute value $|\Delta|$ and adjoint $\widetilde{J}^*$ of operators are briefly reviewed in Appendix A. While the above definition might seem fairly abstract at first, it is in fact a natural setting to investigate structural perturbations as Figure 2 exemplifies. In our current setting, the Hilbert spaces in Definition 4.3 are node-signal spaces $\mathcal{H} = \ell^2(G)$, $\widetilde{\mathcal{H}} = \ell^2(\widetilde{G})$ of different graphs. The notion of ω-δ-closeness is then useful, as it allows to compare filters defined on different graphs but obtained from applying the same function to the respective graph-operators:

Lemma 4.4. In the setting of Definition 4.3 let Δ and $\widetilde{\Delta}$ be ω-δ-close and satisfy $\|\Delta\|_{op}, \|\widetilde{\Delta}\|_{op} \leqslant K$ for some $K > 0$. If $g : \mathbb{C} \to \mathbb{C}$ is holomorphic on the disk $B_{K+1}(0)$ of radius $(K + 1)$, there is a constant $C_g \geqslant 0$ so that

$$\|g(\widetilde{\Delta})J - Jg(\Delta)\|_{\mathrm{op}} \leqslant C_g \cdot \delta$$

with C_g depending on g, ω and K.

An explicit characterization of C_g together with a proof of this result is presented in Appendix F. Lemma 4.4 is our main tool in establishing our next result, proved in Appendix G, which captures stability under vertex-set non-preserving perturbations:

= vertex-weight 1
= vertex-weight 2
= edge-weight $1/\delta^2$

Figure 2: Prototypical Example of δ-unitary-equivalent Node Signal Spaces with (-1)-12δ-close Laplacians. Details in Appendix E.

Theorem 4.5. Let $\Phi_N, \widetilde{\Phi}_N$ be admissible scattering transforms based on a common module sequence Ω_N and differing operator sequences $\mathscr{D}_N, \widetilde{\mathscr{D}}_N$. Assume that there are identification operators $J_n : \ell^2(G_n) \to \ell^2(\widetilde{G}_n)$, $\widetilde{J}_n : \ell^2(\widetilde{G}_n) \to \ell^2(G_n)$ ($0 \leqslant n \leqslant N$) so that the respective signal spaces are δ-unitarily equivalent, the respective normal operators $\Delta_n, \widetilde{\Delta}_n$ are ω-δ-close as well as bounded (in norm) by $K > 0$ and the connecting operators satisfy $\|\widetilde{P}_n J_{n-1} f - J_n P_n f\|_{\ell^2(\widetilde{G}_n)} \leqslant \delta\|f\|_{\ell^2(G_{n-1})}$. For the common module sequence Ω_N assume that the non-linearities satisfy $\|\rho_n(J_n f) - J_n\rho_n(f)\|_{\ell^2(\widetilde{G}_n)} \leqslant \delta\|f\|_{\ell^2(G_n)}$ and that the constants C_{χ_n} and $\{C_{g_{\gamma_n}}\}_{\gamma_n \in \Gamma_N}$ associated through Lemma 4.4 to the functions of the generalized frames in each layer satisfy $C_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_N} C_{g_{\gamma_n}}^2 \leqslant D^2$ for some $D > 0$. Denote the operator that the family $\{J_n\}_n$ of identification operators induce on $\mathscr{F}_N$ through concatenation by $\mathscr{J}_N : \mathscr{F}_N \to \widetilde{\mathscr{F}}_N$. Then we have with $K_N = \sqrt{(8^N - 1)(2D^2 + 12)/7}$ that

$$\|\widetilde{\Phi}_N(J_0 f) - \mathscr{J}_N\Phi_N(f)\|_{\widetilde{\mathscr{F}}_N} \leqslant K_N \cdot \delta \cdot \|f\|_{\ell^2(G)}, \quad \forall f \in \ell^2(G).$$

If additionally $\|\widetilde{P}_n J_{n-1} f - J_n P_n f\|_{\ell^2(\widetilde{G}_n)} = 0$ or $\|\rho_n(J_n f) - J_n\rho_n(f)\|_{\ell^2(\widetilde{G}_n)} = 0$ holds in each layer, then we have $K_N = \mathcal{O}(4^{N/2})$; if both additional equations hold, we have $K_N = \mathcal{O}(2^{N/2})$.

5 Energy Decay and Expressivity

Since stability alone does not ensure usefulness (e.g. a constant all-zero map would be both perfectly stable and perfectly useless), we are also interested in guarantees about the expressivity of scattering transforms. We analyse this question in the limit of infinite network depth and provide conditions under which $\|f\|_{\ell^2(G)} \lesssim \|\Phi_\infty(f)\|_{\mathscr{F}_\infty}$. This then implies that the only signal mapped to the zero-feature in $\mathscr{F}_\infty$ is in fact the zero signal in $\ell^2(G)$. To achieve this, we consider the energy W_N

$$W_N(f) := \sum_{(\gamma_N, \dots, \gamma_1) \in \Gamma^N} \|U[\gamma_N] \circ \dots \circ U[\gamma_1](f)\|_{\ell^2(G_N)}^2$$

stored in the network at layer N. We assume that in each layer $L_n^- |\cdot| \leqslant |\rho_n(\cdot)| \leqslant L_n^+ |\cdot|$ and $R_n^- \|f\|_{\ell^2(G_{n-1})} \leqslant \|P_n(f)\|_{\ell^2(G_n)} \leqslant R_n^+ \|f\|_{\ell^2(G_{n-1})}$ with all multiplicative constants positive. In Appendix H we then prove the following result, generalizing corresponding Euclidean results [39]:

Lemma 5.1. The energy W_N stored in layer N satisfies

$$C_N^- \|f\|_{\ell^2(G)}^2 \leqslant \|\Phi_N(f)\|_{\mathscr{F}_N} + W_N(f) \leqslant C_N^+ \|f\|_{\ell^2(G)}^2,$$

with constants $C_N^- := \prod_{i=1}^N \min\{1, A_i(L_i^- R_i^-)^2\}$ and $C_N^+ := \prod_{i=1}^N \max\{1, B_i(L_i^+ R_i^+)^2\}$.

184 Using this, our next result (proved in Appendix I) ensures decay of the energy W_N as we progress
185 through the layers.

186 **Theorem 5.2.** Let Φ_∞ be a generalized admissible graph scattering transform based on a an operator
187 sequence $\mathscr{D}_\infty = (P_n, \Delta_n)_{n=1}^\infty$ and a module sequence Ω_∞ with each $\rho_n(\cdot) \geqslant 0$. Assume in each
188 layer $n \geqslant 1$ that there is an eigenvector ψ_n of Δ_n with solely positive entries; denote the smallest
189 entry by $m_n := \min_{i \in G_n} \psi_n[i]$ and the eigenvalue corresponding to ψ_n by λ_n. Quantify the
190 'spectral-gap' opened up at this eigenvalue through neglecting the output-generating function by
191 $\eta_n := \sum_{\gamma_n \in \Gamma_n} |g_{\gamma_n}(\lambda_n)|^2$ and assume $B_n m_n \geqslant \eta_n$. We then have

$$W_N(f) \leqslant C_N^+ \cdot \left[\prod_{n=1}^N \left(1 - \left(m_n - \frac{\eta_n}{B_n} \right) \right) \right] \cdot \|f\|_{\ell^2(G)}^2.$$

192 The positivity-assumptions on the eigenvectors ψ_n can e.g. be ensured if they are chosen to lie in the
193 lowest lying eigenspace of a graph Laplacian. With this and Lemma 5.1 we immediately find:

194 **Corollary 5.3.** In the setting of Theorem 5.2, the generalized scattering transform satisfies $\Phi_\infty^{-1}(0) =$
195 $\{0\}$ if $C_N^\pm \to C^\pm$ for some positive constants $C^\pm$ and $\sum_{n=1}^N (m_n - \eta_n/B_n) \to \infty$ as $N \to \infty$.

6 Graph-Level Feature Aggregation

197 To solve tasks such as graph classification or regression over multiple graphs, we need to represent
198 graphs of varying sizes and connectivity in a common feature space. Given a scattering transform
199 Φ_N, we thus need to find a stability preserving map from the feature space $\mathscr{F}_N$ to some Euclidean
200 space that is independent of any vertex set cardinalities. Since the feature space $\mathscr{F}_N$ is a large direct
201 sum of smaller spaces (cf. (1)), we simply construct such maps on each summand independently and
202 then concatenate them into a map on $\mathscr{F}_N$.

6.1 General non-linear feature aggregation

204 Our main tool in passing to graph-level features is a non-linear map $N_p^G : \ell^2(G) \to \mathbb{R}^p$ given as

$$N_p^G(f) = \frac{1}{\sqrt{p}} (\|f\|_{\ell^1(G)}/\sqrt{\mu_G}, \|f\|_{\ell^2(G)}, \|f\|_{\ell^3(G)}, ..., \|f\|_{\ell^p(G)})^\top, \tag{2}$$

205 with $\mu_G := \sum_{i \in G} \mu_i$ and $\|f\|_{\ell^q(G)} := (\sum_{i \in G} |f_i|^q \mu_i)^{1/q}$. Our inspiration to use this map stems from
206 the standard case where all $\mu_i = 1$: For $p \geqslant |G|$, the vector $|f| = ((|f_1|, ..., |f_G|)^\top$ can then be
207 recovered from $N_p^G(f)$ up to permutation of indices [24]. Hence, employing N_p^G (with $p \geqslant |G|$) to
208 aggregate node-information into graph-level information, we lose the minimal necessary information
209 about node permutation (clearly $N_p^G(f) = N_p^G(\Pi f)$ for any permutation matrix Π) and beyond that
210 only information about the complex phase (respectively the sign in the real case) in each entry of f.

Figure 3: Graph Level Scattering

Given a scattering transform Φ_N mapping from $\ell^2(G)$ to the feature space $\mathscr{F}_N = \oplus_{n=1}^N \left(\ell^2(G_n) \right)^{|\Gamma^{n-1}|}$, we obtain a corresponding map Ψ_N mapping from $\ell^2(G)$ to $\mathscr{R}_N = \oplus_{n=1}^N (\mathbb{R}^{p_n})^{|\Gamma^{n-1}|}$ by concatenating the feature map Φ_N with the operator that the family of non-linear maps $\{N_{G_n}^{p_n}\}_{n=1}^N$ induces on $\mathscr{F}_N$ by concatenation. Similarly we obtain the map $\widetilde{\Psi}_N : \ell^2(\widetilde{G}) \to \mathscr{R}_N$ by concatenating the map $\widetilde{\Phi}_N : \ell^2(\widetilde{G}) \to \oplus_{n=1}^N \left(\ell^2(\widetilde{G}_n) \right)^{|\Gamma^{n-1}|}$ with the operator induced by the family $\{N_{\widetilde{G}_n}^{p_n}\}_{n=1}^N$. The feature space $\mathscr{R}_N$ is completely determined by path-sets Γ^N and used maximal p-norm indices p_n. It no longer depends on cardinalities of vertex sets of any graphs, allowing to compare (signals on) varying graphs with each other.

Most of the results of the previous sections are then readily transferred to the setting of graph level features (c.f. Appendix J):

Theorem 6.1. Assuming admissibility, we have $\|\Psi_N(f) - \Psi_N(h)\|_{\mathscr{R}_N} \leqslant \|f - h\|_{\ell^2(G)}$ for all $f, h \in \ell^2(G)$. With the conditions and notation of Theorem 4.2 we have

$$\|\Psi_N(f) - \widetilde{\Psi}_N(f)\|_{\mathscr{R}_N} \leqslant \sqrt{2(2^N - 1)} \cdot D \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

Additionally, in the setting of Theorem 4.5, assuming that for each $n \leqslant N$ the identification operator J_n satisfies $\left|\|J_n f\|_{\ell^1(\widetilde{G}_n)}/\sqrt{\mu_{\widetilde{G}_n}} - \|f\|_{\ell^1(G_n)}/\sqrt{\mu_{G_n}}\right|, \left|\|J_n f\|_{\ell^k(\widetilde{G}_n)} - \|f\|_{\ell^k(G_n)}\right| \leqslant \delta \cdot K \cdot \|f\|_{\ell^2(G_n)}$ for $2 \leqslant k \leqslant p_n$ implies ($\forall f \in \ell^2(G)$)

$$\|\widetilde{\Psi}_N(J_0 f) - \Psi_N(f)\|_{\mathscr{R}_N} \leqslant \sqrt{2} \cdot \sqrt{K_N^2 + K^2} \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

Furhermore, under the assumptions of Corollary 5.3 $\Psi_\infty(f) = 0$ implies $f = 0$.

6.2 Low-pass feature aggregation

While the spectrum-free aggregation scheme of the previous section is especially adapted to settings where there are no high-level spectral properties remaining constant under graph perturbations (e.g. if we choose Δ_n to be the ajacency matrix W_n), there are also settings in which graph scattering transforms amend themselves to spectral aggregation schemes: Many commonly utilized operators, such as normalized and un-normalized graph Laplacians, have a somewhat 'stable' spectral theory: Eigenvalues are always real, non-negative, the lowest-lying eigenvalue is equal zero and simple (if the graph is connected).

In this section we shall thus assume that each mentioned normal operator Δ_n ($\widetilde{\Delta}_n$) has these spectral properties. We denote the lowest lying normalized eigenvector (which is generically determined up to a complex phase) by ψ_{Δ_n} and denote by $M_{G_n}^{|\langle \cdot, \cdot \rangle|} : \ell^2(G_n) \to \mathbb{C}$ the map given by $M_{G_n}^{|\langle \cdot, \cdot \rangle|}(f) = |\langle \psi_{\Delta_n}, f \rangle_{\ell^2(G_n)}|$. The absolute value around the inner product is introduced to absorb the phase-ambiguity in the choice of ψ_{Δ_n}. Given a scattering transform Φ_N mapping from $\ell^2(G)$ to the feature space $\mathscr{F}_N$, we obtain a corresponding map $\Psi_N^{|\langle \cdot, \cdot \rangle|}$ mapping from $\ell^2(G)$ to $\mathscr{C}_N = \oplus_{n=1}^N \mathbb{C}^{|\Gamma^{n-1}|}$ by concatenating the feature map Φ_N with the operator that the family of maps $\{M_{G_n}^{|\langle \cdot, \cdot \rangle|}\}_{n=1}^N$ induces on $\mathscr{F}_N$ by concatenation. As detailed in Appendix J.2, this map inherits stability properties in complete analogy to Theorem 6.1.

7 Higher Order Scattering

Node signals capture information about nodes in isolation. However, one might also want to analyse or incorporate information about binary, ternary or even higher order relations between nodes, such as distances or angles between nodes representing atoms in a molecule. This can be formalized by considering tensorial input signals: A 2-tensor on a graph G is simply an element of $\mathbb{C}^{|G| \times |G|}$, a 3-tensor an element of $\mathbb{C}^{|G| \times |G| \times |G|}$ and so forth. For fixed k, we equip the space of k-tensors with an inner product according to $\langle f, g \rangle = \sum_{i_1, \ldots, i_k = 1}^{|G|} \overline{f_{i_1, \ldots, i_k}} g_{i_1, \ldots, i_k} \mu_{i_1, \ldots, i_k}$ and denote the resulting inner-product space by $\ell^2(G^k)$. Setting e.g. node-weights μ_i and edge weights μ_{ik} to one, the adjacency matrix W as well as normalized or un-normalized graph Laplacians constitute self-adjoint operators on $\ell^2(G^2)$, where they act by matrix multiplication. We can then follow the recipe laid out Section 3 in constructing k^{th}-order scattering transforms; all that we need are a module sequence Ω_N and an operator sequence $\mathscr{D}_N^k := (P_n^k, \Delta_n^k)_{n=1}^N$, where now $P_n^k : \ell^2(G_{n-1}^k) \to \ell^2(G_n^k)$ and $\Delta_n^k : \ell^2(G_n^k) \to \ell^2(G_n^k)$. We denote the resulting feature map by Φ_N^k and write $\mathscr{F}_N^k$ for the corresponding feature space. The node-level stability results of the preceding sections then readily translate to higher order scattering transforms; details are given in Appendix K. The map N_p^G introduced in (2) can also be adapted to aggregate higher-order tensorial features into graph level features: With $\|f\|_q := (\sum_{i_1, \ldots, i_k \in G} |f_{i_1, \ldots, i_k}|^q \mu_{i_1, \ldots, i_k})^{1/q}$ and $\mu_{G^k} := \sum_{i_1 \ldots i_k = 1}^{|G|} \mu_{i_1, \ldots, i_k}$, we define $N_p^{G^k}(f) = (\|f\|_{\ell^1(G^k)}/\sqrt{\mu_{G^k}}, \|f\|_{\ell^2(G^k)}, \|f\|_{\ell^3(G^k)}, \ldots, \|f\|_{\ell^p(G^k)})^\top/\sqrt{p}$. Given a feature map Φ_N^k with feature space $\mathscr{F}_N = \oplus_{n=1}^N (\ell^2(G_n^k))^{|\Gamma^{n-1}|}$, we obtain a corresponding map Ψ_N^k mapping from $\ell^2(G^k)$ to $\mathscr{R}_N = \oplus_{n=1}^N (\mathbb{R}^{p_n})^{|\Gamma^{n-1}|}$ by concatenating Φ_N^k with the map that the family of non-linear maps $\{N_{G_n^k}^{p_n}\}_{n=1}^N$ induces on $\mathscr{F}_N$ by concatenation. The resulting map Ψ_N^k again has stability properties analogous to the node level case, as detailed in Appendix K.

8 Experimental Results

We showcase that even upon selecting a fairly simple architecture conforming to our general theory, our generalized graph scattering networks are able to outperform both wavelet-based scattering transforms and leading graph-networks under different circumstances. To aid visual clarity when comparing results, we colour-code the best-performing method in green, the second-best performing in yellow and the third-best performing method in orange respectively.

Social Network Graph Classification: To facilitate contact between our generalized graph scattering networks, and the wider literature, we combine a network conforming to our general theory with the low pass aggregation scheme of Section 6.2 and a Euclidean support vector machine with RBF-kernel (GGSN+EK). For this hybrid architecture, classification accuracies under 10-fold cross validation on five common social network graph datasets are compared with performances of popular graph kernel approaches, leading deep learning methods as well as geometric wavelet scattering (GS-SVM) [12]. We chose a generalized scattering architecture of depth $N = 4$. As normal-operators, we utilize in each layer the un-normalized graph Laplacian $\mathcal{L} = D - W$ scaled by its largest eigenvalue ($\Delta = \mathcal{L}/\lambda_{max}(\mathcal{L})$). Filters are chosen as $\frac{1}{2}(\sin(\pi/2 \cdot \Delta), [\cos(\pi/2 \cdot \Delta) - \psi_\Delta \psi_\Delta^\top], \sin(\pi \cdot \Delta), [\cos(\pi \cdot \Delta) - \psi_\Delta \psi_\Delta^\top])$, which allows to specify the output generating function solely by demanding $\chi(0) = 1$ and $\chi(\lambda) = 0$ on all other eigenvalues of Δ. Here ψ_Δ is the normalized vector of all ones (satisfying $\Delta \psi_\Delta = 0$). Non-linearities are chosen as modulus. Appendix L provides more details and explains how this architecture conforms to our general framework. As evident from Table 1, our network consistently achieves higher accuracies than the geometric wavelet scattering transform of [12], with the performance gap becoming significant on the more complex REDDIT datasets, finally reaching a relative mean performance increase of more than 10% on REDDIT-12K. This indicates the liberating power of transcending the graph wavelet setting. While on comparatively smaller and somewhat simpler datasets there is a performance gap between our static architecture and fully trainable networks, this gap closes on more complex datasets: While P-Poinc outperforms our method on IMDB datasets for example, the roles are reversed on REDDIT datasets. On REDDIT-B our approach trails only GIN, with the difference in accuracies not being significant. On REDDIT-5K our method comes in third, with the gap to the second best method (GIN) being statistically insignificant. On REDDIT-12K we generate state of the art results.

Table 1: Classification Accuracies on Social Network Datasets

Method	Classification Accuracies [%]					
	COLLAB	**IMDB-B**	**IMDB-M**	**REDDIT-B**	**REDDIT-5K**	**REDDIT-12K**
WL [34]	77.82±1.45	71.60±5.16	N/A	78.52±2.01	50.77 ± 2.02	34.57 ± 1.32
Graphlet [35]	73.42±2.43	65.40±5.95	N/A	77.26±2.34	39.75 ± 1.36	25.98 ± 1.29
DGK [44]	73.00±0.20	66.90±0.50	44.50±0.50	78.00±0.30	41.20 ± 0.10	32.20 ± 0.10
DGCNN [48]	73.76±0.49	70.03±0.86	47.83±0.85	N/A	48.70 ± 4.54	N/A
PSCN [27]	72.60±2.15	71.00±2.29	45.23±2.84	86.30±1.58	49.10 ± 0.70	41.32 ± 0.42
P-Poinc [20]	N/A	81.86±4.26	57.31±4.27	79.78±3.21	51.71 ± 3.01	42.16 ± 3.41
S2S-N2N-PP [16]	81.75±0.80	73.80±0.70	51.19±0.50	86.50±0.80	52.28 ± 0.50	42.47 ± 0.10
GSN-e [3]	85.5 ± 1.2	77.8 ± 3.3	54.3 ± 3.3	N/A	N/A	N/A
WKPI-kC[49]	N/A	75.1 ± 1.1	49.5 ± 0.4	N/A	59.5 ± 0.6	48.4 ± 0.5
GIN [43]	80.20±1.90	75.10±5.10	52.30±2.80	92.40±2.50	57.50 ± 1.50	N/A
GS-SVM [12]	79.94±1.61	71.20±3.25	48.73±2.32	89.65±1.94	53.33 ± 1.37	45.23 ± 1.25
GGSN+EK [OURS]	80.34±1.68	73.20±3.76	49.47±2.27	91.60±1.97	56.89 ± 2.24	49.03 ± 1.58

Regression of Quantum Chemical Energies: In order to showcase the prowess of both our higher order scattering scheme and our spectrum-agnostic aggregation method of Section 6.1, we combine these building blocks into a hybrid architecture which we then apply in combination with kernel methods (2GGST + EK) to the task of atomization energy regression on QM7. This is a comparatively small dataset of 7165 molecular graphs, taken from the 970 million strong molecular database GDB-13 [2]. Each graph in QM7 represents an organic molecule, with nodes corresponding to individual atoms. Beyond the node-level information of atomic charge, there is also edge level information characterising interaction strengths between individual nodes/atoms available. This is encoded into so called Coulomb matrices (see e.g. [32] or Appendix L) of molecular graphs, which for us serve a dual purpose: On the one hand we consider a Coulomb matrix as an edge-level input signal on a given graph. On the other hand, we also treat it as an adjacency matrix from which we build up a graph Laplacian $\mathcal{L}$. Our normal operator is then chosen as $\Delta = \mathcal{L}/\lambda_{max}(\mathcal{L})$ again. Connecting operators are

Figure 4: Atomization Energy as a Function of primary Principal Components of Scattering Features

set to the identity, while non-linearities are fixed to $\rho_{n \geqslant 1}(\cdot) = |\cdot|$. Filters are chosen as $(\sin(\pi/2 \cdot \Delta), \cos(\pi/2 \cdot \Delta), \sin(\pi \cdot \Delta), \cos(\pi \cdot \Delta))$ acting through matrix multiplication. Output generating functions are set to the identity and depth is $N = 4$. Graph level features are aggregated via the map $N_5^{G^2}$ of Section 7. Generated feature vectors are combined with node node level scattering features (based on atomic charge) into composite feature vectors; plotted in Figure 4. As is visually evident, even when reduced to the low-dimensional subspace of their first three principal components, the generated scattering features are able to aptly resolve the atomization energy of the molecules. This aptitude is also reflected in Table 2, comparing our approach with leading graph-based learning methods trained with ten-fold cross validation on node and (depending on the model) edge level information.

Our method is the best performing. We significantly outperform the next best model (DTNN), producing less than half of its mean absolute error (MAE). Errors of other methods are at least one — sometimes two — orders of magnitude greater. In part, this performance discrepancy might be explained by the hightened suitability of our scattering transform for environments with somewhat limited training-data availability. Here we speculate that the additional performance gap might be explained by the fact that our graph shift operator Δ carries the same information as the Coulomb matrix (a proven molecular graph descriptor in itself [32]). Additionally, our filters being infinite series' in powers of the underlying normal operator allows for rapid dispersion of information across underlying molecular graphs, as opposed to e.g. the filters in GraphConv or SchNet, which do not incorporate such higher powers.

Table 2: Comparison of Methods

Method	MAE [kcal/mol]
AttentiveFP [42]	66.2 ± 2.8
DMPNN [46]	105.8 ± 13.2
DTNN [41]	8.2 ± 3.9
GraphConv [18]	118.9 ± 20.2
GROVER (base)[31]	72.5 ± 5.9
MPNN [13]	113.0 ± 17.2
N-GRAM[22]	125.6 ± 1.5
PAGTN (global) [6]	47.8 ± 3.0
PhysChem [47]	59.6 ± 2.3
SchNet [33]	74.2 ± 6.0
Weave [17]	59.6 ± 2.3
2GGST+EK [OURS]	3.4 ± 0.3

9   Discussion of Results in Context

Leaving behind the traditional reliance on graph wavelets, we developed a theoretically well founded framework for the design and analysis of (generalized) graph scattering networks; allowing for varying branching rations, non-linearities and filter banks. We provided spectrum independent stability guarantees, covering changes in input signals and for the first time also arbitrary normal perturbations in the underlying graph-shift-operators. After introducing a new framework to quantify vertex-set non-preserving changes in graph domains, we were able to obtain spectrum-independent stability guarantees for this setting too. We provided conditions for energy decay in scattering networks and discussed their relation to asymptotic expressivity. Then we introduced a new method of graph-level feature aggregation, discussed stability properties of the resulting graph-level scattering transform and extended scattering networks to higher order input data. Our numerical experiments showed that a simple scattering transform conforming to our framework is able to outperform the traditional graph-wavelet based approach to graph scattering in social network graph classification tasks. On complex datasets our method is also competitive with current fully trainable methods, ouperforming all competitors on REDDIT-12K. Additionally, higher order graph scattering transforms significantly outperform current leading graph-based learning methods in predicting atomization energies on QM7. A reasonable critique of scattering networks as tractable models for general graph convolutional networks is their inability to emulate non-tree-structured network topologies. While transcending the wavelet setting has arguably diminished the conceptual gap between the two architectures, this structural difference persists in our work. Additionally we note that despite a provided promising example, it is not yet clear whether the newly introduced graph-perturbation framework can aptly provide stability guarantees to all reasonable coarse-graining procedures. Exploring this question is ongoing work. Finally, we caution against an over-interpretation of established mathematical guarantees: Such guarantees certainly do not negate biases that may be inherent to utilized datasets.

References

[1] Uri Alon and Eran Yahav. On the bottleneck of graph neural networks and its practical implications. In *International Conference on Learning Representations*, 2021.

[2] L. C. Blum and J.-L. Reymond. 970 million druglike small molecules for virtual screening in the chemical universe database GDB-13. *J. Am. Chem. Soc.*, 131:8732, 2009.

[3] Giorgos Bouritsas, Fabrizio Frasca, Stefanos P Zafeiriou, and Michael Bronstein. Improving graph neural network expressivity via subgraph isomorphism counting. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 2022.

[4] Joan Bruna and Stéphane Mallat. Invariant scattering convolution networks. *IEEE Trans. Pattern Anal. Mach. Intell.*, 35(8):1872–1886, 2013.

[5] Joan Bruna, Wojciech Zaremba, Arthur Szlam, and Yann Lecun. Spectral networks and locally connected networks on graphs. In *International Conference on Learning Representations (ICLR2014), CBLS, April 2014*, 2014.

[6] Benson Chen, Regina Barzilay, and Tommi Jaakkola. Path-augmented graph transformer network, 2019.

[7] Paul Civin, Nelson Dunford, and Jacob T. Schwartz. Linear operators. part i: General theory. *American Mathematical Monthly*, 67:199, 1960.

[8] Ronald R. Coifman and Mauro Maggioni. Diffusion wavelets. *Applied and Computational Harmonic Analysis*, 21(1):53–94, 2006. Special Issue: Diffusion Maps and Wavelets.

[9] Michaël Defferrard, Xavier Bresson, and Pierre Vandergheynst. Convolutional neural networks on graphs with fast localized spectral filtering. *Advances in neural information processing systems*, 29, 2016.

[10] Fernando Gama, Alejandro Ribeiro, and Joan Bruna. Diffusion scattering transforms on graphs. In *7th International Conference on Learning Representations, ICLR 2019, New Orleans, LA, USA, May 6-9, 2019*. OpenReview.net, 2019.

[11] Fernando Gama, Alejandro Ribeiro, and Joan Bruna. Stability of graph scattering transforms. In Hanna M. Wallach, Hugo Larochelle, Alina Beygelzimer, Florence d'Alché-Buc, Emily B. Fox, and Roman Garnett, editors, *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019, December 8-14, 2019, Vancouver, BC, Canada*, pages 8036–8046, 2019.

[12] Feng Gao, Guy Wolf, and Matthew J. Hirn. Geometric scattering for graph data analysis. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning, ICML 2019, 9-15 June 2019, Long Beach, California, USA*, volume 97 of *Proceedings of Machine Learning Research*, pages 2122–2131. PMLR, 2019.

[13] Justin Gilmer, Samuel S Schoenholz, Patrick F Riley, Oriol Vinyals, and George E Dahl. Neural message passing for quantum chemistry. In *International conference on machine learning*, pages 1263–1272. PMLR, 2017.

[14] David K Hammond, Pierre Vandergheynst, and Rémi Gribonval. Wavelets on graphs via spectral graph theory. *Applied and Computational Harmonic Analysis*, 30(2):129–150, 2011.

[15] Vassilis N. Ioannidis, Siheng Chen, and Georgios B. Giannakis. Pruned graph scattering transforms. In *International Conference on Learning Representations*, 2020.

[16] Yu Jin and Joseph F. JáJá. Learning graph-level representations with gated recurrent neural networks. *CoRR*, abs/1805.07683, 2018.

[17] Steven Kearnes, Kevin McCloskey, Marc Berndl, Vijay Pande, and Patrick Riley. Molecular graph convolutions: moving beyond fingerprints. *Journal of computer-aided molecular design*, 30(8):595–608, 2016.

[18] Thomas N. Kipf and Max Welling. Semi-supervised classification with graph convolutional networks. In *5th International Conference on Learning Representations, ICLR 2017, Toulon, France, April 24-26, 2017, Conference Track Proceedings*. OpenReview.net, 2017.

[19] Nils M. Kriege, Pierre-Louis Giscard, and Richard C. Wilson. On valid optimal assignment kernels and applications to graph classification. In Daniel D. Lee, Masashi Sugiyama, Ulrike von Luxburg, Isabelle Guyon, and Roman Garnett, editors, *Advances in Neural Information Processing Systems 29: Annual Conference on Neural Information Processing Systems 2016, December 5-10, 2016, Barcelona, Spain*, pages 1615–1623, 2016.

[20] Panagiotis Kyriakis, Iordanis Fostiropoulos, and Paul Bogdan. Learning hyperbolic representations of topological features. In *International Conference on Learning Representations*, 2021.

[21] Junhyun Lee, Inyeop Lee, and Jaewoo Kang. Self-attention graph pooling. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 3734–3743. PMLR, 09–15 Jun 2019.

[22] Shengchao Liu, Mehmet F Demirel, and Yingyu Liang. N-gram graph: Simple unsupervised representation for graphs, with applications to molecules. *Advances in neural information processing systems*, 32, 2019.

[23] Stéphane Mallat. Group invariant scattering. *Communications on Pure and Applied Mathematics*, 65(10):1331–1398, 2012.

[24] Hana Melánová, Bernd Sturmfels, and Rosa Winter. Recovery from power sums. *Experimental Mathematics*, 0(0):1–10, 2022.

[25] Barry Simon Michael Reed. *Methods of modern mathematical physics, Volume 1*. Academic Press, 1981.

[26] Yimeng Min, Frederik Wenkel, and Guy Wolf. Scattering gcn: Overcoming oversmoothness in graph convolutional networks. *Advances in Neural Information Processing Systems*, 33:14498–14508, 2020.

[27] Mathias Niepert, Mohamed Ahmed, and Konstantin Kutzkov. Learning convolutional neural networks for graphs. In Maria Florina Balcan and Kilian Q. Weinberger, editors, *Proceedings of The 33rd International Conference on Machine Learning*, volume 48 of *Proceedings of Machine Learning Research*, pages 2014–2023, New York, New York, USA, 20–22 Jun 2016. PMLR.

[28] Chao Pan, Siheng Chen, and Antonio Ortega. Spatio-temporal graph scattering transform, 2020.

[29] Michael Perlmutter, Feng Gao, Guy Wolf, and Matthew Hirn. Understanding graph neural networks with asymmetric geometric scattering transforms, 2019.

[30] Olaf. Post. *Spectral Analysis on Graph-like Spaces / by Olaf Post.* Lecture Notes in Mathematics, 2039. Springer Berlin Heidelberg, Berlin, Heidelberg, 1st ed. 2012. edition, 2012.

[31] Yu Rong, Yatao Bian, Tingyang Xu, Weiyang Xie, Ying Wei, Wenbing Huang, and Junzhou Huang. Self-supervised graph transformer on large-scale molecular data. *Advances in Neural Information Processing Systems*, 33:12559–12571, 2020.

[32] M. Rupp, A. Tkatchenko, K.-R. Müller, and O. A. von Lilienfeld. Fast and accurate modeling of molecular atomization energies with machine learning. *Physical Review Letters*, 108:058301, 2012.

[33] Kristof Schütt, Pieter-Jan Kindermans, Huziel Enoc Sauceda Felix, Stefan Chmiela, Alexandre Tkatchenko, and Klaus-Robert Müller. Schnet: A continuous-filter convolutional neural network for modeling quantum interactions. *Advances in neural information processing systems*, 30, 2017.

[34] Nino Shervashidze, Pascal Schweitzer, Erik Jan van Leeuwen, Kurt Mehlhorn, and Karsten M. Borgwardt. Weisfeiler-lehman graph kernels. *Journal of Machine Learning Research*, 12(77):2539–2561, 2011.

[35] Nino Shervashidze, SVN Vishwanathan, Tobias Petri, Kurt Mehlhorn, and Karsten Borgwardt. Efficient graphlet kernels for large graph comparison. In David van Dyk and Max Welling, editors, *Proceedings of the Twelth International Conference on Artificial Intelligence and Statistics*, volume 5 of *Proceedings of Machine Learning Research*, pages 488–495, Hilton Clearwater Beach Resort, Clearwater Beach, Florida USA, 16–18 Apr 2009. PMLR.

[36] David I Shuman, Christoph Wiesmeyr, Nicki Holighaus, and Pierre Vandergheynst. Spectrum-adapted tight graph wavelet and vertex-frequency frames. *IEEE Transactions on Signal Processing*, 63(16):4223–4235, 2015.

[37] Jake Topping, Francesco Di Giovanni, Benjamin Paul Chamberlain, Xiaowen Dong, and Michael M. Bronstein. Understanding over-squashing and bottlenecks on graphs via curvature, 2021.

[38] Wihler T.P. On the hölder continuity of matrix functions for normal matrices. *Journal of inequalities in pure and applied mathematics*, 10(4), Dec 2009.

[39] Thomas Wiatowski. *Harmonic analysis of deep convolutional neural networks*. PhD thesis, ETH Zurich, Switzerland, August 2017.

[40] Thomas Wiatowski and Helmut Bölcskei. A mathematical theory of deep convolutional neural networks for feature extraction. *IEEE Transactions on Information Theory*, 64:1845–1866, 2018.

[41] Zhenqin Wu, Bharath Ramsundar, Evan N. Feinberg, Joseph Gomes, Caleb Geniesse, Aneesh S. Pappu, Karl Leswing, and Vijay S. Pande. Moleculenet: a benchmark for molecular machine learning† †electronic supplementary information (esi) available. see doi: 10.1039/c7sc02664a. *Chemical Science*, 9:513 – 530, 2018.

[42] Zhaoping Xiong, Dingyan Wang, Xiaohong Liu, Feisheng Zhong, Xiaozhe Wan, Xutong Li, Zhaojun Li, Xiaomin Luo, Kaixian Chen, Hualiang Jiang, and Mingyue Zheng. Pushing the boundaries of molecular representation for drug discovery with the graph attention mechanism. *Journal of Medicinal Chemistry*, 63(16):8749–8760, 2020. PMID: 31408336.

[43] Keyulu Xu, Weihua Hu, Jure Leskovec, and Stefanie Jegelka. How powerful are graph neural networks? In *7th International Conference on Learning Representations, ICLR 2019, New Orleans, LA, USA, May 6-9, 2019*. OpenReview.net, 2019.

[44] Pinar Yanardag and S.V.N. Vishwanathan. Deep graph kernels. In *Proceedings of the 21th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD '15, page 1365–1374, New York, NY, USA, 2015. Association for Computing Machinery.

[45] Pinar Yanardag and S.V.N. Vishwanathan. Deep graph kernels. In *Proceedings of the 21th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD '15, page 1365–1374, New York, NY, USA, 2015. Association for Computing Machinery.

[46] Kevin Yang, Kyle Swanson, Wengong Jin, Connor Coley, Philipp Eiden, Hua Gao, Angel Guzman-Perez, Timothy Hopper, Brian Kelley, Miriam Mathea, Andrew Palmer, Volker Settels, Tommi Jaakkola, Klavs Jensen, and Regina Barzilay. Analyzing learned molecular representations for property prediction. *Journal of Chemical Information and Modeling*, 59(8):3370–3388, 2019. PMID: 31361484.

[47] Shuwen Yang, Ziyao Li, Guojie Song, and Lingsheng Cai. Deep molecular representation learning via fusing physical and chemical information. *Advances in Neural Information Processing Systems*, 34, 2021.

[48] Muhan Zhang, Zhicheng Cui, Marion Neumann, and Yixin Chen. An end-to-end deep learning architecture for graph classification. In Sheila A. McIlraith and Kilian Q. Weinberger, editors, *Proceedings of the Thirty-Second AAAI Conference on Artificial Intelligence, (AAAI-18), the 30th innovative Applications of Artificial Intelligence (IAAI-18), and the 8th AAAI Symposium on Educational Advances in Artificial Intelligence (EAAI-18), New Orleans, Louisiana, USA, February 2-7, 2018*, pages 4438–4445. AAAI Press, 2018.

[49] Qi Zhao and Yusu Wang. Learning metrics for persistence-based summaries and applications for graph classification. *Advances in Neural Information Processing Systems*, 32, 2019.

[50] Dongmian Zou and Gilad Lerman. Graph convolutional neural networks via scattering. *Applied and Computational Harmonic Analysis*, 49(3):1046–1074, nov 2020.

A Some Concepts in Linear Algebra

In the interest of self-containedness, we provide a brief review of some concepts from linear algebra utilized in this work that might potentially be considered more advanced. Presented results are all standard; a very thorough reference is [25].

Hilbert Spaces: To us, a Hilbert space — often denoted by $\mathcal{H}$ — is a vector space over the complex numbers which also has an inner product — often denoted by $\langle \cdot, \cdot \rangle_{\mathcal{H}}$. Prototypical examples are given by the Euclidean spaces $\mathbb{C}^d$ with inner product $\langle x, y \rangle_{\mathbb{C}^d} := \sum_{i=1}^d \overline{x}_i y_i$. Associated to an inner product is a norm, denoted by $\| \cdot \|_{\mathcal{H}}$ and defined by $\|x\|_{\mathcal{H}} := \sqrt{\langle x, x \rangle_{\mathcal{H}}}$ for $x \in \mathcal{H}$.

Direct Sums of Spaces: Given two potentially different Hilbert spaces $\mathcal{H}$ and $\widehat{\mathcal{H}}$, one can form their direct sum $\mathcal{H} \oplus \widehat{\mathcal{H}}$. Elements of $\mathcal{H} \oplus \widehat{\mathcal{H}}$ are vectors of the form (a, b), with $a \in \mathcal{H}$ and $b \in \widehat{\mathcal{H}}$. Addition and scalar multiplication are defined in the obvious way by

$$(a, b) + \lambda(c, d) := (a + \lambda c, b + \lambda d)$$

for $a, c \in \mathcal{H}$, $b, d \in \widehat{\mathcal{H}}$ and $\lambda \in \mathbb{C}$. The inner product on the direct sum is defined by

$$\langle (a, b), (c, d) \rangle_{\mathcal{H} \oplus \widehat{\mathcal{H}}} := \langle a, c \rangle_{\mathcal{H}} + \langle b, d \rangle_{\widehat{\mathcal{H}}}.$$

As is readily checked, this implies that the norm $\| \cdot \|_{\mathcal{H} \oplus \widehat{\mathcal{H}}}$ on the direct sum is given by

$$\|(a, b)\|^2_{\mathcal{H} \oplus \widehat{\mathcal{H}}} := \|a\|^2_{\mathcal{H}} + \|b\|^2_{\widehat{\mathcal{H}}}.$$

Standard examples of direct sums are again the Euclidean spaces, where one has $\mathbb{C}^d = \mathbb{C}^n \oplus \mathbb{C}^m$ if $m + n = d$, as is easily checked. One might also consider direct sums with more than two summands, writing $\mathbb{C}^d = \oplus_{i=1}^d \mathbb{C}$ for example. In fact, one might also consider infinite sums of Hilbert spaces: The space $\oplus_{i=1}^\infty \mathcal{H}_i$ is made up of those elements $a = (a_1, a_2, a_3, ...)$ with $a_i \in \mathcal{H}_i$ for which the norm

$$\|a\|^2_{\oplus_{i=1}^\infty \mathcal{H}_i} := \sum_{i=1}^\infty \|a_i\|^2_{\mathcal{H}_i}$$

is finite. This means for example that the vector $(1, 0, 0, 0, ...)$ is in $\oplus_{i=1}^\infty \mathbb{C}$, while $(1, 1, 1, 1, ...)$ is not.

Direct Sums of Maps: Suppose we have two collections of Hilbert spaces $\{\mathcal{H}_i\}_{i=1}^\Gamma$, $\{\widetilde{\mathcal{H}}_i\}_{i=1}^\Gamma$ with $\Gamma \in \mathbb{N}$ or $\Gamma = \infty$. Suppose further that for each $i \leqslant \Gamma$ (resp. $i < \Gamma$) we have a (not necessarily linear) map $J_i : \mathcal{H}_i \to \widetilde{\mathcal{H}}_i$. Then the collection $\{J_i\}_{i=1}^\Gamma$ of these 'component' maps induce a 'composite' map

$$\mathscr{J} : \oplus_{i=1}^\Gamma \mathcal{H}_i \longrightarrow \oplus_{i=1}^\Gamma \widetilde{\mathcal{H}}_i$$

between the direct sums. Its value on an element $a = (a_1, a_2, a_3, ...) \in \oplus_{i=1}^\Gamma \mathcal{H}_i$ is defined by

$$\mathscr{J}(a) = (J_1(a_1), J_2(a_2), J_3(a_3), ...) \in \oplus_{i=1}^\Gamma \widetilde{\mathcal{H}}_i.$$

Strictly speaking, one has to be a bit more careful in the case where $\Gamma = \infty$ to ensure that $\|\mathscr{J}(a)\|_{\oplus_{i=1}^\infty \widetilde{\mathcal{H}}_i} \neq \infty$. This can however be ensured if we have $\|J_i(a_i)\|_{\widetilde{\mathcal{H}}_i} \leqslant C\|a_i\|_{\mathcal{H}_i}$ for all $1 \leqslant i$ and some C independent of all i, since then $\|\mathscr{J}(a)\|_{\oplus_{i=1}^\infty \widetilde{\mathcal{H}}_i} \leqslant C\|a\|_{\oplus_{i=1}^\infty \mathcal{H}_i} \leqslant \infty$. If each J_i is a linear operator, such a C exists precisely if the operator norms (defined below) of all J_i are smaller than some constant.

Operator Norm: Let $J : \mathcal{H} \to \widetilde{\mathcal{H}}$ be a linear operator between Hilbert spaces. We measure its 'size' by what is called the operator norm, denoted by $\| \cdot \|_{op}$ and defined by

$$\|J\|_{op} := \sup_{\psi \in \mathcal{H}, \|\psi\|_{\mathcal{H}} = 1} \frac{\|A\psi\|_{\widetilde{\mathcal{H}}}}{\|\psi\|_{\mathcal{H}}}.$$

Adjoint Operators Let $J : \mathcal{H} \to \widetilde{\mathcal{H}}$ be a linear operator from the Hilbert space $\mathcal{H}$ to the Hilbert space $\widetilde{\mathcal{H}}$. Its adjoint $J^* : \widetilde{\mathcal{H}} \to \mathcal{H}$ is an operator mapping in the opposite direction. It is uniquely determined by demanding that

$$\langle Jf, u \rangle_{\widetilde{\mathcal{H}}} = \langle f, J^* u \rangle_{\mathcal{H}}$$

holds true for arbitrary $f \in \mathcal{H}$ and $u \in \widetilde{\mathcal{H}}$.

Normal Operators: If a linear operator $\Delta : \mathcal{H} \to \mathcal{H}$ maps from and to the same Hilbert space, we can compare it directly with its adjoint. If $\Delta\Delta^* = \Delta^*\Delta$, we say that the operator Δ is normal. Special instances of normal operators are self-adjoint operators, for which we have the stronger property $\Delta = \Delta^*$. If an operator is normal, there are unitary maps $U : \mathcal{H} \to \mathcal{H}$ diagonalizing Δ as

$$U^* \Delta U = \operatorname{diag}(\lambda_1, ... \lambda_n),$$

with eigenvalues in $\mathbb{C}$. We call the collection of eigenvalues the spectrum $\sigma(\Delta)$ of Δ. If $\dim \mathcal{H} = d$, we may write $\sigma(\Delta) = \{\lambda\}_{i=1}^d$. It is a standard exercise to verify that each eigenvalue satisfies $|\lambda_i| \leq \|\Delta\|_{op}$. Associated to each eigenvalue is an eigenvector ϕ_i. The collection of all (normalized) eigenvectors forms an orthonormal basis of $\mathcal{H}$. We may then write

$$\Delta f = \sum_{i=1}^d \lambda_i \langle \phi_i, f \rangle_{\mathcal{H}} \phi_i.$$

Resolvent of a (normal) Operator: Given a normal operator Δ on some Hilbert space $\mathcal{H}$, we have that the operator $(\Delta - z) : \mathcal{H} \to \mathcal{H}$ is invertible precisely if $z \neq \sigma(\Delta)$. In this case we write

$$R(z, \Delta) = (\Delta - z)^{-1}$$

and call this operator the **resolvent** of Δ at z. It can be proved that the norm of the resolvent satisfies

$$\|R(z, \Delta)\|_{op} = \frac{1}{dist(z, \sigma(\Delta))},$$

where $dist(z, \sigma(\Delta))$ denotes the minimal distance between z and any eigenvalue of Δ.

Functional Calculus: Given a normal operator $\Delta : \mathcal{H} \to \mathcal{H}$ on a Hilbert space of dimension d and a complex function $g : \mathbb{C} \to \mathbb{C}$, we can define another normal operator obtained from applying the function g to Δ by

$$g(\Delta)f = \sum_{i=1}^f g(\lambda_i)\langle \phi_i, f \rangle_{\mathcal{H}} \phi_i.$$

For example if $g(\cdot) = |\cdot|$, we obtain the absolute value $|\Delta|$ of Δ by specifying for all $f \in \mathcal{H}$ that

$$|\Delta|f = \sum_{i=1}^d |\lambda_i|\langle \phi_i, f \rangle_{\mathcal{H}} \phi_i.$$

Similarly we find (if $z \notin \sigma(\Delta)$ and for $f \in \mathcal{H}$)

$$\frac{1}{\Delta - z} = \sum_{i=1}^d \frac{1}{\lambda_i - z}\langle \phi_i, f \rangle_{\mathcal{H}} \phi_i = (\Delta - z)^{-1} = R(z, \Delta)$$

where we think of the left-hand-side as applying a function to Δ, while we think of the right-hand-side as inverting the operator $(\Delta - z)$. This now allows us to apply tools from complex analysis also to operators: If a function g is analytic (i.e. can be expanded into a power series), we have

$$g(\lambda) = -\frac{1}{2\pi i} \oint_S \frac{g(z)}{\lambda - z} dz$$

for any circle $S \subseteq \mathbb{C}$ encircling λ by Cauchy's integral formula. Thus, if we chose S large enough to encircle the entire spectrum $\sigma(\Delta)$, we have

$$g(\Delta)f = -\sum_{i=1}^{d} \frac{1}{2\pi i} \oint_S \frac{g(z)}{\lambda_i - z} dz \langle \phi_i, f \rangle_{\mathcal{H}} \phi_i = -\frac{1}{2\pi i} \oint_S g(z) R(z, \lambda) dz.$$

Frobenius Norm: Given a finite dimensional Hilbert space $\mathcal{H}$ with inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$, and an orthonormal basis $\{\phi_i\}_{i=1}^{d}$, we define the trace of an operator $A : \mathcal{H} \to \mathcal{H}$ as

$$Tr(A) := \sum_{k=1}^{d} \langle \phi_k, A\phi_k \rangle_{\mathcal{H}}.$$

It is a standard exercise to show that this is independent of the choice of orthonormal basis. The associated Frobenius inner product on the space of operators is then given as

$$\langle B, A \rangle_F := Tr(B^*A) \sum_{k=1}^{d} \langle \phi_k, B^*A\phi_k \rangle_{\mathcal{H}}.$$

Hence the Frobenius norm of an operator is determined by

$$\|A\|_F^2 = Tr(A^*A) = \sum_{k=1}^{d} \langle \phi_k, A^*A\phi_k \rangle_{\mathcal{H}}.$$

It is a standard exercise to verify that we have $\|A\|_{op} \leqslant \|A\|_F$. Since the trace is independent of the choice of orthonormal basis, the Frobenius norm is invariant under unitary transformations. More precisely, if $U, V : \mathcal{H} \to \mathcal{H}$ are unitary, we have

$$\|UAV\|_F^2 = \|A\|_F^2.$$

Frobenius norms can be used to transfer Lipschitz continuity properties of complex functions to the setting of functions applied to normal operators:

Lemma A.1. Let $g : \mathbb{C} \to \mathbb{C}$ be Lipschitz continuous with Lipschitz constant D_g. This implies

$$\|g(X) - g(Y)\|_F \leqslant D_g \cdot \|X - Y\|_F.$$

for normal operators X, Y on $\mathcal{H}$.

Proof. This proof is taken (almost) verbatim from [38]. For an operator $A : \mathcal{H} \to \mathcal{H}$ denote by A_{ij} its matrix representation with respect to the orthonormal basis $\{\phi_i\}_{i=1}^{d}$:

$$A_{ij} := \langle \phi_i, A\phi_j \rangle_{\mathcal{H}}.$$

We then have

$$\|A\|_F^2 = \sum_{i,j=1}^{d} |A_{ij}|^2$$

as a quick calculation shows. Let now U, W be unitary (with respect to the inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$) operators diagonalizing the normal operators X and Y as

$$V^*XV = \text{diag}(\lambda_1, ...\lambda_n) =: D(X)$$
$$W^*YW = \text{diag}(\mu_1, ...\mu_n) =: D(Y).$$

Since the Frobenius norm is invariant under unitary transformations we find

$$
\begin{aligned}
\|g(X) - g(Y)\|_F^2 &= \|g(VD(X)V^*) - g(WD(Y)W^*)\|_F^2 \\
&= \|Vg(D(X))V^* - Wg(D(Y))W^*\|_F^2 \\
&= \|W^*Vg(D(X)) - g(D(Y))W^*V\|_F^2 \\
&= \sum_{i,j=1}^{d} |(W^*Vg(D(X)) - g(D(Y))W^*V)_{ij}|^2 \\
&= \sum_{i,j=1}^{d} \left| \sum_{k=1}^{n} [W^*V]_{ik}[g(D(X))]_{kj} - [g(D(Y))]_{ik}[W^*V]_{kj} \right|^2 \\
&= \sum_{i,j=1}^{d} |[W^*V]_{ij}|^2 |g(\lambda_j) - g(\mu_i)|^2 \\
&\leqslant \sum_{i,j=1}^{d} |[W^*V]_{ij}|^2 D_g^2 |\lambda_j - \mu_i|^2 \\
&= D_g^2 \sum_{i,j=1}^{d} \left| \sum_{k=1}^{n} [W^*V]_{ik}[D(X)]_{kj} - [D(Y)]_{ik}[W^*V]_{kj} \right|^2 \\
&= D_g^2 \|X - Y\|_F^2.
\end{aligned}
$$

$\square$

B Proof of Theorem 2.1

Theorem B.1. Let $\Delta : \ell^2(G) \to \ell^2(G)$ be normal. If the family $\{g_i(\cdot)\}_{i \in I}$ of bounded continuous functions satisfies $A \leqslant \sum_{i \in I} |g_i(c)|^2 \leqslant B$ for all c in the spectrum $\sigma(\Delta)$, we have $(\forall f \in \ell^2(G))$

$$
A\|f\|_{\ell^2(G)}^2 \leqslant \sum_{i \in I} \|g_i(\Delta)f\|_{\ell^2(G)}^2 \leqslant B\|f\|_{\ell^2(G)}^2.
$$

Proof. Writing the normalized eigenvalue-eigenvector sequence of Δ as $(\lambda_i, \phi_i)_{i=1}^{|G|}$, we simply note

$$
\sum_{i \in I} \sum_{k=1}^{|G|} |\langle g_i(\lambda_k)\phi_k, f\rangle_{\ell^2(G)}|^2 = \sum_{k=1}^{|G|} \left(\sum_{i \in I} |g_i(\lambda_k)|^2 \right) |\langle \phi_k, f\rangle_{\ell^2(G)}|^2.
$$

Now under the assumption, we can estimate the sum in brackets by A from below and by B from above. Then we need only use Bessel's (in)equality to prove

$$
A\|f\|^2 \leqslant \sum_{i \in \hat{I}} \sum_{k=1}^{|G|} |\langle g_i(\lambda_k)\phi_k, f\rangle_{\ell^2(G)}|^2 \leqslant B\|f\|^2.
$$

$\square$

C Proof of Theorem 4.1

Theorem C.1. With the notation of Section 3 and setting $B_0 = 1$, we have:

$$
\|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N}^2 \leqslant \left(1 + \sum_{n=1}^{N} \max\{[B_n(L_n^+ R_n^+)^2 - 1], 0\} \prod_{k=0}^{n-1} B_k(R_k^+ L_k^+)^2 \right) \|f - h\|_{\ell^2(G)}^2
$$

To streamline the argumentation let us first introduce some notation:

Notation C.2. Let us denote paths in Γ^N as $q := (\gamma_N, ..., \gamma_1)$. For $f \in \ell^2(G)$ let us write

$$f_q := U[\gamma_N] \circ ... \circ U[\gamma_1](f).$$

Proof. By Definition, we have

$$\|\Phi_N(f) - \Phi_N(g)\|^2_{\mathscr{F}_N} = \sum_{n=1}^{N} \left(\sum_{q \in \Gamma^{n-1}} \|V_n(f_q) - V_n(h_q)\|^2_{\ell^2(G_n)} \right)$$

$$= \sum_{n=1}^{N} \underbrace{\left(\sum_{q \in \Gamma^{n-1}} \|\chi_n(\Delta_n)\rho_n(P_n(f_q)) - \chi_n(\Delta_n)\rho_n(P_n(h_q))\|^2_{\ell^2(G_n)} \right)}_{=:a_n}.$$

We proceed in two steps:

Our initial goal is to upper bound a_n as

$$a_n \leqslant B_n(L_n^+ R_n^+)^2 \cdot b_{n-1} - b_n \equiv (b_{n-1} - b_n) + \left[B_n(L_n^+ R_n^+)^2 - 1 \right] \cdot b_{n-1} \tag{3}$$

for $b_n := \sum_{q \in \Gamma^n} \|f_q - h_q\|^2_{\ell^2(G_n)}$ with $b_0 = \|f - h\|^2_{\ell^2(G)}$. To achieve this we note that (3) is equivalent to

$$a_n + b_n \leqslant B_n(L_n^+ R_n^+)^2 \cdot b_{n-1}$$

which upon unraveling definitions may be written as

$$\sum_{q \in \Gamma^{n-1}} \|\chi_n(\Delta_n)\rho_n(P_n((f_q))) - \chi_n(\Delta_n)\rho_n(P_n(h_q)\|^2_{\ell^2(G_n)} + \sum_{\hat{q} \in \Gamma^n} \|f_{\hat{q}} - h_{\hat{q}}\|^2_{\ell^2(G_n)}$$
$$\leqslant B_n(L_n^+ R_n^+)^2 \sum_{q \in \Gamma^{n-1}} \|f_q - h_q\|^2_{\ell^2(G_{n-1})}. \tag{4}$$

To establish (4), we note, that in the sum over paths of length n, any $\hat{q} \in \Gamma^n$ can uniquely be written as $\hat{q} = (\gamma_n, q)$, with the path $q \in \Gamma^{n-1}$ of length $(n-1)$ determined by

$$\hat{q} = (\gamma_n, \underbrace{\gamma_{n-1}, ..., \gamma_1}_{=:q}).$$

With this we find

$$\sum_{\hat{q} \in \Gamma^n} \|f_{\hat{q}} - h_{\hat{q}}\|^2_{\ell^2(G_n)} = \sum_{\gamma_n \in \Gamma_n} \sum_{q \in \Gamma^{n-1}} \|g_{\gamma_n}(\Delta_n)\rho_n(P_n((f_q))) - g_{\gamma_n}(\Delta_n)\rho_n(P_n(h_q))\|^2_{\ell^2(G_n)}.$$

Thus we can rewrite the left hand side of (4) as

$$\sum_{q \in \Gamma^{n-1}} \|\chi_n(\Delta_n)\rho_n(P_n((f_q))) - \chi_n(\Delta_n)\rho_n(P_n(h_q)\|^2_{\ell^2(G_n)} + \sum_{\hat{q} \in \Gamma^n} \|f_{\hat{q}} - h_{\hat{q}}\|^2_{\ell^2(G_n)}$$

$$= \sum_{q \in \Gamma^{n-1}} \left(\|\chi_n(\Delta_n)\rho_n(P_n(f_q) - \chi_n(\Delta_n)\rho_n(P_n(h_q)\|^2_{\ell^2(G_n)} \right.$$

$$\left. + \sum_{\gamma_n \in \Gamma_n} \|g_{\gamma_n}(\Delta_n)\rho_n(P_n((f_q))) - g_{\gamma_n}(\Delta_n)\rho_n(P_n(h_q))\|^2_{\ell^2(G_n)} \right)$$

$$=:\star$$

The fact that in each layer the function $\{\chi_n(\cdot)\} \bigcup \{g_{\gamma_n}(\cdot)\}_{\gamma_n \in \Gamma_n}$ form a generalized frame with upper frame constant B_n implies by Theorem 2.1, that we can further bound this as

$$\star \leqslant B_n \sum_{q \in \Gamma^{n-1}} \|\rho_n(P_n(f_q) - \rho_n(P_n(h_q))\|^2_{\ell^2(G_n)}.$$

Using the Lipschitz continuity of ρ_n and P_n, we arrive at the desired expression (4).

Having established that

$$a_n \leqslant (b_{n-1} - b_n) + \left[B_n(L_n^+ R_n^+)^2 - 1 \right] \cdot b_{n-1}$$

holds true, we note that we can establish

$$b_{n-1} \leqslant \prod_{k=1}^{n-1} B_k (L_k^+ R_k^+)^2 b_{n-2}$$

arguing similarly as in the case of (4) by using (for $f \in \ell^2(G_{n-1})$)

$$\sum_{\gamma_{n-1} \in \Gamma_{n-1}} \|g_{\gamma_{n-1}}(\Delta_{n-1}) f\|_{\ell^2(G_{n-1})}^2 \leqslant \|\chi_{n-1}(\Delta_{n-1}) f\|_{\ell^2(G_{n-1})}^2 + \sum_{\gamma \in \Gamma} \|g_{\gamma_{n-1}}(\Delta_{n-1}) f\|_{\ell^2(G_{n-1})}^2$$

together with the frame property and Lipschitz continuities. We then iterate this inequality and recall that $b_0 = \|f - h\|_{\ell^2(G)}^2$. Using the fact that

$$\sum_{n=1}^{N} (b_{n-1} - b_n) = b_0 - b_N \leqslant b_0,$$

we finally find

$$\|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N}^2 \leqslant \left(1 + \sum_{n=1}^{N} \max\{[B_n(L_n^+ R_n^+)^2 - 1], 0\} \prod_{k=0}^{n-1} B_k(R_k^+ L_k^+)^2 \right) \|f - h\|_{\ell^2(G)}^2.$$

$\square$

D Proof or Theorem 4.2

Theorem D.1. Let Φ_N and $\widetilde{\Phi}_N$ be two admissible scattering transforms based on the same module sequence Ω_N and operator sequences $\mathscr{D}_N$, $\widetilde{\mathscr{D}}_N$ with the same connecting operators ($P_n = \widetilde{P}_n$) in each layer. Assume that the respective normal operators satisfy $\|\Delta_n - \widetilde{\Delta}_n\|_F \leqslant \delta$ for some $\delta > 0$. Further assume that the functions $\{g_{\gamma_n}\}_{\gamma_n \in \Gamma_n}$ and χ_n in each layer are Lipschitz continuous with associated Lipschitz constants satisfying $L_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_n} L_{g_{\gamma_n}}^2 \leqslant D^2$ for all $n \leqslant N$ and some $D > 0$. Then we have for all $f \in \ell^2(G)$

$$\|\widetilde{\Phi}_N(f) - \Phi_N(f)\|_{\mathscr{F}} \leqslant \sqrt{2(2^N - 1)} \cdot D \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

Notation D.2. Let us denote scattering propagators based on operators Δ_n and connecting operators P_n by U_n and scattering propagators based on operators $\widetilde{\Delta}_n$ by $\widetilde{U}_n$. Similarly, to Notation C.2, let us then write (with $q = (\gamma_N, ..., \gamma_1)$)

$$\widetilde{f}_q := \widetilde{U}_n[\gamma_n] \circ ... \circ \widetilde{U}_1[\gamma_1](f).$$

Proof. By definition we have

$$\|\Phi_N(f) - \widetilde{\Phi}_N(f)\|_{\mathscr{F}_N}^2 = \sum_{n=1}^{N} \underbrace{\left(\sum_{q \in \Gamma^{n-1}} \|\chi_n(\Delta_n)\rho_n(P_n((f_q))) - \chi_n(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(G_n)}^2 \right)}_{=:a_n}.$$

We define $b_n := \sum_{q \in \Gamma^n} \|f_q - \widetilde{f}_q\|_{\ell^2(G_n)}^2$, with $b_0 = \|f - h\|_{\ell^2(G)}^2 = 0$ and note

$$a_n + b_n = \sum_{q \in \Gamma^{n-1}} \left(\|\chi_n(\Delta_n)\rho_n(P_n(f_q)) - \chi_n(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(G_n)}^2 \right.$$

$$\left. + \sum_{\gamma_n \in \Gamma_n} \|g_{\gamma_n}(\Delta_n)\rho_n(P_n((f_q))) - g_{\gamma_n}(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(G_n)}^2 \right).$$

Using (with $|a + b|^2 \leqslant 2(|a|^2 + |b|^2)$)

$$\frac{1}{2}\|g_{\gamma_n}(\Delta_n)\rho_n(P_n(f_q)) - g_{\gamma_n}(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))\|^2_{\ell^2(G_n)}$$

$$\leqslant \|[g_{\gamma_n}(\Delta_n) - g_{\gamma_n}(\widetilde{\Delta}_n)]\rho_n(P_n(f_q))\|^2_{\ell^2(G_n)}$$

$$+ \|g_{\gamma_n}(\widetilde{\Delta}_n)[\rho_n(P_n((f_q))) - \rho_n(P_n(\widetilde{f}_q))]\|^2_{\ell^2(G_n)}$$

$$\leqslant \|[g_{\gamma_n}(\Delta_n) - g_{\gamma_n}(\widetilde{\Delta}_n)]\|^2_\infty \cdot \|\rho_n(P_n(f_q))\|^2_{\ell^2(G_n)}$$

$$+ \|g_{\gamma_n}(\widetilde{\Delta}_n)[\rho_n(P_n(f_q)) - \rho_n(P_n(\widetilde{f}_q))]\|^2_{\ell^2(G_n)},$$

and

$$\|[g_{\gamma_n}(\Delta_n) - g_{\gamma_n}(\widetilde{\Delta}_n)]\|^2_\infty \leqslant \|[g_{\gamma_n}(\Delta_n) - g_{\gamma_n}(\widetilde{\Delta}_n)]\|^2_F \leqslant L^2_{g_\gamma} \cdot \delta^2$$

(c.f. Lemma A.1), we find

$$a_n + b_n \leqslant 2 \sum_{q \in \Gamma^{n-1}} \left(L^2_{\chi_n} + \sum_{\gamma_n \in \Gamma_n} L_{g^2_{\gamma_n}} \right) (L^+_n R^+_n)^2 \|\rho_n(P_n(f_q))\|_{\ell^2(G_n)}$$

$$+ 2 \sum_{q \in \Gamma^{n-1}} B_n \|\rho_n(P_n(f_q)) - \rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(G_n)}.$$

Using $L^2_{\chi_n} + \sum_{\gamma_n \in \Gamma_n} L^2_{\gamma_n} \leqslant D^2$, we then infer (using the admissibility assumption $L^+_n, R^+_n, B_n \leqslant 1$)

$$a_n \leqslant (b_{n-1} - b_n) + b_{n-1} + 2D^2\delta^2\|f\|_{\ell^2(G)}.$$

Using similar arguments, we find

$$b_{n-1} \leqslant 2D^2\delta^2\|f\|^2_{\ell^2(G)} + 2b_{n-2} \leqslant 2D^2\delta^2\|f\|^2_{\ell^2(G)} + 4D^2\delta^2\|f\|^2_{\ell^2(G)} + 4b_{n-3}$$

$$\leqslant \left(\sum_{k=1}^{n-1} 2^k \right) D^2\delta^2\|f\|^2_{\ell^2(G)} = (2^n - 2)D^2\delta^2\|f\|^2_{\ell^2(G)}.$$

Thus we now know

$$a_n \leqslant 2^n D^2\delta^2\|f\|^2_{\ell^2(G)} + (b_{n-1} - b_n)$$

In total we find

$$\sum_{n=1}^{N} a_n \leqslant 2(2^N - 1)D^2\delta^2\|f\|^2_{\ell^2(G)},$$

where we have estimated the sum over $(b_{n-1} - b_n)$ by zero from above again. This establishes the claim. $\qquad\square$

E Prototypical Example illustrating ω-δ Closeness and δ-Unitary Equivalence

To investigate the example of Figure 2, we label the vertices of the respective graphs as depicted in Figure 5. We denote the left graph by G and the right graph by $\widetilde{G}$. The node-weights on $\widetilde{G}$ are given as $\widetilde{\mu}_i = 1$ for $1 \leqslant i \leqslant 7$, while on G the weights are given as $\mu_i = 1$ for $1 \leqslant i \leqslant 5$ while $\mu_6 = 2$. We then consider the respective un-normalized graph Laplacians $\Delta : \ell^2(G) \to \ell^2(G)$ and $\widetilde{\Delta} : \ell^2(\widetilde{G}) \to \ell^2(\widetilde{G})$, which for a given adjacency matrix W on a graph signal space $\ell^2(G)$ with node weights $\{\mu_i\}_i$ is given as

$$(\Delta f)_i = \frac{1}{\mu_i} \sum_j W_{ij}(f_i - f_j).$$

Such operators are positive and hence $|\Delta| = \Delta$ (similarly for $\widetilde{\Delta}$). We now need to find operators $J : \ell^2(G) \to \ell^2(\widetilde{G})$ and $\widetilde{J} : \ell^2(\widetilde{G}) \to \ell^2(G)$ satisfying the conditions of Definition 4.3. To construct J, we define a family $\{\psi_i\}_{i=1}^6$ of vectors on $\ell^2(\widetilde{G})$ as

$$\psi_1 = (1,0,0,0,0,0,0), \quad \psi_2 = (0,1,0,0,0,0,0),$$
$$\psi_3 = (0,0,1,0,0,0,0), \quad \psi_4 = (0,0,0,1,0,0,0),$$
$$\psi_5 = (0,0,0,0,1,0,0), \quad \psi_6 = (0,0,0,0,0,1,1).$$

Figure 5: Indexing on the respective graphs

The map $J : \ell^2(G) \to \ell^2(\widetilde{G})$ is then defined as

$$Jf := \sum_{i=1}^6 f_i \psi_i,$$

for any $f \in \ell^2(G)$. We take $\widetilde{J} : \ell^2(\widetilde{G}) \to \ell^2(G)$ to be its adjoint ($\widetilde{J} := J^*$), which determined explicitly by

$$(\widetilde{J}u)_i = \frac{1}{\mu_i} \langle \psi_i, u \rangle_{\ell^2(\widetilde{G})}$$

for any $u \in \ell^2(\widetilde{G})$ We shall now first check the conditions for δ-quasi unitary equivalence, which we list again for convenience; now adapted to our current setting:

$$\|Jf\|_{\ell^2(\widetilde{G})} \leqslant 2\|f\|_{\ell^2(G)}, \quad \|(J - \widetilde{J}^*)f\|_{\ell^2(\widetilde{G})} \leqslant \delta \|f\|_{\ell^2(G)},$$
$$\|f - \widetilde{J}Jf\|_{\ell^2(G)}^2 \leqslant \delta^2 \left(\|f\|_{\ell^2(G)}^2 + \langle f, \Delta \, f \rangle_{\ell^2(G)} \right), \quad \|u - J\widetilde{J}u\|_{\ell^2(\widetilde{G})}^2 \leqslant \delta^2 \left(\|u\|_{\ell^2(\widetilde{G})}^2 + \langle u, \widetilde{\Delta} \, u \rangle_{\ell^2(\widetilde{G})} \right).$$

We first note that since $\widetilde{J} = J^*$, we have $\|(J - \widetilde{J}^*)f\|_{\ell^2(\widetilde{G})} = 0$. Next we note

$$\|Jf\|_{\ell^2(\widetilde{G})}^2 = \sum_{i=1}^7 |(Jf)_i|^2 = |f_6|^2 + \sum_{i=1}^6 |f_i|^2 = \sum_{i=1}^6 \mu_i = \|f\|_{\ell^2(G)}^2.$$

Furthermore we note

$$(\widetilde{J}Jf)_i = \sum_{k=1}^6 f_k \underbrace{\frac{1}{\mu_i} \langle \psi_i, \psi_k \rangle_{\ell^2(\widetilde{G})}}_{=\delta_{ik}} = f_i$$

and hence $\|f - \widetilde{J}Jf\|_{\ell^2(G)}^2 = 0$. It remains to control $\|u - J\widetilde{J}u\|_{\ell^2(\widetilde{G})}^2$. We note

$$\widetilde{J}u = (u_1, u_2, u_3, u_4, u_5, (u_5 + u_6)/2)^\top$$

and thus

$$J\widetilde{J}u = (u_1, u_2, u_3, u_4, u_5, (u_6 + u_7)/2, (u_6 + u_7)/2)^\top,$$

671 Which implies

$$u - J\tilde{J}u = (0,0,0,0,0,(u_7 - u_6)/2, (u_6 - u_7)/2)^\top,$$

672 and thus

$$\|u - J\tilde{J}u\|^2_{\ell^2(\tilde{G})} = 2\frac{|u_6 - u_7|^2}{4} = \frac{|u_6 - u_7|^2}{2}.$$

673 We have

$$\langle u, \tilde{\Delta}\, u\rangle_{\ell^2(\tilde{G})} = \frac{1}{2}\sum_{i,j=1}^d \widetilde{W}_{ij}|u_i - u_j|^2.$$

674 Since $\widetilde{W}_{67} = 1/\delta^2$ by assumption, we have

$$\|u - J\tilde{J}u\|^2_{\ell^2(\tilde{G})} = \frac{1}{2}|u_6 - u_7|^2 = \frac{1}{2}\frac{\delta^2}{\delta^2}|u_6 - u_7|^2 = \frac{1}{2}\delta^2\widetilde{W}_{67}|u_6 - u_7|^2$$

$$\leqslant \frac{1}{2}\delta^2\sum_{i,j=1}^d \widetilde{W}_{ij}|u_i - u_j|^2 = \delta^2\langle u, \tilde{\Delta}\, u\rangle_{\ell^2(\tilde{G})}$$

$$\leqslant \delta^2\left(\|u\|^2_{\ell^2(\tilde{G})} + \langle u, \tilde{\Delta}\, u\rangle_{\ell^2(\tilde{G})}\right).$$

675 Thus we have proven δ-unitary-equivalence and it remains to establish (-1)-12δ closeness. Com-
676 bining Proposition 4.4.12. and Theorem 4.4.15 of [30], instead of bounding $\|(\tilde{R}J - JR)f\|_{\ell^2(\tilde{G})} \leqslant$
677 $12\delta\|f\|_{\ell^2(G)}$ directly, we may instead establish that there are operators $J^1 : \ell^2(G) \to \ell^2(\tilde{G})$,
678 $\widetilde{J^1} : \ell^2(\tilde{G}) \to \ell^2(G)$ satisfying

$$\|J^1 f - Jf\|_{\ell^2(\tilde{G})} \leqslant \delta^2\left(\|f\|_{\ell^2(G)} + \langle f, \Delta,\, f\rangle_{\ell^2(G)}\right), \tag{5}$$

679

$$\|\widetilde{J^1}u - \tilde{J}u\|_{\ell^2(G)} \leqslant \delta^2\left(\|u\|_{\ell^2(\tilde{G})} + \langle f, \tilde{\Delta},\, u\rangle_{\ell^2(\tilde{G})}\right), \tag{6}$$

680 and

$$\langle J^1 f, \tilde{\Delta}\, u\rangle_{\ell^2(\tilde{G})} = \langle f, \Delta\, \widetilde{J^1}u\rangle_{\ell^2(G)}. \tag{7}$$

681 We chose $J^1 = J$ and determine $\widetilde{J^1}$ by setting (for $(1 \leqslant i \leqslant 6)$)

$$(\widetilde{J^1}u)_i = u_i.$$

682 Thus (5) is clearly satisfied. For (6) we note that we have

$$(\tilde{J}u - \widetilde{J^1}u) = (0,0,0,0,0,(u_7 - u_6)/2).$$

683 Thus we have

$$\|\widetilde{J^1}u - \tilde{J}u\|_{\ell^2(G)} = \frac{1}{2}|u_6 - u_7|^2 \leqslant \delta^2\left(\|u\|^2_{\ell^2(\tilde{G})} + \langle u, \tilde{\Delta}\, u\rangle_{\ell^2(\tilde{G})}\right)$$

684 as before. It remains to establish (7). We have

$$\langle f, \Delta\, \widetilde{J^1}u\rangle_{\ell^2(G)} = \sum_{i,j=1}^6 \overline{f_i}W_{ij}(u_i - u_j),$$

685 while we have

$$\langle J^1 f, \tilde{\Delta}\, u\rangle_{\ell^2(\tilde{G})} = \sum_{i=1}^6 \overline{f_i}\cdot\langle\psi_i, \tilde{\Delta}\, u\rangle_{\ell^2(\tilde{G})}$$

$$= \sum_{i,j=1}^5 \overline{f_i}W_{ij}(U_j - u_i) + \overline{f_6}\cdot\langle\psi_6, \tilde{\Delta}\, u\rangle_{\ell^2(\tilde{G})}.$$

686 We have (with all node-weights on $\ell^2(G)$ equal to unity)

$$\langle \psi_6, \widetilde{\Delta}\, u\rangle_{\ell^2(\widetilde{G})} = (\Delta\, u)_6 + (\Delta\, u)_7 = \left(\sum_j W_{6j}(f_6 - f_j) + \frac{1}{\delta^2}(f_6 - f_7)\right) + \left(\frac{1}{\delta^2}(f_7 - f_6)\right)$$

$$= \left(\sum_j W_{6j}(f_6 - f_j) + \frac{1}{\delta^2}(f_6 - f_7)\right)$$

687 And thus

$$\langle J^1 f, \widetilde{\Delta}\, u\rangle_{\ell^2(\widetilde{G})} = \sum_{i,j=1}^{6} \overline{f_i} W_{ij}(u_i - u_j) = \langle f, \Delta\, \widetilde{J^1} u\rangle_{\ell^2(G)}$$

688 which proves the claim.

F Proof of Lemma 4.4

689

690 **Lemma F.1.** In the setting of Definition 4.3 let Δ and $\widetilde{\Delta}$ be ω-δ-close and satisfy $\|\Delta\|_{op}, \|\widetilde{\Delta}\|_{op} \leqslant K$
691 for some $K > 0$. If $g : \mathbb{C} \to \mathbb{C}$ is holomorphic on the disk $B_{K+1}(0)$ of radius $(K+1)$, there is a
692 constant $C_g \geqslant 0$ so that

$$\|g(\widetilde{\Delta})J - Jg(\Delta)\|_{op} \leqslant C_g \cdot \delta$$

693 with C_g depending on g, ω and K.

694 *Proof.* Without loss of generality, let us assume that $K > |\omega|$. Let us denote the circle of radius r in
695 $\mathbb{C}$ by S_r. For any holomorphic function g and (normal) operator Δ whose spectrum is enclosed by
696 the circle S_r, we can express the operator $g(\Delta)$ as

$$g(\Delta) = -\frac{1}{2\pi i} \oint_{S_r} \frac{g(z)}{\Delta - z} dz$$

697 as discussed in Appendix A (see also [7] for more details). Note that in our case the resolvent
698 $R(z, \Delta) = (\Delta - z)^{-1}$ is well defined for $|z| \geqslant K$, since with our assumptions all eigenvalues are
699 within the circle of radius K. Additionally note that we have

$$dist(z, \sigma(\Delta)) \geqslant dist(z, S_K) = |z| - K$$

700 if $|z| \geqslant K$. The same holds true after replacing Δ with $\widetilde{\Delta}$. Since for any normal operator Δ we have

$$\|R(z, \Delta)\|_{op} = 1/dist(z, \sigma(\Delta)),$$

701 we find

$$|R(z, \widetilde{\Delta})\|_{op}, \|R(z, \Delta)\|_{op} \leqslant 1/(|z| - K).$$

702 To quantify the difference $\|R(z, \widetilde{\Delta})J - JR(z, \Delta)\|_{op}$ in terms of the difference $\|\widetilde{R}(\omega)J -$
703 $JR(\omega)\|_{op} \leqslant \delta$, we define the function

$$\gamma_0(z) := 1 + \frac{|z - \omega|}{|z| - K},$$

704 for which

$$\|R(z, \widetilde{\Delta})J - JR(z, \Delta)\|_{op} \leqslant \gamma_0(z)^2 \|R(\omega, \widetilde{\Delta})J - JR(\omega, \widetilde{\Delta})\|_{op}$$

705 holds, as proved (in more general form) in Lemma 4.5.9 in [30]. Since on S_{K+1} we have and
706 $|z - \omega| \leqslant 2K + 1$ hence $\gamma_0(z) \leqslant 2(K+1)$, we find

$$\|g(\widetilde{\Delta})J - Jg(\Delta)\|_{op} = \left\| \frac{1}{2\pi i} \oint_{S_{K+1}} g(z)\left(R(z, \widetilde{\Delta}) - R(z, \Delta)\right) dz \right\|_{op}$$

$$\leqslant \frac{1}{2\pi} \oint_{S_{K+1}} |g(z)| \left\|R(z, \widetilde{\Delta}) - R(z, \Delta)\right\|_{op} dz$$

$$\leqslant 2\frac{(K+1)^2}{\pi} \left(\oint_{S_{K+1}} |g(z)| dz\right) \cdot \|R(\omega, \widetilde{\Delta})J - JR(\omega, \widetilde{\Delta})\|_{op}.$$

707 Thus we may set

$$C_g := 2\frac{(K+1)^2}{\pi} \oint\limits_{S_{K+1}} |g(z)|dz.$$

708 □

G Proof of Theorem 4.5

710 **Theorem G.1.** Let $\Phi_N, \widetilde{\Phi}_N$ be admissible scattering transforms based on a common module
711 sequence Ω_N and differing operator sequences $\mathscr{D}_N, \widetilde{\mathscr{D}}_N$. Assume that there are identifica-
712 tion operators $J_n : \ell^2(G_n) \to \ell^2(\widetilde{G}_n)$, $\widetilde{J}_n : \ell^2(\widetilde{G}_n) \to \ell^2(G_n)$ ($0 \leqslant n \leqslant N$) so that
713 the respective signal spaces are δ-unitarily equivalent, the respective normal operators $\Delta_n, \widetilde{\Delta}_n$
714 are ω-δ-close as well as bounded (in norm) by $K > 0$ and the connecting operators satisfy
715 $\|\widetilde{P}_n J_{n-1}f - J_n P_n f\|_{\ell^2(\widetilde{G}_n)} \leqslant \delta\|f\|_{\ell^2(G_{n-1})}$. For the common module sequence Ω_N assume that
716 the non-linearities satisfy $\|\rho_n(J_n f) - J_n \rho_n(f)\|_{\ell^2(\widetilde{G}_n)} \leqslant \delta\|f\|_{\ell^2(G_n)}$ and that the constants C_{χ_n} and
717 $\{C_{g_{\gamma_n}}\}_{\gamma_n \in \Gamma_N}$ associated through Lemma 4.4 to the functions of the generalized frames in each layer
718 satisfy $C_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_N} C_{g_{\gamma_n}}^2 \leqslant D^2$ for some $D > 0$. Denote the operator that the family $\{J_n\}_n$ of
719 identification operators induce on $\mathscr{F}_N$ through concatenation by $\mathscr{J}_N : \mathscr{F}_N \to \widetilde{\mathscr{F}}_N$. Then we have
720 with $K_N = \sqrt{(8^N - 1)(2D^2 + 12)/7}$ that

$$\|\widetilde{\Phi}_N(J_0 f) - \mathscr{J}_N \Phi_N(f)\|_{\widetilde{\mathscr{F}}_N} \leqslant K_N \cdot \delta \cdot \|f\|_{\ell^2(G)}, \quad \forall f \in \ell^2(G).$$

721 If additionally $\|\widetilde{P}_n J_{n-1}f - J_n P_n f\|_{\ell^2(\widetilde{G}_n)} = 0$ or $\|\rho_n(J_n f) - J_n \rho_n(f)\|_{\ell^2(\widetilde{G}_n)} = 0$ holds in each
722 layer, then we have $K_N = \mathcal{O}(4^{N/2})$; if both additional equations hold, we have $K_N = \mathcal{O}(2^{N/2})$.

723 **Notation G.2.** Let us denote scattering propagators based on operators Δ_n and P_n by U_n and
724 scattering propagators based on operators $\widetilde{\Delta}_n$ and $\widetilde{P}_n$ by $\widetilde{U}_n$. Similarly, to Notation D.2 and , let us
725 then write (with $q = (\gamma_N, ..., \gamma_1)$)

$$\widetilde{f}_q := \widetilde{U}_n[\gamma_n] \circ ... \circ \widetilde{U}_1[\gamma_1](J_0 f).$$

726 *Proof.* By definition we have

$$\|\mathscr{J}\Phi_N(f) - \widetilde{\Phi}_N(J_0 f)\|_{\widetilde{\mathscr{F}}_N}^2 = \sum_{n=1}^{N} \underbrace{\left(\sum_{q \in \Gamma^{n-1}} \|J_n \chi_n(\Delta_n)\rho_n(P_n(f_q)) - \chi_n(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(\widetilde{G}_n)}^2 \right)}_{=:a_n}.$$

727 We define $b_n := \sum_{q \in \Gamma^n} \|J_n f_q - \widetilde{f}_q\|_{\ell^2(\widetilde{G}_n)}^2$, with $b_0 = \|J_0 f - J_0 f\|_{\ell^2(\widetilde{G})}^2 = 0$ and note

$$a_n + b_n = \sum_{q \in \Gamma^{n-1}} \left(\|J_n \chi_n(\Delta_n)\rho_n(P_n(f_q) - \chi_n(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(\widetilde{G}_n)}^2 \right.$$

$$\left. + \sum_{\gamma_n \in \Gamma_n} \|J_n g_{\gamma_n}(\Delta_n)\rho_n(P_n((f_q))) - g_{\gamma_n}(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(\widetilde{G}_n)}^2 \right).$$

728 Using

$$\frac{1}{2}\|J_n g_{\gamma_n}(\Delta_n)\rho_n(P_n((f_q))) - g_{\gamma_n}(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(\widetilde{G}_n)}^2$$

$$\leqslant \|[J_n g_{\gamma_n}(\Delta_n) - g_{\gamma_n}(\widetilde{\Delta}_n)J_n]\rho_n(P_n(f_q))\|_{\ell^2(\widetilde{G}_n)}^2$$

$$+ \|g_{\gamma_n}(\widetilde{\Delta}_n)[J_n \rho_n(P_n((f_q))) - \rho_n(P_n(\widetilde{f}_q))]\|_{\ell^2(\widetilde{G}_n)}^2$$

$$\leqslant \|[J_n g_{\gamma_n}(\Delta_n) - g_{\gamma_n}(\widetilde{\Delta}_n)J_n]\|_{op} \cdot \|\rho_n(P_n(f_q))\|_{\ell^2(\widetilde{G}_n)}^2$$

$$+ \|g_{\gamma_n}(\widetilde{\Delta}_n)[J_n \rho_n(P_n((f_q))) - \rho_n(P_n(\widetilde{f}_q))]\|_{\ell^2(\widetilde{G}_n)}^2,$$

729 and $\|[g_{\gamma_n}(\Delta_n) - g_{\gamma_n}(\widetilde{\Delta}_n)]\|_\infty \leqslant C_{g_\gamma}^2 \cdot \delta^2$ (c.f. Lemma 4.4), we find

$$
\begin{aligned}
a_n + b_n \leqslant{} & 2 \sum_{q \in \Gamma^{n-1}} \left(C_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_n} C_{g_{\gamma_n}^2} \right) (L_n^+ R_n^+)^2 \|\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(\widetilde{G}_n)} \\
& + 2 \sum_{q \in \Gamma^{n-1}} B_n \|J_n \rho_n(P_n(f_q)) - \rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(\widetilde{G}_n)} \\
\leqslant{} & 2 \sum_{q \in \Gamma^{n-1}} \left(C_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_n} C_{g_{\gamma_n}^2} \right) (L_n^+ R_n^+)^2 \|\rho_n(P_n(\widetilde{f}_q))\|_{\ell^2(\widetilde{G}_n)} \\
& + 4\|f\|_{\ell^2(G)} \delta^2 + 8\|f\|_{\ell^2(G)} \delta^2 + 8 b_{n-1},
\end{aligned}
$$

730 where the second inequality arises from permuting the identification operator J_n through non-linearity
731 and connecting operator. Using $C_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_n} C_{\gamma_n}^2 \leqslant D^2$, we then infer

$$
a_n \leqslant (b_{n-1} - b_n) + 7 b_{n-1} + (2D^2 + 12)\delta^2 \|f\|_{\ell^2(G)}^2.
$$

732 Using similar arguments, we find

$$
\begin{aligned}
b_{n-1} \leqslant{} & (2D^2 + 12)\delta^2 \|f\|_{\ell^2(G)}^2 + 8 b_{n-2} \leqslant (1 + 8)(2D^2 + 12)\delta^2 \|f\|_{\ell^2(G)}^2 + 8^2 b_{n-3} \\
\leqslant{} & \left(\sum_{k=1}^{n-1} 8^{k-1} \right) (2D^2 + 12)\delta^2 \|f\|_{\ell^2(G)}^2 = \frac{1}{56}(8^n - 8)(2D^2 + 12)\delta^2 \|f\|_{\ell^2(G)}^2.
\end{aligned}
$$

733 In total we find

$$
\sum_{n=1}^{N} a_n \leqslant \underbrace{(b_0 - b_N)}_{\leqslant 0} + (2D^2 + 12)\delta^2 \|f\|_{\ell^2(G)}^2 \sum_{n=1}^{N} 8^{n-1} \leqslant \frac{1}{7}(8^N - 1)(2D^2 + 12)\delta^2 \|f\|_{\ell^2(G)}^2.
$$

734 If one of the additional equations holds, we find

$$
a_n + b_n \leqslant (b_{n-1} - b_n) + 3 b_{n-1} + (2D^2 + 4)\delta^2 \|f\|_{\ell^2(G)}^2.
$$

735 and

$$
\begin{aligned}
b_{n-1} \leqslant{} & (2D^2 + 4)\delta^2 \|f\|_{\ell^2(G)}^2 + 4 b_{n-2} \leqslant (1 + 4)(2D^2 + 4)\delta^2 \|f\|_{\ell^2(G)}^2 + 4^2 b_{n-3} \\
\leqslant{} & \left(\sum_{k=1}^{n-1} 4^{k-1} \right) (2D^2 + 4)\delta^2 \|f\|_{\ell^2(G)}^2 = \frac{1}{12}(4^n - 4)(2D^2 + 4)\delta^2 \|f\|_{\ell^2(G)}^2.
\end{aligned}
$$

736 Thus

$$
\sum_{n=1}^{N} a_n \leqslant \underbrace{(b_0 - b_N)}_{\leqslant 0} + (2D^2 + 4)\delta^2 \|f\|_{\ell^2(G)}^2 \sum_{n=1}^{N} 4^{n-1} \leqslant \frac{4}{3}(4^N - 1)(2D^2 + 4)\delta^2 \|f\|_{\ell^2(G)}^2.
$$

737 If both additional equations hold, we find

$$
a_n + b_n \leqslant (b_{n-1} - b_n) + b_{n-1} + (2D^2)\delta^2 \|f\|_{\ell^2(G)}^2.
$$

738 and

$$
\begin{aligned}
b_{n-1} \leqslant{} & 2D^2 \delta^2 \|f\|_{\ell^2(G)}^2 + 2 b_{n-2} \leqslant (1 + 2) 2D^2 \delta^2 \|f\|_{\ell^2(G)}^2 + 2^2 b_{n-3} \\
\leqslant{} & \left(\sum_{k=1}^{n-1} 2^{k-1} \right) 2D^2 \delta^2 \|f\|_{\ell^2(G)}^2 = \frac{1}{2}(2^n - 2) 2D^2 \delta^2 \|f\|_{\ell^2(G)}^2,
\end{aligned}
$$

739 which then implies

$$
\sum_{n=1}^{N} a_n \leqslant \underbrace{(b_0 - b_N)}_{\leqslant 0} + 2D^2 \delta^2 \|f\|_{\ell^2(G)}^2 \sum_{n=1}^{N} 2^{n-1} \leqslant (2^N - 1) 2D^2 \delta^2 \|f\|_{\ell^2(G)}^2.
$$

740 $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\quad \square$

 # H   Proof of Lemma 5.1

Lemma H.1. The energy W_N stored in layer N satisfies

$$C_N^-\|f\|_{\ell^2(G)}^2 \leqslant \|\Phi_N(f)\|_{\mathscr{F}_N} + W_N(f) \leqslant C_N^+\|f\|_{\ell^2(G)}^2,$$

with constants $C_N^- := \prod_{i=1}^N \min\{1, A_i(L_i^- R_i^-)^2\}$ and $C_N^+ := \prod_{i=1}^N \max\{1, B_i(L_i^+ R_i^+)^2\}$.

Proof.

$$\min\{1, A_1(L_1^- R_1^-)^2\}\|f\|_{\ell^2(G)}^2$$
$$=A_1(L_1^- R_1^-)^2\|f\|_{\ell^2(G)}^2$$
$$=A_1\|\rho_1(P_1(f))\|_{\ell^2(G_1)}^2$$
$$\leqslant \sum_{\gamma_1 \in \Gamma_1} \|g_{\gamma_1}(\Delta_1)\rho_1(P_1(f))\|_{\ell^2(G_1)}^2 + \|\chi_1(\Delta_1)\rho_1(P_1(f))\|_{\ell^2(G_1)}^2$$
$$= \sum_{q \in \Gamma^1} \|U[q](f)\|_{\ell^2(G_1)}^2 + \|\chi_1(\Delta_1)\rho_1(P_1(f))\|_{\ell^2(G_1)}^2$$
$$=\|\chi_1(\Delta_1)\rho_1(P_1(f))\|_{\ell^2(G_1)}^2 + W_1(f),$$

and similarly

$$\|\chi_1(\Delta_1)\rho_1(P_1(f))\|_{\ell^2(G_1)}^2 + W_1(f)$$
$$= \sum_{q \in \Gamma^1} \|U[q](f)\|_{\ell^2(G_1)}^2 + \|\chi_1(\Delta_1)\rho_1(P_1(f))\|_{\ell^2(G_1)}^2$$
$$\leqslant B_1(L_1^+ R_1^+)^2\|f\|_{\ell^2(G)}^2.$$

This yields the starting point for our induction. Now for the inductive step assume the claim holds up until layer $N-1$. Then we have

$$C_{N-1}^-\|f\|_{\ell^2(G)}^2 \leqslant \sum_{n=1}^{N-1}\left(\sum_{q \in \Gamma^{n-1}} \|\chi_n(\Delta_n)f_q\|_{\ell^2(G_n)}^2\right) + W_{N-1}(f) \leqslant C_{N-1}^+\|f\|_{\ell^2(G)}^2.$$

using Notation C.2. We note

$$\sum_{n=1}^N\left(\sum_{q \in \Gamma^{n-1}} \|\chi_n(\Delta_n)\rho_n(P_n(f_q))\|_{\ell^2(G_n)}^2\right) + W_N$$
$$=\sum_{n=1}^{N-1}\left(\sum_{q \in \Gamma^{n-1}} \|\chi_n(\Delta_n)\rho_n(P_n(f_q))\|_{\ell^2(G_n)}^2\right) + \sum_{q \in \Gamma^{N-1}} \|\chi_N(\Delta_N)\rho_N(P_N(f_q))\|_{\ell^2(G_N)}^2$$
$$+ \sum_{q \in \Gamma^N} \|f_q\|_{\ell^2(G_N)}^2.$$

Every path $\tilde{q} \in \Gamma^N$ may be written as $q = (\gamma_n, q)$, for some $\gamma_n \in \Gamma_n$ and $q \in \Gamma^{N-1}$. Thus we have

$$\sum_{q \in \Gamma^N} \|f_q\|_{\ell^2(G_N)}^2 = \sum_{q \in \Gamma^{N-1}} \sum_{\gamma_N \in \Gamma_N} \|g_{\gamma_N}(\Delta_N)P_N(\rho_N(f_q))\|_{\ell^2(G_N)}^2$$

749 Inserting this in the above equation yields

$$\sum_{n=1}^{N} \left(\sum_{q \in \Gamma^{n-1}} ||\chi_n(\Delta_n)\rho_n(P_n(f_q))||^2_{\ell^2(G_n)} \right) + W_N$$

$$= \sum_{n=1}^{N-1} \left(\sum_{q \in \Gamma^{n-1}} ||\chi_n(\Delta_n)\rho_n(P_n(f_q))||^2_{\ell^2(G_n)} \right)$$

$$+ \sum_{q \in \Gamma^{N-1}} \underbrace{\left(||\chi_N(\Delta_N)\rho_N(P_N(f_q))||^2_{\ell^2(G_{n-1})} + \sum_{\gamma_n \in \Gamma_N} ||g_{\gamma_N}(\Delta_N)P_N(\rho_N(f_q)||^2_{\ell^2(G_N)} \right)}_{=:\beta(f_q)}.$$

750 We have

$$(L_N^- R_N^-)^2 A_N ||f_q||^2_{\ell^2(G_{n-1})} \leqslant \beta(f_q) \leqslant (L_N^+ R_N^+)^2 B_N ||f_q||^2_{\ell^2(G_{n-1})},$$

751 by the operator frame property. With this we find:

$$\min\{1, (L_N^- R_N^-)^2 A_N\} \left(\sum_{n=1}^{N-1} \left(\sum_{q \in \Gamma^{n-1}} ||\chi_n(\Delta_n)\rho_n(P_n(f_q))||^2_{\ell^2(G_n)} \right) + W_{N-1} \right)$$

$$\leqslant \sum_{n=1}^{N} \sum_{q \in \Gamma^{n-1}} ||\chi_n(\Delta_n)\rho_n(P_n(f_q))||^2_{\ell^2(G_n)} + W_N$$

$$\leqslant \max\{1, (L_N^- R_N^-)^2 B_N\} \left(\sum_{n=1}^{N-1} \left(\sum_{q \in \Gamma^n} ||\chi_n(\Delta_n)U[q](f)||^2_{\ell^2(G_n)} \right) + W_{N-1} \right),$$

752 after unravelling the definition

$$W_{N-1}(f) \equiv \sum_{q \in \Gamma^N} ||f_q||^2_{\ell^2(G_{n-1})}.$$

753 The induction hypothesis together with the definition of $C_N^\pm$ now yields the claim.

754 □

I Proof of Lemma 5.2 and Corollary 5.3

756 **Theorem I.1.** Let Φ_∞ be a generalized admissible graph scattering transform based on a an operator
757 sequence $\mathscr{D}_\infty = (P_n, \Delta_n)_{n=1}^\infty$ and a module sequence Ω_∞ with each $\rho_n(\cdot) \geqslant 0$. Assume in each
758 layer $n \geqslant 1$ that there is an eigenvector ψ_n of Δ_n with solely positive entries; denote the smallest
759 entry by $m_n := \min_{i \in G_n} \psi_n[i]$. Denote the eigenvalue corresponding to ψ_n by λ_n. Quantify the
760 'spectral-gap' opened up at this eigenvalue through neglecting the output-generating function by
761 $\eta_n := \sum_{\gamma_n \in \Gamma_n} |g_{\gamma_n}(\lambda_n)|^2$ and assume $B_n m_n \geqslant \eta_n$. We then have

$$W_N(f) \leqslant C_N^+ \cdot \left[\prod_{n=1}^{N} \left(1 - \left(m_n^2 - \frac{\eta_n}{B_n} \right) \right) \right] \cdot ||f||^2_{\ell^2(G)}.$$

762 *Proof.* Denote the spectral projection (i.e. the orthogonal projection projecting to the space of
763 eigenvectors) onto the eigenspace corresponding to λ_n by P_c^n.

764   Then we have

$$
\begin{aligned}
W_N(f) &= \sum_{q\in\Gamma^{N-1}}\sum_{\gamma_N\in\Gamma_N} \|g_{\gamma_N}(\Delta_N)\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}\\
&= \sum_{q\in\Gamma^{N-1}}\sum_{\gamma_N\in\Gamma_N} \|g_{\gamma_N}(\Delta_N)(1-P_c^N)\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}\\
&\quad + \sum_{q\in\Gamma^{N-1}}\sum_{\gamma_N\in\Gamma_N} \|g_{\gamma_N}(\Delta_N)P_c^N\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}\\
&\leqslant \sum_{q\in\Gamma^{N-1}} B_N\|(1-P_c^N)\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}\\
&\quad + \sum_{q\in\Gamma^{N-1}} \eta_N\|P_c^N\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}\\
&\leqslant \sum_{q\in\Gamma^{N-1}} B_N\|(1-P_c^N)\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}\\
&\quad + \sum_{q\in\Gamma^{N-1}} \eta_N\|\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}.
\end{aligned}
$$

765   By orthogonality of the spectral projection, we then have

$$
\|(1-P_c^N)\rho_N(P_n(f_q))\|^2_{\ell^2(G_N)} = \|\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)} - \|P_c^N\rho_N(P_n(f_q))\|^2_{\ell^2(G_N)}.
$$

766   Furthermore, we have

$$
|\langle \psi^N, \rho_N(P_n(f_q))\rangle_{\ell^2(G_N)}|^2 \leqslant \|P_c^N\rho_N(P_n(f_q))\|^2_{\ell^2(G_N)}
$$

767   with equality if the multiplicity of λ^N is exactly one. With this we find

$$
\begin{aligned}
\|(1-P_c^N)\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)} &= \|\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)} - \|P_c^N\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}\\
&\leqslant \|\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)} - |\langle\psi^N,\rho_N(P_N(f_q))\rangle_{\ell^2(G_N)}|^2.
\end{aligned}
$$

768   Since the image of ρ_N is contained in $\mathbb{R}_+$ by assumption, we have

$$
\begin{aligned}
|\langle\psi^N,\rho_N(P_N(f_q))\rangle_{\ell^2(G_N)}|^2 &= \left|\sum_{i=1}^{|G_N|}\rho_N(P_N(f_q))_i(\psi_N)_i\mu_i\right|^2\\
&\geqslant \left|\sum_{i=1}^{|G_N|}|\rho_N(P_N(f_q))_i|\mu_i\right|^2\cdot m_N^2\\
&\geqslant \left|\sum_{i=1}^{|G_N|}|\rho_N(P_N(f_q))_i|^2\mu_i^2\right|\cdot m_N^2\\
&\geqslant \left|\sum_{i=1}^{|G_N|}|\rho_N(P_N(f_q))_i|^2\mu_i\right|\cdot m_N^2\\
&\geqslant \|\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}\cdot m_N^2
\end{aligned}
$$

769   Here the second to last inequality follows since in any finite dimensional vector space, the 1-norm is
770   larger than the 2-norm ($\|f\|_1 \geqslant \|f\|_2$) and all weights are assumed to satisfy $\mu_i \geqslant 1$. Thus we now
771   know

$$
\|(1-P_c^N)\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)} \leqslant \left(1-m_N^2\right)\|\rho_N(P_N(f_q))\|^2_{\ell^2(G_N)}.
$$

772   Inserting this in our estimate for $W_N(f)$ we find

$$W_N(f) \leqslant \left(1 - \left(m_N^2 - \frac{\eta_n}{B_n}\right)\right) L_N^+ R_N^+ B_N \cdot W_{N-1}(f)$$

$$\leqslant C_N^+ \prod_{n=1}^{N} \left(1 - \left(m_N^2 - \frac{\eta_n}{B_n}\right)\right) ||f||_{\ell^2(G)}^2.$$

$\square$

Taking N to infinity, we know that C_N^+ converges to something larger than zero by assumption. For products of the form $\prod_{n=0}^{N}(1 - q_n)$ with $0 \leqslant q_n < 1$ it is a standard exercise to prove that the limit is non-zero precisely if the sum over the q_n converges. But this is excluded by our assumption in Corollary 5.3.

J   Stability of Graph Level Feature Aggregation

J.1   Proof of Theorem 6.1

Theorem J.1. Assuming admissibility, we have $\|\Psi_N(f) - \Psi_N(g)\|_{\mathscr{R}_N} \leqslant \|f - g\|_{\ell^2(G)}$. With the conditions and notation of Theorem 4.2 we have

$$\|\Psi_N(f) - \widetilde{\Psi}_N(f)\|_{\mathscr{R}_N} \leqslant \sqrt{2(2^N - 1)} \cdot D \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

Additionally, in the setting of Theorem 4.5, assuming that for each $n \leqslant N$ the identification operator J_n satisfies $\left|\|J_n f\|_{\ell^1(\widetilde{G}_n)}/\sqrt{\mu_{\widetilde{G}_n}} - \|f\|_{\ell^1(G_n)}/\sqrt{\mu_{G_n}}\right|, \left|\|J_n f\|_{\ell^k(\widetilde{G}_n)} - \|f\|_{\ell^k(G_n)}\right| \leqslant \delta \cdot K \cdot \|f\|_{\ell^2(G_n)}$ $(2 \leqslant k \leqslant p_n)$ implies $(\forall f \in \ell^2(G))$

$$\|\widetilde{\Psi}_N(J_0 f) - \Psi_N(f)\|_{\mathscr{R}_N} \leqslant \sqrt{2} \cdot \sqrt{K_N^2 \cdot + K^2} \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

Furhermore, under the assumptions of Corollary 5.3 $\Psi_\infty(f) = 0$ implies $f = 0$.

Proof. Let $f, h \in \ell^2(G)$. To prove the first two claims, it suffices to prove

$$\|\Psi_N(f) - \Psi_N(h)\|_{\mathscr{R}_N} \leqslant \|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N},$$

and

$$\|\Psi_N(f) - \widetilde{\Psi}_N(f)\|_{\mathscr{R}_N} \leqslant \|\Phi_N(f) - \widetilde{\Phi}_N(f)\|_{\mathscr{F}_N}.$$

Both statements follow immediately, as soon as we have proved

$$\|N_p^G(f) - N_p^G(h)\|_{\mathbb{R}^p} \leqslant \|f - h\|_{\ell^2(G)}$$

for arbitrary choices of p and G. To this end we note that for $p \geqslant 2$ we have $\|f\|_{\ell^p(G)} \leqslant \|f\|_{\ell^2(G)}$ by the monotonicity of p-norms, while we have $\|f\|_{\ell^1(G)} \leqslant \sqrt{\mu_G} \cdot \|f\|_{\ell^2(G)}$ by Hölder's inequality. With this we find

$$\|N_p^G(f) - N_p^G(h)\|_{\mathbb{R}^p}^2 = \frac{1}{p}\left(\frac{1}{\mu_G}\|f\|_{\ell^1(G)} - \|h\|_{\ell^1(G)}|^2 + \sum_{i=2}^{p}\|f\|_{\ell^i(G)} - \|h\|_{\ell^i(G)}|^2\right)$$

$$\leqslant \frac{1}{p}\left(\frac{1}{\mu_G}\|f - h\|_{\ell^1(G)}|^2 + \sum_{i=2}^{p}\|f - h\|_{\ell^i(G)}|^2\right)$$

$$\leqslant \frac{1}{p} \cdot p \cdot \|f - h\|_{\ell^2(G)}|^2$$

$$= \|f - h\|_{\ell^2(G)}.$$

where we have employed the reverse triangle inequality in the first step.

793 To prove the second claim, we note that we have

$$\|\Psi_N(f) - \widetilde{\Psi}_N(J_0 f)\|^2_{\mathscr{R}_N}$$

$$= \sum_{n=1}^{N} \left(\sum_{q \in \Gamma^{n-1}} \|N_{p_n}^{G_n}(\underbrace{\chi_n(\Delta_n)\rho_n(P_n((f_q)))}_{=:x_q}) - N_{p_n}^{\widetilde{G}_n}(\underbrace{\chi_n(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))}_{=:\widetilde{x}_q})\|^2_{\mathbb{R}^{p_n}} \right)$$

$$\leqslant 2 \sum_{n=1}^{N} \left(\sum_{q \in \Gamma^{n-1}} \|N_{p_n}^{\widetilde{G}_n}(J_n x_q) - N_{p_n}^{\widetilde{G}_n}(\widetilde{x}_q)\|_{\mathbb{R}^{p_n}} \right)$$

$$+ 2 \sum_{n=1}^{N} \left(\sum_{q \in \Gamma^{n-1}} \|N_{p_n}^{\widetilde{G}_n}(J_n x_q) - N_{p_n}^{G_n}(x_q)\|_{\mathbb{R}^{p_n}} \right)$$

$$= 2 \|\mathscr{J}\Phi_N(f) - \widetilde{\Phi}_N(J_0 f)\|^2_{\mathscr{F}_N}$$

$$+ 2 \sum_{n=1}^{N} \left(\sum_{q \in \Gamma^{n-1}} \|N_{p_n}^{\widetilde{G}_n}(J_n x_q) - N_{p_n}^{G_n}(x_q)\|_{\mathbb{R}^{p_n}} \right).$$

794 Thus it remains to bound the last expression. We have

$$\|N_p^{\widetilde{G}_n}(J_n x_q) - N_{p_n}^{G_n}(x_q)\|_{\mathbb{R}^{p_n}}$$

$$= \frac{1}{p_n} \left(\left| \frac{1}{\sqrt{\mu_{G_n}}} \|f\|_{\ell^1(G)} - \frac{1}{\sqrt{\mu_{\widetilde{G}_n}}} \|J_n f\|_{\ell^1(\widetilde{G})} \right|^2 + \sum_{i=2}^{p_n} |\|f\|_{\ell^i(G)} - \|J_n f\|_{\ell^i(\widetilde{G})}|^2 \right)$$

$$\leqslant K^2 \cdot \delta^2 \cdot \|x_q\|^2_{\ell^2(G_n)}.$$

795 By our results of Appendix C and since we assume admissibility, we have

$$\sum_{n=1}^{N} \sum_{q \in \Gamma^{n-1}} \|x_q\|^2_{\ell^2(G_n)} \leqslant \|f\|^2_{\ell^2(G)}.$$

796 Thus in total

$$\|\Psi_N(f) - \widetilde{\Psi}_N(J_0 f)\|^2_{\mathscr{F}_N} \leqslant 2 \|\mathscr{J}\Phi_N(f) - \widetilde{\Phi}_N(J_0 f)\|^2_{\mathscr{F}_N} + 2K\delta \|f\|_{\ell^2(G)},$$

797 from which our stability claim follows.

798 It remains to prove that the assumptions of Corollary 5.3 $\Psi_\infty(f) = 0$ imply $f = 0$. But since
799 $N_p^G(f) = 0$ implies $f = 0$, this is clear. $\qquad\square$

J.2 Statement and Proof of the Analogue of Theorem 6.1 for Low-Pass feature Aggregation

801 The main assumption we have in this section is that each operator Δ_n (and $\widetilde{\Delta}_n$) has a simple lowest
802 lying eigenvalue equal to zero. We denote the associated eigenvector (determined up to a complex
803 phase) by ψ_{Δ_n} and the associated spectral projection to the lowest lying eigenvalue by P_{Δ_n}. It acts
804 as

$$P_{\Delta_n} f \equiv \psi_{\Delta_n} \langle \psi_{\Delta_n}, f \rangle_{\ell^2(G_n)}.$$

805 Now we are ready to state our main stability result under these circumstances:

806 **Theorem J.2.** Assuming admissibility, we have $\|\Psi_N^{|\langle \cdot, \cdot \rangle|}(f) - \Psi_N^{|\langle \cdot, \cdot \rangle|}(g)\|_{\mathscr{C}_N} \leqslant \|f - g\|_{\ell^2(G)}$. With
807 the conditions and notation of Theorem 4.2 and under the additional assumption $\|(P_{\Delta_n} - P_{\widetilde{\Delta}_n})\|_{op} \leqslant$
808 $K \cdot \delta$ for $n \leqslant N$ and some $K \geqslant 0$, we have

$$\|\Psi_N^{|\langle \cdot, \cdot \rangle|}(f) - \widetilde{\Psi}_N^{|\langle \cdot, \cdot \rangle|}(f)\|_{\mathscr{C}_N} \leqslant \sqrt{2} \cdot \sqrt{2(2^N - 1) + K^2} \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

809 In the setting of Theorem 4.5 and under the additional assumption $|\|P_{\Delta_n} f\|_{\ell^2(G_n)} -$
810 $\|P_{\widetilde{\Delta}_n} J_n f\|_{\ell^2(\widetilde{G}_n)}| \leqslant K\delta \|f\|_{\ell^2(G_n)}$ for all $f \in \ell^2(G_n)$ ($n \leqslant N$), we have

$$\|\widetilde{\Psi}_N^{|\langle \cdot, \cdot \rangle|}(J_0 f) - \Psi_N^{|\langle \cdot, \cdot \rangle|}(f)\|_{\mathscr{C}_N} \leqslant \sqrt{2} \cdot \sqrt{K_N^2 \cdot + K^2} \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

Proof. Let $f, h \in \ell^2(G)$. To prove the first claim, it suffices to prove

$$\|\Psi_N^{|\langle\cdot,\cdot\rangle|}(f) - \Psi_N^{|\langle\cdot,\cdot\rangle|}(h)\|_{\mathscr{C}_N} \leqslant \|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N}.$$

This immediately follows from the fact that for all $f \in \ell^2(G_n)$

$$|\langle\psi_{\Delta_n}, f\rangle_{\ell^2(G_n)}|^2 \leqslant \|\psi_{\Delta_n}\|_{\ell^2(G_n)}^2 \cdot \|f\|_{\ell^2(G_n)}^2$$

by Hölder's inequality.

The next claim we want to prove is that we have for all $f \in \ell^2(G)$

$$\|\Psi_N^{|\langle\cdot,\cdot\rangle|}(f) - \widetilde{\Psi}_N^{|\langle\cdot,\cdot\rangle|}(f)\|_{\mathscr{C}_N} \leqslant \sqrt{2} \cdot \sqrt{2(2^N - 1) + K^2} \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

We note

$$\|\Psi_N^{|\langle\cdot,\cdot\rangle|}(f) - \widetilde{\Psi}_N^{|\langle\cdot,\cdot\rangle|}(f)\|_{\mathscr{C}_N}^2$$

$$= \sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| |\langle\psi_{\Delta_n}, \underbrace{\chi_n(\Delta_n)\rho_n(P_n(f_q))}_{=:x_q}\rangle_{\ell^2(G_n)}| - |\langle\psi_{\widetilde{\Delta}_n}, \underbrace{\chi_n(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))}_{\widetilde{x}_q}\rangle_{\ell^2(G_n)}| \right|^2 \right)$$

$$= \sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| \|P_{\Delta_n}x_q\|_{\ell^2(G_n)} - \|P_{\widetilde{\Delta}_n}\widetilde{x}_q\|_{\ell^2(G_n)} \right|^2 \right)$$

$$\leqslant \sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \|P_{\Delta_n}x_q - P_{\widetilde{\Delta}_n}\widetilde{x}_q\|_{\ell^2(G_n)}^2 \right)$$

$$\leqslant 2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \|P_{\widetilde{\Delta}_n}(x_q - \widetilde{x}_q)\|_{\ell^2(G_n)}^2 \right) + 2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \|(P_{\Delta_n} - P_{\widetilde{\Delta}_n})x_q\|_{\ell^2(G_n)}^2 \right)$$

$$\leqslant 2\|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N}^2 + 2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \|(P_{\Delta_n} - P_{\widetilde{\Delta}_n})x_q\|_{\ell^2(G_n)}^2 \right)$$

Hence we need to bound the expression "$\|(P_{\Delta_n} - P_{\widetilde{\Delta}_n})x_q\|_{\ell^2(G_n)}^2$". We note

$$\|(P_{\Delta_n} - P_{\widetilde{\Delta}_n})x_q\|_{\ell^2(G_n)}^2 \leqslant \|(P_{\Delta_n} - P_{\widetilde{\Delta}_n})\|_{op} \cdot \|x_q\|_{\ell^2(G_n)}^2$$
$$\leqslant K^2 \cdot \delta^2 \cdot \|x_q\|_{\ell^2(G_n)}^2$$

and thus

$$\|\Psi_N^{|\langle\cdot,\cdot\rangle|}(f) - \widetilde{\Psi}_N^{|\langle\cdot,\cdot\rangle|}(f)\|_{\mathscr{C}_N}^2$$

$$\leqslant 2\|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N}^2 + 2K^2 \cdot \delta^2 \cdot \sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \|\chi_n(\Delta_n)\rho_n(P_n((f_q)))\|_{\ell^2(G_n)}^2 \right)$$

$$\leqslant 2\|\Phi_N(f) - \Phi_N(h)\|_{\mathscr{F}_N}^2 + 2K^2 \cdot \delta^2 \cdot \|f\|_{\ell^2(G)}^2$$

and the claim follows.

Finally we want to prove

$$\|\widetilde{\Psi}_N^{|\langle\cdot,\cdot\rangle|}(J_0f) - \Psi_N^{|\langle\cdot,\cdot\rangle|}(f)\|_{\mathscr{C}_N} \leqslant \sqrt{2} \cdot \sqrt{K_N^2 \cdot + K^2} \cdot \delta \cdot \|f\|_{\ell^2(G)}.$$

We note

$$\|\Psi_N^{|\langle\cdot,\cdot\rangle|^2}(f) - \widetilde{\Psi}_N^{|\langle\cdot,\cdot\rangle|}(f)\|_{\mathscr{C}_N}$$

$$= \sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| |\langle \psi_{\Delta_n}, \underbrace{\chi_n(\Delta_n)\rho_n(P_n((f_q)))}_{=:x_q}\rangle_{\ell^2(G_n)}| - |\langle \psi_{\widetilde{\Delta}_n}, \underbrace{\chi_n(\widetilde{\Delta}_n)\rho_n(P_n(\widetilde{f}_q))}_{\widetilde{x}_q}\rangle_{\ell^2(\widetilde{G}_n)}| \right|^2 \right)$$

$$= \sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| \|P_{\Delta_n}x_q\|_{\ell^2(G_n)} - \|P_{\widetilde{\Delta}_n}\widetilde{x}_q\|_{\ell^2(\widetilde{G}_n)} \right|^2 \right)$$

$$\leqslant \sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| \|P_{\Delta_n}x_q\|_{\ell^2(G_n)} - \|P_{\widetilde{\Delta}_n}J_n x_q\|_{\ell^2(\widetilde{G}_n)} + \|P_{\widetilde{\Delta}_n}J_n x_q\|_{\ell^2(\widetilde{G}_n)} - \|P_{\widetilde{\Delta}_n}\widetilde{x}_q\|_{\ell^2(\widetilde{G}_n)} \right|^2 \right)$$

$$\leqslant 2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| \|P_{\widetilde{\Delta}_n}J_n x_q\|_{\ell^2(\widetilde{G}_n)} - \|P_{\widetilde{\Delta}_n}\widetilde{x}_q\|_{\ell^2(\widetilde{G}_n)} \right|^2 \right)$$

$$+2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| \|P_{\Delta_n}x_q\|_{\ell^2(G_n)} - \|P_{\widetilde{\Delta}_n}J_n x_q\|_{\ell^2(\widetilde{G}_n)} \right|^2 \right)$$

$$\leqslant 2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| \|P_{\widetilde{\Delta}_n}J_n x_q\|_{\ell^2(\widetilde{G}_n)} - \|P_{\widetilde{\Delta}_n}\widetilde{x}_q\|_{\ell^2(\widetilde{G}_n)} \right|^2 \right)$$

$$+2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| \|P_{\Delta_n}x_q\|_{\ell^2(G_n)} - \|P_{\widetilde{\Delta}_n}J_n x_q\|_{\ell^2(\widetilde{G}_n)} \right|^2 \right)$$

$$\leqslant 2\|\mathscr{J}\Phi_N(f) - \widetilde{\Phi}_N(J_0 f)\|_{\widetilde{\mathscr{F}}_N}^2 + 2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} \left| \|P_{\Delta_n}x_q\|_{\ell^2(G_n)} - \|P_{\widetilde{\Delta}_n}J_n x_q\|_{\ell^2(\widetilde{G}_n)} \right|^2 \right)$$

$$\leqslant 2\|\mathscr{J}\Phi_N(f) - \widetilde{\Phi}_N(J_0 f)\|_{\widetilde{\mathscr{F}}_N}^2 + 2\sum_{n=1}^N \left(\sum_{q\in\Gamma^{n-1}} K^2 \cdot \delta^2 \|x_q\|_{\ell^2(G_n)}^2 \right)$$

$$\leqslant 2\|\mathscr{J}\Phi_N(f) - \widetilde{\Phi}_N(J_0 f)\|_{\widetilde{\mathscr{F}}_N}^2 + 2K^2 \cdot \delta^2 \|f\|_{\ell^2(G)}^2.$$

which proves the claim. $\qquad\square$

In establishing expressivity, we have to be a tiny bit more careful:

Theorem J.3. In the setting of of Corollary 5.3, assume that in each layer n, the output generating function χ_n of the underlying scattering transform satisfies $\chi_n(0) \neq 0$ and $\chi_n(\lambda_i) = 0$ for ordered non-zero eigenvalues $\lambda_2 \leqslant ... \leqslant \lambda_{|G_n|}$ of the operator Δ_n. Then $\Psi_\infty^{|\langle\cdot,\cdot\rangle|}(f) = 0$ implies $f = 0$.

Proof. Under these assumptions, we do not lose any information by projecting to ψ_{Δ_n} in each $\ell^2(G_n)$, since the image of $\chi_n(\Delta_n)$ is already contained in the one-dimensional space generated by the lowest lying eigenvector ψ_{Δ_n}. $\qquad\square$

K   Stability of Higher Order Scattering

In this section, we list stability theorems for higher order scattering transforms. As the respective proofs are identical to the corresponding results for the node setting, we do not repeat them here.

Theorem K.1. With the notation of Section 7, we have for all $f, h \in \ell^2(G^k)$:

$$\|\Phi_N^k(f) - \Phi_N^k(h)\|_{\mathscr{F}_N^k}^2 \leqslant \left(1 + \sum_{n=1}^N \max\{[B_n - 1], [B_n(L_n^+ R_n^+)^2 - 1], 0\} \prod_{\ell=1}^{n-1} B_\ell\right) \|f - h\|_{\ell^2(G^k)}^2$$

Theorem K.2. Let Φ_N and $\widetilde{\Phi}_N$ be two admissible scattering transforms based on the same module sequence Ω_N and operator sequences $\mathscr{D}_N^k, \widetilde{\mathscr{D}}_N^k$ with the same connecting operators $(P_n^k = \widetilde{P}_n^k)$ in each layer. Assume that the respective normal operators satisfy $\|\Delta_n^k - \widetilde{\Delta}_n^k\|_F \leqslant \delta$ for some $\delta > 0$. Further assume that the functions $\{g_{\gamma_n}\}_{\gamma_n \in \Gamma_n}$ and χ_n in each layer are Lipschitz continuous with associated Lipschitz constants satisfying $L_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_n} L_{g_{\gamma_n}}^2 \leqslant D^2$ for all $n \leqslant N$ and some $D > 0$. Then we have for all $f \in \ell^2(G^k)$

$$\|\widetilde{\Phi}_N^k(f) - \Phi_N^k(f)\|_{\mathscr{F}_N^k} \leqslant \sqrt{2(2^N - 1)} \cdot D \cdot \delta \cdot \|f\|_{\ell^2(G^k)}.$$

Theorem K.3. Let $\Phi_N^k, \widetilde{\Phi}_N^k$ be admissible scattering transforms based on a common module sequence Ω_N and differing operator sequences $\mathscr{D}_N^k, \widetilde{\mathscr{D}}_N^k$. Assume that there are identification operators $J_n : \ell^2(G_n^k) \to \ell^2(\widetilde{G}_n^k)$, $\widetilde{J}_n : \ell^2(\widetilde{G}_n) \to \ell^2(G_n^k)$ $(0 \leqslant n \leqslant N)$ so that the respective signal spaces are δ-unitarily equivalent, the respective normal operators $\Delta_n^k, \widetilde{\Delta}_n^k$ are ω-δ-close as well as bounded (in norm) by $K > 0$ and the connecting operators satisfy $\|\widetilde{P}_n J_{n-1} f - J_n P_n f\|_{\ell^2(\widetilde{G}_n)} \leqslant \delta \|f\|_{\ell^2(G_{n-1})}$. For the common module sequence Ω_N assume that the non-linearities satisfy $\|\rho_n(J_n f) - J_n \rho_n(f)\|_{\ell^2(\widetilde{G}_n^k)} \leqslant \delta \|f\|_{\ell^2(G_n^k)}$ and that the constants C_{χ_n} and $\{C_{g_{\gamma_n}}\}_{\gamma_n \in \Gamma_N}$ associated through Lemma 4.4 to the functions of the generalized frames in each layer satisfy $C_{\chi_n}^2 + \sum_{\gamma_n \in \Gamma_N} C_{g_{\gamma_n}}^2 \leqslant D^2$ for some $D > 0$. Denote the operator that the family $\{J_n\}_n$ of identification operators induce on $\mathscr{F}_N^k$ through concatenation by $\mathscr{J}_N : \mathscr{F}_N^k \to \widetilde{\mathscr{F}}_N^k$. Then we have

$$\|\widetilde{\Phi}_N^k(J_0 f) - \mathscr{J}_N \Phi_N^k(f)\|_{\widetilde{\mathscr{F}}_N^k} \leqslant K_N \cdot \delta \cdot \|f\|_{\ell^2(G^k)}, \quad \forall f \in \ell^2(G^k),$$

with $K_N = \sqrt{(8^N - 1)(2D^2 + 12)/7}$. If additionally $\|\widetilde{P}_n J_{n-1} f - J_n P_n f\|_{\ell^2(\widetilde{G}_n^k)} = 0$ or $\|\rho_n(J_n f) - J_n \rho_n(f)\|_{\ell^2(\widetilde{G}_n^k)} = 0$ holds in each layer, then we have $K_N = \mathcal{O}(4^{N/2})$; if both additional equations hold, we have $K_N = \mathcal{O}(2^{N/2})$.

Theorem K.4. Assuming admissibility, we have $\|\Psi_N^k(f) - \Psi_N^k(h)\|_{\mathscr{R}_N} \leqslant \|f - h\|_{\ell^2(G^k)}$ for all $f, h \in \ell^2(G)$. With the conditions and notation of Theorem K.2 we have

$$\|\Psi_N^k(f) - \widetilde{\Psi}_N^k(f)\|_{\mathscr{R}_N} \leqslant \sqrt{2(2^N - 1)} \cdot D \cdot \delta \cdot \|f\|_{\ell^2(G^k)}.$$

Additionally, in the setting of Theorem K.3, assuming that for each $n \leqslant N$ the identification operator J_n satisfies $\left|\|J_n f\|_{\ell^1(\widetilde{G}_n^k)} / \sqrt{\mu_{\widetilde{G}_n^k}} - \|f\|_{\ell^1(G_n^k)} / \sqrt{\mu_{G_n^k}}\right|, \left|\|J_n f\|_{\ell^r(\widetilde{G}_n^k)} - \|f\|_{\ell^r(G_n^k)}\right| \leqslant \delta \cdot K \cdot \|f\|_{\ell^2(G_n^k)}$ for $2 \leqslant r \leqslant p_n$ implies $(\forall f \in \ell^2(G^k))$

$$\|\widetilde{\Psi}_N(J_0 f) - \Psi_N(f)\|_{\mathscr{R}_N} \leqslant \sqrt{2} \cdot \sqrt{K_N^2 + K^2} \cdot \delta \cdot \|f\|_{\ell^2(G^k)}.$$

L Additional Details on Experiments

Here we provide additional details on utilized scattering architectures, training procedures, datasets and (performance of) other methods our approach is being compared to. Irrespective of task, our models are trained on an NVIDIA DGX A100 architecture utilizing between two and eight NVIDIA Tesla A100 GPUs with 80GB memory each. Running 10-fold cross validation for the respective experiments took at most 71 hours (which was needed for social network graph classification on REDDIT-12K).

L.1 Social Network Graph Classification

Datasets: The data we are working with is taken from [45]. In particular this work introduced six social network datasets extracted from from scientific collaborations (COLLAB), movie collaborations (IMDB-B, IMDB-M) and Reddit discussion threads (REDDIT-B, REDDIT-5K, REDDIT-12K).

Data is anonymised and contains no content that might be considered offensive. Each graph carries a class label, and the goal is to predict this label. Some basic properties of these datasets are listed in Table 3 below.

Table 3: Social Network Dataset Characteristics

Attributes:	COLLAB	IMDB- B	IMDB-M	REDDIT-B	REDDIT-5K	REDDIT-12K
Graphs	5K	1K	1.5K	2K	5K	12K
Nodes	372.5K	19.8K	19.5K	859.2K	2.5M	4.7M
Edges	49.1M	386.1K	395.6	4M	11.9M	21.8M
Maximum Degree	2k	540	352	12.2K	8K	12.2K
Minimum Degree	4	4	4	4	4	4
Average Degree	263	39	40	9	9	9
Target Labels	3	2	3	2	5	11
Disconnected Graphs	No	No	No	Yes	Yes	Yes

These datasets contain graph structures, however they don't contain associated weights or graph signals. Having unspecified weights simply means that the adjacency matrix W from which we construct the graph Laplacian

$$\mathcal{L} = D - W$$

on which our operator Δ is based simply has each entry corresponding to an edge set to unity. If no edge is present between vertices i and j, the entry W_{ij} is set to zero. It remains to solve the problem of the missing input signals. Our strategy is to generate signals reflecting the geometry of the underlying graph. We do this by utilizing features that associate to each node a number that characterizes its role or importance within its local environment or within the entire graph. We briefly describe them here:

1. **Degree**: The degree of a node is the number of edges incident at this node.

2. **Eccentricity**: For a connected graph, the eccentricity of a node is the maximum distance from this node to all other nodes. On a disconnected graph it is not defined.

3. **Clustering**: For unweighted graphs the clustering $c(u)$ of a node u is the fraction of possible triangles through that node that actually exist. It is calculated as

$$c(u) = \frac{2T(u)}{\deg(u)(\deg(u) - 1)}.$$

4. **Number of triangles**: The number of triangles containing the given node as a vertex.

5. **Core number**: A k-core is a maximal subgraph that only contains nodes of degree k or more. The core number of a node is the largest value k of a k-core containing that node.

6. **Clique number**: A clique is a subset of vertices of an undirected graph such that every two distinct vertices in the clique are adjacent. This input assigns the number of cliques the nodes participates in to each node.

7. **Pagerank**: This returns the PageRank of the respective nodes in the graph. PageRank computes a ranking of the nodes in the graph based on the structure of the edges. Originally it was designed as an algorithm to rank web pages.

For the first three datasets listed in Table 3 we utilize all listed input features. For the latter three datasets we have to refrain from using eccentricity as an input signal, as these datasets contain graphs that have multiple non-connected graph components.

Scattering Architecture: We chose a generalized scattering architecture of depth $N = 4$. As normal-operators, we utilize in each layer the un-normalized graph Laplacian $\mathcal{L} = D - W$ scaled by its largest eigenvalue ($\Delta = \mathcal{L}/\lambda_{max}(\mathcal{L})$). Filters are chosen as $\frac{1}{2}(\sin(\pi/2 \cdot \Delta), [\cos(\pi/2 \cdot \Delta) - \psi_\Delta \psi_\Delta^\top], \sin(\pi \cdot \Delta), [\cos(\pi \cdot \Delta) - \psi_\Delta \psi_\Delta^\top])$, which allows to specify the output generating function solely by demanding $\chi(0) = 1$ and $\chi(\lambda) = 0$ on all other eigenvalues of Δ. Here ψ_Δ is the normalized vector of all ones (satisfying $\Delta \psi_\Delta = 0$). Connecting operators are chosen as the identity, while we set $\rho_{n \geqslant 1}(\cdot) = |\cdot|$. This scattering architecture is then applied to each of these input signal

individually. For each input signal, this returns a feature vector with $1 + 4 + 16 + 64 = 85$ entries. These individual feature vectors are then concatenated into one final composite feature vector for each graph. Concerning applicable theoretical results, we note the following:

Let us first fix a single graph and let us assume the graph is connected. The associated graph Laplacian has zero as lowest lying eigenvalue. Denote the second lowest eigenvalue by $\lambda_2(\mathcal{L})$. The second lowest eigenvalue of our normal operator Δ is then given by $\lambda_2(\Delta) = \lambda_2(\mathcal{L})/\lambda_{max}(\mathcal{L}))$. Let χ be any function satisfying $\chi(\lambda) \leqslant 1$, $\chi(0) = 1$ and $\chi(\lambda) = 0$ if $\lambda \geqslant \lambda_2(\Delta)/2$. Our architecture then yields the same result as an architecture with χ as output generating function and filter bank given as $\frac{1}{2}(\sin(\pi/2 \cdot \Delta), \sqrt{\cos^2(\pi/2 \cdot \Delta) - \chi(\Delta)}, \sin(\pi \cdot \Delta), \sqrt{\cos^2(\pi \cdot \Delta) - \chi(\Delta)})$, where we interpret $\sqrt{\cos^2(\cdot)} = \cos(\cdot)$ (whenever $\chi = 0$). This is an architecture based on a generalized frame with frame constants $A = B = 1$ (as defined in Section 2). Checking conditions of the various theorems, we see that the resulting scattering transform is admissible and if it were extended to depth $N = \infty$ instead of being truncated at depth $N = 4$ would satisfy the conditions of Theorem 5.2 with $\eta_n = 0$. As connecting operators are chosen as the identity and we have $\Delta_n = \Delta$ for all $n \leqslant N$, this means that the conditions for Corollary 5.3 are also fulfilled. To check for stability under operator perturbations, we note that

$$\left[\cos(\pi/2 \cdot \Delta) - \psi_\Delta \psi_\Delta^\top\right] - \left[\cos(\pi/2 \cdot \widetilde{\Delta}) - \psi_{\widetilde{\Delta}} \psi_{\widetilde{\Delta}}^\top\right] = \cos(\pi/2 \cdot \Delta) - \cos(\pi/2 \cdot \widetilde{\Delta})$$

since $\psi_\Delta = \psi_{\widetilde{\Delta}}$. Thus we also have

$$\left\|\left[\cos(\pi/2 \cdot \Delta) - \psi_\Delta \psi_\Delta^\top\right] - \left[\cos(\pi/2 \cdot \widetilde{\Delta}) - \psi_{\widetilde{\Delta}} \psi_{\widetilde{\Delta}}^\top\right]\right\|_F \leqslant L_{\cos(\pi/2 \cdot)} \|\Delta - \widetilde{\Delta}\|_F.$$

Since the output generating filter is χ_Δ chosen to evaluate to one only on the lowest lying eigenvalue (i.e. zero, which is stable under perturbations) and to zero on higher lying eigenvalues, this introduces no additional instability. In total this implies that in the setting of Theorem 4.2 our architecture enjoys the same stability guarantees as one that is based on a filterbank given by $\frac{1}{2}(\sin(\pi/2 \cdot \Delta), \cos(\pi/2 \cdot \Delta), \sin(\pi \cdot \Delta), \cos(\pi \cdot \Delta))$ and an output generating function with negligible Lipschitz constant. Similar considerations apply in the vertex-set non-conserving setting.

Training Procedure: We train RBF kernel support vector classifiers on our composite scattering features. We fix $\epsilon = 0.1$. The hyperparameter γ scaling the exponent is chosen from

$$G_{\text{pool}} := \{0.00001, 0.0001, 0.001, 0.01, 0.1, 1, 10, 100\},$$

while we pick the C that controls the error our slack variables introduce among

$$C_{\text{pool}} := \{0.001, 0.01, 0.1, 1, 10, 25, 50, 100, 1000\}.$$

We could simply implement the training of the RBF-classifier on our composite scattering features by dividing each social-network dataset into 10 folds, then iteratively choosing one fold for testing and among the remaining 9 folds randomly choosing one for validation (i.e. for tuning the hyperparameters). Instead, following [12] (whose code is released under an Apache license and on which we partially built), we take a slightly different approach: We still randomly split our dataset into 10 folds. Among the 10 folds, we iteratively pick one for testing. Say we have picked the n^{th} fold for testing. Then there are 9 remaining folds. We iteratively pick the m_n^{th} (with $1 \leqslant m_n \leqslant 9$) of the remaining 9 folds for choosing hyperparameters. This leaves 8 folds on which we train our model for each choice of hyper parameter in $C_{\text{pool}} \times G_{\text{pool}}$. The resulting classifiers are all evaluated on the m_n^{th} fold. The one that performs best is retained as classifier m_n. As m_n varies between 1 and 9 (still for fixed n), this yields a set $\{f_{m_n} : 1 \leqslant m_n \leqslant 9\}$ of nine classifiers. From these we build the classifier f_n, whose classification result is obtained from a majority vote among the nine classifiers in $\{f_{m_n} : 1 \leqslant m_n \leqslant 9\}$. Then we evaluate the performance of f_n on the n^{th} fold to obtain the n^{th} estimation of how well our model performs. As n varies from one to ten, we built the mean and variance of the performances of the classifiers f_n on the n^{th} fold expressed as the percentage of correct classifications.

Reference Methods: To allow for a comparison of our results to the literature, typical classification accuracies for graph algorithms on social network datasets are displayed in Table 1. Following the standard format of reporting classification accuracies, they are presented in the format (Accuracy $\pm$

standard deviation). If results are not reported for a dataset, we denote this as not available (N/A). The first three rows of Table 1 display results for graph kernel methods; namely Weisfeiler-Lehman graph kernels (WL, [34]), Graphlet kernels (Graphlet, [35]) and deep graph kernels (DGK, [44]). The subsequent rows display results for geometric deep learning algorithms: Deep graph convolutional neural networks (DGCNN,[48]), Patchy-san (PSCN (with k=10), [27]), recurrent neural network autoencoders (S2S-N2N-PP, [16]) and graph isomorphism networks (GIN [43]). These results are taken from [12]. Additionally we compare with P-Poinc [20], which embeds nodes into a hyperbolic space (the Poincare ball, to be precise), GSN-e [3] which combines message passing with structural features extracted via subgraph isomorphism and WKPI-kC [49] which utilizes a weighted kernel within its metric learning framework. The second to last row (GS-SVM [12]) provides a result that is also based on a method that combines a static scattering architecture with a support vector machine. Its filters are based on graph wavelets built from differences between lazy random walks that have propagated at different time scales.

L.2 Regression of Quantum Chemical Energies

Dataset: The dataset we consider is the QM7 dataset, introduced in [2, 32]. This dataset contains descriptions of 7165 organic molecules, each with up to seven heavy atoms, with all non-hydrogen atoms being considered heavy. A molecule is represented by its Coulomb matrix C^{Clmb}, whose off-diagonal elements

$$C_{ij}^{\text{Clmb}} = \frac{Z_i Z_j}{|R_i - R_j|} \tag{8}$$

correspond to the Coulomb-repulsion between atoms i and j, while diagonal elements encode a polynomial fit of atomic energies to nuclear charge [32]:

$$C_{ii}^{\text{Clmb}} = \frac{1}{2} Z_i^{2.4}$$

For each atom in any given molecular graph, the individual Cartesian coordinates R_i and the atomic charge Z_i are also accessible individually. To each molecule an atomization energy - calculated via density functional theory - is associated. The objective is to predict this quantity, the performance metric is mean absolute error. Numerically, atomization energies are negative numbers in the range -600 to -2200. The associated unit is [*kcal/mol*].

Scattering Architecture: Off-diagonal entries in the Coulomb Matrix clearly represent an inverse distance. A weight of zero can then heuristically be thought of as the inverse distance between two infinitely separated atoms. After calculating the degree matrix D associated to C, we obtain the graph Laplacian once more as $\mathcal{L} = D - C$ and set our normal operator to

$$\Delta = \frac{\mathcal{L}}{\lambda_{\max}(\mathcal{L})}.$$

If we continuously vary the distances in (8), staying clear of zero, then the adjacency matrix and hence the graph Laplacian $\mathcal{L}$ varies continuously. As long as we avoid complete degeneracy, the largest eigenvalue $\lambda_{\max}(\mathcal{L})$ will remain positive. This implies that our normal operator Δ varies continuously under changes of the inter-atomic distances, which implies that our feature vector also varies continuously, as distances are changed. Connecting operators are set to the identity, while non-linearities are fixed to $\rho_{n \geqslant 1}(\cdot) = |\cdot|$. Filters are chosen as $(\sin(\pi/2 \cdot \Delta), \cos(\pi/2 \cdot \Delta), \sin(\pi \cdot \Delta), \cos(\pi \cdot \Delta))$ acting through matrix multiplication. The output generating functions are set to the identity as well. Graph level features are aggregated via the map $N_5^{G^2}(\cdot)$ of Section 7; slightly modified to neglect the normalizing factor in the first entry for improved convenience in numerical implementability. As weights μ_{ij} for our second-order feature space are set to unity and molecular graphs in QM7 contain at most 23-molecules, we note that $\sqrt{\mu_{G^2}} \leqslant \sqrt{23^2} = 23$. Going through the proofs of our graph-level stability results, we see that they remain valid after multiplying each stability constant by 23. The Coulomb matrix (divided by a factor of 10 as this empirically improved performance) is then also utilized as an edge level input signal. Node level features are obtained by applying the above architecture to the node level information provided by the respective atomic charges $\{Z_i\}$ on each graph. We aggregate to graph level features using N_5^G (cf. Section 6.1), again

neglecting the normalizing factor in the first entry for improved convenience in implementing. The network depth is set to $N = 4$ in both cases. We then concatenate graph level features obtained from node- and edge level input into a composite scattering feature vector.

Training Procedure: The QM7 dataset comes with a precomputed partition into five subsets; each containing a representative amount of heavy and light molecules covering the entire complexity range of QM7. To allow for 10-fold cross validation, we further dissect each of these subsets into two smaller datasets, one containing graphs indexed by an even number, one containing graphs indexed by an odd number. On these 10-subsets, we then perform 10-fold cross validation. Among the 10 folds, we iteratively pick one for testing. Say we have picked the n^{th} fold for testing. Then there are 9 remaining folds. We iteratively pick the m_n^{th} (with $1 \leqslant m_n \leqslant 9$) of the remaining 9 folds for choosing hyperparameters. This leaves 8 folds on which we train our model for each choice of hyper parameter in $C_{\text{pool}} \times G_{\text{pool}}$. This yields 8 regression models, which we average to built our final predictor for the n^{th} run. This mean absolute error of this predictor is then evaluated on the n^{th} fold which was retained for testing. As n varies from one to ten, we built the mean and variance of the performances of the generated regression models. To reduce training and evaluation time, we simplified our sets of hyperparametes to only consist of single values. We have set $G_{\text{pool}} = \{0.00003\}$ and $G_{\text{pool}} = \{400000\}$ which were found to perform well enough.

Reference Methods: We comprehensively evaluate our method against 11 popular baselines and state of the art approaches. Among these methods are graph convolutional methods such as GraphConv [18], Weave [17] or SchNet [33]. MPNN [13] and its variant DMPNN [46] are models considering edge features during message passing. AttentiveFP [42] is an extension of the graph attention framework, while N-Gram [22] is a pretrained method. Results for these methods as well as for GROVER are taken from [31]. PhysChem [47] learns molecular representations via fusing physical and chemical information. Deep Tensor Neural Networks (DTNN [41]) are adaptable extensions of the Coulomb Matrix featurizer mapping atom numbers to trainable embeddings which are then updated based on distance information and other (node-level) atomic features. Finally Path-Augmented Graph Transformer Networks (PATGN, [6]) exploit the connectivity structure of the data in a global attention mechanism.