Distribution-free inference for regression: discrete, continuous, and in between

Abstract

In data analysis problems where we are not able to rely on distributional assumptions, what types of inference guarantees can still be obtained? Many popular methods, such as holdout methods, cross-validation methods, and conformal prediction, are able to provide distribution-free guarantees for predictive inference, but the problem of providing inference for the underlying regression function (for example, inference on the conditional mean $\mathbb{E}\left[Y|X\right]$) is more challenging. In the setting where the features X are continuously distributed, recent work has established that any confidence interval for $\mathbb{E}\left[Y|X\right]$ must have non-vanishing width, even as sample size tends to infinity. At the other extreme, if X takes only a small number of possible values, then inference on $\mathbb{E}\left[Y|X\right]$ is trivial to achieve. In this work, we study the problem in settings in between these two extremes. We find that there are several distinct regimes in between the finite setting and the continuous setting, where vanishing-width confidence intervals are achievable if and only if the effective support size of the distribution of X is smaller than the square of the sample size.

1 Introduction

Consider a regression problem, where our aim is to model the distribution of a response variable $Y \in \mathbb{R}$ based on the information carried by features $X \in \mathcal{X}$. Given training data $(X_1, Y_1), \ldots, (X_n, Y_n)$, we aim to build a fitted model to estimate the conditional distribution of $Y \mid X$, or some summary of this distribution such as the conditional mean or conditional median. In this type of setting, our goals are to simultaneously perform two tasks, estimation and inference—that is, we want to accurately estimate the conditional distribution, and we also want a reliable way of quantifying our uncertainty about this estimate.

To make this concrete, suppose the training data $\{(X_i, Y_i)\}$ are drawn i.i.d. from some unknown distribution P on $\mathbb{R}^d \times \mathbb{R}$, and we want to estimate the true conditional mean, $\mu_P(x) := \mathbb{E}[Y|X = x]$, of this distribution. Given the training data, we construct a fitted regression function $\widehat{\mu} : \mathbb{R}^d \to \mathbb{R}$ using any algorithm, for instance, a parametric method such as least squares or a nonparametric procedure such as a Gaussian kernel method. For many regression algorithms, assuming certain conditions on the underlying distribution P will ensure an accurate estimate of μ_P; however, unless we are able to verify these assumptions, we cannot be confident that the corresponding error rates will indeed lead to a valid confidence interval for μ_P. The goal of *distribution-free inference* is to provide inference guarantees—in this case, confidence intervals for $\mu_P(X_{n+1})$ at a newly observed feature vector X_{n+1}—that are valid universally over any underlying distribution P.

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

1.1 Our contributions

In this work, we study the problem of constructing a confidence interval $\widehat{C}_n$ that satisfies the following property:

Definition 1. *An algorithm $\widehat{C}_n$ provides a distribution-free $(1 - \alpha)$-confidence interval for the conditional mean if it holds that*

$$\mathbb{P}_{(X_i, Y_i) \overset{\text{iid}}{\sim} P} \left\{ \mu_P(X_{n+1}) \in \widehat{C}_n(X_{n+1}) \right\} \geq 1 - \alpha \text{ for all distributions } P \text{ on } (X, Y) \in \mathbb{R}^d \times [0, 1].$$

Recent work by Barber [2020], Gupta et al. [2020], Medarametla and Candès [2021] proves that distribution-free coverage properties similar to Definition 1 lead to fundamental limits on the accuracy of inference—if we compute a distribution-free confidence interval $\widehat{C}_n$ on data drawn from a distribution P where the marginal distribution P_X is *nonatomic* (meaning that there are no point masses, i.e., $\mathbb{P}_{P_X} \{X = x\} = 0$ for all points $x \in \mathbb{R}^d$), then $\widehat{C}_n(X_{n+1})$ cannot have vanishing length as sample size n tends to infinity, regardless of the smoothness of P, or any other "nice" properties of this distribution.

Our new results examine the possibility of constructing confidence intervals $\widehat{C}_n$ that are both distribution-free (satisfy Definition 1) and have vanishing length, when P_X may be discrete, nonatomic, or a mixture of the two. We find that the hardness of this problem can be characterized by the *effective support size* of P_X—essentially, how many points $x \in \mathbb{R}^d$ are needed to capture most of the mass of P_X (for example, if P_X is uniform over M points, then its effective support size is $\mathcal{O}(M)$).

Our main theoretical results show that there are two regimes. If the effective support size is $\gg n^2$, then P_X essentially behaves like a nonatomic distribution because in a sample of size n, with high probability all the X values are only observed once; in this regime, we find that the average length of $\widehat{C}_n(X_{n+1})$ is bounded away from zero, i.e., no distribution-free confidence interval can have vanishing length. If instead the effective support size is $\ll n^2$, then it becomes possible for $\widehat{C}_n(X_{n+1})$ to have vanishing length, and in particular, the minimum possible length scales as $\frac{M^{1/4}}{n^{1/2}}$ for effective support size M—interestingly, vanishing length is possible even when M is larger than n, meaning that precise distribution-free inference for $\mathbb{E}[Y|X]$ is possible even if most X values were never observed in the training set.

1.2 Related work

The problem of distribution-free inference has been studied extensively in the context of *predictive inference*, where the goal is to provide a confidence band for the response value Y_{n+1} given a new feature vector X_{n+1}. The prediction problem is fundamentally different from the goal of covering the conditional mean—in particular, by splitting the data and using a holdout set, we can empirically validate the coverage level of any constructed predictive band. Methods such as conformal prediction (see, e.g., Vovk et al. [2005], Papadopoulos et al. [2002], Lei et al. [2018], Vovk et al. [2018]) or jackknife+ [Barber et al., 2021, Kim et al., 2020] can ensure valid distribution-free predictive inference without the need to split the data set (thus avoiding reducing the sample size).

The problem of *distribution-free coverage for the conditional mean or median* of Y given X has been studied by Barber [2020], Gupta et al. [2020] (for the mean) and Medarametla and Candès [2021] (for the median). As mentioned above, these works establish that, if the marginal distribution P_X of the feature vector X is nonatomic, then it is impossible for a distribution-free confidence interval $\widehat{C}_n$ to have vanishing length even as $n \to \infty$; in particular, if $\widehat{C}_n$ is guaranteed to cover the mean or median for all distributions P, then it also must be a predictive interval, that is, it must cover Y itself (which implies its length cannot be vanishing, since Y is inherently noisy). These impossibility results are connected to earlier results on the impossibility of adaptation to smoothness, in the nonparametric inference literature—specifically, if μ_P is β-Hölder smooth, then it is possible to build a confidence interval of length $\mathcal{O}(n^{-\frac{\beta}{2\beta+d}})$ if β is known (e.g., using k-nearest-neighbors with an appropriately chosen k), but this cannot be achieved when β is unknown (see, e.g., Giné and Nickl [2016, Section 8.3] for an overview of results of this type).

While the results above establish the challenges for distribution-free inference when the features X are nonatomic, at the other extreme we can consider scenarios where X has a discrete distribution. In

this setting, the problem of estimating μ_P is related to the *discrete distribution testing*, where the aim is to test properties of a discrete distribution—for instance, we might wish to test equality of two distributions where we draw samples from each [Chan et al., 2014, Acharya et al., 2014, Diakonikolas and Kane, 2016, Canonne et al., 2015]; to test whether a sample is drawn from a known distribution P or not [Diakonikolas and Kane, 2016, Acharya et al., 2015, Valiant and Valiant, 2017, Diakonikolas et al., 2018], or drawn from any distribution belonging to a class $\mathcal{P}$ or not [Acharya et al., 2015, Canonne et al., 2018]; or to estimate certain characteristics of a distribution based on a sample, such as its entropy or its support size [Valiant and Valiant, 2011b,a, Acharya et al., 2014]. The methods developed in this literature are closely related to the problem of distribution-free confidence intervals, as we will see later on when we construct upper bounds in Section 3.

2 Main results: lower bound

Before presenting our main result, we begin with several definitions. For any distribution P_X on $X \in \mathcal{R}^d$ (i.e., the marginal distribution of X under the joint distribution P on (X, Y)), we first define the *effective support size* of P_X at tolerance level $\gamma \in [0, 1)$:

$$M_\gamma(P_X) = \min \left\{ |\mathcal{X}| \ : \ \mathcal{X} \subset \mathbb{R}^d \text{ and } \mathbb{P}_{P_X} \{ X \in \mathcal{X} \} \geq 1 - \gamma \right\},$$

where $|\mathcal{X}|$ denotes the cardinality of the set $\mathcal{X}$. In particular, if P_X is a distribution supported on M points, then $M_\gamma(P_X) \leq M$ for any γ. If instead P_X Is nonatomic, then $M_\gamma(P_X) = \infty$ for all γ.

Next, for any distribution P on $(X, Y) \in \mathbb{R}^d \times [0, 1]$, we define

$$\sigma_{P,\beta}^2 = \text{ the } \beta\text{-quantile of } \text{Var}_P(Y|X), \text{ under the distribution } X \sim P_X.$$

With these definitions in place, the next result establishes a lower bound on the length of any distribution-free confidence interval $\widehat{C}_n$.

Theorem 1. *Fix any $\alpha > 0$, and let $\widehat{C}_n$ be a distribution-free $(1 - \alpha)$-confidence interval (i.e., satisfying Definition 1). Then for any distribution P on $\mathbb{R}^d \times \mathbb{R}$, for any $\beta > 0$ and $\gamma > \alpha + \beta$,*

$$\mathbb{E}\left[\text{Leb}\left(\widehat{C}_n(X_{n+1}) \right) \right] \geq \tfrac{1}{3}\sigma_{P,\beta}^2 (\gamma - \alpha - \beta)^{1.5} \cdot \min\left\{ \frac{\left(M_\gamma(P_X)\right)^{1/4}}{n^{1/2}}, 1 \right\},$$

where the expected value is taken over data points $(X_i, Y_i) \overset{\text{iid}}{\sim} P$, for $i = 1, \ldots, n + 1$.

Here Leb denotes the Lebesgue measure on $\mathbb{R}$, and so $\mathbb{E}\left[\text{Leb}\left(\widehat{C}_n(X_{n+1}) \right) \right]$ measures the expected length of the confidence interval constructed at a new test point X_{n+1}.

2.1 Special cases

To help interpret this result, we now examine its implications in several special cases.

Uniform discrete features If P_X is a uniform distribution over M points, then for any $\gamma > 0$ the effective support size is $M_\gamma(P_X) = \lceil (1 - \gamma)M \rceil$. Therefore, Theorem 1 implies that for any P with nonatomic marginal P_X,

$$\mathbb{E}\left[\text{Leb}\left(\widehat{C}_n(X_{n+1}) \right) \right] \geq \tfrac{1}{3}\sigma_{P,\beta}^2 (\gamma - \alpha - \beta)^{1.5} (1 - \gamma)^{0.25} \cdot \min\left\{ \frac{M^{1/4}}{n^{1/2}}, 1 \right\}$$

for any $\beta \in (0, \gamma - \alpha)$. In particular, we see that $M \gg n^2$ implies a *constant* lower bound on the width of any distribution-free confidence interval, while $M \ll n^2$ allows for the possibility of a *vanishing* width for a distribution-free confidence interval.

Binary response If the response Y is known to be binary (i.e., $Y \in \{0, 1\}$), we can consider relaxing the requirement of distribution-free coverage to only include distributions of this type—that is, we require

$$\mathbb{P}_{(X_i, Y_i) \overset{\text{iid}}{\sim} P} \left\{ \mu_P(X_{n+1}) \in \widehat{C}_n(X_{n+1}) \right\} \geq 1 - \alpha \text{ for all distributions } P \text{ on } \mathbb{R}^d \times \{0, 1\}. \quad (1)$$

(Compare to Definition 1, where the coverage property is required to hold for all distributions P on $\mathbb{R}^d \times [0,1]$, i.e., for a broader class of distributions.) However, it turns out that relaxing the requirement does not improve the lower bound. Specifically, suppose we have an algorithm to construct a confidence interval $\widehat{C}_n$ satisfying (1) (but not required to satisfy Definition 1). We can easily convert $\widehat{C}_n$ into a method that does satisfy Definition 1—given data $(X_1, Y_1), \ldots, (X_n, Y_n)$, for each $i = 1, \ldots, n$ draw a binary response $\tilde{Y}_i \sim \text{Bernoulli}(Y_i)$. Then we clearly have n i.i.d. draws from a distribution on $(X, \tilde{Y}) \in \mathbb{R}^d \times \{0,1\}$, where $\mathbb{E}[\tilde{Y} \mid X] = \mathbb{E}[Y \mid X] = \mu_P(X)$. After running our algorithm to construct $\widehat{C}_n$ on the new data $(X_1, \tilde{Y}_1), \ldots, (X_n, \tilde{Y}_n)$, the binary distribution-free coverage property (1) satisfied by $\widehat{C}_n$ ensures that this modified procedure satisfies Definition 1.

To summarize, then, we see that the problem of distribution-free coverage is equally hard for the binary response case ($Y \in \{0,1\}$) as for the more general bounded response case ($Y \in [0,1]$).

Nonatomic features We now consider the setting where the distribution of X is nonatomic, i.e., $\mathbb{P}\{X = x\} = 0$ for all x. (In particular, this includes the continuous case, where X has a continuous distribution on $\mathbb{R}^d$.) In this case, for any $\gamma > 0$ the effective support size is $M_\gamma(P_X) = \infty$. Therefore, Theorem 1 implies that for any P with nonatomic marginal P_X,

$$\mathbb{E}\left[\text{Leb}\left(\widehat{C}_n(X_{n+1})\right)\right] \geq \tfrac{1}{3}\sigma_{P,\beta}^2(1 - \alpha - \beta)^{1.5}$$

for any $\beta \in (0, 1 - \alpha)$. In particular, this lower bound does not depend on n, and so the width of any distribution-free confidence interval is non-vanishing even for arbitrarily large sample size n (as long as $\sigma_{P,\beta}^2$ is positive).

In case of a binary response, where P is a distribution on $\mathbb{R}^d \times \{0,1\}$ with nonatomic marginal distribution P_X, Barber [2020] establishes that any distribution-free confidence interval for μ must satisfy a lower bound, which is a function only of P and does not depend on n (and, in particular, does not vanish as $n \to \infty$). In this sense, our new result can be viewed as a generalization of this work, since the nonvanishing minimum length for nonatomic P_X is a consequence of our result.

2.2 Adding knowledge of P_X

One way we might try to weaken the notion of distribution-free coverage would be to allow assumptions about the marginal distribution P_X, while remaining assumption-free for the function μ_P determining the conditional mean. In other words, we might weaken Definition 1 to require coverage over all distributions P for which $P_X = P_X^*$, for a known marginal distribution P_X^* (or, all distributions P for which P_X satisfies some assumed property). Interestingly, the lower bound in Theorem 1 remains the same even under this much milder definition of validity—we will see in the proof that knowledge of P_X does not affect the lower bound, since the argument relies only on our uncertainty about the conditional distribution of $Y|X$.

2.3 Bounded or unbounded?

The lower bound established in Theorem 1 assumes distribution-free coverage for distributions with a bounded response Y—that is, Definition 1 requires coverage to hold for distributions where the response Y is supported on $[0,1]$ (although no other assumptions are placed on P). Would it be possible for us to instead consider the general case, where P is an unknown distribution on $\mathbb{R}^d \times \mathbb{R}$? The following result shows that this more general question is not meaningful:

Proposition 1. *Suppose an algorithm $\widehat{C}_n$ satisfies*

$$\mathbb{P}_{(X_i, Y_i) \overset{\text{iid}}{\sim} P}\left\{\mu_P(X_{n+1}) \in \widehat{C}_n(X_{n+1})\right\} \geq 1 - \alpha \text{ for all distributions } P \text{ on } \mathbb{R}^d \times \mathbb{R}.$$

Then for all distributions P, for all $y \in \mathbb{R}$ it holds that

$$\mathbb{P}\left\{y \in \widehat{C}_n(X_{n+1}))\right\} \geq 1 - \alpha,$$

where the probability is calculated with respect to data points $(X_i, Y_i) \overset{\text{iid}}{\sim} P$, for $i = 1, \ldots, n+1$.

This means that if we require $\widehat{C}_n$ to have distribution-free coverage over distributions with *unbounded* response, we will inevitably return confidence intervals that contain *every point in the real line* a substantial portion of the time. (In particular, we will of course have infinite expected width, $\mathbb{E}\left[\text{Leb}\left(\widehat{C}_n(X_{n+1})\right)\right] = \infty$.) For this reason, we therefore restrict our attention to the setting where the response Y takes values in $[0, 1]$, since clearly an unbounded Y cannot result in any meaningful distribution-free inference.

3 Main results: upper bound

We next construct an algorithm that, for certain "nice" distributions P, can achieve a confidence interval length that matches the rate of the lower bound. In order to run the algorithm, we will need two main ingredients as input:

1. A hypothesized support set $\{x^{(1)}, x^{(2)}, \dots\} \subset \mathbb{R}^d$ for the marginal P_X, and

2. A hypothesized mean function $\mu : \mathbb{R}^d \to [0, 1]$.

We emphasize that the coverage guarantee provided by our method does not rely in any way on the accuracy of these initial guesses—the constructed confidence interval will satisfy distribution-free validity (Definition 1) even if these initial parameters are chosen in a completely uninformed way. However, the length of the resulting confidence interval will be affected; high accuracy in these initial guesses can be expected to result in a shorter confidence interval. In particular, the hypothesized support set $\{x^{(1)}, x^{(2)}, \dots\}$ should aim to list the highest-probability values of X early in the list, while the hypothesized mean function μ should aim to be as close to the true conditional mean μ_P as possible. (Our theoretical results below will make these goals more precise.)

Given the hypothesized support and hypothesized mean function, to run our algorithm, we first choose parameters $\gamma, \delta > 0$ satisfying $\gamma + \delta < \alpha$, and then compute the following steps.

- **Step 1: estimate the effective support size.** First, we compute an upper bound on the support size needed to capture γ of the probability under P_X:

$$\widehat{M}_\gamma = \min\left\{ m : \sum_{i=1}^n \mathbb{1}\left\{ X_i \in \{x^{(1)}, \dots, x^{(m)}\} \right\} \geq (1-\gamma)n + \sqrt{\frac{n \log(2/\delta)}{2}} \right\}.$$

Applying the Hoeffding inequality to the $\text{Binom}(n, \gamma)$ distribution, we see that

$$\mathbb{P}\left\{ \widehat{M}_\gamma \geq M_\gamma(P_X) \right\} \geq 1 - \delta/2. \tag{2}$$

- **Step 2: estimate error at each repeated X value.** Next, for each $m = 1, 2, \dots$, let $n_m = \sum_{i=1}^n \mathbb{1}\left\{ X_i = x^{(m)} \right\}$ denote the number of times $x^{(m)}$ was observed, and let

$$N_{\geq 2} = \sum_{m=1}^\infty \mathbb{1}\left\{ n_m \geq 2 \right\} \tag{3}$$

be the number of X values observed at least twice. For each m with $n_m \geq 2$, let $\bar{y}_m = \frac{1}{n_m} \sum_{i=1}^n Y_i \cdot \mathbb{1}\left\{ X_i = x^{(m)} \right\}$ and $s_m^2 = \frac{1}{n_m-1} \sum_{i=1}^n (Y_i - \bar{y}_m)^2 \cdot \mathbb{1}\left\{ X_i = x^{(m)} \right\}$ be the sample mean and sample variance of the corresponding Y values. Define

$$Z = \sum_{\substack{m=1,2,\dots \\ \text{s.t. } n_m \geq 2}} (n_m - 1) \cdot \left((\bar{y}_m - \mu(x^{(m)}))^2 - n_m^{-1} s_m^2 \right). \tag{4}$$

This construction is inspired by analogous statistics appearing in the literature for testing discrete distributions—for instance, the work of Chan et al. [2014]. To see the intuition behind this construction, we observe that $\{Y_i : X_i = x^{(m)}\}$ is a collection of i.i.d. observations with mean $\mu_P(x^{(m)})$, and therefore, conditional on n_m (with $n_m \geq 2$), we have

$$\mathbb{E}\left[\bar{y}_m\right] = \mu_P(x^{(m)}) \text{ and } \text{Var}\left(\bar{y}_m\right) = n_m^{-1}\mathbb{E}\left[s_m^2\right],$$

and therefore $\mathbb{E}\left[(\bar{y}_m - \mu(x^{(m)}))^2 - n_m^{-1} s_m^2\right] = (\mu(x^{(m)}) - \mu_P(x^{(m)}))^2$ is an estimate of our error at this X value.

- **Step 3: construct the confidence interval.** Finally, we define our confidence interval as follows. Let

$$\widehat{\Delta} = \sqrt{\frac{2\widehat{M_\gamma} + n}{n(n-1)}} \cdot \sqrt{4Z_+ + 4\sqrt{N_{\geq 2} \cdot 2/\delta} + 16/\delta},$$

where Z_+ denotes $\max\{Z, 0\}$. Then for each $x \in \mathbb{R}^d$, we define

$$\widehat{C}_n(x) = \left[\max\left\{0, \mu(x) - \frac{\widehat{\Delta}}{\alpha - \delta - \gamma}\right\}, \min\left\{1, \mu(x) + \frac{\widehat{\Delta}}{\alpha - \delta - \gamma}\right\}\right]. \quad (5)$$

(In practice, we might split the data into two sets of size n, using half of the data to construct our estimate μ and/or to construct the hypothesized support, and then using the remaining data to compute the confidence interval.)

With this construction in place, we now verify that it yields a valid distribution-free confidence interval.

Theorem 2. *The confidence interval constructed in* (5) *is a distribution-free* $(1 - \alpha)$-*confidence interval (i.e., $\widehat{C}_n$ satisfies Definition 1).*

Next, we will see how this construction is able to match the rate of the lower bound established in Theorem 1—specifically, in a scenario where the hypothesized support set and mean function are "chosen well", i.e., are a good approximation to the properties of the true distribution P. For interpretability and intuition, we only consider a simple case—where the marginal P_X is approximately uniform over some finite subset of the hypothesized support, and the hypothesized function μ has uniformly bounded error.

Theorem 3. *Suppose the distribution P on $(X, Y) \in \mathbb{R}^d \times \mathbb{R}$ has marginal P_X that is supported on $\{x^{(1)}, \ldots, x^{(M)}\}$ and satisfies $\mathbb{P}_{P_X}\left\{X = x^{(m)}\right\} \leq \eta/M$ for all m, and suppose that P has conditional mean $\mu_P : \mathbb{R}^d \to \mathbb{R}$ that satisfies*

$$\left|\mu_P(X) - \mu(X)\right| \leq \text{err}_\mu \text{ almost surely over } X \sim P_X.$$

Then the confidence interval constructed in (5) *satisfies*

$$\mathbb{E}\left[\text{Leb}(\widehat{C}_n)\right] \leq c\left(\text{err}_\mu + \frac{M^{1/4}}{n^{1/2}}\right),$$

where c depends only on the parameters α, δ, γ, η.

4 Discussion

Our main result, Theorem 1, shows that the problem of constructing distribution-free confidence intervals for a conditional mean has hardness characterized by the effective support size $M_\gamma(P_X)$ of the feature distribution; distribution-free confidence intervals may have vanishing length if the sample size is at least as large as the square root of the effective support size, but must have length bounded away from zero if the sample size is smaller. The rate of the lower bound on length, scaling as $\min\{\frac{(M_\gamma(P_X))^{1/4}}{n^{1/2}}, 1\}$, is achievable in certain settings—Theorems 2 and 3 establish that distribution-free confidence intervals may achieve this length if we have a good hypothesis μ for μ_P. Of course, the specific construction used for these matching bounds may not be optimal—both in terms of constant factors that may inflate its length, and in terms of the range of settings in which it is able (up to constants) to match the lower bound. Improving this construction to provide a practical and accurate algorithm is an important question for future work.

More broadly, all inference methods must inherently involve a tradeoff between the strength of the guarantees, and the precision of the resulting answers. In this present work, we consider a universally strong guarantee (i.e., coverage of the conditional mean for *all* distributions P), which results in precise inference (i.e., vanishing-length confidence intervals) for only *some* distributions P, namely, those with effective support size $\ll n^2$. This tradeoff may not be desirable in practice, since in an applied setting we might instead prefer to relax the required coverage properties for more challenging distributions P in order to allow for more precise answers; an overly strict definition of validity may

inhibit our ability to extract meaningful information from robust inference methods. In practice, we may be satisfied with a validity condition that yields weaker guarantees in a nonatomic setting, but still yields the stronger coverage guarantee in the achievable regime where $M_\gamma(P_X) \ll n^2$. In future work, we aim to study whether this more adaptive type of validity definition, which is weaker than distribution-free coverage, may enables us to build confidence intervals that have vanishing length even in the nonatomic setting.

A Proofs

In this section we will prove Proposition 1 and Theorem 1. The proofs of Theorems 2 and 3, as well as of all the supporting lemmas appearing in this section, are given in the Supplementary Materials.

A.1 Proof of Proposition 1

The proof is simple, and follows a standard argument—we will need to consider replacing P with a distribution that places vanishing probability on some extremely large value. Fix any distribution P, and any $y \in \mathbb{R}$. For any fixed $\epsilon > 0$, define a new distribution Q as follows:

$$\text{Draw } X \sim P_X, \text{ then draw } Y|X \sim (1-\epsilon)P_{Y|X} + \epsilon\delta_{\epsilon^{-1}y - (\epsilon^{-1}-1)\mu_P(X)},$$

where $P_{Y|X}$ is the conditional distribution of $Y|X$ under P, and δ_t denotes the point mass at t. Then we can trivially calculate that $d_{\text{TV}}(P^n \times P_X, Q^n \times Q_X) \le n\epsilon$. Therefore,

$$\mathbb{P}_{P^n \times P_X}\left\{y \in \widehat{C}_n(X_{n+1})\right\} \ge \mathbb{P}_{Q^n \times Q_X}\left\{y \in \widehat{C}_n(X_{n+1})\right\} - n\epsilon.$$

On the other hand, the distribution Q has conditional mean

$$\mu_Q(x) = (1-\epsilon)\mu_P(x) + \epsilon\left(\epsilon^{-1}y - (\epsilon^{-1}-1)\mu_P(x)\right) = y,$$

and so the conditional mean $\mu_Q(X_{n+1})$ is equal to y almost surely. Therefore,

$$\mathbb{P}_{Q^n \times Q_X}\left\{y \in \widehat{C}_n(X_{n+1})\right\} = \mathbb{P}_{Q^n \times Q_X}\left\{\mu_Q(X_{n+1}) \in \widehat{C}_n(X_{n+1})\right\} \ge 1 - \alpha,$$

where the last step holds since $\widehat{C}_n$ must satisfy distribution-free coverage and, therefore, must satisfy coverage with respect to Q. Since $\epsilon > 0$ is arbitrarily small, this completes the proof.

A.2 Proof of Theorem 1

To prove the theorem, we will need several supporting lemmas:

Lemma 1. *Let Q be any distribution on $[0,1]$ with variance σ^2. Then we can write Q as a mixture of two distributions Q_0, Q_1 on $[0,1]$ such that*

$$Q = 0.5Q_0 + 0.5Q_1 \text{ and } \mathbb{E}_{Q_1}[X] - \mathbb{E}_{Q_0}[X] \ge 2\sigma^2.$$

Lemma 2. *Let P_X be any distribution on $\mathbb{R}^d$, and let $\mathbb{R}^d = \mathcal{X}_1 \cup \mathcal{X}_2 \cup \ldots$ be a fixed partition. Define a distribution P_0 on $(X, Z) \in \mathbb{R}^d \times \{0, 1\}$ as:*

$$\text{Draw } X \sim P_X, \text{ and draw } Z \sim \text{Bernoulli}(0.5), \text{ independently from } X.$$

For any fixed sequence $a = (a_1, a_2, \dots)$ of signs $a_1, a_2, \dots \in \{\pm 1\}$, and any fixed $\epsilon_1, \epsilon_2, \dots \in [0, 0.5]$, define a distribution P_a on on $(X, Z) \in \mathbb{R}^d \times \{0, 1\}$ as:

$$\text{Draw } X \sim P_X, \text{ and conditional on } X, \text{ draw } Z|X \in \mathcal{X}_m \sim \text{Bernoulli}(0.5 + a_m \cdot \epsilon_m).$$

Finally define $\tilde{P}_0 = (P_0)^n$ (i.e., n i.i.d. draws from P_0), and define $\tilde{P}_1$ as the mixture distribution:

- *Draw $A_1, A_2, \dots \overset{\text{iid}}{\sim} \text{Unif}\{\pm 1\}$.*

- *Conditional on $A_1, A_2, \dots$, draw $(X_1, Z_1), \dots, (X_n, Z_n) \overset{\text{iid}}{\sim} P_A$.*

Then

$$d_{\mathrm{TV}}(\tilde{P}_0, \tilde{P}_1) \leq 2n \sqrt{\sum_{m=1}^{\infty} \epsilon_m^4 \cdot \mathbb{P}_{P_X} \{X \in \mathcal{X}_m\}^2}.$$

We are now ready to prove the theorem. Define $\mathcal{X}_1 = \{x \in \mathbb{R}^d : \mathbb{P}_{P_X} \{X = x\} > \frac{1}{M_\gamma(P_X)}\}$. We must have $|\mathcal{X}_1| < M_\gamma(P_X)$ since P_X is a probability measure, and therefore, by definition of the effective support size, we must have $\mathbb{P}_{P_X} \{X \in \mathcal{X}_1\} < 1 - \gamma$. On the set $\mathbb{R}^d \backslash \mathcal{X}_1$, any point masses of the distribution P_X must each have probability $\leq 1/M_\gamma(P_X)$, by definition of $\mathcal{X}_1$; P_X may also have a nonatomic component. Applying Dudley et al. [2011, Proposition A.1], we can partition $\mathbb{R}^d \backslash \mathcal{X}_1$ into countably many sets, $\mathcal{X}_2 \cup \mathcal{X}_3 \cup \ldots$, such that $\mathbb{P}_{P_X} \{X \in \mathcal{X}_m\} \leq 1/M_\gamma(P_X)$ for all $m \geq 2$. From this point on, write $p_m = \mathbb{P}_{P_X} \{X \in \mathcal{X}_m\}$.

For each x in the support of P_X, let $P_{Y|X=x}$ denote the conditional distribution of Y given $X = x$. By Lemma 1, we can construct distributions $P_{Y|X=x}^1$ and $P_{Y|X=x}^0$ such that

$$P_{Y|X=x} = 0.5 P_{Y|X=x}^1 + 0.5 P_{Y|X=x}^0 \text{ and } \mathbb{E}_{P_{Y|X=x}^1}[Y] - \mathbb{E}_{P_{Y|X=x}^0}[Y] \geq 2\sigma_P^2(x),$$

where $\sigma_P^2(x) = \mathrm{Var}(Y \mid X = x)$ is the variance of $P_{Y|X=x}$.

Next fix any $\epsilon \in (0, 0.5]$. For any vector $a = (a_1, a_2, \ldots)$ of signs $a_1, a_2, \cdots \in \{\pm 1\}$, define the distribution P_a over (X, Y) as follows:

- Draw $X \sim P_X$, i.e., the same as the marginal distribution of X under P.
- Conditional on X, draw Y as

$$Y \mid X = x \sim \begin{cases} P_{Y|X=x}, & \text{if } x \in \mathcal{X}_1, \\ (0.5 + a_m \epsilon) \cdot P_{Y|X=x}^1 + (0.5 - a_m \epsilon) \cdot P_{Y|X=x}^0, & \text{if } x \in \mathcal{X}_m \text{ for } m \geq 2. \end{cases}$$

In other words, P_a differs from P in that, conditional on $X = x \in \mathcal{X}_m$ for any $m \geq 2$, the distribution of the variable Y is perturbed to be slightly more likely (if $a_m = +1$) or slightly less likely (if $a_m = -1$) to be drawn from $P_{Y|X=x}^1$ rather than from $P_{Y|X=x}^0$. Finally, we define a mixture distribution P_{mix} on $(X_1, Y_1), \ldots, (X_n, Y_n)$ as:

- Draw $A_1, A_2, \ldots \overset{\mathrm{iid}}{\sim} \mathrm{Unif}\{\pm 1\}$.
- Conditional on $A_1, A_2, \ldots$, draw $(X_1, Y_1), \ldots, (X_n, Y_n) \overset{\mathrm{iid}}{\sim} P_A$.

The remainder of the proof will center on the fact that, for ϵ chosen appropriately, it is impossible to distinguish between P^n (i.e., the data points are drawn i.i.d. from P) and P_{mix} (i.e., the data points are drawn i.i.d. from P_A, for some random A)—in the Supplementary Materials, we verify that we can apply Lemma 2 with $\epsilon_1 = 0$ and $\epsilon_2 = \epsilon_3 = \cdots = \epsilon$ to obtain

$$d_{\mathrm{TV}}(P_{\mathrm{mix}}, P^n) \leq 2n \sqrt{\sum_{m \geq 2} \epsilon^4 p_m^2} \leq \frac{2\epsilon^2 n}{\sqrt{M_\gamma(P_X)}}, \tag{6}$$

where the last step holds since $p_m \leq 1/M_\gamma(P_X)$ for all $m \geq 2$, by definition.

From this point on, at a high level, since the conditional mean of $Y|X$ differs by $\mathcal{O}(\epsilon)$ between P and P_A, this means that our confidence interval for μ_P will need to have width at least $\mathcal{O}(\epsilon)$, as long as ϵ is chosen to make the total variation distance sufficiently small. Now we formalize this argument. For any P_a, since $\widehat{C}_n$ satisfies distribution-free coverage, we have

$$\mathbb{P}_{(P_a)^n \times P_X} \left\{ \mu_{P_a}(X_{n+1}) \in \widehat{C}_n(X_{n+1}) \right\} \geq 1 - \alpha.$$

We can also calculate, for each $m \geq 2$ and each $x \in \mathcal{X}_m$ that lies in the support of P_X,

$$\mu_{P_a}(x) = (0.5 + a_m \epsilon) \mathbb{E}_{P_{Y|X=x}^1}[Y] + (0.5 - a_m \epsilon) \mathbb{E}_{P_{Y|X=x}^0}[Y]$$

$$= 0.5 \left(\mathbb{E}_{P_{Y|X=x}^1}[Y] + \mathbb{E}_{P_{Y|X=x}^0}[Y] \right) + a_m \epsilon \left(\mathbb{E}_{P_{Y|X=x}^1}[Y] - \mathbb{E}_{P_{Y|X=x}^0}[Y] \right)$$

$$= \mu_P(x) + a_m \epsilon \Delta(x),$$

297 where we write $\Delta(x) = \left(\mathbb{E}_{P^1_{Y|X=x}} [Y] - \mathbb{E}_{P^0_{Y|X=x}} [Y] \right)$. In particular, if $X_{n+1} \notin \mathcal{X}_1$, then

$$\mu_{P_a}(X_{n+1}) \in \widehat{C}_n(X_{n+1}) \;\Rightarrow\; \{\mu_P(X_{n+1}) \pm \epsilon \Delta(X_{n+1})\} \cap \widehat{C}_n(X_{n+1}) \neq \emptyset.$$

298 Therefore,

$$\mathbb{P}_{(P_a)^n \times P_X} \left\{ \{\mu_P(X_{n+1}) \pm \epsilon \Delta(X_{n+1})\} \cap \widehat{C}_n(X_{n+1}) \neq \emptyset \right\}$$

$$\geq \mathbb{P}_{(P_a)^n \times P_X} \left\{ \mu_{P_a}(X_{n+1}) \in \widehat{C}_n(X_{n+1}) \text{ and } X_{n+1} \notin \mathcal{X}_1 \right\}$$

$$\geq \mathbb{P}_{(P_a)^n \times P_X} \left\{ \mu_{P_a}(X_{n+1}) \in \widehat{C}_n(X_{n+1}) \right\} - \mathbb{P}_{(P_a)^n \times P_X} \left\{ X_{n+1} \in \mathcal{X}_1 \right\}$$

$$\geq (1 - \alpha) - (1 - \gamma) = \gamma - \alpha.$$

299 Since this bound holds for all sign vectors $a = (a_1, a_2, \dots)$, and since P_{mix} is a mixture of distribu-
300 tions $(P_a)^n$, we therefore have

$$\mathbb{P}_{P_{\text{mix}} \times P_X} \left\{ \{\mu_P(X_{n+1}) \pm \epsilon \Delta(X_{n+1})\} \cap \widehat{C}_n(X_{n+1}) \neq \emptyset \right\} \geq \gamma - \alpha.$$

301 By our total variation bound above, therefore,

$$\mathbb{P}_{P^n \times P_X} \left\{ \{\mu_P(X_{n+1}) \pm \epsilon \Delta(X_{n+1})\} \cap \widehat{C}_n(X_{n+1}) \neq \emptyset \right\} \geq \gamma - \alpha - \frac{2\epsilon^2 n}{\sqrt{M_\gamma(P_X)}}.$$

302 Since $\Delta(X_{n+1}) \geq 2\sigma_P^2(X_{n+1})$ by Lemma 1, by definition of $\sigma_{P,\beta}^2$ we therefore have

$$\mathbb{P}_{P^n \times P_X} \left\{ \Delta(X_{n+1}) \geq 2\sigma_{P,\beta}^2 \text{ and } \{\mu_P(X_{n+1}) \pm \epsilon \Delta(X_{n+1})\} \cap \widehat{C}_n(X_{n+1}) \neq \emptyset \right\}$$

$$\geq \gamma - \alpha - \beta - \frac{2\epsilon^2 n}{\sqrt{M_\gamma(P_X)}}.$$

303 Now fix some $\epsilon_0 \in [0, 0.5]$. We calculate

$$\mathbb{E}_{P^n \times P_X} \left[\text{Leb} \left(\widehat{C}_n(X_{n+1}) \right) \right] = \mathbb{E}_{P^n \times P_X} \left[\int_{t=-\infty}^{\infty} \mathbb{1} \left\{ t \in \widehat{C}_n(X_{n+1}) \right\} \, \mathrm{d}t \right]$$

$$\geq \mathbb{E}_{P^n \times P_X} \left[\Delta(X_{n+1}) \cdot \int_{\epsilon=-\epsilon_0}^{\epsilon_0} \mathbb{1} \left\{ \mu_P(X_{n+1}) + \epsilon \Delta(X_{n+1}) \in \widehat{C}_n(X_{n+1}) \right\} \, \mathrm{d}\epsilon \right]$$

$$\geq \mathbb{E}_{P^n \times P_X} \left[\Delta(X_{n+1}) \cdot \int_{\epsilon=0}^{\epsilon_0} \mathbb{1} \left\{ \{\mu_P(X_{n+1}) \pm \epsilon \Delta(X_{n+1})\} \cap \widehat{C}_n(X_{n+1}) \neq \emptyset \right\} \, \mathrm{d}\epsilon \right]$$

$$\geq 2\sigma_{P,\beta}^2 \int_{\epsilon=0}^{\epsilon_0} \mathbb{P}_{P^n \times P_X} \left\{ \Delta(X_{n+1}) \geq 2\sigma_{P,\beta}^2, \, \{\mu_P(X_{n+1}) \pm \epsilon \Delta(X_{n+1})\} \cap \widehat{C}_n(X_{n+1}) \neq \emptyset \right\} \, \mathrm{d}\epsilon$$

$$\geq 2\sigma_{P,\beta}^2 \int_{\epsilon=0}^{\epsilon_0} \gamma - \alpha - \beta - \frac{2\epsilon^2 n}{\sqrt{M_\gamma(P_X)}} \, \mathrm{d}\epsilon = 2\sigma_{P,\beta}^2 \left[\epsilon_0(\gamma - \alpha - \beta) - \frac{2\epsilon_0^3 n}{3\sqrt{M_\gamma(P_X)}} \right].$$

304 Finally, choosing $\epsilon_0 = \min \left\{ \left(\frac{(\gamma - \alpha - \beta)\sqrt{M_\gamma(P_X)}}{2n} \right)^{1/2}, 0.5 \right\}$ yields the desired lower bound.

References

306 Jayadev Acharya, Clément L Canonne, and Gautam Kamath. A chasm between identity and
307   equivalence testing with conditional queries. *arXiv preprint arXiv:1411.7346*, 2014.

308 Jayadev Acharya, Constantinos Daskalakis, and Gautam Kamath. Optimal testing for properties of
309   distributions. *arXiv preprint arXiv:1507.05952*, 2015.

310 Rina Foygel Barber. Is distribution-free inference possible for binary regression? *Electronic Journal*
311   *of Statistics*, 14(2):3487–3524, 2020.

Rina Foygel Barber, Emmanuel J Candes, Aaditya Ramdas, and Ryan J Tibshirani. Predictive inference with the jackknife+. *The Annals of Statistics*, 49(1):486–507, 2021.

Clément L Canonne, Dana Ron, and Rocco A Servedio. Testing probability distributions using conditional samples. *SIAM Journal on Computing*, 44(3):540–616, 2015.

Clément L Canonne, Ilias Diakonikolas, Themis Gouleakis, and Ronitt Rubinfeld. Testing shape restrictions of discrete distributions. *Theory of Computing Systems*, 62(1):4–62, 2018.

Siu-On Chan, Ilias Diakonikolas, Paul Valiant, and Gregory Valiant. Optimal algorithms for testing closeness of discrete distributions. In *Proceedings of the twenty-fifth annual ACM-SIAM symposium on Discrete algorithms*, pages 1193–1203. SIAM, 2014.

Ilias Diakonikolas and Daniel M Kane. A new approach for testing properties of discrete distributions. In *2016 IEEE 57th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 685–694. IEEE, 2016.

Ilias Diakonikolas, Themis Gouleakis, John Peebles, and Eric Price. Sample-optimal identity testing with high probability. In *45th International Colloquium on Automata, Languages, and Programming (ICALP 2018)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2018.

Richard M Dudley, Rimas Norvaiša, and Rimas Norvaiša. *Concrete functional calculus*. Springer, 2011.

Evarist Giné and Richard Nickl. *Mathematical foundations of infinite-dimensional statistical models*, volume 40. Cambridge University Press, 2016.

Chirag Gupta, Aleksandr Podkopaev, and Aaditya Ramdas. Distribution-free binary classification: prediction sets, confidence intervals and calibration. *arXiv preprint arXiv:2006.10564*, 2020.

Byol Kim, Chen Xu, and Rina Foygel Barber. Predictive inference is free with the jackknife+-after-bootstrap. *arXiv preprint arXiv:2002.09025*, 2020.

Jing Lei, Max G?Sell, Alessandro Rinaldo, Ryan J Tibshirani, and Larry Wasserman. Distribution-free predictive inference for regression. *Journal of the American Statistical Association*, 113(523): 1094–1111, 2018.

Dhruv Medarametla and Emmanuel J Candès. Distribution-free conditional median inference. *arXiv preprint arXiv:2102.07967*, 2021.

Harris Papadopoulos, Kostas Proedrou, Volodya Vovk, and Alex Gammerman. Inductive confidence machines for regression. In *European Conference on Machine Learning*, pages 345–356. Springer, 2002.

Gregory Valiant and Paul Valiant. Estimating the unseen: an n/log (n)-sample estimator for entropy and support size, shown optimal via new clts. In *Proceedings of the forty-third annual ACM symposium on Theory of computing*, pages 685–694, 2011a.

Gregory Valiant and Paul Valiant. The power of linear estimators. In *2011 IEEE 52nd Annual Symposium on Foundations of Computer Science*, pages 403–412. IEEE, 2011b.

Gregory Valiant and Paul Valiant. An automatic inequality prover and instance optimal identity testing. *SIAM Journal on Computing*, 46(1):429–455, 2017.

Vladimir Vovk, Alex Gammerman, and Glenn Shafer. *Algorithmic learning in a random world*. Springer Science & Business Media, 2005.

Vladimir Vovk, Ilia Nouretdinov, Valery Manokhin, and Alexander Gammerman. Cross-conformal predictive distributions. In *Conformal and Probabilistic Prediction and Applications*, pages 37–51. PMLR, 2018.