Scalable Estimation of Nonparametric Markov Networks with Mixed-Type Data

Abstract

Markov network characterizes the conditional independence structure, or Markov property, among a set of random variables. Existing work focuses on specific families of distributions (e.g., exponential families) and/or certain structures of the graph, and most of them can only handle variables of a single data type (continuous or discrete). In this work, we generalize the characterization of the conditional independence structure to handle *general distributions for all data types* (i.e., continuous, discrete, and mixed-type) with *general functional relations among variables*, thus giving rise to a Markov network structure learning algorithm in one of the most general settings. To deal with the computational challenge of the problem, especially for large graphs, we unify all cases under the same umbrella of a regularized score matching framework. We validate the theoretical results experimentally and demonstrate the scalability of the approach–it produces the estimated Markov network over up to 5000 nodes within one hour on CPUs. We further discuss the implication of the proposed approach in causal discovery.

1 Introduction

Markov networks (also known as *undirected graphical models* or *Markov random fields*) represent the conditional dependencies among random variables. They provide clear semantics in a graphical manner to cope with the uncertainty in probability theory, with a wide application in fields including physics (Cimini et al., 2019), chemistry (Dodani et al., 2016), biology (Jaimovich et al., 2006), and sociology (Carrington et al., 2005). The undirected nature of edges also allows cyclic, overlapping, or hierarchical interactions (Shen et al., 2009). To estimate the Markov network from observational data, existing work focuses on certain parametric families of distributions, and a majority of them study the Gaussian case. By assuming that the variables are from a multivariate Gaussian distribution, the dependencies can be well represented by the support of the precision, or inverse covariance, matrix according to Hammersley-Clifford theorem (Besag, 1974; Grimmett, 1973). Together with various statistical estimators (e.g., the *graphical lasso* (Friedman et al., 2008) and *neighborhood selection* (Meinshausen & Bühlmann, 2006)), this connection between the precision matrix and graphical structure has been well exploited in the Gaussian case in the past decades (Yuan, 2010; Ravikumar et al., 2011). However, methods for Gaussian graphical models might fail to correctly capture dependencies between variables deviating from Gaussian or including nonlinearity.

While non-Gaussianity may be more common in real-world data generating process, not many results are available for Markov network structure learning on data that do not follow a multivariate Gaussian distribution. In the *discrete* setting, Ravikumar et al. (2010) showed that a binary Ising model can be recovered by neighborhood selection using ℓ_1 penalized logistic regression; Loh & Wainwright (2013) encoded extra structural relations in the proposed generalized covariance matrix to model the dependencies for Markov networks with certain structures (e.g., tree structures or graphs with only singleton separator sets) among variables from exponential families. Several approaches allowed estimation for non-Gaussian continuous variables while most of them assumed parametric assumptions such as the exponential families (Yang et al., 2015; Lin et al., 2016; Suggala et al., 2017) or Gaussian copulas (Liu et al., 2009; 2012; Harris & Drton, 2013). These methods illustrate the possibility of reliable Markov network estimation in several non-Gaussian cases, but still, the models are restricted to specific parametric families of distributions and/or structures of conditional independencies.

Concerned with describing Markov properties of non-Gaussian data with general *continuous* distributions, Morrison et al. (2017); Baptista et al. (2021) used the *generalized precision matrix* to encode the conditional independence structure. Specifically, the approach is based on a theorem that the zero pattern of the Hessian of the log-density determines the conditional independencies between non-Gaussian continuous variables (Spantini et al., 2018). A method based on transport map (SING) is then designed to estimate the data density from samples, and the structure is derived from the estimated density. This approach achieves consistent Markov network structure recovery in a general non-Gaussian continuous setting. However, with the *generalized precision matrix*, it cannot handle discrete or mixed-type data. In addition, density estimation, especially for non-Gaussian data, can be computationally challenging for large graphs, limiting the scalability of this approach.

To deal with these remaining obstacles, we provide a scalable estimation framework for nonparametric Markov networks. For the continuous data, we adopt a characterization of conditional independence for general distributions based on an observation by Spantini et al. (2018). Besides, we propose necessary and sufficient conditions for the conditional independence relations between variables in both general discrete and mixed-type settings as well as the corresponding characterizations of the structure. Moreover, our work does not constrain the distribution to be of specific families, such as exponential families. To the best of our knowledge, this illustrates the feasibility of Markov network structure learning in probably the most general nonparametric setting, where the data could be continuous, discrete, or even mixed-type and does not need to be from specific distributions. Besides, it is worth noting that there are no specific assumptions on the functional relations among variables.

Furthermore, we put all these cases under the same umbrella of the estimation framework based on regularized score matching, as an extension of the score matching framework (Hyvärinen & Dayan, 2005). Different from previous approach (SING) that applies transport map to estimate the data density for general continuous distributions, our framework allows us to only estimate the model score function parameterized by a deep model, from which the characterization matrix of the Markov network structure could be directly calculated. To facilitate the estimation process, we also exploit suitable penalties on the characterization matrix to encourage constantly sparse entries. Besides, we adopt recent advancements on score matching (Song et al., 2020) to further scale up the process. Our method therefore narrows the gap between reliable structure learning and scalable deep learning techniques. We validate the theoretical results experimentally, and the scalability has been illustrated on graphs with up to 5000 nodes. Finally, we discuss how to extend the proposed Markov network estimation approach to perform causal discovery (Spirtes et al., 2000; Pearl, 2000).

2 GENERALIZED CHARACTERIZATION OF CONDITIONAL INDEPENDENCE

Suppose that we observe a collection of random variables $\mathbf{X} = (X_1, \ldots, X_d)$. Our goal is to discover the underlying Markov network structure. Specifically, it is an undirected graph $\mathcal{G}$ comprising a set of vertices $\mathbf{V} = \{1, \ldots, d\}$ and edges $\mathbf{E}$. The edges $\mathbf{E}$ encode the conditional independence relations or the global Markov property: for any disjoint subsets $\mathbf{A}$, $\mathbf{B}$, and $\mathbf{C}$ in the vertices set $\mathbf{V}$ such that $\mathbf{C}$ separates $\mathbf{A}$ and $\mathbf{B}$, $\mathbf{X}_{\mathbf{A}}$ and $\mathbf{X}_{\mathbf{B}}$ are conditionally independent given $\mathbf{X}_{\mathbf{C}}$, i.e., $\mathbf{X}_{\mathbf{A}} \perp\!\!\!\perp \mathbf{X}_{\mathbf{B}} \mid \mathbf{X}_{\mathbf{C}}$.[1] Throughout this paper, we use an uppercase letter to denote a random variable and a lowercase letter with subscripts to denote the value of a random variable (e.g., $X_i = x_i$ for the value of X_i). For a discrete variable, say X_i, we denote its support by $\{x_{i1}, \ldots, x_{iM_i}\}$, where M_i is its cardinality.

As an alternative characterization of the conditional independence relations encoded by the graph, the pairwise Markov property requires that every pair of non-adjacent variables in the graph are conditionally independent given the remaining variables. That is, for any $i \neq j$, an edge between X_i and X_j is absent if and only if X_i and X_j are conditionally independent given the remaining variables, i.e., $X_i \perp\!\!\!\perp X_j \mid \mathbf{X}_{\mathbf{V} \setminus (i,j)}$. According to Lauritzen (1996), the pairwise Markov property is equivalent to the global one when the density is strictly positive. In order to estimate nonparametric Markov networks in this setting, we explore generalized characterizations of conditional independence in all types of data (i.e., continuous, discrete, and mixed-type) without distributional constraints. We start from learning conditional independence structures in continuous data with a procedure inspired by Spantini et al. (2018), and then propose new characterizations for discrete and mixed-type data.

---

[1] For any set $\mathbf{S} \subset \mathbf{V}$, we write $\mathbf{X}_{\mathbf{S}} = \{X_i : i \in \mathbf{S}\}$.

Ideally, we would like to construct a matrix Ω that satisfies the following desiderata so that it can be used to characterize the Markov network structure: **(a)** For any $i \neq j$, if $\Omega_{i,j} = 0$, then $X_i \perp\!\!\!\perp X_j \mid X_{\mathbf{V} \setminus (i,j)}$; **(b)** The probability measure is *not restricted* to be from specific families but only needs to be strictly positive; **(c)** The undirected graph $\mathcal{G}$ is *not restricted* to be of certain structures; **(d)** For continuous variables, the density is smooth; **(e)** For discrete variables, the cardinality is not restricted; **(f)** To enable practical estimation procedure, matrix Ω is differentiable.

Property **(a)** is the characterization of the pairwise Markov property. Properties **(b)** and **(c)** differentiate our work from most previous works such as those assuming Gaussianity or/and certain structures of the conditional independence. Properties **(d)** and **(e)** further raise the difficulty of our task, because, in addition to not being restricted to a specific family of distributions, our characterization Ω has to be available for all data types (i.e., continuous, discrete, and mixed-type) with mild assumptions. For discrete variables, Property **(e)** removes the limitation of cardinality, thus differentiating our work from those focusing on the binary Ising model. Property **(f)** allows us to incorporate an ℓ_1 regularization term in the estimation procedure and make use of gradient-based optimization.

2.1 CHARACTERIZATION FOR CONTINUOUS DATA

We aim to find the necessary and sufficient condition for $X_i \perp\!\!\!\perp X_j \mid \mathbf{X}_{\mathbf{V} \setminus \{i,j\}}$. By definition, if X_i is conditionally independent of X_j given all remaining variable $\mathbf{X}_{\mathbf{V} \setminus \{i,j\}}$, we could factor the probability density function (PDF) $p_{\mathbf{X}}$ as follows

$$p_{\mathbf{X}}(\mathbf{x}) = p_{X_i \mid \mathbf{X}_{\mathbf{V} \setminus \{i,j\}}}(x_i \mid \mathbf{x}_{\mathbf{V} \setminus \{i,j\}}) p_{X_j \mid \mathbf{X}_{\mathbf{V} \setminus \{i,j\}}}(x_j \mid \mathbf{x}_{\mathbf{V} \setminus \{i,j\}}) p_{\mathbf{X}_{\mathbf{V} \setminus \{i,j\}}}(\mathbf{x}_{\mathbf{V} \setminus \{i,j\}}). \quad (1)$$

Together with the assumption that $p_{\mathbf{X}}$ is smooth, we have

$$\frac{\partial^2 \log p_{\mathbf{X}}}{\partial x_i \partial x_j} = 0. \quad (2)$$

Conversely, the solution of Eq. (2) is given by $\log p_{\mathbf{X}}(\mathbf{x}) = g(x_{1:i-1}, x_{i+1:n}) + h(x_{1:j-1}, x_{j+1:n})$, which yields Eq. (1) (Lauritzen, 1996). This connection between pairwise conditional independence and cross derivatives of the log density has been observed in Spantini et al. (2018). Methods based on it have also been proposed recently (Morrison et al., 2017; Baptista et al., 2021). Following Baptista et al. (2021), one could characterize conditional dependence between X_i and X_j in the continuous distribution as

$$\Omega_{ij}^{[c]} := \left(\mathbb{E}_p \left[f_{i,j}^{[c]}(\mathbf{x})^2 \right] \right)^{\frac{1}{2}}, \quad (3)$$

where $f_{i,j}^{[c]}(\cdot)$ denotes the LHS of Eq. (2). The group structure of it could help achieve simultaneous sparse approximation (Yuan & Lin, 2006; Huang & Zhang, 2010) when being applied as an ℓ_1 regularizer in the estimation procedure, as we will describe in Sec. 3. We also apply the same group structures for both the discrete and mixed-type cases and skip the reintroduction of them for brevity. The characterization of the Markov property is as follows.

Corollary 1. *Assume*

    i. $\mathbf{X} = (X_1, \ldots, X_d)$ *is a set of continuous variable.*

    ii. *The PDFs of* $\mathbf{X}$ *are strictly positive and smooth.*

    iii. *The characterization matrix* $\Omega^{[c]}$ *is defined according to Eq.* (3).

Then for any $i \neq j$, $\Omega_{i,j}^{[c]} = 0$ *implies* $X_i \perp\!\!\!\perp X_j \mid \mathbf{X}_{\mathbf{V} \setminus \{i,j\}}$.

The proof is shown in Appx. A.1. It is worth noting that Cor. 1 also covers the Gaussian case, where the cross-derivatives of the log-density correspond to entries in the inverse covariance matrix (Drton et al., 2008), thus generalizing previous work assuming Gaussianity. Hence, the support of matrix Ω characterizes conditional independence between continuous variables for general distributions.

2.2 CHARACTERIZATION FOR DISCRETE DATA

Since most of the previous work focuses on the Gaussian setting, and a few works for non-Gaussian distribution are mostly restricted to the exponential family, the characterization for continuous data

discussed in Sec. 2.1 has broadened the scope of reliable Markov network learning. However, the characterization may not be applied for discrete data as the gradient is undefined. In this section, we provide such a characterization of Markov network structure in the discrete case. Similar to the continuous case, a key ingredient of the proposed characterization is the necessary and sufficient conditions of conditional independence for discrete data, which we establish in the following theorem.

Theorem 1. *Denote* $\mathbf{V}$ *as a set of discrete variables and* $X_i, X_j \in \mathbf{V}$*. For brevity, denote* $\mathbf{V} \backslash \{X_i, X_j\}$ *as* $\mathbf{Z}$*. Let* $\{x_{i1}, \ldots, x_{iM_i}\}$ *and* $\{x_{j1}, \ldots, x_{jM_j}\}$ *be the support of variables* X_i *and* X_j*. Denote* z *as any value(s) of* $\mathbf{Z}$*. Then,* $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$ *if and only if, for all* $k \in [M_i]$ *and* $l \in [M_j]$ *with* $k \neq 1$ *and* $l \neq 1$*, we have*

$$(\log m(x_{i1}, x_{j1}, z) - \log m(x_{ik}, x_{j1}, z)) - (\log m(x_{i1}, x_{jl}, z) - \log m(x_{ik}, x_{jl}, z)) = 0. \quad (4)$$

The proof is provided in Appx. A.2. Note that we denote $m(x_{ik}, x_{jl}, z)$ as the joint probability mass function (PMF) of $\{X_i, X_j, \mathbf{Z}\}$, simplified from $m_{X_i, X_j, \mathbf{Z}}(x_{ik}, x_{jl}, z)$. Based on Thm. 1, we propose the characterization of conditional independence for discrete data $\Omega_{i,j}^{[d]}$ as follows:

$$\Omega_{i,j}^{[d]} := \mathbb{E}_m \left[\sum_{k,l} f^{[d]}(x_{i1}, x_{ik}, x_{j1}, x_{jl}, z)^2 \right], \quad (5)$$

where $f^{[d]}(x_{i1}, x_{ik}, x_{j1}, x_{jl}, z)$ denotes the LHS of Eq. (4). The support of the matrix above satisfies the pairwise Markov property and characterizes the Markov network structure, formally stated below with its proof in Appx. A.3.

Corollary 2. *Assume*

    i. $\mathbf{X} = (X_1, \ldots, X_d)$ *is a set of discrete variable.*

    ii. *The PMFs of* $\mathbf{X}$ *are strictly positive.*

    iii. *The characterization matrix* $\Omega^{[d]}$ *is defined according to Eq.* (5)*.*

Then for any $i \neq j$*,* $\Omega_{i,j}^{[d]} = 0$ *implies* $\mathbf{x}_i \perp\!\!\!\perp \mathbf{x}_j \mid \mathbf{x}_{\mathbf{V} \backslash \{i,j\}}$*.*

Therefore, we have a characterization matrix $\Omega^{[d]}$ to represent the conditional independence structure for discrete data. It is worth noting that, unlike the generalized covariance matrix in Loh & Wainwright (2012) that only applies for certain structures among variables from exponential families, the proposed characterization matrix $\Omega^{[d]}$ encodes the Markov properties for general discrete distributions without any structural constraints. Also, compared with Ravikumar et al. (2010), Thm. 1 could be applied to general graphical models apart from binary Ising models and does not rely on the structural condition. It also does not limit the cardinalities of discrete variables. Hence, Theorem 1 sheds light on characterizing arbitrary conditional independence structures for general discrete distributions.

2.3 CHARACTERIZATION FOR MIXED-TYPE DATA

In the previous sections, we have presented characterizations of conditional independence structures for both general continuous and discrete distribution. However, it is common for real-world datasets to have mixtures of continuous and discrete variables. Unfortunately, most works focus on either continuous or discrete data, and previous results for mixed-type data are mostly based on conditional Gaussian distribution (Lauritzen et al., 1989; Edwards, 1990; Lauritzen, 1996; Fellinghauer et al., 2013; Lee & Hastie, 2015; Cheng et al., 2017). Similar to the continuous and discrete settings, in this section, we introduce a novel characterization of the pairwise Markov property for general distributions with mixed data-types. We first provide necessary and sufficient conditions of conditional independence for mixed-type data in the following theorem, with proof given in Appx. A.4.

Theorem 2. *Denote* $\mathbf{V}$ *as a set of mixed-type variables and* $X_i, X_j \in \mathbf{V}$*, where* X_i *is discrete and* X_j *is continuous. Let* $\{x_{i1}, \ldots, x_{iM_i}\}$ *be the support of variables* X_i*. For brevity, denote* $\mathbf{V} \backslash \{X_i, X_j\}$ *as* $\mathbf{Z}$*. Denote* z *as any value(s) of* $\mathbf{Z}$ *and* x_j *as any value of the continuous variable* X_j*. Then,* $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$ *if and only if, for all* $k \in [M_i]$ *with* $k \neq 1$*, we have*

$$\frac{\partial \log \left(p_{X_j, \mathbf{Z} \mid X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right)}{\partial x_j} - \frac{\partial \log \left(p_{X_j, \mathbf{Z} \mid X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right)}{\partial x_j} = 0. \quad (6)$$

Based on Thm. 2, we propose to characterize the conditional independence between X_i and X_j given all remaining variables with $\Omega_{i,j}^{[m]}$ defined as

$$\Omega_{i,j}^{[m]} := \begin{cases} \mathbb{E}_\pi \left[f_{i,j}^{[c]}(\mathbf{x})^2 \right] & \text{if } X_i \in \mathbf{X}_c, X_j \in \mathbf{X}_c \\ \mathbb{E}_\pi \left[\sum_{k,l} f^{[d]}(x_{i1}, x_{ik}, x_{j1}, x_{jl}, z)^2 \right] & \text{if } X_i \in \mathbf{X}_d, X_j \in \mathbf{X}_d \\ \mathbb{E}_\pi \left[\sum_k f^{[m]}(x_i, x_{j1}, x_{jk}, z)^2 \right] & \text{if } X_i \in \mathbf{X}_c, X_j \in \mathbf{X}_d \\ \mathbb{E}_\pi \left[\sum_k f^{[m]}(x_j, x_{i1}, x_{ik}, z)^2 \right] & \text{if } X_i \in \mathbf{X}_d, X_j \in \mathbf{X}_c, \end{cases} \tag{7}$$

where $f^{[m]}$ denotes LHS of Eq. (6) and π is the probability function. $\mathbf{X}_c$ and $\mathbf{X}_d$ are sets of continuous and discrete variables, respectively. Its characterization of Markov property is as follows

Corollary 3. *Assume*

  i. $\mathbf{X} = (X_1, \ldots, X_d)$ *is a set of variables containing both continuous and discrete variables.*

  ii. *For continuous variables, the PDFs are strictly positive and smooth.*

  iii. *For discrete variables, the PMFs are strictly positive.*

  iv. *The characterization matrix $\Omega^{[m]}$ is defined according to Eq. (7).*

Then for any $i \neq j$, $\Omega_{i,j}^{[m]} = 0$ implies $X_i \perp\!\!\!\perp X_j \mid \mathbf{X}_{\mathbf{V} \setminus \{i,j\}}$.

The proof is given in Appx. A.5. The proposed characterization matrix $\Omega^{[m]}$ encodes the pairwise Markov property for the mixed-type data. More general than previous works, it does not require specific families of distributions, structures of the underlying graph, or cardinality of discrete variables.

3 SCALABLE ESTIMATION WITH REGULARIZED SCORE MATCHING

In Sec. 2, we provide characterizations of conditional independencies for general distribution in continuous, discrete, and mixed-type settings. Based on the introduced necessary and sufficient conditions, these characterizations generalize previous work and establish one of the foundations for nonparametric estimation of Markov network structures with minimal assumptions.

Having said that, besides general characterizations of the Markov property with theoretical guarantees, a scalable estimation framework is necessary for reliable and practical structure learning. Hence, we introduce a regularized score matching-based framework for all considered settings (i.e., general distributions of continuous, discrete, and mixed-type variables), of which the scalability is illustrated by results on graphs with up to 5000 nodes, as shown in Fig. 2.

3.1 ESTIMATION FOR CONTINUOUS DATA

We start with the continuous setting. Denote $p(\mathbf{x}; \theta)$ as a parameterized density model with a parameter vector θ. The goal is to estimate parameter θ from the observation $\mathbf{x}$. We aim to optimize the following objective function, which is based on Fisher divergence:

$$O_c(\theta) = \frac{1}{2} \int_{\mathbf{x} \in \mathbb{R}^d} p(\mathbf{x}) \|\nabla_\mathbf{x} \log p(\mathbf{x}; \theta) - \nabla_\mathbf{x} \log p(\mathbf{x})\|^2 d\mathbf{x} + \rho_\lambda(\Omega^{[c]}), \tag{8}$$

where $\rho_\lambda(\cdot)$ denotes a sparsity penalty function and λ is the penalty parameter. $\Omega^{[c]}$ is defined in Eq. 3 as our characterization of the conditional independence structure for continuous data. If we assume the model is not degenerate, where different values of θ correspond to different PDFs, the asymptotic consistency of the optimization has been shown in Thm. 2 by Hyvärinen & Dayan (2005). We impose a sparsity penalty owing to finite-sampling errors in practice.

Also with a trick in Hyvärinen & Dayan (2005); Pham & Garat (1997), one can remove the data log-density $\log p_\mathbf{X}$ from Eq. (8) by optimizing the following equation, which is equivalent to Eq. (8):

$$O_c(\theta) = \int_{\mathbf{x} \in \mathbb{R}^n} p(\mathbf{x}) \sum_{i=1}^d \left[\frac{1}{2} \|\nabla_{x_i} \log p(\mathbf{x}; \theta)\|^2 + H_{x_i}(\log p(\mathbf{x}; \theta)) \right] dx + \rho_\lambda(\Omega^{[c]}), \tag{9}$$

where H denotes the Hessian. The proof is directly based on Hyvärinen & Dayan (2005) and we include it (Lemma 1) in Appx. A.6.1 for completeness. It is worth noting that previous work on Markov network structure learning with general continuous distribution (SING (Morrison et al., 2017; Baptista et al., 2021)) applies a transport map to estimate data density from samples, which can be computationally challenging for non-Gaussian data with a large number of variables. Thus, it may not be scalable as suggested by Fig. 1 and Table 1. To avoid this, the proposed regularized score matching allows us to optimize the objective function by only estimating the model score function. Moreover, the estimated model score function directly leads to the characterization matrix $\Omega^{[c]}$ by taking further derivatives, thus efficiently giving rise to the estimated Markov network structure.

3.2 ESTIMATION FOR DISCRETE DATA

For the estimation in the discrete case, one can not directly apply the method introduced for the continuous case since the gradient, on which the continuous score function is based, is not defined for discrete data. An intuitive solution is to replace the gradient with a general linear operator $\mathcal{L}$ (Lyu, 2012). For instance, Eq. (8) can be reformulated as follows

$$O_d(\theta) = \frac{1}{2} \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \left\| \frac{\mathcal{L}(m(\mathbf{x}; \theta))}{m(\mathbf{x}; \theta)} - \frac{\mathcal{L}(m_{\mathbf{X}}(\mathbf{x}))}{m_{\mathbf{X}}(\mathbf{x})} \right\|^2 + \rho_\lambda(\Omega^{[d]}), \tag{10}$$

where m denotes PMF. In this formulation, $\mathcal{L}(\cdot)$ is a generalized version of the score function for discrete data. As shown in Lyu (2012), $O_d(\theta)$ keeps the computational advantages of score matching for continuous data, i.e., the normalizing partition is canceled out and the formulation could be transformed to an expectation of functions of the unnormalized model. In order to guarantee the consistency of score matching based on Eq. (10), the linear operator $\mathcal{L}(\cdot)$ needs to be "complete".

Definition 1 (Completeness (Lyu, 2012)). *A linear operator $\mathcal{L}(\cdot)$ is complete if $\frac{\mathcal{L}(p(\mathbf{x}))}{p(\mathbf{x})} = \frac{\mathcal{L}(q(\mathbf{x}))}{q(\mathbf{x})}$ implies $p(\mathbf{x}) = q(\mathbf{x})$ almost everywhere, where $p(\mathbf{x})$ and $q(\mathbf{x})$ are two PMFs.*

According to Defn. 1, Lyu (2012) used the marginalization operator $\mathcal{M}(\cdot) : \mathcal{F}^1 \mapsto \mathcal{F}^d$ as a choice for $\mathcal{L}(\cdot)$, which is defined as

$$\mathcal{M}(f(\mathbf{x})) = \begin{pmatrix} \vdots \\ \mathcal{M}_i(f(\mathbf{x})) \\ \vdots \end{pmatrix} = \begin{pmatrix} \vdots \\ \sum_{\mathbf{x}} f(\mathbf{x}) \\ \vdots \end{pmatrix}, \tag{11}$$

where $f \in \mathcal{F}^1$. We can observe that $\mathcal{M}_i(f(\mathbf{x}))$ is the marginal density of $\mathbf{x}^{\backslash i}$, where $\mathbf{x}^{\backslash i}$ denotes the vector $\mathbf{x}$ after dropping the i-th element (i.e., marginalization). The completeness of $\mathcal{M}(\cdot)$ has been shown in Brook (1964), also included as Lemma 3 in Lyu (2012). Then we have

$$O_d(\theta) = \frac{1}{2} \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \left\| \frac{\mathcal{M}(m(\mathbf{x}; \theta))}{m(\mathbf{x}; \theta)} - \frac{\mathcal{M}(m_{\mathbf{X}}(\mathbf{x}))}{m_{\mathbf{X}}(\mathbf{x})} \right\|^2 + \rho_\lambda(\Omega^{[d]}). \tag{12}$$

Thus, it is plausible for us to replace the gradient with $\mathcal{M}(\cdot)$ for discrete data. However, one key advantage of regularized score matching is that it does not have to explicitly estimate the data density (i.e., $p_X(\mathbf{x})$ in Theorem 1). As shown by Lyu (2012), we can also optimize Eq. (12) in a similar way, which is equivalent to optimizing the following equation

$$O_d(\theta) = \frac{1}{2} \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^d \sum_{x_{ij}} \left(\frac{m(\sim x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)}{m(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)} \right)^2 + \rho_\lambda(\Omega^{[d]}), \tag{13}$$

where $m(\sim x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)$ denotes the conditional PMF for the i-th entry of $\mathbf{x}$ taking values other than x_{ij}. The lemma (Lemma 2) is formalized with its proof in Appx. A.6.2 for completeness. Based on Thm. 1 and Thm. 2, similar to the continuous case, we can estimate Markov network structures for general distributions in the discrete setting under the same umbrella of regularized score matching.

3.3 Estimation for mixed-type data

For mixed-type data, we define the objective function as follows

$$O_m(\theta) = \mathbb{E}_{\mathbf{x}} \left[\sum_i s_i(\mathbf{x}; \theta) \right] + \rho_\lambda(\Omega^{[m]}), \qquad (14)$$

where

$$s_i(\mathbf{x}; \theta) \coloneqq \begin{cases} \frac{1}{2} \|\nabla_{x_i} \log \pi(\mathbf{x}; \theta)\|^2 + H_{x_i}(\log \pi(\mathbf{x}; \theta)) & X_i \in \mathbf{X}_c \\ \frac{1}{2} \sum_{x_{ij}} \left(\frac{\pi(\sim x_{ij} | \mathbf{x}^{\backslash i}; \theta)}{\pi(x_{ij} | \mathbf{x}^{\backslash i}; \theta)} \right)^2 & X_i \in \mathbf{X}_d, \end{cases} \qquad (15)$$

where the density π is strictly positive. Basically, $O_m(\theta)$ is a regularized version of the combination of the objective functions for the continuous and discrete cases. Because $\Omega^{[m]}$ also encodes the dependencies between continuous and discrete variables, we could estimate the support of it for mixed-type data without assuming group structures of data types. The consistency is supported by the following corollary, where we define $O'_m(\theta)$ as $O_m(\theta) - \rho_\lambda(\Omega^{[m]})$.

Corollary 4. *Assume*

    i. *The data density $\pi_{\mathbf{X}}(\cdot)$ is equal to $\pi(\cdot; \theta^*)$ for some θ^*.*

    ii. *The data density $\pi_{\mathbf{X}}(\cdot)$ and model density $\pi(\cdot; \theta)$ are strictly positive. $\pi_{\mathbf{X}}(\cdot)$ and $\pi(\cdot; \theta)$ is differentiable and twice-differentiable, respectively, w.r.t. continuous variables. For some θ^*, $\pi_{\mathbf{X}}(\cdot) = \pi(\cdot; \theta^*)$ and no other parameter value gives a density that is equal to $\pi(\cdot; \theta^*)$ almost everywhere.*

    iii. *The expectations $E_{\mathbf{x}} \left[\|\log \pi(\mathbf{x}; \theta)\|^2 \right]$ and $E_{\mathbf{x}} \left[\|\log \pi_{\mathbf{X}}(\mathbf{x})\|^2 \right]$ are finite for any θ, and $\pi_{\mathbf{X}}(\mathbf{x}) \log \pi(\mathbf{x}; \theta)$ goes to zero for any θ when $\|\mathbf{x}\| \to \infty$.*

Then $O'_m(\theta) = 0$ implies $\theta = \theta^*$.

Cor. 4 follows from Lemma 1 (Hyvärinen & Dayan, 2005) and Lemma 2 (Lyu, 2012) included in Appx. A.6. Together with Thm. 2, one could estimate Markov network structures for mixed-type data in the general setting.

3.4 Sparsity regularization

By minimizing the objective function $O(\theta) \in \{O_c(\theta), O_d(\theta), O_m(\theta)\}$, our goal is essentially to perform a model selection task, i.e., to learn of the support of $\Omega \in \{\Omega^{[c]}, \Omega^{[d]}, \Omega^{[m]}\}$. Here, using ℓ_0 penalty may be computationally infeasible because it leads to a discrete optimization problem that is difficult to solve. Following previous works (Tibshirani, 1996; Friedman et al., 2008), we adopt the ℓ_1 regularizer $\rho_\lambda(\Omega) = \lambda \|\Omega\|_1$. In particular, the high-dimensional support recovery of ℓ_1 regularizer has been extensively studied in the literature; for instance, see Wainwright (2009) for variable selection and Ravikumar et al. (2008; 2011) for Gaussian graphical model selection. Although ℓ_1 regularizer induces sparsity, it may lead to bias in the resulting solution and thereby worsen the performance (Fan & Li, 2001; Breheny & Huang, 2011). This is because the ℓ_1 norm increases linearly with the absolute value of nonzero entries, which is different from ℓ_0 norm that is constant for nonzero entries. Therefore, we also experiment with smoothly clipped absolute deviation (SCAD) penalty (Fan & Li, 2001), minimax concave penalty (MCP) (Zhang, 2010), and adaptive ℓ_1 penalty (Zou, 2006) in this work, which helps remedy the bias issue of ℓ_1 regularization. Specifically, SCAD and MCP penalties may be interpreted as a hybrid of ℓ_0 and ℓ_1 penalty, while adaptive ℓ_1 penalty reweighs the penalty coefficient λ by the initial estimate of Ω without regularization. Furthermore, the support recovery of ℓ_1 penalty relies on the incoherence condition in various cases (Wainwright, 2009; Ravikumar et al., 2008; 2011), which may be a rather strong assumption in practice, whereas the SCAD and MCP penalties do not (Loh & Wainwright, 2017). We adopt the SCAD penalty according to experimental results (Fig. 3 in Appx. B).

4 TOWARDS NONPARAMETRIC CAUSAL DISCOVERY

In this section, we briefly discuss the implication of our proposed Markov network estimation method in causal discovery, of which the goal is to learn graphical models with causal interpretations.

The major classes of approaches for causal discovery are constraint-based approaches that utilize conditional independence tests and score-based approaches that optimize a specific score function. Among them, PC (Spirtes & Glymour, 1991) with kernel-based conditional independence test (Zhang et al., 2012) and GES (Chickering, 2002) with generalized score (Huang et al., 2018) are able to handle nonparametric cases with assumptions such as causal sufficiency. Both of these approaches rely on kernel methods whose computational complexity is cubic w.r.t. the number of samples. Therefore, the running time could be long if the sample size is large. Furthermore, when the number of variables is large, the search procedure may involve computing the kernel-based conditional independence test or score function many times, which therefore may also increase the running time.

As shown by Loh & Bühlmann (2014); Ng et al. (2021) in the linear Gaussian case, the Markov network (i.e., the support of the inverse covariance matrix of the distribution) is guaranteed to be the super-structure of the ground truth directed acyclic graph (DAG) under a specific type of faithfulness assumption. That is, the super-structure contains all edges of the true DAG. Using this idea, they showed that the Markov network may be used to restrict the search space of score-based approaches for causal discovery, which improves the scalability. Their works focus only on the linear case and adopt classical methods like graphical Lasso (Friedman et al., 2008) to estimate the Markov network. In this work, the nonparametric Markov network estimated by our proposed procedure could potentially be used as a super-structure to restrict the search space for nonparametric causal discovery methods, i.e., (kernel-based) PC and GES. Similar to (Loh & Bühlmann, 2014; Ng et al., 2021), this may help reduce the running time and improve the scalability of these methods.

5 EXPERIMENTS

Setup. Following previous works, we conduct experiments for the continuous case on the Butterfly distribution (Morrison et al., 2017; Baptista et al., 2021), where r i.i.d. pairs of random variables (P_i, Q_i) are defined as $P_i \sim \mathcal{N}(0, 1)$ and $Q_i = W_i P_i$ with $W_i \sim \mathcal{N}(0, 1)$ and $W_i \perp\!\!\!\perp Q_i$. This is a commonly used example for two variables to be uncorrelated but dependent. We replace the Gaussian distribution with the multinomial distribution for the discrete case and mix two different types of pairs for the mixed-type case. The proportion of different types of variables for the mixed case is uniformly sampled. All experiments are on 12 CPU cores with 24 GB RAM.

Considered methods. We consider the following representative methods for comparison: *(KCI)* Kernel-based Conditional Independence test (KCI) (Zhang et al., 2012) is a well-known nonparametric test for conditional independence. We adopt KCI with the Incremental Association Markov Blanket (IAMB) algorithm (Tsamardinos et al., 2003) to learn the Markov network structure in our settings; *(GS)* We denote GS as Greedy Equivalence Search (GES) with Generalilzed Score (GS). Generalized Score (Huang et al., 2018) characterizes general (conditional) independence relationships between random variables without assuming particular model classes. It has been proved that GES search procedure with Generalized Score is asymptotically optimal (Huang et al., 2018). Because this procedure returns CPDAGs, we moralize the results to learn the Markov network structure; *(SING)* Sparsity Identification in Non-Gaussian distributions (SING) (Morrison et al., 2017; Baptista et al., 2021) is an algorithm designed for the estimation of Markov networks in non-Gaussian continuous distributions. It applies a transport map to estimate the data density; *(GLASSO)* Graphical Lasso (GLASSO) is a classical sparse penalized estimator for the inverse covariance matrix.

Results. We first conduct comparisons in general distributions for all data types (i.e., discrete, continuous, and mixed-type) with different numbers of variables and sample sizes. Among the considered methods, both KCI and GS are available for the estimation of Markov network structures for general distributions with all data types. Therefore, we compare them with our method in all settings. SING can only deal with continuous data and is therefore only applied in the continuous setting. GLASSO is efficient but only with Gaussian distributions that its estimation is supported with theoretical guarantees. We adopt it as a baseline in our comparisons. We use Hamming distance between the estimated graph and the ground truth graph as the metric. All results are of 5 trials with different random seeds. The missing results are either due to timeout (i.e., > 1 day) or OOM.

(a) Continuous, $n = 1000$ (b) Discrete, $n = 1000$ (c) Mixed-type, $n = 1000$

(d) Continuous, $n = 10000$ (e) Discrete, $n = 10000$ (f) Mixed-type, $n = 10000$

Figure 1: Hamming distances w.r.t. different settings, sample sizes (n), and numbers of variables.

From Fig. 1 and Table 1, one can observe that KCI, GS, and our method can almost recover the true structures with general distributions in all data types. At the same time, KCI and GS cannot finish running within one day when $n = 10000$, and their running time for graphs with 12 variables are significantly longer than our method. Besides, SING performs well on the continuous case. However, it cannot scale with more than 12 variables when $n = 1000$ (OOM) and 6 variables when $n = 10000$ (OOM or timeout). GLASSO is remarkably fast but fail to accurately recover the structure in the general settings. This is to be expected since GLASSO is designed for Gaussian graphical models.

Table 1: Running time for 12 variables. $n = 1000$.

Method	Ours	KCI	GS	SING	GLASSO
Time (s)	62.9	247	9536.5	4020.3	4.4

We also conduct experiments on large graphs, with the number of continuous variables from 200 to 5000 and a sample size of 1000. From Fig. 2, we can see that the running time is approximately linear w.r.t. the number of variables. Additionally, all experiments are conducted on CPUs while our regualrized score matching-based framework could be easily deployed on GPUs, thus making advantage of recent advances for computation, especially for deep models. This suggests the potential of even further improvement on the scalability of our method.

Figure 2: Running time for large graphs.

6 DISCUSSION

We provide a scalable estimation framework based on regularized score matching for nonparametric Markov network structures. We first introduce necessary and sufficient conditions of conditional independence between variables in general distributions for all data types (i.e., continuous, discrete, and mixed-type) without specific assumptions on functional relations among variables. Besides, we propose the corresponding characterizations of the structure accordingly. Then, we unify all these cases under the same umbrella of the estimation framework based on regularized score matching. Appropriate penalties on the characterization matrix are introduced to promote constantly sparse entries for stable estimation. We validate our theoretical claims experimentally in various settings, and the scalability of our approach is suggested by results on graphs with up to 5000 variables. Finally, we discuss the implication of our results in causal discovery and include the related extension as future work.

REFERENCES

Ricardo Baptista, Youssef Marzouk, Rebecca E Morrison, and Olivier Zahm. Learning non-gaussian graphical models via hessian scores and triangular transport. *arXiv preprint arXiv:2101.03093*, 2021.

Julian Besag. Spatial interaction and the statistical analysis of lattice systems. *Journal of the Royal Statistical Society: Series B (Methodological)*, 36(2):192–225, 1974.

Patrick Breheny and Jian Huang. Coordinate descent algorithms for nonconvex penalized regression, with applications to biological feature selection. *The Annals of Applied Statistics*, 5(1):232–253, 2011.

D Brook. On the distinction between the conditional probability and the joint probability approaches in the specification of nearest-neighbour systems. *Biometrika*, 51(3/4):481–483, 1964.

Peter J Carrington, John Scott, and Stanley Wasserman. *Models and methods in social network analysis*, volume 28. Cambridge university press, 2005.

Jie Cheng, Tianxi Li, Elizaveta Levina, and Ji Zhu. High-dimensional mixed graphical models. *Journal of Computational and Graphical Statistics*, 26(2):367–378, 2017.

David M. Chickering. Optimal structure identification with greedy search. *Journal of Machine Learning Research*, 3(Nov):507–554, 2002.

Giulio Cimini, Tiziano Squartini, Fabio Saracco, Diego Garlaschelli, Andrea Gabrielli, and Guido Caldarelli. The statistical physics of real-world networks. *Nature Reviews Physics*, 1(1):58–71, 2019.

Sheel C Dodani, Gert Kiss, Jackson KB Cahn, Ye Su, Vijay S Pande, and Frances H Arnold. Discovery of a regioselectivity switch in nitrating p450s guided by molecular dynamics simulations and markov models. *Nature chemistry*, 8(5):419–425, 2016.

Mathias Drton, Bernd Sturmfels, and Seth Sullivant. *Lectures on algebraic statistics*, volume 39. Springer Science & Business Media, 2008.

David Edwards. Hierarchical interaction models. *Journal of the Royal Statistical Society: Series B (Methodological)*, 52(1):3–20, 1990.

Jianqing Fan and Runze Li. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American statistical Association*, 96(456):1348–1360, 2001.

Bernd Fellinghauer, Peter Bühlmann, Martin Ryffel, Michael Von Rhein, and Jan D Reinhardt. Stable graphical model estimation with random forests for discrete, continuous, and mixed variables. *Computational Statistics & Data Analysis*, 64:132–152, 2013.

Jerome Friedman, Trevor Hastie, and Robert Tibshirani. Sparse inverse covariance estimation with the graphical lasso. *Biostatistics*, 9(3):432–441, 2008.

Geoffrey R Grimmett. A theorem about random fields. *Bulletin of the London Mathematical society*, 5(1):81–84, 1973.

Naftali Harris and Mathias Drton. Pc algorithm for nonparanormal graphical models. *Journal of Machine Learning Research*, 14(11), 2013.

Biwei Huang, Kun Zhang, Yizhu Lin, Bernhard Schölkopf, and Clark Glymour. Generalized score functions for causal discovery. In *Proceedings of the 24th ACM SIGKDD international conference on knowledge discovery & data mining*, pp. 1551–1560, 2018.

Junzhou Huang and Tong Zhang. The benefit of group sparsity. *The Annals of Statistics*, 38(4): 1978–2004, 2010.

Aapo Hyvärinen and Peter Dayan. Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6(4), 2005.

Ariel Jaimovich, Gal Elidan, Hanah Margalit, and Nir Friedman. Towards an integrated protein–protein interaction network: A relational markov network approach. *Journal of Computational Biology*, 13(2):145–164, 2006.

Steffen L Lauritzen. *Graphical models*, volume 17. Clarendon Press, 1996.

Steffen L Lauritzen, Anders Holst Andersen, David Edwards, Karl G Jöreskog, and Søren Johansen. Mixed graphical association models [with discussion and reply]. *Scandinavian Journal of Statistics*, pp. 273–306, 1989.

Jason D Lee and Trevor J Hastie. Learning the structure of mixed graphical models. *Journal of Computational and Graphical Statistics*, 24(1):230–253, 2015.

Lina Lin, Mathias Drton, and Ali Shojaie. Estimation of high-dimensional graphical models using regularized score matching. *Electronic journal of statistics*, 10(1):806, 2016.

Han Liu, John Lafferty, and Larry Wasserman. The nonparanormal: Semiparametric estimation of high dimensional undirected graphs. *Journal of Machine Learning Research*, 10(10), 2009.

Han Liu, Fang Han, and Cun-hui Zhang. Transelliptical graphical models. *Advances in neural information processing systems*, 25, 2012.

Po-Ling Loh and Peter Bühlmann. High-dimensional learning of linear causal networks via inverse covariance estimation. *Journal of Machine Learning Research*, 15(88):3065–3105, 2014.

Po-Ling Loh and Martin J Wainwright. Structure estimation for discrete graphical models: Generalized covariance matrices and their inverses. *Advances in Neural Information Processing Systems*, 25, 2012.

Po-Ling Loh and Martin J. Wainwright. Structure estimation for discrete graphical models: Generalized covariance matrices and their inverses. *The Annals of Statistics*, 41(6):3022–3049, 2013.

Po-Ling Loh and Martin J. Wainwright. Support recovery without incoherence: A case for nonconvex regularization. *The Annals of Statistics*, 45(6):2455–2482, 2017.

Siwei Lyu. Interpretation and generalization of score matching. *arXiv preprint arXiv:1205.2629*, 2012.

Nicolai Meinshausen and Peter Bühlmann. High-dimensional graphs and variable selection with the lasso. *The annals of statistics*, 34(3):1436–1462, 2006.

Rebecca Morrison, Ricardo Baptista, and Youssef Marzouk. Beyond normality: Learning sparse probabilistic graphical models in the non-gaussian setting. *Advances in neural information processing systems*, 30, 2017.

Ignavier Ng, Yujia Zheng, Jiji Zhang, and Kun Zhang. Reliable causal discovery with improved exact search and weaker assumptions. In *Advances in Neural Information Processing Systems*, 2021.

J. Pearl. *Causality: Models, Reasoning, and Inference*. Cambridge University Press, Cambridge, 2000.

Dinh Tuan Pham and Philippe Garat. Blind separation of mixture of independent sources through a quasi-maximum likelihood approach. *IEEE transactions on Signal Processing*, 45(7):1712–1725, 1997.

Pradeep Ravikumar, Garvesh Raskutti, Martin J Wainwright, and Bin Yu. Model selection in Gaussian graphical models: High-dimensional consistency of ℓ_1-regularized MLE. In *Advances in Neural Information Processing Systems*, 2008.

Pradeep Ravikumar, Martin J Wainwright, and John D Lafferty. High-dimensional ising model selection using 1-regularized logistic regression. *The Annals of Statistics*, 38(3):1287–1319, 2010.

Pradeep Ravikumar, Martin J Wainwright, Garvesh Raskutti, and Bin Yu. High-dimensional covariance estimation by minimizing ℓ_1-penalized log-determinant divergence. *Electronic Journal of Statistics*, 5:935–980, 2011.

Huawei Shen, Xueqi Cheng, Kai Cai, and Mao-Bin Hu. Detect overlapping and hierarchical community structure in networks. *Physica A: Statistical Mechanics and its Applications*, 388(8): 1706–1712, 2009.

Yang Song, Sahaj Garg, Jiaxin Shi, and Stefano Ermon. Sliced score matching: A scalable approach to density and score estimation. In *Uncertainty in Artificial Intelligence*, pp. 574–584. PMLR, 2020.

Alessio Spantini, Daniele Bigoni, and Youssef Marzouk. Inference via low-dimensional couplings. *The Journal of Machine Learning Research*, 19(1):2639–2709, 2018.

Peter Spirtes and Clark Glymour. An algorithm for fast recovery of sparse causal graphs. *Social Science Computer Review*, 9:62–72, 1991.

Peter Spirtes, Clark N Glymour, Richard Scheines, and David Heckerman. *Causation, prediction, and search*. MIT press, 2000.

Arun Suggala, Mladen Kolar, and Pradeep K Ravikumar. The expxorcist: Nonparametric graphical models via conditional exponential densities. *Advances in neural information processing systems*, 30, 2017.

Robert Tibshirani. Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society. Series B (Methodological)*, 58(1):267–288, 1996.

Ioannis Tsamardinos, Constantin F Aliferis, Alexander R Statnikov, and Er Statnikov. Algorithms for large scale markov blanket discovery. In *FLAIRS conference*, volume 2, pp. 376–380. St. Augustine, FL, 2003.

Martin J. Wainwright. Sharp thresholds for high-dimensional and noisy sparsity recovery using ℓ_1 -constrained quadratic programming (Lasso). *IEEE Transactions on Information Theory*, 55(5): 2183–2202, 2009.

Eunho Yang, Pradeep Ravikumar, Genevera I Allen, and Zhandong Liu. Graphical models via univariate exponential family distributions. *The Journal of Machine Learning Research*, 16(1): 3813–3847, 2015.

Ming Yuan. High dimensional inverse covariance matrix estimation via linear programming. *The Journal of Machine Learning Research*, 11:2261–2286, 2010.

Ming Yuan and Yi Lin. Model selection and estimation in regression with grouped variables. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 68(1):49–67, 2006.

Cun-Hui Zhang. Nearly unbiased variable selection under minimax concave penalty. *The Annals of Statistics*, 38(2):894–942, 2010.

Kun Zhang, Jonas Peters, Dominik Janzing, and Bernhard Schölkopf. Kernel-based conditional independence test and application in causal discovery. *arXiv preprint arXiv:1202.3775*, 2012.

Hui Zou. The adaptive lasso and its oracle properties. *Journal of the American statistical association*, 101(476):1418–1429, 2006.

A PROOFS

A.1 PROOF OF COROLLARY 1

Corollary 1. *Assume*

    i. $\mathbf{X} = (X_1, \ldots, X_d)$ *is a set of continuous variable.*

    ii. The PDFs of $\mathbf{X}$ *are strictly positive and smooth.*

    iii. The characterization matrix $\Omega^{[c]}$ *is defined according to Eq.* (3).

Then for any $i \neq j$, $\Omega_{i,j}^{[c]} = 0$ *implies* $X_i \perp\!\!\!\perp X_j \mid \mathbf{X}_{\mathbf{V} \setminus \{i,j\}}$.

Proof. According to Eq. (3), it is clear that when $\Omega_{i,j}^{[c]} = 0$, we have $\frac{\partial^2 \log p_{\mathbf{x}}}{\partial x_i \partial x_j} = 0$, which is a necessary and sufficient condition of $x_i \perp\!\!\!\perp x_j \mid x_{\mathbf{V} \setminus \{i,j\}}$ for strictly positive, smooth, and continuous distributions as shown in Sec. 2.1.

It is worth noting that it also applies in the Gaussian case, where $\mathbf{X} \sim \mathcal{N}(\mu, \Sigma)$ is a Gaussian vector with mean μ and non-singular covariance Σ. In this case, we have

$$P_{\mathbf{X}}(\mathbf{x}) \propto \exp\left(\frac{-(\mathbf{x} - \mu)^\top \Sigma^{-1} (\mathbf{x} - \mu)}{2}\right), \tag{16}$$

which implies

$$\frac{\partial^2 \log p_{\mathbf{X}}}{\partial x_i \partial x_j} = (\Sigma^{-1})_{i,j}^2, \tag{17}$$

where $(\Sigma^{-1})_{i,j}$ denotes the corresponding entry of the inverse covariance matrix. This well-known property of Gaussian distribution was also shown in Baptista et al. (2021); Drton et al. (2008). Because the inverse covariance matrix encodes the conditional independence structure when variables are from Gaussian distributions, $\Omega^{[c]}$ characterizes the Markov property for the Gaussian case. $\square$

A.2 PROOF OF THEOREM 1

Theorem 1. *Denote* $\mathbf{V}$ *as a set of discrete variables and* $X_i, X_j \in \mathbf{V}$. *For brevity, denote* $\mathbf{V} \setminus \{X_i, X_j\}$ *as* $\mathbf{Z}$. *Let* $\{x_{i1}, \ldots, x_{iM_i}\}$ *and* $\{x_{j1}, \ldots, x_{jM_j}\}$ *be the support of variables* X_i *and* X_j. *Denote* z *as any value(s) of* $\mathbf{Z}$. *Then,* $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$ *if and only if, for all* $k \in [M_i]$ *and* $l \in [M_j]$ *with* $k \neq 1$ *and* $l \neq 1$, *we have*

$$(\log m(x_{i1}, x_{j1}, z) - \log m(x_{ik}, x_{j1}, z)) - (\log m(x_{i1}, x_{jl}, z) - \log m(x_{ik}, x_{jl}, z)) = 0. \tag{4}$$

Proof. **Sufficient condition.** Without loss of generality, let us consider three discrete variables, i.e., $\{X_i, X_j, Z\}$. Let $\{x_{i1}, \ldots, x_{iM_i}\}$ and $\{x_{j1}, \ldots, x_{jM_j}\}$ be the support of variables X_i and X_j, respectively.

Consider the case that the finite difference of the discrete score function of X_i w.r.t. X_j equals zero.

When $X_i = x_{i1}$, $X_j = x_{j1}$, and differences are considered w.r.t. to $x_{ik'}$ and $x_{jl'}$, we have

$$\begin{aligned}(\log m(x_{i1}, x_{j1}, z) - \log m(x_{ik'}, x_{j1}, z)) \\ - (\log m(x_{i1}, x_{jl'}, z) - \log m(x_{ik'}, x_{jl'}, z)) = 0,\end{aligned} \tag{18}$$

where $m(x_{i1}, x_{j1}, z)$ is the joint PMF simplified from $m_{X_i, X_j, \mathbf{Z}}\{x_{i1}, x_{j1}, z\}$. By iterating all possible differences w.r.t. X_j, for all l in $\{2, \ldots, M_j\}$, we have

$$\begin{aligned}(\log m(x_{i1}, x_{j1}, z) - \log m(x_{ik'}, x_{j1}, z)) \\ - (\log m(x_{i1}, x_{jl}, z) - \log m(x_{ik'}, x_{jl}, z)) = 0.\end{aligned} \tag{19}$$

Define the discrete score function of X_i as $g(x_{i1}, x_{ik'}, \gamma) = \log m(x_{i1}, \gamma) - \log m(x_{ik'}, \gamma)$, where γ denotes other variables. Eq. 19 means $g(x_{i1}, x_{ik'}, \gamma)$ doe not take the value of X_j as an argument when the LHS of Eq. 19 equals zero. As a result, Eq. 19 could be formulated as

$$\log m(x_{i1}, x_{j1}, z) = \log m(x_{ik'}, x_{j1}, z) + g(x_{i1}, x_{ik'}, \gamma). \tag{20}$$

Then by iterating all possible differences w.r.t. X_i, for all k in $I_{x_i} = \{2, \ldots, M_i\}$, we have

$$\log m(x_{i1}, x_{j1}, z) = \log m(x_{ik}, x_{j1}, z) + g(x_{i1}, x_{ik}, \gamma). \tag{21}$$

By summation, we have

$$\begin{aligned}
(N-1) \log m(x_{i1}, x_{j1}, z) &= \sum_{k \in I_{x_i}} \left(\log m(x_{ik}, x_{j1}, z) + g(x_{i1}, x_{ik}, \gamma) \right) \\
&= \sum_{k \in I_{x_i}} \log m(x_{ik}, x_{j1}, z) + \sum_{k \in I_{x_i}} g(x_{i1}, x_{ik}, \gamma),
\end{aligned} \tag{22}$$

which implies that

$$\begin{aligned}
M_i \log m(x_{i1}, x_{j1}, z) &= \sum_{k=1}^{M_i} \log m(x_{ik}, x_{j1}, z) + \sum_{k \in I_{x_i}} g(x_{i1}, x_{ik}, \gamma), \\
\log m(x_{i1}, x_{j1}, z) &= \frac{1}{M_i} \left(\sum_{k=1}^{M_i} \log m(x_{ik}, x_{j1}, z) + \sum_{k \in I_{x_i}} g(x_{i1}, x_{ik}, \gamma) \right).
\end{aligned} \tag{23}$$

Because $\sum_{k=1}^{M_i} \log m(x_{ik}, x_{j1}, z)$ covers all possible values of X_i, this term does not depend on the specific value of X_i. Besides, the other term $\sum_{k \in I_{x_i}} g(x_{i1}, x_{ik}, \gamma)$ does not depend on X_j. It is worth noting that X_{i1} could be any value of X_i w.o.l.g.. Therefore, we could see that when the finite difference of the discrete score function of X_i w.r.t. to X_j equal to zero (after some aggregation of samples), $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$.

Necessary condition. When $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$, we could decompose $m(x_i, x_j, z)$ as $m_{X_i \mid \mathbf{Z}}(x_i \mid z) m_{X_j \mid \mathbf{Z}}(x_j \mid z) m_{\mathbf{Z}}(z)$. This implies that, for all k in $\{2, \ldots, M_i\}$ and l in $\{2, \ldots, M_j\}$, we have

$$\begin{aligned}
&(\log m(x_{i1}, x_{j1}, z) - \log m(x_{ik}, x_{j1}, z)) \\
&\quad - (\log m(x_{i1}, x_{jl}, z) - \log m(x_{ik}, y_{jl}, z)) \\
&= \big(\log \left(m_{X_i \mid \mathbf{Z}}(x_{i1} \mid z) m_{X_j \mid \mathbf{Z}}(x_{j1} \mid z) m_{\mathbf{Z}}(z) \right) \\
&\quad - \log \left(m_{X_i \mid \mathbf{Z}}(x_{ik} \mid z) m_{X_j \mid \mathbf{Z}}(x_{j1} \mid z) m_{\mathbf{Z}}(z) \right) \\
&\quad - \big(\log \left(m_{X_i \mid \mathbf{Z}}(x_{i1} \mid z) m_{X_j \mid \mathbf{Z}}(x_{jl} \mid z) m_{\mathbf{Z}}(z) \right) \\
&\qquad - \log \left(m_{X_i \mid \mathbf{Z}}(x_{ik} \mid z) m_{X_j \mid \mathbf{Z}}(x_{jl} \mid z) m_{\mathbf{Z}}(z) \right) \big) \\
&= \big((\log m_{X_i \mid \mathbf{Z}}(x_{i1} \mid z) + \log m_{X_j \mid \mathbf{Z}}(x_{j1} \mid z) + \log m_{\mathbf{Z}}(z)) \\
&\quad - (\log m_{X_i \mid \mathbf{Z}}(x_{ik} \mid z) + \log m_{X_j \mid \mathbf{Z}}(x_{j1} \mid z) + \log m_{\mathbf{Z}}(z)) \big) \\
&\quad - \big((\log m_{X_i \mid \mathbf{Z}}(x_{i1} \mid z) + \log m_{X_j \mid \mathbf{Z}}(x_{jl} \mid z) + \log m_{\mathbf{Z}}(z)) \\
&\qquad - (\log m_{X_i \mid \mathbf{Z}}(x_{ik} \mid z) + \log m_{X_j \mid \mathbf{Z}}(x_{jl} \mid z) + \log m_{\mathbf{Z}}(z)) \big) \\
&= 0.
\end{aligned} \tag{24}$$

Therefore, when $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$, the finite difference of the discrete score function of X_i w.r.t. to X_j equals zero.

The proof is complete. □

A.3 PROOF OF COROLLARY 2

Corollary 2. *Assume*

    i. $\mathbf{X} = (X_1, \ldots, X_d)$ *is a set of discrete variable.*

    ii. The PMFs of $\mathbf{X}$ are strictly positive.

    iii. The characterization matrix $\Omega^{[d]}$ is defined according to Eq. (5).

Then for any $i \neq j$, $\Omega_{i,j}^{[d]} = 0$ implies $\mathbf{x}_i \perp\!\!\!\perp \mathbf{x}_j \mid \mathbf{x}_{\mathbf{V} \setminus \{i,j\}}$.

Proof. According to Eq. (5), we have

$$\Omega_{i,j}^{[d]} := \mathbb{E}_m \left[\sum_{k,l} f^{[d]}(x_{i1}, x_{ik}, x_{j1}, x_{jl}, z)^2 \right], \tag{25}$$

where $f^{[d]}(x_{i1}, x_{ik}, x_{j1}, x_{jl}, z)$ denotes the LHS of Eq. (4), i.e.,

$$\begin{aligned}
& f^{[d]}(x_{i1}, x_{ik}, x_{j1}, x_{jl}, z) \\
& = (\log m(x_{i1}, x_{j1}, z) - \log m(x_{ik}, x_{j1}, z)) \\
& \quad - (\log m(x_{i1}, x_{jl}, z) - \log m(x_{ik}, x_{jl}, z)).
\end{aligned} \tag{26}$$

Thus, if $\Omega_{i,j}^{[d]} = 0$, we must have

$$\begin{aligned}
& (\log m(x_{i1}, x_{j1}, z) - \log m(x_{ik}, x_{j1}, z)) \\
& \quad - (\log m(x_{i1}, x_{jl}, z) - \log m(x_{ik}, x_{jl}, z)) = 0,
\end{aligned} \tag{27}$$

for all $k \in [M_i]$ and $l \in [M_j]$, where M_i and M_j denote the cardinalities of X_i and X_j, respectively. Based on Theorem. 1, we have $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$.

The proof is complete. $\qquad\square$

A.4 PROOF OF THEOREM 2

Theorem 2. *Denote $\mathbf{V}$ as a set of mixed-type variables and $X_i, X_j \in \mathbf{V}$, where X_i is discrete and X_j is continuous. Let $\{x_{i1}, \ldots, x_{iM_i}\}$ be the support of variables X_i. For brevity, denote $\mathbf{V} \backslash \{X_i, X_j\}$ as $\mathbf{Z}$. Denote z as any value(s) of $\mathbf{Z}$ and x_j as any value of the continuous variable X_j. Then, $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$ if and only if, for all $k \in [M_i]$ with $k \neq 1$, we have*

$$\frac{\partial \log \left(p_{X_j, \mathbf{Z}|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right)}{\partial x_j} - \frac{\partial \log \left(p_{X_j, \mathbf{Z}|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right)}{\partial x_j} = 0. \tag{6}$$

Proof. **Sufficient condition.** Without loss of generality, let us consider three variables, i.e., $\{X_i, X_j, Z\}$:

$$\begin{aligned}
x_i & \in \{x_{i1}, \ldots, x_{iM_i}\} \\
x_j & \in \mathbb{R},
\end{aligned} \tag{28}$$

where we set $X_i = x_i$ as the discrete variable and $X_j = x_j$ as the continuous variable w.l.o.g. Note that we do not constraint the type of $Z = z$ here but set Z as continuous for brevity.

Consider the case that the finite difference of the score function of X_j w.r.t. X_i equals zero. Also, we define p as the p.d.f. and m as the p.m.f.. We first consider the difference between x_{i1} and $x_{ik'}$.

$$\frac{\partial \log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right)}{\partial x_j} - \frac{\partial \log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{ik'}) m_{X_i}(x_{ik'}) \right)}{\partial x_j} = 0. \tag{29}$$

By iterating all possible differences w.r.t. x_i, for all k in $I_{x_i} = \{2, \ldots, M\}$, we have

$$\frac{\partial \log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right)}{\partial x_j} - \frac{\partial \log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right)}{\partial x_j} = 0, \tag{30}$$

which is equivalent to

$$\frac{\partial \log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right)}{\partial x_j} = \frac{\partial \log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right)}{\partial x_j}. \tag{31}$$

Then by integrating on both sides w.r.t. X_j, we have

$$\log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right) = \log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right) + C_k, \tag{32}$$

where C_k is a constant. We then apply a summation as follows

$$\begin{aligned}
& (M_i - 1) \log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right) \\
& = \sum_{k \in I_{x_i}} \left(\log \left(p_{X_j, Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right) + C_k \right),
\end{aligned} \tag{33}$$

which implies that

$$\log \left(p_{X_j,Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right)$$

$$= \frac{1}{M_i} \left(\sum_{k=1}^{M_i} \log \left(p_{X_j,Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right) + \sum_{k \in I_{x_i}} C_k \right). \tag{34}$$

Because $\sum_{k=1}^{M_i} \log \left(p_{X_j,Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right)$ covers all possible values of k, this term does not depend on the specific value of X_i. Besides, C_k does not depend on X_j. Therefore, by iterating all possible differences of X_i, we could see that when the finite difference of the score function of X_j w.r.t. X_i equals zero (after some aggregation of samples), $X_i \perp\!\!\!\perp X_j \mid Z$.

It is noteworthy that another "symmetric" case, where the derivative of the discrete score function of X_i w.r.t. X_j equals zero, is as follows

$$\frac{\partial \left(\log \left((p_{X_j,Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1})) - \log \left(p_{X_j,Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right) \right) \right)}{\partial x_j} = 0, \tag{35}$$

which is equivalent to

$$\frac{\partial \log \left(p_{X_j,Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right)}{\partial x_j} - \frac{\partial \log \left(p_{X_j,Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right)}{\partial x_j} = 0. \tag{36}$$

Thus, we only need to consider Eq. 36, which is the case that the finite difference of the discrete score function of X_i w.r.t. X_j equals zero.

Necessary condition. When $X_i \perp\!\!\!\perp X_j \mid Z$, we could decompose $p_{X_j,Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1})$ as $m_{X_i|Z}(x_{i1} \mid z) p_{X_j|Z}(x_j \mid z) p_Z(z)$. This implies that, for all k in $\{2, \ldots, M_i\}$, we have

$$\frac{\partial \log \left(p_{X_j,Z|X_i}(x_j, z \mid x_{i1}) m_{X_i}(x_{i1}) \right)}{\partial x_j} - \frac{\partial \log \left(p_{X_j,Z|X_i}(x_j, z \mid x_{ik}) m_{X_i}(x_{ik}) \right)}{\partial x_j}$$

$$= \frac{\partial \log \left(m_{X_i|Z}(x_{i1} \mid z) p_{X_j|Z}(x_j \mid z) p_Z(z) \right)}{\partial x_j}$$

$$\quad - \frac{\partial \log \left(m_{X_i|Z}(x_{ik} \mid z) p_{X_j|Z}(x_j \mid z) p_Z(z) \right)}{\partial x_j}$$

$$= \frac{\partial \left(\log m_{X_i|Z}(x_{i1} \mid z) + \log p_{X_j|Z}(x_j \mid z) + \log p_Z(z) \right)}{\partial x_j} \tag{37}$$

$$\quad - \frac{\partial \left(\log m_{X_i|Z}(x_{ik} \mid z) + \log p_{X_j|Z}(x_j \mid z) + \log p_Z(z) \right)}{\partial x_j}$$

$$= \frac{\partial \log p_{X_j|Z}(x_j \mid z)}{\partial x_j} - \frac{\partial \log p_{X_j|Z}(x_j \mid z)}{\partial x_J}$$

$$= 0.$$

Therefore, when $X_i \perp\!\!\!\perp X_j \mid Z$, the finite difference of the score function of X_j w.r.t. to X_i equal to zero.

The proof is complete. $\qquad \square$

A.5 Proof of Corollary 3

Corollary 3. *Assume*

i. $\mathbf{X} = (X_1, \ldots, X_d)$ *is a set of variables containing both continuous and discrete variables.*

ii. For continuous variables, the PDFs are strictly positive and smooth.

iii. For discrete variables, the PMFs are strictly positive.

iv. The characterization matrix $\Omega^{[m]}$ is defined according to Eq. (7).

Then for any $i \neq j$, $\Omega^{[m]}_{i,j} = 0$ implies $X_i \perp\!\!\!\perp X_j \mid \mathbf{X}_{\mathbf{V}\setminus\{i,j\}}$.

Proof. According to Eq. (7), we have

$$
\Omega^{[m]}_{i,j} := \begin{cases}
\mathbb{E}_\pi\left[f^{[c]}_{i,j}(\mathbf{x})^2\right] & \text{if } X_i \in \mathbf{X}_c, X_j \in \mathbf{X}_c \\
\mathbb{E}_\pi\left[\sum_{k,l} f^{[d]}(x_{i1}, x_{ik}, x_{j1}, x_{jl}, z)^2\right] & \text{if } X_i \in \mathbf{X}_d, X_j \in \mathbf{X}_d \\
\mathbb{E}_\pi\left[\sum_k f^{[m]}(x_i, x_{j1}, x_{jk}, z)^2\right] & \text{if } X_i \in \mathbf{X}_c, X_j \in \mathbf{X}_d \\
\mathbb{E}_\pi\left[\sum_k f^{[m]}(x_j, x_{i1}, x_{ik}, z)^2\right] & \text{if } X_i \in \mathbf{X}_d, X_j \in \mathbf{X}_c,
\end{cases}
\tag{38}
$$

where $\mathbf{X}_c$ and $\mathbf{X}_d$ are the sets of continuous and discrete variables, respectively. We have already proved the first two cases (i.e., $\{X_i \in \mathbf{X}_c, X_j \in \mathbf{X}_c\}$ and $\{X_i \in \mathbf{X}_d, X_j \in \mathbf{X}_d\}$) in the proofs of Cor. 1 and Cor. 2, respectively. So here we will focus on the other two cases. We start from the third case, where $\{X_i \in \mathbf{X}_c, X_j \in \mathbf{X}_d\}$. We have

$$
f^{[m]}(x_i, x_{j1}, x_{jk}, z) = \frac{\partial \log\left(p_{X_i, \mathbf{Z}|X_j}(x_i, z \mid x_{j1}) m_{X_j}(x_{j1})\right)}{\partial x_i}
$$
$$
- \frac{\partial \log\left(p_{X_i, \mathbf{Z}|X_j}(x_i, z \mid x_{jk}) m_{X_j}(x_{jk})\right)}{\partial x_i}.
\tag{39}
$$

Thus, if $\Omega^{[m]}_{i,j} = 0$ for $\{X_i \in \mathbf{X}_c, X_j \in \mathbf{X}_d\}$, we must have

$$
\frac{\partial \log\left(p_{X_i, \mathbf{Z}|X_j}(x_i, z \mid x_{j1}) m_{X_j}(x_{j1})\right)}{\partial x_i}
$$
$$
- \frac{\partial \log\left(p_{X_i, \mathbf{Z}|X_j}(x_i, z \mid x_{jk}) m_{X_j}(x_{jk})\right)}{\partial x_i} = 0,
\tag{40}
$$

for all $k \in [M_j]$, where M_j denotes the cardinality of X_j. Thus, according to Theorem 2, we have $X_i \perp\!\!\!\perp X_j \mid \mathbf{Z}$ if $\Omega^{[m]}_{i,j} = 0$ for $\{X_i \in \mathbf{X}_c, X_j \in \mathbf{X}_d\}$.

The similar derivation applies for the last case, where $\{X_i \in \mathbf{X}_d, X_j \in \mathbf{X}_c\}$. $\qquad\square$

A.6 PROOF OF COROLLARY 4

We first introduce the following lemmas and their proofs for completeness.

A.6.1 PROOF OF LEMMA 1

Lemma 1. *[directly from Thm. 1 in (Hyvärinen & Dayan, 2005)] Assume*

i. $\mathbf{X} = (X_1, \ldots, X_d)$ *is a set of continuous variables.*

ii. The data PDF $p_{\mathbf{X}}(\mathbf{x})$ is differentiable. The model PDF $(\mathbf{x}; \theta)$ is twice-differentiable. Both of them are strictly positive.

iii. The expectations $E_{\mathbf{x}}\left\{\|\log p(\mathbf{x}; \theta)\|^2\right\}$ and $E_{\mathbf{x}}\left\{\|\log p_{\mathbf{X}}(\mathbf{x})\|^2\right\}$ are finite for any θ, and $p_{\mathbf{X}}(\mathbf{x}) \log p(\mathbf{x}; \theta)$ goes to zero for any θ when $\|\mathbf{x}\| \to \infty$.

Then Eq. (8) is equivalent to

$$
O_c(\theta) = \int_{\mathbf{x} \in \mathbb{R}^n} p_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \left[\frac{1}{2}\|\nabla_{x_i} \log p(\mathbf{x}; \theta)\|^2 + H_{x_i}(\log p(\mathbf{x}; \theta))\right] d\mathbf{x} + \rho_\lambda(\Omega^{[c]}).
\tag{41}
$$

Proof. Based on Eq. (8), we have

$$
O_c(\theta) = \frac{1}{2} \int p_{\mathbf{x}}(\mathbf{x}) |\nabla_{\mathbf{x}} \log p(\mathbf{x}; \theta) - \nabla_{\mathbf{x}} \log p_{\mathbf{x}}(\mathbf{x})|^2 d\mathbf{x} + \rho_\lambda(\Omega^{[c]}),
\tag{42}
$$

where $\rho_\lambda(\cdot)$ denotes a sparsity penalty function and λ is the penalty parameter. This is equivalent to

$$O_c(\theta) = \frac{1}{2} \int p_{\mathbf{X}}(\mathbf{x}) \left(\|\nabla_{\mathbf{x}} \log p(\mathbf{x}; \theta)\|^2 + \|\nabla_{\mathbf{x}} \log p_{\mathbf{X}}(\mathbf{x})\|^2 \right.$$
$$\left. - 2 \left(\nabla_{\mathbf{x}} \log p_{\mathbf{X}}(\mathbf{x})\right)^\top \left(\nabla_{\mathbf{x}} \log p(\mathbf{x}; \theta)\right) d\mathbf{x} + \rho_\lambda(\Omega^{[c]}). \tag{43}$$

We first consider the integral for the following part

$$- \int p_{\mathbf{X}}(\mathbf{x}) \left(\nabla_{\mathbf{x}} \log p_{\mathbf{X}}(\mathbf{x})\right)^\top \left(\nabla_{\mathbf{x}} \log p(\mathbf{x}; \theta)\right) d\mathbf{x}, \tag{44}$$

by which we could obtain

$$- \sum_i \int p_{\mathbf{X}}(\mathbf{x}) \left(\nabla_{x_i} \log p_{\mathbf{X}}(\mathbf{x})\right) \left(\nabla_{x_i} \log p(\mathbf{x}; \theta)\right) d\mathbf{x}$$
$$= - \sum_i \int \left(\nabla_{x_i} p_{\mathbf{X}}(\mathbf{x})\right) \left(\nabla_{x_i} \log p(\mathbf{x}; \theta)\right) d\mathbf{x}$$
$$= - \sum_i \int \left[\int \nabla_{x_i} p_{\mathbf{X}}(\mathbf{x}) \left(\nabla_{x_i} \log p(\mathbf{x}; \theta)\right) dx_1 \right] d(x_1, \dots, x_d) \tag{45}$$
$$\overset{(\star)}{=} - \sum_i \int \left[\lim_{a \to \infty, b \to -\infty} \left[p_{\mathbf{X}}(a, x_2, \dots, x_d) \nabla_{x_i} \log p(a, x_2, \dots, x_d, \theta) \right. \right.$$
$$- p_{\mathbf{X}}(b, x_2, \dots, x_n) \nabla_{x_i} \log p(b, x_2, \dots, x_d, \theta)]$$
$$\left. - \int \frac{\partial^2 \log p_{\mathbf{X}}}{\partial x_i{}^2} p_{\mathbf{X}}(\mathbf{x}) dx_1 \right] d(x_2, \dots, x_d),$$

where Eq. $(\star)$ is because if we assume f and g are both differential, we have

$$\frac{\partial f(\mathbf{x}) g(\mathbf{x})}{\partial x_i} = f(\mathbf{x}) \frac{\partial g(\mathbf{x})}{\partial x_1} + g(\mathbf{x}) \frac{\partial f(\mathbf{x})}{\partial x_1}. \tag{46}$$

For $i \neq 1$, the cases follow similarly. Because we assume $p_{\mathbf{X}}(\mathbf{x}) \log p(\mathbf{x}; \theta)$ goes to zero for any θ when $\|\mathbf{x}\| \to \infty$, the limit is zero. Thus, we have proven that

$$- \sum_i \int p_{\mathbf{X}}(\mathbf{x}) \nabla_{x_i} \log p_{\mathbf{X}}(\mathbf{x}) \nabla_{x_i} \log p(\mathbf{x}; \theta)) d\mathbf{x} = \sum_i \int \frac{\partial^2 \log p_{\mathbf{X}}}{\partial x_i{}^2} p_{\mathbf{X}}(\mathbf{x}) d\mathbf{x}, \tag{47}$$

By injecting it into Eq. (43), we obtain

$$O_c(\theta) = \int p_{\mathbf{X}}(\mathbf{x}) \left[\frac{1}{2} \|\nabla_{\mathbf{x}} \log p(\mathbf{x}; \theta)\|^2 + \frac{1}{2} \|\nabla_{\mathbf{x}} \log p_{\mathbf{X}}(\mathbf{x})\|^2 \right.$$
$$\left. + \text{tr}\left(H_{\mathbf{x}}(\log p(\mathbf{x}; \theta))\right) \right] d\mathbf{x} + \rho_\lambda(\Omega^{[c]}). \tag{48}$$

Because $\frac{1}{2} \|\nabla_{\mathbf{x}} \log p_{\mathbf{X}}(\mathbf{x})\|^2$ does not depend on θ, we could ignore it. Then we have

$$O_c(\theta) = \int \sum_{i=1}^d \left[\frac{1}{2} \|\nabla_{x_i} \log p(\mathbf{x}; \theta)\|^2 + H_{x_i}(\log p(\mathbf{x}; \theta)) \right] d\mathbf{x} + \rho_\lambda(\Omega^{[c]}). \tag{49}$$

Thus, the proof is complete. $\square$

A.6.2 PROOF OF LEMMA 2

Lemma 2. *[directly from (Lyu, 2012)] Assume*

  i. $\mathbf{X} = (X_1, \dots, X_d)$ *is a set of discrete variables.*

  ii. The data PMF $m_{\mathbf{X}}(\mathbf{x})$ *and the model PMF* $m(\mathbf{x}; \theta)$ *are strictly positive.*

Then Eq. (12) *is equivalent to*

$$O_d(\theta) = \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \sum_{x_{ij}} \left(\frac{m(\sim x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)}{m(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)} \right)^2 + \rho_\lambda(\Omega^{[d]}), \tag{50}$$

where $m(\sim x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)$ *denotes the conditional PMF for the i-th entry of $\mathbf{x}$ taking values other than* x_{ij}.

Proof. Based on Eq. (12), we have

$$O_d(\theta) = \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \left\| \frac{\mathcal{M}(m(\mathbf{x}; \theta))}{m(\mathbf{x}; \theta)} - \frac{\mathcal{M}(m_{\mathbf{X}}(\mathbf{x}))}{m_{\mathbf{X}}(\mathbf{x})} \right\|^2 + \rho_\lambda(\Omega^{[d]}), \tag{51}$$

which implies

$$
\begin{aligned}
O_d(\theta) &= \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \left\| \frac{\mathcal{M}_i(m(\mathbf{x}; \theta))}{m(\mathbf{x}; \theta)} - \frac{\mathcal{M}_i(m_{\mathbf{X}}(\mathbf{x}))}{m_{\mathbf{X}}(\mathbf{x})} \right\|^2 + \rho_\lambda(\Omega^{[d]}) \\
&\overset{(\star)}{=} \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \left[\left(\frac{\mathcal{M}_i(m(\mathbf{x}; \theta))}{m(\mathbf{x}; \theta)} \right)^2 - 2\mathcal{M}_i \left(\frac{\mathcal{M}_i(m(\mathbf{x}; \theta))}{m(\mathbf{x}; \theta)} \right) \right] + \rho_\lambda(\Omega^{[d]}) \\
&= \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \sum_{x_{ij}} \left[\frac{1}{m\left(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta\right)^2} - \frac{2}{m\left(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta\right)} \right] + \rho_\lambda(\Omega^{[d]}) \\
&= \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \sum_{x_{ij}} \frac{1 - 2m\left(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta\right)}{m\left(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta\right)^2} + \rho_\lambda(\Omega^{[d]}) \\
&= \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \sum_{x_{ij}} \left(\frac{1 - 2m\left(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta\right) + m\left(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta\right)^2}{m\left(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta\right)^2} - 1 \right) + \rho_\lambda(\Omega^{[d]}) \\
&= \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \sum_{x_{ij}} \left(\frac{1 - m(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)}{m(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)} \right)^2 - M_i d + \rho_\lambda(\Omega^{[d]}) \\
&\overset{(\circ)}{=} \sum_{\mathbf{x}} m_{\mathbf{X}}(\mathbf{x}) \sum_{i=1}^{d} \sum_{x_{ij}} \left(\frac{m(\sim x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)}{m(x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)} \right)^2 + \rho_\lambda(\Omega^{[d]}),
\end{aligned}
\tag{52}
$$

where $m(\sim x_{ij} \mid \mathbf{x}^{\backslash i}; \theta)$ *denotes the conditional PMF for the i-th entry of $\mathbf{x}$ taking values other than* x_{ij}. *We denote the cardinality of X_i as M_i. Eq. $(\star)$ is due to that $\left(\frac{\mathcal{M}_i(m_{\mathbf{X}}(\mathbf{x}))}{m_{\mathbf{X}}(\mathbf{x})} \right)^2$ does not take θ as an argument. Eq. $(\circ)$ is because $M_i d$ is a constant.*

The proof is complete. $\qquad\square$

Then the corollary follows from these lemmas, which is included as follows.

Corollary 4. *Assume*

    i. *The data density $\pi_{\mathbf{X}}(\cdot)$ is equal to $\pi(\cdot; \theta^*)$ for some θ^*.*

    ii. *The data density $\pi_{\mathbf{X}}(\cdot)$ and model density $\pi(\cdot; \theta)$ are strictly positive. $\pi_{\mathbf{X}}(\cdot)$ and $\pi(\cdot; \theta)$ is differentiable and twice-differentiable, respectively, w.r.t. continuous variables. For some θ^*, $\pi_{\mathbf{X}}(\cdot) = \pi(\cdot; \theta^*)$ and no other parameter value gives a density that is equal to $\pi(\cdot; \theta^*)$ almost everywhere.*

    iii. *The expectations $E_{\mathbf{x}} \left[\| \log \pi(\mathbf{x}; \theta) \|^2 \right]$ and $E_{\mathbf{x}} \left[\| \log \pi_{\mathbf{X}}(\mathbf{x}) \|^2 \right]$ are finite for any θ, and $\pi_{\mathbf{X}}(\mathbf{x}) \log \pi(\mathbf{x}; \theta)$ goes to zero for any θ when $\|\mathbf{x}\| \to \infty$.*

Then $O'_m(\theta) = 0$ implies $\theta = \theta^*$.

Proof. The $O'_m(\theta)$ is defined as follows

$$O'_m(\theta) = \mathbb{E}_{\mathbf{x}}\left[\sum_i s_i(\mathbf{x};\theta)\right], \tag{53}$$

where

$$s_i(\mathbf{x};\theta) \coloneqq \begin{cases} \frac{1}{2}\|\nabla_{x_i}\log\pi(\mathbf{x};\theta)\|^2 + H_{x_i}(\log\pi(\mathbf{x};\theta)) & X_i \in \mathbf{X}_c \\ \frac{1}{2}\sum_{x_{ij}}\left(\frac{\pi(\sim x_{ij}|\mathbf{x}^{\backslash i};\theta)}{\pi(x_{ij}|\mathbf{x}^{\backslash i};\theta)}\right)^2 & X_i \in \mathbf{X}_d, \end{cases} \tag{54}$$

where the probability function π is strictly positive. According to Lemma 1 and Lemma 2, both cases of $s_i(\cdot;\theta)$ in $O'_m(\theta)$ are equivalent to $\frac{1}{2}\mathbb{E}\left[\left\|\frac{g(\pi(\cdot;\theta))}{\pi(\cdot;\theta)} - \frac{g(\pi(\cdot))}{\pi(\cdot)}\right\|^2\right]$, where g denotes the gradient operator for continuous variables or the marginalization operator for discrete variables. If $O'_m(\theta) = 0$, $s_i(x;\theta)$ must equal to zero for any i. Because of Brook's Lemma (Brook, 1964), which is also included in Lyu (2012) as Lemma 3, the marginalization operator $\mathcal{M}$ is complete (Defn. 1). Thus, for the discrete variables, we could replace the gradient in the continuous score function with the marginalization operator while preserving local consistency as that for the continuous variables, which is shown by Theorem 2 in Hyvärinen & Dayan (2005). □

B SUPPLEMENTARY EXPERIMENTAL RESULTS

To explore the effect of different regularization functions, we compare the results of our method with different sparsity penalties, which are shown in Figure 3. The experiments are conducted with the number of continuous variables ranging from 4 to 20 and the sample size of 1000. One could observe that SCAD and MCP outperform other penalties, while SCAD performs slightly better than MCP in general. Adaptive ℓ_1 (Zou, 2006) also illustrates its advantage compared to the original ℓ_1 penalty. This suggests the importance of appropriate penalty functions.

Figure 3: Hamming distances w.r.t. different sparsity penalty functions and numbers of variables, $n = 1000$.