Distributed inference for high-dimensional convoluted rank regression

Abstract

Convoluted rank regression is recently developed as a powerful tool to deal with outliers and heavy-tailed noise data. However, there is still a lack of suitable methods for convoluted rank regression in a distributed system, which is becoming very common nowadays. To solve this issue, we investigate the distributed inference of the convoluted rank regression for high-dimensional data. We introduce two debiased estimators which are communication-efficient and computation-efficient, respectively. For these two introduced estimators, we derive the asymptotic representations and establish the Gaussian approximation results. To further reduce ℓ_1 and ℓ_2 errors, we also introduce sparse debiased estimators. Finally, simulation studies and real data analysis are conducted to illustrate our proposed methods.

Keywords: Debiased estimator, distributed data, high-dimensional convoluted rank regression

Mathematics Subject Classification (2020): 62J05

1 Introduction

With the rapid advancement of information technology, modern datasets are often characterized not only by high dimensionality but also by sheer size. Conventional centralized methods, where all datasets are stored and processed in a central computing facility, are often impractical in applications. Privacy or security constraints also make it challenging to process the entire dataset on a single central machine. Consequently, distributed computing systems become increasingly popular. Large-scale high-dimensional datasets are often characterized by outliers and heavy-tailed noise, posing new computational and storage challenges to conventional statistical methods. We are dedicated to advancing robust statistical inference procedures for high-dimensional distributed datasets.

There are many developments to deal with outliers and heavy-tailed noise. Actually quantile regression and Huber regression are two typical choices for robustness considerations. Recent developments about these two regression models can be found in Belloni and Chernozhukov (2011), Wang et al. (2012), Fan et al. (2017), and Sun et al. (2020). Besides, rank regression recently receives a lot of attention due to its both robustness and high efficiency. Wang et al. (2020) made a deep theoretical investigation of high-dimensional sparse penalized rank regression. The non-smooth nature of the rank regression loss function limits the rank regression's popularity. Notably, Zhou et al. (2024) adopted the convoluted smoothing technique (Fernandes et al., 2021) and introduced a novel convoluted rank regression (CRR). They further proposed

corresponding penalized estimators and derived their estimation error bounds. Since the penalized convoluted rank regression estimators in Zhou et al. (2024) cannot be directly used to make inference, Cai et al. (2025) further introduced a debiased estimator and developed inference procedures. However, when the sample size is very large, the computation of high-dimensional convoluted rank regression is still a big problem.

Directly aggregating all data encounters practical obstacles, including storage limitations, communication overhead, and privacy concerns. To address these issues, one-shot methods are commonly employed. For these methods, estimators computed on local machines are transmitted to a central site and aggregated to form a global estimator, as in Zhang et al. (2013), Rosenblatt and Nadler (2016) and Fan et al. (2019). The most popular and direct aggregation method is simple averaging, which averages the estimators obtained on each local machine, as in Zhang et al. (2013). Besides of averaging, some other aggregating methods have been considered, such as meta estimators considering the variance of each local estimator developed in Liu et al. (2015) and Kullback-Leibler divergence method in Liu and Ihler (2014). Although one-shot approaches have the lowest communication costs, they suffer from several disadvantages, as discussed by Zhou et al. (2023). Alternatively there are some communication-iterative methods allowing a reasonable number of iterations, as in Shamir et al. (2014), Jordan et al. (2019) and Fan et al. (2023). Shamir et al. (2014) introduced a distributed approximate Newton-type method, which avoids transferring Hessian matrices across sites. Jordan et al. (2019) developed a communication-efficient surrogate likelihood framework for distributed data, where the key idea is to update the Hessian matrix on a single site only. For other important methods, kindly see the review by Gao et al. (2022) and Zhou et al. (2023).

In this paper, we investigate distributed inference methods for the high-dimensional convoluted rank regression. Motivated by the debiased techniques (see, for instance, Zhang and Zhang (2014); Van de Geer et al. (2014); Javanmard and Montanari (2014)), we develop debiased procedures for distributed convoluted rank regression. We make the following contributions.

- Firstly, we propose an averaging debiased estimator, which is one-shot approach and thus communication-efficient. The procedure involves obtaining debiased convoluted rank regression estimators on each machine and then averaging them.

- Secondly, we introduce a one-step debiased estimator, which is computation-efficient and requires only an estimator of the Hessian matrix on a single machine.

- Thirdly, we develop sparse averaging debiased estimator and sparse one-step debiased estimator, which significantly reduce the ℓ_1 and ℓ_2 errors of the averaging debiased estimator and the one-step debiased estimator.

- Lastly, we establish theoretical results for our introduced estimators. For the averaging debiased estimator and the one-step debiased estimator, we derive their asymptotic representation, Gaussian approximation, and ℓ_∞ error bound. Meanwhile, for the sparse debiased estimators, we obtain their ℓ_1 and ℓ_2 error bounds.

The remainder of the paper is organized as follows. In Section 2, we develop statistical inference procedures based on the averaging debiased estimator. In Section 3, we develop statistical

inference procedures based on the one-step debiased estimator. Section 4 reports numerical results from Monte Carlo studies. A real data example is presented in Section 5. Conclusions and discussions are provided in Section 6. All proofs and additional simulation studies are given in the Supplementary Material.

Notations: For a vector $\boldsymbol{x} = (x_1, \ldots, x_p)^\top$, let $\|\boldsymbol{x}\|_r = (|x_1|^r + \cdots + |x_p|^r)^{1/r}$ for $r \geq 1$ and $\|\boldsymbol{x}\|_\infty = \max_{1 \leq i \leq p} |x_i|$. For a matrix $\boldsymbol{A} = (a_{ij})_{n \times n}$, define $\|\boldsymbol{A}\|_r = \sup_{\|\boldsymbol{x}\|_r = 1} \|\boldsymbol{A}\boldsymbol{x}\|_r$, $\|\boldsymbol{A}\|_{\max} = \max_{1 \leq i,j \leq n} |a_{ij}|$ and $\|\boldsymbol{A}\|_\infty = \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|$. $\lambda_{\min}(\boldsymbol{B})$ and $\lambda_{\max}(\boldsymbol{B})$ are the smallest and largest eigenvalues of a positive semidefinite matrix $\boldsymbol{B}$. The sub-Gaussian norm of a sub-Gaussian random variable X is defined as $\|X\|_{\psi_2} = \inf\{t > 0 : \mathbb{E}[\exp(X^2/t^2)] \leq 2\}$. The sub-exponential norm of a sub-exponential random variable Y is defined as $\|Y\|_{\psi_1} = \inf\{t > 0 : \mathbb{E}(\exp(|Y|/t)) \leq 2\}$. A p-dimensional random vector $\boldsymbol{X}$ is sub-Gaussian if and only if $\sup_{\|\boldsymbol{u}\|_2 = 1} \|\boldsymbol{u}^\top \boldsymbol{X}\|_{\psi_2} < \infty$. Let $\|\boldsymbol{X}\|_{\psi_2} = \sup_{\|\boldsymbol{u}\|_2 = 1} \|\boldsymbol{u}^\top \boldsymbol{X}\|_{\psi_2}$ be the sub-Gaussian norm of $\boldsymbol{X}$. For two numerical sequences $\{x_n\}$ and $\{y_n\}$, $x_n = O(y_n)$ if there exists $c > 0$ such that $|x_n| \leq c|y_n|$, and $x_n = o(y_n)$ if $\lim_{n \to \infty} x_n/y_n = 0$. For a sequence of random variables $\{X_n\}$, $X_n = O_p(1)$ if $\lim_{t \to \infty} \limsup_{n \to \infty} \mathbb{P}(|X_n| \geq t) = 0$, and $X_n = o_p(1)$ if X_n convergences to 0 in probability.

2 Inference based on averaging debiased estimator

Consider the following linear regression model:

$$Y_i = \boldsymbol{X}_i^\top \boldsymbol{\beta}^* + \epsilon_i,$$

where $\{Y_i\}_{i=1}^n$ are independent and identically distributed (i.i.d.) responses, $\{\epsilon_i\}_{i=1}^n$ are random errors, and $\{\boldsymbol{X}_i\}_{i=1}^n$ are i.i.d. p-dimensional random vectors with $\mathbb{E}(\boldsymbol{X}_i) = \boldsymbol{0}$ and $\text{Var}(\boldsymbol{X}_i) = \boldsymbol{\Sigma} > 0$. The random vector $\boldsymbol{X}_i$ is independent of ϵ_i. The unknown parameter of interest is $\boldsymbol{\beta}^* \in \mathbb{R}^p$. Let $\mathcal{S} = \{j : \beta_j^* \neq 0\}$ and $s = |\mathcal{S}|$.

Hettmansperger and McKean (2010) considered the following rank regression estimator for $\boldsymbol{\beta}^*$:

$$\arg \min_{\boldsymbol{\beta} \in \mathbb{R}^p} \frac{1}{n(n-1)} \sum_{i \neq j}^n |(Y_i - \boldsymbol{X}_i^\top \boldsymbol{\beta}) - (Y_j - \boldsymbol{X}_j^\top \boldsymbol{\beta})|.$$

They noted that the rank regression estimator performs well when the error distribution is heavy-tailed. However, the non-smooth nature of the rank loss function poses a significant computational challenge. Zhou et al. (2024) proposed CRR, which reduces the computational burden of the rank regression estimator:

$$\arg \min_{\boldsymbol{\beta} \in \mathbb{R}^p} \frac{1}{n(n-1)} \sum_{i \neq j}^n L_h\big((Y_i - Y_j) - (\boldsymbol{X}_i - \boldsymbol{X}_j)^\top \boldsymbol{\beta}\big),$$

where $L_h(u) = \int_{-\infty}^{+\infty} |u - v| K_h(v) \, dv$ with $K_h(v) = h^{-1} K(v/h)$. Here, $K(v)$ is a kernel function, and h is a bandwidth. A smooth kernel function makes the convoluted rank loss function smooth. Define

$$\boldsymbol{\beta}_h^* = \arg \min_{\boldsymbol{\beta} \in \mathbb{R}^p} \mathbb{E}[L_h\big((Y_i - Y_j) - (\boldsymbol{X}_i - \boldsymbol{X}_j)^\top \boldsymbol{\beta}\big)].$$

Now consider a distributed setting with M local machines. Assume that the dataset, with sample size N, is equally distributed across the M machines. Let $n = N/M$ denote the sample size on each machine. Let $\{\boldsymbol{X}_i^{(k)}, Y_i^{(k)}\}_{i=1}^n$ be the observations stored on machine k, and let $\epsilon_i^{(k)}$ be the corresponding error term, where $1 \leq k \leq M$. On the k-th machine, the model is given by

$$Y_i^{(k)} = \boldsymbol{X}_i^{(k)\top} \boldsymbol{\beta}^* + \epsilon_i^{(k)}.$$

The corresponding loss function on the k-th machine is

$$\mathcal{L}_k(\boldsymbol{\beta}) = \frac{1}{n(n-1)} \sum_{i \neq j}^n L_h\big((Y_i^{(k)} - Y_j^{(k)}) - (\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)})^\top \boldsymbol{\beta}\big).$$

When the dimension p is much larger than the sample size n, we typically consider regularized estimation on each local machine. Specifically, on the k-th machine, the regularized estimator is given by

$$\widehat{\boldsymbol{\beta}}_{kh} = \arg\min_{\beta \in \mathbb{R}^p} \Big\{ \mathcal{L}_k(\boldsymbol{\beta}) + \sum_{l=1}^p p_\lambda(\|\beta_l\|) \Big\}. \tag{2.1}$$

Here, $p_\lambda(\cdot)$ is a penalty function and λ is the corresponding tuning parameter.

To obtain a final estimator of $\boldsymbol{\beta}^*$, we consider the average of the $\widehat{\boldsymbol{\beta}}_{kh}$'s. However, the regularized estimators $\widehat{\boldsymbol{\beta}}_{kh}$'s are generally biased (Zhang and Huang, 2008; Zhang and Zhang, 2014). Aggregating these estimators does not typically reduce the bias (Zhao et al., 2019). Motivated by the debiased procedures introduced by Zhang and Zhang (2014) and Van de Geer et al. (2014), we also consider the debiased estimator for each machine and conduct statistical inference based on the average of the debiased estimators.

Now we begin to introduce our debiased estimator. Let

$$\boldsymbol{J}_h = \mathbb{E}\big[L_h''(\epsilon_i^{(k)} - \epsilon_j^{(k)})(\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)})(\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)})^\top\big] = 2\mathbb{E}\big[L_h''(\epsilon_i^{(k)} - \epsilon_j^{(k)})\big]\boldsymbol{\Sigma}.$$

The sample version of $\boldsymbol{J}_h$ on the k-th machine is

$$\widehat{\boldsymbol{J}}_{kh} = \frac{1}{n(n-1)} \sum_{i \neq j}^n L_h''(\widehat{\epsilon}_i^{(k)} - \widehat{\epsilon}_j^{(k)})(\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)})(\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)})^\top.$$

In high dimensional settings, $\widehat{\boldsymbol{J}}_{kh}$ is not invertible. To estimate $\boldsymbol{J}_h^{-1}$, we adopt the method in Cai et al. (2011) and define $\widehat{\boldsymbol{W}}_{kh}$ as follows:

$$\begin{aligned} \widehat{\boldsymbol{W}}_{kh} = \arg\min_{\boldsymbol{W} \in \mathbb{R}^{p \times p}} \ & \|\boldsymbol{W}\|_\infty \\ \text{s.t.} \ & \|\boldsymbol{W}\widehat{\boldsymbol{J}}_{kh} - \boldsymbol{I}\|_{\max} \leq \gamma_n, \end{aligned} \tag{2.2}$$

where γ_n is a predetermined tuning parameter. From the proof of Cai et al. (2025), it is known that $\|\widehat{\boldsymbol{W}}\widehat{\boldsymbol{J}}_{kh} - \boldsymbol{I}\|_{\max} = O_p(s\log^2(p)/\sqrt{n})$. Thus a suitable choice of γ_n is $\gamma_n = O(s\log^2(p)/\sqrt{n})$.

Then the debiased CRR on the k-th machine is:

$$\widetilde{\boldsymbol{\beta}}_{kh} = \widehat{\boldsymbol{\beta}}_{kh} + \widehat{\boldsymbol{W}}_{kh}\frac{1}{n(n-1)}\sum_{i\neq j}^{n} L_h'(\widehat{\epsilon}_i^{(k)} - \widehat{\epsilon}_j^{(k)})(\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)}), \qquad (2.3)$$

where $\widehat{\epsilon}_j^{(k)} = Y_j^{(k)} - (\boldsymbol{X}_j^{(k)})^\top \widehat{\boldsymbol{\beta}}_{kh}$. The averaging debiased estimator (ADE) is the average of $\widehat{\boldsymbol{\beta}}_{kh}$ across all machines:

$$\widetilde{\boldsymbol{\beta}}_{\text{ade}} = \frac{1}{M}\sum_{k=1}^{M} \widetilde{\boldsymbol{\beta}}_{kh}. \qquad (2.4)$$

To establish theoretical results for the ADE, some technical assumptions are introduced.

- (A1) Assume that $\widehat{\boldsymbol{\beta}}_{kh}$ satisfies the following conditions for $1 \le k \le M$: $\|\widehat{\boldsymbol{\beta}}_{kh} - \boldsymbol{\beta}^*\|_1 = O_p(s\sqrt{\log p/n})$ and $\|\widehat{\boldsymbol{\beta}}_{kh} - \boldsymbol{\beta}^*\|_2 = O_p(\sqrt{s\log p/n})$. Further, assume that the kernel function $K(\cdot)$ is upper bounded, non-negative, symmetric about 0, satisfies $\int_{\mathbb{R}} K(t)\,dt = 1$ and $\int_{-\infty}^{\infty} |t|K(t)\,dt < \infty$. Additionally, assume that $K(\cdot)$ has a bounded derivative.

- (A2) Assume that $\boldsymbol{X}_i$ is a p-dimensional sub-Gaussian random vector with $\|\boldsymbol{X}_i\|_{\psi_2} \le M_0$, and the eigenvalues of the covariance matrix $\boldsymbol{\Sigma}$ satisfy $c_{\boldsymbol{\Sigma}} \le \lambda_{\min}(\boldsymbol{\Sigma}) \le \lambda_{\max}(\boldsymbol{\Sigma}) \le C_{\boldsymbol{\Sigma}}$ for some positive constants $c_{\boldsymbol{\Sigma}}$ and $C_{\boldsymbol{\Sigma}}$.

- (A3) Assume that $\boldsymbol{J}_h^{-1} = (\boldsymbol{\omega}_1, \ldots, \boldsymbol{\omega}_p)^\top$ satisfies $\max_{1 \le i \le p} \sum_{j=1}^{p} |\omega_{ij}|^q < C_\omega$ for some $q \in (0,1)$ and positive constant C_ω.

- (A4) Let $f(\cdot)$ be the probability density function of ϵ_i. Assume that $f(\cdot)$ is continuous and there exists a positive constant δ_1 such that $\inf_{v \in [-\delta_1, \delta_1]} f(v) > 0$. Let $g(\cdot)$ be the probability density function of $\epsilon_i - \epsilon_j$. Assume that $|g(x) - g(y)| \le L_0|x-y|$, $\forall x, y \in \mathbb{R}$, for some constant $L_0 > 0$. Further assume that $0 < \underline{g} \le g(0) \le \overline{g} < \infty$.

Assumptions (A1)-(A2) are standard conditions on the penalized estimator, kernel function, and covariance matrix, see for instance Zhou et al. (2024). The error bound of $\widehat{\boldsymbol{\beta}}_{kh}$ has been derived by Cai et al. (2025) under some mild assumptions on the density function of ϵ_i, the penalty function $p_\lambda(\cdot)$, and the distribution of $\boldsymbol{X}_i$. Assumption (A3) asks $\boldsymbol{J}_h^{-1}$ to be sparse in terms of ℓ_p norm of matrix row space. Similar conditions were widely imposed by many authors; see Bickel and Levina (2008), Cai et al. (2011), and Cai et al. (2025). Since the error ϵ and $\boldsymbol{X}$ are independent, Assumption (A3) can be reduced to sparsity condition on $\boldsymbol{\Sigma}^{-1}$. Assumption (A4) is a mild condition for the density of the error term. Notably no moment condition is required for the error term.

The next two theorems depict the asymptotic behavior of the ADE. We first present the asymptotic representation for the ADE, which shows that $\sqrt{N}(\widetilde{\boldsymbol{\beta}}_{\text{ade}} - \boldsymbol{\beta}^*)$ can be expressed as a high-dimensional U-statistic up to an $o_p(1)$ term.

Theorem 2.1. *Let $a = (5 - 4q)/(2 - 2q)$. Under Assumption (A1)-(A4), the ADE satisfies*

$$\|\sqrt{N}(\widetilde{\boldsymbol{\beta}}_{ade} - \boldsymbol{\beta}^*) - \sqrt{N}\boldsymbol{J}_h^{-1}\frac{1}{Mn(n-1)}\sum_{k=1}^{M}\sum_{i\neq j}^{n} L_h'(\epsilon_i^{(k)} - \epsilon_j^{(k)})(\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)})\|_\infty$$

$$= O_p\big(M^{1/2}\big[s\log^a(p)/\sqrt{n}\big]^{1-q} + M^{1/2}s^2\log^{5/2}(p)/\sqrt{n}\big).$$

We can then further obtain the Gaussian approximation result for the ADE presented in the following theorem.

Theorem 2.2. *Under Assumption (A1)-(A4), suppose that $M\log^7(pn)n^{r-1} = O(1)$ for some $r \in (0,1)$, $M^{1/2}[s\log^b(p)/\sqrt{n}]^{1-q} + M^{1/2}s^2\log^3(p)/\sqrt{n} = o(1)$ with $b = (3-2q)/(1-q)$. Then there is a p-dimensional Gaussian random vector $\boldsymbol{Z}_n$ with mean zero and covariance matrix*

$$\boldsymbol{\Sigma}_{\boldsymbol{z}} = \frac{\mathbb{E}\big[\mathbb{E}^2\big(L_h'(\epsilon_i - \epsilon_j) \mid \epsilon_i\big)\big]}{\big[\mathbb{E}L_h''(\epsilon_i - \epsilon_j)\big]^2}\boldsymbol{\Sigma}^{-1} = \frac{2\mathbb{E}\big[\mathbb{E}^2\big(L_h'(\epsilon_i - \epsilon_j) \mid \epsilon_i\big)\big]}{\mathbb{E}L_h''(\epsilon_i - \epsilon_j)}\boldsymbol{J}_h^{-1},$$

such that the ADE satisfies

$$\sup_{t \in \mathbb{R}}\big|\mathbb{P}(\sqrt{N}\|\widetilde{\boldsymbol{\beta}}_{ade} - \boldsymbol{\beta}^*\|_\infty \le t) - \mathbb{P}(\|\boldsymbol{Z}_n\|_\infty \le t)\big| = o(1).$$

Theorem 2.2 establishes the Gaussian approximation for $\sqrt{N}\|\widetilde{\boldsymbol{\beta}}_{\mathrm{ade}} - \boldsymbol{\beta}^*\|_\infty$. The theorem implies that $\sqrt{N}(\widetilde{\boldsymbol{\beta}}_{\mathrm{ade}} - \boldsymbol{\beta}^*)$ asymptotically follows a high-dimensional Gaussian distribution, represented by the random vector $\boldsymbol{Z}_n$.

From the above Theorem, we can construct confidence interval for each β_j^*. Actually on each machine, we can estimate covariance matrix $\boldsymbol{\Sigma}_{\boldsymbol{z}}$ by

$$\widehat{\boldsymbol{\Sigma}}_{\boldsymbol{z}}^{(k)} = \frac{2(n-1)\sum_{i=1}^n\big\{\sum_{j=1}^n L_h'(\widehat{\epsilon}_i^{(k)} - \widehat{\epsilon}_j^{(k)})\big\}^2}{n^2\sum_{i=1}^n\sum_{j\neq i}^n L_h''(\widehat{\epsilon}_i^{(k)} - \widehat{\epsilon}_j^{(k)})}\,\widehat{\boldsymbol{W}}_{kh},$$

Let $\boldsymbol{\Sigma}_{\boldsymbol{z}} = (\sigma_{ij})_{i,j}^p$. We estimate σ_{jj} by averaging the results from M machines, denoted as $\widehat{\sigma}_{jj}$. The confidence interval for each β_j^* can be constructed as follows:

$$\bigg\{\widetilde{\boldsymbol{\beta}}_{\mathrm{ade},j} - z_{\alpha/2}\sqrt{\frac{\widehat{\sigma}_{jj}}{N}}, \quad \widetilde{\boldsymbol{\beta}}_{\mathrm{ade},j} + z_{\alpha/2}\sqrt{\frac{\widehat{\sigma}_{jj}}{N}}\bigg\}. \tag{2.5}$$

Here $z_{\alpha/2}$ is the $\alpha/2$-quantile of the standard normal distribution.

Besides the above results, we can also obtain the ℓ_∞ error bound of the ADE. To be precise, we have the following theorem.

Theorem 2.3. *Under the assumptions in Theorem 2.2, we have that*

$$\|\widetilde{\boldsymbol{\beta}}_{ade} - \boldsymbol{\beta}^*\|_\infty = O_p(\sqrt{\log(p)/N}).$$

3 Inference based on one-step debiased estimator

To calculate the averaging debiased estimator, we have to calculate each $\widehat{\boldsymbol{W}}_{kh}$ for $1 \le k \le M$, which is computationally expensive. To reduce the computational cost, we introduce a one-step debiased estimator (ODE) in this section. The ODE is more computation-efficient, as it only

requires the calculation of $\widehat{\boldsymbol{W}}_{1h}$. Specifically, the ODE is defined as follows:

$$\widetilde{\boldsymbol{\beta}}_{\text{ode}} = \widehat{\boldsymbol{\beta}}_{1h} + \widehat{\boldsymbol{W}}_{1h} \frac{1}{Mn(n-1)} \sum_{k=1}^{M} \sum_{i \neq j}^{n} L_h'(\breve{\epsilon}_i^{(k)} - \breve{\epsilon}_j^{(k)})(\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)}), \tag{3.1}$$

where $\breve{\epsilon}_j^{(k)} = Y_j^{(k)} - (\boldsymbol{X}_j^{(k)})^\top \widehat{\boldsymbol{\beta}}_{1h}$.

We make some discussions about the construction of the ODE. Note that the debiased estimator on the first machine is given as follows

$$\widetilde{\boldsymbol{\beta}}_1 = \widehat{\boldsymbol{\beta}}_{1h} + \widehat{\boldsymbol{W}}_{1h} \frac{1}{n(n-1)} \sum_{i \neq j}^{n} L_h'(\widehat{\epsilon}_i^{(1)} - \widehat{\epsilon}_j^{(1)})(\boldsymbol{X}_i^{(1)} - \boldsymbol{X}_j^{(1)}),$$

where $\widehat{\epsilon}_j^{(1)} = Y_j^{(1)} - (\boldsymbol{X}_j^{(1)})^\top \widehat{\boldsymbol{\beta}}_{1h}$. However, this debiased estimator cannot achieve root-N convergence rate since there are only n units on the first machine. Instead of only using the gradient information on the first machine, the ODE aggregates all the gradient information from all the machines which makes it achieve the optimal convergence rate. On the other hand, different from the ADE, the ODE requires only one estimator of $\boldsymbol{J}_h^{-1}$, while the ADE requires $M \, \widehat{\boldsymbol{W}}_{kh}$'s. When M is large, the computation can be heavy.

The next two theorems depict the asymptotic behavior for the ODE. Similar to the ADE, we firstly give the asymptotic representation for the ODE, suggesting that $\sqrt{N}(\widetilde{\boldsymbol{\beta}}_{\text{ode}} - \boldsymbol{\beta}^*)$ can be expressed as a high-dimensional U-statistic up to an $o_p(1)$ term.

Theorem 3.1. *Let $a = (5-4q)/(2-2q)$. Under Assumption (A1)-(A3), the ODE satisfies*

$$\|\sqrt{N}(\widetilde{\boldsymbol{\beta}}_{ode} - \boldsymbol{\beta}^*) - \sqrt{N}\boldsymbol{J}_h^{-1} \frac{1}{Mn(n-1)} \sum_{k=1}^{M} \sum_{i \neq j}^{n} L_h'(\epsilon_i^{(k)} - \epsilon_j^{(k)})(\boldsymbol{X}_i^{(k)} - \boldsymbol{X}_j^{(k)})\|_\infty$$
$$= O_p(M^{1/2}[s\log^a(p)/\sqrt{n}]^{1-q} + M^{1/2}s^2\log^{5/2}(p)/\sqrt{n}).$$

Notice that the asymptotic representation for the ODE is the same as that for the ADE, which means the error bound of the ODE is asymptotically the same as the ADE. But the ODE is more computation-efficient, while the ADE is more communication-efficient. We can further get the Gaussian approximation result for the ODE by the following theorem.

Theorem 3.2. *Under Assumption (A1)-(A4), suppose $M\log^7(pn)n^{r-1} = O(1)$ for some $r \in (0,1)$, $M^{1/2}[s\log^b(p)/\sqrt{n}]^{1-q} + M^{1/2}s^2\log^3(p)/\sqrt{n} = o(1)$ with $b = (3-2q)/(1-q)$. Then there is a p-dimensional Gaussian random vector $\boldsymbol{Z}_n$ with mean zero and covariance matrix*

$$\boldsymbol{\Sigma}_z = \frac{\mathbb{E}\left[\mathbb{E}^2\left(L_h'(\epsilon_i - \epsilon_j) \mid \epsilon_i\right)\right]}{\left[\mathbb{E}L_h''(\epsilon_i - \epsilon_j)\right]^2} \boldsymbol{\Sigma}^{-1} = \frac{2\mathbb{E}\left[\mathbb{E}^2\left(L_h'(\epsilon_i - \epsilon_j) \mid \epsilon_i\right)\right]}{\mathbb{E}L_h''(\epsilon_i - \epsilon_j)} \boldsymbol{J}_h^{-1},$$

such that the ODE satisfies

$$\sup_{t \in \mathbb{R}} |\mathbb{P}(\sqrt{N}\|\widetilde{\boldsymbol{\beta}}_{ode} - \boldsymbol{\beta}^*\|_\infty \leq t) - \mathbb{P}(\|\boldsymbol{Z}_n\|_\infty \leq t)| = o(1).$$

Derived from the above theorem, we construct the confidence interval given by the ODE for each β_j^* as follows. Now we estimate covariance matrix $\boldsymbol{\Sigma}_z$ only based on the first machine, that

is

$$\widehat{\boldsymbol{\Sigma}}_z = \frac{2(n-1)\sum_{i=1}^n \big\{ \sum_{j=1}^n L_h'(\check{\epsilon}_i^{(1)} - \check{\epsilon}_j^{(1)}) \big\}^2}{n^2 \sum_{i=1}^n \sum_{j\neq i}^n L_h''(\check{\epsilon}_i^{(1)} - \check{\epsilon}_j^{(1)})} \, \widehat{\boldsymbol{W}}_{1h},$$

The confidence interval for each β_j^* can be constructed as follows:

$$\Big\{ \widetilde{\boldsymbol{\beta}}_{\mathrm{ode},j} - z_{\alpha/2}\sqrt{\frac{\widehat{\boldsymbol{\Sigma}}_{z,jj}}{N}}, \quad \widetilde{\boldsymbol{\beta}}_{\mathrm{ode},j} + z_{\alpha/2}\sqrt{\frac{\widehat{\boldsymbol{\Sigma}}_{z,jj}}{N}} \Big\}. \tag{3.2}$$

Similarly, the ODE also satisfies the same error bound as the ADE.

Theorem 3.3. *Under the assumptions in Theorem 3.2, we have that*

$$\|\widetilde{\boldsymbol{\beta}}_{ode} - \boldsymbol{\beta}^*\|_\infty = O_p\Big(\sqrt{\frac{\log p}{N}}\Big).$$

Finally, note that the debiased estimators (the ADE and the ODE) do not exhibit sparsity. To address this, a hard thresholding approach can be applied to obtain the sparse averaging debiased estimator (SADE) and the sparse one-step debiased estimator (SODE).

Let c be a threshold level. We respectively define SADE and SODE as

$$\widetilde{\beta}_{\mathrm{ade},j}^c = \widetilde{\beta}_{\mathrm{ade},j}\,\mathbb{1}\{|\widetilde{\beta}_{\mathrm{ade},j}| > c\}, \quad \widetilde{\beta}_{\mathrm{ode},j}^c = \widetilde{\beta}_{\mathrm{ode},j}\,\mathbb{1}\{|\widetilde{\beta}_{\mathrm{ode},j}| > c\}.$$

Here $\mathbb{1}\{\cdot\}$ is the indicator function. The advantage of the sparse aggregated estimators is that they reduce both the ℓ_1 and ℓ_2 errors of the debiased estimators. We obtain the following result which establishes the estimation error bound of SADE and SODE.

Theorem 3.4. *Let $c = c_0\sqrt{\log p/N}$ with a sufficiently large c_0, and let $\widetilde{\boldsymbol{\beta}}^c$ be either SADE or SODE. Under the assumptions of Theorem 3.2, we have*

$$\|\widetilde{\boldsymbol{\beta}}^c - \boldsymbol{\beta}^*\|_\infty = O_p\Big(\sqrt{\frac{\log p}{N}}\Big), \quad \|\widetilde{\boldsymbol{\beta}}^c - \boldsymbol{\beta}^*\|_2 = O_p\Big(\sqrt{\frac{s\log p}{N}}\Big), \quad \|\widetilde{\boldsymbol{\beta}}^c - \boldsymbol{\beta}^*\|_1 = O_p\Big(s\sqrt{\frac{\log p}{N}}\Big).$$

When we choose the threshold as $c = c_0\sqrt{\log p/N}$, the sparse debiased estimators achieve the same convergence rate as the global regularized estimator obtained by using all observations in both ℓ_1 and ℓ_2 norms.

4 Numerical simulation

In this section, we conduct numerical simulations to evaluate the finite sample performance of our proposed procedures. We adopt the Epanechnikov kernel as the kernel function, $K(u) = 3(1-u^2)\mathbb{1}\{-1 \leq u \leq 1\}/4$, and the loss function is defined by

$$L_h(u) = \begin{cases} u, & u \geq h, \\ 3u^2 h^{-1}/4 - u^4 h^{-3}/8 + 3h/8, & -h < u < h, \\ -u, & u \leq -h. \end{cases}$$

Here we fix $h = 1$.

We generate data from a linear model as follows:

$$Y_i = \boldsymbol{X}_i^\top \boldsymbol{\beta}^* + \epsilon_i,$$

where $\boldsymbol{\beta}^* = (\sqrt{3}, \sqrt{3}, \sqrt{3}, 0, \ldots, 0)^\top$. To evaluate the performance under different true coefficients, we further consider different settings of $\boldsymbol{\beta}$, and the corresponding results are illustrated in Tables S.9-S.12 in the Supplementary Material. The predictor $\boldsymbol{X}_i$ is from the multivariate normal distribution $\mathbf{N}_p(\mathbf{0}_p, \boldsymbol{\Sigma})$. Furthermore, the random error term ϵ_i is independent with $\boldsymbol{X}_i$. Two scenarios are considered for the distribution of ϵ_i:

- **Scenario 1:** ϵ_i is simulated from standard normal distribution, i.e. $\epsilon_i \sim N(0, 1)$.

- **Scenario 2:** ϵ_i follows the mixture gaussian distribution $\tau N(0, 1) + (1 - \tau)N(0, 9)$ with $\tau \sim B(1, 0.95)$.

Additional results for other error distributions, including the t-distribution, the Cauchy distribution, and several asymmetric distributions, are reported in Tables S.1-S.8 in the Supplementary Material. There are two settings for the correlation structure of $\boldsymbol{X}$:

- **Setting 1:** $\boldsymbol{\Sigma}$ is Toeplitz with $\Sigma_{ij} = 0.3^{|i-j|}$.

- **Setting 2:** $\boldsymbol{\Sigma}$ is block diagonal, that is, $\Sigma_{ii} = 1$ for $i = 1, \ldots, p$ and $\Sigma_{ij} = 0.3\mathbb{1}_{\{|i-j|=1\}}$ for $i \neq j$.

Besides, we estimate the unknown parameter $\boldsymbol{\beta}^*$ using the local linear approximation algorithm as described in Algorithm 1 of Zhou et al. (2024).

We compute the centralized debiased estimator (CDE) obtained by a single machine using all observations, ADE, ODE, SADE, and SODE. Our first goal is to assess the accuracy of CDE, ADE, SADE, ODE, and SODE using the ℓ_∞ norm and the ℓ_2 norm under different scenarios and settings. Secondly, we aim to evaluate the confidence intervals based on ADE and ODE under various circumstances. Specifically, we are interested in conducting inference on the coefficients corresponding to certain index set $\mathcal{G}$. Different choices of $\mathcal{G}$ are taken into consideration: $\mathcal{G}_1 = \{1, 2, \ldots, \lfloor p/2 \rfloor\}$, $\mathcal{G}_2 = \{\lfloor p/2 \rfloor + 1, \lfloor p/2 \rfloor + 2, \ldots, p\}$, and $\mathcal{G}_3 = \{1, 2, \ldots, p\}$. We evaluate the average coverage probability (ACP) and the average length (AL) of the 95% confidence intervals. Specifically, the ACP is defined as $\text{ACP} := \sum_{j \in \mathcal{G}} |\mathcal{G}|^{-1}\text{CP}_j$, where CP_j denotes the coverage probability of the 95% confidence interval for β_j^*. Similarly, the AL is given by $\text{AL} := \sum_{j \in \mathcal{G}} |\mathcal{G}|^{-1}\text{AL}_j$, where AL_j represents the length of the 95% confidence interval for β_j^*. All results are calculated based on 500 simulation runs.

For the threshold c in sparse aggregated estimators, we consider two approaches. The first approach is to directly set $c = c_0\sqrt{\log(p)/N}$ with $c_0 = 1.5$. For similar adoption, see also Hou et al. (2023). The other is a rule suggested by Zhao et al. (2019). That is, let $s_{\max} = \max_{1 \leq k \leq M} \|\widehat{\boldsymbol{\beta}}_{kh}\|_0$, order $|\widetilde{\beta}|_{(1)} \geq \cdots \geq |\widetilde{\beta}|_{(p)}$, and set $c = |\widetilde{\beta}|_{(s_{\max}+1)}$. Here $|\widetilde{\beta}|_{(l)}$ represents the l-th largest element of the absolute value version of the ADE or the ODE. The simulation results based on this rule are reported in Figures S.1-S.3 in the Supplementary Material.

First, we consider sample sizes $N =$ 1000, 2000, 3000, 4000, with $p = 500$ and fix the number of samples in each local machine to be $n =$ 200. It is seen from Figure 1 that the errors of all the estimators decrease when the total sample size increases. In terms of ℓ_∞ error, the differences among all estimators are small. In terms of ℓ_2 error, SADE and SODE perform better than CDE, ADE, and ODE which are not sparse. In other words, thresholding is effective in reducing the ℓ_2 error.

In the second set of simulations, we vary $M = 1, 5, 10, 20$ while fixing $N = 2000$ and $p = 500$. Performance generally deteriorates with increasing M. Figure 2 shows that, the thresholding is quite effective in reducing the ℓ_2 error.

In the third set of simulations, we set $N = 2000$, $M = 1, 10$ ($M = 1$ corresponds to the centralized estimator) and $p = 100, 300, 500, 700$. Figure 3 shows the errors of the distributed estimators with $M = 10$. As expected, when the dimension p increases, the errors become larger. Sparse aggregated estimators reduce the ℓ_2 error, while the ℓ_∞ error changes little.

In the final set of simulations, we vary $M = 5, 10, 15, 20$ and $p = 100, 200$ for constructing confidence intervals by fixing the number of samples in each local machine to be $n = 200$. Corresponding numerical results are presented in Tables 1-2. The results in Tables 1-2 provide strong evidence that corroborates our established theoretical results. Actually for all scenarios, settings, and index sets, the coverage probabilities for ADE and ODE are all close to the nominal level 95%. The average lengths of these confidence intervals are also very short.

The simulations are carried out on Windows with R version 4.3.2, using an 8core CPU and 16 GB of memory. For the third set of simulations with $p = 100$, for example, the CDE requires about 31.9 hours to finish all 500 repetitions, while the ADE with $M = 10$ requires about 7.26 hours and the ODE requires about 1.18 hours. This verifies that the ODE is relatively computation-efficient.

Figure 1: The plot of ℓ_∞ and ℓ_2 errors for $N \in \{1000, 2000, 3000, 4000\}$ under different scenarios and settings. The dashdot line with cross points, dashed line with left triangle points, dashed line with up triangle points, dotted line with down triangle points, and dotted line with diamond points respectively represent the results corresponding to CDE, ADE, SADE, ODE, and SODE.

Figure 2: The plot of ℓ_∞ and ℓ_2 errors for $M \in \{1, 5, 10, 20\}$ under different scenarios and settings ($M = 1$ corresponds to the CDE). The dashed line with left triangle points, dashed line with up triangle points, dotted line with down triangle points, and dotted line with diamond points respectively represent the results corresponding to ADE, SADE, ODE, and SODE.

Figure 3: The plot of ℓ_∞ and ℓ_2 errors for $p \in \{100, 300, 500, 700\}$ under different scenarios and settings. The dashdot line with cross points, dashed line with left triangle points, dashed line with up triangle points, dotted line with down triangle points, and dotted line with diamond points respectively represent the results corresponding to CDE, ADE, SADE, ODE, and SODE.

Table 1: Empirical average coverage probability (ACP) and average length (AL) of confidence intervals for ADE and ODE at the 95% level under **Scenario 1** and **Scenario 2** with $p = 100$.

		M	Scenario 1 $\mathcal{G}_1$ ACP	AL	$\mathcal{G}_2$ ACP	AL	$\mathcal{G}_3$ ACP	AL	Scenario 2 $\mathcal{G}_1$ ACP	AL	$\mathcal{G}_2$ ACP	AL	$\mathcal{G}_3$ ACP	AL
Setting 1	ADE	5	0.946	0.142	0.946	0.142	0.947	0.142	0.947	0.144	0.947	0.144	0.947	0.144
		10	0.944	0.101	0.944	0.101	0.945	0.101	0.949	0.102	0.949	0.102	0.947	0.102
		15	0.945	0.082	0.945	0.082	0.945	0.082	0.945	0.083	0.945	0.083	0.945	0.083
		20	0.944	0.071	0.944	0.071	0.945	0.071	0.949	0.072	0.949	0.072	0.947	0.072
	ODE	5	0.944	0.133	0.944	0.133	0.944	0.133	0.940	0.133	0.948	0.133	0.944	0.133
		10	0.938	0.094	0.945	0.094	0.942	0.094	0.941	0.094	0.943	0.094	0.942	0.094
		15	0.936	0.077	0.943	0.077	0.939	0.077	0.935	0.077	0.942	0.077	0.938	0.077
		20	0.930	0.066	0.943	0.066	0.937	0.066	0.933	0.067	0.944	0.067	0.939	0.067
Setting 2	ADE	5	0.943	0.151	0.943	0.151	0.942	0.151	0.941	0.152	0.941	0.152	0.942	0.152
		10	0.941	0.107	0.941	0.107	0.943	0.107	0.944	0.108	0.944	0.108	0.943	0.108
		15	0.945	0.087	0.945	0.087	0.947	0.087	0.946	0.088	0.946	0.088	0.946	0.088
		20	0.944	0.075	0.944	0.075	0.944	0.075	0.943	0.076	0.943	0.076	0.943	0.076
	ODE	5	0.944	0.141	0.944	0.141	0.944	0.141	0.937	0.141	0.944	0.141	0.941	0.141
		10	0.938	0.099	0.945	0.099	0.942	0.099	0.935	0.100	0.943	0.100	0.941	0.100
		15	0.935	0.081	0.947	0.081	0.941	0.081	0.934	0.081	0.944	0.081	0.941	0.081
		20	0.930	0.070	0.945	0.070	0.938	0.070	0.927	0.070	0.942	0.071	0.934	0.071

Table 2: Empirical average coverage probability (ACP) and average length (AL) of CIs for ADE and ODE at the 95% level under **Scenario 1** and **Scenario 2** with $p = 200$.

		M	Scenario 1 $\mathcal{G}_1$ ACP	AL	$\mathcal{G}_2$ ACP	AL	$\mathcal{G}_3$ ACP	AL	Scenario 2 $\mathcal{G}_1$ ACP	AL	$\mathcal{G}_2$ ACP	AL	$\mathcal{G}_3$ ACP	AL
Setting 1	ADE	5	0.944	0.144	0.944	0.144	0.947	0.142	0.944	0.146	0.944	0.146	0.943	0.146
		10	0.941	0.102	0.941	0.102	0.945	0.101	0.943	0.103	0.943	0.103	0.943	0.103
		15	0.942	0.083	0.942	0.083	0.945	0.082	0.943	0.084	0.943	0.084	0.943	0.084
		20	0.941	0.072	0.941	0.072	0.945	0.071	0.942	0.073	0.942	0.073	0.942	0.073
	ODE	5	0.944	0.131	0.946	0.130	0.945	0.131	0.946	0.131	0.944	0.131	0.945	0.131
		10	0.939	0.092	0.941	0.092	0.940	0.092	0.940	0.093	0.941	0.092	0.941	0.093
		15	0.939	0.075	0.942	0.075	0.939	0.075	0.940	0.076	0.941	0.076	0.941	0.076
		20	0.936	0.065	0.942	0.065	0.939	0.065	0.937	0.065	0.941	0.065	0.939	0.065
Setting 2	ADE	5	0.946	0.153	0.946	0.153	0.945	0.153	0.945	0.155	0.945	0.155	0.944	0.155
		10	0.947	0.109	0.947	0.109	0.947	0.108	0.944	0.109	0.944	0.109	0.944	0.109
		15	0.946	0.089	0.946	0.089	0.944	0.089	0.944	0.089	0.944	0.089	0.944	0.089
		20	0.946	0.077	0.946	0.077	0.944	0.077	0.945	0.077	0.945	0.077	0.944	0.077
	ODE	5	0.945	0.138	0.941	0.138	0.943	0.138	0.943	0.139	0.945	0.139	0.944	0.139
		10	0.943	0.098	0.945	0.098	0.944	0.098	0.941	0.098	0.945	0.098	0.944	0.098
		15	0.938	0.080	0.943	0.080	0.941	0.080	0.938	0.080	0.943	0.080	0.941	0.080
		20	0.938	0.069	0.942	0.069	0.942	0.069	0.937	0.069	0.942	0.069	0.941	0.069

5 Real data analysis

We illustrate our proposed method through studying the Genotype-Tissue Expression(GTEx) data, referred to as GTEx data, which has been analyzed by Li et al. (2022). We focus on the central nervous systems MODULE_137. This dataset is obtained from `https://www.gsea-msigdb.org/gsea/msigdb/cards/MODULE_137.html`. After removing missing values, it includes information on 536 genes of 17,382 observations. The dataset comprises genes measured across 13 brain tissues, such as the Amygdala, Anterior Cingulate Cortex, and Frontal Cortex. Specifically, we are interested in predicting the expression level of gene Dishevelled Segment Polarity Protein 1 (DVL1). DVL1 is a conserved hub of the Wnt pathway. Its activity is regulated by post-translational modifications, including TTLL11-mediated polyglutamylation (Kravec et al., 2024). Perturbation of DVL1 changes β-catenin signaling and cancer-related traits. Activation of DVL1 promotes stemness and metastasis, whereas knockdown reduces these features, making its expression a clinically relevant and interpretable endpoint (Yin et al., 2021).

We set gene DVL1 as the response variable and the remaining 535 genes as predictors. The large sample size 17,382 makes the computation not sufficient for a central analysis. To this end, the 17,382 observations were randomly divided across $M = 10$ machines. We apply the feature screening method of Li et al. (2012) and retain the top $n/\log(n)$ genes. These selected genes are then used to predict the expression of DVL1.

Our primary objective is to construct confidence intervals for the coefficients of the predictors. Table 3 reports the 95% confidence intervals for the coefficients of five most significant predictors. For both ADE and ODE, every confidence interval excludes zero. This suggests a positive association with DVL1 expression. We observe that the gene WDR7 is the most important gene for the expression of DVL1. This gene is also observed in Crummy et al. (2019). Other genes such as TRIM23, EXD2, ACTL6B, and RALGPS1 have also been highlighted in biomedical research, for example in Broderick et al. (2016). These consistent observations increase our confidence in the predictors identified by our method and provide useful directions for future study.

Table 3: Confidence intervals at the confidence level 95% in GTEx data.

	ADE	ODE
WDR7	(37.884, 39.268)	(36.354, 37.691)
TRIM23	(13.962, 14.555)	(16.566, 17.179)
EXD2	(11.576, 12.084)	(11.841, 12.364)
ACTL6B	(8.403, 8.698)	(11.890, 12.207)
RALGPS1	(8.089, 8.589)	(8.396, 8.908)

WDR7 denotes WD Repeat Domain 7. TRIM23 denotes Tripartite Motif Containing 23. EXD2 denotes Exonuclease 3'5' Domain-Containing 2. ACTL6B denotes Actin-Like 6B. RALGPS1 denotes Ral GEF with PH Domain and SH3-binding Motif 1.

6 Conclusions and discussions

In this paper, we introduce two debiased estimators for high-dimensional convoluted rank regression in distributed setting. The averaging debiased estimator is a one-shot method and thus is communication-efficient, while the one-step debiased estimator computes the Hessian matrix only on a single machine and thus is computation-efficient. For these two debiased estimators, we derive their asymptotical representation, Gaussian approximation, and ℓ_∞ error bound. Finally, a thresholding approach is involved to construct sparse averaging debiased estimator and sparse one-step debiased estimator, which significantly reduce the ℓ_1 and ℓ_2 errors.

There are several possible extensions for future research. Firstly, we may consider investigating distributed inference for high-dimensional semi-parametric convoluted rank regression. Secondly, the development of Byzantine-robust distributed procedure is an important issue. Further in this paper, we assume that there is a central machine which can communicate the estimators or the gradients. However, in practice, decentralized systems are also very common. We will explore these issues in near future.

References

Alexandre Belloni and Victor Chernozhukov. ℓ_1-penalized quantile regression in high-dimensional sparse models. *Ann. Stat.*, 39(1):82–130, 2011.

Peter J Bickel and Elizaveta Levina. Covariance regularization by thresholding. *Ann. Stat.*, 36(6):2577–2604, 2008.

Ronan Broderick, Jadwiga Nieminuszczy, Hannah T Baddock, Rajashree A Deshpande, Opher Gileadi, Tanya T Paull, Peter J McHugh, and Wojciech Niedzwiedz. Exd2 promotes homologous recombination by facilitating dna end resection. *Nature cell biology*, 18(3):271–280, 2016.

Leheng Cai, Xu Guo, Heng Lian, and Liping Zhu. Statistical inference for high-dimensional convoluted rank regression. *J. Am. Stat. Assoc.*, accepted, 2025.

Tony Cai, Weidong Liu, and Xi Luo. A constrained ℓ_1 minimization approach to sparse precision matrix estimation. *J. Am. Stat. Assoc.*, 106(494):594–607, 2011.

Ellen Crummy, Muralidharan Mani, John C Thellman, and Thomas FJ Martin. The priming factor caps1 regulates dense-core vesicle acidification by interacting with rabconnectin3β/wdr7 in neuroendocrine cells. *J. Biol. Chem.*, 294(24):9402–9415, 2019.

Jianqing Fan, Quefeng Li, and Yuyan Wang. Estimation of high dimensional mean regression in the absence of symmetry and light tail assumptions. *J. R. Stat. Soc. Ser. B (Stat. Methodol.)*, 79(1):247–265, 2017.

Jianqing Fan, Dong Wang, Kaizheng Wang, and Ziwei Zhu. Distributed estimation of principal eigenspaces. *Ann. Stat.*, 47(6):3009, 2019.

Jianqing Fan, Yongyi Guo, and Kaizheng Wang. Communication-efficient accurate statistical estimation. *J. Am. Stat. Assoc.*, 118(542):1000–1010, 2023.

Marcelo Fernandes, Emmanuel Guerre, and Eduardo Horta. Smoothing quantile regressions. *J. Bus. Econ. Stat.*, 39(1):338–357, 2021.

Yuan Gao, Weidong Liu, Hansheng Wang, Xiaozhou Wang, Yibo Yan, and Riquan Zhang. A review of distributed statistical inference. *Stat. Theory Relat. Fields*, 6(2):89–99, 2022.

Thomas P Hettmansperger and Joseph W McKean. *Robust nonparametric statistical methods.* CRC press, 2010.

Zhaohan Hou, Wei Ma, and Lei Wang. Sparse and debiased lasso estimation and inference for high-dimensional composite quantile regression with distributed data. *TEST*, 32(4):1230–1250, 2023.

Adel Javanmard and Andrea Montanari. Confidence intervals and hypothesis testing for high-dimensional regression. *J. Mach. Learn. Res.*, 15(1):2869–2909, 2014.

Michael I Jordan, Jason D Lee, and Yun Yang. Communication-efficient distributed statistical inference. *J. Am. Stat. Assoc.*, 114:668–681, 2019.

Marek Kravec, Ondrej Šedo, Jana Nedvědová, Miroslav Micka, Marie Šulcová, Nikodém Zezula, Kristína Gömöryová, David Potěšil, Ranjani Sri Ganji, Sara Bologna, et al. Carboxy-terminal polyglutamylation regulates signaling and phase separation of the dishevelled protein. *EMBO J.*, 43(22):5635–5666, 2024.

Runze Li, Wei Zhong, and Liping Zhu. Feature screening via distance correlation learning. *J. Am. Stat. Assoc.*, 107(499):1129–1139, 2012.

Sai Li, T Tony Cai, and Hongzhe Li. Transfer learning for high-dimensional linear regression: Prediction, estimation and minimax optimality. *J. R. Stat. Soc. Ser. B (Stat. Methodol.)*, 84(1):149–173, 2022.

Dungang Liu, Regina Y Liu, and Minge Xie. Multivariate meta-analysis of heterogeneous studies using only summary statistics: efficiency and robustness. *J. Am. Stat. Assoc.*, 110(509):326–340, 2015.

Qiang Liu and Alexander Ihler. Distributed estimation, information loss and exponential families. In *Proceedings of the 27th International Conference on Neural Information Processing Systems - Volume 1*, NIPS'14, page 10981106, Cambridge, MA, USA, 2014. MIT Press.

Jonathan D Rosenblatt and Boaz Nadler. On the optimality of averaging in distributed statistical learning. *Inf. Inference: J. IMA*, 5(4):379–404, 2016.

Ohad Shamir, Nati Srebro, and Tong Zhang. Communication-efficient distributed optimization using an approximate newton-type method. In Eric P. Xing and Tony Jebara, editors, *Proceedings of the 31st International Conference on Machine Learning*, volume 32 of *Proceedings of Machine Learning Research*, pages 1000–1008, Bejing, China, 22–24 Jun 2014. PMLR.

Qiang Sun, Wen-Xin Zhou, and Jianqing Fan. Adaptive huber regression. *J. Am. Stat. Assoc.*, 115(529):254–265, 2020.

Sara Van de Geer, Peter Bühlmann, Yaacov Ritov, and Ruben Dezeure. On asymptotically optimal confidence regions and tests for high-dimensional models. *Ann. Stat.*, 42(3)(3):1166–1202, 2014.

Lan Wang, Yichao Wu, and Runze Li. Quantile regression for analyzing heterogeneity in ultra-high dimension. *J. Am. Stat. Assoc.*, 107(497):214–222, 2012.

Lan Wang, Bo Peng, Jelena Bradic, Runze Li, and Yunan Wu. A tuning-free robust and efficient approach to high-dimensional regression. *J. Am. Stat. Assoc.*, 115(532):1700–1714, 2020.

Haofan Yin, Tianxiao Gao, Jinye Xie, Zhijian Huang, Xiaoyan Zhang, Fengyu Yang, Weiwei Qi, Zhonghan Yang, Ti Zhou, Guoquan Gao, et al. Fubp1 promotes colorectal cancer stemness and metastasis via dvl1-mediated activation of wnt/β-catenin signaling. *Mol. Oncol.*, 15(12): 3490–3512, 2021.

Cun-Hui Zhang and Jian Huang. The sparsity and bias of the lasso selection in high-dimensional linear regression. *Ann. Stat.*, 36(4):1567–1594, 2008.

Cun-Hui Zhang and Stephanie S Zhang. Confidence intervals for low dimensional parameters in high dimensional linear models. *J. R. Stat. Soc. Ser. B (Stat. Methodol.)*, 76(1):217–242, 2014.

Yuchen Zhang, Martin J Wainwright, and John C Duchi. Communication-efficient algorithms for statistical optimization. *J. Mach. Learn. Res.*, 14(68):3321–3363, 2013.

Weihua Zhao, Fode Zhang, and Heng Lian. Debiasing and distributed estimation for high-dimensional quantile regression. *IEEE Trans. Neural Netw. Learn. Syst.*, 31(7):2569–2577, 2019.

Le Zhou, Boxiang Wang, and Hui Zou. Sparse convoluted rank regression in high dimensions. *J. Am. Stat. Assoc.*, 119(546):1500–1512, 2024.

Ling Zhou, Ziyang Gong, and Pengcheng Xiang. Distributed computing and inference for big data. *Annu. Rev. Stat. Its Appl.*, 11:533–551, 2023.