EQUIVARIANT ENERGY-GUIDED SDE FOR INVERSE MOLECULAR DESIGN

ABSTRACT

Inverse molecular design is critical in material science and drug discovery, where the generated molecules should satisfy certain desirable properties. In this paper, we propose *equivariant energy-guided stochastic differential equations* (EEGSDE), a flexible framework for controllable 3D molecule generation under the guidance of an energy function in diffusion models. Formally, we show that EEGSDE naturally exploits the geometric symmetry in 3D molecular conformation, as long as the energy function is invariant to orthogonal transformations. Empirically, under the guidance of designed energy functions, EEGSDE significantly improves the baseline on QM9, in inverse molecular design targeted to quantum properties and molecular structures. Furthermore, EEGSDE is able to generate molecules with multiple target properties by combining the corresponding energy functions linearly.

1 INTRODUCTION

The discovery of new molecules with desired properties is critical in many fields, such as the drug and material design (Hajduk & Greer, 2007; Mandal et al., 2009; Kang et al., 2006; Pyzer-Knapp et al., 2015). However, brute-force search in the overwhelming molecular space is extremely challenging. Recently, inverse molecular design (Zunger, 2018) provides an efficient way to explore the molecular space, which directly predicts promising molecules that exhibit desired properties.

A natural way of inverse molecular design is to train a conditional generative model (Sanchez-Lengeling & Aspuru-Guzik, 2018). Formally, it learns a distribution of molecules conditioned on certain properties from data, and new molecules are predicted by sampling from the distribution with the condition set to desired properties. Among them, *equivariant diffusion models* (EDM) (Hoogeboom et al., 2022) leverage the current state-of-art diffusion models (Ho et al., 2020), which involves a forward process to perturb data and a reverse process to generate 3D molecules conditionally or unconditionally. While EDM generates stable and valid 3D molecules, we argue that a single conditional generative model is insufficient for generating accurate molecules that exhibit desired properties (see Table 1 and Table 2 for an empirical verification).

In this work, we propose *equivariant energy-guided stochastic differential equations* (EEGSDE), a flexible framework for controllable 3D molecule generation under the guidance of an energy function in diffusion models. EEGSDE formalizes the generation process as an equivariant stochastic differential equation, and plug in energy functions to improve the controllability of generation. Formally, we show that EEGSDE naturally exploits the geometric symmetry in 3D molecular conformation, as long as the energy function is invariant to orthogonal transformations.

We apply EEGSDE to various applications by carefully designing task-specific energy functions. When targeted to quantum properties, EEGSDE is able to generate more accurate molecules than EDM, e.g., reducing the mean absolute error by more than 30% on the dipole moment property. When targeted to specific molecular structures, EEGSDE better capture the structure information in molecules than EDM, e.g, improving the similarity to target structures by more than 10%. Furthermore, EEGSDE is able to generate molecules targeted to multiple properties by combining the corresponding energy functions linearly. These demonstrate that our EEGSDE enables a flexible and controllable generation of molecules, providing a smart way to explore the chemical space.

$$\boxed{z_T} \text{---} \mathrm{d}z = [f(t)z + g(t)^2(\frac{1}{\sqrt{\beta_{t|0}}}\epsilon_\theta(z,t) + \underbrace{(\nabla_x E(z,c,t) - \overline{\nabla_x E(z,c,t)}, \nabla_h E(z,c,t)))}]\mathrm{d}t + g(t)\mathrm{d}(\tilde{w}_x, \tilde{w}_h) \text{---} \boxed{z_0}$$

Equivariant Energy Guidance

Figure 1: Overview of our EEGSDE. EEGSDE iteratively generates molecules with desired properties (represented by the condition c) by adopting the guidance of energy functions in each step. As the energy function is invariant to rotational transformation R, its gradient (i.e., the energy guidance) is equivariant to R, and therefore the distribution of generated samples is invariant to R.

2 RELATED WORK

Diffusion models are initially proposed by Sohl-Dickstein et al. (2015). Recently, they are better understood in theory by connecting it to score matching and stochastic differential equations (SDE) (Ho et al., 2020; Song et al., 2020). After that, diffusion models have shown strong empirical performance in many applications Dhariwal & Nichol (2021); Ramesh et al. (2022); Chen et al. (2020); Kong et al. (2020). There are also variants proposed to improve or accelerate diffusion models (Nichol & Dhariwal, 2021; Vahdat et al., 2021; Dockhorn et al., 2021; Bao et al., 2022b;a; Salimans & Ho, 2022; Lu et al., 2022).

Guidance is a technique to control the generation process of diffusion models. Initially, Song et al. (2020); Dhariwal & Nichol (2021) use classifier guidance to generate samples belonging to a class. Then, the guidance is extended to CLIP (Radford et al., 2021) for text to image generation, and semantic-aware energy (Zhao et al., 2022) for image-to-image translation. Prior guidance methods focus on image data, and are nontrivial to apply to molecules, since they do not consider the geometric symmetry. In contrast, our work proposes a general guidance framework for 3D molecules, which leverage the geometric symmetry of molecules.

Molecule generation. Several works attempt to model molecules as 3D objects via deep generative models Nesterov et al. (2020); Gebauer et al. (2019); Satorras et al. (2021a); Hoffmann & Noé (2019); Hoogeboom et al. (2022). Among them, the most relevant one is the equivariant diffusion model (EDM) (Hoogeboom et al., 2022), which generates molecules in an iterative denoising manner. Benefiting from recent advances of diffusion models, EDM is stable to train and is able to generate high quality molecules. We provide a formal description of EDM in Section 3.

Some other methods generate simplified representations of molecules, such as 1D SMILES strings (Weininger, 1988) and 2D graphs of molecules. These include variational autoencoders (Kusner et al., 2017; Dai et al., 2018; Jin et al., 2018; Simonovsky & Komodakis, 2018; Liu et al., 2018), normalizing flows (Madhawa et al., 2019; Zang & Wang, 2020; Luo et al., 2021), generative adversarial networks (Bian et al., 2019; Assouel et al., 2018), and autoregressive models (Popova et al., 2019; Flam-Shepherd et al., 2021).

Inverse molecular design. Generative models have been applied to inverse molecular design. For example, conditional autoregressive models (Gebauer et al., 2022) and EDM (Hoogeboom et al., 2022) directly generate 3D molecules with desired quantum properties. Gebauer et al. (2019)

also finetune pretrained generative models on a biased subset to generate 3D molecules with small HOMO-LUMO gaps. In contrast to these conditional generative models, our work further proposes a guidance method, a flexible way to control the generation process of molecules.

Some other methods apply optimization methods to search molecules with desired properties, such as reinforcement learning (Zhou et al., 2019; You et al., 2018) and genetic algorithms (Jensen, 2019; Nigam et al., 2019). These optimization methods generally consider the 1D SMILES strings or 2D graphs of molecules, and the 3D information is not provided.

3 BACKGROUND

3D representation of molecules. Suppose a molecule has M atoms and let $\boldsymbol{x}^i \in \mathbb{R}^n$ ($n = 3$ in general) be the coordinate of the ith atom. The collection of coordinates $\boldsymbol{x} = (\boldsymbol{x}^1, \ldots, \boldsymbol{x}^M) \in \mathbb{R}^{Mn}$ determines the *conformation* of the molecule. In addition to the coordinate, each atom is also associated with an atom feature, e.g., the atom type. We use $\boldsymbol{h}^i \in \mathbb{R}^d$ to represent the atom feature of the ith atom, and use $\boldsymbol{h} = (\boldsymbol{h}^1, \ldots, \boldsymbol{h}^M) \in \mathbb{R}^{Md}$ to represent the collection of atom features in a molecule. We use a tuple $\boldsymbol{z} = (\boldsymbol{x}, \boldsymbol{h})$ to represent a molecule, which contains both the 3D geometry information and the atom feature information.

Equivariance and invariance. Suppose $\boldsymbol{R}$ is a transformation. A distribution $p(\boldsymbol{x}, \boldsymbol{h})$ is said to be *invariant* to $\boldsymbol{R}$, if $p(\boldsymbol{x}, \boldsymbol{h}) = p(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h})$ holds for all $\boldsymbol{x}$ and $\boldsymbol{h}$. Here $\boldsymbol{R}\boldsymbol{x} = (\boldsymbol{R}\boldsymbol{x}^1, \ldots, \boldsymbol{R}\boldsymbol{x}^M)$ is applied to each coordinate. A function $(\boldsymbol{a}^x, \boldsymbol{a}^h) = \boldsymbol{f}(\boldsymbol{x}, \boldsymbol{h})$ that have two components $\boldsymbol{a}^x, \boldsymbol{a}^h$ in its output is said to be *equivariant* to $\boldsymbol{R}$, if $\boldsymbol{f}(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}) = (\boldsymbol{R}\boldsymbol{a}^x, \boldsymbol{a}^h)$ holds for all $\boldsymbol{x}$ and $\boldsymbol{h}$. A function $\boldsymbol{f}(\boldsymbol{x}, \boldsymbol{h})$ is said to be *invariant* to $\boldsymbol{R}$, if $\boldsymbol{f}(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}) = \boldsymbol{f}(\boldsymbol{x}, \boldsymbol{h})$ holds for all $\boldsymbol{x}$ and $\boldsymbol{h}$.

Zero CoM subspace. It has been shown that the invariance to translational and rotational transformations is an important factor for the success of 3D molecule modeling (Köhler et al., 2020; Xu et al., 2022). However, the translational invariance is impossible for a distribution in the full space $\mathbb{R}^{Mn}$ (Satorras et al., 2021a). Nevertheless, we can view two collections of coordinates $\boldsymbol{x}$ and $\boldsymbol{y}$ as equivalent if $\boldsymbol{x}$ can be translated from $\boldsymbol{y}$, since the translation doesn't change the identity of a molecule. Such an equivalence relation partitions the whole space $\mathbb{R}^{Mn}$ into disjoint equivalence classes. Indeed, all elements in the same equivalence classes represent the same conformation, and we can use the element with zero center of mass (CoM), i.e., $\frac{1}{M} \sum_{i=1}^{M} \boldsymbol{x}^i = \boldsymbol{0}$, as the specific representation. These elements collectively form the *zero CoM linear subspace* X (Xu et al., 2022; Hoogeboom et al., 2022), and the rest of the paper always uses elements in X to represent conformations.

Equivariant graph neural network. Satorras et al. (2021b) propose *equivariant graph neural networks* (EGNNs), which incorporate the equivariance inductive bias into neural networks. Specifically, $(\boldsymbol{a}^x, \boldsymbol{a}^h) = \text{EGNN}(\boldsymbol{x}, \boldsymbol{h})$ is a composition of L *equivariant convolutional layers*. The l-th layer takes the tuple $(\boldsymbol{x}_l, \boldsymbol{h}_l)$ as the input and outputs an updated version $(\boldsymbol{x}_{l+1}, \boldsymbol{h}_{l+1})$, as follows:

$$\boldsymbol{m}^{ij} = \Phi_m(\boldsymbol{h}_l^i, \boldsymbol{h}_l^j, \|\boldsymbol{x}_l^i - \boldsymbol{x}_l^j\|_2^2, e^{ij}; \boldsymbol{\theta}_m), \ w^{ij} = \Phi_w(\boldsymbol{m}^{ij}; \boldsymbol{\theta}_w), \ \boldsymbol{h}_{l+1}^i = \Phi_h(\boldsymbol{h}_l^i, \sum_{j \neq i} w^{ij} \boldsymbol{m}^{ij}; \boldsymbol{\theta}_h),$$

$$\boldsymbol{x}_{l+1}^i = \boldsymbol{x}_l^i + \sum_{j \neq i} \frac{\boldsymbol{x}_l^i - \boldsymbol{x}_l^j}{\|\boldsymbol{x}_l^i - \boldsymbol{x}_l^j\|_2 + 1} \Phi_x(\boldsymbol{h}_l^i, \boldsymbol{h}_l^j, \|\boldsymbol{x}_l^i - \boldsymbol{x}_l^j\|_2^2, e^{ij}; \boldsymbol{\theta}_x),$$

where $\Phi_m, \Phi_w, \Phi_h, \Phi_x$ are parameterized by fully connected neural networks with parameters $\boldsymbol{\theta}_m, \boldsymbol{\theta}_w, \boldsymbol{\theta}_h, \boldsymbol{\theta}_x$ respectively, and e^{ij} are optional feature attributes. We can verify that these layers are equivariant to orthogonal transformations, which include rotational transformations as special cases. As their composition, the EGNN is also equivariant to orthogonal transformations. Furthermore, let $\text{EGNN}^h(\boldsymbol{x}, \boldsymbol{h}) = \boldsymbol{a}^h$, i.e., the second component in the output of the EGNN. Then $\text{EGNN}^h(\boldsymbol{x}, \boldsymbol{h})$ is invariant to orthogonal transformations.

Equivariant diffusion models (EDM) (Hoogeboom et al., 2022) are a variant of diffusion models (Ho et al., 2020) for molecule data. EDMs gradually inject noise to the molecule data $\boldsymbol{z} = (\boldsymbol{x}, \boldsymbol{h})$ via a forward process

$$q(\boldsymbol{z}_{1:N}|\boldsymbol{z}_0) = \prod_{n=1}^{N} q(\boldsymbol{z}_n|\boldsymbol{z}_{n-1}), \ q(\boldsymbol{z}_n|\boldsymbol{z}_{n-1}) = \mathcal{N}_X(\boldsymbol{x}_n|\sqrt{\alpha_n}\boldsymbol{x}_{n-1}, \beta_n)\mathcal{N}(\boldsymbol{h}_n|\sqrt{\alpha_n}\boldsymbol{h}_{n-1}, \beta_n), \ (1)$$

where α_n and β_n represent the noise schedule and satisfy $\alpha_n + \beta_n = 1$, and $\mathcal{N}_X$ represent the Gaussian distribution in the zero CoM subspace X (see its formal definition in Appendix A.2). Let $\overline{\alpha}_n = \alpha_1 \alpha_2 \cdots \alpha_n$, $\overline{\beta}_n = 1 - \overline{\alpha}_n$ and $\tilde{\beta}_n = \beta_n \overline{\beta}_{n-1} / \overline{\beta}_n$. To generate samples, the forward process is reversed using a Markov chain:

$$p(\boldsymbol{z}_{0:N}) = p(\boldsymbol{z}_N) \prod_{n=1}^{N} p(\boldsymbol{z}_{n-1}|\boldsymbol{z}_n), p(\boldsymbol{z}_{n-1}|\boldsymbol{z}_n) = \mathcal{N}_X(\boldsymbol{x}_{n-1}|\boldsymbol{\mu}_n^x(\boldsymbol{z}_n), \tilde{\beta}_n)\mathcal{N}(\boldsymbol{h}_{n-1}|\boldsymbol{\mu}_n^h(\boldsymbol{z}_n), \tilde{\beta}_n). \quad (2)$$

Here $p(\boldsymbol{z}_N) = \mathcal{N}_X(\boldsymbol{x}_N|\boldsymbol{0}, 1)\mathcal{N}(\boldsymbol{h}_N|\boldsymbol{0}, 1)$. The mean $\boldsymbol{\mu}_n(\boldsymbol{z}_n) = (\boldsymbol{\mu}_n^x(\boldsymbol{z}_n), \boldsymbol{\mu}_n^h(\boldsymbol{z}_n))$ is parameterized by a noise prediction network $\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}_n, n)$, and is trained using a MSE loss, as follows:

$$\boldsymbol{\mu}_n(\boldsymbol{z}_n) = \frac{1}{\sqrt{\alpha_n}}(\boldsymbol{z}_n - \frac{\beta_n}{\sqrt{\overline{\beta}_n}}\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}_n, n)), \quad \min_{\boldsymbol{\theta}} \mathbb{E}_n \mathbb{E}_{q(\boldsymbol{z}_0, \boldsymbol{z}_n)} w(t) \|\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}_n, n) - \boldsymbol{\epsilon}_n\|^2,$$

where $\boldsymbol{\epsilon}_n = \frac{\boldsymbol{z}_n - \sqrt{\overline{\alpha}_n}\boldsymbol{z}_0}{\sqrt{\overline{\beta}_n}}$ is the standard Gaussian noise injected to $\boldsymbol{z}_0$ and $w(t)$ is the weight term.

Hoogeboom et al. (2022) show that the distribution of generated samples $p(\boldsymbol{z}_0)$ is invariant to rotational transformations if the noise prediction network is equivariant to orthogonal transformations. In Section 4.2, we extend this proposition to the SDE formulation of molecular diffusion modelling.

Hoogeboom et al. (2022) also present a conditional version of EDM for inverse molecular design by adding an extra input of the condition to the noise prediction network.

4 Equivariant Energy-Guided SDE

In this part, we introduce our equivariant energy-guided SDE (EEGSDE), which is illustrated in Figure 1. EEGSDE is based on the SDE formulation of molecular diffusion modeling, which is described in Section 4.1 and Section 4.2. Then, we formally present our EEGSDE in Section 4.3.

We provide proof and derivations in Appendix A.

4.1 SDE in the Product Space

Recall that a molecule is represented as a tuple $\boldsymbol{z} = (\boldsymbol{x}, \boldsymbol{h})$, where $\boldsymbol{x} = (\boldsymbol{x}^1, \ldots, \boldsymbol{x}^M) \in X$ represents the conformation and $\boldsymbol{h} = (\boldsymbol{h}^1, \ldots, \boldsymbol{h}^M) \in \mathbb{R}^{Md}$ represents atom features. Here $X = \{\boldsymbol{x} \in \mathbb{R}^{Mn} : \frac{1}{M}\sum_{i=1}^{M} \boldsymbol{x}^i = \boldsymbol{0}\}$ is the zero CoM subspace as mentioned in Section 3, and d is the dimension of features. We first introduce a continuous-time diffusion process $\{\boldsymbol{z}_t\}_{0 \le t \le T}$ in the product space $X \times \mathbb{R}^{Md}$, which gradually adds noise to both $\boldsymbol{x}$ and $\boldsymbol{h}$. This can be described by the following forward SDE:

$$\mathrm{d}\boldsymbol{z} = f(t)\boldsymbol{z}\mathrm{d}t + g(t)\mathrm{d}(\boldsymbol{w}_x, \boldsymbol{w}_h), \quad \boldsymbol{z}_0 \sim q(\boldsymbol{z}_0), \quad (3)$$

where $f(t)$ and $g(t)$ are two scalar functions, $\boldsymbol{w}_x$ and $\boldsymbol{w}_h$ are independent standard Wiener processes in X and $\mathbb{R}^{Md}$ respectively, and the SDE starts from the data distribution $q(\boldsymbol{z}_0)$. Note that $\boldsymbol{w}_x$ can be constructed by subtracting the CoM of a standard Wiener process $\boldsymbol{w}$ in $\mathbb{R}^{Mn}$, i.e., $\boldsymbol{w}_x = \boldsymbol{w} - \overline{\boldsymbol{w}}$, where $\overline{\boldsymbol{w}} = \frac{1}{M}\sum_{i=1}^{M} \boldsymbol{w}^i$ is the CoM of $\boldsymbol{w} = (\boldsymbol{w}^1, \ldots, \boldsymbol{w}^M)$. It can be shown that the SDE has a linear Gaussian transition kernel $q(\boldsymbol{z}_t|\boldsymbol{z}_s) = q(\boldsymbol{x}_t|\boldsymbol{x}_s)q(\boldsymbol{h}_t|\boldsymbol{h}_s)$ from $\boldsymbol{x}_s$ to $\boldsymbol{x}_t$, where $0 \le s < t \le T$. Specifically, there exists two scalars $\alpha_{t|s}$ and $\beta_{t|s}$, s.t., $q(\boldsymbol{x}_t|\boldsymbol{x}_s) = \mathcal{N}_X(\boldsymbol{x}_t|\sqrt{\alpha_{t|s}}\boldsymbol{x}_s, \beta_{t|s})$ and $q(\boldsymbol{h}_t|\boldsymbol{h}_s) = \mathcal{N}(\boldsymbol{h}_t|\sqrt{\alpha_{t|s}}\boldsymbol{h}_s, \beta_{t|s})$. Here $\mathcal{N}_X$ denotes the Gaussian distribution in the subspace X, and see Appendix A.2 for its formal definition. Indeed, the forward process of EDM in Eq. (1) is a discretization of the forward SDE in Eq. (3).

To generate molecules, we reverse the diffusion process in Eq. (3) from T to 0. Such a time reversal forms another a SDE, which can be represented by both the *score function form* and the *noise prediction form*:

$$\mathrm{d}\boldsymbol{z} = [f(t)\boldsymbol{z} - g(t)^2 \underbrace{(\nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{z}) - \overline{\nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{z})}, \nabla_{\boldsymbol{h}} \log q_t(\boldsymbol{z}))}_{\text{score function form}}]\mathrm{d}t + g(t)\mathrm{d}(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h),$$

$$= [f(t)\boldsymbol{z} + \frac{g(t)^2}{\sqrt{\beta_{t|0}}} \underbrace{\mathbb{E}_{q(\boldsymbol{z}_0|\boldsymbol{z}_t)}\boldsymbol{\epsilon}_t}_{\text{noise prediction form}}]\mathrm{d}t + g(t)\mathrm{d}(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h), \quad \boldsymbol{z}_T \sim q_T(\boldsymbol{z}_T). \quad (4)$$

Here $q_t(z)$ is the marginal distribution of z_t, $\nabla_x \log q_t(z)$ is the gradient of $\log q_t(z)$ w.r.t. x[1], $\overline{\nabla_x \log q_t(z)} = \frac{1}{M} \sum_{i=1}^{M} \nabla_{x^i} \log q_t(z)$ is the CoM of $\nabla_x \log q_t(z)$, dt is the infinitesimal negative timestep, $\tilde{w}_x$ and $\tilde{w}_h$ are independent reverse-time standard Wiener processes in X and $\mathbb{R}^{Md}$ respectively, and $\epsilon_t = \frac{z_t - \sqrt{\alpha_{t|0}} z_0}{\sqrt{\beta_{t|0}}}$ is the standard Gaussian noise injected to z_0. Compared to the original SDE introduced by Song et al. (2020), our reverse SDE in Eq. (4) additionally subtracts the CoM of $\nabla_x \log q_t(z)$. This ensures x_t always stays in the zero CoM subspace as time flows back.

To sample from the reverse SDE in Eq. (4), we use a noise prediction network $\epsilon_\theta(z_t, t)$ to estimate $\mathbb{E}_{q(z_0|z_t)} \epsilon_t$, through minimizing the following MSE loss

$$\min_{\theta} \mathbb{E}_t \mathbb{E}_{q(z_0, z_t)} w(t) \|\epsilon_\theta(z_t, t) - \epsilon_t\|^2,$$

where t is uniformly sampled from $[0, T]$, and $w(t)$ controls the weight of the loss term at time t. Note that the noise ϵ_t is in the product space $X \times \mathbb{R}^{Md}$, so we subtract the CoM of the predicted noise of x_t to ensure $\epsilon_\theta(z_t, t)$ is also in the product space.

Substituting $\epsilon_\theta(z_t, t)$ into Eq. (4), we get an approximate reverse-time SDE parameterized by θ:

$$dz = [f(t)z + \frac{g(t)^2}{\sqrt{\beta_{t|0}}} \epsilon_\theta(z, t)]dt + g(t)d(\tilde{w}_x, \tilde{w}_h), \quad z_T \sim p_T(z_T), \tag{5}$$

where $p_T(z_T) = \mathcal{N}_X(x_T | \mathbf{0}, 1) \mathcal{N}(h_T | \mathbf{0}, 1)$ is a Gaussian prior in the product space that approximates $q_T(z_T)$. We define $p_\theta(z_0)$ as the marginal distribution of Eq. (5) at time $t = 0$, which is the distribution of our generated samples. Similarly to the forward process, the reverse process of EDM in Eq. (2) is a discretization of the reverse SDE in Eq. (5).

4.2 Equivariant SDE

To leverage the geometric symmetry in 3D molecular conformation, $p_\theta(z_0)$ should be invariant to translational and rotational transformations. As mentioned in Section 3, the translational invariance of $p_\theta(z_0)$ is already satisfied by considering the zero CoM subspace. The rotational invariance can be satisfied if the noise prediction network is equivariant to orthogonal transformations, as summarized in the following theorem:

Theorem 1. *Let $(\epsilon_\theta^x(z_t, t), \epsilon_\theta^h(z_t, t)) = \epsilon_\theta(z_t, t)$, where $\epsilon_\theta^x(z_t, t)$ and $\epsilon_\theta^h(z_t, t)$ are the predicted noise of x_t and h_t respectively. If for any orthogonal transformation $R \in \mathbb{R}^{n \times n}$, $\epsilon_\theta(z_t, t)$ is equivariant to R, i.e., $\epsilon_\theta(Rx_t, h_t, t) = (R\epsilon_\theta^x(x_t, h_t, t), \epsilon_\theta^h(x_t, h_t, t))$, and $p_T(z_T)$ is invariant to R, i.e., $p_T(Rx_T, h_T) = p_T(x_T, h_T)$, then $p_\theta(z_0)$ is invariant to any rotational transformation.*

As mentioned in Section 3, the EGNN satisfies the equivariance constraint, and we parameterize $\epsilon_\theta(z_t, t)$ using an EGNN following Hoogeboom et al. (2022). See details in Appendix C.

4.3 Equivariant Energy-Guided SDE

Now we describe *equivariant energy-guided SDE* (EEGSDE), which guides the generated molecules of Eq. (5) towards desired properties c by leveraging a time-dependent energy function $E(z, c, t)$:

$$dz = [f(t)z + g(t)^2(\frac{1}{\sqrt{\beta_{t|0}}} \epsilon_\theta(z, t)$$
$$+ \underbrace{(\nabla_x E(z, c, t) - \overline{\nabla_x E(z, c, t)}, \nabla_h E(z, c, t))}_{\text{energy gradient taken in the product space}})]dt + g(t)d(\tilde{w}_x, \tilde{w}_h), \ z_T \sim p_T(z_T), \tag{6}$$

which defines a distribution $p_\theta(z_0|c)$ conditioned on the property c. Here the CoM $\overline{\nabla_x E(z, c, t)}$ of the gradient is subtracted to keep the SDE in the product space, which ensures the translational invariance of $p_\theta(z_0|c)$. Besides, the rotational invariance is satisfied by using energy invariant to orthogonal transformations, as summarized in the following theorem:

---

[1] While $q_t(z)$ is defined in $X \times \mathbb{R}^{Md}$, its domain can be extended to $\mathbb{R}^{Mn} \times \mathbb{R}^{Md}$ and the gradient is valid. See Remark 1 in Appendix A.2 for details.

Algorithm 1 Sample from EEGSDE using the Euler-Maruyama method

---

Require: Number of steps N
  $\Delta t = \frac{T}{N}$
  $\boldsymbol{z} \leftarrow (\boldsymbol{x} - \overline{\boldsymbol{x}}, \boldsymbol{h})$, where $\boldsymbol{x} \sim \mathcal{N}(\boldsymbol{0}, 1), \boldsymbol{h} \sim \mathcal{N}(\boldsymbol{0}, 1)$ {Sample from the prior $p_T(\boldsymbol{z}_T)$}
  for $i = N$ to 1 **do**
    $t \leftarrow i\Delta t$
    $\boldsymbol{g}_x \leftarrow \nabla_{\boldsymbol{x}} E(\boldsymbol{z}, c, t), \boldsymbol{g}_h \leftarrow \nabla_{\boldsymbol{h}} E(\boldsymbol{z}, c, t)$ {Calculate the gradient of the energy function}
    $\boldsymbol{g} \leftarrow (\boldsymbol{g}_x - \overline{\boldsymbol{g}_x}, \boldsymbol{g}_h)$ {Subtract the CoM of the gradient}
    $\boldsymbol{F} \leftarrow f(t)\boldsymbol{z} + g(t)^2(\frac{1}{\sqrt{\beta_{t|0}}}\boldsymbol{\epsilon_\theta}(\boldsymbol{z}, t) + \boldsymbol{g})$
    $\boldsymbol{\epsilon} \leftarrow (\boldsymbol{\epsilon}_x - \overline{\boldsymbol{\epsilon}_x}, \boldsymbol{\epsilon}_h)$, where $\boldsymbol{\epsilon}_x \sim \mathcal{N}(\boldsymbol{0}, 1), \boldsymbol{\epsilon}_h \sim \mathcal{N}(\boldsymbol{0}, 1)$
    $\boldsymbol{z} \leftarrow \boldsymbol{z} - \boldsymbol{F}\Delta t + g(t)\sqrt{\Delta t}\boldsymbol{\epsilon}$ {Update $\boldsymbol{z}$ according to Eq. (6)}
  end for
  return $\boldsymbol{z}$

Theorem 2. *Suppose the assumptions in Theorem 1 hold and $E(\boldsymbol{z}, c, t)$ is invariant to any orthogonal transformation $\boldsymbol{R}$, i.e., $E(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}, c, t) = E(\boldsymbol{x}, \boldsymbol{h}, c, t)$. Then $p_{\boldsymbol{\theta}}(\boldsymbol{z}_0|c)$ is invariant to any rotational transformation.*

Note that we can also use a conditional model $\boldsymbol{\epsilon_\theta}(\boldsymbol{z}, c, t)$ in Eq. (6). See Appendix B for details.

To sample from $p_{\boldsymbol{\theta}}(\boldsymbol{z}_0|c)$, various solvers can be used for Eq. (6), such as the Euler-Maruyama method (Song et al., 2020) and the Analytic-DPM sampler (Bao et al., 2022b;a). In Algorithm 1, we present the Euler-Maruyama method as an example.

Compared with Eq. (5), Eq. (6) additionally performs a gradient descent on the energy function. This encourages generated molecules to have a low energy. Thus, we can design the energy function according to the consistency between the molecule $\boldsymbol{z}$ and the property c. In the rest of the paper, we specify the choice of energy functions, and show these energies improve controllable molecule generation targeted to quantum properties, molecular structures, and even a combination of them.

5   Generating Molecules with Desired Quantum Properties

Let $c \in \mathbb{R}$ be a certain quantum property. To generate molecules with the desired property, we set the energy function as the squared error between the predicted property and the desired property:

$$E(\boldsymbol{z}_t, c, t) = s|g(\boldsymbol{z}_t, t) - c|^2, \tag{7}$$

where $g(\boldsymbol{z}_t, t)$ is a time-dependent property prediction model, and s is the scaling factor controlling the strength of the guidance. Specifically, we parameterize $g(\boldsymbol{z}_t, t)$ using the second component in the output of EGNN (see Section 3) followed by a decoder (Dec):

$$g(\boldsymbol{z}_t, t) = \text{Dec}(\text{EGNN}^h(\boldsymbol{x}_t, \boldsymbol{h}'_t)), \quad \boldsymbol{h}'_t = \text{concatenate}(\boldsymbol{h}_t, t), \tag{8}$$

where the concatenation is performed on each atom feature, and the decoder consists of a node-wise multilayer perceptron (MLP), a sum pooling layer and an MLP for the final property prediction based on Satorras et al. (2021b). The training objective of $g(\boldsymbol{z}_t, t)$ is available in Appendix D. This parameterization ensures that the energy function $E(\boldsymbol{z}_t, c, t)$ is invariant to orthogonal transformations, and thus the distribution of generated samples is also invariant according to Theorem 2.

We evaluate on the QM9 dataset (Ramakrishnan et al., 2014), which contains quantum properties and coordinates of $\sim$130k organic molecules with up to nine heavy atoms from (C, N, O, F). We follow Hoogeboom et al. (2022) for training and evaluation settings. **Data split:** We split QM9 into training, validation and test sets, which include 100K, 18K and 13K samples respectively. The training set is further divided into two non-overlapping halves D_a, D_b equally. **Training setting:** We train the noise prediction network and the energy function on D_b. **Evaluation setting:** We report the mean absolute error (MAE), which is evaluated by a property classifier network ϕ_c (Satorras et al., 2021b) trained on the other half D_a. We report the loss of ϕ_c on D_b as the lower bound (L-bound) (Hoogeboom et al., 2022) of the MAE. We also report the novelty of generated molecules, as well as the atom stability (AS) and molecule stability (MS), which measure the validity of generated molecules. Since the noise prediction network and the energy function are trained on D_b,

Table 1: Generation performance targeted to a single quantum property (its unit is indicated in the parentheses after it). We report the mean absolute error (MAE), as well as the novelty, the atom stability (AS), and the molecule stability (MS). We reproduce L-bound and EDM results, which is consistent with Hoogeboom et al. (2022) (see Appendix F.2). "#Atoms" uses public results from Hoogeboom et al. (2022).

Method	MAE↓	Novelty↑	AS↑	MS↑	Method	MAE↓	Novelty↑	AS↑	MS↑
C_v ($\frac{\text{cal}}{\text{mol}}$K)					μ (D)				
U-bound	6.873	-	-	-	U-bound	1.611	-	-	-
#Atoms	1.971	-	-	-	#Atoms	1.053	-	-	-
EDM	1.066	83.64	**98.25**	80.50	EDM	1.138	84.04	98.14	80.04
EEGSDE (s=1)	1.038	83.71	98.21	**80.61**	EEGSDE (s=0.5)	0.935	84.06	**98.20**	**80.05**
EEGSDE (s=5)	0.981	83.75	98.12	79.84	EEGSDE (s=1)	0.853	**84.65**	98.13	79.66
EEGSDE (s=10)	**0.939**	**83.92**	98.06	79.21	EEGSDE (s=2)	**0.772**	**84.65**	98.08	79.07
L-bound	0.040	-	-	-	L-bound	0.043	-	-	-
$\Delta\varepsilon$ (meV)					$\varepsilon_{\text{HOMO}}$ (meV)				
U-bound	1460	-	-	-	U-bound	644	-	-	-
#Atoms	866	-	-	-	#Atoms	426	-	-	-
EDM	666	84.44	**98.27**	81.83	EDM	373	84.38	**98.24**	**79.97**
EEGSDE (s=0.5)	577	83.80	98.11	80.73	EEGSDE (s=0.1)	362	**84.82**	98.23	79.88
EEGSDE (s=1)	544	84.22	98.05	79.65	EEGSDE (s=0.5)	319	84.38	98.15	79.37
EEGSDE (s=3)	**485**	**85.87**	97.72	77.01	EEGSDE (s=1)	**302**	84.68	98.10	79.20
L-bound	65	-	-	-	L-bound	39	-	-	-
α (Bohr^3)					$\varepsilon_{\text{LUMO}}$ (meV)				
U-bound	8.98	-	-	-	U-bound	1451	-	-	-
#Atoms	3.86	-	-	-	#Atoms	813	-	-	-
EDM	2.74	84.30	98.13	79.32	EDM	595	84.57	98.21	81.01
EEGSDE (s=0.5)	2.65	84.01	**98.21**	80.46	EEGSDE (s=0.5)	525	84.03	**98.29**	81.44
EEGSDE (s=1)	2.61	**84.83**	98.15	80.09	EEGSDE (s=1)	496	84.40	**98.29**	**81.49**
EEGSDE (s=3)	**2.50**	83.91	98.21	80.79	EEGSDE (s=3)	**442**	85.08	98.14	80.20
L-bound	0.09	-	-	-	L-bound	36	-	-	-

Table 2: Generation performance with multiple quantum properties.

Method	MAE1↓	MAE2↓	Novelty↑	AS↑	MS↑
C_v ($\frac{\text{cal}}{\text{mol}}$K), μ (D)					
EDM	1.075	1.163	85.45	**97.99**	77.02
EEGSDE (s_1=10, s_2=1)	**0.982**	**0.916**	**85.51**	97.58	73.94
$\Delta\varepsilon$ (meV), μ (D)					
EDM	682	1.138	85.04	**97.96**	75.95
EEGSDE (s_1=s_2=1)	**566**	**0.866**	**85.98**	97.65	72.91
α (Bohr^3), μ (D)					
EDM	2.76	1.161	85.47	**98.03**	77.83
EEGSDE (s_1=s_2=1.5)	**2.56**	**0.859**	**85.50**	97.97	77.14

Table 3: Generation performance targeted to structures from test set on QM9 dataset.

Method	Similarity↑
EDM	0.670
EEGSDE (s=0.1)	0.695
EEGSDE (s=0.5)	0.738
EEGSDE (s=1)	**0.748**

the novelty is also evaluated on D_b, which leads to a higher value than that evaluated on the whole dataset (Hoogeboom et al., 2022). We provide additional experimental settings in Appendix E.2.

By default, EEGSDE uses a conditional noise prediction network in Eq. (6), since we find it generates more accurate molecules than using an unconditional one (see Appendix F.1 for an ablation study). We compare our EEGSDE to EDM, which only adopts a conditional noise prediction network. We also compare two additional baselines "U-bound" and "#Atoms" from Hoogeboom et al. (2022) (see Appendix E.1 for details).

Following Hoogeboom et al. (2022), we consider six quantum properties in QM9: polarizability α, highest occupied molecular orbital energy $\varepsilon_{\text{HOMO}}$, lowest unoccupied molecular orbital energy

Figure 2: Examples of generated molecules targeted to specific structures (unseen during training). Compared to EDM, the molecular structures of EEGSDE align better with the target structures.

Figure 3: Visualization of the effect of the scaling factor.

$\varepsilon_{\text{LUMO}}$, HOMO-LUMO gap $\Delta\varepsilon$, dipole moment μ and heat capacity C_v. Firstly, we generate molecules targeted to one of these six properties. As shown in Table 1, with the energy guidance, our EEGSDE has a significantly better MAE than the EDM on all properties. Remarkably, with a proper scaling factor s, the MAE of EEGSDE is reduced by more than 25% compared to EDM on properties $\Delta\varepsilon$, $\varepsilon_{\text{HOMO}}$, and more than 30% on μ. Note that the energy guidance does not affect the novelty, the atom stability and molecule stability much (always within 6% compared to EDMs).

We further generate molecules targeted to multiple quantum properties by combining energies linearly. As shown in Table 2, our EEGSDE still has a significantly better MAE than the EDM, and does not affect the novelty, the atom stability and molecule stability much.

As a result, our EEGSDE is able to explore the chemical space in a guided way to generate promising molecules, which may benefit applications such as drug and material discovery.

6 GENERATING MOLECULES WITH TARGET STRUCTURES

Following Gebauer et al. (2022), we use the molecular fingerprint to encode the structure information of a molecule. The molecular fingerprint $c = (c_1, \ldots, c_L)$ is a series of bits that capture the presence or absence of substructures in the molecule. To generate molecules with a specific structure, we set the energy function as the squared error $E(\boldsymbol{z}_t, c, t) = s\|m(\boldsymbol{z}_t, t) - c\|^2$ between a time-dependent multi-label classifier $m(\boldsymbol{z}_t, t)$ and the corresponding fingerprint c. Here s is the scaling factor. In initial experiments, we also try binary cross entropy loss for the energy, but we find it causes the generation process unstable. The multi-label classifier is parameterized by EGNN as

$$m(\boldsymbol{z}_t, t) = \sigma(\text{Dec}(\text{EGNN}^h(\boldsymbol{x}_t, \boldsymbol{h}_t'))), \quad \boldsymbol{h}_t' = \text{concatenate}(\boldsymbol{h}_t, t).$$

The multi-label classifier has the same backbone to the property prediction model in Eq. (8), except that the decoder outputs a vector of dimension L, and the sigmoid function σ is adopted for multi-label classification. The training objective of $m(\boldsymbol{z}_t, t)$ is available in Appendix D. Similarly to Eq. (8), the EGNN in the multi-label classifier guarantees the invariance of the distribution of generated samples according to Theorem 2.

Figure 4: Generate molecules targeted to both the quantum property α and the molecular structure.

We train the noise prediction network and the energy function on the whole training set of QM9. We evaluate the Tanimoto similarity (Gebauer et al., 2022) between unseen target structures from test set and structures of generated molecules, which is computed by comparing fingerprints of two structures. We provide more experimental details in Appendix E.3.

As shown in Table 3, EEGSDE significantly improves the similarity between target structures and generated structures compared to EDM. Also note in a proper range, a larger scaling factor results in a better similarity, and EEGSDE with s=1 improves the similarity by more than 10% compared to EDM. In Figure 2, we plot generated molecules of EDM and EEGSDE (s=1) targeted to specific structures, where our EEGSDE aligns better with them. We further visualize the effect of the scaling factor in Table 3, where the generated structures align better as the scaling factor grows. These results demonstrate that our EEGSDE captures the structure information in molecules well.

We also perform experiments on the more challenging GEOM-Drug (Axelrod & Gomez-Bombarelli, 2022) dataset, and we train the conditional EDM baseline using the default setting of Hoogeboom et al. (2022). We find the EDM baseline has a similarity of 0.166, which is much lower than QM9. We hypothesize this is because molecules in GEOM-Drug has much more atoms than QM9 with a more complex structure, and the default setting in Hoogeboom et al. (2022) is suboptimal. For example, the EDM on GEOM-Drug has a smaller number of parameters than the EDM on QM9 (15M v.s. 26M), which is insufficient to capture the structure information. Nevertheless, our EEGSDE still improves the similarity by $\sim$%17. We provide more results on GEOM-Drug in Appendix F.3.

Finally, we demonstrate that our EEGSDE is a flexible framework to generate molecules targeted to multiple properties, which is often the practical case. We additionally target to the quantum property α (polarizability) by combining the energy function for molecular structures in this section and the energy function for quantum properties in Section 5. Here we choose $\alpha = 100 \, \text{Bohr}^3$, which is a relatively large value, and we expect it to encourage less isometrically shaped structures. As shown in Figure 4, the generated molecule aligns better with the target structure as the scaling factor s_1 grows, and meanwhile a ring substructure in the generated molecule vanishes as the scaling factor for polarizability s_2 grows, leading to a less isometrically shaped structure, which is as expected.

7 CONCLUSION

This work presents equivariant energy-guided SDE (EEGSDE), a flexible framework for controllable 3D molecule generation under the guidance of an energy function in diffusion models. EEGSDE naturally exploits the geometric symmetry in 3D molecular conformation, as long as the energy function is invariant to orthogonal transformations. EEGSDE significantly improves the EDM baseline in inverse molecular design targeted to quantum properties and molecular structures. Furthermore, EEGSDE is able to generate molecules with multiple target properties by combining the corresponding energy functions linearly.

ETHICS STATEMENT

Inverse molecular design is critical in fields like material science and drug discovery. Our EEGSDE is a flexible framework for inverse molecular design and thus might benefit these fields. Currently the negative consequences are not obvious.

REPRODUCIBILITY STATEMENT

Our code is included in the supplementary material. The implementation of our experiment is described in Section 5 and Section 6. Further details such as the hyperparameters of training and model backbones are provided in Appendix E. We provide complete proofs and derivations of all theoretical results in Appendix A.

REFERENCES

Rim Assouel, Mohamed Ahmed, Marwin H Segler, Amir Saffari, and Yoshua Bengio. Defactor: Differentiable edge factorization-based probabilistic graph generation. *arXiv preprint arXiv:1811.09766*, 2018.

Simon Axelrod and Rafael Gomez-Bombarelli. Geom, energy-annotated molecular conformations for property prediction and molecular generation. *Scientific Data*, 9(1):1–14, 2022.

Fan Bao, Chongxuan Li, Jiacheng Sun, Jun Zhu, and Bo Zhang. Estimating the optimal covariance with imperfect mean in diffusion probabilistic models. *arXiv preprint arXiv:2206.07309*, 2022a.

Fan Bao, Chongxuan Li, Jun Zhu, and Bo Zhang. Analytic-dpm: an analytic estimate of the optimal reverse variance in diffusion probabilistic models. *arXiv preprint arXiv:2201.06503*, 2022b.

Yuemin Bian, Junmei Wang, Jaden Jungho Jun, and Xiang-Qun Xie. Deep convolutional generative adversarial network (dcgan) models for screening and design of small molecules targeting cannabinoid receptors. *Molecular pharmaceutics*, 16(11):4451–4460, 2019.

Nanxin Chen, Yu Zhang, Heiga Zen, Ron J Weiss, Mohammad Norouzi, and William Chan. Wavegrad: Estimating gradients for waveform generation. *arXiv preprint arXiv:2009.00713*, 2020.

Hanjun Dai, Yingtao Tian, Bo Dai, Steven Skiena, and Le Song. Syntax-directed variational autoencoder for structured data. *arXiv preprint arXiv:1802.08786*, 2018.

Prafulla Dhariwal and Alex Nichol. Diffusion models beat gans on image synthesis. *arXiv preprint arXiv:2105.05233*, 2021.

Tim Dockhorn, Arash Vahdat, and Karsten Kreis. Score-based generative modeling with critically-damped langevin diffusion. *arXiv preprint arXiv:2112.07068*, 2021.

Daniel Flam-Shepherd, Kevin Zhu, and Alán Aspuru-Guzik. Keeping it simple: Language models can learn complex molecular distributions. *arXiv preprint arXiv:2112.03041*, 2021.

Niklas Gebauer, Michael Gastegger, and Kristof Schütt. Symmetry-adapted generation of 3d point sets for the targeted discovery of molecules. *Advances in neural information processing systems*, 32, 2019.

Niklas WA Gebauer, Michael Gastegger, Stefaan SP Hessmann, Klaus-Robert Müller, and Kristof T Schütt. Inverse design of 3d molecular structures with conditional generative neural networks. *Nature communications*, 13(1):1–11, 2022.

Philip J Hajduk and Jonathan Greer. A decade of fragment-based drug design: strategic advances and lessons learned. *Nature reviews Drug discovery*, 6(3):211–219, 2007.

Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. *arXiv preprint arXiv:2006.11239*, 2020.

Moritz Hoffmann and Frank Noé. Generating valid euclidean distance matrices. *arXiv preprint arXiv:1910.03131*, 2019.

Emiel Hoogeboom, Victor Garcia Satorras, Clément Vignac, and Max Welling. Equivariant diffusion for molecule generation in 3d. In *International Conference on Machine Learning*, pp. 8867–8887. PMLR, 2022.

Jan H Jensen. A graph-based genetic algorithm and generative model/monte carlo tree search for the exploration of chemical space. *Chemical science*, 10(12):3567–3572, 2019.

Wengong Jin, Regina Barzilay, and Tommi Jaakkola. Junction tree variational autoencoder for molecular graph generation. In *International conference on machine learning*, pp. 2323–2332. PMLR, 2018.

Kisuk Kang, Ying Shirley Meng, Julien Breger, Clare P Grey, and Gerbrand Ceder. Electrodes with high power and high capacity for rechargeable lithium batteries. *Science*, 311(5763):977–980, 2006.

Jonas Köhler, Leon Klein, and Frank Noé. Equivariant flows: exact likelihood generative learning for symmetric densities. In *International conference on machine learning*, pp. 5361–5370. PMLR, 2020.

Zhifeng Kong, Wei Ping, Jiaji Huang, Kexin Zhao, and Bryan Catanzaro. Diffwave: A versatile diffusion model for audio synthesis. *arXiv preprint arXiv:2009.09761*, 2020.

Matt J Kusner, Brooks Paige, and José Miguel Hernández-Lobato. Grammar variational autoencoder. In *International conference on machine learning*, pp. 1945–1954. PMLR, 2017.

Qi Liu, Miltiadis Allamanis, Marc Brockschmidt, and Alexander Gaunt. Constrained graph variational autoencoders for molecule design. *Advances in neural information processing systems*, 31, 2018.

Cheng Lu, Yuhao Zhou, Fan Bao, Jianfei Chen, Chongxuan Li, and Jun Zhu. Dpm-solver: A fast ode solver for diffusion probabilistic model sampling in around 10 steps. *arXiv preprint arXiv:2206.00927*, 2022.

Youzhi Luo, Keqiang Yan, and Shuiwang Ji. Graphdf: A discrete flow model for molecular graph generation. In *International Conference on Machine Learning*, pp. 7192–7203. PMLR, 2021.

Kaushalya Madhawa, Katushiko Ishiguro, Kosuke Nakago, and Motoki Abe. Graphnvp: An invertible flow model for generating molecular graphs. *arXiv preprint arXiv:1905.11600*, 2019.

Soma Mandal, Sanat K Mandal, et al. Rational drug design. *European journal of pharmacology*, 625(1-3):90–100, 2009.

Vitali Nesterov, Mario Wieser, and Volker Roth. 3dmolnet: a generative network for molecular structures. *arXiv preprint arXiv:2010.06477*, 2020.

Alex Nichol and Prafulla Dhariwal. Improved denoising diffusion probabilistic models. *arXiv preprint arXiv:2102.09672*, 2021.

AkshatKumar Nigam, Pascal Friederich, Mario Krenn, and Alán Aspuru-Guzik. Augmenting genetic algorithms with deep neural networks for exploring the chemical space. *arXiv preprint arXiv:1909.11655*, 2019.

Mariya Popova, Mykhailo Shvets, Junier Oliva, and Olexandr Isayev. Molecularrnn: Generating realistic molecular graphs with optimized properties. *arXiv preprint arXiv:1905.13372*, 2019.

Edward O Pyzer-Knapp, Changwon Suh, Rafael Gómez-Bombarelli, Jorge Aguilera-Iparraguirre, and Alán Aspuru-Guzik. What is high-throughput virtual screening? a perspective from organic materials discovery. *Annu. Rev. Mater. Res.*, 2015.

Alec Radford, Jong Wook Kim, Chris Hallacy, Aditya Ramesh, Gabriel Goh, Sandhini Agarwal, Girish Sastry, Amanda Askell, Pamela Mishkin, Jack Clark, et al. Learning transferable visual models from natural language supervision. In *International Conference on Machine Learning*, pp. 8748–8763. PMLR, 2021.

Raghunathan Ramakrishnan, Pavlo O Dral, Matthias Rupp, and O Anatole Von Lilienfeld. Quantum chemistry structures and properties of 134 kilo molecules. *Scientific data*, 1(1):1–7, 2014.

Aditya Ramesh, Prafulla Dhariwal, Alex Nichol, Casey Chu, and Mark Chen. Hierarchical text-conditional image generation with clip latents. *arXiv preprint arXiv:2204.06125*, 2022.

Tim Salimans and Jonathan Ho. Progressive distillation for fast sampling of diffusion models. *arXiv preprint arXiv:2202.00512*, 2022.

Benjamin Sanchez-Lengeling and Alán Aspuru-Guzik. Inverse molecular design using machine learning: Generative models for matter engineering. *Science*, 361(6400):360–365, 2018.

Victor Garcia Satorras, Emiel Hoogeboom, Fabian B Fuchs, Ingmar Posner, and Max Welling. E (n) equivariant normalizing flows. *arXiv preprint arXiv:2105.09016*, 2021a.

Vıctor Garcia Satorras, Emiel Hoogeboom, and Max Welling. E (n) equivariant graph neural networks. In *International conference on machine learning*, pp. 9323–9332. PMLR, 2021b.

Martin Simonovsky and Nikos Komodakis. Graphvae: Towards generation of small graphs using variational autoencoders. In *International conference on artificial neural networks*, pp. 412–422. Springer, 2018.

Jascha Sohl-Dickstein, Eric Weiss, Niru Maheswaranathan, and Surya Ganguli. Deep unsupervised learning using nonequilibrium thermodynamics. In *International Conference on Machine Learning*, pp. 2256–2265. PMLR, 2015.

Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. *arXiv preprint arXiv:2011.13456*, 2020.

Arash Vahdat, Karsten Kreis, and Jan Kautz. Score-based generative modeling in latent space. *arXiv preprint arXiv:2106.05931*, 2021.

David Weininger. Smiles, a chemical language and information system. 1. introduction to methodology and encoding rules. *Journal of chemical information and computer sciences*, 28(1):31–36, 1988.

Minkai Xu, Lantao Yu, Yang Song, Chence Shi, Stefano Ermon, and Jian Tang. Geodiff: A geometric diffusion model for molecular conformation generation. *arXiv preprint arXiv:2203.02923*, 2022.

Jiaxuan You, Bowen Liu, Zhitao Ying, Vijay Pande, and Jure Leskovec. Graph convolutional policy network for goal-directed molecular graph generation. *Advances in neural information processing systems*, 31, 2018.

Chengxi Zang and Fei Wang. Moflow: an invertible flow model for generating molecular graphs. In *Proceedings of the 26th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*, pp. 617–626, 2020.

Min Zhao, Fan Bao, Chongxuan Li, and Jun Zhu. Egsde: Unpaired image-to-image translation via energy-guided stochastic differential equations. *arXiv preprint arXiv:2207.06635*, 2022.

Zhenpeng Zhou, Steven Kearnes, Li Li, Richard N Zare, and Patrick Riley. Optimization of molecules via deep reinforcement learning. *Scientific reports*, 9(1):1–10, 2019.

Alex Zunger. Inverse design in search of materials with target functionalities. *Nature Reviews Chemistry*, 2(4):1–16, 2018.

A   DERIVATIONS

A.1   ZERO CoM SUBSPACE

The zero CoM subspace $X = \{\boldsymbol{x} \in \mathbb{R}^{Mn} : \sum_{i=1}^{M} \boldsymbol{x}^i = \boldsymbol{0}\}$ is a $(M-1)n$ dimensional subspace of $\mathbb{R}^{Mn}$. Therefore, there exists an isometric isomorphism ϕ from $\mathbb{R}^{(M-1)n}$ to X, i.e., ϕ is a linear bijection from $\mathbb{R}^{(M-1)n}$ to X, and $\|\phi(\hat{\boldsymbol{x}})\|_2 = \|\hat{\boldsymbol{x}}\|_2$ for all $\hat{\boldsymbol{x}} \in \mathbb{R}^{(M-1)n}$. We use $A_\phi \in \mathbb{R}^{Mn \times (M-1)n}$ represent the matrix corresponding to ϕ, so we have $\phi(\hat{\boldsymbol{x}}) = A_\phi \hat{\boldsymbol{x}}$. An important property of the isometric isomorphism is that $A_\phi A_\phi^\top \boldsymbol{x} = \boldsymbol{x} - \overline{\boldsymbol{x}}$ for all $\boldsymbol{x} \in \mathbb{R}^{Mn}$. We show the proof as following.

Proposition 1. *Suppose ϕ is an isometric isomorphism from $\mathbb{R}^{(M-1)n}$ to X, and let $A_\phi \in \mathbb{R}^{Mn \times (M-1)n}$ be the matrix corresponding to ϕ. Then we have $A_\phi A_\phi^\top \boldsymbol{x} = \boldsymbol{x} - \overline{\boldsymbol{x}}$ for all $\boldsymbol{x} \in \mathbb{R}^{Mn}$, where $\overline{\boldsymbol{x}} = \frac{1}{M} \sum_{i=1}^{M} \boldsymbol{x}^i$.*

Proof. We consider a new subspace of $\mathbb{R}^{Mn}$, $X^\perp = \{\boldsymbol{x} \in \mathbb{R}^{Mn} : \boldsymbol{x} \perp X\}$, i.e., the orthogonal component of X. We can verify that $X^\perp = \{\boldsymbol{x} \in \mathbb{R}^{Mn} : \boldsymbol{x}_1 = \boldsymbol{x}_2 = \cdots = \boldsymbol{x}_M\}$, and $X^\perp$ is n dimensional. Thus, there exists an isometric isomorphism ψ from $\mathbb{R}^n$ to $X^\perp$. Let $A_\psi \in \mathbb{R}^{Mn \times n}$ represent the matrix corresponding to ψ. Then we define $\lambda(\hat{\boldsymbol{x}}, \hat{\boldsymbol{y}}) = \phi(\hat{\boldsymbol{x}}) + \psi(\hat{\boldsymbol{y}})$, where $\hat{\boldsymbol{x}} \in \mathbb{R}^{(M-1)n}$ and $\hat{\boldsymbol{y}} \in \mathbb{R}^n$. The image of λ is $\{\boldsymbol{x} + \boldsymbol{y} : \boldsymbol{x} \in X, \boldsymbol{y} \in X^\perp\} = \mathbb{R}^{Mn}$. Therefore, λ is a linear bijection from $\mathbb{R}^{Mn}$ to $\mathbb{R}^{Mn}$, and the matrix corresponding to λ is $A_\lambda = [A_\phi, A_\psi]$. Furthermore, λ is an isometric isomorphism, since $\|\lambda(\hat{\boldsymbol{x}}, \hat{\boldsymbol{y}})\|^2 = \|\phi(\hat{\boldsymbol{x}})\|^2 + \|\psi(\hat{\boldsymbol{y}})\|^2 + 2\langle \phi(\hat{\boldsymbol{x}}), \psi(\hat{\boldsymbol{y}}) \rangle = \|\hat{\boldsymbol{x}}\|^2 + \|\hat{\boldsymbol{y}}\|^2 = \|(\hat{\boldsymbol{x}}^\top, \hat{\boldsymbol{y}}^\top)^\top\|^2$. This means A_λ is orthogonal transformation. Therefore, $A_\lambda A_\lambda^\top = A_\phi A_\phi^\top \boldsymbol{x} + A_\psi A_\psi^\top = \boldsymbol{I}$ and $A_\phi A_\phi^\top \boldsymbol{x} + A_\psi A_\psi^\top \boldsymbol{x} = \boldsymbol{x}$. Since $A_\phi A_\phi^\top \boldsymbol{x} \in X$ and $A_\psi A_\psi^\top \boldsymbol{x} \in X^\top$, we can conclude that $A_\phi A_\phi^\top \boldsymbol{x}$ is the orthogonal projection of $\boldsymbol{x}$ to X, which is exactly $\boldsymbol{x} - \overline{\boldsymbol{x}}$. $\qquad\square$

Since $\mathbb{R}^{(M-1)n}$ and X are two intrinsically equivalent spaces, an equivalence of distributions in these two spaces can also be established, as shown in the following propositions.

Proposition 2. *Suppose $\boldsymbol{x}$ is a random vector distributed in X, and $\hat{\boldsymbol{x}} = \phi^{-1}(\boldsymbol{x})$ is its equivalent representation in $\mathbb{R}^{(M-1)n}$. If $\boldsymbol{x} \sim q(\boldsymbol{x})$, then $\hat{\boldsymbol{x}} \sim \hat{q}(\hat{\boldsymbol{x}})$, where $\hat{q}(\hat{\boldsymbol{x}}) = q(\phi(\hat{\boldsymbol{x}}))$.*

Proposition 3. *Suppose $\boldsymbol{x}, \boldsymbol{y}$ is are two random vectors distributed in X, and $\hat{\boldsymbol{x}} = \phi^{-1}(\boldsymbol{x}), \hat{\boldsymbol{y}} = \phi^{-1}(\boldsymbol{y})$ are their equivalent representations in $\mathbb{R}^{(M-1)n}$. If $\boldsymbol{x}|\boldsymbol{y} \sim q(\boldsymbol{x}|\boldsymbol{y})$, then $\hat{\boldsymbol{x}}|\hat{\boldsymbol{y}} \sim \hat{q}(\hat{\boldsymbol{x}}|\hat{\boldsymbol{y}})$, where $\hat{q}(\hat{\boldsymbol{x}}|\hat{\boldsymbol{y}}) = q(\phi(\hat{\boldsymbol{x}})|\phi(\hat{\boldsymbol{y}}))$.*

A.2   SDE IN THE PRODUCT SPACE

Definition 1 (Gaussian distributions in the zero CoM subspace). *Suppose $\boldsymbol{\mu} \in X$. Let $\mathcal{N}_X(\boldsymbol{x}|\boldsymbol{\mu}, \sigma^2) := (2\pi\sigma^2)^{-(M-1)n/2} \exp(-\frac{1}{2\sigma^2} \|\boldsymbol{x} - \boldsymbol{\mu}\|_2^2)$, which is the isotropic Gaussian distribution with mean $\boldsymbol{\mu}$ and variance σ^2 in the zero CoM subspace X.*

Proposition 4 (Transition kernels of a SDE in the zero CoM subspace). *Suppose $d\boldsymbol{x} = f(t)\boldsymbol{x}dt + g(t)d\boldsymbol{w}_x$, $\boldsymbol{x}_0 \sim q(\boldsymbol{x}_0)$ is a SDE in the zero CoM subspace. Then the transition kernel from $\boldsymbol{x}_s$ to $\boldsymbol{x}_t$ $(0 \leq s < t \leq T)$ can be expressed as $q(\boldsymbol{x}_t|\boldsymbol{x}_s) = \mathcal{N}_X(\boldsymbol{x}_t|\sqrt{\alpha_{t|s}}\boldsymbol{x}_s, \beta_{t|s})$, where $\alpha_{t|s} = \exp(2\int_s^t f(\tau)d\tau)$ and $\beta_{t|s} = \alpha_{t|s} \int_s^t g(\tau)^2/\alpha_{\tau|s}d\tau$ are two scalars determined by $f(\cdot)$ and $g(\cdot)$.*

Proof. Firstly, we map the process $\{\boldsymbol{x}_t\}_{t=0}^T$ in the zero CoM subspace to the equivalent space $\mathbb{R}^{(M-1)n}$ through the isometric isomorphism ϕ introduced in Appendix A.1. This produces a new process $\{\hat{\boldsymbol{x}}_t\}_{t=0}^T$, where $\hat{\boldsymbol{x}}_t = \phi^{-1}(\boldsymbol{x}_t)$. By applying ϕ^{-1} to the SDE in the zero CoM subspace, we know $\hat{\boldsymbol{x}}_t = \phi^{-1}(\boldsymbol{x}_t)$ satisfies the following SDE in $\mathbb{R}^{(M-1)n}$:

$$d\hat{\boldsymbol{x}} = f(t)\hat{\boldsymbol{x}}dt + g(t)d\hat{\boldsymbol{w}}, \quad \hat{\boldsymbol{x}}_0 \sim \hat{q}(\hat{\boldsymbol{x}}_0), \qquad (9)$$

where $\hat{\boldsymbol{w}}$ is the standard Wiener process in $\mathbb{R}^{(M-1)n}$ and $\hat{q}(\hat{\boldsymbol{x}}_0) = q(\phi(\hat{\boldsymbol{x}}_0))$. According to Song et al. (2020), the transition of Eq. (9) from $\hat{\boldsymbol{x}}_s$ to $\hat{\boldsymbol{x}}_t$ $(s < t)$ can be expressed as $\hat{q}(\hat{\boldsymbol{x}}_t|\hat{\boldsymbol{x}}_s) = \mathcal{N}(\hat{\boldsymbol{x}}_t|\sqrt{\alpha_{t|s}}\hat{\boldsymbol{x}}_s, \beta_{t|s}\boldsymbol{I})$, where $\alpha_{t|s} = \exp(2\int_s^t f(\tau)d\tau)$ and $\beta_{t|s} = \alpha_{t|s} \int_s^t g(\tau)^2/\alpha_{\tau|s}d\tau$ are two

scalars determined by $f(\cdot)$ and $g(\cdot)$. According to Proposition 3, we know the transition kernel $q(\boldsymbol{x}_t|\boldsymbol{x}_s)$ from $\boldsymbol{x}_s$ to $\boldsymbol{x}_t$ satisfies

$$
\begin{aligned}
q(\boldsymbol{x}_t|\boldsymbol{x}_s) &= \hat{q}(\phi^{-1}(\boldsymbol{x}_t)|\phi^{-1}(\boldsymbol{x}_s)) = (2\pi\beta_{t|s})^{-(M-1)n/2}\exp(-\frac{1}{2\beta_{t|s}}\|\phi^{-1}(\boldsymbol{x}_t) - \sqrt{\alpha_{t|s}}\phi^{-1}(\boldsymbol{x}_s)\|_2^2) \\
&= (2\pi\beta_{t|s})^{-(M-1)n/2}\exp(-\frac{1}{2\beta_{t|s}}\|\phi^{-1}(\boldsymbol{x}_t - \sqrt{\alpha_{t|s}}\boldsymbol{x}_s)\|_2^2) \qquad \text{// linearity of } \phi^{-1} \\
&= (2\pi\beta_{t|s})^{-(M-1)n/2}\exp(-\frac{1}{2\beta_{t|s}}\|\boldsymbol{x}_t - \sqrt{\alpha_{t|s}}\boldsymbol{x}_s\|_2^2) \qquad \text{// norm preserving of } \phi^{-1} \\
&= \mathcal{N}_X(\boldsymbol{x}_t|\sqrt{\alpha_{t|s}}\boldsymbol{x}_s, \beta_{t|s}).
\end{aligned}
$$

$\square$

Proposition 5 (Transition kernels of the SDE in the product space). *Suppose* $\mathrm{d}\boldsymbol{z} = f(t)\boldsymbol{z}\mathrm{d}t + g(t)\mathrm{d}(\boldsymbol{w}_x, \boldsymbol{w}_h), \boldsymbol{z}_0 \sim q(\boldsymbol{z}_0)$ *is the SDE in the product space* $X \times \mathbb{R}^{Md}$, *as introduced in Eq. (3). Then the transition kernel from* $\boldsymbol{z}_s$ *to* $\boldsymbol{z}_t$ $(0 \leq s < t \leq T)$ *can be expressed as* $q(\boldsymbol{z}_t|\boldsymbol{z}_s) = q(\boldsymbol{x}_t|\boldsymbol{x}_s)q(\boldsymbol{h}_t|\boldsymbol{h}_s)$, *where* $q(\boldsymbol{x}_t|\boldsymbol{x}_s) = \mathcal{N}_X(\boldsymbol{x}_t|\sqrt{\alpha_{t|s}}\boldsymbol{x}_s, \beta_{t|s})$ *and* $q(\boldsymbol{h}_t|\boldsymbol{h}_s) = \mathcal{N}(\boldsymbol{h}_t|\sqrt{\alpha_{t|s}}\boldsymbol{h}_s, \beta_{t|s})$. *Here* $\alpha_{t|s}$ *and* $\beta_{t|s}$ *are defined as in Proposition 4.*

Proof. Since $\boldsymbol{w}_x$ and $\boldsymbol{w}_h$ are independent to each other, the transition kernel from $\boldsymbol{z}_s$ to $\boldsymbol{z}_t$ can be factorized as $q(\boldsymbol{z}_t|\boldsymbol{z}_s) = q(\boldsymbol{x}_t|\boldsymbol{x}_s)q(\boldsymbol{h}_t|\boldsymbol{h}_s)$, where $q(\boldsymbol{x}_t|\boldsymbol{x}_s)$ is the transition kernel of $\mathrm{d}\boldsymbol{x} = f(t)\boldsymbol{x}\mathrm{d}t + g(t)\mathrm{d}\boldsymbol{w}_x$ and $q(\boldsymbol{h}_t|\boldsymbol{h}_s)$ is the transition kernel of $\mathrm{d}\boldsymbol{h} = f(t)\boldsymbol{h}\mathrm{d}t + g(t)\mathrm{d}\boldsymbol{w}_h$. According to Proposition 4 and Song et al. (2020), we have $q(\boldsymbol{x}_t|\boldsymbol{x}_s) = \mathcal{N}_X(\boldsymbol{x}_t|\sqrt{\alpha_{t|s}}\boldsymbol{x}_s, \beta_{t|s})$ and $q(\boldsymbol{h}_t|\boldsymbol{h}_s) = \mathcal{N}(\boldsymbol{h}_t|\sqrt{\alpha_{t|s}}\boldsymbol{h}_s, \beta_{t|s})$. $\square$

Remark 1. *The marginal distribution of* $\boldsymbol{z}_t$ *is*

$$
q_t(\boldsymbol{z}_t) = \int_Z \int_{\mathbb{R}^{Md}} q(\boldsymbol{z}_0)q(\boldsymbol{x}_t|\boldsymbol{x}_0)q(\boldsymbol{h}_t|\boldsymbol{h}_0)\mathrm{d}\boldsymbol{h}_0\lambda(\mathrm{d}\boldsymbol{x}_0),
$$

where λ *is the Lebesgue measure in the zero CoM subspace* X. *While* $q_t(\boldsymbol{z}_t)$ *is a distribution in* $X \times \mathbb{R}^{Md}$, *it has a natural differentiable extension to* $\mathbb{R}^{Mn} \times \mathbb{R}^{Md}$, *since* $q(\boldsymbol{x}_t|\boldsymbol{x}_0) = \mathcal{N}_X(\boldsymbol{x}_t|\sqrt{\alpha_{t|s}}\boldsymbol{x}_s, \beta_{t|s})$ *has a differentiable extension to* $\mathbb{R}^{Mn}$ *according to Definition 1. Thus, we can take gradient of* $q_t(\boldsymbol{z}_t)$ *w.r.t.* $\boldsymbol{x}_t$ *in the whole space* $\mathbb{R}^{Mn}$.

Proposition 6 (Time reversal of the SDE in the product space). *Suppose* $\mathrm{d}\boldsymbol{z} = f(t)\boldsymbol{z}\mathrm{d}t + g(t)\mathrm{d}(\boldsymbol{w}_x, \boldsymbol{w}_h), \boldsymbol{z}_0 \sim q(\boldsymbol{z}_0)$ *is the SDE in the product space* $X \times \mathbb{R}^{Md}$, *as introduced in Eq. (3). Then its time reversal satisfies the following reverse-time SDE, which can be represented by both the score function form and the noise prediction form:*

$$
\mathrm{d}\boldsymbol{z} = [f(t)\boldsymbol{z} - g(t)^2 \underbrace{(\nabla_{\boldsymbol{x}}\log q_t(\boldsymbol{z}) - \overline{\nabla_{\boldsymbol{x}}\log q_t(\boldsymbol{z})}, \nabla_{\boldsymbol{h}}\log q_t(\boldsymbol{z}))}_{\text{score function form}}]\mathrm{d}t + g(t)\mathrm{d}(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h),
$$

$$
= [f(t)\boldsymbol{z} + \frac{g(t)^2}{\sqrt{\beta_{t|0}}} \underbrace{\mathbb{E}_{q(\boldsymbol{z}_0|\boldsymbol{z}_t)}\boldsymbol{\epsilon}_t}_{\text{noise prediction form}}]\mathrm{d}t + g(t)\mathrm{d}(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h), \quad \boldsymbol{z}_T \sim q_T(\boldsymbol{z}_T),
$$

where $\tilde{\boldsymbol{w}}_x$ *and* $\tilde{\boldsymbol{w}}_h$ *are reverse-time standard Wiener processes in* X *and* $\mathbb{R}^{Md}$ *respectively, and* $\boldsymbol{\epsilon}_t = \frac{\boldsymbol{z}_t - \sqrt{\alpha_{t|0}}\boldsymbol{z}_0}{\sqrt{\beta_{t|0}}}$ *is the standard Gaussian noise in* $X \times \mathbb{R}^{Md}$ *injected to* $\boldsymbol{z}_0$.

Furthermore, we have $\nabla_{\boldsymbol{x}}\log q_t(\boldsymbol{x}) - \overline{\nabla_{\boldsymbol{x}}\log q_t(\boldsymbol{x})} = -\frac{1}{\sqrt{\beta_{t|0}}}\mathbb{E}_{q(\boldsymbol{x}_0|\boldsymbol{x}_t)}\boldsymbol{\epsilon}_t$, *where* $\boldsymbol{\epsilon}_t = \frac{\boldsymbol{x}_t - \sqrt{\alpha_{t|0}}\boldsymbol{x}_0}{\sqrt{\beta_{t|0}}}$ *is the standard Gaussian noise in the zero CoM subspace injected to* $\boldsymbol{x}_0$.

Proof. Let $\hat{\boldsymbol{z}}_t = (\hat{\boldsymbol{x}}_t, \boldsymbol{h}_t)$, where $\hat{\boldsymbol{x}}_t = \phi^{-1}(\boldsymbol{x}_t)$, introduced in the proof of Proposition 4. Then $\{\hat{\boldsymbol{z}}_t\}_{t=0}^T$ is a process in $\mathbb{R}^{(M-1)n} \times \mathbb{R}^{Md}$ determined by the following SDE

$$
\mathrm{d}\hat{\boldsymbol{z}} = f(t)\hat{\boldsymbol{z}}\mathrm{d}t + g(t)\mathrm{d}\hat{\boldsymbol{w}}, \quad \hat{\boldsymbol{z}}_0 \sim \hat{q}(\hat{\boldsymbol{z}}_0), \tag{10}
$$

where $\hat{\boldsymbol{w}}$ is the standard Wiener process in $\mathbb{R}^{(M-1)n} \times \mathbb{R}^{Md}$ and $\hat{q}(\hat{\boldsymbol{z}}_0) = q(\phi(\hat{\boldsymbol{x}}_0), \boldsymbol{h})$.

According to Song et al. (2020), Eq. (10) has a time reversal:

$$d\hat{\boldsymbol{z}} = [f(t)\hat{\boldsymbol{z}} - g(t)^2 \nabla_{\hat{\boldsymbol{z}}} \log \hat{q}_t(\hat{\boldsymbol{z}})]dt + g(t)d\tilde{\boldsymbol{w}}, \quad \hat{\boldsymbol{z}}_T \sim \hat{q}_T(\hat{\boldsymbol{z}}_T), \tag{11}$$

where $\hat{q}_t(\boldsymbol{z})$ is the marginal distribution of $\hat{\boldsymbol{z}}_t$, which satisfies $\hat{q}_t(\hat{\boldsymbol{z}}_t) = q_t(\phi(\hat{\boldsymbol{x}}_t), \boldsymbol{h}_t)$ according to Proposition 2, and $\tilde{\boldsymbol{w}}$ is the reverse-time standard Wiener process in $\mathbb{R}^{(M-1)n} \times \mathbb{R}^{Md}$.

Then we apply the linear transformation $T\hat{\boldsymbol{z}} = (\phi(\hat{\boldsymbol{x}}), \boldsymbol{h})$ to Eq. (11), which maps $\hat{\boldsymbol{z}}_t$ back to $\boldsymbol{z}_t$. This yields

$$d\boldsymbol{z} = [f(t)\boldsymbol{z} - g(t)^2 T(\nabla_{\hat{\boldsymbol{z}}} \log \hat{q}_t(\hat{\boldsymbol{z}}))]dt + g(t)d(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h), \quad \boldsymbol{z}_T \sim q_T(\boldsymbol{z}_T). \tag{12}$$

Here $\tilde{\boldsymbol{w}}_x$ and $\tilde{\boldsymbol{w}}_h$ are reverse-time standard Wiener processes in X and $\mathbb{R}^{Md}$ respectively, and $T(\nabla_{\hat{\boldsymbol{z}}} \log \hat{q}_t(\hat{\boldsymbol{z}}))$ can be expressed as $(\phi(\nabla_{\hat{\boldsymbol{x}}} \log \hat{q}_t(\hat{\boldsymbol{z}})), \nabla_{\boldsymbol{h}} \log \hat{q}_t(\hat{\boldsymbol{z}}))$.

Since $\nabla_{\hat{\boldsymbol{x}}} \log \hat{q}_t(\hat{\boldsymbol{z}}) = A_\phi^\top \nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{x}, \boldsymbol{h}) = A_\phi^\top \nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{z})$, we have $\phi(\nabla_{\hat{\boldsymbol{x}}} \log \hat{q}_t(\hat{\boldsymbol{z}})) = A_\phi A_\phi^\top \nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{z})$, where A_ϕ represents the matrix corresponding to ϕ. According to Proposition 1, we have $\phi(\nabla_{\hat{\boldsymbol{x}}} \log \hat{q}_t(\hat{\boldsymbol{z}})) = \nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{z}) - \overline{\nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{z})}$. Besides, $\nabla_{\boldsymbol{h}} \log \hat{q}_t(\hat{\boldsymbol{z}}) = \nabla_{\boldsymbol{h}} \log q_t(\boldsymbol{z})$. Thus, Eq. (12) can be written as

$$d\boldsymbol{z} = [f(t)\boldsymbol{z} - g(t)^2(\nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{z}) - \overline{\nabla_{\boldsymbol{x}} \log q_t(\boldsymbol{z})}, \nabla_{\boldsymbol{h}} \log q_t(\boldsymbol{z}))]dt + g(t)d(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h),$$

which is the score function form of the reverse-time SDE.

We can also write the score function in Eq. (11) as $\nabla_{\hat{\boldsymbol{z}}} \log \hat{q}_t(\hat{\boldsymbol{z}}) = \mathbb{E}_{\hat{q}(\hat{\boldsymbol{z}}_0|\hat{\boldsymbol{z}}_t)} \nabla_{\hat{\boldsymbol{z}}} \log \hat{q}(\hat{\boldsymbol{z}}_t|\hat{\boldsymbol{z}}_0) = -\frac{1}{\sqrt{\beta_{t|0}}} \mathbb{E}_{\hat{q}(\hat{\boldsymbol{z}}_0|\hat{\boldsymbol{z}}_t)} \hat{\boldsymbol{\epsilon}}_t$, where $\hat{\boldsymbol{\epsilon}}_t = \frac{\hat{\boldsymbol{z}}_t - \sqrt{\alpha_{t|0}} \hat{\boldsymbol{z}}_0}{\sqrt{\beta_{t|0}}}$ is the standard Gaussian noise injected to $\hat{\boldsymbol{z}}_0$. With this expression, we have $T(\nabla_{\hat{\boldsymbol{z}}} \log \hat{q}_t(\hat{\boldsymbol{z}})) = -\frac{1}{\sqrt{\beta_{t|0}}} \mathbb{E}_{\hat{q}(\hat{\boldsymbol{z}}_0|\hat{\boldsymbol{z}}_t)} T(\hat{\boldsymbol{\epsilon}}_t) = -\frac{1}{\sqrt{\beta_{t|0}}} \mathbb{E}_{q(\boldsymbol{z}_0|\boldsymbol{z}_t)} \boldsymbol{\epsilon}_t$, where $\boldsymbol{\epsilon}_t = \frac{\boldsymbol{z}_t - \sqrt{\alpha_{t|0}} \boldsymbol{z}_0}{\sqrt{\beta_{t|0}}}$ is the standard Gaussian noise in $X \times \mathbb{R}^{Md}$ injected to $\boldsymbol{z}_0$. Thus, Eq. (12) can also be written as

$$d\boldsymbol{z} = [f(t)\boldsymbol{z} + \frac{g(t)^2}{\sqrt{\beta_{t|0}}} \mathbb{E}_{q(\boldsymbol{z}_0|\boldsymbol{z}_t)} \boldsymbol{\epsilon}_t]dt + g(t)d(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h),$$

which is the noise prediction form of the reverse-time SDE.

$\square$

A.3 EQUIVARIANCE

Theorem 1. *Let* $(\boldsymbol{\epsilon}_{\boldsymbol{\theta}}^x(\boldsymbol{z}_t, t), \boldsymbol{\epsilon}_{\boldsymbol{\theta}}^h(\boldsymbol{z}_t, t)) = \boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}_t, t)$*, where* $\boldsymbol{\epsilon}_{\boldsymbol{\theta}}^x(\boldsymbol{z}_t, t)$ *and* $\boldsymbol{\epsilon}_{\boldsymbol{\theta}}^h(\boldsymbol{z}_t, t)$ *are the predicted noise of* $\boldsymbol{x}_t$ *and* $\boldsymbol{h}_t$ *respectively. If for any orthogonal transformation* $\boldsymbol{R} \in \mathbb{R}^{n \times n}$*,* $\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}_t, t)$ *is equivariant to* $\boldsymbol{R}$*, i.e.,* $\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{R}\boldsymbol{x}_t, \boldsymbol{h}_t, t) = (\boldsymbol{R}\boldsymbol{\epsilon}_{\boldsymbol{\theta}}^x(\boldsymbol{x}_t, \boldsymbol{h}_t, t), \boldsymbol{\epsilon}_{\boldsymbol{\theta}}^h(\boldsymbol{x}_t, \boldsymbol{h}_t, t))$*, and* $p_T(\boldsymbol{z}_T)$ *is invariant to* $\boldsymbol{R}$*, i.e.,* $p_T(\boldsymbol{R}\boldsymbol{x}_T, \boldsymbol{h}_T) = p_T(\boldsymbol{x}_T, \boldsymbol{h}_T)$*, then* $p_{\boldsymbol{\theta}}(\boldsymbol{z}_0)$ *is invariant to any rotational transformation.*

Proof. Suppose $\boldsymbol{R} \in \mathbb{R}^{n \times n}$ is an orthogonal transformation. Let $\boldsymbol{z}_t^{\boldsymbol{\theta}} = (\boldsymbol{x}_t^{\boldsymbol{\theta}}, \boldsymbol{h}_t^{\boldsymbol{\theta}})$ $(0 \leq t \leq T)$ be the process determined by Eq. (5). Let $\boldsymbol{y}_t^{\boldsymbol{\theta}} = \boldsymbol{R}\boldsymbol{x}_t^{\boldsymbol{\theta}}$ and $\boldsymbol{u}_t^{\boldsymbol{\theta}} = (\boldsymbol{y}_t^{\boldsymbol{\theta}}, \boldsymbol{h}_t^{\boldsymbol{\theta}})$. We use $p_t^{\boldsymbol{u}}(\boldsymbol{u}_t)$ and $p_t^{\boldsymbol{z}}(\boldsymbol{z}_t)$ to denote the distributions of $\boldsymbol{u}_t^{\boldsymbol{\theta}}$ and $\boldsymbol{z}_t^{\boldsymbol{\theta}}$ respectively, and they satisfy $p_t^{\boldsymbol{u}}(\boldsymbol{y}_t, \boldsymbol{h}_t) = p_t^{\boldsymbol{z}}(\boldsymbol{R}^{-1}\boldsymbol{y}_t, \boldsymbol{h}_t)$. By applying the transformation $T\boldsymbol{z} = (\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h})$ to Eq. (5), we know the new process $\{\boldsymbol{u}_t^{\boldsymbol{\theta}}\}_{t=0}^T$ satisfies the following SDE:

$$d\boldsymbol{u} = Td\boldsymbol{z} = [f(t)\boldsymbol{u} + \frac{g(t)^2}{\sqrt{\beta_{t|0}}} (\boldsymbol{R}\boldsymbol{\epsilon}_{\boldsymbol{\theta}}^x(\boldsymbol{z}, t), \boldsymbol{\epsilon}_{\boldsymbol{\theta}}^h(\boldsymbol{z}, t))]dt + g(t)d(\boldsymbol{R}\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h)$$

$$= [f(t)\boldsymbol{u} + \frac{g(t)^2}{\sqrt{\beta_{t|0}}} (\boldsymbol{\epsilon}_{\boldsymbol{\theta}}^x(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}, t), \boldsymbol{\epsilon}_{\boldsymbol{\theta}}^h(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}, t))]dt + g(t)d(\boldsymbol{R}\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h) \quad \text{// by equivariance}$$

$$= [f(t)\boldsymbol{u} + \frac{g(t)^2}{\sqrt{\beta_{t|0}}} \boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{u}, t)]dt + g(t)d(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h), \quad \text{// Winner process is isotropic}$$

and its initial distribution is $p_T^{\boldsymbol{u}}(\boldsymbol{u}_T) = p_T^{\boldsymbol{u}}(\boldsymbol{y}_T, \boldsymbol{h}_T) = p_T^{\boldsymbol{z}}(\boldsymbol{R}^{-1}\boldsymbol{y}_T, \boldsymbol{h}_T) = p_T(\boldsymbol{R}^{-1}\boldsymbol{y}_T, \boldsymbol{h}_T) = p_T(\boldsymbol{y}_T, \boldsymbol{h}_T) = p_T(\boldsymbol{u}_T)$. Thus, the SDE of $\{\boldsymbol{u}_t^{\boldsymbol{\theta}}\}_{t=0}^T$ is exactly the same to that of $\{\boldsymbol{z}_t^{\boldsymbol{\theta}}\}_{t=0}^T$.

This indicates that the distribution of $\boldsymbol{u}_0^{\boldsymbol{\theta}}$ is the same to the distribution of $\boldsymbol{z}_0^{\boldsymbol{\theta}}$, i.e., $p_0^u(\boldsymbol{u}_0) = p_0^z(\boldsymbol{u}_0) = p_{\boldsymbol{\theta}}(\boldsymbol{u}_0)$. Also note that $p_0^u(\boldsymbol{u}_0) = p_0^z(\boldsymbol{R}^{-1}\boldsymbol{y}_0, \boldsymbol{h}_0) = p_{\boldsymbol{\theta}}(\boldsymbol{R}^{-1}\boldsymbol{y}_0, \boldsymbol{h}_0)$. Thus, $p_{\boldsymbol{\theta}}(\boldsymbol{u}_0) = p_{\boldsymbol{\theta}}(\boldsymbol{R}^{-1}\boldsymbol{y}_0, \boldsymbol{h}_0)$, and consequently $p_{\boldsymbol{\theta}}(\boldsymbol{R}\boldsymbol{x}_0, \boldsymbol{h}_0) = p_{\boldsymbol{\theta}}(\boldsymbol{x}_0, \boldsymbol{h}_0)$. This means $p_{\boldsymbol{\theta}}(\boldsymbol{z}_0)$ is invariant to any orthogonal transformation, which includes rotational transformations as special cases. $\qquad\square$

Theorem 2. *Suppose the assumptions in Theorem 1 hold and $E(\boldsymbol{z}, c, t)$ is invariant to any orthogonal transformation $\boldsymbol{R}$, i.e., $E(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}, c, t) = E(\boldsymbol{x}, \boldsymbol{h}, c, t)$. Then $p_{\boldsymbol{\theta}}(\boldsymbol{z}_0|c)$ is invariant to any rotational transformation.*

Proof. Suppose $\boldsymbol{R} \in \mathbb{R}^{n \times n}$ is an orthogonal transformation. Taking gradient to both side of $E(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}, c, t) = E(\boldsymbol{x}, \boldsymbol{h}, c, t)$ w.r.t. $\boldsymbol{x}$, we get

$$\boldsymbol{R}^\top \nabla_{\boldsymbol{y}} E(\boldsymbol{y}, \boldsymbol{h}, c, t)|_{\boldsymbol{y}=\boldsymbol{R}\boldsymbol{x}} = \nabla_{\boldsymbol{x}} E(\boldsymbol{x}, \boldsymbol{h}, c, t).$$

Multiplying $\boldsymbol{R}$ to both sides, we get

$$\nabla_{\boldsymbol{y}} E(\boldsymbol{y}, \boldsymbol{h}, c, t)|_{\boldsymbol{y}=\boldsymbol{R}\boldsymbol{x}} = \boldsymbol{R}\nabla_{\boldsymbol{x}} E(\boldsymbol{x}, \boldsymbol{h}, c, t).$$

Let $\boldsymbol{\phi}(\boldsymbol{z}, c, t) = (\nabla_{\boldsymbol{x}} E(\boldsymbol{z}, c, t) - \overline{\nabla_{\boldsymbol{x}} E(\boldsymbol{z}, c, t)}, \nabla_{\boldsymbol{h}} E(\boldsymbol{z}, c, t))$. Then we have

$$\begin{aligned}
\boldsymbol{\phi}(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}, c, t) =& (\nabla_{\boldsymbol{y}} E(\boldsymbol{y}, \boldsymbol{h}, c, t) - \overline{\nabla_{\boldsymbol{y}} E(\boldsymbol{y}, \boldsymbol{h}, c, t)}, \nabla_{\boldsymbol{h}} E(\boldsymbol{y}, \boldsymbol{h}, c, t))|_{\boldsymbol{y}=\boldsymbol{R}\boldsymbol{x}} \\
=& (\boldsymbol{R}\nabla_{\boldsymbol{x}} E(\boldsymbol{x}, \boldsymbol{h}, c, t) - \boldsymbol{R}\overline{\nabla_{\boldsymbol{x}} E(\boldsymbol{x}, \boldsymbol{h}, c, t)}, \nabla_{\boldsymbol{h}} E(\boldsymbol{R}\boldsymbol{x}, \boldsymbol{h}, c, t)) \\
=& (\boldsymbol{R}(\nabla_{\boldsymbol{x}} E(\boldsymbol{x}, \boldsymbol{h}, c, t) - \overline{\nabla_{\boldsymbol{x}} E(\boldsymbol{x}, \boldsymbol{h}, c, t)}), \nabla_{\boldsymbol{h}} E(\boldsymbol{x}, \boldsymbol{h}, c, t)).
\end{aligned}$$

Thus, $\boldsymbol{\phi}(\boldsymbol{z}, c, t)$ is equivariant to $\boldsymbol{R}$. Let $\hat{\boldsymbol{\epsilon}}_{\boldsymbol{\theta}}(\boldsymbol{z}, c, t) = \boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}, t) + \sqrt{\beta_{t|0}}\boldsymbol{\phi}(\boldsymbol{z}, c, t)$, which is a linear combination of two equivariant functions and is also equivariant to $\boldsymbol{R}$.

Then, Eq. (6) can be written as

$$\mathrm{d}\boldsymbol{z} = [f(t)\boldsymbol{z} + \frac{g(t)^2}{\sqrt{\beta_{t|0}}}\hat{\boldsymbol{\epsilon}}_{\boldsymbol{\theta}}(\boldsymbol{z}, c, t)]\mathrm{d}t + g(t)\mathrm{d}(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h), \quad \boldsymbol{z}_T \sim p_T(\boldsymbol{z}_T).$$

According to Theorem 1, we know its marginal distribution at time $t = 0$, i.e., $p_{\boldsymbol{\theta}}(\boldsymbol{z}_0|c)$, is invariant to any rotational transformation.

$\qquad\square$

B CONDITIONAL NOISE PREDICTION NETWORKS

In Eq. (6), we can alternatively use a conditional noise prediction network $\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}, c, t)$ for a stronger guidance, as follows

$$\begin{aligned}
\mathrm{d}\boldsymbol{z} = [f(t)\boldsymbol{z} + g(t)^2(&\frac{1}{\sqrt{\beta_{t|0}}}\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}, c, t) \\
&+ (\nabla_{\boldsymbol{x}} E(\boldsymbol{z}, c, t) - \overline{\nabla_{\boldsymbol{x}} E(\boldsymbol{z}, c, t)}, \nabla_{\boldsymbol{h}} E(\boldsymbol{z}, c, t)))]\mathrm{d}t + g(t)\mathrm{d}(\tilde{\boldsymbol{w}}_x, \tilde{\boldsymbol{w}}_h), \ \boldsymbol{z}_T \sim p_T(\boldsymbol{z}_T).
\end{aligned}$$

The conditional noise prediction network is trained similarly to the unconditional one, using the following MSE loss

$$\min_{\boldsymbol{\theta}} \mathbb{E}_t \mathbb{E}_{q(c, \boldsymbol{z}_0, \boldsymbol{z}_t)} w(t)\|\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}_t, c, t) - \boldsymbol{\epsilon}_t\|^2.$$

C PARAMETERIZATION OF NOISE PREDICTION NETWORKS

We parameterize the noise prediction network following Hoogeboom et al. (2022), and we provide the specific parameterization for completeness. For the unconditional model $\boldsymbol{\epsilon}_{\boldsymbol{\theta}}(\boldsymbol{z}, t)$, we first concatenate each atom feature $\boldsymbol{h}^i$ and t, which gives $\boldsymbol{h}^{i\prime} = (\boldsymbol{h}^i, t)$. Then we input $\boldsymbol{x}$ and $\boldsymbol{h}' = (\boldsymbol{h}^{1\prime}, \dots, \boldsymbol{h}^{M\prime})$ to the EGNN as follows

$$(\boldsymbol{a}^x, \boldsymbol{a}^{h\prime}) = \mathrm{EGNN}(\boldsymbol{x}, \boldsymbol{h}') - (\boldsymbol{x}, \boldsymbol{0}).$$

Finally, we subtract the CoM of $\boldsymbol{a}^x$, and gets the parameterization of $\epsilon_{\boldsymbol{\theta}}(\boldsymbol{z}, t)$:

$$\epsilon_{\boldsymbol{\theta}}(\boldsymbol{z}, t) = (\boldsymbol{a}^x - \overline{\boldsymbol{a}^x}, \boldsymbol{a}^h),$$

where $\boldsymbol{a}^h$ comes from discarding the last component of $\boldsymbol{a}^{h\prime}$ that corresponds to the time.

For the conditional model $\epsilon_{\boldsymbol{\theta}}(\boldsymbol{z}, c, t)$, we additionally concatenate c to the atom feature $\boldsymbol{h}^i$, i.e., $\boldsymbol{h}^{i\prime} = (\boldsymbol{h}^i, t, c)$, and other parts in the parameterization remain the same.

D TRAINING OBJECTIVES OF ENERGY FUNCTIONS

Time-dependent property prediction model. Since the quantum property is a scalar, we train the time-dependent property prediction model $g(\boldsymbol{z}_t, t)$ using the ℓ_1 loss

$$\mathbb{E}_t \mathbb{E}_{q(c, \boldsymbol{z}_0, \boldsymbol{z}_t)} |g(\boldsymbol{z}_t, t) - c|,$$

where t is uniformly sampled from $[0, T]$.

Time-dependent multi-label classifier. Since the fingerprint is a bit map, predicting it can be viewed as a multi-label classification task. Thus, we use a time dependent multi-label classifier model $m(\boldsymbol{z}_t, t)$, and train it using the binary cross entropy loss

$$\mathbb{E}_t \mathbb{E}_{q(c, \boldsymbol{x}_0, \boldsymbol{x}_t)} \sum_{l=1}^{L} c_l \log m_l(\boldsymbol{z}_t, t) + (1 - c_l) \log(1 - m_l(\boldsymbol{z}_t, t)),$$

where t is uniformly sampled from $[0, T]$, and $m_l(\boldsymbol{z}_t, t)$ is the l-th component of $m(\boldsymbol{z}_t, t)$.

E EXPERIMENTAL DETAILS

E.1 THE "U-BOUND" AND "#ATOMS" BASELINES

The "U-bound" and "#Atoms" baselines are from Hoogeboom et al. (2022). The "U-bound" baseline shuffles the labels in D_b and then calculate the loss of ϕ_c on it, which can be regarded as an upper bound of the MAE. The "#Atoms" baseline predicts a quantum property c using only the number of atoms in a molecule.

E.2 GENERATING MOLECULES WITH DESIRED QUANTUM PROPERTIES

For the noise prediction network, we use the same setting with EDM Hoogeboom et al. (2022) for a fair comparison, where the models is trained $\sim$2000 epochs with a batch size of 64, a learning rate of 0.0001 with the Adam optimizer and an exponential moving average (EMA) with a rate of 0.9999.

The EGNN used in the energy function has 192 hidden features and 7 layers. We train 2000 epochs with a batch size of 128, a learning rate of 0.0001 with the Adam optimizer and an exponential moving average (EMA) with a rate of 0.9999.

E.3 GENERATING MOLECULES WITH TARGET STRUCTURES

QM9. For the backbone of the noise prediction network, we use a three-layer MLP with 768, 512 and 192 hidden nodes as embedding to encode the fingerprint and add the output of it with the embedding of atom features $\boldsymbol{h}$, which is then fed into the following EGNN. The EGNN has 256 hidden features and 9 layers. We train it 1500 epoch with a batch size of 64, a learning rate of 0.0001 with the Adam optimizer and an exponential moving average (EMA) with a rate of 0.9999. The energy function is trained with 1750 epoch with a batch size of 128, a learning rate of 0.0001 with the Adam optimizer and an exponential moving average (EMA) with a rate of 0.9999. Its EGNN has 192 hidden features and 7 layers.

GEOM-Drug. We use the same data split of GEOM-Drug with Hoogeboom et al. (2022), where training, validation and test set include 554K, 70K and 70K samples respectively. We train the noise

prediction network and the energy function on the training set and evaluate the Tanimoto similarity on the randomly selected 10k molecules from the test set. For the noise prediction network, we use the recommended hyperparameters of EDM (Hoogeboom et al., 2022), where the EGNN has 256 hidden features and 4 layers, and the other part is the same as that in QM9. We train 10 epoch with a batch size of 64, a learning rate of 0.0001 with the Adam optimizer and an exponential moving average (EMA) with a rate of 0.9999. The backbone of the energy function is the same as that in QM9 and we train the energy function for 14 epoch.

F ADDITIONAL RESULTS

F.1 ABLATION STUDY ON NOISE PREDICTION NETWORKS AND ENERGY GUIDANCE

We perform an ablation study on the conditioning, i.e., use conditional or unconditional noise prediction networks, and the energy guidance. When neither the conditioning nor the energy guidance is adopted, a single unconditional model can't perform conditional generation, and therefore we only report its atom stability and the molecule stability. As shown in Table 4, Table 5 and Table 6, the conditional noise prediction network improves the MAE compared to the unconditional one, and the energy guidance improves the MAE compared to a sole conditional model. Both the conditioning and the energy guidance do not affect the atom stability and the molecule stability much.

Table 4: Effects of conditioning and energy guidance on a single quantum property μ (D).

Conditional	Guidance	Scale	MAE↓	Novelty↑	AS↑	MS↑
×	×	-	-	82.92	98.37	81.74
×	✓	0.1	1.415	83.60	98.33	81.16
×	✓	0.5	1.241	83.84	98.27	80.75
✓	-	-	1.138	84.04	98.14	80.04
✓	✓	0.1	1.071	84.36	98.18	80.08
✓	✓	0.5	0.935	84.06	98.20	80.05

Table 5: Effects of conditioning and energy guidance on a single quantum property C_v ($\frac{\text{cal}}{\text{mol}}$K).

Conditional	Guidance	Scale	MAE↓	Novelty ↑	AS↑	MS↑
×	×	-	-	82.92	98.37	81.74
×	✓	1	2.360	84.15	98.36	81.44
×	✓	10	1.459	84.25	98.07	79.06
✓	×	-	1.066	83.64	98.25	80.50
✓	✓	1	1.038	83.71	98.21	80.61
✓	✓	10	0.939	83.92	98.06	79.21

Table 6: Effects of conditioning and energy guidance on multiple quantum properties C_v, μ.

Conditional	Guidance	Scale	MAE1↓	MAE2↓	Novelty↑	AS↑	MS↑
×	×	-	-	-	82.92	98.37	81.74
×	✓	1,0.1	2.381	1.421	83.81	98.31	80.52
×	✓	10,1	1.501	1.157	84.11	98.02	78.20
✓	×	-	1.075	1.163	85.45	97.99	77.02
✓	✓	1,0.1	1.053	1.097	85.20	97.96	77.08
✓	✓	10,1	0.982	0.916	85.51	97.58	73.94

F.2 REPRODUCE

We compare our reproduced results and the original results of EDM (Hoogeboom et al., 2022) in Table 7. The results are consistent.

Table 7: The reproduced results of EDM and L-bound are consistent with the original ones.

Quantum property	α	$\Delta\varepsilon$	$\varepsilon_{\text{HOMO}}$	$\varepsilon_{\text{LUMO}}$	μ	C_v
Unit	Bohr^3	meV	meV	meV	D	$\frac{\text{cal}}{\text{mol}}$K
EDM (Hoogeboom et al., 2022)	2.76	655	356	584	1.111	1.101
EDM (reproduce)	2.79	674	371	593	1.118	1.054
L-bound (Hoogeboom et al., 2022)	0.10	64	39	36	0.043	0.040
L-bound (reproduce)	0.09	65	39	36	0.043	0.040

F.3 GENERATING MOLECULES WITH TARGET STRUCTURES ON GEOM-DRUG

We provide the Tanimoto similarity between target structures and generated structures in Table 8. The conditional EDM baseline is trained following the default setting of Hoogeboom et al. (2022). We plot generated molecules on GEOM-Drug in Figure 5, and it can be observed that the atom types of generated molecules with EEGSDE often match the target better than EDM.

Table 8: Generation performance targeted to structures from test set on GEOM-Drugs dataset.

Method	EDM	EEGSDE (s=0.1)	EEGSDE (s=0.5)
Similarity↑	0.166	0.186	**0.194**

Figure 5: Generated molecules on GEOM-Drug.