LIPSCHITZ REGULARIZED DEEP NEURAL NETWORKS CONVERGE AND GENERALIZE

ABSTRACT

Generalization of deep neural networks (DNNs) is an open problem which, if solved, could impact the reliability of deep neural network architectures. In this paper, we show that if the usual fidelity term used in training DNNs is augmented by a Lipschitz regularization term, then the networks converge and generalize. The convergence is in the limit as the number of data points, $n \to \infty$, while also allowing the network to grow as needed to fit the data. Two regimes are identified: in the case of clean labels, we prove convergence to the label function which corresponds to zero loss, in the case of corrupted labels we prove convergence to a regularized label function which is the solution of a limiting variational problem. In both cases, a convergence rate is also provided.

1 INTRODUCTION

While deep neural networks networks (DNNs) give more accurate predictions than other machine learning methods (LeCun et al., 2015), they lack some of the performance guarantees of these methods. One step towards performance guarantees for DNNs is a proof of convergence with a rate, which could lead to quantitative error estimates. In this paper, we present such a result, for Lipschitz regularized DNNs. A proof of generalization follows as a consequence of convergence.

We begin by establishing the notation for our problem. We consider the classification problem to fix ideas, although regularization can apply to other problems as well.

Definition 1.1. Assume the data is normalized so that the data space is $X = [0,1]^d$. Write $\mathcal{D}_n = x_1, \ldots, x_n$ for the training data. Assume $\mathcal{D}_n$ is a sequence of *i.i.d.* random variables on X sampled from the probability distribution ρ. We consider the classification problem with m labels which are imbedded into the probability simplex, the label space, $Y \subset \mathbb{R}^D$. Write $u_0 : X \to Y$ for the map from data to label space, so that $y_i = u_0(x_i)$.

Our results point towards improved generalization results using Lipschitz regularization, which we define now.

Definition 1.2. Choose norms $\| \cdot \|_Y$, and $\| \cdot \|_X$ on X and Y, respectively. The Lipschitz constant (in these norms) of a function $u : X_0 \subset X \to Y$ is given by

$$\mathrm{Lip}(u; X_0) = \sup_{x_1, x_2 \in X_0} \frac{\|u(x_1) - u(x_2)\|_Y}{\|x_1 - x_2\|_X}$$

When X_0 is all of X, we write $\mathrm{Lip}(u; X) = \mathrm{Lip}(u)$. The Lipschitz constant of the data is $\mathrm{Lip}(u_0; \mathcal{D}_n)$.

Write $u(x; w)$ for the map from the input to data to the last layer of the network.[1] We consider the variational problem with Lipschitz regularization

$$\min_{u:X \to Y} J^n[u] = \frac{1}{n} \sum_{i=1}^{n} \ell(u(x_i; w), y_i) + \lambda \max(\mathrm{Lip}(u) - L_0, 0) \tag{1}$$

The first term in (1) is the usual averaged loss on the training data $\mathcal{D}_n$. The second term in (1) the Lipschitz regularization term: the excess Lipschitz constant of the map u, compared to the constant

---

[1] We apologize for not using the standard notation f for the last layer!

L_0. Choosing $\lambda > 0$ in (1) introduces a Lipschitz regularization penalty to the standard learning problem, which corresponds to $\lambda = 0$. In theory we take $L_0 = \text{Lip}(u_0)$, the Lipschitz constant of the data on the whole data manifold (discussed below).

Remark 1.3. In practice, $\text{Lip}(u_0)$ can be estimated by the Lipschitz constant of the data, $\text{Lip}(u_0; \mathcal{D}_n)$. In fact, for many common data sets, the Lipschitz constant is very small and the Lipschitz constants of networks is much larger (Finlay & Oberman, 2018), so in practice we can set $L_0 = 0$. However for clean data, if we wish to recover u_0 exactly, we need $L_0 > 0$.

If u is differentiable[2], then the boundedness of X implies

$$\text{Lip}(u; X) = \max_{x \in X} \|\nabla u(x; w)\|_{X,Y}$$

where $\|M\|_{X,Y}$ is the matrix norm induced by the norms on X and Y. In practise, we take the ∞-norm for Y and the 2-norm for X, and use explicit formulas for the $\|M\|_{\infty,2}$ norm, see §B. The Lipschitz constant of u, $\text{Lip}(u; X)$, can thus be estimated from below by the maximum of the norm of the gradient on a minibatch

$$\max_{i \in I} \|\nabla_x u(x_i; w)\|_{X,Y} \leq \text{Lip}(u; X) \tag{2}$$

Our analysis will apply to the problem (1) which is *convex* in u, and does not depend explicitly on the weights, w. Of course, once u is restricted to a fixed neural network architecture, the corresponding minimization problem becomes non-convex in the weights. Our analysis can avoid the dependence on the weights because we make the assumption that there are enough parameters so that u can exactly fit the training data. The assumption is justified by Zhang et al. (2016). As we send $n \to \infty$ for convergence, we require that the network also grow, in order to continue to satisfy this assumption. Our results apply to other non-parametric methods in this regime.

In Figure 1 we illustrate the solution of (1) (with $L_0 = 0$), using synthetic one dimensional data. In this case, the labels $\{-1, 0, 1\}$ are embedded naturally into $Y = \mathbb{R}$, and $\lambda = 0.1$. Notice that the solution matches the labels exactly on a subset of the data. In the second part of the figure, we show a solution with corrupted labels which introduce a large Lipschitz constant, in this case, the solution reduces the Lipschitz constant, thereby correcting the errors.

Figure 1: Synthetic labelled data and Lipschitz regularized solution u. Left: The solution value matches the labels exactly on a large portion of the data set. Right: 10% of the data is corrupted by incorrect labels; the regularized solution corrects the errors.

1.1 RELATED WORK AND APPLICATIONS

Generalization bounds have been obtained previously by using the Lipschitz constant of a network (Bartlett, 1997), as well as by using more general stability results (Bousquet & Elisseeff, 2002). More recently, (Bartlett et al., 2017) proposed the Lipschitz constant of the network as a candidate

---

[2]We follow the common practise and treat the the architecture as differentiable for training purposes. To be rigorous we can use the L^∞ norm and appeal to Rachemacher's Theorem.

measure for the Rademacher complexity, which is a measure of generalization (Shalev-Shwartz & Ben-David, 2014, Chapter 26). Also, Cranko et al. (2018) showed that Lipschtiz regularization can be viewed as a special case of distributional robustness. Unlike other recent contributions such as (Hardt et al., 2015), our analysis does not depend on the training method. In fact, our analysis has more in common with inverse problems in image processing, such as Total Variation denoising and inpainting (Bertalmio et al., 2000; Rudin et al., 1992). For further discussion, see Appendix C.

The estimate of $\text{Lip}(u; X)$ provided by (2) can be quite different from the the Tychonoff gradient regularization (Drucker & Le Cun, 1992),

$$\frac{1}{|I|} \sum_{i \in I} \|\nabla_x u(x_i)\|^2$$

since (2) corresponds to a maximum of the values of the norms, and the previous equation corresponds to average of the values. In fact, recent work on semi-supervised learning suggests that higher p-norms of the gradient are needed for generalization when the data manifold is not well approximated by the data (El Alaoui et al., 2016; Calder, 2017; Kyng et al., 2015; Slepcev & Thorpe, 2017). In Figure 2 we compare to the problems in Figure 1 using Tychonoff regularization. The Tychonoff regularization is less effective at correcting errors. The effect is more pronounced in higher dimensions.

Figure 2: Synthetic labelled data and Tychonoff regularized solution u. Left: The solution value matches the labels exactly on a large portion of the data set. Right: 10% of the data is corrupted by incorrect labels; the regularized solution is not as effective at correcting errors. The effect is more pronounced in higher dimenions.

An upper bound for the Lipschitz constant of the model is given by the norm of the product of the weight matrices (Szegedy et al., 2013, Section 4.3). Let $w = (w^1, \ldots, w^J)$ be the weight matrices for each layer. Then

$$\text{Lip}(u; X) \leq \Pi_{j=1}^{J} \|w^i\|. \tag{3}$$

Regularization of the network using methods based on (3) has been implemented recently in (Gouk et al., 2018) and (Yoshida & Miyato, 2017). Because the upper bound in (3) does not take into account the coefficients in weight matrices which are zero due to the activation functions, the gap in the inequality can be off by factors of many orders of magnitude for deep networks (Finlay & Oberman, 2018).

Implementing (2) can be accomplished using backpropragation in the x variable on each label, which can become costly for m large. Special architectures could also be used to implement Lipschitz regularization, for example, on a restricted architecture, Liao et al. (2018) renormalized the weight matrices of each layer to be norm 1. In practise, the computational cost can be reduced by regularizing the loss applied to the model $\ell \circ u$, instead

$$\max_{i \in I} \|\nabla_x \ell(u(x_i))\| \leq \text{Lip}(\ell \circ u; X) \tag{4}$$

which requires only one backpropagation. Lipschitz regularization may help with adversarial examples (Szegedy et al., 2013) (Goodfellow et al., 2014) which poses a problem for model reliability (Goodfellow et al., 2018). Since the Lipschitz constant L_ℓ of the loss, ℓ, controls the norm of a perturbation

$$\|\ell(u(x_i + \epsilon v)) - \ell(u(x_i))\|_Y \leq \epsilon L_\ell \|v\|_X$$

maps with smaller Lipschitz constants may be more robust to adversarial examples. Finlay & Oberman (2018) implemented Lipschitz regularization of the loss, as in (4), and achieved better robustness against adversarial examples, compared to adversarial training (Goodfellow et al., 2014) alone.

Lipschitz regularization may also improve stability of GANs. 1-Lipschitz networks are also important for Wasserstein-GANs (Arjovsky et al., 2017) (Arjovsky & Bottou, 2017). In (Wei et al., 2018) the gradient penalty away from norm 1 is implemented, augmented by a penalty around perturbed points, with the goal of improved stability. Spectral regularization for GANs was implemented in (Miyato et al., 2018).

Problem 1.4. Consider the following problem inspired by (Zhang et al., 2016). Suppose we are given a labelled data set, which has Lipchitz constant $\text{Lip}(\mathcal{D}) = \mathcal{O}(1)$, where

$$\text{Lip}(\mathcal{D}) = \max_{i \neq j} \frac{\|y_i - y_j\|}{\|x_i - x_j\|} \tag{5}$$

Consider making copies of 10 percent of the data, adding a vector of norm ϵ to the perturbed data points, and changing the label. Call the new *dirty* data set $\tilde{\mathcal{D}}$. The new data set has $\text{Lip}(\tilde{\mathcal{D}}) = \mathcal{O}(1/\epsilon)$. However, the distribution of the values on the right hand side of (5) are mostly below $\text{Lip}(\mathcal{D})$ with a small fraction $\mathcal{O}(1/\epsilon)$, since the duplicated images are ϵ apart but with different labels.

2 LIPSCHITZ REGULARIZATION AND CONVERGENCE

2.1 LIMITING PROBLEM

The variational problem (1) admits Lipschitz continuous minimizers, but in general the minimizers are not unique. When $L_0 = \text{Lip}(u_0)$, it is clear that u_0, is a solution of (1): both the loss term and the regularization term are zero when applied to u_0. In addition, any L_0-Lipschitz extension of $u_0|_{\mathcal{D}_n}$ is also a minimizer of (1), so solutions are not unique.

Let u_n be any solution of the Lipschitz regularized variational problem (1). We study the limit of u_n as $n \to \infty$. Since the empirical probability measures ρ_n converge to the data distribution ρ, the continuum variational problem corresponding to (1) is

$$\min_{u: X \to Y} J[u] \equiv L[u; \rho] + \lambda \max(\text{Lip}(u) - L_0, 0), \tag{6}$$

where in (6) we have introduced the following notation.

Definition 2.1. Given the loss function, ℓ, a map $u : X \to Y$, and a probability measure, μ, supported on X, define

$$L[u, \mu] = \mathbb{E}_{x \sim \mu}[\ell(u(x), u_0(x))] = \int_X \ell(u(x), u_0(x)) d\mu(x)$$

to be the expectation of the loss with respect to the measure. In particular, the *generalization loss* of the map $u : X \to Y$ is given by $L[u, \rho]$. Write $L[u, \mathcal{D}_n] := L[u, \rho_n]$ for the average loss on the data set $\mathcal{D}_n$, where $\rho_n := \frac{1}{n} \sum \delta_{x_i}$ is the empirical measure corresponding to $\mathcal{D}_n$.

Remark 2.2. Generalization is defined in (Goodfellow et al., 2016, Section 5.2) as the expected value of the loss function on a new input sampled from the data distribution. As defined, the full generalization error includes the training data, but it is of measure zero, so removing it does not change the value.

2.2 LOSS FUNCTION ASSUMPTIONS

We introduce the following assumption on the loss function.

Assumption 2.3 (Loss function). The function $\ell : Y \times Y \to \mathbb{R}$ is a *loss* function if it satisfies (i) $\ell \geq 0$, (ii) $\ell(y_1, y_2) = 0$ if and only if $y_1 = y_2$, and (iii) ℓ is strictly convex in y_1.

Example 2.4 ($\mathbb{R}^D$ with L^2 loss). Set $Y = \mathbb{R}^D$, and let each label be a basis vector. Set $\ell(y_1, y_2) = \|y_1 - y_2\|_2^2$ to be the L^2 loss.

Example 2.5 (Classification). In classification problems, the output of the network is a probability vector on the labels. Thus $Y = \Delta_p$, the p-dimensional probability simplex, and each label is mapped to a basis vector. The cross-entropy loss is given by $\ell^{KL}(y, z) = -\sum_{i=1}^{p} z_i \log(y_i/z_i)$. For labels, $\ell^{KL}(y, e_k) = -\log(y_k)$.

Example 2.6. In the classification setting, it is often the case that the softmax function

$$\mathrm{softmax}(z)_j = \frac{e^{z_j}}{\sum_{k=1}^{D} e^{z_k}} \tag{7}$$

is combined with the cross-entropy loss. In this paper, we choose to absorb softmax into the last layer of the DNN, and assume the output $u(x)$ of the network lies in the probability simplex. If the output, z, of the second to last layer of the DNN, which is the input to softmax in (7), lies in a compact set, i.e., $|z_j| \leq C$ for all i and some $C > 0$, then $\mathrm{softmax}(z)_j \geq e^{-2C}$, and so the range of softmax lies in the set

$$A := \{y \in \mathbb{R}^D \, : \, y_i \geq e^{-2C} \text{ and } y_1 + \cdots + y_D = 1\},$$

which is strictly interior to the probability simplex. Restricted to A, the cross-entropy loss ℓ^{KL} is strongly convex and Lipschitz continuous, which is required in Theorems 2.11 and 2.15 below.

In our analysis, it is slightly more convenient to define the *regularized cross entropy loss* with parameter $\epsilon > 0$ by

$$\ell_\epsilon^{KL}(y, z) = -\sum_{i=1}^{p} z_i \log\left(\frac{y_i + \epsilon}{z_i}\right).$$

For classification problems, where $z = e_k$, we have $\ell_\epsilon^{KL}(y, e_k) = -\log(y_k + \epsilon)$, which is Lipschitz and strongly convex for any $0 \leq y_i \leq 1$ within the probability simplex. Thus, the regularized cross entropy ℓ_ϵ^{KL} satisfies the strong convexity and Lipschitz regularity required by Theorems 2.11 and 2.15 on the whole probability simplex. We note that while $\ell_\epsilon^{KL}(e_k, e_k) = -\log(1 + \epsilon) \neq 0$, as is required by Assumption 2.3 (ii), it is trivial to apply the theory by adding the constant $C = \log(1+\epsilon)$ to the loss.

2.3 CONVERGENCE RESULT

Here, we show that solutions of the random variational problem (1) converge to solutions of (6). We make the standard manifold assumption (Chapelle et al., 2006), and assume the data distribution ρ is a probability density supported on a compact, smooth, m-dimensional manifold $\mathcal{M}$ embedded in $X = [0,1]^d$, where $m \ll d$. We denote the probability density again by $\rho : \mathcal{M} \to [0, \infty)$. Hence, the data $\mathcal{D}_n$ is a sequence $x_1, \ldots, x_n$ of *i.i.d.* random variables on $\mathcal{M}$ with probability density ρ. Associated with the random sample we have the closet point projection map $\sigma_n : X \to \{x_1, \ldots, x_n\} \subset X$ that satisfies

$$\|x - \sigma_n(x)\|_X = \min_{1 \leq i \leq n} \{\|x - x_i\|_X\}$$

for all $x \in X$. We recall that $W^{1,\infty}(X; Y)$ is the space of Lipschitz mappings from X to Y. Throughout this section, $C, c > 0$ denote positive constants depending only on $\mathcal{M}$, and we assume $C \geq 1$ and $0 < c < 1$. We follow the analysis tradition of allowing the particular values of C and c to change from line to line.

We establish that that minimizers of (6) are unique on $\mathcal{M}$, which follows from the strict convexity of the loss restricted to the data manifold $\mathcal{M}$, in Theorem A.1. See also Figure 3 which shows how the solutions need not be unique off the data manifold.

Our first convergence result is in the case where $\mathrm{Lip}[u_0] \leq L_0$, and so the Lipschitz regularizer is not fully active. This corresponds to the case of clean labels. We state our result in generality, for approximate minimizers of (1), and specialize to the case $\mathrm{Lip}[u_0] \leq L_0$ in Remark 2.8.

Theorem 2.7 (Convergence result). *Assume* $\inf_{x \in \mathcal{M}} \rho(x) > 0$. *For any* $t > 0$, *with probability at least* $1 - Ct^{-1}n^{-(ct-1)}$ *every sequence* $u_n \in W^{1,\infty}(X;Y)$ *with zero empircal loss* $L[u_0, \rho_n] = 0$ *satisfies*

$$\|u_0 - u_n\|_{L^\infty(\mathcal{M};Y)} \leq C(L_0 + \mathrm{Lip}[u_n]) \left(\frac{t \log(n)}{n} \right)^{1/m}.$$

Remark 2.8. If $u_n \in W^{1,\infty}(X;Y)$ is any sequence of minimizers of (1) and $\mathrm{Lip}[u_0] \leq L_0$, then $J[u_n] \leq J[u_0] = 0$. Thus, $\mathrm{Lip}[u_n] \leq L_0$ and Theorem 2.7 applies to the sequence u_n, yielding

$$\|u_0 - u_n\|_{L^\infty(\mathcal{M};Y)} \leq CL_0 \left(\frac{t \log(n)}{n} \right)^{1/m}.$$

It is important to note that Theorem 2.7 does not requires u_n to be minimizers of (1)—we just require zero epmpirical loss, which is often acheived in practice (Zhang et al., 2016). This allows for approximation errors in solving (1) on the whole domain X, due to the restriction that u must be expressed via a Deep Neural Network.

As an immediate corollary, we can prove that the generalization loss converges to zero, and so we obtain generalization.

Corollary 2.9. *Assume that for some* $q \geq 1$ *the loss* ℓ *satisfies*

$$\ell(y, y_0) \leq C \|y - y_0\|_Y^q \quad \text{for all } y_0, y \in Y. \tag{8}$$

Then under the assumptions of Theorem 2.7

$$L[u_n, \rho] \leq C(L_0 + \mathrm{Lip}[u_n])^q \left(\frac{t \log(n)}{n} \right)^{q/m}$$

holds with probability at least $1 - Ct^{-1}n^{-(ct-1)}$.

Proof. By (8), we can bound the generalization loss as follows

$$L[u_n, \rho] = \int_{\mathcal{M}} \ell(u_n(x), u_0(x))\rho(x) \, dVol(x) \leq C \|u_n - u_0\|_{L^\infty(\mathcal{M};Y)}^q.$$

The proof is completed by invoking Theorem 2.7. $\qquad\square$

We now turn to the proof of Theorem 2.7, which requires a bound on the distance between the closest point projection σ_n and the identity. The result is standard in probability, and we include it for completeness in Lemma 2.10 proved in §A.1. We refer the interested reader to (Penrose et al., 2003) for more details.

Lemma 2.10. *Suppose that* $\inf_{\mathcal{M}} \rho > 0$. *Then for any* $t > 0$

$$\|Id - \sigma_n\|_{L^\infty(\mathcal{M};X)} \leq C \left(\frac{t \log(n)}{n} \right)^{1/m}$$

with probability at least $1 - Ct^{-1}n^{-(ct-1)}$.

We now give the proof of Theorem 2.7.

Proof of Theorem 2.7. Since $L[u_n, \rho_n] = 0$ we have $u_0(x_i) = u_n(x_i)$ for all $1 \leq i \leq n$. Thus for any $x \in X$ we have

$$\|u_0(x) - u_n(x)\|_Y = \|u_0(x) - u_0(\sigma_n(x)) + u_0(\sigma_n(x)) - u_n(\sigma_n(x)) + u_n(\sigma_n(x)) - u_n(x)\|_Y$$

$$\leq \|u_0(x) - u_0(\sigma_n(x))\|_Y + \|u_n(\sigma_n(x)) - u_n(x)\|_Y$$
$$\leq (L_0 + \mathrm{Lip}[u_n])\|x - \sigma_n(x)\|_X.$$

Therefore, we deduce

$$\|u_0 - u_n\|_{L^\infty(\mathcal{M};Y)} \leq (L_0 + \mathrm{Lip}[u_n])\|Id - \sigma_n\|_{L^\infty(\mathcal{M};X)}.$$

The proof is completed by invoking Lemma 2.10. $\qquad\square$

Figure 3: Non-uniqueness of the solution off the data manifold: in the middle areas off the data manifold where the Lipschitz constant is attained, the solution is unique. In the outer area off the data manifold, the solution is not unique.

2.4 CONVERGENCE FOR NOISY LABELS

We now address problems with some degree of uncertainty. Recall the discussion in Problem 1.4 of the setting where labels are corrupted, which is also illustrated in the right hand side of Figure 1. We assume the true labels u_{true} are corrupted by additive error of the form

$$u_0 = u_{\text{true}} + e$$

for an error function $e : X \to Y$. We assume the true labels u_{true} have a small Lipschtiz constant L_0, while the error vector e has a much larger Lipschitz constant, leading to $\text{Lip}(u_0) \gg L_0$. A simple model is described in Problem 1.4.

In this case, it is desirable to fit some, but not all, of the data. In particular, we wish to fit the subset of the labels that generates the low Lipschitz constant L_0, while hopefully ignoring the rest, which correspond to pairs of labels that generate a high Lipschitz constant. In this setting, the Lipschitz regularizer is active, leading to smoothing of the noisy labels as depicted on the right side of Figure 1. Our main results (Theorems 2.11 and 2.15) show that minimizers of J^n converge to minimizers of J almost surely as the number of training points n tends to ∞.

Theorem 2.11. *Suppose that $\inf_{\mathcal{M}} \rho > 0$, $\ell : Y \times Y \to \mathbb{R}$ is Lipschitz, and let $u^* \in W^{1,\infty}(X;Y)$ be any minimizer of* (6). *Then with probability one*

$$u_n \longrightarrow u^* \quad \text{uniformly on } \mathcal{M} \text{ as } n \to \infty, \tag{9}$$

where u_n is any sequence of minimizers of (1). *Furthermore, every uniformly convergent subsequence of u_n converges on X to a minimizer of* (6).

Remark 2.12. In Theorem 2.11 and Theorem 2.15 below, the sequence u_n does not, in general, converge on the whole domain X. The important point is that the sequence converges on the data manifold $\mathcal{M}$, and solves the variational problem (6) off of the manifold, which ensures that the output of the DNN is stable with respect to the input. See Figure 3.

The proof of Theorem 2.11 requires a preliminary Lemma. Let $H_L(X;Y)$ denote the collection of L-Lipschitz functions $w : X \to Y$.

Lemma 2.13. *Suppose that $\inf_{\mathcal{M}} \rho > 0$, and $\dim(\mathcal{M}) = m$. Then for any $t > 0$*

$$\sup_{w \in H_L(X;Y)} \left| \frac{1}{n} \sum_{i=1}^{n} w(x_i) - \int_{\mathcal{M}} w \rho \, dVol(x) \right| \le CL \left(\frac{t \log(n)}{n} \right)^{\frac{1}{m+2}} \tag{10}$$

holds with probability at least $1 - 2t^{-\frac{m}{m+2}} n^{-(ct-1)}$.

The estimate (10) is called a discrepancy result (Talagrand, 2006; Györfi et al., 2006), and is a uniform version of concentration inequalities. We include a simple proof in Section A.2.

Proof of Theorem 2.11. By Lemma 2.13 the event that

$$\lim_{n \to \infty} \sup_{w \in H_L(X;Y)} |L[w, \rho_n] - L[w, \rho]| = 0 \tag{11}$$

for all Lipschitz constants $L > 0$ has probability one. For the rest of the proof we restrict ourselves to this event.

Let $u_n \in W^{1,\infty}(X;Y)$ be a sequence of minimizers of (1), and let $u^* \in W^{1,\infty}(X;Y)$ be any minimizer of (6). Then since

$$\lambda(\mathrm{Lip}(u_n) - L_0) \leq J^n[u_n] \leq J^n[u_0] = \lambda(\mathrm{Lip}(u_0) - L_0)$$

we have $\mathrm{Lip}(u_n) \leq \mathrm{Lip}(u_0) =: L$ for all n. By the Arzelà-Ascoli Theorem (Rudin, 1976) there exists a subsequence u_{n_j} and a function $u \in W^{1,\infty}(X;Y)$ such that $u_{n_j} \to u$ uniformly as $n_j \to \infty$. Note we also have $\mathrm{Lip}(u) \leq \liminf_{j \to \infty} \mathrm{Lip}(u_{n_j})$. Since

$$\begin{aligned}
|L[u_n, \rho_n] - L[u, \rho]| &\leq |L[u_n, \rho_n] - L[u, \rho_n]| + |L[u, \rho_n] - L[u, \rho]| \\
&\leq C\|u_n - u\|_{L^\infty(\mathcal{M};Y)} + \sup_{w \in H_L(X;Y)} |L[w, \rho_n] - L[w, \rho]|
\end{aligned}$$

it follows from (11) that $L[u_{n_j}, \rho_{n_j}] \to L[u, \rho]$ as $j \to \infty$. It also follows from (11) that $J^n[u^*] \to J[u^*]$ as $n \to \infty$. Therefore

$$\begin{aligned}
J[u^*] &= \lim_{n \to \infty} J^n[u^*] \\
&\geq \liminf_{n \to \infty} J^n[u_n] \\
&= \liminf_{n \to \infty} L[u_n, \rho_n] + \lambda \max(\mathrm{Lip}(u_n) - L_0, 0) \\
&= \lim_{n \to \infty} L[u_n, \rho_n] + \liminf_{n \to \infty} \lambda \max(\mathrm{Lip}(u_n) - L_0, 0) \\
&\geq L[u, \rho] + \lambda \max(\mathrm{Lip}(u) - L_0, 0) = J[u].
\end{aligned}$$

Therefore, u is a minimizer of J. By Theorem A.1, $u = u^*$ on $\mathcal{M}$, and so $u_{n_j} \to u^*$ uniformly on $\mathcal{M}$ as $j \to \infty$.

Now, suppose that (9) does not hold. Then there exists a subsequence u_{n_j} and $\delta > 0$ such that

$$\max_{x \in \mathcal{M}} |u_{n_j}(x) - u^*(x)| > \delta$$

for all $j \geq 1$. However, we can apply the argument above to extract a further subsequence of u_{n_j} that converges uniformly on $\mathcal{M}$ to u^*, which is a contradiction. This completes the proof. $\square$

Finally, we prove a rate in the case where the loss ℓ is strongly convex in the first variable.

Definition 2.14. We say that ℓ is strongly convex with parameter $\theta > 0$ if

$$\ell(ty_1 + (1-t)y_2, y_0) + \frac{\theta}{2}t(1-t)\|y_1 - y_2\|_Y^2 \leq t\ell(y_1, y_0) + (1-t)\ell(y_2, y_0) \tag{12}$$

for all $y_0, y_1, y_2 \in Y$ and $0 \leq t \leq 1$.

We note that when ℓ is twice differentiable, this notion of strong convexity is equivalent to assuming $\nabla_{y_1}^2 \ell \geq \theta I$. The definition in equation (12) is useful for non-smooth functions, such as the Lipschitz semi-norm present in $J[u]$.

Our final result is the following L^2 convergence rate in the strongly convex setting.

Theorem 2.15. *Suppose that $\ell : Y \times Y \to \mathbb{R}$ is Lipschitz and strongly convex and let $L = \mathrm{Lip}(u_0)$. Then for any $t > 0$, with probability at least $1 - 2t^{-\frac{m}{m+2}} n^{-(ct-1)}$ all minimizing sequences u_n of (1) and all minimizers u^* of (6) satisfy*

$$\frac{\theta}{2} \int_{\mathcal{M}} \|u_n - u^*\|_Y^2 \rho \, dVol(x) \leq CL \left(\frac{t \log(n)}{n} \right)^{\frac{1}{m+2}}.$$

Before proving Theorem 2.15, we require a preliminary lemma.

Lemma 2.16. *If $u^* \in W^{1,\infty}(X; Y)$ is a minimizer of* (6) *and $u \in W^{1,\infty}(X; Y)$ then*

$$\frac{\theta}{2} \int_{\mathcal{M}} \|u - u^*\|_Y^2 \rho \, dVol(x) \le J[u] - J[u^*].$$

Proof. We use Proposition A.4 with $u_1 = u^*$ and $u_2 = u$ to obtain

$$J[tu^* + (1-t)u] + \frac{\theta}{2}t(1-t) \int_{\mathcal{M}} \|u^* - u\|_Y^2 \rho \, dVol(x) \le tJ[u^*] + (1-t)J[u].$$

Since $J[tu^* + (1-t)u] \ge J[u^*]$

$$J[u^*] + \frac{\theta}{2}t(1-t) \int_{\mathcal{M}} \|u^* - u\|_Y^2 \rho \, dVol(x) \le tJ[u^*] + (1-t)J[u],$$

and so

$$\frac{\theta}{2}t \int_{\mathcal{M}} \|u^* - u\|_Y^2 \rho \, dVol(x) \le J[u] - J[u^*].$$

Setting $t = 1$ completes the proof. $\qquad\square$

Proof of Theorem 2.15. Let $L = \text{Lip}(u_0)$. By Lemma 2.13

$$\sup_{w \in H_L(X;Y)} |L[w, \rho_n] - L[w, \rho]| \le CL \left(\frac{t \log(n)}{n} \right)^{\frac{1}{m+2}} \tag{13}$$

holds with probability at least $1 - 2t^{-\frac{m}{m+2}} n^{-(ct-1)}$ for any $t > 0$. Let us assume for the rest of the proof that (13) holds.

As in the proof of Theorem 2.11, we have $\text{Lip}(u_n) \le L$ and $\text{Lip}(u^*) \le L$, and so

$$|J^n[u^*] - J[u^*]|, |J^n[u_n] - J[u_n]| \le CL \left(\frac{t \log(n)}{n} \right)^{\frac{1}{m+2}}.$$

Therefore

$$J[u_n] - J[u^*] = J^n[u_n] - J[u^*] + J[u_n] - J^n[u_n] \le CL \left(\frac{t \log(n)}{n} \right)^{\frac{1}{m+2}}.$$

By Lemma 2.16 we deduce

$$\frac{\theta}{2} \int_{\mathcal{M}} \|u_n - u^*\|_Y^2 \rho \, dVol(x) \le CL \left(\frac{t \log(n)}{n} \right)^{\frac{1}{m+2}},$$

which completes the proof. $\qquad\square$

REFERENCES

Martin Arjovsky and Léon Bottou. Towards principled methods for training generative adversarial networks. *arXiv preprint arXiv:1701.04862*, 2017.

Martin Arjovsky, Soumith Chintala, and Léon Bottou. Wasserstein gan. *arXiv preprint arXiv:1701.07875*, 2017.

Gunnar Aronsson, Michael Crandall, and Petri Juutinen. A tour of the theory of absolutely minimizing functions. *Bulletin of the American mathematical society*, 41(4):439–505, 2004.

Gilles Aubert and Pierre Kornprobst. *Mathematical problems in image processing: partial differential equations and the calculus of variations*, volume 147. Springer Science & Business Media, 2006.

Peter L Bartlett. For valid generalization the size of the weights is more important than the size of the network. In *Advances in neural information processing systems*, pp. 134–140, 1997.

Peter L Bartlett, Dylan J Foster, and Matus J Telgarsky. Spectrally-normalized margin bounds for neural networks. In *Advances in Neural Information Processing Systems*, pp. 6240–6249, 2017.

Marcelo Bertalmio, Guillermo Sapiro, Vincent Caselles, and Coloma Ballester. Image inpainting. In *Proceedings of the 27th annual conference on Computer graphics and interactive techniques*, pp. 417–424. ACM Press/Addison-Wesley Publishing Co., 2000.

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.

Olivier Bousquet and André Elisseeff. Stability and generalization. *Journal of machine learning research*, 2(Mar):499–526, 2002.

Andrea Braides. *Gamma-convergence for Beginners*, volume 22. Clarendon Press, 2002.

Jeff Calder. Consistency of lipschitz learning with infinite unlabeled data and finite labeled data. *arXiv preprint arXiv:1710.10364*, 2017.

Olivier Chapelle, Bernhard Scholkopf, and Alexander Zien. *Semi-supervised learning*. MIT, 2006.

Zac Cranko, Simon Kornblith, Zhan Shi, and Richard Nock. Lipschitz networks and distributional robustness. *arXiv preprint arXiv:1809.01129*, 2018.

Bernard Dacorogna. *Direct methods in the calculus of variations*, volume 78. Springer Science & Business Media, 2007.

Harris Drucker and Yann Le Cun. Improving generalization performance using double backpropagation. *IEEE Transactions on Neural Networks*, 3(6):991–997, 1992.

Ahmed El Alaoui, Xiang Cheng, Aaditya Ramdas, Martin J Wainwright, and Michael I Jordan. Asymptotic behavior of\ell_p-based laplacian regularization in semi-supervised learning. In *Conference on Learning Theory*, pp. 879–906, 2016.

Christopher Elion and Luminita A Vese. An image decomposition model using the total variation and the infinity laplacian. In *Computational Imaging V*, volume 6498, pp. 64980W. International Society for Optics and Photonics, 2007.

Lawrence C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, 1998. ISBN 0-8218-0772-2.

Chris Finlay and Adam M Oberman. Improved robustness to adversarial examples using lipschitz regularization of the loss, 2018.

Ian Goodfellow, Yoshua Bengio, and Aaron Courville. *Deep Learning*. MIT Press, 2016. http://www.deeplearningbook.org.

Ian Goodfellow, Patrick McDaniel, and Nicolas Papernot. Making machine learning robust against adversarial inputs. *Communications of the ACM*, 61(7):56–66, June 2018. ISSN 00010782. doi: 10.1145/3134599. URL http://dl.acm.org/citation.cfm?doid=3234519.3134599.

Ian J Goodfellow, Jonathon Shlens, and Christian Szegedy. Explaining and harnessing adversarial examples. *arXiv preprint arXiv:1412.6572*, 2014.

Henry Gouk, Eibe Frank, Bernhard Pfahringer, and Michael Cree. Regularisation of neural networks by enforcing lipschitz continuity. *arXiv preprint arXiv:1804.04368*, 2018.

Laurence Guillot and Carole Le Guyader. Extrapolation of vector fields using the infinity laplacian and with applications to image segmentation. In Xue-Cheng Tai, Knut Mørken, Marius Lysaker, and Knut-Andreas Lie (eds.), *Scale Space and Variational Methods in Computer Vision*, pp. 87–99, Berlin, Heidelberg, 2009. Springer Berlin Heidelberg. ISBN 978-3-642-02256-2.

László Györfi, Michael Kohler, Adam Krzyzak, and Harro Walk. *A distribution-free theory of nonparametric regression*. Springer Science & Business Media, 2006.

Moritz Hardt, Benjamin Recht, and Yoram Singer. Train faster, generalize better: Stability of stochastic gradient descent. *arXiv preprint arXiv:1509.01240*, 2015.

Roger A Horn, Roger A Horn, and Charles R Johnson. *Matrix analysis*. Cambridge university press, 1990.

William B Johnson and Joram Lindenstrauss. Extensions of lipschitz mappings into a hilbert space. *Contemporary mathematics*, 26(189-206):1, 1984.

Rasmus Kyng, Anup Rao, Sushant Sachdeva, and Daniel A Spielman. Algorithms for lipschitz learning on graphs. In *Conference on Learning Theory*, pp. 1190–1223, 2015.

Yann LeCun, Yoshua Bengio, and Geoffrey Hinton. Deep learning. *nature*, 521(7553):436, 2015.

Qianli Liao, Brando Miranda, Andrzej Banburski, Jack Hidary, and Tomaso Poggio. A surprising linear relationship predicts test performance in deep networks. *arXiv preprint arXiv:1807.09659*, 2018.

Edward James McShane. Extension of range of functions. *Bulletin of the American Mathematical Society*, 40(12):837–842, 1934.

Takeru Miyato, Toshiki Kataoka, Masanori Koyama, and Yuichi Yoshida. Spectral normalization for generative adversarial networks. *arXiv preprint arXiv:1802.05957*, 2018.

Mathew Penrose et al. *Random geometric graphs*. Number 5. Oxford university press, 2003.

Thomas Pock, Daniel Cremers, Horst Bischof, and Antonin Chambolle. Global solutions of variational models with convex regularization. *SIAM Journal on Imaging Sciences*, 3(4):1122–1145, 2010.

Leonid I Rudin, Stanley Osher, and Emad Fatemi. Nonlinear total variation based noise removal algorithms. *Physica D: nonlinear phenomena*, 60(1-4):259–268, 1992.

Walter Rudin. *Principles of mathematical analysis*. McGraw-hill New York, 1976.

Shai Shalev-Shwartz and Shai Ben-David. *Understanding Machine Learning: From Theory to Algorithms*. Cambridge University Press, 2014. doi: 10.1017/CBO9781107298019.

Dejan Slepcev and Matthew Thorpe. Analysis of p-laplacian regularization in semi-supervised learning. *arXiv preprint arXiv:1707.06213*, 2017.

Christian Szegedy, Wojciech Zaremba, Ilya Sutskever, Joan Bruna, Dumitru Erhan, Ian Goodfellow, and Rob Fergus. Intriguing properties of neural networks. *arXiv preprint arXiv:1312.6199*, 2013.

Michel Talagrand. *The generic chaining: upper and lower bounds of stochastic processes*. Springer Science & Business Media, 2006.

AN Tikhonov and V Ya Arsenin. *Solutions of Ill-Posed Problems*. Winston and Sons, New York, 1977.

Xiang Wei, Boqing Gong, Zixia Liu, Wei Lu, and Liqiang Wang. Improving the improved training of wasserstein gans: A consistency term and its dual effect. *arXiv preprint arXiv:1803.01541*, 2018.

Yuichi Yoshida and Takeru Miyato. Spectral norm regularization for improving the generalizability of deep learning. *arXiv preprint arXiv:1705.10941*, 2017.

Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. *arXiv:1611.03530*, 2016.

A PROOFS

A.1 PROOFS FOR CLEAN LABELS

In this section we provide the proof of results stated in §2.3.

Theorem A.1. *Suppose the loss function satisfies Assumption 2.3. If $u, v \in W^{1,\infty}(X; Y)$ are two minimizers of (6) and $\inf_{\mathcal{M}} \rho > 0$ then $u = v$ on $\mathcal{M}$.*

Proof. Let $w = (u + v)/2$. Then

$$
\begin{aligned}
J[w] &= \int_{\mathcal{M}} \ell \left(\tfrac{1}{2}u + \tfrac{1}{2}v, u_0 \right) \rho \, dVol(x) + \lambda \max \left(\text{Lip} \left(\tfrac{1}{2}u + \tfrac{1}{2}v \right), 0 \right) \\
&\leq \int_{\mathcal{M}} \left[\tfrac{1}{2}\ell(u, u_0) + \tfrac{1}{2}\ell(v, u_0) \right] \rho \, dVol(x) + \lambda \max \left(\tfrac{1}{2}\text{Lip}(u) + \tfrac{1}{2}\text{Lip}(v), 0 \right) \\
&\leq \int_{\mathcal{M}} \left[\tfrac{1}{2}\ell(u, u_0) + \tfrac{1}{2}\ell(v, u_0) \right] \rho \, dVol(x) + \lambda \left[\tfrac{1}{2}\max(\text{Lip}(u), 0) + \tfrac{1}{2}\max(\text{Lip}(v), 0) \right] \\[4pt]
&= \tfrac{1}{2}J[u] + \tfrac{1}{2}J[v] = \min_{u} J[u].
\end{aligned}
$$

Therefore, w is a minimizer of J and so we have equality above, which yields

$$
\int_{\mathcal{M}} \left[\tfrac{1}{2}\ell(u, u_0) + \tfrac{1}{2}\ell(v, u_0) \right] \rho \, dVol(x) = \int_{\mathcal{M}} \ell \left(\tfrac{1}{2}u + \tfrac{1}{2}v, u_0 \right) \rho \, dVol(x).
$$

Since ℓ is strictly convex in its first argument, it follows that $u = v$ on $\mathcal{M}$. □

Proof of Lemma 2.10 of §2.3. There exists $\epsilon_{\mathcal{M}}$ such that for any $0 < \epsilon \leq \epsilon_{\mathcal{M}}$, we can cover $\mathcal{M}$ with N geodesic balls $B_1, B_2, \ldots, B_N$ of radius ϵ, where $N \leq C\epsilon^{-m}$ and C depends only on $\mathcal{M}$ (Györfi et al., 2006). Let Z_i denote the number of random variables $x_1, \ldots, x_n$ falling in B_i. Then $Z_i \sim B(n, p_i)$, where $p_i = \int_{B_i} \rho(x) \, dVol(x)$. Since $\rho \geq \theta > 0$ and $Vol(B_i) \geq c\epsilon^m$ we have $p_i \geq c\epsilon^m$. Let A_n denote the event that at least one B_i is empty (i.e., $Z_i = 0$ for some i). Then by the union bound we deduce

$$
\begin{aligned}
\mathbb{P}(A_n) &\leq \sum_{i=1}^{N} \mathbb{P}(Z_i = 0) \\
&\leq C\epsilon^{-d}(1 - c\epsilon^m)^n \\
&= C \exp\left(n \log(1 - c\epsilon^m) - \log(\epsilon^m) \right) \\
&\leq C \exp\left(-cn\epsilon^m - \log(\epsilon^m) \right).
\end{aligned}
$$

Choose $0 < \epsilon \leq \epsilon_{\mathcal{M}}$ in the form $n\epsilon^m = t \log(n)$ with $t \leq n\epsilon_{\mathcal{M}}^m / \log(n)$. Then

$$
\mathbb{P}(A_n) \leq Ct^{-1} \exp\left(-(ct - 1)\log(n) \right).
$$

In the event that A_n does not occur, then each B_i has at least one point, and so $|x - \sigma_n(x)| \leq C\epsilon$ for all $x \in \mathcal{M}$. Therefore

$$
\|\text{Id} - \sigma_n\|_{L^\infty(\mathcal{M};X)} \leq C\epsilon = C \left(\frac{t \log(n)}{n} \right)^{1/m}
$$

with probability at least $1 - Ct^{-1} \exp\left(-(ct - 1)\log(n) \right)$. Since $\|\text{Id} - \sigma_n\|_{L^\infty(\mathcal{M};X)} \leq C\sqrt{d}$, the result holds for $t \geq n\epsilon_{\mathcal{M}}^m / \log(n)$, albeit with a larger constant C. □

A.2 PROOFS FOR NOISY LABELS

Here, we give the proofs of lemmas and propositions required in Section 2.4. We first give the proof of Lemma 2.13. A key tool in the proof is Bernstein's inequality (Boucheron et al., 2013), which we recall now for the reader's convenience. For $X_1, \ldots, X_n$ *i.i.d.* with variance $\sigma^2 = \mathbb{E}[(X_i - \mathbb{E}[X_i])^2]$, if $|X_i| \leq M$ almost surely for all i then Bernstein's inequality states that for any $\epsilon > 0$

$$
\mathbb{P}\left(\left| \frac{1}{n} \sum_{i=1}^{n} X_i - \mathbb{E}[X_i] \right| > \epsilon \right) \leq 2 \exp\left(-\frac{n\epsilon^2}{2\sigma^2 + 4M\epsilon/3} \right).
$$

Proof of Lemma 2.13. We note that it is sufficient to prove the result for $w \in H_L(X;Y)$ with $\int_{\mathcal{M}} w\rho \, dVol(x) = 0$. In this case, we have $w(x) = 0$ for some $x \in \mathcal{M}$, and so $\|w\|_{L^\infty(X;Y)} \leq CL$. We also write m in place of m in the proof for simplicity.

We first give the proof for $\mathcal{M} = X = [0,1]^m$. We partition X into hypercubes $B_1, \ldots, B_N$ of side length $h > 0$, where $N = h^{-m}$. Let Z_j denote the number of $x_1, \ldots, x_n$ falling in B_j. Then Z_j is a Binomial random variable with parameters n and $p_j = \int_{B_j} \rho \, dx \geq ch^m$. By the Bernstein inequality we have for each j that

$$\mathbb{P}\left(\left| \frac{1}{n} Z_j - \int_{B_j} \rho \, dx \right| > \epsilon \right) \leq 2 \exp\left(-cnh^{-m}\epsilon^2\right) \tag{14}$$

provided $0 < \epsilon \leq h^m$. Therefore, we deduce

$$\frac{1}{n} \sum_{i=1}^{n} w(x_i) \leq \frac{1}{n} \sum_{j=1}^{N} Z_j \max_{B_j} w$$

$$\overset{(14)}{\leq} \sum_{j=1}^{N} \left(\int_{B_j} \rho \, dx + \epsilon \right) \max_{B_j} w$$

$$\leq \sum_{j=1}^{N} \max_{B_j} w \int_{B_j} \rho \, dx + CLh^{-m}\epsilon$$

$$\leq \sum_{j=1}^{N} (\min_{B_j} w + CLh) \int_{B_j} \rho \, dx + CLh^{-m}\epsilon$$

$$\leq \sum_{j=1}^{N} \int_{B_j} w\rho \, dx + CLh^{-m}(h^{m+1} + \epsilon)$$

$$= \int_{X} w\rho \, dx + CL(h + h^{-m}\epsilon)$$

holds with probability at least $1 - 2h^{-m} \exp\left(-cnh^{-m}\epsilon^2\right)$ for any $0 < \epsilon \leq h^m$. Choosing $\epsilon = h^{m+1}$ we have that

$$\left| \frac{1}{n} \sum_{i=1}^{n} w(x_i) - \int_X w\rho \, dx \right| \leq CLh$$

holds for all $u \in H_L(X;Y)$ with probability at least $1 - 2h^{-m} \exp\left(-cnh^{m+2}\right)$, provided $h \leq 1$. By selecting $nh^{m+2} = t\log(n)$

$$\sup_{w \in H_L(X;Y)} \left| \frac{1}{n} \sum_{i=1}^{n} w(x_i) - \int_{\mathcal{M}} w\rho \, dVol(x) \right| \leq CL \left(\frac{t\log(n)}{n} \right)^{\frac{1}{m+2}}$$

holds with probability at least $1 - 2t^{-\frac{m}{m+2}} n^{-(ct-1)}$ for $t \leq n/\log(n)$. Since we have $\|w\|_{L^\infty(X;Y)} \leq CL$, the estimate

$$\sup_{w \in H_L(X;Y)} \left| \frac{1}{n} \sum_{i=1}^{n} w(x_i) - \int_{\mathcal{M}} w\rho \, dVol(x) \right| \leq CL,$$

trivially holds, and hence we can allow $t > n/\log(n)$ as well.

We sketch here how to prove the result on the manifold $\mathcal{M}$. We cover $\mathcal{M}$ with k geodesic balls of radius $\epsilon > 0$, denoted $B_{\mathcal{M}}(x_1, \epsilon), \ldots, B_{\mathcal{M}}(x_k, \epsilon)$, and let $\varphi_1, \ldots, \varphi_k$ be a partition of unity subordinate to this open covering of $\mathcal{M}$. For $\epsilon > 0$ sufficiently small, the Riemannian exponential map $\exp_x : B(0, \epsilon) \subset T_x\mathcal{M} \to \mathcal{M}$ is a diffeomorphism between the ball $B(0, r) \subset T_x\mathcal{M}$ and the geodesic ball $B_{\mathcal{M}}(x, \epsilon) \subset \mathcal{M}$, where $T_x\mathcal{M} \cong \mathbb{R}^m$. Furthermore, the Jacobian of $\exp_x$ at $v \in B(0, r) \subset T_x\mathcal{M}$, denoted by $J_x(v)$, satisfies (by the Rauch Comparison Theorem)

$$(1 + C|v|^2)^{-1} \leq J_x(v) \leq 1 + C|v|^2.$$

Therefore, we can run the argument above on the ball $B(0, r) \subset \mathbb{R}^m$ in the tangent space, lift the result to the geodesic ball $B_{\mathcal{M}}(x_i, \epsilon)$ via the Riemannian exponential map $\exp_x$, and apply the bound

$$\left| \frac{1}{n} \sum_{i=1}^n w(x_i) - \int_{\mathcal{M}} w\rho \, dVol(x) \right| \leq \sum_{j=1}^k \left| \frac{1}{n} \sum_{i=1}^n \varphi_j(x_i) w(x_i) - \int_{\mathcal{M}} \varphi_j w \rho \, dVol(x) \right|$$

to complete the proof. $\qquad\square$

Remark A.2. The exponent $1/(m + 2)$ is not optimal, but affords a very simple proof. It is possible to prove a similar result with the optimal exponent $1/m$ in dimension $m \geq 3$, but the proof is significantly more involved. We refer the reader to (Talagrand, 2006) for details.

Remark A.3. The proof of Theorem 2.11 shows that (1) Γ-converges to (6) almost surely as $n \to \infty$ in the $L^\infty(X; Y)$ topology. Γ-convergence is a notion of convergence for functionals that ensures minimizers along a sequence of functionals converge to a minimizer of the Γ-limit. While we do not use the language of Γ-convergence here, the ideas are present in the proof of Theorem 2.11. We refer to (Braides, 2002) for details on Γ-convergence.

We finally give a proposition useful in the proof of Lemma 2.16.

Proposition A.4. *If ℓ is strongly convex with parameter $\theta > 0$ then*

$$J[tu_1 + (1-t)u_2] + \frac{\theta}{2}t(1-t) \int_{\mathcal{M}} \|u_1 - u_2\|_Y^2 \rho \, dVol(x) \leq tJ[u_1] + (1-t)J[u_2]$$

for all $u_1, u_2 \in W^{1,\infty}(X; Y)$ and $0 \leq t \leq 1$.

Proof. We compute

$$\begin{aligned} &J[tu_1 + (1-t)u_2] \\ &= \int_{\mathcal{M}} \ell(tu_1 + (1-t)u_2, u_0)\rho \, dVol(x) + \lambda \max\left(\mathrm{Lip}(tu_1 + (1-t)u_2), 0\right) \\ &\leq tJ[u_1] + (1-t)J[u_2] - \frac{\theta}{2}t(1-t) \int_{\mathcal{M}} \|u_1 - u_2\|_Y^2 \rho \, dVol(x), \end{aligned}$$

which completes the proof. $\qquad\square$

B   INDUCED MATRIX NORMS

In some cases, we can take advantage of explicit formulas for matrix norms, which makes the estimates in (3) an explicit function of the weights. Define the induced matrix norm by

$$\|M\|_{p,q} = \sup_x \frac{\|Mx\|_q}{\|x\|_p}$$

Then the following matrix norms formulas hold (see (Horn et al., 1990, Chapter 5.6.4))

$$\|M\|_{\infty,\infty} = \max_i \sum_j |m_{ij}|, \qquad \|M\|_{1,1} = \max_j \sum_i |m_{ij}|$$

$$\|M\|_{1,\infty} = \max_{i,j} |m_{ij}|, \quad \|M\|_{2,\infty} = \max_i \sqrt{\sum_j m_{ij}^2}$$

C   VARIATIONAL PROBLEMS IN IMAGE PROCESSING AND LIPSCHITZ
EXTENSIONS

The variational problem (1) can be interpreted as a relaxation of the Lipschitz Extension problem.

$$\begin{cases} \min_{u:X \to Y} \mathrm{Lip}[u] \\ \quad \text{subject to } u(x) = u_0(x) \text{ for } x \in \mathcal{D} \end{cases} \tag{LE}$$

Figure 4: Comparison of different regularization methods. Lipschitz regularization preserves most of the labels (Figure 1). Tychonoff regularization smooths the solution (left). Total Variation regularization shifts the label values towards the mean (right).

for $\mathcal{D} \subset X$. The problem (LE) has more that one solution. Two classical results giving explicit solutions in one dimension go back to Kirzbaum and to McShane (McShane, 1934). However solving (LE) is not practical for large scale problems. There has be extensive work on the Lipschitz Extension problem, see, (Johnson & Lindenstrauss, 1984), for example. More recently, optimal Lipschitz extensions have been studied, with connections to Partial Differential Equations, see (Aronsson et al., 2004). We can interpret (1) as a relaxed version of (LE), where λ^{-1} is a parameter which replaces the unknown Lagrange multiplier for the constraint.

Variational problems are fundamental tools in mathematical approaches to image processing (Aubert & Kornprobst, 2006) and inverse problems more generally. Without regularization inverse problems can be ill-posed. The general form of the problem is

$$J[u] = L[u; u_0] + \lambda R[\nabla u] \tag{15}$$

which combines a loss or *fidelity* functional, $L[u, u_0]$, which depends on the values of u and the reference image u_0, and a *regularization* functional, $R[\nabla u]$, which depends on the gradient, ∇u. The parameter λ determines the relative strength of the two terms which emphasize fidelity versus regularization.

Example C.1. For example, a typical fidelity term is the standard least-squares $L[u, u_0] = \|u - u_0\|_{L^2(D)}^2$. The regularization $\||\nabla u(x)|\|_{L^2(D)}^2$ corresponds to the classical Tychonov regularization (Tikhonov & Arsenin, 1977), $R[\nabla u] = \||\nabla u(x)|\|_{L^1(D)}$ is the Total Variation regularization model of Rudin, Osher and Fatemi (Rudin et al., 1992).

Lipschitz regularization in not nearly as common. It appears in image processing in (Pock et al., 2010, §4.4) (Elion & Vese, 2007) and (Guillot & Le Guyader, 2009). Variational problems of the form (15) can be studied by the direct method in the calculus of variations (Dacorogna, 2007). The problem (15) can be discretized to obtain a finite dimensional convex convex optimization problem. The variational problem can also be studied by finding the first variation, which is a Partial Differential Equation (Evans, 1998), which can then be solved numerically. Both approaches are discussed in (Aubert & Kornprobst, 2006).

In Figure 4 we compare different regularization terms, in one dimension. The difference between the regularizers is more extreme in higher dimensions.