# Integrating Prescriptions into Calendars: Design Considerations

Category: Research

## ABSTRACT

Optimizing the adherence to medical prescriptions remains an ongoing challenge in our health care systems. Among the many reasons for non-adherence (including socioeconomic status) are forgetfulness and the need to manage dynamically evolving busy lives. Electronic medication reminder apps have become broadly available to aid this situation. However, these apps are rarely integrated with other tools used for managing day to day activities, i.e., general electronic Calendars. In this paper, we investigate the design of calendars that can be used for management of medication prescriptions by patients. The proposed designs include mechanisms for visualizing medication entries and communicating attributes related to medication prescriptions. The designs also include ways of annotating conflicts arising from scheduling of medication entries that violate temporal constraints specified in the prescriptions. A user study was conducted to evaluate the calendar designs. Results show that there is potential for the utilization of calendars to manage medication prescriptions.

**Index Terms:** Visualization—Visualization design and evaluation methods—Calendar Design—Prescription Schedules.

## 1 INTRODUCTION

A prescription is a common and important form of medical intervention that is used in modern clinical settings. It comes as a recommendation from an healthcare provider to a patient [1]. It indicates actions such as taking medications, following a diet, or executing physical exercises [2]. When agreed upon between the patient and the healthcare provider, the patient is expected to follow a prescription [3]. The extent to which this implementation corresponds to the agreed upon recommendation is known as *adherence* [2]. Non-adherence to prescriptions is a significant problem in healthcare [2,4]. Adherence rates average 50% and account for 33-69% of hospital re-admissions, resulting into billions of dollars per year [5,6].

Non-adherence is a multidimensional phenomenon, influenced by a complex interplay between socioeconomic, healthcare system-related, medical condition-related, therapy-related and patient-related factors [7]. Patient-related factors of non-adherence include fear of side effects, cost implications, too many medications, failure to perceive benefit, and mistrust of prescribing physicians [8–10]. Non-adherence can also result from forgetfulness and inadvertent omissions that result from failure to make lifestyle changes that take the prescriptions into consideration [11, 12]. Forgetfulness is often addressed through reminders that trigger alarms when a patient is supposed to take medications [6, 10, 13–15]. But when the problem involves lifestyle dynamics, reminders often fall short of addressing the problem. Consider a patient who manages a busy schedule of dynamically changing day-to-day commitments using an application (typically a calendar), while using a separate medication reminder app. What happens when a medication reminder is triggered during the time when the patient is busy? How does that patient reschedule the reminder taking into considerations other activities already planned for? How does one resolve clashes between other activities and medication administration without violating the constraints specified in the prescription? An integrated redesign of these apps is necessary to address these questions.

Electronic calendars have become instrumental in the management of daily activities [16–18]. They are used to coordinate interactions among individual schedules of family or team members and can convey meaning and values behind the priorities of scheduling [19]. Calendars have been used to visualize temporal trends that include everyday activities such as energy use in work places, fitness tracking, and work routines [20–22]. In healthcare, unit-of-use packaging that incorporates a simple day or date feature have been used in efforts to help manage prescriptions and improve adherence by prompting patients to maintain the prescribed dosing schedule [23]. While such prescription managers come as stand-alone tools, we look at the possibility of integrating prescription management into the main calendar already used by the patient. Such integration comes with challenges. The first challenge is how to render the prescription entries so that the user can differentiate between a normal calendar entry and one that is part of a prescription. The second challenge is how to ensure that the patient rescheduling a prescription entry does so within the safety envelop of the prescription.

In this paper we investigate this problem by researching how reminders and medication management can be incorporated into mainstream calendars [24, 25]. Our design goal is to integrate prescriptions into calendars with support for safety checks that avoid (or at least indicate) changes to schedules that no longer adhere to current prescriptions (**DG1**). We consider ways of rendering medication entries so that they are easily identifiable and that all medication-related information is clearly conveyed. We explore ways of rendering unsafe drug-interactions that may arise when medication administration times are being scheduled or changed. This is an important aspect to consider because administration times that conflict with a prescription are considered a form of non-adherence [26]. To our knowledge, incorporating prescriptions and reminders into calendars has not been studied before.

Our contribution is twofold. First, we propose the design of calendars that support medication prescriptions. The calendars allow for the scheduling of medication alongside other everyday activities and provide a way for rendering and resolving scheduling conflicts when they are raised by unsafe drug interactions due to violation of administration constraints specified in the prescription. Second, we present the results of a qualitative study with twelve participants interacting with the calendar designs. Results indicate that calendars can be designed to support the integration of medication prescriptions and that people would generally be favorable to using a calendar to manage their medications, if the design does not deviate from familiar calendars. Results also show that conflicts arising from unsafe rescheduling can be rendered within the calendars. These results inform five additional design goals that an integrated calendar should address: the use of familiar design (DG2), avoiding clutter (DG3), allowing for personalization (DG3), supporting personal reflection (DG5), and highlighting for user attention (DG6).

In the remainder of this paper, we present a review of the relevant literature. We present the data format of medication prescriptions and basic usability requirements for an on-calendar prescription visualization, before introducing three such designs. Then, we report on the study design, methodology and results of these designs before discussing a series of design considerations for integrating prescriptions into calendars.

## 2 RELATED WORK

We review the literature relevant to on-calendar prescription visualization. This includes work related to medication reminders, visualization of schedules, and on-calendar visualizations.

Figure 1: Calendar design with daily medication summaries that giver users a preview (Name and number of times) of the medication to be administered on a particular day.

## 2.1 Medication Reminders

Reminders are among the most common technological interventions to improve adherence to medication [4, 27, 28]. Reminders can take many forms, including interventions of caregivers through video and voice calls [29] and text messages [30], smart pill boxes [10], and computer applications [13]. Focusing on systems that visually represent reminder events to stay within the scope of this research, we identified two relevant technological interventions.

The first one is a health literacy tool called Medication Calendar [14]. It was designed to improve antihypertensive medication adherence. The Medication Calendar provides a graphical view of medication to be taken during a given section of a day. Its layout shows Morning, Afternoon, Evening, and Bedtime as columns and the medications as rows. It displays (in text format) the medication name; the time of day that it should be consumed, the number of times daily that it should be taken; the dose of medication administered; and clinical indication for medication. The application then triggers reminders when it is time to take a medication.

The second relevant intervention is a tool for representing graphically enhanced interventions in coronary heart disease [31] The tool they designed also shows Morning, Afternoon, Evening, and Bedtime as columns, the list of medication as rows; an additional column indicates what the medication is for; andthe row headers include the name of the medication, the dosage, and the time of day that it should be consumed. The novelty in this work is that the cells are filled with graphical images of the corresponding medication.

These two contributions provide foundational design elements to consider for visualizing medications and their reminders in a calendar design. Specifically, the layout (columns and rows) and design concepts (graphical representation of drug entities) provide a starting point for design.

## 2.2 Visualization of schedules

To introduce medication reminders and schedules in general calendar applications, we turn to related work on visualization of schedules. Defined broadly, visualization of schedules involves the representation of planned events on a timeline that depicts a sequence of events (where each event has a single time point) or interval event (continuous quantitative time-series data) [32]. The events are often represented as bars that span multiple time points or nodes that are attached to single time points.

Gantt charts [33] and Pert charts [34] are two standard ways of representing event schedules that have been extended (addressing issues such as visual clutter and event prioritization) to solve planning problems in the field of Engineering (e.g., [35–37]). While not directly about schedule visualization, in the healthcare context a large number of systems have been developed for visual analysis of patient's cohorts temporal data (that also consist of series of temporal events). Lifeline [38, 39], Lifeflow [40], Prima [41] and TimeSpan [42] are some of the works that have explored visualization of timelines to show patients' information such as medication histories, hospital visitations, and treatment processes.

In this research, we focus on i) visualizing a single patient's data, and ii) visualizing events on a calendar – a two-dimensional chart where one dimension shows days of a particular year and the other dimension shows time of day [43]. Therefore we then turn to work on on-calendar visualization.

## 2.3 On-Calendar Visualization

Two dominant models are used to visualize events on a calendar: the radial model and the linear model [44]. With the radial model, data points are positioned along a circle, ellipse, or spiral [45]. Lines are usually drawn from the center of the circle to its circumference at equal adjacent spacing to represent time points. Rings drawn from the center of the circle extending outwards are added to further divide the temporal dimension. Popularized by William Playfair [46] and Florence Nightingale [47], radial calendars have proven effective in visualizing univariate calendar events. Variations of these include Radar Bars [48] and Radar Plots [49]. Radial calendars have been used to show daily consumption of provisions of everyday supplies for a whole calendar year by varying the color and size of marks along the rings [44]; to visualize personal data obtained from different body sensors [50] by positioning stacked bars on a single ring with 24 partitions for a 24-hour day; or quantified-self data [51] by showing streamgraphs and heatmaps of multi-year data along a spiral. Although hailed as state-of-the-art [48] and space-efficient [52], the radial model has limits when it comes to representing multivariate data and data without a defined time limit.

The linear model does not suffer from the same limitations as the radial model. It entails the conventional calendar layout with rows and columns representing days and times of day (or vice versa), and each cell representing a time interval of a particular day [53, 54]. Huang et al. incorporated users' quantitative feedback data, collected from daily activities using Fitbit, within a personal digital calendar [21]. Basing their design on Google calendar [54], they overlaid the calendar with horizontal line graphs to show the person's level of activity each day. Wijk et al. used a linear calendar to show both the power demand and employee presence at a research facility [20] using color hue, height, and color saturation. MineTime Insight [55] is a tool for improving short and long-term scheduling decisions. It uses a calendar design to show multiple coordinated views for the exploration of personal calendar data. This tool however, concentrates on the analysis of calendar data and not on augmenting the calendar with non-calendar data. Research has also investigated shared calendars [56], and more recently, how calendars can be used to enhance team communication and collaboration [57]. In the latter, they augmented a calendar with visualization support for the exploration of conversation data generated by team messaging platforms like Slack. They used color to identify keywords in messages, size to indicate word frequency, and lines to show connections between messages within the channel.

Incorporating non-standard data, such as fitness data or daily activities, on a calendar view has been studied at length [44,50,55–57]. However, these efforts do not fully support the visualization of

medication schedules requirements because most research efforts i) focus on representing univariate and quantitative data; ii) tend to ignore the standard calendar events already present; and iii) usually supports events with two attributes (time and a quantitative attribute) [24, 38, 42]. Visualizing events with a temporal and a quantitative attribute is usually achieved through adding line graphs [21], shading [57], or bars [50], to the calendar, and those are the only visual elements added to the calendar view. Our work differs from this scenario because in our context, events have more than two attributes (allowed placement interval, preferred time, duration, and other descriptive units), are to be presented alongside the normal calendar events, and mechanisms are needed to visually communicate conflicts between events. To the best of our knowledge, there is no existing calendar design that presents a way of visually communicating conflicts that may arise between two or more events, and visualizing prescriptions with their temporal constraints alongside other activities in a standard calendar has not been investigated.

## 3 MEDICATION PRESCRIPTIONS

To design an on-calendar visualization of prescription, we need to i) understand what the characteristics of a prescription are; and ii) what usability requirement such a system should support.

### 3.1 Prescription Characteristics

According to Ethier et al. [58], a prescription is made of the following six building blocks which define the modalities of administration for a given drug: 1) drug prescription item (specifying actions related to one or several drugs), 2) drug administration specification (specifying the drug product), 3) drug course specification (specifying duration, initiation, and termination), 4) drug dosage specification (specifying the dosage of a drug), 5) drug dose administration specification (administration instructions), and 6) drug dispensing specification (specifying the dispensing of a drug product). Kumar et al. [59] summarizes these building blocks as superscription (directive to take), inscription (name and dose), subscription (directions to the pharmacists), and signature (Instructions for Patient), and Fox [60] as drug name, drug dose, drug dose units, drug dose frequency, duration (comprising start and end date), and indication. Like others (e.g. [14, 31]), we adopt the latter classification by Fox [60] because of the simplicity and directness of the naming convention.

Following this classification, the design of an on-calendar prescription visualization should support medication entries with the following attributes: Drug name (D1), Drug dose (D2), Drug dose frequency (D3), Duration (D4), and Drug indications (D5).

As an example of how these data items exist in a prescription, let us consider the following scenario presented by Diemert et al [61].

> *Ms. Smith's physician has prescribed 6mg of Warfarin to be taken orally once daily to address her deep vein thrombosis. Additionally, she was prescribed 15 mg of long-acting Morphine orally twice daily and 600 mg of Ibuprofen to be taken three times a day as needed with food to help manage her pain.*

The scenario above specifies three different medications: Warfarin, Morphine, and Ibuprofen. The classification above describes the prescribed Ibuprofen, for example, with a name (Ibuprofen - D1), dosage (6mg - D2), frequency (three times a day - D3), duration (no end specified - D4), and indication (with food - D5). In the rest of the document, we use the term *medication* more often than the term *drug* because it refers more broadly to an entry in a prescription [61]; it also includes prescription of physical activity, for example.

### 3.2 Usability Requirements

An on-calendar prescription visualization should support tasks that relate to reading calendar entries (e.g., reading calendar events and accessing events' information) and to managing calendar entries (e.g., adding, editing and deleting calendar entries). Because such a system requires adding visual information about prescriptions on top of an existing calendar, the first step in the design process is to understand how to support tasks related to reading calendar entries - that include information about prescriptions and their potential conflicts. Below we introduce basic requirements for reading tasks related to reading calendar entries.

A design that integrates prescription visualization to a calendar must be presented on a calendar. Therefore, such a design should satisfy the following *basic* usability requirement:

> $R_{layout}$: *The user should be able to correctly and efficiently read the calendar's temporal features. These features include the calendar's current day, month and year; the days of the week; and the times of the day.*

The addition of medication entries in the calendar introduces new visual elements. Medication entries also communicate more details (D1 - D5) than the standard title, time, and location of standard calendar events. Therefore, a design that integrates prescription visualization to a calendar should satisfy the following *prescription-related* usability requirement:

> $R_{medic}$: *The user should be able to accurately and efficiently identify the calendar's medication entries. They should also be able to read the entries' name (D1), dosage (D2), frequency (D3), duration (D4), and indications (D5).*

Prescriptions come with constraints associated with drug dosage units, frequency and indications. Constraints can occur within the same medication and between different medications. An example of within-medication constraints is found in the the prescription *Take 600 mg of Ibuprofen three times a day as needed with food*, that has three constraints: 1) that 600mg should be taken at a given time; 2) that the maximum number of intakes per day is three; and 3) that the drug must be taken with food. But medications are often more complex. Consider for example the following prescription: *Take Tenofovir 1 tablet once daily and Metformin 500mg once daily. Take Tenofovir and Metformin at least 6 hours apart.* This example includes within-medication constraints: both Tenofovir and Metformin must be be taken once daily, and the dosage is 1 tablet for the former and 500mg for the latter. This example also includes a between-medication constraint: the two medications should be taken at least 6 hours apart. A violation of constraints (within- or between-medications) is what we call a scheduling conflict. In this paper, we focus on conflicts that deal with restrictions in the scheduling times of medications that may be either unsafe or ineffective when taken together. The nature of these conflicts falls into two categories: too close (those with lower limit time constraints) and too apart (those with upper limit time constraints). Therefore, a design that integrates prescription visualization to a calendar should satisfy the following *conflict-related* usability requirement:

> $R_{conflict}$: *The users should be able to correctly identify conflicting medication entries in the calendar. They should also be able to name the medications involved in the conflict, identify the nature of the conflict, and envision actions that may be taken to resolve the conflict.*

These three usability requirements are basic requirements that an on-calendar prescription visualization should satisfy. Next, we present three calendar designs that satisfy these requirements.

## 4 ON-CALENDAR PRESCRIPTION VISUALIZATION DESIGN

In this section, we propose three designs that show the data (D1 - D5) and meet the usability requirements $R_{layout}$, $R_{medic}$ and $R_{conflict}$.

Figure 2: $Design_A$ with colored vertical bars used for medication entries and gray rectangular entries used for other activities.

Figure 3: $Design_B$ with marked rectangles bearing a colored outline used to represent medication entries. Rectangular entries with dashed outlines represent Physical Activities. Blue full-width rectangular entries show other activities.

## 4.1 Ideation Session

We started the design process with an ideation session with eight researchers with background in human-computer interaction and visualization. The goal of the session was to explore layout variations and features that a calendar that integrates medication prescriptions should have. We asked the researchers to sketch calendars that show all the data (D1 - D5) and satisfy the given requirements. The ideation session lasted 30 minutes.

We identified several design dimensions and their variations from the different sketches that were produced: the layout of the calendar (linear or cyclic), the positioning of the days and times of the day (on the left or at the top), the shape of drug entries (rectangular, cylindrical, or circular), and the orientation of the calendar (vertical or horizontal). Various sizes, colors, and shapes were also used in the designs. Connecting lines were predominantly used to denote the presence (and absence) of conflicts.

We created three designs ($Design_A$, $Design_B$, $Design_C$) by considering i) variations according to these design dimensions, ii) the constraint of compatibility with already existing calendars, and iii) the intention to remain as close as possible to the design of regular medication schedules. The three designs cover a range of design variations regarding layout, representation of medication entries, and representation of conflicts. This will allow us to assess the usefulness of design variations at a component level, rather than at an overall design level. Next, we structure the presentation of these three design variations according to the usability requirements.

## 4.2 Calendar Layout ($R_{layout}$)

We employed the linear layout [44] for all designs. We discarded the radial layout to stay as close to both regular calendars [53, 54] and medication schedules [62] as possible. $Design_A$ and $Design_B$ show the days of the week at the top of the calendar and the time of day on the left, like most digital calendars. $Design_C$ shows the days of the week on the left and time of day at the top, like most medication schedules. All designs showed the day of the week (in abbreviated form), the month of the year and the year.

To differentiate headers from entry cells, headers have a grey background whereas cells have a white background. Each design highlights the current day: $Design_A$ does so with a bold and bigger font size like in MS Outlook [53]), $Design_B$ with a red font color, and $Design_C$ with a grey background like in Google Calendar [54].

## 4.3 Calendar Entries ($R_{medic}$)

All three designs show entries using rectangular shapes like most digital calendars do. However, design variations in terms of position, color and size were explored with each calendar design.

$Design_A$, shown in Figure 2, maintains the layout of existing calendars such as Google Calendar [54] and MS Outlook Calendar [53].

Calendar entries are shown like in standard calendars: the height of rectangular entries indicates their duration, their color hue indicates their type, or category, as set by the user, and their name is conveyed with a label. In this design, we also represent medication entries with rectangles (or bars), whose vertical position and height indicates start and end of the allowed administration period for the medication. Medication entries have an embossed horizontal marker placed at some point along the bar to indicate the preferred administration period (at which point the reminder would trigger if programmed). Preferred administration time of a medication entry is shown with higher opacity and allowed administrative time with lower opacity. Color hue encodes the type of medication.

$Design_A$ supports two sorts of medication entries: drug or physical activity. Each drug entry in the calendar is labelled with the name of the drug and suffixed with bracketed drug dosage. The suffix -WF indicates that the drug should be administered with food. Physical activity entries have a full-color fill, a dashed border, and a label indicating the name of the activity. All other calendar entries are represented with rectangles filled with different shades of grey.

$Design_B$, shown in Figure 3, maintains the layout of existing calendars and maintains the standard way of representing calendar entries. As such, this design consists of not altering the representation of the existing base calendar and its entries, and to *overlay* a representation of medication entries. $Design_B$ has the same layout as $Design_A$. Medication entries, like other entries, are shown with rectangular shapes. They have no fill color, have an outline color hue that conveys the type of medication, a solid outline or a dashed outline if they represent drug or physical activity respectively, and a label that displays the name and dosage of the medication. A filled circle and protruding vertical bar (to illustrate a spoon) in the top-left corner of the rectangle indicates whether the medication should be taken with food or not. A small filled rectangular inset indicates the period of the day when the medication should be administered through its vertical position and height. No marker indicates that the medication should to be taken during the hour on which the entry is positioned; a full-height marker that it can be administered at any time of the day; and a partial-size marker indicates the period of the day that it should be administered (top for morning, middle for afternoon, and bottom for evening). This encoding makes it possible to visually convey larger time periods for administration without cluttering the calendar with very large rectangles.

$Design_C$, shown in Figure 4, has a different layout than $Design_A$ and $Design_B$ as it maintains the layout of medication schedules. It shows days of the week on the left and times of the day in columns at the top, giving a layout with daily entries that read from left to right (as opposed to top to bottom). To each medication entry within a day corresponds one row in the calendar. The first column of a row shows

Figure 4: $Design_C$ with medication entries represented as sliders that communicate allowed administer time and recommended administration time. Each medication entry has its own row. Rectangular entries with and without rugged side-edges represent physical activities and other activities, respectively.

the name of the medication, the number of times the medication should be administered that day, and a green banana icon indicates that it should be administered with food. The medication dosage and administration time are represented with a slider. The position and length of the slider indicate the period in which that medication can be taken. The rectangular buckle indicates the preferred time.

Two additional rows are displayed: one for physical activities, and one for all other, standard calendar entries. They are visually differentiated by using different types of edges. The start and width of a block indicates the start and duration of the activity.

### 4.4 Conflict Visualization ($R_{conflict}$)

Visualization of conflicts between medication entries is shown differently in the three designs, considering i) that conflicts can be of two types (too-apart or too-close) and ii) the results from the ideation session (color and connecting lines were dominantly used to denote conflicts). Figure 5 shows how conflicts are shown in each design.

Some designs featured rectangular enclosures for conflicting pairs. For B and C, we used lines to show conflicts and match conflicting entries because lines do not overwhelm the calendar (and would generally be effective in connecting points as employed in graphs [63]). For A, we matched the conflicts using numbers.

$Design_A$ stays as close to the features already supported in the calendars as possible. As such, we used fill and stroke to denote the presence of conflicts between two entries. An entry that is flagged as being part of a conflict has its block filled with red (for too close) or its outline changed to red (for too apart). Numeric labels indicate the conflict identifier i.e., two entries marked with the same number are part of the same conflict.

$Design_B$ indicates conflicts between entries with connecting red dashed lines – an approach that is effective in showing connection between points as employed in node-link diagrams [63]. Start and end of line markers convey conflict type: a filled circle with a cross mark inset indicates a too-close conflict, and an outlined circle with a cross mark inset indicates a too-apart conflict.

$Design_C$ indicates conflicts using connecting lines like in $Design_B$. However, in this design the line style encodes type of conflict (solid line with arrows pointing outwards for too-close and dotted line with arrows pointing inwards for too-apart). Start and end of line markers are filled circles with a cross mark inside.

## 5 STUDY RATIONALE AND METHODOLOGY

We designed three on-prescription calendars with the aim to study design variations of layout, representation of entries, and representation of conflict. The goal of this study is to assess the quality of design variations, rather than the quality of each design as a whole.

We hosted the three designs on a public server and conducted remote moderated think-aloud sessions via Zoom. Participants were asked to complete a series of tasks with each design variation. The

Figure 5: Conflict representations: (a) filled and outlined entries for $Design_A$, (b) dashed lines with filled and outlined circles for $Design_B$, and (c) directed solid and dotted lines for $Design_C$.

order in which the designs were presented to the participants was counterbalanced and each participant was randomly assigned an order. Each session lasted for about one hour. The study was approved by our institutional Ethics Board.

### 5.1 Participants

Our recruitment was informed by the focus of this research on active people with busy daily-life schedules and who manage medication prescriptions. The recruitment was targeted at people who were between the ages of 35 and 65, who were taking (or helping someone taking) medication regularly, and who considered themselves to have a reasonably busy schedule. We recruited participants through social media platforms, university mailing lists, and posters displayed on campus. Those who were interested reached out to the researcher and were recruited on a first-come-first-serve basis. Participants were compensated with a CAD20 Amazon e-gift card. Twelve participants (P) were recruited and completed the study (10 female, 2 male). They were aged between 35 and 54 years old. All participants were either taking and/or helping someone taking multiple medications: nine were on medication, two were both on medication and helping another person on medication, and one was only helping another person on medication. There were five students, two nurses, three information and communication technology workers, one human resource consultant, and one professor. The frequency with which participants used a calendar to manage their schedule was daily (9), very often (2) and none (1 – this participant indicated frequent use of Medication Administration Record).

### 5.2 Study Procedure

The study began with the participant agreeing to the consent form. Then, they were asked to provide demographic information and were provided a url to access the designs from their browser. They went through a familiarization phase during which they could study the guide (legend), the layout, and the content of the three designs. The participants were encouraged to think aloud throughout the session.

The participants were then asked to complete a series of tasks with each design. The participant performed 16 tasks: 5 addressing for $R_{layout}$, 7 addressing $R_{medic}$, and 4 addressing $R_{conflict}$. Table 1 summarizes these tasks. After they had completed each task, participants were asked whether there were features of the design that they found either useful or not useful to complete that task. After tasks $T_{MED;entry}$ and $T_{MED;day}$ under $R_{medic}$, and tasks $T_{CON;entry}$ and $T_{CON;type}$ under $R_{conflict}$, participants were asked to evaluate the difficulty of performing the task. We picked these tasks because other tasks within each usability requirement are dependent of them. Once a participant had completed all tasks for a given design, they were asked for design suggestions before moving to the next design.

At the end of the session, participants were asked which design they preferred overall and how likely they were to adopt any of the designs for the management of their medications. They were given an opportunity to suggest improvement to each design in order to better suit their Medication management routines and asked if they had any concluding remarks before the session ended.

Table 1: Tasks participants performed with each of the designs, grouped by usability requirement each task addresses. The last two columns indicate which measures were collected for each task.

| Usability Requirement | Task Identifier | Task Description | Correctness | Completion Time |
|---|---|---|---|---|
| $R_{layout}$ | $T_{LAYOUT;year}$ | Reading the current year | x | x |
| | $T_{LAYOUT;month}$ | Reading the current month | x | x |
| | $T_{LAYOUT;week}$ | Locating the day of the week | x | x |
| | $T_{LAYOUT;day}$ | Identifying the current day | x | x |
| | $T_{LAYOUT;hour}$ | Locating the hour of the day | x | x |
| $R_{medic}$ | $T_{MED;entry}$ | Identifying medication and non-medication entries present in the calendar | x | |
| | $T_{MED;day}$ | Reading medication to be administered on a given day | x | x |
| | $T_{MED;repeat}$ | Reading the number of times a medication is repeated in a given day | x | x |
| | $T_{MED;cycle}$ | Reading other days of the week when a given medication was administered | x | x |
| | $T_{MED;dosage}$ | Reading the dosage of a given medication | x | x |
| | $T_{MED;food}$ | Indicating whether a given medication is to be taken with food or not | x | |
| | $T_{MED;slots}$ | Identifying alternative slots for a given medication entry | x | |
| $R_{conflict}$ | $T_{CON;entry}$ | Identifying conflicts that are present in the calendar | x | |
| | $T_{CON;med}$ | Naming the medication involved in the conflict | x | |
| | $T_{CON;type}$ | Determining the type of conflict | x | |
| | $T_{CON;resolve}$ | Suggesting actions for resolving an identified conflict | x | |

## 5.3 Data Collection

We recorded audio and video streams of the sessions. This came from the recording of the interview sessions. The audio component also included three verbally administered questionnaires.

The *demographics questionnaire* was completed at the beginning of the session. The *difficulty questionnaire* asked participants to indicate the difficulty: (i) of differentiating medication entries from non-medication entries ($T_{MED;entry}$), (ii) of reading the medication to be taken at a given time ($T_{MED;day}$), (iii) of identifying the presence of conflicts in the schedule ($T_{CON;entry}$), and (iv) of knowing the nature of the conflict ($T_{CON;type}$). Answers to these questions were provided on a 7-point Likert scale (Very Easy, Easy, Somewhat Easy, Unsure, Somewhat Difficult, Difficult, and Very Difficult). The *adoption questionnaire* was for evaluating the likelihood of using a calendar that employs visualizations to show schedules that include visualization of medication prescriptions alongside other activities. Answers to these questions were provided on a 7-point Likert scale (Very Likely, Likely, Somewhat Likely, Unsure, Somewhat Unlikely, Unlikely, and Very Unlikely).

We recorded the correctness (measure of success), and the completion time for tasks whose answer was succinct. We do not report completion time for tasks that had participants provide descriptive answers that could range from one or two seconds to dozens of seconds, as this would report the time it takes to provide a description and not the time it takes to find the answer to a question.

## 5.4 Data Analysis

We performed a qualitative analysis of participants' think-aloud, reactions, opinions, and suggestions. We transcribed the entire sessions using Otter [64] and used NVivo [65] for the analysis. We employed three rounds of coding: first open coding (identifying any interesting concepts), then selective coding (grouping the concepts), and finally axial coding (relating the concepts) [66].

In the open coding stage, one researcher analyzed the transcribed data and coded the data according to the task addressed. The codes from this round were grouped according to the design referenced.

The second round of coding involved carefully examining the text to identify emerging concepts across designs. A single participant's data was selectively coded by the same researcher to gain insights into the depth of user sentiments on various aspects of the designs. A second researcher who was not involved in the study independently coded the data for the same participant. Notes were then compared

and the categorization and naming conventions used for the codes were harmonized. This stage resulted in 34 codes.

These codes were further analyzed in a round of axial coding to identify and define relationships between them. This analysis resulted into three categories: *temporal features of the calendar*, *design of medication entries*, and *rendering of conflicts*. These findings will be discussed in details under the results section. We structure the report of the study results under these three themes.

To complement the qualitative analysis, we analyzed the quantitative data (correctness and completion time) using descriptive statistics. We used this approach because the small sample size and the small amount of data collected yields too low statistical power to confidently draw conclusions between design variations. We conducted the quantitative analysis using Tableau Software [67] and relied on the median as the measure of central tendency.

## 6 STUDY RESULTS

We first report the study results for each task using completion rate and completion time as well as contextualizing qualitative feedback. Then, we present participant's preferences regarding the different designs and individual design elements.

### 6.1 Task-based Results

We break down the task-based results according to our three task categories: those that deal with the temporal features of the calendar and address $R_{layout}$; those that deal with the design of medication entries and address $R_{medic}$; and those that deal with the rendering and conflicts and address $R_{conflict}$. Figure 6 shows task completion (correctness) for all tasks and for each design. In addition, we report on completion time for all tasks addressing $R_{layout}$ and for four of the seven tasks addressing $R_{medic}$ (see Figure 7).

#### 6.1.1 Reading Temporal Features

Here we present the results for the five tasks that addressed $R_{layout}$. Correctness and completion time were recorded for all five tasks.

$T_{LAYOUT;year}$: All 12 participants successfully performed this task with $Design_A$, 11 with $Design_B$, and 12 with $Design_C$. Median completion time was 1.0 seconds with $Design_A$, 2.5 seconds with $Design_B$, and 1.0 seconds for $Design_C$). No comments were made regarding this task.

Figure 6: A summary of task completion for the three designs A, B, and C. Tasks in $R_{conflict}$ had the most successful completion rate. $T_{MED;slots}$ was the most failed task with Design B recording the lowest score from the entire study.

Figure 7: Box-plots showing task completion times for five layout tasks ($R_{layout}$- Left) and four medication tasks ($R_{medic}$- right) for the three designs. Each box-plot shows the minimum time (lower whisker), first quartile (Q1), median, third Quartile (Q3), and maximum time (Upper Whisker).

$T_{LAYOUT;month}$: 11 participants successfully performed this task with $Design_A$, 12 with $Design_B$, and 9 with $Design_C$. Median completion time was 1.0 seconds with $Design_A$ and $Design_B$, and 2.0 seconds with $Design_C$. Five participants (P3, P4, P6, P8, and P9) commented that showing the month with a number in $Design_C$ made reading the date difficult because they could not tell which number represented the month and which one represented the day. For example, P6 said *"I don't like the fact that March is represented by a number. I find that confusing when I've got two numbers slashed beside each other."*.

$T_{LAYOUT;week}$: All 12 participants successfully performed this task with $Design_A$, $Design_B$, and $Design_C$. Median completion time was 1.5 seconds with $Design_A$ and $Design_B$, and 1.0 seconds with $Design_C$. No comments were made regarding this task.

$T_{LAYOUT;day}$: 10 participants successfully performed this task with $Design_A$, 12 with $Design_B$, and 12 with $Design_C$. Median completion time was 2.0 seconds with $Design_A$, 1.5 seconds with $Design_B$, and 1.0 seconds with $Design_C$. Two participants (P3 and P6) thought the larger font size used to indicate the current date in $Design_A$ was an error. For example, P6 said *"That's why the 21 is so big. I was gonna say there's an error. There's something weird going on with the 21. Maybe Friday was a special day and it gets to be big."* P3 proposed that instead of just showing the current day, there should also be an indication of the current hour, saying *"So right now we're in the 10 o'clock block. So to have maybe a dotted outline around that block, just to let the user know, this is the time of day."* P3 also suggested that as opposed to an indication of the current day on the header, the entire column should be highlighted, and that the header should be floating to remain in place when scrolling.

$T_{LAYOUT;hour}$: All 12 participants successfully performed this task with $Design_A$, $Design_B$, and $Design_C$. Median completion time was 1.0 seconds with $Design_A$, 1.5 seconds with $Design_B$, and 1.0 seconds with $Design_C$. No comments were made regarding this task.

### 6.1.2 Reading Medication Entries

Here we present the results for the seven tasks that addressed $R_{medic}$. Correctness was recorded for all seven tasks, and completion time was recorded for those with non-descriptive answers ($T_{MED;day}$, $T_{MED;repeat}$, $T_{MED;cycle}$, and $T_{MED;dosage}$).

$T_{MED;entry}$: All 12 participants successfully performed this task with $Design_A$, $Design_B$, and $Design_C$. Median completion time was X seconds with $Design_A$, X seconds with $Design_B$, and X seconds with $Design_C$. Once they had provided their answer, participants were asked to describe which visual variables (among color, size, position, and shape) they relied on to provide their answer. Participants relied on color only (6 with $Design_A$, 2 with $Design_B$, 3 with $Design_C$), shape only (3 with $Design_A$, 5 with $Design_B$, 7 with $Design_C$), size and shape (1 with $Design_A$), shape and color (1 with $Design_A$, 4 with $Design_B$), and shape and position (1 with $Design_C$). One participant did not rely on any visual variable to provide their answers with all three designs. P6, P7 and P4 used the name of the medication to identify medication entries in $Design_A$, $Design_B$, and $Design_C$ respectively. For example, P7 said after they had completed the task with $Design_B$, *"I can tell which ones the medication entries are by the names of the medications. I also noted that there was a dosage size or a number of tablets."*. One participant (P9) indicated that the difference between medication and non-medication entries was not clear until they read the legend, saying *"For the first moment it was*

*not so clear. But after [reading the legend] the shape is thinner than the [one for] regular meetings."*

$T_{MED;day}$: All 12 participants successfully performed this task with $Design_A$, $Design_B$, and $Design_C$. Median completion time was 10.0 seconds with $Design_A$ and $Design_B$, and 7.0 seconds with $Design_C$.

To complete this task with $Design_A$, participants had to vertically scroll to the end of the day to see all entries. Three participants (P3, P5, and P7) complained about this. For example, P3 said *"I find it's a lot of scrolling down. It would be helpful if there was a way to condense it or to make it possible to see the entire calendar available in terms of morning, afternoon, and evening."*, and P8 said *"The time frames are a bit big. So it makes like I said, it really makes it scroll off that you can't see it all in one consolidated view"*.

With $Design_B$, participants were expected to use the daily medication summaries provided at the top. Three participants (P1, P6, and P9) found the daily summaries helpful in performing this task. For example, P9 said *"Yeah, I like the idea of having the first row on the calendar dedicated only for the medications that needs to be taken. I think it brings an overall idea [of] what should be taken during that day.".* Three participants (P3, P4, and P5) also complained about the lines demarcating days not being clear. For example, P4 said *"I have a harder time differentiating the calendar component the days, because there's not a strong border between the days of the week.".*

$T_{MED;repeat}$: All 12 participants successfully performed this task with $Design_A$, 11 with $Design_B$, and 12 with $Design_C$. Median completion time was 4.0 seconds with $Design_A$, 2.0 seconds with $Design_B$, and 1.0 seconds with $Design_C$.

To complete this task with $Design_A$, participants had to scroll to count all the times the medication appeared on a given day. Similar complaints about the need to scroll as for $T_{MED;day}$ were made by P3, P5, and P8.

With $Design_B$, participants could count the number of dots beside the name of the medication in the daily summaries at the top. 5 participants (P3, P6, P7, P9, and P10) complained that the dots were not intuitive. For example, P6 said *"So why is there three circles for purple and one circle for green and one circle for Orange? I'm not sure what that means. It's not immediately intuitive"* P3, P7, and P9 thought that the dots indicated the maximum number of pills that was supposed to be taken per day, not the actual number of pills that should be taken. For example, P7 said *"And then these numbers in the circles next to the names of the medicines, I don't know if that means the maximum number of times allowed per day. So the numbers don't really make sense to me."*

No participant had trouble with $Design_C$, where the number was directly indicated beside the name of the medication.

$T_{MED;cycle}$: All 12 participants successfully performed this task with $Design_A$, $Design_B$, and $Design_C$. Median completion time was 10.0 seconds with $Design_A$, 3.0 seconds with $Design_B$, and 4.0 seconds with $Design_C$. With $Design_A$, they again had to scroll through the entire week; with $Design_B$ they could rely on the daily summaries; and with $Design_C$ the information was readily accessible on the row headers. No comments were made regarding this task.

$T_{MED;dosage}$: All 12 participants successfully performed this task with $Design_A$, $Design_B$, and $Design_C$. Median completion time was 3.0 seconds with $Design_A$, 1.5 seconds with $Design_B$, and 2.0 seconds with $Design_C$. Four participants (P1, P4, P7, and P10) commented on the inconsistency in the unit used for the dosage, saying that only milligrams (mg) should be used. For example, P10 said *"It's a little bit inconsistent that it's milligrams of Metformin, but not for the Advil [...] those could be in milligrams too.".*

$T_{MED;food}$: Eight participants successfully performed this task with $Design_A$, 10 with $Design_B$, and 12 with $Design_C$. 10 participants commented that an indicator for take-with-food medication should use a food icon (as used in $Design_C$). For example, with $Design_B$ P3

said *"I'd love to see a symbol that is food-related, as opposed to a slider-style tab.".* P7 indicated that the *WF* (for "With Food") suffix used in $Design_A$ could also be read as "Without Food". P7 also thought that when a food icon is used (such as in $Design_C$), it should be a realistic icon and alternatively be personalizable, saying *"But I would want to make it look more like a banana. If that's supposed to be banana, they're green, but bananas are typically yellow. [...] It'd be fun if someone could choose the emoji they want to use or the picture they want to use to indicate this action".*

$T_{MED;slots}$: Eight participants successfully performed this task with $Design_A$, three with $Design_B$, and 10 with $Design_C$. To complete this task, participants could rely on the bars that indicate allowed medication intake times with $Design_A$ and $Design_C$. With $Design_B$, the marker on the medication entry indicated the allowed time.

As most participants were not successful in completing this task with $Design_B$, they provided ample feedback on this design. Five participants said the design does not support the task, for example, P1 *"it doesn't show any other time of the day that you can take it".* Three participants (P5, P11, and P12) said they can reschedule in any free slot, for example, P9 said *"It seems that 7am is a possibility because there is no other and there is no indication of conflicts.".* P10 commented they would move it to a slot and observe if a conflict was flagged, saying *"I don't know. I think I would just move it and see if a conflict came up.".* P8 participants indicated that use use of the bar to indicate allowed schedule times could also be read as extended release time, saying *"That would mean that it's something that it's an extended release. Warfarin is not an extended release."* P10 and P6 reasoned that a medication that is supposed to be taken at a specific time point should not occupy a full hour on the calendar. For example, P10 said *"I really don't like this fact that it says 6am on the side and then it makes it a block of time."*

### 6.1.3 Reading Conflict Annotations

Here we present the results for the four tasks that addressed $R_{conflict}$. Correctness was recorded for all four tasks, but not completion time because they all asked for descriptive answers. All 12 participants successfully performed $T_{CON;entry}$, $T_{CON;med}$, and $T_{CON;type}$ with $Design_A$, $Design_B$, and $Design_C$. All 12 participants also successfully performed $T_{CON;resolve}$ with $Design_A$ and $Design_C$, and 11 were successful with $Design_B$.

While all the participants managed to identify the conflicts, for some, their representations were not immediately intuitive. For example, P4 said *"At first glance, it did not seem clear. It looked unfriendly just because there's red and axes and stuff. But talking through it with you, and I was trying to explain the differences, it made more sense."* For P8, the fill and outline used for $Design_A$ was not as clear at with the other two designs. They said *"I think switching the coloring doesn't necessarily work to bring out the conflict because you lose the fact that you expect to see this green with blue border for Advil, and now I'm instead seeing this red with green border. Is that a different pill?"* Two participants (P3 and P6) thought the design should not allow users to schedule in slots that would cause conflict, for example P6 said *"You're supposed to take one tablet every four to six hours. They shouldn't have scheduled it there.".* Four participants (P1, P10, P3, and P8) said the designs should show available slots and flag an error when the user tries to schedule a medication entry an illegal slot. For example, P10 said *"So I don't know if there wouldn't be like, the potential spots, if you're going to ask them to move it like where you could have included the constraints to make them blank spots in there."* The four participants (P3, P8, P7, and P10) that noticed that information about the conflict was available when hovering over the conflicting entries with $Design_A$ and $Design_C$ were positive about the feature. For example, P7 said *"when you hover over it, it gives you more information about the conflict that you're having. I do like that."*

Figure 8: Participant's evaluation of the difficulty of performing tasks $T_{MED;entry}$, $T_{MED;day}$, $T_{CON;entry}$, and $T_{CON;type}$ for the three designs.

### 6.1.4 Task Difficulty

Figure 8 shows participant's evaluation of the difficulty of performing tasks $T_{MED;entry}$, $T_{MED;day}$, $T_{CON;entry}$, and $T_{CON;type}$.

## 6.2 Design Preferences

When asked about the likelihood of using a calendar that integrates medication entries, eight answered positively (1 somehow likely, 3 likely, 4 very likely) and four answered negatively (1 somehow unlikely, 1 unlikely, 2 very unlikely). When asked which design they preferred, 1 answered $Design_A$, 10 answered $Design_B$, and 1 answered $Design_C$. Below we present participants' rationale for their reservations and preferences.

Four participants had **reservations** to adopt such a calendar. They provided different reasons for this. P6 wanted to avoid having a constant reminder of medication in their calendar. They said *"I live by calendars. But I have so much going on in my calendar as it is, if I start putting my medication in there as well. It gets way too much emphasis."* P7 was satisfied with their current paper-based system. They said *"Because I have a system that I enjoy using on my own. I have my pills in a pill case. I have a list that I can pull up at any time, that will remind me how many times per day what the dosages per day."* P8 did not see the need of a calendar for medication, because in their case all their medications are to be taken at the same time daily. They said *"I have my medications [which] I take once a day. I make sure that myself, it's the first thing I do when I get up in the morning. It's one of the first activities."*

**Familiarity** (or unfamiliarity) was one reason why participants opted for one design as opposed to another. This was the case for five participants (P7, P6, P5, P4, and P10). For example P7 commented about $Design_C$, *"I haven't actually used a calendar that looked like this in quite some time. And so right now, for me, this isn't a design that's useful"*. P5 said about $Design_B$, *"I think this is more of a calendar view that I'm used to seeing items in slots like this. So that's familiar."*, and P4 said about $Design_B$ *"This one reminds me a little bit more of a Google Calendar, which I'm familiar with."*

**Clutter**, or lack thereof, was another reason for favoring a design raised by 6 participants (P1, P2, P5, P6, P8, and P9) indicated that $Design_A$ and $Design_C$ were too cluttered. For example, P8 said about $Design_C$: *"My first thing seeing it, it's very cluttered. It's got a lot of information on it."* P6 thought that medication entries were given too much space at the expense of other entries, saying *"Well, I don't like again, but there's so much emphasis on the medication, it's almost like the medication takes precedence. And everything else that you've got going on your life is almost insignificant."* P1 suggested toning down medication entries, saying *"instead of having the name of the medication, just put four tablets."* P2 found that too much color was used in $Design_C$, saying *"I think there are a lot of colors here, which makes me confused."* P6 found that the squares used for $Design_A$ could be made smaller, saying *"We [should] make the squares a bit smaller somehow or less predominant, less colorful."*

One (P10) participant raised **privacy** concerns. They suggested that the entries should be discrete enough to avoid exposing information in the event that someone else accessed the calendar. *"So I don't know if it would be something that was like, more discrete or if someone was to accidentally see my calendar and like, saw that I was taking all these medications or something like that."*

## 6.3 Other considerations

Participants described a few features they thought were missing in the designs. These include the timing of reminder, the inclusion of notation for medication that should be taken as needed, and the inclusion of over the counter medication. P8 said that there should be feedback mechanism embedded into the system so that the calendar should confirm that the user has taken the medication. They asked: *"I guess to all this calendar stuff is when you need to take it. And then did you actually get confirmation of taking it?"*

## 7 DISCUSSION AND DESIGN GOALS

Results from our study highlight that prescriptions can be incorporated into mainstream calendars for easy management of medication prescriptions (**DG1**) and that people on multiple prescriptions would be willing to adopt a calendar system that supports the inclusion of medication administration data. Results also show that the design of such a calendar should not deviate from the way conventional calendars are designed and that medication entries should be integrated with other non-medication calendar entries. They should not occupy too much space and should be dismissible by users.

Participants preferred $Design_B$ over $Design_A$ and $Design_C$. $Design_B$ is the design that leverages features from both conventional calendars and prescription schedules. It resulted in better task completion correctness than $Design_A$ and $Design_C$ for layout related tasks ($R_{layout}$). $Design_C$, with its vertical layout, resulted in better correctness than $Design_A$ and $Design_B$ for medication entry related tasks ($R_{medic}$). The three designs led to similar correctness for conflict related tasks ($R_{conflict}$). In terms of task completion time, $Design_A$ and $Design_C$ performed better than $Design_B$ for layout related tasks while $Design_C$ outperformed both $Design_A$ and $Design_B$ for medication entry related tasks.

Below, we discuss the results under the five following themes that emerged from analyzing the qualitative data: use familiar design, avoid clutter, allow for personalization, support personal reflection, and highlight for attention. Each theme informs a new design goals for integrating prescriptions into calendars.

### 7.1 Use Familiar Design

*The design of a calendar should not deviate from calendar interfaces that users are familiar with (**DG2**).* This was observed in various aspects of the design such as layout, medication entries, and icons used to annotate entries such as those which should be taken with food. Over 80% of the participants preferred $Design_B$ because of its probable similitude to already existing calendars. This was surprising to us because $Design_A$ was the design that was intended to resemble existing calendars. The sidelining of $Design_A$ can be attributed mainly to the height of medication entries which spanned the entire allowed administration period. This, according to participants, introduced too much clutter. Results indicate that the preferred layout should be vertically oriented with days of the week at the top and times of the day on the left.

The dosage used on the medication entry should be one that users are familiar with. The unit used should also be consistent with the ones used in the prescriptions. It should show the actual quantity (e.g., milligrams) as opposed to relative classifications such as number of pills or tablets. This observation also applies to the icon used to denote medication that should be taken with food. Realistic

icons that are related to food should be used instead of custom designed icons. In this case, bananas are more effective in indicating the take-with-food action than an icon resembling a utensil such as spoon. Such icon should be positioned together with the entry and not as part of medication summaries.

When dealing with conflicts, arrows are effective in communicating the suggested conflict resolution action. End-of-line arrows can be used to indicate that the medication entries which have been scheduled too close together should be taken apart and vice-versa. The calendar should also have support for indication that a given entry is optional. Such support targets entries such as prescription medication that should be taken as needed and non-prescription medication that is bought over the counter. These design decisions are influenced by everyday activities that users are familiar with.

## 7.2 Avoid Clutter

*The design of the calendar should avoid design elements that introduce clutter to the calendar (DG3).* Design elements such as colors, sliders, labels, and markers should be carefully employed to avoid overwhelming the calendar. One of the reasons why $Design_B$ was preferred is because it is less cluttered: medication entries can be rendered effectively using position, shape, and size. The size of a medication entry should be as small as possible so as not to occupy too much space. Size should not be used to indicate either allowed or preferred administration time of a medication entry, and the size of the entry should also be uniform regardless of the length of the allowed period of administration. Using shapes with a colored outline and transparent fill was associated with less noise and hence preferred by the users. While the slider design was effective in communicating both the allowed and preferred administration period, it made the entry occupy a lot of calendar space and was also misread by some participants as an indicator of delayed release for certain medications. Sliders should thus be avoided. Familiar icons such as tablets can be used to indicate medication entries.

Color should be made less dominant and should not be used as the primary identifier for medication entries. While solid dull color was effective in indicating busy slots, using color fill for medication entries was cluttering the designs. The amount of medication identification information which is visually present in the calendar entry should also be minimized. Medication labels, including name and dosage, tend to overflow the containing entry. Labels were a source of confusion as to which entry they referred to when multiple entries occupied the same cell. They should be abstracted from the overview and instead be made available as details on demand.

Conflict overlays were easily identifiable on all the designs. The user-preferred way to represent such conflicts is to use indicators for the position of medication entries that are involved in the conflict. The connectors for conflicting pairs should avoid thick solid lines, that create clutter. Instead, thin or dotted lines should be employed. While lines are effective in connecting conflicting entries, employing line style to indicate the nature of the conflict is contrariwise. Different line styles may appear similar at a distance and hence fail in communicating the intended message. Line style should be carefully employed to imply an action.

## 7.3 Allow for Personalization

*The design of the calendar should have provision for users to personalize some of its features (DG4).* Such desirable personalizations include adding color to medication entries, choosing icons to be used for medication that should be administered with food, deciding which medication information to display on the entry, and also choosing whether to use the inbuilt entry shape or substitute it for other Emojis or Icons. Personalization should also cover data privacy. The design of the calendar should have features that will protect the users' sensitive data from unauthorized access. This is even more so in situations where the calendar is accessed by more than one person, such as is the case when caregivers are granted access to a calendar. Users should be able to hide features of the calendar that they do not wish anyone without privilege to see.

## 7.4 Support Personal Reflection

*Medication entries should have separate designs for entries that are future and those that are past (DG5).* Past entries should allow for reflection of past medication-taking behavior by confirming whether the user took the medication or not. The design should therefore have a way of letting users confirm that they have taken the medication to actualize personal reflection.

## 7.5 Highlight For Attention

*The calendar should highlight entries to which the users' attention should be drawn on any given day (DG6).* This should start from the calendar layout. The top of the calendar with the year, month, and weekday labels should be floating so that they are always visible when scrolling vertically. Day separators should be clearly defined and the entire current days should be highlighted as opposed to formatting the day's header. The current hour should also be highlighted. The legend did not do much to help participant understand the meaning of entries, as participants were able to read and correctly interpret the design without referring back to the legend. Less emphasis should therefore be placed on the legend. Making the legend available on request is another option.

Medication entries should have a marker that communicates the time that their reminder will be triggered. The marker should not communicate a time range (as we used in all three designs), but a point in time, such as a minute, when the reminder is triggered. The calendar should have daily summaries. These summaries should be used to give an overview of the list of medications to be administered every day. They should only contain the name of the medication. Their design should be such that the user can decide whether to add or remove them from the display. When dealing with conflicts, emphasis should be placed on the position of the conflicting entries and not the connectors. If present, lines that connect conflicting entries should have less emphasis.

On-calendar conflict representation should not be used as the main indication of an error after a rescheduling activity. The user should instead be notified of the impending conflict beforehand. Participants preferred that normal, dismissible error messages be displayed and show the full information regarding the conflicts being introduced by the action. When rescheduling medication entries, cells that are either safe or unsafe should be communicated to the user by highlighting them using a color that implies either success or warning (e.g., green for the former and yellow for the latter). Although some participants felt that the design should not allow them to schedule an entry in the space that is likely to cause a conflict, there might be situations where this possibility would be desirable. The user should, in this case, be guided on possible moves that will resolve the conflict. This can be done by shading or using an outline for all the cells to which an entry may be rescheduled to resolve the conflict, and letting users configure the amount of warnings and error messages they want to receive.

The successful implementation of these design goals will lead not only to a calendar that expressively incorporates all prescription data but to one that is compatible with everyday use of those calendars. Such a calendar would employ design elements that users are familiar with and with little clutter. The calendar would also be personalizable, provide highlighting for principal features, and allow for personal reflection.

## 7.6 Limitations

One limitation of our study is its relatively small sample size. While 12 participants is appropriate for the qualitative analysis of collected data, more participants are required to make task-based statistical

comparisons between designs. That being said, the purpose of this study was exploratory. Findings from this study allow us to then turn to high-fidelity prototyping of calendar designs and to conduct such quantitative task-based study.

Another limitation is that since the study was online, we did not have the privilege of observing participant's full activity cycles. It is likely that remote sessions also lead to participants employing less think-aloud than when participating in person. We also constrained participation to people between the age of 35 and 65 who were either on multiple prescription medications or played the role of caregivers to others on multiple medications. While this allowed us to capture insights for that specific population, these insights do not necessarily generalize to other populations. Our calendar designs were also suited for relatively large screens such as laptops and tablets and were not evaluated on mobile devices. Finally, the study was only limited to tasks that relate to reading calendar entries. In the future, tasks such as adding and modifying medication entries should be included.

## 8 CONCLUSION AND RECOMMENDATIONS

In this research, we explored the possibility of integrating prescriptions into calendars (DG1). We considered ways of rendering medication entries so that they were easily identifiable and that all medication related information was clearly conveyed. We also considered ways of rendering unsafe drug interactions that might arise when medication administration times were being scheduled or changed. We designed three calendars that leverage features available in both conventional calendars and medication schedules. We conducted a study with twelve participants, to evaluate the effectiveness of the calendar designs. Results show that calendars can be designed to support medication prescriptions. Eight out of twelve participants indicated that they were between somehow likely and very likely to use such a calendar to manage their medications. Feedback from the participants indicated that the design of such a calendar should not deviate from familiar calendars. Findings from our study inform five additional design goals that an integrated calendar should address: using of familiar design (DG2), avoiding clutter (DG3), allowing for personalization (DG4), supporting personal reflection (DG5), and highlighting for user attention (DG6). With support for these design goals, people who have busy schedules and are on multiple medications would benefit from having their medication management tasks integrated into their everyday digital calendars.

## REFERENCES

[1] Sujatha Dyasanoor and Ayeesha Urooge. Insight into quality of prescription writing-an instituitional study. *Journal of clinical and diagnostic research: JCDR*, 10(3):ZC61, 2016.

[2] Eduardo Sabaté. Evidence for action World Health Organization 2003. *World Health Organization*, 2003.

[3] Simon Diemert, Jens H Weber, and Morgan Price. An engagement model for medication management: From prescription to description and conscription. *ITCH*, pages 81–86, 2017.

[4] Robby Nieuwlaat, Nancy Wilczynski, Tamara Navarro, Nicholas Hobson, Rebecca Jeffery, Arun Keepanasseril, Thomas Agoritsas, Niraj Mistry, Alfonso Iorio, Susan Jack, et al. Interventions for enhancing medication adherence. *Cochrane database of systematic reviews*, (11), 2014.

[5] Lars Osterberg and Terrence Blaschke. Adherence to medication. *The New England Journal of Medicine*, 353(5):487–497., 2005.

[6] Marie T. Brown, Jennifer Bussell, Suparna Dutta, Katherine Davis, Shelby Strong, and Suja Mathew. Medication Adherence: Truth and Consequences. *American Journal of the Medical Sciences*, 351(4):387–399, 2016.

[7] Neil R. Poulter, Claudio Borghi, Gianfranco Parati, Atul Pathak, Diana Toli, Bryan Williams, and Roland E. Schmieder. Medication adherence in hypertension. *Journal of hypertension*, 38(4):579–587, 2020.

[8] American Medical Association. 8 reasons patients don't take their medications — American Medical Association. https://www.ama-assn.org/delivering-care/patient-support-advocacy/8-reasons-patients-dont-take-their-medications.

[9] Nancy Tran, Janet M. Coffman, Kaharu Sumino, and Michael D. Cabana. Patient reminder systems and asthma medication adherence: A systematic review. *Journal of Asthma*, 51(5):536–543, 2014.

[10] AJ Cross, Ra Elliott, K Petrie, L Kuruvilla, and J George. Adherence in older adults prescribed multiple medications. *Cochrane Database of Systematic Reviews*, 5(5), 2020.

[11] Jia Rong Wu, Debra K. Moser, Terry A. Lennie, and Patricia V. Burkhart. Medication Adherence in Patients Who Have Heart Failure: a Review of the Literature. *Nursing Clinics of North America*, 43(1):133–153, mar 2008.

[12] Lars Osterberg and Terrence Blaschke. Adherence to medication. *Pediatriya*, 55(2):68–69, 2015.

[13] Q Liu, K Abba, Alejandria Mm, D Sinclair, Balanag Vm, and Lansang Mad. Reminder systems to improve patient adherence to tuberculosis clinic appointments for diagnosis and treatment ( Review ) Summary Of Findings For The Main Comparison. *Cochrane Database Syst Rev.*, 11:1–59, 2014.

[14] Leah L. Zullig, Felicia McCant, S. Dee Melnyk, Susanne Danus, and Hayden B. Bosworth. A health literacy pilot intervention to improve medication adherence using Meducation® technology. *Patient Education and Counseling*, 95(2):288–291, may 2014.

[15] Yuhan Luo, Peiyi Liu, and Eun Kyoung Choe. Co-Designing Food Trackers with Dietitians: Identifying Design Opportunities for Food Tracker Customization. *Proceedings of the 2019 CHI Conference on Human Factors in Computing Systems*, 2019.

[16] Carman Neustaedter, AJ Bernheim Brush, and Saul Greenberg. The calendar is crucial: Coordination and awareness through the family calendar. *ACM Transactions on Computer-Human Interaction (TOCHI)*, 16(1):1–48, 2009.

[17] Carman Neustaedter and AJ Bernheim Brush. " linc-ing" the family: the participatory design of an inkable family calendar. In *Proceedings of the SIGCHI conference on Human Factors in computing systems*, pages 141–150, 2006.

[18] Andy Crabtree, Terry Hemmings, Tom Rodden, and John Mariani. Informing the development of calendar systems for domestic use. In *ECSCW 2003*, pages 119–138. Springer, 2003.

[19] Predrag Klasnja, Andrea Hartzler, Christopher Powell, and Wanda Pratt. Supporting cancer patients' unanchored health information management with mobile technology. In *AMIA Annual Symposium Proceedings*, volume 2011, page 732. American Medical Informatics Association, 2011.

[20] Jarke J. van Wijk and Edward R. van Selow. Cluster and calendar based visualization of time series data. *Proceedings of the IEEE Symposium on Information Visualization*, pages 4–9, 1999.

[21] Dandan Huang, Melanie Tory, and Lyn Bartram. A Field Study of On-Calendar Visualizations. *arXiv preprint arXiv:1706.01123.*, 2017.

[22] Earo Wang, Dianne Cook, and Rob J. Hyndman. Calendar-Based Graphics for Visualizing People's Daily Schedules. *Journal of Computational and Graphical Statistics*, 29(3):490–502, jul 2020.

[23] Barbara K. Zedler, Priyanka Kakad, Susan Colilla, Lenn Murrelle, and Nirav R. Shah. Does Packaging with a Calendar Feature Improve Adherence to Self-Administered Medication for Long-Term Use? A Systematic Review. *Clinical Therapeutics*, 33(1):62–73, jan 2011.

[24] Catherine Plaisant, Aaron Clamage, Hilary Browne Hutchinson, Benjamin B. Bederson, and Allison Druin. Shared family calendars. *ACM Transactions on Computer-Human Interaction (TOCHI)*, 13(3):313–346, sep 2006.

[25] Carman Neustaedter, Lyn Bartram, and Aaron Mah. Everyday activities and energy consumption: How families understand the relationship. *Conference on Human Factors in Computing Systems - Proceedings*, pages 1183–1192, 2013.

[26] Marie T. Brown and Jennifer K. Bussell. Medication adherence: WHO cares? *Mayo Clinic Proceedings*, 86(4):304–314, 2011.

[27] Sandra van Dulmen, Emmy Sluijs, Liset Van Dijk, Denise de Ridder, Rob Heerdink, and Jozien Bensing. Patient adherence to medical treatment: a review of reviews. *BMC health services research*, 7(1):1–

13, 2007.

[28] Vicki S Conn, Todd M Ruppar, Maithe Enriquez, and Pam Cooper. Medication adherence interventions that target subjects with adherence problems: systematic review and meta-analysis. *Research in Social and Administrative Pharmacy*, 12(2):218–246, 2016.

[29] Holly B. Jimison, Stuart Hagler, Gregorij Kurillo, Ruzena Bajcsy, and Misha Pavel. Remote health coaching for interactive exercise with older adults in a home environment. *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, EMBS*, 2015-Novem:5485–5488, 2015.

[30] DeKoekkoek Tracy, Given Barbara, Given Charles, Ridenour Kimberly, Schueller Monica, and Spoelstra Sandra. mHealth SMS text messaging interventions and to promote medication adherence: An integrative review. *Journal of Clinical Nursing*, 24(19-20):2722–2735, 2015.

[31] Sunil Kripalani, Brian Schmotzer, and Terry A. Jacobson. Improving Medication Adherence through Graphically Enhanced Interventions in Coronary Heart Disease (IMAGE-CHD): a randomized controlled trial. *Journal of general internal medicine*, 27(12):1609–1617, jul 2012.

[32] Matthew Brehmer, Bongshin Lee, Benjamin Bach, Nathalie Henry Riche, and Tamara Munzner. Timelines Revisited: A Design Space and Considerations for Expressive Storytelling. *IEEE TRANSACTIONS ON VISUALIZATION AND COMPUTER GRAPHICS*, 1, 2016.

[33] Wallace. Clark. The Gantt chart: A working tool of management. *Ronald Press Company*, 1992.

[34] T. I. Kirkpatrick and N. R. Clark. Pert as an Aid to Logic Design. *IBM Journal of Research and Development*, 10(2):135–141, apr 2010.

[35] Alan D. Russell and Asad Udaipurwala. Construction Schedule Visualization. *Congress on Computing in Civil Engineering, Proceedings*, pages 167–178, 2002.

[36] Hyounseok Moon, Nashwan Dawood, and Leenseok Kang. Development of workspace conflict visualization system using 4D object of work schedule. *Advanced Engineering Informatics*, 28(1):50–65, jan 2014.

[37] Jaemin Jo, Jaeseok Huh, Jonghun Park, Bohyoung Kim, and Jinwook Seo. Livegantt: Interactively visualizing a large manufacturing schedule. *IEEE Transactions on Visualization and Computer Graphics*, 20(12):2329–2338, 2014.

[38] C. Plaisant, R. Mushlin, A. Snyder, J. Li, D. Heller, and B. Shneiderman. LifeLines: using visualization to enhance navigation and analysis of patient records. *Proceedings / AMIA ... Annual Symposium. AMIA Symposium*, pages 76–80, 1998.

[39] Brett Milash, Catherine Plaisant, and Anne Rose. LifeLines: visualizing personal histories. *Conference on Human Factors in Computing Systems - Proceedings*, pages 392–393, 1996.

[40] Krist Wongsuphasawat, John Alexis Guerra Gómez, Catherine Plaisant, Taowei David Wang, Shneiderman Ben, and Meirav Taieb-Maimon. LifeFlow: Visualizing an overview of event sequences. *Conference on Human Factors in Computing Systems - Proceedings*, pages 1747–1756, 2011.

[41] D. L. Gresh, D. A. Rabenhorst, A. Shabo, and S. Slavin. PRIMA: A case study of using information visualization techniques for patient record analysis. *Proceedings of the IEEE Visualization Conference*, pages 509–512, 2002.

[42] Mona Hosseinkhani Loorak, Charles Perin, Noreen Kamal, Michael Hill, Mona Hosseinkhani Loorak, Charles Perin, Noreen Kamal, Michael Hill, Sheelagh Carpendale Times, Mona Hosseinkhani Loorak, Charles Perin, Noreen Kamal, Michael Hill, and Sheelagh Carpendale. TimeSpan : Using Visualization to Explore Temporal Multidimensional Data of Stroke Patients To cite this version : HAL Id : hal-01618608 TimeSpan : Using Visualization to Explore Temporal Multi-dimensional Data of Stroke Patients. *IEEE transactions on visualization and computer graphics 22.1*, 2017.

[43] In Merriam-Webster.com. Calendar definition. `https://www.merriam-webster.com/dictionary/calendar`, 2022. Accessed: 2022-05-04.

[44] Catarina Macas and Penousal Machado. Radial calendar of consumption. *Information Visualisation - Biomedical Visualization, Visualisation on Built and Rural Environments and Geometric Modelling and Imaging, IV 2018*, pages 103–109, dec 2018.

[45] Wolfgang Mueller, Marc Weber Marc, and Alexa Wolfgang Müller. Visualizing Time-Series on Spirals. In *IEEE Symposium on Information Visualization*, 2001.

[46] Ian Spence. William Playfair and the psychology of graphs. *Proceedings of the American Statistical Association*, Section on Statistical Graphics:2426–2436, 2006.

[47] Florence Nightingale. Notes on matters affecting the health, efficiency, and hospital administration of the British army: founded chiefly on the experience of the late war. *Harrison and Sons*, St. Martin's Lane, WC, 1987.

[48] Paolo Buono, Maria Francesca Costabile, and Rosa Lanzilotti. A circular visualization of people's activities in distributed teams. *Journal of Visual Languages & Computing*, 25(6):903–911, dec 2014.

[49] M. Joan Saary. Radar plots: a useful way for presenting multivariate health care data. *Journal of Clinical Epidemiology*, 61(4):311–317, apr 2008.

[50] Bruno Pagno and Luciana Nedel. Everyday Visualization. *IEEE VIS 2015 workshop*, 2015.

[51] Charles Perin. Cycles and (A)Symmetry — Exploring the Design of Shareable Personal Visualizations. In IEEE, editor, *IEEE VIS Arts Program Annotated Portfolios*, Berlin, Germany, October 2018. IEEE.

[52] Geoffrey M. Draper, Geoffrey M. Draper, Yarden Livnat, and Richard F. Riesenfeld. A Survey of Radial Methods for Information Visualization. *IEEE transactions on visualization and computer graphics*, 2009.

[53] Microsoft. Outlook calendar. `https://outlook.office.com/calendar`, 2022. Accessed: 2022-05-04.

[54] Google. Google calendar. `https://calendar.google.com/calendar`, 2022. Accessed: 2022-05-04.

[55] Marco Ancona, Marilou Beyeler, Markus Gross, and Tobias Gunther. MineTime Insight: Visualizing Meeting Habits to Promote Informed Scheduling Decisions. *IEEE Transactions on Visualization and Computer Graphics*, 27(3):1986–1999, mar 2021.

[56] Joe Tullio, Jeremy Goecks, Elizabeth D. Mynatt, and David H. Nguyen. Augmenting shared personal calendars. *Proceedings of the 15th annual ACM symposium on User interface software and technology.*, page 11, 2002.

[57] Siwei Fu, Jian Zhao, Hao Fei Cheng, Haiyi Zhu, and Jennifer Marlow. T-Cal: Understanding Team Conversation Data with Calendar-based Visualization. *Proceedings of the 2018 CHI Conference on Human Factors in Computing Systems*, 2018.

[58] Jean François Ethier, François Goyer, Paul Fabry, and Adrien Barton. The Prescription of Drug Ontology 2.0 (PDRO): More Than the Sum of Its Parts. *International Journal of Environmental Research and Public Health 2021, Vol. 18, Page 12025*, 18(22):12025, nov 2021.

[59] Ajay Kumar, Krishna Kumar Pandey, Saurabh Jain, Indu Dangi, Shalini Chowdary, Omprakash Choubitker, and RS Pawar. IDEAL DRUG PRESCRIPTION WRITING. *Ajay et al. World Journal of Pharmacy and Pharmaceutical Sciences*, 8:634, 2019.

[60] Andy Fox. Safer Prescribing Workbook. https://www.uhs.nhs.uk/media/suhtideal /doctors/saferprescribingworkbook/section1-prescriptionwriting.pdf, 2022. Accessed: 2022-06-09.

[61] Diemert Simon. A mathematical basis for Medication Prescription and Adherence. *Journal of Chemical Information and Modeling*, 53(9):1689–1699, 2013.

[62] Sunil Kripalani, Rashanda Robertson, Melissa H Love-Ghaffari, Laura E Henderson, Jessica Praska, Akilah Strawder, Marra G Katz, and Terry A Jacobson. Development of an illustrated medication schedule as a low-literacy patient education tool. *Patient education and counseling*, 66(3):368–377, 2007.

[63] Douglas Brent West et al. *Introduction to graph theory*, volume 2. Prentice hall Upper Saddle River, 2001.

[64] Inc Otter.ai. Otter. `https://otter.ai`, 2022. Accessed: 2022-05-04.

[65] QSR International Pty Ltd. Nvivo.

[66] Jonathan Lazar, Jinjuan Heidi Feng, and Harry Hochheiser. *Research methods in human-computer interaction*. Morgan Kaufmann, 2017.

[67] Tableau Software LLC. Tableau desktop.