Convergence Analysis of Linear Coupling with Inexact Proximal Operator

Abstract

Linear coupling is recently proposed to accelerate first-order algorithms by linking gradient descent and mirror descent together, which is able to achieve the accelerated convergence rate for first-order algorithms. This work focuses on the convergence analysis of linear coupling for convex composite minimization when the proximal operator cannot be exactly computed. It is of particular interest to study the convergence of linear coupling because it not only achieves the accelerated convergence rate for first-order algorithm but also works for generic norms. We present convergence analysis of linear coupling by allowing the proximal operator to be computed up to a certain precision. Our analysis illustrates that the accelerated convergence rate of linear coupling with inexact proximal operator can be preserved if the error sequence of inexact proximal operator decreases in a sufficiently fast rate. More importantly, our analysis leads to better bounds than existing works on inexact proximal operator. Experiment results on several real-world datasets verify our theoretical results.

1 INTRODUCTION

In this work, we consider the convex composite minimization of the form

$$\min_{\mathbf{x} \in \mathbb{R}^d} f(\mathbf{x}) \stackrel{\text{def}}{=} g(\mathbf{x}) + h(\mathbf{x}), \tag{1}$$

where function g is convex or μ-strongly convex, and L-smooth, and h is convex but possibly non-smooth function [Nesterov, 2005]. Various machine learning problems can be formulated in the form of (1), where g defines a convex loss function for training examples, and h regularizes the model to promote a specified structure [Bach et al., 2011, Sra et al.,

2012, Jenatton et al., 2011]. For instance, it is well-known that $h(\mathbf{x}) = \|\mathbf{x}\|_1$ can be used to induce sparse structure for $\mathbf{x}$ [Tibshirani, 1996]. In addition, constrained optimization problem can also be formulated in the form of (1) through reformulation. Specifically, for any convex optimization problem with constraint $\mathbf{x} \in \mathcal{C}$, it can be reformulated as (1) by defining $h(\mathbf{x}) \stackrel{\text{def}}{=} I_{\mathcal{C}}(\mathbf{x})$ as the indicator function of the convex set $\mathcal{C}$, where $I_{\mathcal{C}}(\mathbf{x}) = 0$ if $\mathbf{x} \in \mathcal{C}$ or $I_{\mathcal{C}}(\mathbf{x}) = \infty$ otherwise.

To solve (1), first-order algorithms are the most popular choice due to their simplicity and generality [Sra et al., 2012, Nesterov, 2013]. First-order algorithms generally assume that the first-order gradient of a smooth function can be queried by a black-box in constant time. Therefore, the complexity for solving a smooth and convex function f is measured by the number of times that first-order gradient is queried to produce a sequence $\{\mathbf{x}_k\}_{k=1}^T$ s.t. $f(\mathbf{x}_T) - f(\mathbf{x}^\star) \leq \epsilon$, where $\mathbf{x}^\star$ is the optimal solution. The gradient descent algorithm has $O(L/\epsilon)$ iteration complexity for L-smooth convex minimization problems, which improves to $O(L/\mu \log(1/\epsilon))$ when the function is μ-strongly convex and L-smooth [Nesterov, 2013]. However, these complexities are not optimal, which leaves big space for improvement [Nemirovsky and Yudin, 1983]. In the seminal work [Nesterov, 1983], Nesterov laid the foundation of accelerated gradient methods (AGDs) for convex and L-smooth functions (i.e. $h(\mathbf{x}) = 0$ in (1)). Specifically, the method proposed in [Nesterov, 1983] successfully improves the complexities of gradient descent to $O(\sqrt{L/\epsilon})$ and $O(\sqrt{L/\mu} \log(1/\epsilon))$ for generally and μ-strongly convex and smooth functions, respectively, which are accelerated and optimal for first-order algorithms [Nemirovsky and Yudin, 1983, Nesterov, 2013]. Since then, accelerated first-order algorithms with the optimal convergence rate have been further developed to solve convex composite minimization problems [Tseng, 2008, Lin et al., 2017].

However, the interpretation of the proof of acceleration and intuitions behind the convergence analysis in Nesterov's AGDs are not clear. Many efforts have been devoted to

present clear interpretation for Nesterov's AGDs, or alternatively to develop new accelerated and interpretable variants with the optimal convergence rate [Bubeck et al., 2015, Su et al., 2014, Krichene et al., 2015, Diakonikolas and Orecchia, 2018]. Recently, Allen-Zhu and Orecchia [Allen Zhu and Orecchia, 2017] proved that the optimal convergence rate can be achieved for smooth and convex optimization problems by *linearly coupling* two fundamental first-order algorithms (namely gradient descent and mirror descent). Later, the extension of linear coupling to generally convex composite minimization was presented in [Rodomanov, 2016].

As a variant of Nesterov's AGDs, linear coupling (LC) elegantly extends AGDs to non-Euclidean norms, which is important for many applications. In [Allen Zhu and Orecchia, 2017, Appendix A.1], Allen-Zhu presented several concrete examples to illustrate the importance of allowing non-Euclidean norms in first-order algorithms. For example, one prefers to use the ℓ_1 norm instead of ℓ_2 norm gradient descent for the saddle point problem $\min_{\mathbf{x} \in \Delta_n} \max_{\mathbf{y} \in \Delta_m} \mathbf{y}^\top \mathbf{A} \mathbf{x}$, where Δ_n and Δ_m denotes the unit simplex in $\mathbb{R}^n$ and $\mathbb{R}^m$, respectively, and $\mathbf{A} \in \mathbb{R}^{m \times n}$ with all the entries are between $[-1, 1]$. In this problem, there are two reasons for choosing ℓ_1 norm gradient descent.

1) In order to apply ℓ_2 norm gradient descent, one needs to further assume the square ℓ_2 norm of each row of $\mathbf{A}$ is upper bounded by 1. Obviously, it is a stronger condition and harder to satisfy than that of ℓ_1 norm.

2) More importantly, even the stronger condition is satisfied, the ℓ_2 norm also leads to a larger value for L, thus the ℓ_1 norm gradient descent has faster convergence [Nesterov, 2005]. Another example is near-linear time maximum flow in which one need to apply ℓ_∞ gradient descent [Kelner et al., 2014].

For convex composite optimization problems (1), the proximal gradient descent [Parikh and Boyd, 2014] performs the following updates:

$$\mathbf{x}_{k+1} = \text{prox}_{\eta_{k+1} h} (\mathbf{x}_k - \eta_{k+1} \nabla g(\mathbf{x}_k)),$$

where $\text{prox}_{\eta h}(\cdot)$ is the proximal operator [Combettes and Pesquet, 2011] of $\eta h(\mathbf{x})$ defined for any scalar $\eta > 0$ as the unique solution of

$$\text{prox}_{\eta h}(\mathbf{y}) = \underset{\mathbf{x} \in \mathbb{R}^d}{\text{argmin}} \left\{ \eta h(\mathbf{x}) + \frac{1}{2} \|\mathbf{x} - \mathbf{y}\|^2 \right\}. \quad (2)$$

If h is considerably simple (e.g., $h(\mathbf{x}) = \|\mathbf{x}\|_1$), there is an analytical solution for $\mathbf{x}_{k+1}$ [Combettes and Pesquet, 2011]. However, in more general cases, it is challenging to obtain the exact solution of the proximal operator possibly due to two reasons.

- First, the proximal operator does not admit an analytical solution. For example, there is no closed form solution for the proximal operator if h is the isotropic total variation regularization [Beck and Teboulle, 2009]. In this case, the proximal operator can only be solved by employing some optimization algorithms up to a certain precision. More details of this example can be found in our *Experiments*.

- Second, it might be computationally expensive to obtain the exact solution. For instance, for optimization problems with an ℓ_1 norm ball constraint (e.g., $h(\mathbf{x}) = I_{\|\mathbf{x}\|_1 \leq r}(\mathbf{x}), r \in \mathbb{R}_+$), the complexity of exactly performing the proximal operator is $O(d \log d)$ [Duchi et al., 2008, Liu and Ye, 2009]. Therefore, it is highly demanding in computation for high-dimensional learning problems as d is considerably large. Indeed, we empirically compare the efficiency of exact and inexact proximal operator of ℓ_1 norm ball constraint. The result suggests that inexact proximal operator outperforms the exact counterpart by carefully controlling the error sequences in this application.

Therefore, it is of particular interest to characterize the convergence behavior of linear coupling, when the proximal operator is not exactly solved. However, existing analyses [Schmidt et al., 2011] on inexact proximal operator only cover the ℓ_2 norm, thus it does not applicable to linear coupling due to non-Euclidean norm. To this end, we present a complete study for the convergence rate of *linear coupling* with inexact proximal operator.

Compared with existing works [Schmidt et al., 2011, Lin et al., 2017, Kulunchakov and Mairal, 2019], our focus is the convergence analysis of linear coupling with inexact proximal operator, which presents new challenges due to the generic Bregman divergence (refer to Definition 3). In particular, the key step for analyzing inexact proximal operator is to bound the subgradient of inexact solution. It is an easy task in the case of squared Euclidean distance as the bounding problem has an analytical solution [Schmidt et al., 2011]. In contrast, it does not admit an analytical form in our case due to the generic Bregman divergence (refer to Definition 1), e.g., Kullback-Leibler divergence. To address it, we present a relaxation method for the bounding problem so that the subgradient still can be bounded (see Lemma 1). More importantly, our analysis leads to tighter bounds (refer to Remarks 2 and 5 for details) than previous works [Schmidt et al., 2011, Lin et al., 2017, Kulunchakov and Mairal, 2019].

2 NOTATION AND PRELIMINARIES

Throughout this paper, vectors and matrices are denoted by lower-case and upper-case boldface characters (e.g., $\mathbf{x}$ and $\mathbf{X}$), respectively. Let $\mathbf{0}$ be a vector or matrix with all its entries equal to 0. For both $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$, their inner product

is denoted by $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^{d} x_i y_i$. Let $\|\cdot\|$ be a generic norm and its dual norm is denoted by $\|\cdot\|_*$ that is defined as $\|\mathbf{y}\|_* = \sup_{\mathbf{x}} \{\langle \mathbf{x}, \mathbf{y} \rangle \mid \|\mathbf{x}\| \leq 1\}$. Let $\|\cdot\|_1$ and $\|\cdot\|_2$ denote the ℓ_1 and ℓ_2 norm, respectively.

Definition 1. *A function f is L-smooth w.r.t. $\|\cdot\|$, if*

$$f(\mathbf{y}) \leq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{L}{2}\|\mathbf{y} - \mathbf{x}\|^2, \ \forall \mathbf{x}, \mathbf{y}.$$

Definition 2. *A function f is μ-strongly convex w.r.t. $\|\cdot\|$, if*

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{\mu}{2}\|\mathbf{y} - \mathbf{x}\|^2, \ \forall \mathbf{x}, \mathbf{y}.$$

Definition 3. *Let $\psi : \mathcal{Q} \to \mathbb{R}$ be a strictly convex and continuously differentiable function. Then, the Bregman divergence is*

$$V_\psi(\mathbf{y}, \mathbf{x}) \stackrel{\text{def}}{=} \psi(\mathbf{y}) - \psi(\mathbf{x}) - \langle \nabla \psi(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle, \forall \mathbf{x}, \mathbf{y} \in \mathcal{Q}.$$

The Definition 3 implies $V_\psi(\mathbf{x}, \mathbf{x}) = 0$, and $V_\psi(\mathbf{y}, \mathbf{x}) \geq \frac{\rho}{2}\|\mathbf{x} - \mathbf{y}\|^2$ if ψ is ρ-strongly convex w.r.t. $\|\cdot\|$. The Bregman divergence includes many well-known examples.

- 1) If $\psi(\mathbf{x}) \stackrel{\text{def}}{=} \frac{1}{2}\|\mathbf{x}\|_2^2$, then $V_\psi(\mathbf{y}, \mathbf{x})$ is the squared Euclidean distance $V_\psi(\mathbf{y}, \mathbf{x}) = \frac{1}{2}\|\mathbf{x} - \mathbf{y}\|_2^2$

- 2) If $\mathcal{Q} \stackrel{\text{def}}{=} \{\mathbf{x} \in \mathbb{R}_+^d \mid \sum_i x_i = 1\}$ and $\psi(\mathbf{x}) \stackrel{\text{def}}{=} \sum_i x_i \log x_i$, then $V_\psi(\mathbf{y}, \mathbf{x})$ becomes the Kullback-Leibler divergence $V_\psi(\mathbf{y}, \mathbf{x}) = \sum_i y_i \log(\frac{y_i}{x_i})$ between two probability distributions $\mathbf{x}$ and $\mathbf{y}$. In particular, $\psi(\mathbf{x})$ in this case is 1-strongly convex w.r.t. $\|\cdot\|_1$ that leads to $V_\psi(\mathbf{y}, \mathbf{x}) \geq \|\mathbf{x} - \mathbf{y}\|_1^2$. In this case, one needs to employ optimization methods that are applicable for ℓ_1 norm. Therefore, this example illustrates the importance of convergence analysis for linear coupling with inexact proximal operator since it works for a generic norm.

Definition 4. *For function $f(\mathbf{x})$, its convex conjugate is defined as*

$$f^*(\mathbf{y}) \stackrel{\text{def}}{=} \sup_{\mathbf{x}} \{\langle \mathbf{x}, \mathbf{y} \rangle - f(\mathbf{x})\}.$$

3 LINEAR COUPLING WITH INEXACT PROXIMAL OPERATOR

For smooth and convex functions, the accelerated convergence rate can be obtained by two non-accelerated algorithms: linearly coupling gradient descent and mirror descent [Allen Zhu and Orecchia, 2017]. In fact, it can also be extended to solve convex composite minimization problem (1) [Rodomanov, 2016]. We assume that g is convex ($\mu = 0$) or μ-strongly convex ($\mu > 0$), and L-smooth w.r.t. $\|\cdot\|$. Here, we present the extension of linear coupling (LC) for (1) when g is either convex ($\mu = 0$) or strongly convex

---

Algorithm 1 LC with Inexact Proximal Operator

1: **Input**: $\mathbf{x}_0, \alpha_0, \mu, L$
2: **Initialization**: $\mathbf{y}_0 \leftarrow \mathbf{x}_0, \mathbf{z}_0 \leftarrow \mathbf{x}_0$
3: **for** $k = 0$ **to** $T - 1$ **do**
4:    **if** $\mu = 0$ **then**
5:       Set $\eta_{k+1} \leftarrow \frac{k+2}{2L}$ and $\tau_k \leftarrow \frac{1}{L\eta_{k+1}}$
6:       Set $\mathbf{w}_{k+1} \leftarrow \mathbf{z}_k$
7:    **else**
8:       Set $\eta_{k+1} \leftarrow \frac{1}{L\alpha_{k+1}}$ and $\tau_k \leftarrow \frac{L\alpha_{k+1} - \mu}{L - \mu}$ where α_{k+1} is obtained via $\alpha_{k+1}^2 = (1 - \alpha_{k+1})\alpha_k^2 + \frac{\mu}{L}\alpha_{k+1}$
9:       Set $\mathbf{w}_{k+1} \leftarrow \frac{\tau_k}{\alpha_{k+1}}\mathbf{z}_k + \left(1 - \frac{\tau_k}{\alpha_{k+1}}\right)\mathbf{y}_k$
10:    **end if**
11:    $\mathbf{x}_{k+1} \leftarrow \tau_k \mathbf{z}_k + (1 - \tau_k)\mathbf{y}_k$
12:    Find a ξ_{k+1}-suboptimal solution $\mathbf{y}_{k+1}$ for (3)
13:    Find a ξ_{k+1}-suboptimal solution $\mathbf{z}_{k+1}$ for (4)
14: **end for**
15: **Output**: $\mathbf{y}_T$

($\mu > 0$) and summarize the high-level idea in Algorithm 1. Missing proofs can be found in the Appendix.

Let $\mathbf{y}_k$ and $\mathbf{z}_k$ be the outputs of gradient descent and mirror descent in the $(k-1)$-th iteration, respectively. The key idea of linear coupling is to combine $\mathbf{y}_k$ and $\mathbf{z}_k$ together by a linear coupling rate τ_k as the starting point for the next iteration such that the accelerated convergence rate can be achieved.

Define $\mathbf{x}_{k+1} \stackrel{\text{def}}{=} \tau_k \mathbf{z}_k + (1 - \tau_k)\mathbf{y}_k$. In linear coupling, the gradient descent performs the following update:

$$\mathbf{y}_{k+1} = \operatorname*{argmin}_{\mathbf{y}} \widetilde{Q}_{k+1}(\mathbf{y}; \mathbf{x}_{k+1}), \tag{3}$$

where $\widetilde{Q}_{k+1}(\mathbf{y}; \mathbf{x}_{k+1})$ is define as

$$\widetilde{Q}_{k+1}(\mathbf{y}; \mathbf{x}_{k+1}) = \langle \nabla g(\mathbf{x}_{k+1}), \mathbf{y} \rangle + h(\mathbf{y}) + \frac{L}{2}\|\mathbf{y} - \mathbf{x}_{k+1}\|^2.$$

Define η_{k+1}, τ_k and $\mathbf{w}_{k+1}$ as in Algorithm 1, the mirror descent performs the following update:

$$\mathbf{z}_{k+1} = \operatorname*{argmin}_{\mathbf{x}} \widehat{Q}_{k+1}(\mathbf{z}; \mathbf{w}_{k+1}), \tag{4}$$

where

$$\widehat{Q}_{k+1}(\mathbf{z}; \mathbf{w}_{k+1}) \stackrel{\text{def}}{=} \langle \nabla g(\mathbf{x}_{k+1}), \mathbf{z} \rangle + h(\mathbf{z}) + \frac{V_\psi(\mathbf{z}, \mathbf{w}_{k+1})}{\eta_{k+1}}.$$

Unlike standard gradient descent and mirror descent, the linear coupling takes the gradient at $\mathbf{x}_{k+1}$ instead of $\mathbf{y}_k$ or $\mathbf{z}_k$ to obtain $\mathbf{y}_{k+1}$ and $\mathbf{z}_{k+1}$. In this way, the gradient descent and mirror descent are coupled together to solve (1), which is able to achieve the optimal convergence rate Allen Zhu and Orecchia [2017].

3.1 THE ξ_{k+1}-SUBOPTIMAL SOLUTION

The proximal operator is involved in both (3) and (4) due to the non-smooth h. It is worth noting that the proximal distance in both (3) and (4) are defined based a generic norm $\|\cdot\|$ instead of $\|\cdot\|_2$. Specifically, they are the squared norm $\|\cdot\|^2$ and the general Bregman divergence $V_\psi(\cdot, \cdot)$, respectively. In contrast, existing works mainly focus on the case of squared Euclidean distance (ℓ_2 norm). Therefore, (3) and (4) are more challenging to solve than existing works. In other words, it is often the case that (3) and (4) can only be solved up to a certain precision. Therefore, it is critical to study the convergence rate of Algorithm 1 by allowing the proximal operator of the form (3) and (4) to be solved approximately.

For the case of inexact proximal operator, we introduce ξ_{k+1}-suboptimal solution to analyze the convergence rate of Algorithm 1. We assume that h is equipped with an oracle such that the proximal operator can be computed up to a certain precision. Specifically, given a non-negative ξ_{k+1}, the oracle is able to produce $\mathbf{y}_{k+1}$ and $\mathbf{z}_{k+1}$ such that

$$\widetilde{Q}_{k+1}(\mathbf{y}_{k+1}; \mathbf{x}_{k+1}) - \min_{\mathbf{y}} \widetilde{Q}_{k+1}(\mathbf{y}; \mathbf{x}_{k+1}) \leq \xi_{k+1}, \quad (5)$$

$$\widehat{Q}_{k+1}(\mathbf{z}_{k+1}; \mathbf{w}_{k+1}) - \min_{\mathbf{z}} \widehat{Q}_{k+1}(\mathbf{z}; \mathbf{w}_{k+1}) \leq \xi_{k+1}. \quad (6)$$

If $\xi_{k+1} = 0$, it implies that the proximal operator is exactly solved for both (3) and (4). Otherwise, it means that the proximal operator is solved up to a certain precision controlled by ξ_{k+1}. Thus, $\mathbf{y}_{k+1}$ and $\mathbf{z}_{k+1}$ are referred to as ξ_{k+1}-suboptimal solutions to (3) and (4), respectively.

We note that the inexactness criteria (5) and (6) are same as that used in [Schmidt et al., 2011, Lin et al., 2017, Kulunchakov and Mairal, 2019]. Whereas such a criterion has limitations, it remains the most standard one for convergence analysis with inexact proximal operator.

Our analysis allows the sub-problems (3) and (4) do not admit closed-form solutions, therefore one needs to compute an approximate solution up to a certain accuracy with some iterative algorithm $\mathcal{M}$. Note that the sub-problems both (3) and (4) are strongly convex even the objective $f(\mathbf{x})$ is not strongly convex. The strong convexity allows us to efficiently obtain a ξ_{k+1}-suboptimal solution via $\mathcal{M}$ with a linear convergence rate [Lin et al., 2017, Kulunchakov and Mairal, 2019].

4 CONVERGENCE ANALYSIS

In this section, we present the convergence analysis for Algorithm 1 when apply it to solve (1). Specifically, we first present properties of a suboptimal solution in Section 4.1. Then, in Sections 4.2 and 4.3, we present specific convergence results of Algorithm 1 for $\mu = 0$ and $\mu > 0$, in Theorems 1 and 2, respectively. For convenience, we also

assume the $\psi(\mathbf{x})$ is 1-strongly convex and ρ-smooth w.r.t. $\|\cdot\|$.

4.1 PROPERTIES OF SUBOPTIMAL SOLUTION

For convergence analysis with inexact proximal operator, the key is to bound the solution of inexact proximal operator and the ξ-subgradient of the inexact solution. Thus, we first introduce the definition of ξ-subgradient, which is a generalization of subgradient.

Definition 5. *[Bertsekas et al., 2003] For convex function $f : \mathbb{R}^d \to \mathbb{R}$ and a non-negative scalar ξ, $\partial_\xi f(\mathbf{x})$ is the ξ-subgradient of f at $\mathbf{x}$ if it holds that, $\forall \mathbf{v} \in \partial_\xi f(\mathbf{x})$,*

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \langle \mathbf{v}, \mathbf{y} - \mathbf{x} \rangle - \xi, \forall \mathbf{y} \in \mathbb{R}^d. \quad (7)$$

Definition 5 implies that $\mathbf{0}$ is a ξ-subgradient of f at $\mathbf{x}$ if $\mathbf{x}$ is a ξ-suboptimal solution of f. In the case of ℓ_2 norm, Schmidt et al. [2011] showed that $\mathbf{v}$ in Definition 5 can be easily bounded as the squared Euclidean distance in (2) leads to an analytical form for $\mathbf{v}$. In contrast, our case is more challenging as it involves $(\nabla \psi)^{-1}$ which generally does not has an analytical form for generic Bregman divergence. To address this problem, we present a relaxation method by exploiting the strongly convexity of ψ so that it still admit an analytical form. We define

$$Q_{k+1}(\mathbf{z}; \mathbf{w}_{k+1}) \overset{\text{def}}{=} \langle \nabla g(\mathbf{x}_{k+1}), \mathbf{z} - \mathbf{w}_{k+1} \rangle + h(\mathbf{z}).$$

If $\mathbf{z}_{k+1}$ is ξ_{k+1}-suboptimal to (4), Lemma 1 provides a ξ_{k+1}-subgradient for $Q_{k+1}(\cdot; \mathbf{w}_{k+1})$ at $\mathbf{z}_{k+1}$.

Lemma 1. *For $\forall k \geq 0$, if $\mathbf{z}_{k+1}$ is a ξ_{k+1}-suboptimal solution to (4) in the sense of (6), then there exists $\boldsymbol{\beta}_{k+1}$ with $\|\boldsymbol{\beta}_{k+1}\|_*^2 \leq 2\rho\xi_{k+1}/\eta_{k+1}$ such that*

$$\frac{\nabla\psi(\mathbf{w}_{k+1}) - \nabla\psi(\mathbf{z}_{k+1})}{\eta_{k+1}} - \boldsymbol{\beta}_{k+1} \in \partial_{\xi_{k+1}} Q_{k+1}(\mathbf{z}_{k+1}; \mathbf{w}_{k+1}).$$

The proof of Lemma 1 is given in Appendix A.1. By using Lemma 1, the following lemma enables us to bound the intermediate results of mirror descent with inexact proximal operator.

Lemma 2. *Under the same setting as in Lemma 1, then there exists $\boldsymbol{\beta}_{k+1}$ with $\|\boldsymbol{\beta}_{k+1}\|_*^2 \leq 2\rho\xi_{k+1}/\eta_{k+1}$ such that*

$$\widehat{Q}_{k+1}(\mathbf{z}_{k+1}; \mathbf{w}_{k+1}) + V_\psi(\mathbf{u}, \mathbf{z}_{k+1})/\eta_{k+1} - \xi_{k+1}$$
$$\leq \widehat{Q}_{k+1}(\mathbf{u}; \mathbf{w}_{k+1}) + \langle \boldsymbol{\beta}_{k+1}, \mathbf{u} - \mathbf{z}_{k+1} \rangle, \forall \mathbf{u}, k \geq 0.$$

The proof of Lemma 2 is given in Section A.2.

4.2 CONVERGENCE RATES OF CONVEX g

We first focus on the case when g is convex. The next lemma derives a characteristic inequality for a specific Lyapunov

function for inexact linear coupling by considering the inexactness of $\mathbf{y}_{k+1}$ and $\mathbf{z}_{k+1}$.

Lemma 3. *Under the same setting as in Lemma 1, for any $k \geq 0$, if g is convex and $\tau_k = 1/L\eta_{k+1}$,*

$$\frac{1}{\tau_k^2}\left(f(\mathbf{y}_{k+1}) - f(\mathbf{x}^\star)\right) + LV_\psi(\mathbf{x}^\star, \mathbf{z}_{k+1})$$
$$\leq \frac{1 - \tau_k}{\tau_k^2}\left(f(\mathbf{y}_k) - f(\mathbf{x}^\star)\right) + LV_\psi(\mathbf{x}^\star, \mathbf{z}_k)$$
$$+ \sqrt{\frac{2\rho L\xi_{k+1}}{\tau_k}}\|\mathbf{x}^\star - \mathbf{z}_{k+1}\| + \frac{1 + \tau_k}{\tau_k^2}\xi_{k+1}. \quad (8)$$

The proof of Lemma 3 is given in Appendix A.3. From the Lyapunov function, we obtain a general convergence result for linear coupling with inexact proximal operator.

Theorem 1. *Under the same setting as in Lemma 1, if g is convex, $\eta_{k+1} = (k+2)/2L$ and $\tau_k = 1/L\eta_{k+1}, \forall k \geq 0$, then $\forall T \geq 1$:*

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq \frac{6\left(LV_\psi(\mathbf{x}^\star, \mathbf{x}_0) + \widetilde{E}_T + \widehat{E}_T\right)}{(T+1)^2}, \quad (9)$$

where $\widetilde{E}_T \stackrel{\text{def}}{=} \sum_{k=1}^T (k+2)^2\xi_k$ and $\widehat{E}_T \stackrel{\text{def}}{=} \left(\sum_{k=1}^T \sqrt{2\rho(k+1)\xi_k}\right)^2$.

The proof of Theorem 1 is given in Appendix A.4.

Remark 1. *If the proximal operator is exact (i.e., $\xi_k = 0, \forall k \geq 1$), it leads to $\widetilde{E}_T = 0$ and $\widehat{E}_T = 0$. Then, Theorem 1 recoveries the accelerated complexity $O(\sqrt{L/\epsilon})$ for convex objectives [Nesterov, 2013].*

Corollary 1. *Consider the same setting as Theorem 1, if $\xi_k \leq \xi$ for all $k \geq 1$, then $\forall T \geq 1$:*

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq \frac{6LV_\psi(\mathbf{x}^\star, \mathbf{x}_0)}{(T+1)^2} + (\theta_1 T + \theta_2)\xi. \quad (10)$$

where $\theta_1 = (6 + 16\rho)$ and $\theta_2 = 15 + 32\rho$.

Remark 2. *Under the same setting as Corollary 1, the [Schmidt et al., 2011, Proposition 4] gives*

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq \frac{6L\|\mathbf{x}_0 - \mathbf{x}^\star\|^2}{(T+1)^2} + \left(12T^2 + 4T + 2\right)\xi. \quad (11)$$

The [Lin et al., 2017, Theorem 3] gives

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq \frac{4L\|\mathbf{x}_0 - \mathbf{x}^\star\|^2}{(T+1)^2} + \frac{9}{2}(T+2)^2\xi. \quad (12)$$

The [Kulunchakov and Mairal, 2019, Proposition 4] gives

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq \frac{2e^{1+\gamma}L\|\mathbf{x}_0 - \mathbf{x}^\star\|^2}{(T+1)^2} + \frac{e^{1+\gamma}}{2\gamma}(T+2)^2\xi. \quad (13)$$

Comparing (10) *with* (11), (12) *and* (13), *our analysis achieves better bound than existing works [Schmidt et al., 2011, Lin et al., 2017, Kulunchakov and Mairal, 2019]. Specifically, our bound on the inexact proximal operator ξ increases as $O(T\xi)$ while it is $O(T^2\xi)$ for existing works.*

Theorem 1 suggests that the inexact proximal operator leads to error accumulation in the convergence result. To preserve the accelerated rate $O(\frac{1}{T^2})$, Theorem 1 implies the error sequence should decrease to 0 at a sufficiently fast rate.

Corollary 2. *Under the same setting as Theorem 1, for any $\delta > 0$, if ξ_k is chosen as $\xi_k \stackrel{\text{def}}{=} \frac{f(\mathbf{x}_0) - f(\mathbf{x}^\star)}{(k+2)^{3+\delta}}$, then $\forall T \geq 1$,*

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq \frac{LV_\psi(\mathbf{x}^\star, \mathbf{x}_0)}{(T+1)^2}\left(1 + \frac{2}{\delta} + \frac{8\rho}{\delta^2}\right). \quad (14)$$

Remark 3. *To preserve the accelerated rate $O(1/k^2)$ for convex objectives, existing works [Schmidt et al., 2011, Lin et al., 2017, Kulunchakov and Mairal, 2019] requires the error sequences decreases at the rate of $O(1/k^{4+\delta})$. In contrast, our analysis suggests $O(1/k^{3+\delta})$ is sufficient. This is consistent with our tighter bound shown in Corollary 1.*

As observed in (14), the objective value converges faster for more accurate proximal operator (i.e., a larger value for δ). However, larger δ also requires more computation time for each iteration.

Comparison with Alternating Direction Method of Multipliers (ADMM). Besides the accelerated proximal method, the ADMM is another popular method for solving (1) due to it is easy to implement [Boyd et al., 2011]. However, it is well-know that the ADMM converges as the rate of $O(1/T)$ for convex objectives [He and Yuan, 2012]. In contrast, the convergence rate of accelerated proximal method is $O(1/T^2)$, that is more desirable for large-scale machine learning problems.

As discussed in [Boyd et al., 2011], the ADMM will converge even the when the subproblems of each iteration are not solved exactly, as long as the approximate solutions satisfy certain suboptimality measures. In other words, the ADMM also suffer from error accumulation if the subproblems cannot be exactly solved, for example, the proximal operator does not admit a closed-form solution. We note our inexactness conditions (5) and (6) are absolute criteria that are same as the one used in [Schmidt et al., 2011, Lin et al., 2017, Kulunchakov and Mairal, 2019]. In contrast, existing works on convergence analysis of the ADMM are mainly based on relative error accuracy [He et al., 2002, Eckstein and Yao, 2018, Alves et al., 2020] that is generally a stronger inexactness condition than ours. If the ADMM takes the inexactness criterion as ours, under the same setting as Corollary 1, one can show that the convergence rate of inexact ADMM is

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq \frac{\nu\|\mathbf{x} - \mathbf{x}^\star\|^2}{T} + \gamma T\xi,$$

where ν and γ are some constants. Comparing it with (10), we observed both ADMM and our result have $O(T\xi)$ error accumulation, however, our result achieves $O(1/T^2)$ convergence rate while it is only $O(1/T)$ for ADMM.

4.3 CONVERGENCE RATES OF STRONGLY CONVEX g

In the last section, we present the convergence result of linear coupling with inexact proximal operator for convex g. Next, we present the convergence result for strongly convex g. As is customary [Nesterov, 2013, Cohen et al., 2018], in this setting we assume that $\|\cdot\| = \|\cdot\|_2$.

We first introduce the analogue of Lemma 3 for strongly convex g. By considering the inexactness of $\mathbf{y}_{k+1}$ and $\mathbf{z}_{k+1}$, the next lemma derives a characteristic inequality for a specific Lyapunov function for strongly convex objectives.

Lemma 4. *Under the same setting as Lemma 1, if g is strongly convex, $\tau_k = \frac{L\alpha_{k+1}-\mu}{L-\mu}$ and $\eta_{k+1} = \frac{1}{L\alpha_{k+1}}$, then $\forall k \geq 0$,*

$$f(\mathbf{y}_{k+1}) - f(\mathbf{x}^\star) + L\alpha_{k+1}^2 V_\psi(\mathbf{x}^\star, \mathbf{z}_{k+1}) \qquad (15)$$
$$\leq \left(1 - \alpha_{k+1}\right)\left(f(\mathbf{y}_k) - f(\mathbf{x}^\star) + L\alpha_k^2 V_\psi(\mathbf{x}^\star, \mathbf{z}_k)\right)$$
$$+ \sqrt{2\rho L \alpha_{k+1}^3 \xi_{k+1}}\|\mathbf{x}^\star - \mathbf{z}_{k+1}\| + \left(1 + \alpha_{k+1}\right)\xi_{k+1}.$$

The proof of Lemma 4 is given in Appendix A.8. From the Lyapunov function, we obtain a general convergence result for linear coupling with inexact proximal operator.

We define $\Delta_k \stackrel{\text{def}}{=} f(\mathbf{y}_k) - f(\mathbf{x}^\star) + L\alpha_k^2 V_\psi\left(\mathbf{x}^\star, \mathbf{z}_k\right), \forall k \geq 0$. Theorem 2 presents the convergence result of linear coupling with inexact proximal operator for strongly convex g.

Theorem 2. *Under the same setting as Lemma 1, if g is strongly convex, $\tau_k = \frac{L\alpha_{k+1}-\mu}{L-\mu}$ and $\eta_{k+1} = \frac{1}{L\alpha_{k+1}}$, then $\forall T \geq 1$:*

$$\Delta_T \leq \Gamma_T \left(\Delta_0 + \sum_{k=1}^{T} \frac{\sqrt{2\rho L \alpha_k^3 \xi_k}\|\mathbf{x}^\star - \mathbf{z}_k\| + 2\xi_k}{\Gamma_k}\right), \quad (16)$$

where $\Gamma_k \stackrel{\text{def}}{=} \prod_{i=1}^{k}(1 - \alpha_k)$. If $\alpha_0 = \sqrt{\frac{\mu}{L}}$, then $\forall T \geq 1$:

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq \left(1 - \sqrt{\frac{\mu}{L}}\right)^T \left(3\widetilde{\Delta}_0 + \widetilde{R}_T + \widehat{R}_T\right), \quad (17)$$

where $\widetilde{\Delta}_0 \stackrel{\text{def}}{=} f(\mathbf{x}_0) - f(\mathbf{x}^\star)$, $\widetilde{R}_T \stackrel{\text{def}}{=} 3\sum_{k=1}^{T}\left(1 - \sqrt{\frac{\mu}{L}}\right)^{-k}\xi_k$ and $\widehat{R}_T \stackrel{\text{def}}{=} 6\rho\sqrt{\frac{\mu}{L}}\left(\sum_{k=1}^{T}\left(1 - \sqrt{\frac{\mu}{L}}\right)^{-k/2}\sqrt{\xi_k}\right)^2$.

The proof of Theorem 2 is given in Appendix A.9.

Remark 4. *If the proximal operator is exact (i.e., $\xi_k = 0, \forall k \geq 1$), it leads to $\widetilde{R}_T = 0$ and $\widehat{R}_T = 0$. Then, (17) recoveries the accelerated complexity $O(\sqrt{L/\mu}\log(1/\epsilon))$ for μ-strongly convex objectives [Nesterov, 2013].*

Corollary 3. *Consider the same setting as Theorem 2, if $\xi_k \leq \xi$ for all $k \geq 1$, then $\forall T \geq 1$:*

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq 3\left(1 - \sqrt{\frac{\mu}{L}}\right)^T \widetilde{\Delta}_0 + (3+24\rho)\sqrt{\frac{L}{\mu}}\xi. \quad (18)$$

Remark 5. *Under the same setting as Corollary 3, the [Schmidt et al., 2011, Proposition 4] gives*

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq 4\left(1 - \sqrt{\frac{\mu}{L}}\right)^T \widetilde{\Delta}_0 + \left(\frac{64L^2}{\mu^2} + 4\sqrt{\frac{\mu}{L}}\right)\xi. \quad (19)$$

The [Lin et al., 2017, Theorem 3] gives

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq 4\left(1 - \sqrt{\frac{\mu}{L}}\right)^T \widetilde{\Delta}_0 + \frac{72L}{\mu}\xi. \quad (20)$$

The [Kulunchakov and Mairal, 2019, Proposition 4] gives

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq 2\left(1 - \frac{1}{2}\sqrt{\frac{\mu}{L}}\right)^T \widetilde{\Delta}_0 + \frac{8L}{\mu}\xi. \quad (21)$$

Comparing (18) with (19), (20) and (21), our analysis achieves better bound than existing works. Specifically, our error bound on ξ is $O(\sqrt{\frac{L}{\mu}}\xi)$, while that are $O(\frac{L^2}{\mu^2}\xi)$ for [Schmidt et al., 2011] and $O(\frac{L}{\mu}\xi)$ for [Lin et al., 2017, Kulunchakov and Mairal, 2019], respectively.

To preserve accelerated rate for Algorithm 1, Theorem 2 implies that the error sequence $\{\xi_k\}_{k\geq 1}$ needs to decrease to 0 at a linear rate.

Corollary 4. *Under the same setting as in Theorem 2, for any $\vartheta \in (0, \sqrt{\mu/L})$, if ξ_k is chosen as $\xi_k \leq \frac{1}{1+2\rho}\widetilde{\Delta}_0(1-\vartheta)^k$, then $\forall T \geq 1$*

$$f(\mathbf{y}_T) - f(\mathbf{x}^\star) \leq (1-\vartheta)^{T+1}\frac{12\widetilde{\Delta}_0}{(\sqrt{\mu/L}-\vartheta)^2}. \quad (22)$$

Similar to the conclusion for convex g, the objective with strongly convex g has a faster convergence speed when the error of inexact proximal operator decreases in a faster rate (i.e., a larger value for ρ).

5 EXPERIMENTS

In this section, we perform two experiments to verify our theoretical results.

5.1 CUR-LIKE FACTORIZATION

We first apply Algorithm 1 to solve the CUR-like factorization optimization problem [Mairal et al., 2011]. For a given

Figure 1: Results of linear coupling and ADMM with inexact proximal operator for CUR-like factorization. Objective function values v.s. number of iterations for different qualities of approximate solution of the proximal operator. From left to right: `a1a`, `secom` and `mushroom`.

matrix $\mathbf{D} \in \mathbb{R}^{m \times n}$, the CUR-like factorization aims to approximate $\mathbf{D}$ by a matrix $\mathbf{X}$ with sparse rows and sparse columns.

$$\min_{\mathbf{X} \in \mathbb{R}^{m \times n}} \frac{1}{2} \|\mathbf{DXD} - \mathbf{D}\|_F^2 + \lambda_1 \sum_{i=1}^{m} \|\mathbf{X}_{i,\cdot}\|_2 + \lambda_2 \sum_{j=1}^{n} \|\mathbf{X}_{\cdot,j}\|_2,$$

where $\mathbf{X}_{i\cdot}$ and $\mathbf{X}_{\cdot,j}$ denote the i-th row and j-th column of $\mathbf{X}$, respectively. The last two terms impose the $\ell_{2,1}$ norms for both row and columns of $\mathbf{X}$, that yields both sparse rows and columns. However, the proximal operator of this regularizer does not admit an analytical solution, even no iterative algorithm can exactly compute the proximity operator.

Following [Schmidt et al., 2011], we approximately compute the proximal operator by a block coordinate descent (BCD) algorithm that is presented by Jenatton et al. [2011] to efficiently obtain an approximate solution of the proximal operator. The BCD alternates between computing the proximal operator with respect to the rows and to the columns. At each iteration, we employ the BCD to solve (3) and (4) until (5) and (6) are satisfied that means both $\mathbf{y}_k$ and $\mathbf{z}_k$ are ξ_k-suboptimal solutions.

Suggested by Corollary 2, we consider a decreased error sequences $\{\xi_k\}_{k \geq 1}$ where $\xi_k \leq 1/k^\alpha$ and the value of α are set to $\alpha = 1, 2, 3, 4$ in our experiments. As discussed before, the accelerated convergence rate of linear coupling can be preserved if $\alpha > 3$.

We perform experiments on four data sets[1]: `mushroom`, `secom`, `a1a` and `musk`. We set $\lambda_1 = 0.01$ and $\lambda_2 = 0.01$ for all four datasets. Rather than assume the Lipschitz constant L is known, we estimate it by line search [Nesterov, 2013]. Specifically, we initialize the value of L as $L = 0.5$ and double it if the following inequality is not satisfied

$$g(\mathbf{y}_k) \leq g(\mathbf{x}_k) + \langle \nabla g(\mathbf{x}_k), \mathbf{y}_k - \mathbf{x}_k \rangle + \frac{L}{2} \|\mathbf{y}_k - \mathbf{x}_k\|^2.$$

[1] The datasets can be downloaded at https://archive.ics.uci.edu/ml/datasets.php.

In our experiments, we observed that this strategy always performs better than a fixed but conservative value for L.

To demonstrate the accelerated convergence rate of linear coupling, we compare it with ADMM. It is well-know that the performance of ADMM is highly dependent on the choice of penalty parameter ϱ. To show the best performance of ADMM, we perform a grid search to find the best value of ϱ and fixed it for all iterations.

Fig. 1 shows the objective function values versus the number of iterations of inexact linear coupling on the three data sets: `a1a`, `secom` and `mushroom`. For linear coupling, the choice of $\xi_k \leq 1/k^4$ achieves the fastest convergence rate according to our analysis (refer to Corollary 2), provides the best empirical performance across all three data sets. However, as the iteration goes, the performance gap between the choices of $\xi_k \leq 1/k^3$ and $\xi_k \leq 1/k^4$ becomes smaller. This is consistent with our theoretical results that the error sequences only need to decrease faster than $O(1/k^3)$ instead of $O(1/k^4)$ [Schmidt et al., 2011, Lin et al., 2017, Kulunchakov and Mairal, 2019].

Furthermore, as observed from Fig. 1, the linear coupling clearly outperforms the ADMM on this task. Here, we show the results of ADMM with $\xi_k \leq 1/k^3$ and $\xi_k \leq 1/k^4$. In fact, we observed the result of ADMM when $\alpha \geq 2$ are very close to each other, that is consistent with the theoretical result. Specifically, the error sequences should decrease faster than $O(1/k^2)$ if the convergence rate of the algorithm is $O(1/k)$, for example, non-accelerated proximal method.

5.2 IMAGE DEBLURRING WITH ISOTROPIC TOTAL VARIATION

For the second experiment, we consider image deblurring with isotropic total variation regularization [Chambolle, 2004, Beck and Teboulle, 2009], which does not admit an analytical solution for the proximal operator [Chambolle and Pock, 2011, Beck and Teboulle, 2009]. For $\mathbf{X} \in \mathbb{R}^{m \times n}$,

Figure 2: Results of linear coupling with inexact proximal operator for image deblurring with isotropic total variation. Objective function values v.s. number of iterations for different qualities of approximate solution of the proximal operator. From left to right: `Lena`, `Boat` and `Football`.

the discrete gradient operator is

$$
(\nabla \mathbf{X})_{ij} = \begin{cases} (X_{ij}-X_{i+1,j}, X_{ij}-X_{i,j+1}) & \text{if } i < m, j < n \\ (0, X_{ij}-X_{i,j+1}) & \text{if } i = m, j < n \\ (X_{ij}-X_{i+1,j}, 0) & \text{if } i < m, j = n \end{cases}
$$

Then, image deblurring with isotropic total variation regularization can be written as [Chambolle and Pock, 2016, Beck and Teboulle, 2009]

$$
\min_{\mathbf{X} \in \mathcal{C}} \frac{1}{2} \|\mathcal{A}(\mathbf{X}) - \mathbf{B}\|_{\mathrm{F}}^2 + \lambda \sum_{i=1}^{m} \sum_{j=1}^{n} \|(\nabla \mathbf{X})_{ij}\|,
$$

where $\mathcal{A} : \mathbb{R}^{m \times n} \to \mathbb{R}^{m \times n}$ is a linear operator representing some blurring processing, $\mathbf{B} \in \mathbb{R}^{m \times n}$ is the blurred and noisy image, and $\lambda > 0$ denotes a regularization parameter. Unlike the ℓ_1 total variation, the proximal operator of isotropic total variation regularization does not admit an analytical solution. Same as before, we employ the BCD algorithm to compute an approximate solution.

We conducted this experiment on three images `Lena`, `Boat` and `Football` from MATLAB image processing toolbox. We first resize each image to 128×128 pixels. To obtain $\mathbf{B}$, the clean image was first blurred by a 5×5 kernel matrix: $\mathbf{S} = \frac{1}{25}\mathbf{I}_{5 \times 5}$, where $\mathbf{I}_{5 \times 5} \in \mathbb{R}^{5 \times 5}$ is an identify matrix, followed by additive Gaussian noise with zero mean and standard deviation 10^{-1}. The regularization parameter λ was set to 0.1 for all three images.

Fig. 2 shows the objective function values versus the number of iterations of inexact linear coupling on the`Lena`, `Boat` and `Football`. Similar trends as the first experiments are observed for this experiment except the performance between $\xi_k \le 1/k^3$ and $\xi_k \le 1/k^4$ are very close to each other.

In this experiment, we did not perform the experiments of ADMM due to it solves the subproblem associated with $\frac{1}{2}\|\mathcal{A}(\mathbf{X})-\mathbf{B}\|_F^2$ is computationally expensive. Nevertheless, the experiment in Section 5.1 already clearly shows the advantages of linear coupling over ADMM.

6 CONCLUSION

Non-smooth regularizations have played irreplaceable roles in machine learning. However, many of them do not admit an analytical solution for proximal operator. As a variant of Nesterov's AGDs, the linear coupling can efficiently solve convex composite minimization with accelerated convergence rates if the proximal operator is exactly computed. To further understand the performance of linear coupling, we present a complete convergence analysis for it with inexact proximal operator. Our analysis suggests that inexact linear coupling still achieves the accelerated convergence rate if the error sequence of inexact proximal operator decreases in a sufficiently fast rate. More importantly, our theoretical results is better than previous works. We empirically verify our theoretical analysis by employing linear coupling with inexact operator to solve CUR-like factorization and image deblurring with isotropic total variation on different datasets. Empirical results clearly support our theoretical analysis.

References

Zeyuan Allen Zhu and Lorenzo Orecchia. Linear coupling: An ultimate unification of gradient and mirror descent. In *Proceedings of the 8th Innovations in Theoretical Computer Science Conference (ITCS)*, volume 67, pages 3:1–3:22, 2017.

Maicon Marques Alves, Jonathan Eckstein, Marina Geremia, and Jefferson G. Melo. Relative-error inertial-relaxed inexact versions of douglas-rachford and ADMM splitting algorithms. *Computational Optimization and Applications*, 75(2):389–422, 2020.

Francis R. Bach, Rodolphe Jenatton, Julien Mairal, and Guillaume Obozinski. Convex optimization with sparsity-inducing norms. *Optimization for Machine Learning*, 170:19–53, 2011.

Amir Beck and Marc Teboulle. Fast gradient-based algorithms for constrained total variation image denoising and deblurring problems. *IEEE Transactions on Image Processing (TIP)*, 18(11):2419–2434, 2009.

Dimitri P. Bertsekas, Angelia Nedi, and Asuman E. Ozdaglar. *Convex Analysis and Optimization*. Athena Scientific, 2003.

Stephen P. Boyd, Neal Parikh, Eric Chu, Borja Peleato, and Jonathan Eckstein. Distributed optimization and statistical learning via the alternating direction method of multipliers. *Foundations and Trends in Machine Learning*, 3 (1):1–122, 2011.

Sébastien Bubeck, Yin Tat Lee, and Mohit Singh. A geometric alternative to nesterov's accelerated gradient descent. *CoRR*, abs/1506.08187, 2015.

Antonin Chambolle. An algorithm for total variation minimization and applications. *Journal of Mathematical Imaging and Vision*, 20(1-2):89–97, 2004.

Antonin Chambolle and Thomas Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. *Journal of Mathematical Imaging and Vision*, 40(1):120–145, 2011.

Antonin Chambolle and Thomas Pock. An introduction to continuous optimization for imaging. *Acta Numerica*, 25: 161–319, 2016.

Michael Cohen, Jelena Diakonikolas, and Lorenzo Orecchia. On acceleration with noise-corrupted gradients. In *Proceedings of the 35th International Conference on Machine Learning (ICML)*, volume 80, pages 1018–1027, 2018.

Patrick L. Combettes and Jean-Christophe Pesquet. Proximal splitting methods in signal processing. In *Fixed-Point Algorithms for Inverse Problems in Science and Engineering*, pages 185–212. 2011.

Jelena Diakonikolas and Lorenzo Orecchia. Accelerated extra-gradient descent: A novel accelerated first-order method. In *Proceedings of the 9th Innovations in Theoretical Computer Science Conference (ITCS)*, volume 94, pages 23:1–23:19, 2018.

John C. Duchi, Shai Shalev-Shwartz, Yoram Singer, and Tushar Chandra. Efficient projections onto the l_1-ball for learning in high dimensions. In *Proceedings of the 25th International Conference on Machine Learning (ICML)*, volume 307, pages 272–279, 2008.

Jonathan Eckstein and Wang Yao. Relative-error approximate versions of douglas-rachford splitting and special cases of the ADMM. *Mathematical Programming*, 170 (2):417–444, 2018.

Bingsheng He and Xiaoming Yuan. On the o(1/n) convergence rate of the douglas-rachford alternating direction method. *SIAM Journal on Numerical Analysis (SINUM)*, 50(2):700–709, 2012.

Bingsheng He, Li-Zhi Liao, Deren Han, and Hai Yang. A new inexact alternating directions method for monotone variational inequalities. *Mathematical Programming*, 92 (1):103–118, 2002.

Rodolphe Jenatton, Jean-Yves Audibert, and Francis R. Bach. Structured variable selection with sparsity-inducing norms. *Journal of Machine Learning Research (JMLR)*, 12:2777–2824, 2011.

Jonathan A. Kelner, Yin Tat Lee, Lorenzo Orecchia, and Aaron Sidford. An almost-linear-time algorithm for approximate max flow in undirected graphs, and its multicommodity generalizations. In *Proceedings of the 25th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 217–226, 2014.

Walid Krichene, Alexandre M. Bayen, and Peter L. Bartlett. Accelerated mirror descent in continuous and discrete time. In *Proceedings of the 29th Annual Conference on Neural Information Processing Systems (NIPS)*, pages 2845–2853, 2015.

Andrei Kulunchakov and Julien Mairal. A generic acceleration framework for stochastic composite optimization. In *Proceedings of the 33rd Annual Conference on Neural Information Processing Systems (NeurIPS)*, pages 12556–12567, 2019.

Hongzhou Lin, Julien Mairal, and Zaïd Harchaoui. Catalyst acceleration for first-order convex optimization: from theory to practice. *Journal of Machine Learning Research (JMLR)*, 18:212:1–212:54, 2017.

Jun Liu and Jieping Ye. Efficient euclidean projections in linear time. In *Proceedings of the 26th International Conference on Machine Learning (ICML)*, volume 382, pages 657–664, 2009.

Julien Mairal, Rodolphe Jenatton, Guillaume Obozinski, and Francis R. Bach. Convex and network flow optimization for structured sparsity. *Journal of Machine Learning Research (JMLR)*, 12:2681–2720, 2011.

Arkadii Nemirovsky and David Yudin. *Problem Complexity and Method Efficiency in Optimization*. Wiley Interscience Series in Discrete Mathematics, 1983.

Yurii Nesterov. A method for solving the convex programming problem with convergence rate $O(1/k^2)$. *Doklady AN SSSR (translated as Soviet Mathematics Doklady)*, 269:543–547, 1983.

Yurii Nesterov. Smooth minimization of non-smooth functions. *Mathematical Programming*, 140(1):127–152, 2005.

Yurii Nesterov. *Introductory lectures on convex optimization: A basic course*. Springer Science & Business Media, 2013.

Neal Parikh and Stephen P. Boyd. Proximal algorithms. *Foundations and Trends in Optimization*, 1(3):127–239, 2014.

Anton Rodomanov. Linear coupling of gradient and mirror descent: Version for composite functions with adaptive estimation of the lipschitz constant. 2016. URL `https://bayesgroup.github.io`.

Mark W. Schmidt, Nicolas Le Roux, and Francis R. Bach. Convergence rates of inexact proximal-gradient methods for convex optimization. In *Proceedings of the 25th Annual Conference on Neural Information Processing Systems (NIPS)*, pages 1458–1466, 2011.

Suvrit Sra, Sebastian Nowozin, and Stephen J. Wright. *Optimization for Machine Learning*. MIT Press, 2012.

Weijie Su, Stephen P. Boyd, and Emmanuel J. Candès. A differential equation for modeling nesterov's accelerated gradient method: Theory and insights. In *Proceedings of the 28th Annual Conference on Neural Information Processing Systems (NIPS)*, pages 2510–2518, 2014.

Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society. Series B*, 58(1):267–288, 1996.

Paul Tseng. On accelerated proximal gradient methods for convex-concave optimization. *Submitted to SIAM Journal on Optimization (SIOPT)*, 2008.