Feature-Robustness, Flatness and Generalization Error for Deep Neural Networks

Abstract

The performance of deep neural networks is often attributed to their automated, task-related feature construction. It remains an open question, though, why this leads to solutions with good generalization, even in cases where the number of parameters is larger than the number of samples. Back in the 90s, Hochreiter and Schmidhuber observed that flatness of the loss surface around a local minimum correlates with low generalization error. For several flatness measures, this correlation has been empirically validated. However, it has recently been shown that existing measures of flatness cannot theoretically be related to generalization: if a network uses ReLU activations, the network function can be reparameterized without changing its output in such a way that flatness is changed almost arbitrarily. This paper proposes a natural modification of existing flatness measures that results in invariance to reparameterization. The proposed measures imply a robustness of the network to changes in the input and the hidden layers. Connecting this feature robustness to generalization leads to a generalized definition of the representativeness of data . With this, the generalization error of a model trained on representative data can be bounded by its feature robustness which depends on our novel flatness measure.

1 Introduction

Neural networks (NNs) have become the state of the art machine learning approach in many applications. An explanation for their superior performance is attributed to their ability to automatically learn suitable features from data. In supervised learning, these features are learned implicitly through minimizing the **empirical error** $\mathcal{E}_{emp}(f, S) = 1/|S| \sum_{(x,y) \in S} \ell(f(x), y)$ for a **training set** $S \subset \mathcal{X} \times \mathcal{Y}$ drawn iid according to a **target distribution** $\mathcal{D} : \mathcal{X} \times \mathcal{Y} \to [0, 1]$, and a **loss function** $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}_+$. Here, $f : \mathcal{X} \to \mathcal{Y}$ denotes the function represented by a neural network.

It is an open question why minimizing the empirical error during deep neural network training leads to good generalization, even though in many cases the number of network parameters is higher than the number of training examples. That is, why deep neural networks have a low **generalization error**

$$\mathcal{E}_{gen} = \mathbb{E}_{(x,y) \sim \mathcal{D}} \left[\ell(f(x), y) \right] - \frac{1}{|S|} \sum_{(x,y) \in S} \ell(f(x), y) \tag{1}$$

which is the difference between expected error on the target distribution $\mathcal{D}$ and the empirical error on a finite dataset $S \subset \mathcal{X} \times \mathcal{Y}$.

It has been proposed that good generalization correlates with flat minima of the non-convex loss surface (Hochreiter & Schmidhuber, 1997; 1995) and this correlation has been empirically validated (Keskar et al., 2016; Novak et al., 2018; Wang et al., 2018). Thus, for deep neural networks trained with stochastic gradient descent (SGD), this could present a (partial) explanation for their generalization performance (Zhang et al., 2016), since minibatch SGD tends to converge to flat local minima (Zhang et al., 2018; Jastrzębski et al., 2017). This idea was elaborated on by Chaudhari et al. (2016) who suggest a new training method that favors flat over sharp minima even at the cost of a slightly higher empirical error – indeed solutions found by this algorithm exhibit better generalization performance. Similarly, Dziugaite & Roy (2017) augment the loss to improve generalization and find that this promotes flat minima. However, as Dinh et al. (2017) remarked, current flatness measures—which are based only on the Hessian of the loss function—cannot

theoretically be related to generalization: For deep neural networks with ReLU activation functions, there are layer-wise reparameterizations that leave the network function unchanged (hence, also the generalization performance), but change any measure derived only from the loss Hessian.

Another, more intuitive explanation for generalization is that the function generalizes well if the extracted features encode a semantic similarity of the input that is robust to small changes—both in the input and the features. This allows to generalize from the training set to novel, sufficiently similar data. Starting from such a concept of robustness with respect to changes of features, we derive a measure of flatness that is invariant under the mentioned reparameterizations and that reduces to the well-known ridge regression penalty in the special case of a linear regression.

This brings three seemingly related properties into our focus: flatness, robustness, and generalization. The exact relationship, however, between flatness of the loss surface around local minima (measuring changes of the empirical error for perturbations in parameter space), robustness (measuring changes of the error for perturbations in either

Figure 1: Illustration of the decomposition of $f = \psi \circ \phi$.

input or feature space), and generalization (performance on unseen data from the target distribution) is not well-understood. This paper provides new insights into this relationship.

The notion of feature robustness proposed in this paper measures the robustness of a function $f = \psi \circ \phi$ (e.g., a neural network) toward local changes in a feature space. That is, f can be split into a composition of functions $f(x) = (\psi \circ \phi)(x)$ for $x \in \mathcal{X}$, $\phi : \mathcal{X} \to \mathbb{R}^m$ and $\psi : \mathbb{R}^m \to \mathcal{Y}$. The function ϕ is considered as a feature extraction, mapping the input $\mathcal{X}$ into a **feature space** $\mathbb{R}^m$, while the function ψ corresponds to the model (e.g., a classifier) with $\mathbb{R}^m$ as its domain (see Figure 1 for illustration). It is the feature space defined by ϕ where we measure robustness toward small perturbations. For neural networks, the activation values of any but the output layer can be viewed as a feature space. A function f is called ϵ**-feature robust** on a dataset $S \subset \mathcal{X} \times \mathcal{Y}$ if small changes in the feature space defined by ϕ do not change the empirical error by more than ϵ. This differs from the notion of robustness defined by Xu & Mannor (2012) using a cover of the sample space, which has been theoretically connected to generalization. Flatness of the loss surface, however, is a local property and we require a more local version of robustness to derive a connection between flatness and robustness. Then, indeed, feature-robustness is upper bounded by the proposed flatness measure. To finally connect the two local properties of robustness and flatness to generalization, we necessarily need a notion describing how representative the given samples are for the true distribution. We define a suitable notion, leading to an upper bound for the generalization error given by feature robustness together with representativeness.

In summary, our contributions are as follows: (i) For models of the form $f(x) = (\psi \circ \phi)(x)$ (e.g. most (deep) neural networks) that split up into a feature extractor ϕ and a model ψ on the feature space defined by ϕ, we define a property of feature robustness that measures the change of the loss function under small perturbations of the features. This property is strongly related to flatness of the loss surface at local minima. (ii) We propose a novel flatness measure. For neural networks with ReLU activation functions, it is invariant under layer-wise reparameterization, addressing a shortcoming of previous measures of flatness. (iii) We define a suitable notion of representativeness of a dataset connecting feature robustness to the generalization error in form of an upper bound. (iv) The proposed flatness measure is empirically shown to strongly correlate with good generalization performance. Thereby, we recover Hessian based quantities as measures of flatness.

2   FEATURE ROBUSTNESS

We will define a notion of robustness in feature space $\mathbb{R}^m$ for the model $f = (\psi \circ \phi) : \mathcal{X} \to \mathcal{Y}$, which depends on a small number $\delta > 0$, a training set S, and a **feature selection** defined by a matrix $A \in \mathbb{R}^{m \times m}$ of **operator norm** $||A|| \leq 1$. In the case of neural networks split into a composition according to Figure 1, traditionally, the activation values $\phi_j(x)$ of neurons are considered as feature

values. The feature value defined by the j-th neuron in the feature space $\phi(x) \in \mathbb{R}^m$ can be written as $\phi_j(x) = \langle \phi(x), e_j \rangle$, where e_j denotes the j-th unit vector and $\langle \cdot, \cdot \rangle$ the scalar product in $\mathbb{R}^m$. However, it was shown by Szegedy et al. (2013) that, for any other direction $v \in \mathbb{R}^m, ||v|| = 1$, the values $\langle \phi(x), v \rangle = \text{proj}_v \phi(x)$ obtained from the projection $\phi(x)$ onto v, can be likewise semantically interpreted as a feature. We can single out the feature defined by v from $\phi(x)$ by multiplication with the projection matrix $E_v = vv^T$. Similarly, multiplication of $\phi(x)$ with a matrix A corresponds to a weighted selection of $rank(A)$-many features in parallel (e.g., projection matrices on d-dimensional subspaces correspond to the selection of d many features). This justifies our terminology considering a matrix A as a feature selection. The same way that, for a sample input x, non-activated neurons $\phi_j(x) = 0$ are considered as non-expressed features, we call a selection of features defined by matrix A as **non-expressed** whenever $A\phi(x) = 0$.

We define our notion of **feature robustness**. In words, feature robustness measures the mean change in loss over a dataset under small changes of features in the feature space. Hereby, a matrix A determines which features shall be perturbed. For each sample, the perturbation is linear in the expression of the feature. Thereby, we only perturb features that are relevant for the output for a given sample and leave feature values unchanged that are not expressed (in the sense explained above). With

$$\mathcal{F}(\delta, S, A) := \frac{1}{|S|} \sum_{(x,y) \in S} \left[\ell(\psi(\phi(x) + \delta A \phi(x)), y) - \ell(f(x), y) \right], \tag{2}$$

the precise definition is given as follows:

Definition 1. *Let $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}_+$ denote a loss function, δ and ϵ two strictly positive (small) real numbers, $S = \{(x_i, y_i) \mid i = 1, \ldots, N\} \subseteq \mathcal{X} \times \mathcal{Y}$ a set, and $A \in \mathbb{R}^{m \times m}$ a matrix such that $||A|| \leq 1$. A model $f(x) = (\psi \circ \phi)(x)$, which is a composition of functions $\phi : \mathcal{X} \to \mathbb{R}^m$ and $\psi : \mathbb{R}^m \to \mathcal{Y}$, is called $((\delta, S, A), \epsilon)$-feature robust, if $|\mathcal{F}(\delta', S, A)| \leq \epsilon$ for all $|\delta'| \leq \delta$.*
More generally, if $\mathcal{A} \subset \mathbb{R}^{m \times m}$ denotes a probability space over matrices such that $||A|| \leq 1$ for all $A \in \mathcal{A}$, then we call the model $((\delta, S, \mathcal{A}), \epsilon)$-feature robust on average over $\mathcal{A}$, if $\mathbb{E}_{A \sim \mathcal{A}} [|\mathcal{F}(\delta', S, A)|] \leq \epsilon$ for all $|\delta'| \leq \delta$.

We will bound feature robustness at local minima for a dataset S uniformly over all feature selections A and dependent on δ. With our interpretation, this corresponds to an upper bound of the change in loss when perturbing features in feature space $\mathbb{R}^m$. In Appendix C.1 we note how feature robustness is related to noise injection in the layer of consideration, which is known to be related to generalization (An, 1996; Bishop, 1995).

3 FEATURE ROBUSTNESS IS CONNECTED TO FLATNESS OF THE LOSS CURVE

Consider a function $f(x, \mathbf{w}) = \psi(\mathbf{w}, \phi(x)) = g(\mathbf{w}\phi(x))$, where ψ is the composition of a twice differentiable function $g : \mathbb{R}^d \to \mathcal{Y}$ and a matrix product with a matrix $\mathbf{w} \in \mathbb{R}^{d \times m}$. As before, $\phi : \mathcal{X} \to \mathbb{R}^m$ can be considered as a feature extractor. We fix a loss function $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}_+$ for supervised learning and let $\mathbf{w}_*$ denote a choice of parameters for which the empirical error $\mathcal{E}_{emp}(\mathbf{w}, S) = \frac{1}{|S|} \sum_{(x,y) \in S} \ell(f(x, \mathbf{w}), y)$, considered as a function on $\mathbf{w}$, is at a local minimum on the training set $S = \{(x_i, y_i) \mid i = 1, \ldots, N\}$. In the following, we write $z = \phi(x)$.

For any matrix $A \in \mathbb{R}^{m \times m}$ we have that

$$\psi(\mathbf{w}, z + \delta A z) = g(\mathbf{w}(z + \delta A z)) = g((\mathbf{w} + \delta \mathbf{w} A)z) = \psi(\mathbf{w} + \delta \mathbf{w} A, z). \tag{3}$$

Therefore,

$$\begin{aligned} \mathcal{F}(\delta, S, A) + \mathcal{E}_{emp}(\mathbf{w}, S) &= \frac{1}{|S|} \sum_{(x,y) \in S} \ell(\psi(\mathbf{w}, z + A\delta z), y) \\ &= \frac{1}{|S|} \sum_{(x,y) \in S} \ell(\psi(\mathbf{w} + \delta \mathbf{w} A, z), y). \end{aligned} \tag{4}$$

The latter is the empirical error $\mathcal{E}_{emp}(\mathbf{w} + \delta \mathbf{w} A, S)$ of the model f on the dataset S at parameters $\mathbf{w} + \delta \mathbf{w} A$. If δ is sufficiently small, then by Taylor expansion of $\mathcal{E}_{emp}(\mathbf{w}, S)$ with respect to

parameters $\mathbf{w}$ around the critical point $\mathbf{w}_*$, we have that

$$
\begin{aligned}
\mathcal{E}_{emp}(\mathbf{w}_* + \delta\mathbf{w}_*A, S) = {}& \mathcal{E}_{emp}(\mathbf{w}_*, S) + \langle \delta\mathbf{w}_*A,\ \nabla\mathcal{E}_{emp}(\mathbf{w}_*, S) \rangle \\
& + \frac{1}{2}\langle \delta\mathbf{w}_*A,\ H\mathcal{E}_{emp}(\mathbf{w}_*, S) \cdot (\delta\mathbf{w}_*A) \rangle + \mathcal{O}(\delta^3||\mathbf{w}_*A||_F^3) \\
= {}& \mathcal{E}_{emp}(\mathbf{w}_*, S) + \frac{\delta^2}{2}\langle \mathbf{w}_*A,\ H\mathcal{E}_{emp}(\mathbf{w}_*, S) \cdot (\mathbf{w}_*A) \rangle + \mathcal{O}(\delta^3||\mathbf{w}_*A||_F^3)
\end{aligned}
\tag{5}
$$

with $H\mathcal{E}_{emp}(\mathbf{w}_*, S)$ denoting the Hessian of the empirical error with respect to $\mathbf{w}$, $\langle \cdot, \cdot \rangle$ the scalar product with vectorized versions of the parameters and $||\mathbf{w}||_F$ the Frobenius norm of $\mathbf{w}$.

Subtracting $\mathcal{E}_{emp}(\mathbf{w}_*, S)$ from (5), maximizing over matrices $||A|| \leq 1$ and using (4), we get that, for any feature selection A, the function (2) defining feature robustness is bounded by

$$
\max_{||A|| \leq 1} \mathcal{F}(\delta, S, A) \leq \frac{\delta^2}{2}||\mathbf{w}_*||_F^2\ \lambda_{max}^H(\mathbf{w}_*) + \mathcal{O}(\delta^3)
\tag{6}
$$

where $\lambda_{max}^H(\mathbf{w}_*)$ denotes the largest eigenvalue of the Hessian $H\mathcal{E}_{emp}(\mathbf{w}_*, S)$ of the empirical error at $\mathbf{w}_*$. Here we used the identity that $\max_{||x||=1} x^T M x = \lambda_{max}^M$ for any symmetric matrix M, and that for matrices of norm $||A|| \leq 1$, we have $||\mathbf{w}_*A||_F \leq ||\mathbf{w}_*||_F$. We show details of the proof of (6) in the appendix. We summarize the connection between feature robustness and flatness in terms of the Hessian in the following theorem.

Theorem 2. *Let $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}_+$ denote a loss function, δ a strictly positive (small) real number, $A \in \mathbb{R}^{m \times m}$ a matrix with $||A|| \leq 1$, and let $f(x, \mathbf{w}) = g(\mathbf{w}\phi(x))$ be a model with g an arbitrary twice differentiable function on a matrix product of parameters $\mathbf{w}$ and the image of x under a (feature) function ϕ. Let $\mathbf{w}_*$ denote a local minimum of the empirical error on a dataset S.*

Then the model $f(\mathbf{w}_*)$ is $((\delta, S, A), \epsilon)$-feature robust for $\epsilon = \frac{\delta^2}{2}||\mathbf{w}_*||_F^2\ \lambda_{max}^H(\mathbf{w}_*) + \mathcal{O}(\delta^3)$.

4 MEASURES OF FLATNESS OF THE LOSS CURVE

Motivated by the relation of feature robustness with the Hessian H, we define a novel measure of flatness. Note that the Hessian is computed with respect to those parameters $\mathbf{w}$ that are applied linearly on the feature space $\phi(\mathcal{X}) \subseteq \mathbb{R}^m$.

Definition 3. *Let $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}_+$ denote a loss function and $f(x, \mathbf{w}) = g(\mathbf{w}\phi(x))$ be a model with $g : \mathbb{R}^m \to \mathcal{Y}$ an arbitrary twice differentiable function on a matrix product of parameters $\mathbf{w}$ and the image of x under a (feature) function $\phi : \mathcal{X} \to \mathbb{R}^m$. Then $\kappa^\phi(\mathbf{w})$ shall denote a flatness measure of the loss surface defined by*

$$
\kappa^\phi(\mathbf{w}) := ||\mathbf{w}||^2 \cdot \lambda_{max}^H(\mathbf{w}).
\tag{7}
$$

Note that small values of $\kappa^\phi(\mathbf{w})$ indicate flatness and high values indicate sharpness.

Linear regression with squared loss In the case of linear regression, $f(x, \mathbf{w}) = \mathbf{w}x \in \mathbb{R}$ ($\mathcal{X} = \mathbb{R}^d$, $g = id$ and $\phi = id$), for any loss function ℓ, we compute second derivatives with respect to the parameters $\mathbf{w} \in \mathbb{R}^d$ as

$$
\frac{\partial^2 \ell}{\partial w_i \partial w_j} = \frac{\partial^2 \ell}{\partial(f(x, \mathbf{w}))^2} x_i x_j
\tag{8}
$$

If ℓ is the squared loss function $\ell(\hat{y}, y) = (\hat{y} - y)^2$, then $\partial^2\ell/\partial\hat{y}^2 = 2$ and the Hessian is independent of the parameters $\mathbf{w}$. In this case, $\kappa^{id} = c \cdot ||\mathbf{w}||^2$ with a constant $c = 2\lambda_{max}(\sum_{x \in S} xx^t)$ and the measure κ^{id} reduces to (a constant multiple of) the well-known Tikhonov (ridge) regression penalty.

Layers of Neural Networks We consider neural network functions

$$
f(x) = \mathbf{w}_L\sigma(\ldots \sigma(\mathbf{w}_2\sigma(\mathbf{w}_1 x + b_1) + b_2)\ldots) + b_L
\tag{9}
$$

of a neural network of L layers with nonlinear activation function σ. We hide a possible non-linearity at the output by integrating it in a loss function ℓ chosen for neural network training. By letting $\phi^l(x) = \sigma(\mathbf{w}_{l-1}\sigma(\ldots \sigma(\mathbf{w}_2\sigma(\mathbf{w}_1 x + b_1) + b_2)\ldots) + b_{l-1})$ denote the output of the composition of the first $l-1$ layers and $g^l(z) = \mathbf{w}_L\sigma(\ldots \sigma(z + b_l)\ldots) + b_L$ the composition of the activation function

of the l-th layer together with the rest of layers, we can write for each layer l, $f(x, \mathbf{w}_l) = g^l(\mathbf{w}_l \phi^l(x))$. Using (7) we obtain for each layer of the neural network a measure of flatness at parameter values $\mathbf{w}$:

$$\kappa^l(\mathbf{w}) := ||\mathbf{w}_l||^2 \cdot \lambda_{max}^{H,l}(\mathbf{w}_l) \tag{10}$$

with $\lambda_{max}^{H,l}(\mathbf{w}_l)$ the largest eigenvalue of the Hessian of the empirical error with respect to the parameters of the l-th layer. By Theorem 2, κ^l is related to small changes of feature values in layer l.

Corollary 4. *Let f denote a neural network function of an L-layer fully connected neural network. For each layer $l, 1 \leq l \leq L$ of size n_l, let $A \in \mathbb{R}^{n_l \times n_l}$ with $||A|| \leq 1$ correspond to feature selections of features in the l-th layer of the neural network. Let $\mathbf{w}_{l*}$ denote weights of the l-th layer at a local minimum of the empirical error.*

Then the neural network is $((\delta, S, A), \epsilon)$- feature robust in layer l at $\mathbf{w}_*$ for $\epsilon = \frac{\delta^2}{2} \kappa^l(\mathbf{w}_*) + \mathcal{O}(\delta^3)$.

For an everywhere well-defined Hessian of the loss function, we assumed our network function to be twice differentiable. With the usual adjustments (equations only hold almost everywhere in parameter space), we can also consider neural networks with ReLU activation functions. In this case, Dinh et al. (2017) noted that a linear reparameterization of one layer, $\mathbf{w}_l \to \lambda \mathbf{w}_l$ for $\lambda > 0$, can lead to the same network function by simultaneously multiplying another layer by the inverse of λ, $\mathbf{w}_k \to 1/\lambda \mathbf{w}_k$, $k \neq l$. Representing the same function, the generalization performance remains unchanged. However, this linear reparameterization changes all common measures of the Hessian of the loss. This constitutes an issue in relating flatness of the loss curve to generalization. We counteract this behavior by the multiplication with $||\mathbf{w}_l||^2$.

Theorem 5. *Let $f = f(\mathbf{w}_1, \mathbf{w}_2, \ldots, \mathbf{w}_L)$ denote a neural network function parameterized by weights $\mathbf{w}_l$ of the l-th layer. Suppose there are positive numbers $\lambda_1, \ldots, \lambda_L$ such that $f(\mathbf{w}_1, \mathbf{w}_2, \ldots, \mathbf{w}_L) = f(\lambda_1 \mathbf{w}_1, \lambda_2 \mathbf{w}_2, \ldots, \lambda_L \mathbf{w}_L)$ for all $\mathbf{w}_l$. Then, with $\mathbf{w} = (\mathbf{w}_1, \mathbf{w}_2, \ldots, \mathbf{w}_L)$ and $\mathbf{w}^\lambda = (\lambda_1 \mathbf{w}_1, \lambda_2 \mathbf{w}_2, \ldots, \lambda_L \mathbf{w}_L)$, we have*

$$\kappa^l(\mathbf{w}) = \kappa^l(\mathbf{w}^\lambda) \text{ for all } 1 \leq l \leq L. \tag{11}$$

We provide a proof in Appendix A.2.

An Averaging Alternative Experimental work (Ghorbani et al., 2019) suggests that the spectrum of the Hessian has a lot of small values and only a few large outliers. In this case, our flatness measure serving as an upper bound for feature robustness is governed by the outlier. However, feature robustness for different feature selections is governed by different eigenvalues of the Hessian, according to (5). We therefore consider the trace as an average of the spectrum. We will show that this tracial averaging corresponds to feature robustness on average over all orthogonal feature selection matrices. The following theorem specifies this connection between feature robustness and the unnormalized trace $Tr(H\mathcal{E}_{emp}(\mathbf{w}_*))$ of the empirical error at a local minimum $\mathbf{w}_*$. The details and the proof can be found in Appendix A.3.

Theorem 6. *Let $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}_+$ denote a loss function, δ a strictly positive (small) real number, and let $f(x, \mathbf{w}) = g(\mathbf{w}\phi(x))$ be a model with g an arbitrary twice differentiable function on a matrix product of parameters $\mathbf{w} \in \mathbb{R}^{d \times m}$ and the image of x under a (feature) function ϕ. Let $\mathbf{w}_*$ denote a local minimum of the empirical error on a dataset S and $O_m \subset \mathbb{R}^{m \times m}$ denote the set of orthogonal matrices. Then, (i) for each feature selection matrix $||A|| \leq 1$ the model $f(\mathbf{w}_*)$ is $((\delta, S, A), \epsilon)$-feature robust for $\epsilon = \frac{\delta^2}{2} ||\mathbf{w}_*||_F^2 Tr(H\mathcal{E}_{emp}(\mathbf{w}_*)) + \mathcal{O}(\delta^3)$, and (ii) the model $f(\mathbf{w}_*)$ is $((\delta, S, O_m), \epsilon)$-feature robust on average over O_m for $\epsilon = \frac{\delta^2}{2m} ||\mathbf{w}_*||_F^2 Tr(H\mathcal{E}_{emp}(\mathbf{w}_*)) + \mathcal{O}(\delta^3)$.*

We therefore consider the unnormalized trace as a suitable and efficiently computable measure of flatness and define for each layer l of a neural network

$$\kappa_{Tr}^l(\mathbf{w}) := ||\mathbf{w}_l||_F^2 \cdot Tr(H\mathcal{E}_{emp}(\mathbf{w}_l, S)). \tag{12}$$

The same arguments as those used to prove Theorem 5 also show the measure κ_{Tr}^l to be independent with respect to the same layer-wise reparameterizations. The analogue of Corollary 4 is as follows.

Corollary 7. *Let f denote a neural network function of an L-layer fully connected neural network. For each layer $l, 1 \leq l \leq L$ of size n_l, let $O_{n_l} \subset \mathbb{R}^{n_l \times n_l}$ denote the set of orthogonal feature selections in the l-th layer of the neural network. Let $\mathbf{w}_{l*} \in \mathbb{R}^{n_{l+1} \times n_l}$ denote weights of the l-th layer at a local minimum of the empirical error. Then the neural network is $((\delta, S, O_n), \epsilon)$- feature robust in layer l on average over O_n at $\mathbf{w}_*$ for $\epsilon = \frac{\delta^2}{2n_l} \kappa_{Tr}^l(\mathbf{w}_*) + \mathcal{O}(\delta^3)$.*

Remark 8. *Other Hessian-based measures have been proposed that are invariant under the given reparameterizations. Liang et al. (2019) consider the Fisher-Rao norm defined by $||\mathbf{w}^T \hat{H}(\mathbf{w})\mathbf{w}||$ where $\hat{H}(\mathbf{w}) = (\nabla_{\mathbf{w}}\ell)(\nabla_{\mathbf{w}}\ell)^T$ denotes the Gauss-Newton approximation of the loss Hessian. Therefore, the Fisher-Rao norm considers the second partial derivative only into the direction defined by the given weight values $\mathbf{w}$. Our measure considers all directions of moving away from a local minimum. In particular, in contrast to these measures, we take the full spectrum of the Hessian into account, which results in a natural measure of flatness around a local minimum. We also came across preprints by Tsuzuku et al. (2019) and Rangamani et al. (2019), which propose a similar measure of flatness. While the first one derives the flatness measure from a PAC Baysian approach, the latter considers the Riemannian metric on the quotient manifold obtained from the equivalence relation given by the refactorization of layers as above.*

5 FEATURE ROBUSTNESS AND GENERALIZATION

In this section we aim to study the relation between flatness, feature robustness and the generalization error (defined in (1)). The connection of flat local minima with generalization in PAC Baysian bounds has been considered in several works (Neyshabur et al., 2017; Tsuzuku et al., 2019; Dziugaite & Roy, 2017). Arora et al. (2018) relates noise injection to generalization under the same setting. The work of McAllester (1998; 1999) and Langford & Caruana (2001) initiated the PAC-Baysian approach to generalization, which measures the generalization error (usually for the 0-1 loss) of stochastic classifiers. This leads to bounds on the expected true error over a distribution of models Q in terms of the expectation of the empirical error over Q and the Kullback–Leibler divergence between Q and some prior distribution P. For example, Neyshabur et al. (2017) use work by McAllester (2003) to derive an inequality relating the generalization error for the 0-1 loss ℓ and expected sharpness $\gamma_\nu := \mathbb{E}_\nu\left[\mathcal{E}_{emp}(\mathbf{w}+\nu, S)\right] - \mathcal{E}_{emp}(\mathbf{w}, S)$ over distribution ν: With probability $(1-\epsilon)$ over datasets S of size N, the expected error $\mathcal{E}_{\mathbf{w}+\nu} := \mathbb{E}_\nu\left[\mathbb{E}_{(x,y)\sim\mathcal{D}}\left[\ell(f(x, \mathbf{w}+\nu), y)\right]\right]$ over the "posterior" distribution $(\mathbf{w}+\nu)$ is bounded by $\mathcal{E}_{emp}(\mathbf{w}, S) + \gamma_\nu + 4\sqrt{\frac{1}{N}\left(\text{KL}(\mathbf{w}+\nu||P) + \ln\frac{2N}{\epsilon}\right)}$. Here, P denotes a "prior" distribution which is fixed before seeing any data and KL denotes the Kullback-Leibler divergence. If we aim to use distributions with local support (as considered in feature robustness and the Taylor expansion relating feature robustness to flatness) with a data-independent prior P, the KL term goes to infinity as the distribution ν becomes increasingly localized. Since feature robustness as a local property is related to generalization (Morcos et al., 2018), we aim to connect the local properties of flatness and feature robustness to generalization of a specific model by following a different approach. Our approach will be independent of the loss function and work in the sample space instead of averages over models in parameter space.

Since feature robustness is a local property in neighborhoods around the points $(x, y) \in S$, to connect feature robustness to generalization we necessarily need an assumption of representativeness of the given data samples S. A simple computation shows that

$$\mathcal{E}_{gen}(f, S) = \mathbb{E}_{A\sim\mathcal{A}}\left[\mathcal{F}(\delta, S, A)\right]$$
$$+ \left(\mathbb{E}_{(x,y)\sim\mathcal{D}}\left[\ell(f(x), y)\right] - \frac{1}{|S|}\sum_{(x_i,y_i)\in S}\mathbb{E}_{A\sim\mathcal{A}}\left[\ell(\psi(\phi(x_i) + \delta A\phi(x_i)), y_i)\right]\right) \quad (13)$$

The first term is exactly feature robustness on average over a probability distribution $\mathcal{A}$ of feature matrices. For the second term, we accordingly define a notion on datasets S that describes how well the loss on the true distribution can be approximated by certain probability distributions. The distributions we consider are composed of a dataset and (local) probability distributions around its points suitably restricted to local distributions λ_i and ν_i centered around the origin 0.

Definition 9. *Let $\psi : \mathbb{R}^m \to \mathcal{Y}$ be a model, $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}_+$ denote a loss function, ϵ a strictly positive (small) real number, and $S = \{(x_i, y_i) \mid i = 1, \ldots, N\} \subseteq \mathcal{X} \times \mathcal{Y}$ a set. Let $\Lambda = (\lambda_i, \nu_i)_{1\leq i\leq N}$ denote a family of pairs of probability distributions on $\mathbb{R}^m \times \mathcal{Y}$, where each λ_i and ν_i have support contained in a neighborhood of the origin 0. (i) The pair (S, Λ) is called*

ε-representative for ψ (with respect to the loss ℓ and distribution $\mathcal{D}$) if $|Rep(S, \Lambda)| \leq \epsilon$, where

$$Rep(S, \Lambda) := \mathbb{E}_{(x,y) \sim \mathcal{D}} \left[\ell(\psi(x), y) \right] - \frac{1}{|S|} \sum_{(x_i, y_i) \in S} \mathbb{E}_{(\xi_x, \xi_y) \sim (\lambda_i \times \nu_i)} \left[\ell(\psi(x_i + \xi_x), y_i + \xi_y) \right].$$

(14)

(ii) With Ω a collection of families Λ as above and $\mathcal{H}$ a hypothesis space, we say that S is **(ϵ, Ω)-representative** *for $\mathcal{H}$ if for all $\psi \in \mathcal{H}$ there is $\Lambda_\psi \in \Omega$ such that (S, Λ_ψ) is ϵ-representative for ψ.*

Interestingly, we naturally derived a definition of representativeness which is a generalization of classical ϵ-representativeness (see e.g. Definition 4.1 in (Shalev-Shwartz & Ben-David, 2014)), justifying the terminology. Indeed, let Λ_0 denote the family of probability distributions where each $\lambda_i = \delta_0$ and $\nu_i = \delta_0$ have full weight on the origin. Then S is $(\epsilon, \{\Lambda_0\})$-representative exactly when S is ϵ-representative in the classical sense. Further, if S is ϵ-representative and S is (ϵ', Ω)-representative for some Ω containing Λ_0, then $\epsilon' \leq \epsilon$.

In our setting of a model $f(x) = (\psi \circ \phi)(x)$, which is split up into a feature extractor ϕ and a model ψ, we consider $(\phi(S), \Lambda)$-representativeness for model ψ and specific choices for $\Lambda = \Lambda_{\delta, \mathcal{A}}$. Here, $\Lambda_{\delta, \mathcal{A}}$ is a family of probability distributions induced by a distribution $\mathcal{A}$ on feature matrices A such that $||A|| \leq \delta$ as follows: We assume that a Borel measure μ_A is defined by a probability distribution $\mathcal{A}$ on matrices $\mathbb{R}^{m \times m}$. We then define Borel measures μ_i on $\mathbb{R}^m$ by $\mu_i(C) = \mu_A(\{A \mid A\phi(x_i) \in C\})$ for Borel sets $C \subseteq \mathbb{R}^m$. Then λ_i is the probability distribution defined by μ_i. We fix the distributions $\nu_i = \delta_0$ and denote the set containing all families of distributions (λ_i, ν_i) that can be generated this way by $\mathfrak{A}_\delta$. The following result is a direct consequence of Equation 13 and our Definition 9.

Theorem 10. *Let $f(x) = (\psi \circ \phi)(x)$ be a model composed of functions $\phi : \mathcal{X} \to \mathbb{R}^m$ and $\psi : \mathbb{R}^m \to \mathcal{Y}$. If f is $((\delta, S, A), \epsilon)$-feature robust for all $||A|| \leq 1$ and $\phi(S)$ is $(\epsilon', \mathfrak{A}_\delta)$-representative for some $\mathcal{H}$ containing ψ, then the generalization error of f is bounded by $\mathcal{E}_{gen}(f, S) \leq \epsilon + \epsilon'$.*

Hence, for generalization we need a model that is feature-robust and training data that is sampled densely enough. In the trivial case with $\mathcal{A} = \delta_0$ the distribution with full weight on the 0-matrix, we can choose $\delta = 0$ to obtain $\epsilon = 0$ and $\mathcal{E}_{gen} \leq \epsilon'$. The more feature robust a model is, the larger δ we can consider to use the flexibility of choosing a nontrivial $\mathcal{A}$ to lower the bound on representativeness and therefore the generalization error. We hope that in future work it will be possible to find suitable distributions $\mathcal{A}$ that lead to computable generalization bounds.

6 EMPIRICAL EVALUATION

In this section we empirically validate the practical usefulness of the proposed flatness measure. A correlation between generalization and Hessian-based flatness measures at local minima has been observed previously, but the results of Dinh et al. (2017) questioned the usefulness of these measures. We show that our measure does not only overcome the theoretical issues, but also preserves the strong correlation with the generalization error. Previous works mostly use accuracy of the trained model on the testing dataset (Rangamani et al., 2019; Keskar et al., 2016) for evaluating the generalization properties of the achieved minimum. Nevertheless this does not directly correspond to the theoretical definition of the generalization error (1). For measuring the generalization error, we employ a Monte Carlo approximation of the target distribution defined by the testing dataset and measure the difference between loss value on this approximation and empirical error. In order to track the correlation of the flatness measure to the generalization error, sufficiently different minima should be achieved by training. The most popular technique is to train the model with small and large batch size (Rangamani et al., 2019; Keskar et al., 2016; Novak et al., 2018; Wang et al., 2018), which we also employed.

A neural network (LeNet5 (LeCun et al.)) is trained on CIFAR10 multiple times until convergence with various training setups. This way, we obtain network configurations in multiple local minima. In particular four different initialization schemes were considered (Xavier normal, Kaiming uniform, uniform in $(-0.1, 0.1)$, normal with $\mu = 0$ and $\sigma^2 = 0.1$), with four different mini-batch sizes (4, 32, 64, 512) and corresponding learning rates to keep the ration between them equal (0.001, 0.008, 0.02, 0.1) for the standard SGD optimizer. Each of the setups was run for 9 times with different random initializations.

Here the generalization error is the difference between summed error values on test samples multiplied by 5 (since the size of the training set is 5 times larger) and summed error values on the training

Figure 2: LeNet5 characteristics after training on CIFAR10. Each color corresponds to a different setup of training, characterized by initialization strategy, mini batch size and learning rate. The setups are ordered in ascending order by the mini batch size, with the largest corresponding to the brightest color of the displayed points.

Figure 3: LeNet5 configurations trained on CI-FAR10 with random reparameterizations. The correlation stays the same for the proposed measure, while it breaks for classic Hessian-based measure.

Figure 4: Robustness and flatness for LeNet5 configurations trained on CIFAR10. Results ordered by flatness, showing that robustness is bound by our flatness measure.

examples. Figure 2 shows the approximated generalization error with respect to the flatness measure (for both κ^l and κ^l_{Tr} with $l = 5$ corresponding to the last hidden layer) for all network configurations. The correlation is significant for both measures, and it is stronger (with $\rho = 0.91$) for κ^5_{Tr}. This indicates that taking into account the full spectrum of the Hessian is beneficial. To investigate the invariance of the proposed measure to reparameterization, we apply the reparameterization discussed in Sec. 4 to all networks using random factors in the interval $[5, 25]$. The impact of the reparameterization on the proposed flatness measure based on the trace in comparison to the traditional one is shown in Figure 3. While the proposed flatness measure is not affected, the one purely based on the Hessian has very weak correlation with the generalization error after the modifications. To verify the relation described by Equation 6, we also compared feature robustness with $\delta = 0.001$ and feature matrices A that have only one non-zero value 1 on the diagonal. Figure 4 shows that up to outliers the robustness is bound by the flatness measure. Additional experiments conducted on MNIST dataset are described in Appendix E, where we obtain correlation factors between the generalization error and tracial flatness κ^l_{Tr} of $0.73, 0.70, 0.72, 0.71$ for the network's hidden layers $l = 1, 2, 3, 4$ respectively.

7 DISCUSSION AND CONCLUSION

We established a theoretical connection between flatness, feature robustness and, under the assumption of representative data, the generalization error. The relation between feature robustness and Hessian-based flatness measures has been established for κ^l, which takes into account the maximum eigenvalue of the Hessian, and κ^l_{Tr}, which uses the trace instead. Empirically, the measure κ^l_{Tr} based on the trace of the Hessian shows a stronger correlation with the generalization error. This is not surprising,

since it takes into account the whole spectrum of the Hessian and every eigenvalue corresponds to a feature selection matrix of feature robustness. The tracial measure can be related to feature robustness by either bounding the maximum eigenvalue of the loss Hessian by its unnormalized trace or by averaging feature robustness over all orthogonal matrices $A \in O_m$. It is interesting to note that strong feature robustness does not exclude the possibility of adversarial examples, first observed by Szegedy et al. (2013), since large changes of loss for individual samples (i.e. adversarial examples) may be hidden in the mean in the definition of feature robustness. In Appendix C.2 we briefly discuss the freedom of perturbing individual points by suitable feature selection matrices A.

In contrast to existing measures of flatness, our proposed measure is invariant to layer-wise reparameterizations of ReLU networks. However, we note that other reparameterizations are possible, e.g., we can use the positive homogeneity and multiply all incoming weights into a single neuron by a positive number $\lambda > 0$ and multiply all outgoing weights of the same neuron by $1/\lambda$. While the Fisher-Rao norm suggested by Liang et al. (2019) is invariant to such reparameterizations, our proposed measures of flatness κ^l and κ^l_{Tr} are in general not. In principle, variations of our flatness measures can be found that are invariant to such reparameterizations as well (see Appendix B) but their analysis, except for some empirical evaluations in Appendix E, is left for future work.

The second term in the generalization bound of Theorem 10 is given by our notion of representativeness. In order to find specific bounds for the ϵ-representativeness of $(S, \mathfrak{A}_\delta)$, a distribution over matrices is required that induces a distribution which is similar to a localized kernel density estimation (KDE). While our notion of representativeness is a generalization of classical representativeness, it remains open whether it is efficiently computable. The more feature robust a model is, the more freedom there is to finding specific distributions over matrices that lead to bounds on the generalization error. In Appendix D we give a computation of representativeness for a KDE with Gaussian kernels.

Taking things together, we proposed a novel and practically useful flatness measure that strongly correlates with the generalization error. We theoretically investigated this connection by relating this measure to feature robustness. This notion of robustness, together with a novel notion of representativeness provides a link to the generalization error. To the best of our knowledge, this yields the first theoretical connection between a notion of robustness, flatness of the loss surface, and generalization error and can help to better understand the performance of deep neural networks.

REFERENCES

Guozhong An. The effects of adding noise during backpropagation training on a generalization performance. *Neural computation*, 8(3):643–674, 1996.

Sanjeev Arora, Rong Ge, Behnam Neyshabur, and Yi Zhang. Stronger generalization bounds for deep nets via a compression approach. *Proceedings of the 34th International Conference on Machine Learning*, 2018.

Chris M Bishop. Training with noise is equivalent to tikhonov regularization. *Neural computation*, 7 (1):108–116, 1995.

Pratik Chaudhari, Anna Choromanska, Stefano Soatto, Yann LeCun, Carlo Baldassi, Christian Borgs, Jennifer T. Chayes, Levent Sagun, and Riccardo Zecchina. Entropy-sgd: Biasing gradient descent into wide valleys. In *ICLR 2017*, 2016.

Laurent Dinh, Razvan Pascanu, Samy Bengio, and Yoshua Bengio. Sharp minima can generalize for deep nets. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pp. 1019–1028. JMLR. org, 2017.

Gintare Karolina Dziugaite and Daniel M Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. *AAAI*, 2017.

Gregory Fasshauer, Fred Hickernell, and Henryk Woźniakowski. On dimension-independent rates of convergence for function approximation with gaussian kernels. *Journal on Numerical Analysis*, 50 (1):247–271, 2012.

Behrooz Ghorbani, Shankar Krishnan, and Ying Xiao. An investigation into neural net optimization via hessian eigenvalue density. *arXiv preprint arXiv:1901.10159*, 2019.

Sepp Hochreiter and Jürgen Schmidhuber. Simplifying neural nets by discovering flat minima. In *Advances in neural information processing systems*, pp. 529–536, 1995.

Sepp Hochreiter and Jürgen Schmidhuber. Flat minima. *Neural Computation*, 9(1):1–42, 1997.

Stanisław Jastrzębski, Zachary Kenton, Devansh Arpit, Nicolas Ballas, Asja Fischer, Yoshua Bengio, and Amos Storkey. Three factors influencing minima in sgd. *arXiv preprint arXiv:1711.04623*, 2017.

Nitish Shirish Keskar, Dheevatsa Mudigere, Jorge Nocedal, Mikhail Smelyanskiy, and Ping Tak Peter Tang. On large-batch training for deep learning: Generalization gap and sharp minima. In *ICLR 2018*, 2016.

Steven G. Krantz and Harold R. Parks. *Geometric integration theory*. Springer Science and Business Media, 2008.

John Langford and Rich Caruana. (not) bounding the true error. In *Proceedings of the 14th International Conference on Neural Information Processing Systems: Natural and Synthetic*, NIPS'01, pp. 809–816, 2001.

Yann LeCun et al. Lenet-5, convolutional neural networks.

Tengyuan Liang, Tomaso Poggio, Alexander Rakhlin, and James Stokes. Fisher-rao metric, geometry, and complexity of neural networks. *International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2019.

David McAllester. Simplified pac-bayesian margin bounds. In *Learning theory and Kernel machines*, pp. 203–215. Springer, 2003.

David A McAllester. Some pac-bayesian theorems. In *Proceedings of the eleventh annual conference on Computational learning theory,*, volume 98, pp. 230–234, 1998.

David A McAllester. Pac-bayesian model averaging. In *Proceedings of the twelfth annual conference on Computational learning theory,*, volume 99, pp. 164–170, 1999.

Ari S Morcos, David GT Barrett, Neil C Rabinowitz, and Matthew Botvinick. On the importance of single directions for generalization. In *ICLR 2018*, 2018.

Behnam Neyshabur, Srinadh Bhojanapalli, David McAllester, and Nati Srebro. Exploring generalization in deep learning. In *Advances in Neural Information Processing Systems*, pp. 5947–5956, 2017.

Roman Novak, Yasaman Bahri, Daniel A Abolafia, Jeffrey Pennington, and Jascha Sohl-Dickstein. Sensitivity and generalization in neural networks: an empirical study. In *ICLR 2018*, 2018.

Akshay Rangamani, Nam H. Nguyen, Abhishek Kumar, Dzung T. Phan, Sang H. Chin, and Trac D. Tran. A scale invariant flatness measure for deep network minima. *arXiv preprint arXiv:1902.02434*, 2019.

Shai Shalev-Shwartz and Shai Ben-David. *Understanding machine learning: From theory to algorithms*. Cambridge university press, 2014.

Christian Szegedy, Wojciech Zaremba, Ilya Sutskever, Joan Bruna, Dumitru Erhan, Ian Goodfellow, and Rob Fergus. Intriguing properties of neural networks. *arXiv preprint arXiv:1312.6199*, 2013.

Yusuke Tsuzuku, Issei Sato, and Masashi Sugiyama. Normalized flat minima: Exploring scale invariant definition of flat minima for neural networks using pac-bayesian analysis. *arXiv preprint arXiv:1901.04653*, 2019.

Huan Wang, Nitish Shirish Keskar, Caiming Xiong, and Richard Socher. Identifying generalization properties in neural networks. *arXiv preprint arXiv:1809.07402*, 2018.

Huan Xu and Shie Mannor. Robustness and generalization. *Machine learning*, 86(3):391–423, 2012.

Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. In *ICLR 2017*, 2016.

Chiyuan Zhang, Qianli Liao, Alexander Rakhlin, Brando Miranda, Noah Golowich, and Tomaso Poggio. Theory of deep learning iib: Optimization properties of sgd. *arXiv preprint arXiv:1801.02254*, 2018.