Simple steps are all you need: Frank-Wolfe and generalized self-concordant functions

Abstract

Generalized self-concordance is a key property present in the objective function of many important learning problems. We establish the convergence rate of a simple Frank-Wolfe variant that uses the open-loop step size strategy $\gamma_t = 2/(t+2)$, obtaining a $O(1/t)$ convergence rate for this class of functions in terms of primal gap and Frank-Wolfe gap, where t is the iteration count. This avoids the use of second-order information or the need to estimate local smoothness parameters of previous work. We also show improved convergence rates for various common cases, e.g., when the feasible region under consideration is uniformly convex or polyhedral.

1 Introduction

Constrained convex optimization is the cornerstone of many machine learning problems. We consider such problems, formulated as:

$$\min_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}), \tag{1.1}$$

where $f : \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$ is a generalized self-concordant function and $\mathcal{X} \subseteq \mathbb{R}^n$ is a compact convex set. When computing projections onto the feasible regions as required in, e.g., projected gradient descent, is prohibitive, Frank-Wolfe (FW) (Frank & Wolfe, 1956) algorithms (a.k.a. Conditional Gradients (CG) (Levitin & Polyak, 1966)) are often the algorithm of choice, relying on Linear Minimization Oracles (LMO) at each iteration to solve Problem (1.1). The analysis of their convergence often relies on the assumption that the gradient is Lipschitz-continuous. This assumption does not necessarily hold for generalized self-concordant functions, an important class of functions for which the growth can be unbounded.

1.1 Related work

In the classical analysis of Newton's method, when the Hessian of f is assumed to be Lipschitz continuous and the function is strongly convex, one arrives at a convergence rate for the algorithm that depends on the Euclidean structure of $\mathbb{R}^n$, despite the fact that the algorithm is affine-invariant. This motivated the introduction of self-concordant functions in Nesterov & Nemirovskii (1994), functions for which the third derivative is bounded by the second-order derivative, with which one can obtain an affine-invariant convergence rate for the aforementioned algorithm. More importantly, many of the barrier functions used

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

in interior-point methods are self-concordant, which extended the use of polynomial-time interior point methods to many settings of interest.

Self-concordant functions have received strong interest in recent years due to the attractive properties that they allow to prove for many statistical estimation settings (Marteau-Ferey et al., 2019; Ostrovskii & Bach, 2021). The original definition of self-concordance has been expanded and generalized since its inception, as many objective functions of interest have self-concordant-like properties without satisfying the strict definition of self-concordance. For example, the logistic loss function used in logistic regression is not strictly self-concordant, but it fits into a class of pseudo-self-concordant functions, which allows one to obtain similar properties and bounds as those obtained for self-concordant functions (Bach et al., 2010). This was also the case in Ostrovskii & Bach (2021) and Tran-Dinh et al. (2015), in which more general properties of these pseudo-self-concordant functions were established. This was fully formalized in Sun & Tran-Dinh (2019), in which the concept of *generalized-self concordant* functions was introduced, along with key bounds, properties, and variants of Newton methods for the unconstrained setting which make use of this property.

Most algorithms that aim to solve Problem (1.1) assume access to second-order information, as this often allows the algorithms to make monotonous progress, remain inside the domain of f, and often, converge quadratically when close enough to the optimum. Recently, several lines of work have focused on using Frank-Wolfe algorithm variants to solve these types of problems in the projection-free setting, for example constructing second-order approxima-tions to a self-concordant f using first and second order information, and minimizing these approximations over $\mathcal{X}$ using the Frank-Wolfe algorithm (Liu et al., 2020). Other approaches, such as the ones presented in Dvurechensky et al. (2020a) (later extended in Dvurechensky et al. (2020b)), apply the Frank-Wolfe algorithm to a generalized self-concordant f, using first and second-order information about the function to guarantee that the step sizes are so that the iterates do not leave the domain of f, and monotonous progress is made. An additional FW variant in that work, in the spirit of Garber & Hazan (2016), utilizes first and second order information about f, along with a Local Linear Optimization Oracle for $\mathcal{X}$, to obtain a linear convergence rate in primal gap over polytopes given in inequality description. The last algorithm presented in Dvurechensky et al. (2020b), the only one that does not use second-order information, uses the FW algorithm with the backtracking line search of Pedregosa et al. (2020) to estimate local smoothness parameters at a given iterate. Other specialized FW algorithms have been developed for specific problems involving generalized self-concordant functions, such as the FW variant developed for marginal inference with concave maximization (Krishnan et al., 2015), or the variant developed in Zhao & Freund (2020) for θ-homogeneous barrier functions, a subset of standard self-concordant functions.

1.2 Contribution (see also Table 1)

Simple FW for generalized self-concordant functions. We show that a small variation of the original Frank-Wolfe algorithm (Frank & Wolfe, 1956) with an open-loop step size of the form $\gamma_t = 2/(t+2)$, where t is the iteration count is all that is needed to achieve a convergence rate of $O(1/t)$ in primal gap; this also answers an open question posed in Dvurechensky et al. (2020b). Our variation ensures monotonous progress while employing an open-loop strategy which, together with the iterates being convex combinations, ensures that we do not leave the domain of f. In contrast to other methods that depend on either a line search or second-order information, our variant uses only first-order information and a domain oracle for $f(\mathbf{x})$. The assumption of the latter oracle is very mild and was also implicitly assumed in the first-order algorithm in Dvurechensky et al. (2020b). As such, our iterations are much cheaper than those in previous work, while essentially achieving the same convergence rates in the general case of Problem (1.1). Moreover, our variant relying on the open-loop step size $\gamma_t = 2/(t+2)$ is adaptive, i.e., does not need to estimate local smoothness parameters and it allows to establish a convergence rate of $O(1/t)$ for the Frank-Wolfe gap as well.

Algorithm	Convergence		Reference	1$^{\text{st}}$-order / LS free?	Requirements
	Primal gap	FW gap			
FW-GSC	$O(1/\varepsilon)$	-	[1, Alg.2]	✗ / ✓	-
B-FW	$O(1/\varepsilon)$	-	[1, Alg.3]	✓ / ✗	DO
FW-LLOO	$O(\log 1/\varepsilon)$	-	[1, Alg.5]	✗ / ✓	polyh. $\mathcal{X}$, LLOO
M-FW	$O(1/\varepsilon)$	$O(1/\varepsilon)$	**This work**	✓ / ✓	DO
B-AFW	$O(\log 1/\varepsilon)$	$O(\log 1/\varepsilon)$	**This work**	✓ / ✗	polyh. $\mathcal{X}$, DO

Table 1: Convergence results for Problem 1.1 in the literature to achieve an ε-optimal solution, in terms of number of iterations. We denote Dvurechensky et al. (2020b) using [1], line search by LS, domain oracle by DO, and local linear optimization oracle by LLOO.

Faster rates in common special cases. We also obtain improved convergence rates when the optimum is contained in the interior of $\mathcal{X} \cap \text{dom}(f)$, or when the set $\mathcal{X}$ is uniformly or strongly convex, using the backtracking line search of Pedregosa et al. (2020). We also show that the Away-Step Frank-Wolfe algorithm (Lacoste-Julien & Jaggi, 2015; Wolfe, 1970) can use the aforementioned line search to achieve linear rates over polytopes.

Numerical experiments. We provide numerical experiments that showcase the performance of the algorithms on generalized self-concordant objectives to complement the theoretical results. In particular, they highlight that the simple step size strategy we propose is competitive with and sometimes outperforms other variants on many instances.

1.3 Preliminaries and Notation

We denote the (potentially non-unique) minimizer of Problem (1.1) by $\mathbf{x}^*$ and we denote the *primal gap* and the *Frank-Wolfe gap* at $\mathbf{x} \in \mathcal{X}$ as $h(\mathbf{x}) \overset{\text{def}}{=} f(\mathbf{x}) - f(\mathbf{x}^*)$ and $g(\mathbf{x}) \overset{\text{def}}{=} \max_{\mathbf{v} \in \mathcal{X}} \langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{v} \rangle$, respectively. We use $\|\cdot\|$, $\|\cdot\|_H$, and $\langle \cdot, \cdot \rangle$ to denote the *Euclidean norm*, the *matrix norm* induced by a symmetric positive definite matrix $H \in \mathbb{R}^{n \times n}$, and the *Euclidean inner product*, respectively. We denote the *diameter* of $\mathcal{X}$ as $D \overset{\text{def}}{=} \max_{\mathbf{x}, \mathbf{y} \in \mathcal{X}} \|\mathbf{x} - \mathbf{y}\|$. Given a non-empty set $\mathcal{X} \subset \mathbb{R}^n$ we refer to its *boundary* as $\text{Bd}(\mathcal{X})$ and to its *interior* as $\text{Int}(\mathcal{X})$. We denote the *probability simplex* of dimension n by Δ_n and the *domain* of f, as $\text{dom}(f) \overset{\text{def}}{=} \{\mathbf{x} \in \mathbb{R}^n, f(\mathbf{x}) < +\infty\}$. Given a compact convex set $C \subseteq \text{dom}(f)$ we denote $L_f^C = \max_{\mathbf{u} \in C, \mathbf{d} \in \mathbb{R}^n} \|\mathbf{d}\|_{\nabla^2 f(\mathbf{u})}^2 / \|\mathbf{d}\|_2^2$ and $\mu_f^C = \min_{\mathbf{u} \in C, \mathbf{d} \in \mathbb{R}^n} \|\mathbf{d}\|_{\nabla^2 f(\mathbf{u})}^2 / \|\mathbf{d}\|_2^2$. We assume access to:

1. **Domain Oracle (DO):** Given $\mathbf{x} \in \mathcal{X}$, return `true` if $\mathbf{x} \in \text{dom}(f)$, `false` otherwise.

2. **First-Order Oracle (FOO):** Given $\mathbf{x} \in \text{dom}(f)$, return $\nabla f(\mathbf{x})$.

3. **Linear Minimization Oracle (LMO):** Given $\mathbf{d} \in \mathbb{R}^n$, return $\text{argmin}_{\mathbf{x} \in \mathcal{X}} \langle \mathbf{x}, \mathbf{d} \rangle$.

The FOO and LMO oracles are standard in the FW literature and the DO oracle is motivated by the properties of generalized self-concordant functions. It is reasonable to assume the availability of such oracles: following the definition of the function codomain, one could simply evaluate f at $\mathbf{x}$ and assert $f(\mathbf{x}) < +\infty$. In many cases, testing the membership of $\mathbf{x} \in \text{dom}(f)$ is computationally less demanding than the function evaluation.

Remark 1.1. Access to a domain oracle is a mild assumption, that was also implicitly assumed in one of the three FW-variants presented in Dvurechensky et al. (2020b) when computing the step size according to the strategy from Pedregosa et al. (2020); see Line 3 in Algorithm 5 in the Appendix. The remaining two variants ensure that $\mathbf{x} \in \text{dom}(f)$ by using second-order information about f, which we explicitly do not rely on.

The following example motivates the use of Frank-Wolfe algorithms in the context of generalized self-concordant functions. We present more examples in the computational results.

Example 1.2 (Intersection of a convex set with a polytope). Consider Problem (1.1) where $\mathcal{X} = \mathcal{P} \cap C$, $\mathcal{P}$ is a polytope over which we can minimize a linear function efficiently, and C is a convex compact set for which one can easily build a barrier function.

(a) Plot of $f(\mathbf{x})$. (b) Plot of $f(\mathbf{x}) + \mu'\Phi_C(\mathbf{x})$. (c) Plot of $f(\mathbf{x}) + \mu\Phi_C(\mathbf{x})$.

Figure 1: Minimizing $f(\mathbf{x})$ over $\mathcal{P} \cap C$, versus minimizing the sum of $f(\mathbf{x})$ and $\Phi_C(\mathbf{x})$ over $\mathcal{P}$ for two different penalty values μ' and μ such that $\mu' \gg \mu$.

Solving a linear optimization problem over $\mathcal{X}$ may be extremely expensive. In light of this, we can incorporate C into the optimization problem through the use of a barrier penalty in the objective function, minimizing instead $f(\mathbf{x}) + \mu\Phi_C(\mathbf{x})$ where $\Phi_C(\mathbf{x})$ is a log-barrier function for C and μ is a parameter controlling the penalization for points closer to $\mathrm{Bd}(C)$. The reformulation of the problem is illustrated in Figure 1. Note that if the original objective function is generalized self-concordant, so is the new objective function. We assume that computing the gradient of $f(\mathbf{x}) + \mu\Phi_C(\mathbf{x})$ is roughly as expensive as computing the gradient for $f(\mathbf{x})$ and solving an LP over $\mathcal{P}$ is inexpensive relative to solving an LP over $\mathcal{P} \cap C$. The μ parameter can be driven down to 0 after a solution converges in a warm-starting procedure similar to interior-point methods, ensuring convergence to the true optimum.

An additional advantage of this transformation of the problem is the solution structure. Running Frank-Wolfe on the set $\mathcal{P} \cap C$ could potentially select a large number of extremal points from $\mathrm{Bd}(C)$ if C is non-polyhedral. In contrast, $\mathcal{P}$ has a finite number of vertices, a small subset of which will be selected throughout the optimization procedure. The same solution as that of the original problem can thus be constructed as a convex combination of a small number of vertices of $\mathcal{P}$, improving sparsity and interpretability in many applications.

The following definition formalizes the setting of Problem (1.1).

Definition 1.3 (Generalized self-concordant function). Let $f \in C^3(\mathrm{dom}(f))$ be a closed convex function with $\mathrm{dom}(f) \subseteq \mathbb{R}^n$ open. Then f is (M, ν) *generalized self-concordant* if:

$$|\langle D^3 f(\mathbf{x})[\mathbf{w}]\mathbf{u}, \mathbf{u}\rangle| \leq M \|\mathbf{u}\|^2_{\nabla^2 f(\mathbf{x})} \|\mathbf{w}\|^{\nu-2}_{\nabla^2 f(\mathbf{x})} \|\mathbf{w}\|^{3-\nu}_2,$$

for any $\mathbf{x} \in \mathrm{dom}(f)$ and $\mathbf{u}, \mathbf{w} \in \mathbb{R}^n$, where $D^3 f(\mathbf{x})[\mathbf{w}] = \lim_{\alpha \to 0} \alpha^{-1}\left(\nabla^2 f(\mathbf{x} + \alpha\mathbf{w}) - \nabla^2 f(\mathbf{x})\right)$.

2 Frank-Wolfe Convergence Guarantees

Algorithm 1 Monotonous Frank-Wolfe (M-FW)

Input: Point $\mathbf{x}_0 \in \mathcal{X} \cap \mathrm{dom}(f)$, function f
Output: Iterates $\mathbf{x}_1, \ldots \in \mathcal{X}$

1: **for** $t = 0$ **to** $\ldots$ **do**
2:     $\mathbf{v}_t \leftarrow \mathrm{argmin}_{\mathbf{v} \in \mathcal{X}} \langle \nabla f(\mathbf{x}_t), \mathbf{v}\rangle$
3:     $\gamma_t \leftarrow 2/(t+2)$
4:     $\mathbf{x}_{t+1} \leftarrow \mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)$
5:     **if** $\mathbf{x}_{t+1} \notin \mathrm{dom}(f)$ **or** $f(\mathbf{x}_{t+1}) > f(\mathbf{x}_t)$ **then**
6:         $\mathbf{x}_{t+1} \leftarrow \mathbf{x}_t$

We establish convergence rates for a Frank-Wolfe variant with an open-loop step size strategy on generalized self-concordant functions. The *Monotonous Frank-Wolfe (M-FW)* algorithm presented in Algorithm 1 is a rather simple, but powerful modification of the standard Frank-Wolfe algorithm, with the only difference that before taking a step, we verify if $\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t) \in \mathrm{dom}(f)$, and if so, we check whether moving to the next iterate provides primal progress. Note, that the open-loop step size rule $2/(t+2)$ does not guarantee monotonous primal progress for the vanilla Frank-Wolfe algorithm in general. If either of these two checks fails, we simply do not move:

the algorithm sets $\mathbf{x}_{t+1} = \mathbf{x}_t$ in Line 6 of Algorithm 1. As customary, we assume short-circuit evaluation of the logical conditions in Algorithm 1, i.e., if the first condition in Line 5 is true, then the second condition is not even checked, and the algorithm directly goes to Line 6. This minor modification of the vanilla Frank-Wolfe algorithm enables us to use the monotonicity of the iterates in the proofs to come, at the expense of one extra function evaluation per iteration. In order to lower bound the progress per iteration we use Proposition 2.1.

Proposition 2.1. *(C.f., (Sun & Tran-Dinh, 2019, Proposition 10)) Given a (M, ν) generalized self-concordant function, then for $\nu \geq 2$, we have that:*

$$f(\mathbf{y}) - f(\mathbf{x}) - \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle \leq \omega_\nu(d_\nu(\mathbf{x} - \mathbf{y})) \|\mathbf{y} - \mathbf{x}\|_{\nabla^2 f(\mathbf{x})}^2, \tag{2.1}$$

where the inequality holds if and only if $d_\nu(\mathbf{x}, \mathbf{y}) < 1$ for $\nu > 2$, and we have that,

$$d_\nu(\mathbf{x}, \mathbf{y}) \stackrel{\text{def}}{=} \begin{cases} M \|\mathbf{y} - \mathbf{x}\| & \text{if } \nu = 2 \\ (\frac{\nu}{2} - 1)M \|\mathbf{y} - \mathbf{x}\|^{3-\nu} \|\mathbf{y} - \mathbf{x}\|_{\nabla^2 f(\mathbf{x})}^{\nu-2} & \text{if } \nu > 2, \end{cases}$$

where:

$$\omega_\nu(\tau) \stackrel{\text{def}}{=} \begin{cases} \frac{e^\tau - \tau - 1}{\tau^2} & \text{if } \nu = 2 \\ \frac{-\tau - \ln(1-\tau)}{\tau^2} & \text{if } \nu = 3 \\ \frac{(1-\tau)\ln(1-\tau)+\tau}{\tau^2} & \text{if } \nu = 4 \\ \left(\frac{\nu-2}{4-\nu}\right) \frac{1}{\tau} \left[\frac{\nu-2}{2(3-\nu)\tau} \left((1-\tau)^{\frac{2(3-\nu)}{2-\nu}} - 1 \right) - 1 \right] & \text{otherwise.} \end{cases}$$

The inequality shown in Equation (2.1) is very similar to the one that we would obtain if the gradient of f were Lipschitz continuous, however, while the Lipschitz continuity of the gradient leads to an inequality that holds globally for all $\mathbf{x}, \mathbf{y} \in \text{dom}(f)$, the inequality in Equation (2.1) only holds for $d_\nu(\mathbf{x}, \mathbf{y}) < 1$. Moreover, there are two other important differences, the norm used in Equation (2.1) is now the norm defined by the Hessian at $\mathbf{x}_t$ instead of the ℓ_2 norm, and the term multiplying the norm is $\omega_\nu(d_\nu(\mathbf{x}, \mathbf{y}))$ instead of $1/2$. We deal with the latter issue by bounding $\omega_\nu(d_\nu(\mathbf{x}, \mathbf{y}))$ with a constant that depends on ν for any $\mathbf{x}, \mathbf{y} \in \text{dom}(f)$ such that $d_\nu(\mathbf{x}, \mathbf{y}) \leq 1/2$, as shown in Remark 2.2.

Remark 2.2. As $d\omega_\nu(\tau)/d\tau > 0$ for $\tau < 1$ and $\nu \geq 2$, then $\omega_\nu(\tau) \leq \omega_\nu(1/2)$ for $\tau \leq 1/2$.

Due to the fact that we use a simple step size $\gamma_t = 2/(t + 2)$, that we make monotonous progress, and we ensure that the iterates are inside $\text{dom}(f)$, careful accounting allows us to bound the number of iterations until $d_\nu(\mathbf{x}_t, \mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)) \leq 1/2$. Before formalizing the convergence rate we first review a lemma that we will need in the proof.

Lemma 2.3. *(C.f., (Sun & Tran-Dinh, 2019, Proposition 7)) Let f be a generalized self concordant function with $\nu > 2$. If $d_\nu(\mathbf{x}, \mathbf{y}) < 1$ and $\mathbf{x} \in \text{dom}(f)$ then $\mathbf{y} \in \text{dom}(f)$. For the case $\nu = 2$ we have that $\text{dom}(f) = \mathbb{R}^n$.*

Putting all these things together allows us to obtain a convergence rate for Algorithm 1.

Theorem 2.4. *Suppose $\mathcal{X}$ is a compact convex set and f is a (M, ν) generalized self-concordant function with $\nu \geq 2$. Then the Monotonous Frank-Wolfe algorithm (Algorithm 1) satisfies:*

$$h(\mathbf{x}_t) \leq \frac{4(T_\nu + 1)}{t + 1} \max \left\{ h(\mathbf{x}_0), L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2) \right\}. \tag{2.2}$$

for $t \geq T_\nu$, where $L_f^{\mathcal{L}_0} = \max\limits_{\mathbf{u} \in \mathcal{L}_0, \mathbf{d} \in \mathbb{R}^n} \|\mathbf{d}\|_{\nabla^2 f(\mathbf{u})}^2 / \|\mathbf{d}\|_2^2$ and T_ν is defined as:

$$T_\nu \stackrel{\text{def}}{=} \begin{cases} \lceil 4MD \rceil - 2 & \text{if } \nu = 2 \\ \left\lceil 2MD(L_f^{\mathcal{L}_0})^{\nu/2-1}(\nu - 2) \right\rceil - 2 & \text{otherwise.} \end{cases} \tag{2.3}$$

Otherwise it holds that $h(\mathbf{x}_t) \leq h(\mathbf{x}_0)$ for $t < T_\nu$.

Proof. Consider the compact set $\mathcal{L}_0 \overset{\text{def}}{=} \{\mathbf{x} \in \text{dom}(f) \cap \mathcal{X} \mid f(\mathbf{x}) \leq f(\mathbf{x}_0)\}$. As the algorithm makes monotonous progress and moves towards points such that $\mathbf{x}_t \in \text{dom}(f)$, then $\mathbf{x}_t \in \mathcal{L}_0$ for $t \geq 0$. This allows us to claim, in a similar fashion as is done in Dvurechensky et al. (2020b), that $\|\mathbf{x}_t - \mathbf{v}_t\|^2_{\nabla^2 f(\mathbf{x}_t)} \leq L_f^{\mathcal{L}_0} D^2$. We then define T_ν as in Equation (2.3). Note that for $t \geq T_\nu$ we have that $d(\mathbf{x}_t, \mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)) \leq 1/2$, and so as $\mathbf{x}_t \in \text{dom}(f)$ we will have $\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t) \in \text{dom}(f)$ for $t \geq T_\nu$, by application of Lemma 2.3. This means that the non-zero step size γ_t will automatically ensure that $\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t) \in \text{dom}(f)$ in Line 5 of Algorithm 1. Moreover, it allows us to use the upper bound on the Bregman divergence between points $\mathbf{x}_t$ and $\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)$ in Proposition 2.1, which holds for $d(\mathbf{x}_t, \mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)) < 1$. With this we can estimate the primal progress we can guarantee for $t \geq T_\nu$ if we move from $\mathbf{x}_t$ to $\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)$:

$$h(\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)) \leq h(\mathbf{x}_t) - \gamma_t g(\mathbf{x}_t) + \gamma_t^2 \omega_\nu\left(d_\nu(\mathbf{x}_t, \mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t))\right)\|\mathbf{v}_t - \mathbf{x}_t\|^2_{\nabla^2 f(\mathbf{x}_t)}$$

$$\leq h(\mathbf{x}_t)(1 - \gamma_t) + \gamma_t^2 L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2),$$

where the second inequality follows from the upper bound on the primal gap via the Frank-Wolfe gap $g(\mathbf{x}_t)$, the application of Remark 2.2 as for $t \geq T_\nu$ we have that $d_\nu(\mathbf{x}_t, \mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)) \leq 1/2$, and from the fact that $\mathbf{x}_t \in \mathcal{L}_0$ for all $t \geq 0$. With the previous chain of inequalities we can bound the primal progress for $t \geq T_\nu$ as

$$h(\mathbf{x}_t) - h(\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)) \geq \gamma_t h(\mathbf{x}_t) - \gamma_t^2 L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2). \tag{2.4}$$

From these facts we can prove the convergence rate shown in Equation (2.2) by induction. The base case $t = T_\nu$ holds trivially by the fact that using monotonicity we have that $h(\mathbf{x}_{T_\nu}) \leq h(\mathbf{x}_0)$. Assuming the claim is true for some $t \geq T_\nu$ we distinguish two cases.

Case $\gamma_t h(\mathbf{x}_t) - \gamma_t^2 L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2) > 0$: Focusing on the first case, we can plug the previous inequality into Equation (2.4) to find that γ_t guarantees primal progress, that is, $h(\mathbf{x}_t) > h(\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t))$ with the step size γ_t, and so we know that we will not go into Line 6 of Algorithm 1, and we have that $h(\mathbf{x}_{t+1}) = h(\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t))$. Thus using the induction hypothesis and plugging in the expression for $\gamma_t = 2/(t+2)$ into Equation (2.4) we have:

$$h(\mathbf{x}_{t+1}) \leq 4 \max\left\{h(\mathbf{x}_0), L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2)\right\}\left(\frac{(T_\nu + 1)t}{(t+1)(t+2)} + \frac{1}{(t+2)^2}\right)$$

$$\leq \frac{4(T_\nu + 1)}{t+2} \max\left\{h(\mathbf{x}_0), L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2)\right\},$$

where we use that $(T_\nu + 1)t/(t+1) + 1/(t+2) \leq T_\nu + 1$ for all $t \geq 0$ and any $t \geq T_\nu$.

Case $\gamma_t h(\mathbf{x}_t) - \gamma_t^2 L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2) \leq 0$: In this case, we cannot guarantee that the step size γ_t provides primal progress by plugging into Equation (2.4), and so we cannot guarantee if a step size of γ_t will be accepted and we will have $\mathbf{x}_{t+1} = \mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)$, or we will simply have $\mathbf{x}_{t+1} = \mathbf{x}_t$, that is, we may go into Line 6 of Algorithm 1. Nevertheless, if we reorganize the expression $\gamma_t h(\mathbf{x}_t) - \gamma_t^2 L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2) \leq 0$, by monotonicity we will have that:

$$h(\mathbf{x}_{t+1}) \leq h(\mathbf{x}_t) \leq \frac{2}{t+2} L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2) \leq \frac{4(T_\nu + 1)}{t+2} \max\left\{h(\mathbf{x}_0), L_f^{\mathcal{L}_0} D^2 \omega_\nu(1/2)\right\}.$$

Where the last inequality holds as $2 \leq 4(T_\nu + 1)$ for any $T_\nu \geq 0$. $\qquad\square$

Remark 2.5. In the case where $\nu = 2$ we can easily bound the primal gap $h(\mathbf{x}_1)$, as in this setting $\text{dom}(f) = \mathbb{R}^n$, which leads to $h(\mathbf{x}_1) \leq L_f^{\mathcal{X}} D^2$ from Equation (2.4), regardless of if we set $\mathbf{x}_1 = \mathbf{x}_0$ or $\mathbf{x}_1 = \mathbf{v}_0$. Moreover, as the upper bound on the Bregman divergence holds for $\nu = 2$ regardless of the value of $d_2(\mathbf{x}, \mathbf{y})$, we can modify the proof of Theorem 2.4 to obtain a convergence rate of the form $h(\mathbf{x}_t) \leq 2/(t+1)L_f^{\mathcal{X}} D^2 w_2(MD)$ for $t \geq 1$, which is reminiscent of the $\mathcal{O}(L_f^{\mathcal{X}} D^2/t)$ rate of the original Frank-Wolfe algorithm for the smooth and convex case.

Note that in the proof of Theorem 2.4 we explicitly use the progress bound from generalized self-concordance as opposed to the progress bound that arises from $L_f^{\mathcal{L}_0}$-smoothness, as there

222 is no straightforward way to bound the number of iterations until the latter progress bound
223 holds indefinitely for all $\mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)$, while there is a straightforward criterion on γ_t that
224 allows us to ensure that the former holds from some point onward (see Remark A.1 for more
225 details). Furthermore, with this simple step size we can also prove a convergence rate for the
226 Frank-Wolfe gap, as shown in Theorem 2.6 (see Theorem A.2 in Appendix for the proof).

227 **Theorem 2.6.** *Suppose $\mathcal{X}$ is a compact convex set and f is a (M, ν) generalized self-concordant*
228 *function with $\nu \geq 2$. Then if the Monotonous Frank-Wolfe algorithm (Algorithm 1) is run for*
229 $T \geq T_\nu + 6$ *iterations, we will have that $\min_{1 \leq t \leq T} g(\mathbf{x}_t) \leq O(1/T)$.*

2.1 Improved convergence guarantees

Algorithm 2 (Monotonous) Frank-Wolfe with Backtrack of Pedregosa et al. (2020)

Input: $\mathbf{x}_0 \in \mathcal{X} \cap \mathrm{dom}(f)$, function f, estimate L_{-1}
Output: Iterates $\mathbf{x}_1, \ldots \in \mathcal{X}$

1: **for** $t = 0$ **to** ... **do**
2: $\quad \mathbf{v}_t \leftarrow \mathrm{argmin}_{\mathbf{v} \in \mathcal{X}} \langle \nabla f(\mathbf{x}_t), \mathbf{v} \rangle$
3: $\quad \gamma_t, L_t \leftarrow \texttt{Backtrack}(f, \mathbf{x}_t, \mathbf{v}_t - \mathbf{x}_t, L_{t-1}, 1)$
4: $\quad \mathbf{x}_{t+1} \leftarrow \mathbf{x}_t + \gamma_t(\mathbf{v}_t - \mathbf{x}_t)$

We will now establish improved convergence rates for various special cases. We first focus on the assumption that $\mathbf{x}^* \in \mathrm{Int}(\mathcal{X} \cap \mathrm{dom}(f))$, obtaining improved rates when we use the FW algorithm coupled with the adaptive step size strategy from Pedregosa et al. (2020) (see Algorithm 5 in Appendix).

The analysis in this case is reminiscent of the analysis of Guélat & Marcotte

241 (1986), and is a reasonable assumption if for example $\mathrm{Bd}(\mathcal{X}) \not\subseteq \mathrm{dom}(f)$, and $\mathrm{Int}(\mathcal{X}) \subseteq$
242 $\mathrm{dom}(f)$. We can upper bound the value of L_t for $t \geq 0$ by $\tilde{L} \stackrel{\mathrm{def}}{=} \max\{\tau L_f^{\mathcal{L}_0}, L_{-1}\}$, where $\tau > 1$
243 is the backtracking parameter and L_{-1} is the initial smoothness estimate in Algorithm 5.

244 **Theorem 2.7.** *Let f be a (M, ν) generalized self-concordant function with $\nu \geq 2$ and let*
245 $\mathrm{dom}(f)$ *not contain straight lines. Furthermore, we denote by $r > 0$ the largest value such that*
246 $\mathcal{B}(\mathbf{x}^*, r) \subseteq \mathcal{X} \cap \mathrm{dom}(f)$. *Then the Frank-Wolfe algorithm with Backtrack (Algorithm 2) achieves*
247 *a convergence rate for $t \geq 1$ of:*

$$h(\mathbf{x}_t) \leq h(\mathbf{x}_0) \left(1 - \frac{\mu_f^{\mathcal{L}_0}}{2\tilde{L}} \left(\frac{r}{D} \right)^2 \right)^t.$$

248 The assumption that $\mathrm{dom}(f)$ does not contain straight lines in Theorem 2.7 is related to the
249 Hessian being positive definite over $\mathrm{dom}(f)$ (see the proof in the Appendix in Theorem A.5).
250 Note that this is a very mild assumption as we can simply modify the function with a very
251 small ℓ_2 regularizer, as e.g., in Nesterov (2012). Next, we recall the definition of uniformly
252 convex sets, used in Kerdreux et al. (2021), which will allow us to to obtain improved
253 convergence rates for the FW algorithm over uniformly convex feasible regions.

254 **Definition 2.8** ((κ, q)-uniformly convex set)**.** Given two positive numbers κ and q, we say the
255 set $\mathcal{X} \subseteq \mathbb{R}^n$ is (κ, q)-*uniformly convex* with respect to a norm $\|\cdot\|$ if for any $\mathbf{x}, \mathbf{y} \in \mathcal{X}$, $0 \leq \gamma \leq 1$,
256 and $\mathbf{z} \in \mathbb{R}^n$ with $\|\mathbf{z}\| = 1$ we have that $\mathbf{y} + \gamma(\mathbf{x} - \mathbf{y}) + \gamma(1 - \gamma) \cdot \kappa \|\mathbf{x} - \mathbf{y}\|^q \mathbf{z} \in \mathcal{X}$.

257 **Theorem 2.9.** *Suppose $\mathcal{X}$ is a compact (κ, q)-strongly convex set and f is a (M, ν) generalized*
258 *self-concordant function with $\nu \geq 2$. Furthermore, assume that $\min_{\mathbf{x} \in \mathcal{X}} \|\nabla f(\mathbf{x})\| \geq C > 0$. Then*
259 *the Frank-Wolfe algorithm with Backtrack (Algorithm 2) achieves a convergence rate of:*

$$h_t \leq \begin{cases} h(\mathbf{x}_0) \left(1 - \frac{1}{2} \min \left\{ 1, \frac{\kappa C}{\tilde{L}} \right\} \right)^t & \text{if } q = 2 \\ \frac{h(\mathbf{x}_0)}{2^t} & \text{if } q > 2, 1 \leq t \leq t_0 \\ \frac{\tilde{L}^{q/(q-2)}/(\kappa C)^{2/(q-2)}}{(1 + (q-2)(t-t_0)/(2q))^{q/(q-2)}} = O\left(t^{-q/(q-2)} \right) & \text{if } q > 2, t > t_0, \end{cases}$$

260 *for $t \geq 1$, where $t_0 = \max \left\{ 1, \left\lfloor \log_{1/2} \left(\frac{(\tilde{L}^q/(\kappa C)^2)^{1/(q-2)}}{h(\mathbf{x}_0)} \right) \right\rfloor \right\}$.*

261 However, in the general case we cannot assume that the norm of the gradient is bounded
262 away from zero over $\mathcal{X}$. We deal with the general case in Theorem 2.10

263 **Theorem 2.10.** *Suppose $\mathcal{X}$ is a compact (κ, q)-strongly convex set and f is a (M, ν) generalized*
264 *self-concordant function with $\nu \geq 2$ for which the domain does not contain straight lines. Then*
265 *the Frank-Wolfe algorithm with Backtrack (Algorithm 2) results in a convergence rate:*

$$h_t \leq \begin{cases} \dfrac{h(\mathbf{x}_0)}{2^t} & \text{if } 1 \leq t \leq t_0 \\ \dfrac{(\breve{L}^q/(\kappa^2 \mu_f^{\mathcal{L}_0}))^{1/(q-1)}}{(1+(q-1)(t-t_0)/(2q))^{q/(q-1)}} = O\left(t^{-q/(q-1)}\right) & \text{if } t > t_0, \end{cases}$$

266 *for $t \geq 1$, where $t_0 = \max\left\{1, \left\lfloor \log_{1/2}\left(\dfrac{(\breve{L}^q/(\kappa^2 \mu_f^{\mathcal{L}_0}))^{1/(q-1)}}{h(\mathbf{x}_0)}\right)\right\rfloor\right\}$.*

267 **Remark 2.11.** Contrary to previous claims, there is no obstacle for the *Away-step Frank-Wolfe*
268 (AFW) algorithm (Guélat & Marcotte, 1986; Lacoste-Julien & Jaggi, 2015) together with
269 the step size strategy in Algorithm 5 to obtain a linear convergence rate in primal and
270 Frank-Wolfe gap when $\mathcal{X}$ is a polytope and f is generalized self-concordant. This is not
271 surprising, as f is strongly convex and smooth over $\mathcal{L}_0$ if dom(f) does not contain straight
272 lines, and monotonicity ensures the feasibility of the iterates. We leave the analysis for this
273 case to Appendix B, and the formal convergence statement to Theorem B.2 and B.3.

3 Computational experiments

275 We showcase the performance of the Monotonous Frank-Wolfe algorithm (`M-FW`), the second-
276 order step size and the LLOO algorithm from Dvurechensky et al. (2020b) (`GSC-FW` and
277 `LLOO`) and the Frank-Wolfe and the Away-Step Frank-Wolfe algorithm with the backtracking
278 stepsize of Pedregosa et al. (2020), denoted by `B-FW` and `B-AFW` respectively. All experi-
279 ments are carried out in `Julia` using the `FrankWolfe.jl` package (Besançon et al., 2021),
280 available under the MIT license and the examples considered extend the ones presented in
281 Dvurechensky et al. (2020b) and Liu et al. (2020). We also use the vanilla FW algorithm
282 denoted by `FW`, which is simply Algorithm 1 without Lines 5 and 6 using the traditional
283 $\gamma_t = 2/(t+2)$ open-loop step size rule. Note that there are no formal convergence guarantees
284 for this algorithm when applied to Problem (1.1). Details on the experiments setup, data
285 and remarks on the considered problems are provided in Appendix C. All figures show the
286 evolution of the $h(\mathbf{x}_t)$ and $g(\mathbf{x}_t)$ against time and number of iterations with a log-log scale.
287 As in Dvurechensky et al. (2020b) we implemented the LLOO based variant only for the
288 portfolio optimization instance over the probability simplex; for the other examples the
289 oracle implementation was less straightforward due to the estimation of parameters.

290 As can be seen in all experiments, the Monotonous Frank-Wolfe algorithm is very competitive,
291 outperforming previously proposed variants in both in progress per iteration and time. The
292 only other algorithm that is sometimes faster is the Away-Step Frank-Wolfe variant as detailed
293 in Remark 2.11, which however depends on an active set, and can induce up to a quadratic
294 overhead, making iterations progressively more expensive; this can be also observed in our
295 experiments as the advantage in time is much less pronounced than in iterations.

296 **Portfolio optimization.** We consider $f(\mathbf{x}) = -\sum_{t=1}^{P} \log(\langle \mathbf{r}_t, \mathbf{x} \rangle)$, where p denotes the number
297 of periods and $\mathcal{X} = \Delta_n$. The results are shown in Figure 2.

298 **Signal recovery with KL divergence.** We apply the aforementioned algorithms to the
299 recovery of a sparse signal from a noisy linear image using the Kullback-Leibler divergence,
300 expressed as $f(\mathbf{x}) = D(W\mathbf{x}, \mathbf{y}) = \sum_{i=1}^{N} \left\{ \langle \mathbf{w}_i, \mathbf{x} \rangle \log\left(\frac{\langle \mathbf{w}_i, \mathbf{x} \rangle}{y_i}\right) - \langle \mathbf{w}_i, \mathbf{x} \rangle + y_i \right\}$, where $\mathbf{w}_i$ is the i^{th}
301 row of a matrix W. In order to promote sparsity and enforce nonnegativity of the solution,
302 we use the unit simplex of radius R as the feasible set $\mathcal{X} = \{\mathbf{x} \in \mathbb{R}_+^d, \|\mathbf{x}\|_1 \leq R\}$. The results
303 are shown in Figure 3. We used the same $M = 1$ choice for the second-order method as in
304 Dvurechensky et al. (2020b) for comparison; its admissibility is unknown (see Remark C.1).

305 **Logistic regression.** We consider a logistic regression task with a design matrix with
306 rows $\mathbf{a}_i \in \mathbb{R}^n$ with $1 \le i \le N$ and a vector $\mathbf{y} \in \{-1, 1\}^N$ and formulate the problem
307 with elastic net regularization, in a similar fashion as is done in Liu et al. (2020), with
308 $f(\mathbf{x}) = 1/N \sum_{i=1}^N \log(1 + \exp(-y_i \langle \mathbf{x}, \mathbf{a}_i \rangle)) + \mu/2 \|\mathbf{x}\|^2$, where μ is a regularization parameter
309 and $\mathcal{X}$ is the ℓ_1 ball of radius ρ. The results can be seen in Figure 4 and Appendix C.

310 **Birkhoff polytope.** All previously considered applications have in common a feasible region
311 possessing computationally inexpensive LMOs (probability/unit simplex and ℓ_1 norm ball).
312 Additionally, each vertex returned from the LMO is highly sparse with at most one non-zero
313 element. To complement the results, we consider the logistic regression problem over the
314 Birkhoff polytope, where the LMO call uses the Hungarian method and is not as inexpensive
315 as in the other examples. The results are shown in Figure 5.

Figure 2: **Portfolio Optimization**: LLOO and GSC-FW perform similarly to FW on a per-iteration
basis but the iterations are computationally more expensive. B-AFW is the fastest method
both in terms iteration and runtime, followed by M-FW which is the only other method to
terminate with the specified dual gap tolerance.

Figure 3: **Signal Recovery**: B-AFW significantly outperforms all other methods. FW and B-FW
perform similarly in dual gap progress and converge slower than M-FW. In terms of primal
gap progress, M-FW and FW perform similarly and outperform B-FW.

Figure 4: **Logistic Regression**: This instance shows that although simple in essence, M-FW
can outperform other methods including B-AFW in terms of convergence. The primal and
dual gaps for B-FW and GSC-FW converge at similar rates against iteration count.

Figure 5: **Birkhoff Polytope**: B-AFW is the fastest-converging method for all measures.
However, the dual gap reaches a plateau due to numerical issues above the termination
threshold, unlike M-FW which reaches the dual gap tolerance. GSC-FW is run for 1000 iterations
only given the longer runtime. Its slow progress is likely due to numerical instabilities in the
Hessian computation which do not occur in first-order methods.

References

Bach, F. et al. Self-concordant analysis for logistic regression. *Electronic Journal of Statistics*, 4:384–414, 2010.

Besançon, M., Carderera, A., and Pokutta, S. FrankWolfe.jl: a high-performance and flexible toolbox for Frank-Wolfe algorithms and conditional gradients. *arXiv preprint arXiv:2104.06675*, 2021.

Csiszar, I. et al. Why least squares and maximum entropy? An axiomatic approach to inference for linear inverse problems. *The annals of statistics*, 19(4):2032–2066, 1991.

Dvurechensky, P., Ostroukhov, P., Safin, K., Shtern, S., and Staudigl, M. Self-concordant analysis of Frank-Wolfe algorithms. In *Proceedings of the 37th International Conference on Machine Learning*, pp. 2814–2824. PMLR, 2020a.

Dvurechensky, P., Safin, K., Shtern, S., and Staudigl, M. Generalized self-concordant analysis of Frank-Wolfe algorithms. *arXiv preprint arXiv:2010.01009*, 2020b.

Frank, M. and Wolfe, P. An algorithm for quadratic programming. *Naval research logistics quarterly*, 3(1-2):95–110, 1956.

Garber, D. and Hazan, E. A linearly convergent variant of the conditional gradient algorithm under strong convexity, with applications to online and stochastic optimization. *SIAM Journal on Optimization*, 26(3):1493–1528, 2016.

Guélat, J. and Marcotte, P. Some comments on Wolfe's 'away step'. *Mathematical Programming*, 35(1):110–119, 1986.

Jaggi, M. Revisiting Frank-Wolfe: Projection-free sparse convex optimization. In *Proceedings of the 30th International Conference on Machine Learning*, pp. 427–435. PMLR, 2013.

Kerdreux, T., d'Aspremont, A., and Pokutta, S. Restarting Frank-Wolfe. In *Proceedings of the 22nd International Conference on Artificial Intelligence and Statistics*, pp. 1275–1283. PMLR, 2019.

Kerdreux, T., d'Aspremont, A., and Pokutta, S. Projection-free optimization on uniformly convex sets. In *Proceedings of the 24th International Conference on Artificial Intelligence and Statistics*, pp. 19–27. PMLR, 2021.

Krishnan, R. G., Lacoste-Julien, S., and Sontag, D. Barrier Frank-Wolfe for Marginal Inference. In *Proceedings of the 28th Conference in Neural Information Processing Systems*. PMLR, 2015.

Lacoste-Julien, S. and Jaggi, M. On the global linear convergence of Frank-Wolfe optimization variants. In *Proceedings of the 29th Conference on Neural Information Processing Systems*, pp. 566–575. PMLR, 2015.

Levitin, E. S. and Polyak, B. T. Constrained minimization methods. *USSR Computational Mathematics and Mathematical Physics*, 6(5):1–50, 1966.

Liu, D., Cevher, V., and Tran-Dinh, Q. A Newton Frank-Wolfe method for constrained self-concordant minimization. *arXiv preprint arXiv:2002.07003*, 2020.

Marron, J. S., Todd, M. J., and Ahn, J. Distance-weighted discrimination. *Journal of the American Statistical Association*, 102(480):1267–1271, 2007.

Marteau-Ferey, U., Ostrovskii, D., Bach, F., and Rudi, A. Beyond least-squares: Fast rates for regularized empirical risk minimization through self-concordance. In *Proceedings of the 32nd Conference on Learning Theory*, pp. 2294–2340. PMLR, 2019.

Nesterov, Y. How to make the gradients small. *Optima. Mathematical Optimization Society Newsletter*, (88):10–11, 2012.

Nesterov, Y. and Nemirovskii, A. *Interior-point polynomial algorithms in convex programming*. SIAM, 1994.

Nesterov, Y. et al. *Lectures on convex optimization*, volume 137. Springer, 2018.

Ostrovskii, D. M. and Bach, F. Finite-sample analysis of M-estimators using self-concordance. *Electronic Journal of Statistics*, 15(1):326–391, 2021.

Pedregosa, F., Negiar, G., Askari, A., and Jaggi, M. Linearly convergent Frank–Wolfe with backtracking line-search. In *Proceedings of the 23rd International Conference on Artificial Intelligence and Statistics*. PMLR, 2020.

Sun, T. and Tran-Dinh, Q. Generalized self-concordant functions: a recipe for Newton-type methods. *Mathematical Programming*, 178(1):145–213, 2019.

Temlyakov, V. Greedy approximation in convex optimization. *Constructive Approximation*, 41 (2):269–296, 2015.

Tran-Dinh, Q., Li, Y.-H., and Cevher, V. Composite convex minimization involving self-concordant-like cost functions. In *Modelling, Computation and Optimization in Information Systems and Management Sciences*, pp. 155–168. Springer, 2015.

Wolfe, P. Convergence theory in nonlinear programming. In *Integer and Nonlinear Programming*, pp. 1–36. North-Holland, Amsterdam, 1970.

Zhao, R. and Freund, R. M. Analysis of the Frank-Wolfe method for logarithmically-homogeneous barriers, with an extension. *arXiv preprint arXiv:2010.08999*, 2020.