Global Convergence and Stability of Stochastic Gradient Descent

Abstract

In machine learning, stochastic gradient descent (SGD) is widely deployed to train models using highly non-convex objectives with equally complex noise models. Unfortunately, SGD theory often makes restrictive assumptions that fail to capture the non-convexity of real problems, and almost entirely ignore the complex noise models that exist in practice. In this work, we demonstrate the restrictiveness of these assumptions using three canonical models in machine learning, then we develop novel theoretical tools to address this shortcoming in two ways. First, we establish that SGD's iterates will either globally converge to a stationary point or diverge under nearly arbitrary nonconvexity and noise models. Under a slightly more restrictive assumption on the joint behavior of the non-convexity and noise model that generalizes current assumptions in the literature, we show that the objective function cannot diverge, even if the iterates diverge. As a consequence of our results, SGD can be applied to a greater range of stochastic optimization problems with confidence about its global convergence behavior and stability.

1 Introduction

Stochastic Gradient Descent (SGD) and its variants are dominant algorithms for solving stochastic optimization problems arising in machine learning, and have expanded their reach to more complex problems from estimating Gaussian Processes [Chen et al., 2020], covariance estimation in stochastic filters [Kim et al., 2021], and systems identification [Hardt et al., 2016, Zhang and Patel, 2020]. Accordingly, understanding the behavior of SGD and its variants has been crucial to their reliable application in machine learning and beyond. As a result, the theory of these methods has greatly advanced, most notably for SGD as it is the basis for, and simplest of, these methods. Indeed, SGD has been analyzed from many perspectives: global convergence analysis [Lei et al., 2019, Gower et al., 2020, Khaled and Richtárik, 2020, Mertikopoulos et al., 2020, Patel, 2021], local convergence analysis [Mertikopoulos et al., 2020], greedy and global complexity analysis [Gower et al., 2020, Khaled and Richtárik, 2020], asymptotic weak convergence [Wang et al., 2021], and saddle point analysis [Fang et al., 2019, Mertikopoulos et al., 2020, Jin et al., 2021].

While all of these perspectives add new dimensions to our understanding of SGD, the global convergence analysis of SGD is the foundation as it dictates whether local analyses, complexity analyses or saddle point analyses are even warranted. As surveyed in Patel [2021], these current global convergence analyses of SGD make a wide variety of assumptions, most commonly: (1) the objective function is bounded from below, (2) the gradient function is globally Lipschitz continuous, (3) the stochastic gradients are unbiased, and (4) the variance of the stochastic gradients are bounded. While the first and third assumption are generally reasonable,[1] the second and fourth assumptions are not

---

[1]See Bottou et al. [2018] for a simple relaxation of unbiased stochastic gradients.

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

generally applicable to machine learning problems as we now demonstrate through three simple examples (see Appendix A for details).

Example 1: Linear Regression. Consider fitting a linear regression model using squared-error loss given independent copies of the example (Y, Z) where for some θ^* and error term ϵ, $Y = \langle Z, \theta^* \rangle + \epsilon$. The next results states that for a very nice linear regression problem, the stochastic gradients violate the bounded variance assumption.

Proposition 1. *There exist random variables Z and ϵ such that Z has bounded fourth moment, and ϵ is mean zero with bounded variance such that the variance of the stochastic gradients evaluated at θ diverges to infinity as $\|\theta\|_2 \to \infty$.*

Example 2: Feed Forward Neural Network. Consider the example (Y, Z) where Y is a binary label and Z is a feature vector. We will attempt to predict Y from Z using a simple multi-layer feed forward network as shown in Fig. 1. The next result states that for a simple distribution over the example space and for a simple, archetype network, the gradient function is not globally Lipschitz continuous *and* the stochastic gradient does not have globally bounded variance.

Figure 1: A diagram of a simple feed forward network for binary classification.

Proposition 2. *Consider the feed forward network in Fig. 1 with σ linear and φ sigmoid trained with a binary cross-entropy loss with a Ridge penalty. There exists a finite, discrete distribution for (Y, Z) such that the risk function's gradient is not globally Lipschitz continuous, nor is the variance of the stochastic gradients bounded.*

Example 3: Recurrent Neural Network. Consider the example (Y, Z_0, Z_1, Z_2, Z_3) where Y is a binary label and $\{Z_0, Z_1, Z_2, Z_3\}$ are sequentially observed. We will attempt to predict Y from Z using a simple recurrent network as shown in Fig. 2. The next result states that for a simple distribution over the example space and for a simple, archetype network, the gradient function violates global Lipschitz continuity *and* the stochastic gradient violates the bounded variance assumption.

Figure 2: A diagram of a recurrent neural network for a binary classification task.

Proposition 3. *Consider the recurrent network in Fig. 1 with σ linear and φ sigmoid trained with a binary cross-entropy loss with a Ridge penalty. There exists a finite, discrete distribution for (Y, Z_0, Z_1, Z_2, Z_3) such that the risk function's gradient is not globally Lipschitz continuous, nor is the variance of the stochastic gradients bounded.*

As these three examples show, global convergence analyses that make use of the aforementioned assumptions do not apply to these canonical examples of machine learning problems. In fact, while the linear regression case can be covered by more recent analyses (see Bottou et al. [2018]), to our

Table 1: A survey of influential, recent global analyses of SGD and their dependence on the two assumptions that are either individually or both violated by the simple examples discussed in Section 1.

Assumption	Works Depending on the Assumption
Global Lipschitz or Hölder Continuity of Gradient	Reddi et al. [2016a], Ma and Klabjan [2017], Zhou et al. [2018], Bassily et al. [2018], Lei et al. [2019], Li and Orabona [2019], Gower et al. [2020], Khaled and Richtárik [2020], Mertikopoulos et al. [2020], Patel [2021], Jin et al. [2021], Wang et al. [2021].
Bounded Variance of Stochastic Gradients	Reddi et al. [2016b], Ma and Klabjan [2017], Majewski et al. [2018], Hu et al. [2019], Bi and Gunn [2019], Zou et al. [2019], Mertikopoulos et al. [2020].[2]

knowledge and as summarized Table 1, *there are no global convergence analyses of SGD* that apply to the neural network training examples.

The Problem:
As a result, we do not have guarantees about how SGD behaves on these simple neural network problems, which calls into question what SGD and its variants are doing on more complicated machine learning models.

Contributions. To address this problem,

1. We relax the global Lipschitz continuous gradient assumption and the bounded variance assumption to assumptions that are applicable to the examples above. Specifically, we relax the global Lipschitz continuity assumption to local α-Hölder continuity of the gradient for $\alpha \in (0, 1]$ (see Assumption 2), which is even a relaxation even for deterministic gradient algorithms (c.f. Nocedal and Wright [2006],Theorems 3.2, 3.8, 4.5, 4.6). For the α in the local Hölder assumption, we also relax the bounded variance assumption to only require that the $(1 + \alpha)$-moment of the stochastic gradient is bounded by an *arbitrary* upper semi-continuous function (see Assumption 4). Our assumption allows stochastic gradients whose noise may not have a variance. Moreover, our assumption generalizes the noise assumption of Bottou et al. [2018], the expected smoothness assumption of Gower et al. [2020] and Khaled and Richtárik [2020], and the noise assumption of Asi and Duchi [2019]. We also point out that we do not require the coercivity or asymptotic flatness assumptions that are commonly considered in the analysis of SGD (e.g., Mertikopoulos et al. [2020], Assumptions 2 and 3).

2. Owing to the relaxation in the assumptions, we cannot apply the standard analysis as the local Hölder constant and the iterate difference are conditionally dependent random variables (see the discussion after Lemma 1). As a result, we innovate a new analysis strategy (see Section 4.1) to prove that, with probability one, either SGD's iterates will converge to a stationary point *or* they diverge (see Theorem 2). Importantly, our new analysis strategy can be broadly applied even to deterministic algorithms to relax the assumptions found in the literature.

3. The divergence component of our Theorem 2 is somewhat disconcerting as we cannot say exactly what happens if the iterates diverge. To ameliorate this concern, we add an additional assumption (see Assumption 5) and introduce another analysis strategy (see Section 4.2) to prove, even if the iterates are diverging, the objective function converges to a finite random variable and the gradient function converges to zero with probability one (see Theorem 3).

Our results *do not* supply a rate of convergence as this is impossible for the broad class of nonconvex functions and the generality of the noise models studied in this work [Wolpert and Macready, 1997]; in other words, we can always construct a nonconvex objective function, a noise process, and choose an initialization such that any rate of convergence statement will be violated. Indeed, we find it remarkable that it is even possible to provide a global convergence statement for such a broad class of nonconvex functions and general noise models.

---

[2]In Majewski et al. [2018], this is implied by their third assumption.

Limitations. We make note of two important limitations in our work. First, we do not consider the important case of nonsmoothness in this work as we require that the gradients of the stochastic optimization function are continuous. However, we note that if the results of Bianchi et al. [2022] are broadly applicable, then SGD never observes a point of nonsmoothness and our results would then be applicable. Second, we do not have a simple interpretation of Assumption 5—though it seems to have a close relative in another analysis (see Wang and Wu [2020])—, nor have we been able to construct a relevant counterexample that can illuminate the limitations of this assumption.

2 Stochastic Optimization

We consider solving the optimization problem

$$\min_{\theta \in \mathbb{R}^p} \{ F(\theta) := \mathbb{E}\left[f(\theta, X) \right] \}, \tag{1}$$

where F maps $\mathbb{R}^p$ into $\mathbb{R}$; f maps $\mathbb{R}^p$ and the co-domain of the random variable X into $\mathbb{R}$; and $\mathbb{E}$ is the expectation operator. As we require gradients, we take F and f to differentiable in θ, and denote its derivatives with respect to θ by $\dot{F}(\theta)$ and $\dot{f}(\theta, X)$. With this notation, we make the following general assumptions about the deterministic portion of the objective function.

Assumption 1. There exists $F_{l.b.} \in \mathbb{R}$ such that $\forall \theta \in \mathbb{R}^p$, $F_{l.b.} \leq F(\theta)$.

Assumption 2. There exists $\alpha \in (0, 1]$ such that $\dot{F}(\theta)$ is locally α-Hölder continuous.

Remark 1. As an example, an empirical risk minimization problem for a deep neural network with twice continuously differentiable activation functions with a twice continuously differentiable loss function will readily satisfy the above conditions.

Assumptions 1 and 2 would even be considered mild in the context of non-convex deterministic optimization, in which it is also common to assume that the objective function well-behaved level sets [e.g., Nocedal and Wright, 2006, Theorems 3.2, 3.8, 4.5, 4.6]. Importantly, Assumption 2 relaxes the common restrictive assumption of globally Hölder continuous gradient functions that is common in other analyses (see Table 1).

Our final step is to make some assumptions about the stochastic portion of the objective function. The first assumption requires the stochastic gradients to be unbiased, which can readily be relaxed [Bottou et al., 2018]. The second assumption allows for a generic noise model for an α-Hölder continuous gradient function, and even allows for the second moment to not exist when $\alpha < 1$ [c.f. Wang et al., 2021, which requires a decomposition of the noise term that we do not require].

Assumption 3. For all $\theta \in \mathbb{R}^p$, $\dot{F}(\theta) = \mathbb{E}[\dot{f}(\theta, X)]$.

Assumption 4. Let $\alpha \in (0, 1]$ be as in Assumption 2. There exists an upper semi-continuous function $G(\theta)$ such that $\mathbb{E}[\|\dot{f}(\theta, X)\|_2^{1+\alpha}] \leq G(\theta)$.

> We will show that Assumptions 1 to 4 are sufficient for a global convergence result (see Theorem 2).

Remark 2. It is entirely possible that $\mathbb{E}[\|\dot{f}(\theta, X)\|_2^{1+\alpha}]$ is (at least) upper semi-continuous, and to set $G(\theta)$ equal to this function. In the case that $\mathbb{E}[\|\dot{f}(\theta, X)\|_2^{1+\alpha}]$ is not upper semi-continuous, it is possible to specify $G(\theta)$ as the upper envelope of $\mathbb{E}[\|\dot{f}(\theta, X)\|_2^{1+\alpha}]$ (i.e., the its limit supremum function). However, it is unlikely that $\mathbb{E}[\|\dot{f}(\theta, X)\|_2^{1+\alpha}]$ nor its upper envelope are easy to specify explicitly, and it is more likely to be able to find an upper bound.

Remark 3. We use Assumption 4 to conclude that on any compact set, the $1 + \alpha$ moment of the stochastic gradient is bounded. Of course, we can assume this directly (i.e., on any compact set, the $1 + \alpha$ moment is bounded), which, at first glance, appears to be a relaxation. However, if we assume that on any compact set, the $1 + \alpha$ moment is bounded, we can use this to construct a $G(\theta)$ that is upper semi-continuous. Thus, the two assumptions are equivalent.

Remark 4. A simple example that shows the utility of Assumption 4 is to optimize $\mathbb{E}[\theta^X]$ where X is an exponential random variable with parameter 1 and $\theta \in [1, u]$ where $u < \exp(1)$. First, it is easy to confirm that the objective function is differentiable and its derivative is globally Lipschitz continuous. Moreover, given that we are on a bounded interval, we conclude that the derivative is

globally α-Hölder continuous for any $\alpha \in (0, 1]$; therefore, we are free to choose the α as we see fit. Now, when $u < \exp(1/2)$, we have that second moment of the stochastic gradient function exists. However, when $\exp(1/2) < u < \exp(1)$, only smaller moments of the stochastic gradient will exist. Specifically, only for $u < \exp(1/(1 + \epsilon))$ with $\epsilon \in (0, 1)$ will the $1 + \epsilon$ moment of the stochastic gradient will exist. Thus, depending on the size of our interval, we may not have the existence of the second moment.

In order to show that the objective function cannot diverge (i.e., to prove stability), we will need an additional assumption. This assumption will relate the gradient function, noise model and variation on the local Hölder constant. To begin, we define the variation on the local Hölder constant. Let $\alpha \in (0, 1]$ be as in Assumption 2 and $\epsilon > 0$ be arbitrary, and define

$$
\mathcal{L}_\epsilon(\theta) = \begin{cases} \sup_\varphi \left\{ \frac{\left\| \dot{F}(\varphi) - \dot{F}(\theta) \right\|_2}{\|\varphi - \theta\|_2^\alpha} : \|\varphi - \theta\|_2 \le (G(\theta) \vee \epsilon)^{\frac{1}{1+\alpha}} \right\} & \text{if this quantity is nonzero} \\ \epsilon & \text{otherwise,} \end{cases} \tag{2}
$$

where $\vee$ indicates the maximum between two quantities. Note, the choice of ϵ is irrelevant, and they can be distinct for the two cases in the definition of $\mathcal{L}_\epsilon$, but we fix them to be the same for simplicity. Importantly, the quantity, $\mathcal{L}_\epsilon$, is defined at every parameter θ under Assumption 2.

With this quantity, we can state a nonintuitive,technical assumption that is needed to prove stability.

Assumption 5. There exists $C_1, C_2, C_3 \ge 0$ such that, $\forall \theta \in \mathbb{R}^p$,

$$
\mathcal{L}_\epsilon(\theta) G(\theta) + \alpha \left(\frac{\left\| \dot{F}(\theta) \right\|_2^{1+\alpha}}{\mathcal{L}_\epsilon(\theta)} \right)^{1/\alpha} \le C_1 + C_2(F(\theta) - F_{l.b.}) + C_3 \left\| \dot{F}(\theta) \right\|_2^2. \tag{3}
$$

Assumption 5 generalizes Assumption 4.3(c) of Bottou et al. [2018], which is satisfied for a large swath of statistical models. Moreover, Assumption 5 generalizes the notion of expected smoothness [see Khaled and Richtárik, 2020, for a history of the assumption], which expanded the optimization problems covered by the theory of Bottou et al. [2018]. Note, Assumption 5 is about the asymptotic properties of the stochastic optimization problem as the left hand side of the inequality in Assumption 5 can be bounded inside of any compact set. Thus, Assumption 5 covers a variety of asymptotic behaviors, such as $\exp(\|\theta\|_2^2)$, $\exp(\|\theta\|_2)$, $\|\theta\|_2^r$ for $r \in \mathbb{R}$, $\log(\|\theta\|_2 + 1)$, and $\log(\log(\|\theta\|_2 + 1) + 1)$. Therefore, Assumption 5 holds for functions with a variety of different asymptotic behaviors.

> We will show that Assumptions 1 to 5 are sufficient for a stability result (see Theorem 3).

Now that we have specified the nature of the stochastic optimization problem, we turn our attention to the algorithm used to solve the problem, namely, stochastic gradient descent (SGD).

3 Stochastic Gradient Descent

SGD starts with an arbitrary initial value, $\theta_0 \in \mathbb{R}^p$, and generates a sequence of iterates $\{\theta_k : k \in \mathbb{N}\}$ according to the rule

$$
\theta_{k+1} = \theta_k - M_k \dot{f}(\theta_k, X_{k+1}), \tag{4}
$$

where $\{M_k : k + 1 \in \mathbb{N}\} \subset \mathbb{R}^{p \times p}$; and $\{X_k : k \in \mathbb{N}\}$ are independent and identically distributed copies of X. Importantly, $\{M_k\}$ cannot be arbitrary, and the following properties specify a generalization of the Robbins and Monro [1951] conditions for matrix-valued learning rates [c.f. Patel, 2021].

The first condition requires a positive learning rate, and imposes symmetry to ensure the existence of real eigenvalues.

Property 1. $\{M_k : k + 1 \in \mathbb{N}\}$ are symmetric, positive definite matrices.

The next two properties are a natural generalization of the Robbins-Monro conditions. Let $\alpha \in (0, 1]$ be as in Assumption 2.

Property 2. Let $\lambda_{\max}(\cdot)$ denote the largest eigenvalue of a symmetric, positive definite matrix. Then, $\sum_{k=0}^{\infty} \lambda_{\max}(M_k)^{1+\alpha} =: S < \infty$.

Property 3. Let $\lambda_{\min}(\cdot)$ denote the smallest eigenvalue of a symmetric, positive definite matrix. Then, $\sum_{k=0}^{\infty} \lambda_{\min}(M_k) = \infty$.

We will show that Properties 1 to 3 are sufficient for a global convergence result (see Theorem 2).

The final property ensures the stability of the condition number of $\{M_k\}$. Note, this property is readily satisfied for scalar learning rates satisfying the Robbins-Monro conditions.

Property 4. Let $\kappa(\cdot)$ denote the ratio of the largest and smallest eigenvalues of a symmetric, positive definite matrix. Then, $\lim_{k\to\infty} \lambda_{\max}(M_k)^{\alpha} \kappa(M_k) = 0$.

We will show that Properties 1 to 4 are sufficient for stability (see Theorem 3).

---

4 Global Convergence & Stability

With the stochastic optimization problem and with stochastic gradient descent (SGD) specified, we now turn our attention to what happens when SGD is applied to a stochastic optimization problem. The key step in the analysis of SGD on any objective function is to establish a bound between the optimality gap at θ_{k+1} with that of θ_k. This step is achieved by using the local Hölder continuity of the gradient function and the fundamental theorem of calculus. Using Assumption 2, we first specify the local Hölder constant.

Definition 1. For any $\theta, \varphi \in \mathbb{R}^p$, define

$$L(\theta, \varphi) = \sup_{\psi} \left\{ \frac{\left\| \dot{F}(\psi) - \dot{F}(\theta) \right\|_2}{\|\psi - \theta\|_2^{\alpha}} : \psi \in \overline{B(\theta, \|\varphi - \theta\|_2)} \right\}, \tag{5}$$

where $B(\theta, r)$ is an open ball around θ of radius $r > 0$. Moreover, for any $R \geq 0$, let L_R be the supremum of $L(\theta, \varphi)$ for any distinct $\theta, \varphi \in \overline{B(0, R)}$.

Remark 5. Note, when the gradient is locally Hölder continuous, L_R is finite for any $R \geq 0$.

With this definition, we can now relate the optimality gap of θ_{k+1} with that of θ_k by using the following result and proved in Appendix B.

Lemma 1. *Suppose Assumptions 1 and 2 hold. Then, for any $\theta, \varphi \in \mathbb{R}^p$,*

$$F(\varphi) - F_{l.b.} \leq F(\theta) - F_{l.b.} + \dot{F}(\theta)'(\varphi - \theta) + \frac{L(\theta, \varphi)}{1 + \alpha} \|\varphi - \theta\|_2^{1+\alpha}. \tag{6}$$

Now, if we simply set $\varphi = \theta_{k+1}$ and $\theta = \theta_k$ in Lemma 1 and try to take expectations to manage the randomness of the stochastic gradient, we will run into the problem that $L(\theta_k, \theta_{k+1})$ and $\|\theta_{k+1} - \theta_k\|_2$ are dependent, and we cannot compute its expectation. In previous work, this technical challenge is waived away by using a global Hölder constant to upper bound $L(\theta_k, \theta_{k+1})$, which is unrealistic even for simple problems (see Section 1).

To address this technical challenge, we innovate two new strategies for handling the dependence between $L(\theta_k, \theta_{k+1})$ and $\|\theta_{k+1} - \theta_k\|_2$. In both strategies, we follow the same general approach:

1. We begin by restricting our analysis to specific events, which will allow us to decouple $L(\theta_k, \theta_{k+1})$ and $\|\theta_{k+1} - \theta_k\|_2$.
2. With these two quantities decoupled, we will develop a recurrence relationship between the optimality gap at θ_{k+1} and that of θ_k.
3. We apply this recurrence relationship with refinements of standard arguments or new ones to derive the desired property about the objective function.

220   4. Finally, we state the generality of the specific events on which we have studied SGD's
221   iterates.

222 Thus, it follows, we will define two distinct series of events for the two strategies. The first strategy,
223 which we refer to as the pseudo-global strategy, will provide the global convergence analysis. The
224 second strategy, which we refer to as the local strategy, will provide the stability result.

4.1   Pseudo-Global Strategy and Global Convergence Analysis

226 For the first strategy, which supplies the global convergence result, we study SGD on the events

$$\mathcal{B}_k(R) := \bigcap_{j=0}^{k} \left\{ \|\theta_j\|_2 \le R \right\}, \; k+1 \in \mathbb{N}, \tag{7}$$

227 for every $R \ge 0$. We now try to control the optimality gap at iteration $k+1$ with that of iteration k,
228 which will result in two cases.

229   1. (Case 1) $\mathcal{B}_{k+1}(R)$ holds. We can bound $L(\theta_k, \theta_{k+1})$ by L_R, and $G(\theta)$ is also bounded in
230   the ball of radius R about the origin (which follows from G being upper semi-continuous
231   in Assumption 4). As a result, we could then proceed with the analysis in a manner that is
232   similar to having a global Hölder constant.
233   2. (Case 2) $\|\theta_{k+1}\|_2 > R$ and $\mathcal{B}_k(R)$ holds. In this case, controlling $L(\theta_k, \theta_{k+1})$ is very
234   challenging and, to our knowledge, was not solved before our work.

235 Our approach for controlling the optimality gap in both cases is supplied in the next lemma, whose
236 proof is in Appendix C.

237 **Lemma 2.** *Let $\{M_k\}$ satisfy Property 1. Suppose Assumptions 1 to 4 hold. Let $\{\theta_k\}$ satisfy (4).*
238 *Then, $\forall R \ge 0$,*

$$\mathbb{E}\left[[F(\theta_{k+1}) - F_{l.b.}] \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] | \mathcal{F}_k \right] \le [F(\theta_k) - F_{l.b.}] \mathbf{1}\left[\mathcal{B}_k(R)\right]$$
$$- \lambda_{\min}(M_k) \left\| \dot{F}(\theta_k) \right\|_2^2 \mathbf{1}\left[\mathcal{B}_k(R)\right] + \frac{L_{R+1} + \partial F_R}{1 + \alpha} \lambda_{\max}(M_k)^{1+\alpha} G_R, \tag{8}$$

239 *where $G_R = \sup_{\theta \in \overline{B(0,R)}} G(\theta) < \infty$ with $G(\theta)$; and $\partial F_R = \sup_{\theta \in \overline{B(0,R)}} \|\dot{F}(\theta)\|_2 (1+\alpha) < \infty$.*

240 With this recursion and standard martingale results [Robbins and Siegmund, 1971, Neveu and Speed,
241 1975, Exercise II.4], the limit of $[F(\theta_k) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_k(R)\right]$ exists with probability one and is finite for
242 every $R \ge 0$. As a result, the limit of $F(\theta_k) - F_{l.b.}$ exists and is finite on the event $\{\sup_k \|\theta_k\|_2 < \infty\}$
243 (see Corollary 1).

244 We can also use Lemma 2 to make a statement about the gradient. Specifically, we can show that the
245 limit infimum of $\mathbb{E}[\|\dot{F}(\theta_k)\|_2^2 \mathbf{1}\left[\mathcal{B}_k(R)\right]]$ must be zero, which is now a standard argument that mimics
246 Zoutendijk's theorem [Nocedal and Wright, 2006, Theorem 3.2]. By Markov's inequality, this result
247 implies that $\|\dot{F}(\theta_k)\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right]$ gets arbitrarily close to 0 infinitely often (see Lemma 9). To show
248 convergence to zero, however, is not standard. Several strategies have been developed, namely those
249 of Li and Orabona [2019], Lei et al. [2019], Mertikopoulos et al. [2020], Patel [2021]. Unfortunately,
250 the approaches of Li and Orabona [2019], Lei et al. [2019] rely intimately on the existence of a global
251 Hölder constant, while that of Mertikopoulos et al. [2020] requires even more restrictive assumptions.
252 Fortunately, the approach of Patel [2021] can be improved and generalized to the current context (see
253 Lemma 10). Thus, we show that $\lim_{k \to \infty} \|\dot{F}(\theta_k)\|_2 = 0$ on $\{\sup_k \|\theta_k\|_2 < \infty\}$ (see Corollary 2).

254 Our final step is to clarify the role of $\{\sup_k \|\theta_k\|_2 < \infty\}$ in the asymptotic behavior of SGD's iterates.
255 At first glance, this event seems to imply that the iterates converge to a point. However, owing to the
256 general nature of the noise, it is also possible, say, that the iterates approach a limit cycle or oscillate
257 between points with the same norm. Even beyond this event, the generality of the noise model may
258 allow for substantial excursions between $F_{l.b.}$ and infinity (c.f., a simple random walk, which has a
259 limit supremum of infinity and a limit infinimum of negative infinity). Thankfully, we can prove that
260 either the iterates converge to a point or they must diverge—a result that we refer to as the Capture
261 Theorem (see Appendix C).

262 **Theorem 1** (Capture Theorem)**.** *Let $\{\theta_k\}$ be defined as in (4), and let $\{M_k\}$ satisfy Properties 1*
263 *and 2. If Assumption 4 holds, then either $\{\lim_{k \to \infty} \theta_k \text{ exists}\}$ or $\{\liminf_{k \to \infty} \|\theta_k\|_2 = \infty\}$ must*
264 *occur.*

By putting together the above arguments and results, we can conclude that either SGD's iterates diverge or SGD's iterates converge to a stationary point. This is formally stated in the following theorem. See Section 1 for a discussion of the practical value of this result.

Theorem 2 (Global Convergence). *Let θ_0 be arbitrary, and let $\{\theta_k : k \in \mathbb{N}\}$ be defined according to (4) with $\{M_k : k + 1\}$ satisfying Properties 1 to 3. Suppose Assumptions 1 to 4 hold. Let $\mathcal{A}_1 = \{\liminf_{k \to \infty} \|\theta_k\|_2 = \infty\}$ and $\mathcal{A}_2 = \{\lim_{k \to \infty} \theta_k \text{ exists}\}$. Then, the following statements hold.*

1. $\mathbb{P}[\mathcal{A}_1] + \mathbb{P}[\mathcal{A}_2] = 1$.
2. On $\mathcal{A}_2$, there exists a finite random variable, $F_{\lim}$, such that $\lim_{k \to \infty} F(\theta_k) = F_{\lim}$ and $\lim_{k \to \infty} \dot{F}(\theta_k) = 0$ with probability one.

Proof. By Theorem 1, we have that $\mathbb{P}[\mathcal{A}_1] + \mathbb{P}[\mathcal{A}_2] = 1$. Then, on $\mathcal{A}_2$, Corollaries 1 and 2 imply that $F(\theta_k) \to F_{\lim}$, which is finite, and $\dot{F}(\theta_k) \to 0$. $\qquad\square$

4.2 Local Strategy and Stability Analysis

While Theorem 2 provides a complete global convergence result, it allows for the possibility of diverging iterates. The possibility of divergent iterates raises the spectre of whether the objective function can also diverge along this sequence. That is, there is a possibility that SGD may be unstable, which would be highly unexpected and undesirable, especially when the objective function is coercive (e.g., has an ℓ^1 penalty on the parameters). To formalize this concept, we define a relevant notion of stability.

Definition 2. Stochastic Gradient Descent is stable if

$$\mathbb{P}\left[\limsup_{k \to \infty} F(\theta_k) = \infty\right] = 0, \tag{9}$$

where $\{\theta_k\}$ satisfy (4).

We now state the events that we will use to decouple the relationship between $L(\theta_k, \theta_{k+1})$ and $\|\theta_{k+1} - \theta_k\|_2$. Unlike in our pseudo-global strategy, we will make use of two sequences of events that are closely related. To define these sequences, we will first need to define stopping times. For every $j + 1 \in \mathbb{N}$, define

$$\tau_j = \min\left\{ k : \begin{array}{l} F(\theta_{k+1}) - F_{l.b.} > F(\theta_k) - F_{l.b.} + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) \\ + \dfrac{\mathcal{L}_\epsilon(\theta_k)}{1 + \alpha} \|\theta_{k+1} - \theta_k\|_2^{1+\alpha}, \text{ and } k > j \end{array} \right\}. \tag{10}$$

Analogously, for every $j + 1 \in \mathbb{N}$, define

$$\nu_j = \min\left\{k : L(\theta_k, \theta_{k+1}) > \mathcal{L}_\epsilon(\theta_k), \text{ and } k > j\right\}. \tag{11}$$

Now, we will use (10) to establish the stability of the objective function, and we will use (11) to show that the gradient function must tend to zero. Just as we did with $\mathcal{B}_k(R)$, we will derive a recursion on the optimality gap over the events $\{\{\tau_j > k\} : k + 1 \in \mathbb{N}\}$. Of course, just as before, the main challenge in deriving a recursive formula is to address $\{\tau_j = k\}$. Our solution is supplied in the following lemma, whose proof is in Appendix D.

Lemma 3. *Let $\{M_k\}$ satisfy Property 1. Suppose Assumptions 1 to 4 hold. Let $\{\theta_k\}$ satisfy (4). Then, for any $j + 1 \in \mathbb{N}$ and $k > j$,*

$$\mathbb{E}\left[(F(\theta_{k+1}) - F_{l.b.})\, \boldsymbol{1}\left[\tau_j > k\right] \middle| \mathcal{F}_k\right] \leq \left(F(\theta_k) - F_{l.b.} - \dot{F}(\theta_k)' M_k \dot{F}(\theta_k)\right) \boldsymbol{1}\left[\tau_j > k - 1\right]$$

$$+ \frac{\lambda_{\max}(M_k)^{1+\alpha}}{1 + \alpha}\left[\mathcal{L}_\epsilon(\theta_k) G(\theta_k) + \alpha \left[\frac{\left\|\dot{F}(\theta_k)\right\|_2^{1+\alpha}}{\mathcal{L}_\epsilon(\theta_k)}\right]^{1/\alpha}\right] \boldsymbol{1}\left[\tau_j > k - 1\right]. \tag{12}$$

From Lemma 3, there is a clear motivation for Assumption 5. Indeed, if we apply Assumption 5, Lemma 3 produces the following simple recursive relationship.

Lemma 4. *If Assumptions 1 to 5, and Properties 1 and 4 hold, and $\{\theta_k\}$ satisfy (4), then there exists a $K \in \mathbb{N}$ such that for any $j + 1 \in \mathbb{N}$ and any $k \geq \min\{K, j+1\}$,*

$$
\begin{aligned}
&\mathbb{E}\left[(F(\theta_{k+1}) - F_{l.b.})\boldsymbol{I}\left[\tau_j > k \right] | \mathcal{F}_k \right] \\
&\leq \left(1 + \lambda_{\max}(M_k)^{1+\alpha} \frac{C_2}{1+\alpha} \right) (F(\theta_k) - F_{l.b.})\boldsymbol{I}\left[\tau_j > k - 1 \right] \\
&\quad - \frac{1}{2}\lambda_{\min}(M_k) \left\| \dot{F}(\theta_k) \right\|_2^2 \boldsymbol{I}\left[\tau_j > k - 1 \right] + \lambda_{\max}(M_k)^{1+\alpha}\frac{C_1}{1+\alpha}.
\end{aligned}
\tag{13}
$$

Just as in the pseudo-global strategy, Lemma 4 can be combined with standard martingale results [Robbins and Siegmund, 1971, Neveu and Speed, 1975, Exercise II.4] to conclude that the limit of $F(\theta_k)$ exists and is finite on the event $\cup_{j=0}^{\infty}\{\tau_j = \infty\}$ (see Corollary 3). Also as in the pseudo-global strategy, by improving on the arguments in Patel [2021], we show that $\lim_k \dot{F}(\theta_k) = 0$ on the event $\cup_{j=0}^{\infty}\{\nu_j = \infty\}$ (see Corollary 4).

Finally, we show that $\cup_{j=0}^{\infty}\{\tau_j = \infty\}$ and $\cup_{j=0}^{\infty}\{\nu_j = \infty\}$ are probability one events (see Theorem 5). In other words, we show that eventually $\mathcal{L}_\epsilon(\theta_k)$ will always dominate $L(\theta_k, \theta_{k+1})$. This statement should not come as a surprise on the event $\{\lim_k \theta_k \text{ exists}\}$, but is slightly surprising that it must also hold on $\{\lim_k \|\theta_k\|_2 = \infty\}$. By combining these results, we can conclude as follows.

Theorem 3 (Stability). *Let θ_0 be arbitrary, and let $\{\theta_k : k \in \mathbb{N}\}$ be defined according to (4) with $\{M_k : k+1\}$ satisfying Properties 1 to 3. Suppose Assumptions 1 to 5 hold. Then,*

1. *There exists a finite random variable, $F_{\lim}$, such that $\lim_{k \to \infty} F(\theta_k) = F_{\lim}$ with probability one;*
2. *$\lim_{k \to \infty} \dot{F}(\theta_k) = 0$ with probability one.*

Proof. Using Corollary 3, we conclude that $\exists F_{\lim}$ that is finite such that $\lim_k F(\theta_k) = F_{\lim}$ on $\cup_{j=0}^{\infty}\{\tau_j = \infty\}$. Using Corollary 4, we conclude that $\lim_k \dot{F}(\theta_k) = 0$ on $\cup_{j=0}^{\infty}\{\nu_j = \infty\}$. Finally, we apply Theorem 5 to conclude that $\mathbb{P}[\cup_{j=0}^{\infty}\{\nu_j = \infty\}] = \mathbb{P}[\cup_{j=0}^{\infty}\{\tau_j = \infty\}] = 1$. $\square$

5 Conclusion

In this work, we studied the global convergence analysis of Stochastic Gradient Descent with diminishing step size. We began our discussion by producing three simple problems for which the common assumptions (i.e., global Hölder continuity, bounded variance) in the SGD literature are violated, and, to our knowledge, there does not exist theory that covers the two simple neural network problems. That is, prior to our work, it was unknown what SGD with arbitrary initialization and diminishing step sizes will do on these two simple neural network problems, which raised the question of what SGD is doing on more complicated learning problems.

Motivated by our example problems, we considered a more general set of assumptions (see Assumptions 2 and 4). Given the generality of our assumptions, we needed to develop a new analysis technique that is of interest beyond this work, and we proved that SGD's iterates either converge to a stationary point or diverge. Thus, we now know how SGD with arbitrary initialization and diminishing step sizes will do on a much larger class of learning problems.

We note that we do not provide rate of convergence results mainly because it is *impossible* for the broad class of functions admitted by our assumptions [Wolpert and Macready, 1997]. We stress that global rates of convergence (e.g., complexity statements) results that exist do not apply to the two simple neural network problems that we supplied at the beginning of this work without making more restrictive assumptions.

We also studied what happens when SGD's iterates can diverge. To this end, we required an additional assumption under which we developed another novel analysis technique and showed that, regardless of SGD's iterates' behavior, the objective function will converge to a finite random variable and the gradient function converges to zero with probability one. Unfortunately, we make an assumption that we were not able to interpret, but we will leave this to future work.

References

H. Asi and J. C. Duchi. Stochastic (approximate) proximal point methods: Convergence, optimality, and adaptivity. *SIAM Journal on Optimization*, 29(3):2257–2290, 2019.

R. Bassily, M. Belkin, and S. Ma. On exponential convergence of sgd in non-convex over-parametrized learning. *arXiv preprint arXiv:1811.02564*, 2018.

J. Bi and S. R. Gunn. A stochastic gradient method with biased estimation for faster nonconvex optimization. In *Pacific Rim International Conference on Artificial Intelligence*, pages 337–349. Springer, 2019.

P. Bianchi, W. Hachem, and S. Schechtman. Convergence of constant step stochastic gradient descent for non-smooth non-convex functions. *Set-Valued and Variational Analysis*, pages 1–31, 2022.

L. Bottou, F. E. Curtis, and J. Nocedal. Optimization methods for large-scale machine learning. *Siam Review*, 60(2):223–311, 2018.

H. Chen, L. Zheng, R. Al Kontar, and G. Raskutti. Stochastic gradient descent in correlated settings: A study on gaussian processes. In *NeurIPS*, 2020.

C. Fang, Z. Lin, and T. Zhang. Sharp analysis for nonconvex sgd escaping from saddle points. In *Conference on Learning Theory*, pages 1192–1234. PMLR, 2019.

R. M. Gower, O. Sebbouh, and N. Loizou. Sgd for structured nonconvex functions: Learning rates, minibatching and interpolation. *arXiv preprint arXiv:2006.10311*, 2020.

M. Hardt, T. Ma, and B. Recht. Gradient descent learns linear dynamical systems. *arXiv preprint arXiv:1609.05191*, 2016.

W. Hu, C. J. Li, L. Li, and J.-G. Liu. On the diffusion approximation of nonconvex stochastic gradient descent. *Annals of Mathematical Sciences and Applications*, 4(1), 2019.

C. Jin, P. Netrapalli, R. Ge, S. M. Kakade, and M. I. Jordan. On nonconvex optimization for machine learning: Gradients, stochasticity, and saddle points. *Journal of the ACM (JACM)*, 68(2):1–29, 2021.

A. Khaled and P. Richtárik. Better theory for sgd in the nonconvex world. *arXiv preprint arXiv:2002.03329*, 2020.

H.-S. Kim, L. Zhang, A. Bienkowski, and K. R. Pattipati. Multi-pass sequential mini-batch stochastic gradient descent algorithms for noise covariance estimation in adaptive kalman filtering. *IEEE Access*, 9:99220–99234, 2021.

Y. Lei, T. Hu, G. Li, and K. Tang. Stochastic gradient descent for nonconvex learning without bounded gradient assumptions. *IEEE transactions on neural networks and learning systems*, 31(10):4394–4400, 2019.

X. Li and F. Orabona. On the convergence of stochastic gradient descent with adaptive stepsizes. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 983–992. PMLR, 2019.

Y. Ma and D. Klabjan. Convergence analysis of batch normalization for deep neural nets. *arXiv preprint arXiv:1705.08011*, 2017.

S. Majewski, B. Miasojedow, and E. Moulines. Analysis of nonsmooth stochastic approximation: the differential inclusion approach. *arXiv preprint arXiv:1805.01916*, 2018.

P. Mertikopoulos, N. Hallak, A. Kavis, and V. Cevher. On the almost sure convergence of stochastic gradient descent in non-convex problems. *arXiv preprint arXiv:2006.11144*, 2020.

J. Neveu and T. Speed. *Discrete-parameter martingales*, volume 10. North-Holland Amsterdam, 1975.

J. Nocedal and S. Wright. *Numerical optimization*. Springer Science & Business Media, 2006.

V. Patel. Stopping criteria for, and strong convergence of, stochastic gradient descent on bottou-curtis-nocedal functions. *Mathematical Programming*, pages 1–42, 2021.

S. Reddi, S. Sra, B. Poczos, and A. J. Smola. Proximal stochastic methods for nonsmooth nonconvex finite-sum optimization. *Advances in neural information processing systems*, 29:1145–1153, 2016a.

S. J. Reddi, A. Hefny, S. Sra, B. Poczos, and A. Smola. Stochastic variance reduction for nonconvex optimization. In *International conference on machine learning*, pages 314–323. PMLR, 2016b.

H. Robbins and S. Monro. A stochastic approximation method. *The annals of mathematical statistics*, pages 400–407, 1951.

H. Robbins and D. Siegmund. A convergence theorem for non negative almost supermartingales and some applications. In *Optimizing methods in statistics*, pages 233–257. Elsevier, 1971.

H. Wang, M. Gürbüzbalaban, L. Zhu, U. Şimşekli, and M. A. Erdogdu. Convergence rates of stochastic gradient descent under infinite noise variance. *arXiv preprint arXiv:2102.10346*, 2021.

Y. Wang and S. Wu. Asymptotic analysis via stochastic differential equations of gradient descent algorithms in statistical and computational paradigms. *Journal of machine learning research*, 21, 2020.

D. H. Wolpert and W. G. Macready. No free lunch theorems for optimization. *IEEE transactions on evolutionary computation*, 1(1):67–82, 1997.

S. Zhang and V. Patel. Stochastic approximation for high-frequency observations in data assimilation. *arXiv preprint arXiv:2011.02672*, 2020.

D. Zhou, P. Xu, and Q. Gu. Stochastic nested variance reduction for nonconvex optimization. *arXiv preprint arXiv:1806.07811*, 2018.

F. Zou, L. Shen, Z. Jie, W. Zhang, and W. Liu. A sufficient condition for convergences of adam and rmsprop. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pages 11127–11135, 2019.

A Details for Counter Examples

A.1 Simple Linear Regression

Here, we prove Proposition 1 for a rather simple problem. Let $Z \in \mathbb{R}$ be a random variable such that $\mathbb{E}\left[Z^2\right] = 1$ and $\mathbb{E}\left[Z^4\right] = 2$. Moreover, let ϵ be an independent random variable with mean zero and variance 1. Finally, let $\theta^* \in \mathbb{R}$ and define $Y = Z\theta^* + \epsilon$. Consider the estimation problem of minimizing $F(\theta)$ where

$$F(\theta) = \frac{1}{2}\mathbb{E}\left[(Z\theta - Y)^2\right] = \frac{1}{2}(\theta - \theta^*)^2 + \frac{1}{2}. \tag{14}$$

Letting $X = (Y, Z)$, let $f(\theta, X) = 0.5(Z\theta - Y)^2$. Now, the variance of $\dot{f}(\theta, X)$ is

$$\mathbb{E}\left[((Z^2 - 1)(\theta - \theta^*) - z\epsilon)^2\right]$$
$$= \mathbb{E}\left[(Z^2 - 1)^2\right](\theta - \theta^*)^2 - 2(\theta - \theta^*)\mathbb{E}\left[(Z^2 - 1)Z\epsilon\right] + \mathbb{E}\left[Z^2\epsilon\right] \tag{15}$$
$$= (\theta - \theta^*)^2 + 1. \tag{16}$$

Clearly, the variance scales with the error in the parameter, which violates the common bounded noise model assumption. In particular, as $|\theta| \to \infty$, the variance diverges.

On the other hand, the simple linear regression problem does satisfy our assumptions. In particular,

1. Assumptions 1 and 3 are easily verified.

2. Given that $\dot{F}$ is globally Lipschitz continuous, it is locally Lipschitz continuous. Therefore, Assumption 2 is satisfied.

3. From the variance calculation of $\dot{f}(\theta, X)$, we conclude

$$\mathbb{E}\left[\dot{f}(\theta, X)^2\right] = 2(\theta - \theta^*)^2 + 1, \tag{17}$$

   which is a continuous function. Hence, Assumption 4 is satisfied.

A.2 Feed Forward Network for Binary Classification

We now prove Proposition 2. Consider the binary classification problem with label Y and feature Z where $(Y, Z) = (0, 0)$ with probability $1/2$ and $(Y, Z) = (1, 1)$ with probability $1/2$. We solve this classification problem using the network shown in Fig. 1 with σ linear and φ sigmoid. We will train this model using the binary cross entropy loss function. Letting $X = (Y, Z)$ and $\theta = (W_1, W_2, W_3, W_4)$,

$$f(\theta, X) = -Y\log(\hat{y}) - (1 - Y)\log(1 - \hat{y}) + \frac{1}{2}\sum_{i=1}^{4} W_i^2, \tag{18}$$

and

$$\hat{y} = \begin{cases} \frac{1}{2} & Z = 0 \\ \frac{1}{1 + \exp(-W_4 W_3 W_2 W_1 Z)} & Z = 1. \end{cases} \tag{19}$$

Moreover,

$$\dot{f}(\theta, X) = \begin{cases} \begin{bmatrix} W_1 \\ W_2 \\ W_3 \\ W_4 \end{bmatrix} & (Y, Z) = (0, 0) \\[2em] \frac{-1}{1 + \exp(W_4 W_3 W_2 W_1)} \begin{bmatrix} W_4 W_3 W_2 \\ W_4 W_3 W_1 \\ W_4 W_2 W_1 \\ W_3 W_2 W_1 \end{bmatrix} + \begin{bmatrix} W_1 \\ W_2 \\ W_3 \\ W_4 \end{bmatrix} & (Y, Z) = (1, 1), \end{cases} \tag{20}$$

and, consequently,

$$\dot{F}(\theta) = \frac{-1/2}{1 + \exp(W_4 W_3 W_2 W_1)} \begin{bmatrix} W_4 W_3 W_2 \\ W_4 W_3 W_1 \\ W_4 W_2 W_1 \\ W_3 W_2 W_1 \end{bmatrix} + \begin{bmatrix} W_1 \\ W_2 \\ W_3 \\ W_4 \end{bmatrix}. \tag{21}$$

We first establish that $\dot{F}(\theta)$ is not globally Lipschitz continuous. With $\theta = (1, -1, W_3, W_3)$ and $\phi = (1, 0, 0, 0)$, it is enough to find a lower bound for the first component of $\dot{F}(\theta) - \dot{F}(\phi)$, denoted by $\dot{F}_1(\theta) - \dot{F}_1(\phi)$. To this end,

$$|\dot{F}_1(\theta) - \dot{F}_1(\phi)| = \frac{0.5 W_3^2}{1 + \exp(-W_3^2)} \geq \frac{1}{4}|W_3 - 0|^2. \tag{22}$$

Thus, $\dot{F}$ is not globally Lipschitz.

To show that the variance is not bounded, we study the variance of the first component of $\dot{f}(\theta, X)$ which we denote by $\dot{f}_1(\theta, X)$. By direct calculation,

$$\mathbb{E}\left[(\dot{f}_1(\theta, X) - \dot{F}_1(\theta))^2\right] = \frac{1}{4} \frac{W_4^2 W_3^2 W_2^2}{[1 + \exp(W_4 W_3 W_2 W_1)]^2}. \tag{23}$$

We again consider $\theta = (1, -1, W_3, W_3)$, then the variance at this value of θ is

$$\frac{1}{4} \frac{W_3^4}{[1 + \exp(-W_3^2)]^2} \geq \frac{1}{16} W_3^4. \tag{24}$$

Therefore, as $W_3 \to \infty$, the variance goes to infinity. That is, the variance of the stochastic gradients is unbounded.

On the other hand, the problem does satisfy our assumptions. In particular,

1. Assumptions 1 and 3 are easily verified.

2. Given that $\dot{F}$ is continuously differentiable, then compactness and continuity of the derivative of $\dot{F}$ imply that it is locally Lipschitz continuous. Therefore, Assumption 2 is satisfied.

3. Given the computation of the variance for the first component, we have $\mathbb{E}[\dot{f}_1(\theta, X)^2]$ is

$$\frac{1}{4} \frac{W_4^2 W_3^2 W_2^2}{[1 + \exp(W_4 W_3 W_2 W_1)]^2} + \dot{F}_1(\theta)^2, \tag{25}$$

which is a continuous function. By repeating this argument for each component, we conclude that Assumption 4 is satisfied.

A.3 Recurrent Neural Network for Binary Classification

Consider observing one of two sequences $(1, 0, 0, 0)$ or $(0, 0, 0, 0)$ with equal probabilities, and suppose that each sequence corresponds to the label 1 or 0, respectively. Now consider Fig. 2 to be a 1-dimensional linear recurrent neural network which reads each element of the sequence and uses a logistic output layer to predict either a label of one or zero. If we fix $H_0 = 0$, $W_2 = 1$ and $W_3 = 1$, then the model predicts the probability of a 1 label as

$$\hat{y}(Z_0, Z_1, Z_2, Z_3) = \frac{\exp(W_1^3 Z_0)}{1 + \exp(W_1^3 Z_0)}. \tag{26}$$

If we use the binary cross entropy loss with ℓ^2 regularization, and let $X = (Y, Z_0, Z_1, Z_2, Z_3)$ and $\theta = W_1$ then

$$f(\theta, X) = -Y \log \hat{y}(Z_0, Z_1, Z_2, Z_3) - (1 - Y) \log[1 - \hat{y}(Z_0, Z_1, Z_2, Z_3)] + \frac{1}{2}\theta^2 \tag{27}$$

$$= -Y\left[\theta^3 Z_0 - \log(1 + \exp(\theta^3 Z_0))\right] + (1 - Y)\log(1 + \exp(\theta^3 Z_0)) + \frac{1}{2}\theta^2 \tag{28}$$

$$= -\theta^3 Z_0 Y + \log(1 + \exp(\theta^3 Z_0)) + \frac{1}{2}\theta^2, \tag{29}$$

and

$$\dot{f}(\theta, X) = -3\theta^2 Z_0 Y + \frac{3\theta^2 Z_0 \exp(\theta^3 Z_0)}{1 + \exp(\theta^3 Z_0)} + \theta. \tag{30}$$

Taking the expectations, we compute

$$F(\theta) = \frac{1}{2} \left[\log(2) + \log(1 + \exp(\theta^3)) - \theta^3 + \theta^2 \right], \tag{31}$$

and

$$\dot{F}(\theta) = \frac{-3\theta^2}{2} \frac{1}{1 + \exp(\theta^3)} + \theta. \tag{32}$$

From this calculation alone, it is easy to see that as $\theta \to -\infty$, $\dot{F}(\theta) \propto -\theta^2$, which is not a Lipschitz continuous function. Moreover, computing the variance of $\dot{f}(\theta, X)$, we recover

$$\mathbb{E}\left[(\dot{f}(\theta, X) - \dot{F}(\theta))^2\right] = \frac{9\theta^4}{4(1 + \exp(\theta^3))^2}, \tag{33}$$

which does not satisfy a bounded variance assumption. Thus, any work that makes either a global Lipschitz bound on the gradient or a global noise model bound fails to apply to this simple recurrent neural network training problem.

On the other hand, the problem does satisfy our assumptions. In particular,

1. Assumptions 1 and 3 are easily verified.

2. Given that $\dot{F}$ is continuously differentiable, then compactness and continuity of the derivative of $\dot{F}$ imply that it is locally Lipschitz continuous. Therefore, Assumption 2 is satisfied.

3. Moreover,

$$\mathbb{E}\left[\dot{f}(\theta, X)^2\right] = \frac{9\theta^4}{4(1 + \exp(\theta^3))^2} + \dot{F}(\theta)^2, \tag{34}$$

which is a continuous function. Hence, Assumption 4 is satisfied.

A.4 Noiseless Feed Forward Network for Binary Classification

Out of interest, we reconsider the second example but construct a different data distribution that produces noiseless stochastic gradient. Consider the binary classification problem with label Y and feature Z where $(Y, Z) = (0, -1)$ with probability $1/2$ and $(Y, Z) = (1, 1)$ with probability $1/2$. We solve this classification problem using the network shown in Fig. 1 with σ linear and φ sigmoid. We will train this model using the binary cross entropy loss function. Letting $X = (Y, Z)$ and $\theta = (W_1, W_2, W_3, W_4)$,

$$f(\theta, X) = -Y \log(\hat{y}) - (1 - Y) \log(1 - \hat{y}) + \frac{1}{2} \sum_{i=1}^{4} W_i^2, \tag{35}$$

and

$$\hat{y} = \frac{1}{1 + \exp(-W_4 W_3 W_2 W_1 Z)}. \tag{36}$$

Moreover,

$$\dot{f}(\theta, X) = Z(\hat{y} - Y) \begin{bmatrix} W_4 W_3 W_2 \\ W_4 W_3 W_1 \\ W_4 W_2 W_1 \\ W_3 W_2 W_1 \end{bmatrix} + \begin{bmatrix} W_1 \\ W_2 \\ W_3 \\ W_4 \end{bmatrix}, \tag{37}$$

and, consequently,

$$\dot{F}(\theta) = \frac{-1}{1 + \exp(W_4 W_3 W_2 W_1)} \begin{bmatrix} W_4 W_3 W_2 \\ W_4 W_3 W_1 \\ W_4 W_2 W_1 \\ W_3 W_2 W_1 \end{bmatrix} + \begin{bmatrix} W_1 \\ W_2 \\ W_3 \\ W_4 \end{bmatrix}. \tag{38}$$

We first establish that $\dot{F}(\theta)$ is not globally Lipschitz continuous. With $\theta = (1, -1, W_3, W_3)$ and $\phi = (1, 0, 0, 0)$, it is enough to find a lower bound for the first component of $\dot{F}(\theta) - \dot{F}(\phi)$, denoted by $\dot{F}_1(\theta) - \dot{F}_1(\phi)$. To this end,

$$|\dot{F}_1(\theta) - \dot{F}_1(\phi)| = \frac{W_3^2}{1 + \exp(-W_3^2)} \geq \frac{1}{2}|W_3 - 0|^2. \tag{39}$$

Thus, $\dot{F}$ is not globally Lipschitz.

On the other hand, the problem does satisfy our assumptions. In particular,

1. Assumptions 1 and 3 are easily verified.

2. Given that $\dot{F}$ is continuously differentiable, then compactness and continuity of the derivative of $\dot{F}$ imply that it is locally Lipschitz continuous. Therefore, Assumption 2 is satisfied.

3. Moreover, $\dot{f}(\theta, Z) = \dot{F}(\theta)$—that is, there $\dot{f}(\theta, Z)$ has zero variance for the distribution that we have constructed. Therefore,

$$\mathbb{E}\left[\dot{f}(\theta, X)^2\right] = \dot{F}(\theta)^2, \tag{40}$$

which is a continuous function. Hence, Assumption 4 is satisfied.

B   Technical Lemmas

Lemma 5 (Lemma 1). *Suppose Assumptions 1 and 2 hold. Then, for any $\theta, \varphi \in \mathbb{R}^p$,*

$$F(\varphi) - F_{l.b.} \leq F(\theta) - F_{l.b.} + \dot{F}(\theta)'(\varphi - \theta) + \frac{L(\theta, \varphi)}{1 + \alpha} \|\varphi - \theta\|_2^{1+\alpha}. \tag{41}$$

Proof. By Taylor's theorem,

$$F(\varphi) - F_{l.b.} = F(\theta) - F_{l.b.} + \int_0^1 \dot{F}(\theta + t(\varphi - \theta))'(\varphi - \theta)dt. \tag{42}$$

Now, add and subtract $\dot{F}(\theta)$ to $\dot{F}(\theta + t(\varphi - \theta))$ in the integral, then apply Assumption 2. By Definition 1,

$$\left\|\dot{F}(\theta + t(\varphi - \theta)) - \dot{F}(\theta)\right\|_2 \leq L(\theta, \varphi)t^\alpha \|\theta - \varphi\|_2^\alpha. \tag{43}$$

We conclude,

$$F(\varphi) - F_{l.b.} \leq F(\theta) - F_{l.b.} + \dot{F}(\theta)'(\varphi - \theta) + L(\theta, \varphi) \|\varphi - \theta\|_2^{1+\alpha} \int_0^1 t^\alpha dt. \tag{44}$$

By computing the integral, the result follows. $\qquad\square$

Lemma 6. *Suppose $\{M_k : k + 1 \in \mathbb{N}\}$ satisfy Properties 1 and 4. Then $\forall C > 0$, $\exists K \in \mathbb{N}$ such that $\forall k \geq K$,*

$$\lambda_{\min}(M_k) - \frac{C}{2}\lambda_{\max}(M_k)^{1+\alpha} \geq \frac{1}{2}\lambda_{\min}(M_k). \tag{45}$$

Proof. Fix $C > 0$. Rearranging the conclusion, we see that it is equivalent to prove that $\exists K \in \mathbb{N}$ such that $\forall k \geq K$, $1/C \geq \lambda_{\max}(M_k)^\alpha \kappa(M_k)$. This follows from Property 4. $\qquad\square$

Lemma 7. *For any $\theta \in \mathbb{R}^p$, $v \in \mathbb{R}$, $L > 0$ and $\alpha \in (0, 1]$,*

$$\frac{L}{1 + \alpha}v^{1+\alpha} - \left\|\dot{F}(\theta)\right\|_2 v \geq -\frac{\alpha}{1 + \alpha}\left[\frac{\left\|\dot{F}(\theta)\right\|_2^{1+\alpha}}{L}\right]^{1/\alpha}. \tag{46}$$

Proof. If we minimize the left hand side of the inequality, we see that a minimum value occurs when $v^\alpha = \left\|\dot{F}(\theta)\right\|_2 / L \geq 0$. Solving for v and plugging this back into the left hand side, we conclude that the inequality holds. $\qquad\square$

 # C Global Convergence Analysis

545 We begin by first deriving a recursive relationship between the optimality gap at iteration $k + 1$ and
546 the optimality gap at iteration k on the events $\{\mathcal{B}_j(R)\}$ as defined in (7) for arbitrary $R \geq 0$. Using
547 this result, we then provide an analysis of the convergence of the objective function. Then, we turn
548 our attention to the gradient function. Note, $B(\theta, r)$ is the open ball around θ of radius r.

C.1 A Recursive Relationship

550 **Lemma 8** (Lemma 2). *Let $\{M_k\}$ satisfy Property 1. Suppose Assumptions 1 to 4 hold. Let $\{\theta_k\}$*
551 *satisfy (4). Then, $\forall R \geq 0$,*

$$
\mathbb{E}\left[\left[F(\theta_{k+1}) - F_{l.b.}\right]\boldsymbol{I}\left[\mathcal{B}_{k+1}(R)\right]\middle|\mathcal{F}_k\right] \leq \left[F(\theta_k) - F_{l.b.}\right]\boldsymbol{I}\left[\mathcal{B}_k(R)\right]
$$
$$
- \lambda_{\min}(M_k)\left\|\dot{F}(\theta_k)\right\|_2^2\boldsymbol{I}\left[\mathcal{B}_k(R)\right] + \frac{L_{R+1} + \partial F_R}{1 + \alpha}\lambda_{\max}(M_k)^{1+\alpha}G_R, \tag{47}
$$

552 *where $G_R = \sup_{\theta \in \overline{B(0,R)}} G(\theta) < \infty$ with $G(\theta)$; and $\partial F_R = \sup_{\theta \in \overline{B(0,R)}}\|\dot{F}(\theta)\|_2(1 + \alpha) < \infty$.*

553 *Proof.* Fix $R \geq 0$. For any $k + 1 \in \mathbb{N}$, the definition of local Hölder continuity implies that L_{R+1} is
554 well defined (see Definition 1). Therefore, Lemma 1 implies

$$
\left[F(\theta_{k+1}) - F_{l.b.}\right]\boldsymbol{1}\left[\mathcal{B}_{k+1}(R + 1)\right]
$$
$$
\leq \left(\left[F(\theta_k) - F_{l.b.}\right] + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{L_{R+1}}{1 + \alpha}\|\theta_{k+1} - \theta_k\|_2^{1+\alpha}\right)\boldsymbol{1}\left[\mathcal{B}_{k+1}(R + 1)\right]. \tag{48}
$$

555 Now, since $\overline{B(0, R)} \subset \overline{B(0, R + 1)}$, it also holds true that

$$
\left[F(\theta_{k+1}) - F_{l.b.}\right]\boldsymbol{1}\left[\mathcal{B}_{k+1}(R)\right]
$$
$$
\leq \left(\left[F(\theta_k) - F_{l.b.}\right] + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{L_{R+1}}{1 + \alpha}\|\theta_{k+1} - \theta_k\|_2^{1+\alpha}\right)\boldsymbol{1}\left[\mathcal{B}_{k+1}(R)\right]. \tag{49}
$$

556 Our goal now is to replace $\mathcal{B}_{k+1}(R)$ on the right hand side by $\mathcal{B}_k(R)$. However, there is a technical
557 difficulty which we must address. First, it follows from the preceding inequality that

$$
\left[F(\theta_{k+1}) - F_{l.b.}\right]\boldsymbol{1}\left[\mathcal{B}_{k+1}(R)\right]
$$
$$
\leq \left(\left[F(\theta_k) - F_{l.b.}\right] + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{L_{R+1}}{1 + \alpha}\|\theta_{k+1} - \theta_k\|_2^{1+\alpha}\right)
$$
$$
\times \left(\boldsymbol{1}\left[\mathcal{B}_{k+1}(R)\right] - \boldsymbol{1}\left[\mathcal{B}_k(R)\right]\right) \tag{50}
$$
$$
+ \left(\left[F(\theta_k) - F_{l.b.}\right] + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{L_{R+1}}{1 + \alpha}\|\theta_{k+1} - \theta_k\|_2^{1+\alpha}\right)\boldsymbol{1}\left[\mathcal{B}_k(R)\right].
$$

558 The first term on the right hand side of the inequality only contributes meaningfully if it is positive.
559 Since $\boldsymbol{1}\left[\mathcal{B}_k(R)\right] \geq \boldsymbol{1}\left[\mathcal{B}_{k+1}(R)\right]$, then two statements hold: (i) $\boldsymbol{1}\left[\mathcal{B}_k(R)\right]\boldsymbol{1}\left[\mathcal{B}_{k+1}(R)\right] = \boldsymbol{1}\left[\mathcal{B}_{k+1}(R)\right]$;
560 and (ii) the first term of the right hand side of (50) is positive if and only if

$$
\left(\left[F(\theta_k) - F_{l.b.}\right] + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{L_{R+1}}{1 + \alpha}\|\theta_{k+1} - \theta_k\|_2^{1+\alpha}\right)\boldsymbol{1}\left[\mathcal{B}_k(R)\right] < 0. \tag{51}
$$

561 By the choice of L_{R+1}, Assumption 1 and Lemma 1 imply that if (51) occurs, then $\|\theta_{k+1}\|_2 > R +$
562 $1 \geq \|\theta_k\|_2 + 1$. By the reverse triangle inequality and (4), if (51) occurs, then $\|M_k\dot{f}(\theta_k, X_{k+1})\|_2 \geq 1$.

Hence,

$$
\left([F(\theta_k) - F_{l.b.}] + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{L_{R+1}}{1+\alpha} \|\theta_{k+1} - \theta_k\|_2^{1+\alpha} \right)
$$
$$
\times \left(\mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] - \mathbf{1}\left[\mathcal{B}_k(R)\right] \right)
$$
$$
\leq \left(-[F(\theta_k) - F_{l.b.}] - \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) - \frac{L_{R+1}}{1+\alpha} \|\theta_{k+1} - \theta_k\|_2^{1+\alpha} \right) \tag{52}
$$
$$
\times \left(\mathbf{1}\left[\mathcal{B}_k(R)\right] - \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \right) \mathbf{1}\left[\mathcal{B}_k(R)\right] \mathbf{1}\left[\left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2 \geq 1\right].
$$

We now compute another coarse upper bound for this inequality. Note, by Assumption 1 and Cauchy-Schwarz,

$$
\left(-[F(\theta_k) - F_{l.b.}] - \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) - \frac{L_{R+1}}{1+\alpha} \|\theta_{k+1} - \theta_k\|_2^{1+\alpha} \right) \tag{53}
$$
$$
\times \left(\mathbf{1}\left[\mathcal{B}_k(R)\right] - \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \right) \mathbf{1}\left[\mathcal{B}_k(R)\right] \mathbf{1}\left[\left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2 \geq 1\right]
$$
$$
\leq \left\| \dot{F}(\theta_k) \right\|_2 \left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \mathbf{1}\left[\left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2 \geq 1\right] \tag{54}
$$
$$
\leq \left\| \dot{F}(\theta_k) \right\|_2 \left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2^{1+\alpha} \mathbf{1}\left[\mathcal{B}_k(R)\right] \tag{55}
$$
$$
\leq \frac{\partial F_R}{1+\alpha} \left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2^{1+\alpha} \mathbf{1}\left[\mathcal{B}_k(R)\right], \tag{56}
$$

where $\partial F_R = \sup_{\theta \in \overline{B(0,R)}} \|\dot{F}(\theta)\|_2 (1+\alpha) < \infty$ given that $\|\dot{F}(\theta)\|_2$ is a continuous function of θ.

Applying this inequality to (50), we conclude

$$
[F(\theta_{k+1}) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_{k+1}(R)\right]
$$
$$
\leq \left([F(\theta_k) - F_{l.b.}] - \dot{F}(\theta_k)' M_k \dot{f}(\theta_k, X_{k+1}) + \frac{L_{R+1} + \partial F_R}{1+\alpha} \left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2^{1+\alpha} \right) \tag{57}
$$
$$
\times \mathbf{1}\left[\mathcal{B}_k(R)\right].
$$

By Assumption 3,

$$
\mathbb{E}\left[[F(\theta_{k+1}) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \big| \mathcal{F}_k \right]
$$
$$
\leq \left([F(\theta_k) - F_{l.b.}] - \dot{F}(\theta_k)' M_k \dot{F}(\theta_k) + \frac{L_{R+1} + \partial F_R}{1+\alpha} \mathbb{E}\left[\left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2^{1+\alpha} \Big| \mathcal{F}_k \right] \right) \tag{58}
$$
$$
\times \mathbf{1}\left[\mathcal{B}_k(R)\right].
$$

Using Property 1 and Assumption 4,

$$
\mathbb{E}\left[[F(\theta_{k+1}) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \big| \mathcal{F}_k \right]
$$
$$
\leq \left([F(\theta_k) - F_{l.b.}] - \lambda_{\min}(M_k) \left\| \dot{F}(\theta_k) \right\|_2^2 + \frac{L_{R+1} + \partial F_R}{1+\alpha} \lambda_{\max}(M_k)^{1+\alpha} G(\theta_k) \right) \mathbf{1}\left[\mathcal{B}_k(R)\right]. \tag{59}
$$

By Assumption 4, G is upper semicontinuous and $\overline{B(0,R)}$ is compact, which implies that G_R is well defined and finite. The result follows. $\qquad\square$

C.2 Objective Function Analysis

Corollary 1. *Let $\{\theta_k\}$ be defined as in* (4) *satisfying Properties* 1 *and* 2. *Suppose Assumptions* 1 *to* 4 *hold. Then, there exists a finite random variable $F_{\lim}$ such that on the event $\{\sup_k \|\theta_k\|_2 < \infty\}$, $\lim_{k\to\infty} F(\theta_k) = F_{\lim}$ with probability one.*

Proof. By Lemma 2, for every $R \geq 0$,

$$
\begin{aligned}
\mathbb{E}\left[[F(\theta_{k+1}) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_{k+1}(R)\right]|\,\mathcal{F}_k\right] \\
\leq [F(\theta_k) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_k(R)\right] + \frac{(L_{R+1} + \partial F_R)G_R}{1 + \alpha}\lambda_{\max}(M_k)^{1+\alpha}.
\end{aligned}
\tag{60}
$$

By Neveu and Speed [1975, Exercise II.4] (cf. Robbins and Siegmund [1971]) and Property 2, $\lim_{k\to\infty}[F(\theta_k) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_k(R)\right]$ converges to a finite random variable with probability one. Since $R \geq 0$ is arbitrary, we conclude that there exists a finite random variable $F_{\lim}$ such that the set $\{\sup_k \|\theta_k\|_2 \leq R\}$ is a subset of $\{\lim_k F(\theta_k) = F_{\lim}\}$ up to a measure zero set. Since the countable union of measure zero sets has measure zero,

$$
\left\{\sup_k \|\theta_k\|_2 < \infty\right\} = \bigcup_{R\in\mathbb{N}}\left\{\sup_k \|\theta_k\|_2 \leq R\right\} \subset \left\{\lim_{k\to\infty} F(\theta_k) = F_{\lim}\right\},
\tag{61}
$$

up to a measure zero set. The result follows. $\qquad\square$

C.3 Gradient Function Analysis

We now prove that the gradient norm evaluated at SGD's iterates must, repeatedly, get arbitrarily close to zero.

Lemma 9. *Let $\{\theta_k\}$ be defined as in (4) satisfying Properties 1 to 3. Suppose Assumptions 1 to 4 hold. Then, $\forall R \geq 0$ and for all $\delta > 0$,*

$$
\mathbb{P}\left[\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta,\ i.o.\right] = 1.
\tag{62}
$$

Proof. By Lemma 2,

$$
\begin{aligned}
\lambda_{\min}(M_k)\mathbb{E}\left[\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right]\right] \leq \mathbb{E}\left[[F(\theta_k) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_k(R)\right]\right] \\
- \mathbb{E}\left[[F(\theta_{k+1}) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_{k+1}(R)\right]\right] + \frac{(L_{R+1} + \partial F_R)G_R}{1 + \alpha}\lambda_{\max}(M_k)^{1+\alpha}.
\end{aligned}
\tag{63}
$$

Taking the sum of this equation for all k from 0 to $j \in \mathbb{N}$, we have

$$
\begin{aligned}
\sum_{k=0}^{j}\lambda_{\min}(M_k)\mathbb{E}\left[\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right]\right] \leq [F(\theta_0) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_0(R)\right] \\
- \mathbb{E}\left[[F(\theta_{j+1}) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_{j+1}(R)\right]\right] + \frac{(L_{R+1} + \partial F_R)G_R}{1 + \alpha}\sum_{k=0}^{j}\lambda_{\max}(M_k)^{1+\alpha}.
\end{aligned}
\tag{64}
$$

By Assumption 1 and Property 2, the right hand side is bounded by

$$
[F(\theta_0) - F_{l.b.}]\mathbf{1}\left[\mathcal{B}_0(R)\right] + \frac{(L_{R+1} + \partial F_R)G_R}{1 + \alpha}S,
\tag{65}
$$

which is finite. Therefore, $\sum_{k=0}^{\infty}\lambda_{\min}(M_k)\mathbb{E}[\|\dot{F}(\theta_k)\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right]]$ is finite. Furthermore, by Property 3, $\liminf_k \mathbb{E}[\|\dot{F}(\theta_k)\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right]] = 0$.

Now, for any $\delta > 0$, Markov's inequality implies that for all $j + 1 \in \mathbb{N}$,

$$
\mathbb{P}\left[\bigcap_{k=j}^{\infty}\left\{\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right] > \delta\right\}\right] \leq \frac{1}{\delta}\min_{j\leq k}\mathbb{E}\left[\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right]\right],
\tag{66}
$$

where the right hand side is zero because $\liminf_k \mathbb{E}[\|\dot{F}(\theta_k)\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right]] = 0$.

As the countable union of measure zero sets has measure zero, we conclude that for all $\delta > 0$,

$$
\mathbb{P}\left[\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta,\ i.o.\right] = 1.
\tag{67}
$$

$\qquad\square$

Unfortunately, Lemma 9 does not guarantee that the gradient norm will be captured within a region of zero. In order to prove this, we first show that it is not possible (i.e., a zero probability event) for the limit supremum and limit infimum of the gradients to be distinct.

Lemma 10. *Let $\{\theta_k\}$ be defined as in* (4) *satisfying Properties 1 and 2. Suppose Assumptions 1 to 4 hold. Then, $\forall R \geq 0$ and for all $\delta > 0$,*

$$\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] > \delta, \left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta, \ i.o.\right] = 0. \tag{68}$$

Proof. Let $\gamma > 0$. Let L_R be as in Definition 1, and G_R be as in Lemma 2. Then, for $\delta > 0$,

$$\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta\right] > \delta + L_R\gamma^\alpha\right] \tag{69}$$

$$= \mathbb{P}\left[\left(\left\|\dot{F}(\theta_{k+1})\right\|_2 - \left\|\dot{F}(\theta_k)\right\|_2 + \left\|\dot{F}(\theta_k)\right\|_2\right) \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right]\right. \tag{70}$$

$$\left. \times \mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta\right] > \delta + L_R\gamma^\alpha\right]. \tag{71}$$

Using the reverse triangle inequality, $\|\dot{F}(\theta_{k+1})\|_2 - \|\dot{F}(\theta_k)\|_2 \leq \|\dot{F}(\theta_{k+1}) - \dot{F}(\theta_k)\|_2$. Now, making use of the restriction to $\mathcal{B}_{k+1}(R)$, $\|\dot{F}(\theta_{k+1}) - \dot{F}(\theta_k)\|_2 \leq L_R\|\theta_{k+1} - \theta_k\|_2^\alpha$. Moreover, on $\|\dot{F}(\theta_k)\|_2 \leq \delta$, $\|\dot{F}(\theta_k)\|_2\mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \leq \delta$. Putting these two observations together,

$$\mathbb{P}\left[\left(\left\|\dot{F}(\theta_{k+1})\right\|_2 - \left\|\dot{F}(\theta_k)\right\|_2 + \left\|\dot{F}(\theta_k)\right\|_2\right) \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right]\right. \tag{72}$$

$$\left. \times \mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta\right] > \delta + L_R\gamma^\alpha\right] \tag{73}$$

$$\leq \mathbb{P}\left[L_R \left\|\theta_{k+1} - \theta_k\right\|_2^\alpha \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta\right] > L_R\gamma^\alpha\right] \tag{74}$$

$$= \mathbb{P}\left[\left\|M_k \dot{f}(\theta_k, X_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta\right] > \gamma\right]. \tag{75}$$

Now, using $\mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right]\right] \leq \mathbf{1}\left[\mathcal{B}_k(R)\right]$ and Markov's inequality,

$$\mathbb{P}\left[\left\|M_k \dot{f}(\theta_k, X_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta\right] > \gamma\right] \tag{76}$$

$$\leq \mathbb{P}\left[\left\|M_k \dot{f}(\theta_k, X_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] > \gamma\right] \tag{77}$$

$$\leq \frac{1}{\gamma^{1+\alpha}} \|M_k\|_2^{1+\alpha} \mathbb{E}\left[\mathbb{E}\left[\left\|\dot{f}(\theta_k, X_{k+1})\right\|_2^{1+\alpha}\middle| \mathcal{F}_k\right] \mathbf{1}\left[\mathcal{B}_k(R)\right]\right]. \tag{78}$$

By Assumption 4, $\mathbb{E}\left[\|\dot{f}(\theta_k, X_{k+1})\|_2^{1+\alpha}\middle| \mathcal{F}_k\right] \leq G(\theta_k)$. Moreover, on $\mathcal{B}_k(R)$, $G(\theta_k) \leq G_R$. Using this in the expectation, we conclude

$$\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] \mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta\right] > \delta + L_R\gamma^\alpha\right] \leq \frac{1}{\gamma^{1+\alpha}} \|M_k\|_2^{1+\alpha} G_R. \tag{79}$$

By Property 2, the sum of the last expression over all $k + 1 \in \mathbb{N}$ is finite. By the Borel-Cantelli lemma, for all $R \geq 0$, $\delta > 0$ and $\gamma > 0$,

$$\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] > \delta + L_R\gamma^\alpha, \left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta, \ i.o.\right] = 0. \tag{80}$$

Since this holds for any $\gamma > 0$, it will hold for every value in a sequence $\gamma_n \downarrow 0$. Since the countable union of measure zero events has measure zero,

$$\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] > \delta, \left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta, \ i.o.\right] = 0. \tag{81}$$

$\square$

We now put together Lemmas 9 and 10 to show that, on the event $\{\sup_k \|\theta_k\|_2 < \infty\}$, $\|\dot{F}(\theta_k)\|_2$ converges to 0 with probability one.

Corollary 2. *Let $\{\theta_k\}$ be defined as in (4) satisfying Properties 1 to 3. Suppose Assumptions 1 to 4 hold. Then, on the event $\{\sup_k \|\theta_k\|_2 < \infty\}$, $\lim_{k\to\infty} \|\dot{F}(\theta_k)\|_2 = 0$.*

Proof. For any $R \geq 0$ and $\delta > 0$, Lemma 9 implies

$$
\begin{aligned}
&\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] > \delta,\ i.o.\right] \\
&= \mathbb{P}\left[\left\{\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] > \delta,\ i.o.\right\} \cap \left\{\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta,\ i.o.\right\}\right].
\end{aligned}
\tag{82}
$$

We see that this latter event is exactly,

$$
\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] > \delta,\ \left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\mathcal{B}_k(R)\right] \leq \delta,\ i.o.\right],
\tag{83}
$$

which, by Lemma 10, is zero with probability one. Therefore, $\mathbb{P}[\|\dot{F}(\theta_{k+1})\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] > \delta,\ i.o.]$ is zero. Letting $\delta_n \downarrow 0$ and noting that the countable union of measure zero sets has measure zero, we conclude $\mathbb{P}[\|\dot{F}(\theta_{k+1})\|_2 \mathbf{1}\left[\mathcal{B}_{k+1}(R)\right] > 0,\ i.o.] = 0$.

Therefore, for all $R \geq 0$, $\{\sup_k \|\theta_k\|_2 \leq R\} \subset \{\lim_{k\to\infty} \|\dot{F}(\theta_k)\|_2 = 0\}$ up to a measure zero set. Since $\{\sup_k \|\theta_k\|_2 < \infty\} = \cup_{R\in\mathbb{N}}\{\sup_k \|\theta_k\|_2 \leq R\}$, the result follows. $\qquad\square$

C.4   Capture Theorem

The final step in our proof is to study the event $\{\sup_k \|\theta_k\| < \infty\}$.

Theorem 4 (Theorem 1). *Let $\{\theta_k\}$ be defined as in (4), and let $\{M_k\}$ satisfy Properties 1 and 2. If Assumption 4 holds, then either $\{\lim_{k\to\infty} \theta_k \text{ exists}\}$ or $\{\liminf_{k\to\infty} \|\theta_k\|_2 = \infty\}$ must occur.*

Proof. Let $\bar{\theta} \in \mathbb{R}^p$. Fix $R \geq 0$ and let $\gamma > 0$. Then,

$$
\begin{aligned}
&\mathbb{P}\left[\left\|\theta_{k+1} - \bar{\theta}\right\|_2 \geq R + \gamma,\ \left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] \\
&= \mathbb{P}\left[\left\|\theta_{k+1} - \bar{\theta}\right\|_2 \mathbf{1}\left[\left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] \geq R + \gamma\right] \\
&= \mathbb{P}\left[\left(\left\|\theta_{k+1} - \bar{\theta}\right\|_2 - \left\|\theta_k - \bar{\theta}\right\|_2 + \left\|\theta_k - \bar{\theta}\right\|_2\right) \mathbf{1}\left[\left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] \geq R + \gamma\right].
\end{aligned}
\tag{84, 85}
$$

Now, $\|\theta_k - \bar{\theta}\|_2 \mathbf{1}\left[\|\theta_k - \bar{\theta}\|_2 \leq R\right] \leq R$. Therefore,

$$
\begin{aligned}
&\mathbb{P}\left[\left(\left\|\theta_{k+1} - \bar{\theta}\right\|_2 - \left\|\theta_k - \bar{\theta}\right\|_2 + \left\|\theta_k - \bar{\theta}\right\|_2\right) \mathbf{1}\left[\left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] \geq R + \gamma\right] \\
&\leq \mathbb{P}\left[\left(\left\|\theta_{k+1} - \bar{\theta}\right\|_2 - \left\|\theta_k - \bar{\theta}\right\|_2\right) \mathbf{1}\left[\left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] + R \geq R + \gamma\right] \\
&\leq \mathbb{P}\left[\left\|\theta_{k+1} - \theta_k\right\|_2 \mathbf{1}\left[\left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] \geq \gamma\right],
\end{aligned}
\tag{86, 87, 88}
$$

where the last line follows by applying the reverse triangle inequality. By using (4) and Markov's inequality,

$$
\begin{aligned}
&\mathbb{P}\left[\left\|\theta_{k+1} - \theta_k\right\|_2 \mathbf{1}\left[\left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] \geq \gamma\right] \\
&\leq \mathbb{P}\left[\left\|M_k \dot{f}(\theta_k, X_{k+1})\right\|_2 \mathbf{1}\left[\left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] \geq \gamma\right] \\
&\leq \frac{1}{\gamma^{1+\alpha}} \left\|M_k\right\|_2^{1+\alpha} \mathbb{E}\left[\mathbb{E}\left[\left\|\dot{f}(\theta_k, X_{k+1})\right\|_2^{1+\alpha}\ \middle|\ \mathcal{F}_k\right] \mathbf{1}\left[\left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right]\right].
\end{aligned}
\tag{89, 90, 91}
$$

By applying Assumption 4, $\mathbb{E}\left[\|\dot{f}(\theta_k, X_{k+1})\|_2^{1+\alpha}\ \middle|\ \mathcal{F}_k\right] \leq G(\theta_k)$. Moreover, on $\|\theta_k - \bar{\theta}\|_2 \leq R$, $G(\theta_k) \leq \sup_{\theta:\|\theta\|_2 \leq R+\|\bar{\theta}\|_2} G(\theta) =: G_{R+\|\bar{\theta}\|_2} < \infty$ since G is upper semi-continuous. Combining these steps,

$$
\mathbb{P}\left[\left\|\theta_{k+1} - \bar{\theta}\right\|_2 \geq R + \gamma,\ \left\|\theta_k - \bar{\theta}\right\|_2 \leq R\right] \leq \frac{1}{\gamma^{1+\alpha}} \left\|M_k\right\|_2^{1+\alpha} G_{R+\|\bar{\theta}\|_2},
\tag{92}
$$

By Property 2, we see that the sum of the probabilities is finite. Together with the Borel-Cantelli lemma, $\mathbb{P}[\|\theta_{k+1} - \bar{\theta}\|_2 \geq R + \gamma,\ \|\theta_k - \bar{\theta}\|_2 \leq R,\ i.o.] = 0$. Since $\gamma > 0$ is arbitrary, we can

show that this statement holds for a countable sequence of $\gamma_n \downarrow 0$. Therefore, $\mathbb{P}[\|\theta_{k+1} - \bar{\theta}\|_2 > R, \ \|\theta_k - \bar{\theta}\|_2 \leq R, \ i.o.] = 0$.

Since R is arbitrary as well, we conclude that for each element in a finite sequence $R_n \uparrow \infty$, $\mathbb{P}[\|\theta_{k+1} - \bar{\theta}\|_2 > R_n, \ \|\theta_k - \bar{\theta}\|_2 \leq R_n, \ i.o.] = 0$. Again, the countable union of measure zero sets is measure zero. Hence, we conclude that $\mathbb{P}[\limsup_k \|\theta_k - \bar{\theta}\|_2 > \liminf_k \|\theta_k - \bar{\theta}\|_2] = 0$. Thus, either $\lim_k \|\theta_k - \bar{\theta}\|_2$ exists and is either infinite or finite.

Moreover, on the event that the limit is finite, since $\bar{\theta}$ is arbitrary, we can choose three distinct values of $\bar{\theta}$ which are not colinear, and, by triangulation, the $\lim_k \theta_k$ converges to a fixed point (up to a set of measure zero). $\qquad\square$

D Stability Analysis

We begin with a recursive relationship on the events $\{\tau_j > k\}$. We use this result to prove that the objective function converges to a finite limit on these events. Then, we use this result to conclude that the gradient function converges to zero on $\{\nu_j > k\}$. Finally, we study the combination of these events to establish that the two statements above hold with probability one.

D.1 A Recursive Relationship

Lemma 11 (Lemma 3)**.** *Let* $\{M_k\}$ *satisfy* Property 1*. Suppose* Assumptions 1 *to* 4 *hold. Let* $\{\theta_k\}$ *satisfy* (4)*. Then, for any* $j + 1 \in \mathbb{N}$ *and* $k > j$*,*

$$\mathbb{E}\left[(F(\theta_{k+1}) - F_{l.b.}) \, \mathbf{1}\left[\tau_j > k\right] \middle| \mathcal{F}_k \right] \leq \left(F(\theta_k) - F_{l.b.} - \dot{F}(\theta_k)' M_k \dot{F}(\theta_k) \right) \mathbf{1}\left[\tau_j > k-1\right]$$

$$+ \frac{\lambda_{\max}(M_k)^{1+\alpha}}{1+\alpha} \left[\mathcal{L}_\epsilon(\theta_k) G(\theta_k) + \alpha \left[\frac{\left\| \dot{F}(\theta_k) \right\|_2^{1+\alpha}}{\mathcal{L}_\epsilon(\theta_k)} \right]^{1/\alpha} \right] \mathbf{1}\left[\tau_j > k-1\right]. \tag{93}$$

Proof. By the construction of τ_j, when $\tau_j > k$, then

$$F(\theta_{k+1}) - F_{l.b.} \leq F(\theta_k) - F_{l.b.} + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{\mathcal{L}_\epsilon(\theta_k)}{1+\alpha} \|\theta_{k+1} - \theta_k\|_2^{1+\alpha}. \tag{94}$$

Using this relationship and using $\mathbf{1}\left[\tau_j > k\right] = \mathbf{1}\left[\tau_j > k-1\right] - \mathbf{1}\left[\tau_j = k\right]$,

$$\mathbb{E}\left[\{ F(\theta_{k+1}) - F_{l.b.} \} \mathbf{1}\left[\tau_j > k\right] \middle| \mathcal{F}_k \right]$$

$$\leq \mathbb{E}\left[\left\{ F(\theta_k) - F_{l.b.} + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{\mathcal{L}_\epsilon(\theta_k)}{1+\alpha} \|\theta_{k+1} - \theta_k\|_2^{1+\alpha} \right\} \mathbf{1}\left[\tau_j > k-1\right] \middle| \mathcal{F}_k \right]$$

$$- \mathbb{E}\left[\left\{ F(\theta_k) - F_{l.b.} + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{\mathcal{L}_\epsilon(\theta_k)}{1+\alpha} \|\theta_{k+1} - \theta_k\|_2^{1+\alpha} \right\} \mathbf{1}\left[\tau_j = k\right] \middle| \mathcal{F}_k \right] \tag{95}$$

For the first term on the right hand side, we can apply Assumptions 3 and 4, Property 1, and (4) to calculate

$$\mathbb{E}\left[\left\{ F(\theta_k) - F_{l.b.} + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{\mathcal{L}_\epsilon(\theta_k)}{1+\alpha} \|\theta_{k+1} - \theta_k\|_2^{1+\alpha} \right\} \mathbf{1}\left[\tau_j > k-1\right] \middle| \mathcal{F}_k \right]$$

$$\leq \left\{ F(\theta_k) - F_{l.b.} - \dot{F}(\theta_k)' M_k \dot{F}(\theta_k) + \frac{\lambda_{\max}(M_k)^{1+\alpha}}{1+\alpha} \mathcal{L}_\epsilon(\theta_k) G(\theta_k) \right\} \mathbf{1}\left[\tau_j > k-1\right]. \tag{96}$$

For the second term on the right hand side of (95), we first note $\mathbf{1}\left[\tau_j = k\right] \leq \mathbf{1}\left[\tau_j > k-1\right]$ which implies $\mathbf{1}\left[\tau_j = k\right] = \mathbf{1}\left[\tau_j = k\right] \mathbf{1}\left[\tau_j > k-1\right]$. Second, the Cauchy-Schwarz inequality and Lemma 7

imply

$$- F(\theta_k)' M_k \dot{f}(\theta_k, X_{k+1}) + \frac{\mathcal{L}_\epsilon(\theta_k)}{1+\alpha} \| M_k f(\theta_k, X_{k+1}) \|_2^{1+\alpha}$$

$$\geq - \| F(\theta_k) \|_2 \left\| M_k \dot{f}(\theta_k, X_{k+1}) \right\|_2 + \frac{\mathcal{L}_\epsilon(\theta_k)}{1+\alpha} \| M_k f(\theta_k, X_{k+1}) \|_2^{1+\alpha} \qquad (97)$$

$$\geq - \frac{\alpha}{1+\alpha} \left[\frac{\left\| \dot{F}(\theta_k) \right\|_2^{1+\alpha}}{\mathcal{L}_\epsilon(\theta_k)} \right]^{1/\alpha} \qquad (98)$$

Hence, using (4),

$$- [F(\theta_k) - F_{l.b.}] - \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) - \frac{\mathcal{L}_\epsilon(\theta_k)}{1+\alpha} \| \theta_{k+1} - \theta_k \|_2^{1+\alpha}$$

$$\leq - [F(\theta_k) - F_{l.b.}] + \frac{\alpha}{1+\alpha} \left[\frac{\left\| \dot{F}(\theta_k) \right\|_2^{1+\alpha}}{\mathcal{L}_\epsilon(\theta_k)} \right]^{1/\alpha} \qquad (99)$$

With the help of Theorem 5, we conclude

$$- \mathbb{E} \left[\left\{ F(\theta_k) - F_{l.b.} + \dot{F}(\theta_k)'(\theta_{k+1} - \theta_k) + \frac{\mathcal{L}_\epsilon(\theta_k)}{1+\alpha} \| \theta_{k+1} - \theta_k \|_2^{1+\alpha} \right\} \mathbf{1}\left[\tau_j = k \right] \bigg| \mathcal{F}_k \right]$$

$$\leq \left\{ - [F(\theta_k) - F_{l.b.}] + \frac{\alpha}{1+\alpha} \left[\frac{\left\| \dot{F}(\theta_k) \right\|_2^{1+\alpha}}{\mathcal{L}_\epsilon(\theta_k)} \right]^{1/\alpha} \right\} \mathbb{P}\left[\tau_j = k | \mathcal{F}_k \right] \mathbf{1}\left[\tau_j > k - 1 \right] \qquad (100)$$

$$\leq \frac{\alpha \lambda_{\max}(M_k)^{1+\alpha}}{1+\alpha} \left[\frac{\left\| \dot{F}(\theta_k) \right\|_2^{1+\alpha}}{\mathcal{L}_\epsilon(\theta_k)} \right]^{1/\alpha} \mathbf{1}\left[\tau_j > k - 1 \right]. \qquad (101)$$

By putting the bounds on the first and second term together, the result follows. □

By applying Assumption 5 to Lemma 3, we have the following simplified form.

Lemma 12 (Lemma 4). *If Assumptions 1 to 5, and Properties 1 and 4 hold, and $\{\theta_k\}$ satisfy (4), then there exists a $K \in \mathbb{N}$ such that for any $j + 1 \in \mathbb{N}$ and any $k \geq \min\{K, j + 1\}$,*

$$\mathbb{E}\left[(F(\theta_{k+1}) - F_{l.b.}) \boldsymbol{I}\left[\tau_j > k \right] | \mathcal{F}_k \right]$$

$$\leq \left(1 + \lambda_{\max}(M_k)^{1+\alpha} \frac{C_2}{1+\alpha} \right) (F(\theta_k) - F_{l.b.}) \boldsymbol{I}\left[\tau_j > k - 1 \right] \qquad (102)$$

$$- \frac{1}{2} \lambda_{\min}(M_k) \left\| \dot{F}(\theta_k) \right\|_2^2 \boldsymbol{I}\left[\tau_j > k - 1 \right] + \lambda_{\max}(M_k)^{1+\alpha} \frac{C_1}{1+\alpha}.$$

Proof. The result follows by first using Assumption 5 in Lemma 3. Then, collecting similar terms, we apply Lemma 6 to find K. □

D.2 Objective Function Analysis

With this recursive formula, we now have the first result.

Corollary 3. *If Assumptions 1 to 5 and Properties 1, 2 and 4 hold, and $\{\theta_k\}$ satisfy (4), then $\lim_{k\to\infty} F(\theta_k)$ exists and is finite on $\cup_{j=0}^\infty \{\tau_j = \infty\}$.*

Proof. By Lemma 4 and Robbins and Siegmund [1971], Neveu and Speed [1975, Exercise II.4], the limit as k goes to infinity of $(F(\theta_k) - F_{l.b.})\mathbf{1}\left[\tau_j > k \right]$ exists with probability one and is integrable. Therefore, on the event $\{\tau_j = \infty\}$, the limit of $F(\theta_k) - F_{l.b.}$ exists and is integrable. As a result, the limit of $F(\theta_k) - F_{l.b.}$ exists and is finite on $\cup_{j=0}^\infty \{\tau_j = \infty\}$. □

678 Additionally, we can state the following useful result.

679 **Lemma 13.** *If Assumptions 1 to 5, and Properties 1, 2 and 4 hold, and $\{\theta_k\}$ satisfy (4), then $\exists K \in \mathbb{N}$*
680 *such that for any $j > K$, $\exists N_j > 0$ for which*

$$\sup_{k > j} \mathbb{E}\left[(F(\theta_k) - F_{l.b.})\mathbf{1}\left[\tau_j > k\right]\right] \leq N_j. \tag{103}$$

681 *Proof.* By Lemma 4, Property 2, and any $k \geq j$,

$$\mathbb{E}\left[(F(\theta_{k+1}) - F_{l.b.})\mathbf{1}\left[\tau_j > k\right]\right] + \frac{C_1}{1+\alpha}\sum_{\ell=k+1}^{\infty}\lambda_{\max}(M_\ell)^{1+\alpha}$$

$$\leq \exp\left(\frac{C_2}{1+\alpha}\lambda_{\max}(M_k)^{1+\alpha}\right)\left[\mathbb{E}\left[(F(\theta_k) - F_{l.b.})\mathbf{1}\left[\tau_j > k-1\right]\right] + \frac{C_1}{1+\alpha}\sum_{\ell=k}^{\infty}\lambda_{\max}(M_\ell)^{1+\alpha}\right]. \tag{104}$$

682 Hence,

$$\mathbb{E}\left[(F(\theta_{k+1}) - F_{l.b.})\mathbf{1}\left[\tau_j > k\right]\right] + \frac{C_1}{1+\alpha}\sum_{\ell=k+1}^{\infty}\lambda_{\max}(M_\ell)^{1+\alpha}$$

$$\leq \exp\left(\frac{C_2}{1+\alpha}\sum_{\ell=j}^{k}\lambda_{\max}(M_\ell)^{1+\alpha}\right)\left[\mathbb{E}\left[(F(\theta_j) - F_{l.b.})\right] + \frac{C_1}{1+\alpha}\sum_{\ell=j}^{\infty}\lambda_{\max}(M_\ell)^{1+\alpha}\right], \tag{105}$$

683 where we have used $\mathbf{1}\left[\tau_j > j-1\right] = 1$. By Property 2, the summation in the exponent is finite,
684 which implies the result. □

D.3 Gradient Function Analysis

686 **Lemma 14.** *If Assumptions 1 to 5, and Properties 1 to 4 hold, and $\{\theta_k\}$ satisfy (4), then, for any*
687 $\delta > 0$,

$$\mathbb{P}\left[\left\|\dot{F}(\theta_k)\right\|_2\mathbf{1}\left[\tau_j > k-1\right] \leq \delta \text{ i.o.}\right] = 1. \tag{106}$$

688 *Proof.* By Lemma 4,

$$\frac{1}{2}\lambda_{\min}(M_k)\mathbb{E}\left[\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\tau_j > k-1\right]\right] \leq \mathbb{E}\left[(F(\theta_k) - F_{l.b.})\mathbf{1}\left[\tau_j > k-1\right]\right]$$

$$- \mathbb{E}\left[(F(\theta_{k+1}) - F_{l.b.})\mathbf{1}\left[\tau_j > k\right]\right] + \frac{C_2}{1+\alpha}\lambda_{\max}(M_k)^{1+\alpha}\mathbb{E}\left[(F(\theta_k) - F_{l.b.})\mathbf{1}\left[\tau_j > k-1\right]\right]$$

$$+ \frac{C_1}{1+\alpha}\lambda_{\max}(M_k)^{1+\alpha}. \tag{107}$$

689 By applying Lemma 13,

$$\frac{1}{2}\lambda_{\min}(M_k)\mathbb{E}\left[\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\tau_j > k-1\right]\right] \leq \mathbb{E}\left[(F(\theta_k) - F_{l.b.})\mathbf{1}\left[\tau_j > k-1\right]\right]$$

$$- \mathbb{E}\left[(F(\theta_{k+1}) - F_{l.b.})\mathbf{1}\left[\tau_j > k\right]\right] + \lambda_{\max}(M_k)^{1+\alpha}\left(\frac{C_2 N_j + C_1}{1+\alpha}\right). \tag{108}$$

690 By summing and using Assumption 1,

$$\frac{1}{2}\sum_{k=j}^{\infty}\lambda_{\min}(M_k)\mathbb{E}\left[\left\|\dot{F}(\theta_k)\right\|_2^2\mathbf{1}\left[\tau_j > k\right]\right]$$

$$\leq \mathbb{E}\left[F(\theta_j) - F_{l.b.}\right] + \frac{C_2 N_j + C_1}{1+\alpha}\sum_{k=j}^{\infty}\lambda_{\max}(M_k)^{1+\alpha}. \tag{109}$$

By Property 2, the right hand side is bounded. Now, by Property 3,

$$\liminf_{k\to\infty} \mathbb{E}\left[\left\|\dot{F}(\theta_k)\right\|_2^2 \mathbf{1}\left[\tau_j > k\right]\right] = 0. \tag{110}$$

Using Markov's inequality, for any $\ell \in \mathbb{N}$ and any $\delta > 0$,

$$\mathbb{P}\left[\bigcap_{k=\ell}^\infty \left\{\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\tau_j > k\right] > \delta\right\}\right] \leq \frac{1}{\delta^2}\min_{k\geq\ell}\mathbb{E}\left[\left\|\dot{F}(\theta_k)\right\|_2^2 \mathbf{1}\left[\tau_j > k\right]\right] = 0. \tag{111}$$

As the countable union of sets of measure zero have measure zero, the result follows. $\square$

Corollary 4. *If Assumptions 1 to 5, and Properties 1 to 4 hold, and $\{\theta_k\}$ satisfy (4), then, on $\cup_{j=0}^\infty\{\nu_j = \infty\}$, $\lim_{k\to\infty}\|\dot{F}(\theta_k)\| = 0$.*

Proof. Let $\delta > 0$ and $\gamma > 0$. Note, $\mathbf{1}\left[\nu_j > k\right] \leq \mathbf{1}\left[\nu_j > k - 1\right]$. Then,

$$\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\nu_j > k\right] > \delta + \mathcal{L}_\epsilon(\theta_k)^{\frac{1}{1+\alpha}}\gamma^{\frac{\alpha}{1+\alpha}}, \left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\nu_j > k - 1\right] \leq \delta\,\Big|\,\mathcal{F}_k\right] \tag{112}$$

$$\leq \mathbb{P}\left[\left(\left\|\dot{F}(\theta_{k+1})\right\|_2 - \left\|\dot{F}(\theta_k)\right\|_2 + \left\|\dot{F}(\theta_k)\right\|_2\right)\mathbf{1}\left[\nu_j > k\right] > \delta + \mathcal{L}_\epsilon(\theta_k)^{\frac{1}{1+\alpha}}\gamma^{\frac{\alpha}{1+\alpha}},\right.$$

$$\left.\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\nu_j > k - 1\right] \leq \delta\,\Big|\,\mathcal{F}_k\right] \tag{113}$$

$$\leq \mathbb{P}\left[\mathcal{L}_\epsilon(\theta_k)\|\theta_{k+1} - \theta_k\|_2^\alpha \mathbf{1}\left[\nu_j > k\right] > \mathcal{L}_\epsilon(\theta_k)^{\frac{1}{1+\alpha}}\gamma^{\frac{\alpha}{1+\alpha}},\right. \tag{114}$$

$$\left.\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\nu_j > k - 1\right] \leq \delta\,\Big|\,\mathcal{F}_k\right] \tag{115}$$

$$\leq \mathbb{P}\left[\mathcal{L}_\epsilon(\theta_k)^{\frac{\alpha}{1+\alpha}}\|\theta_{k+1} - \theta_k\|_2^\alpha \mathbf{1}\left[\nu_j > k\right] > \gamma^{\frac{\alpha}{1+\alpha}}, \left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\nu_j > k - 1\right] \leq \delta\,\Big|\,\mathcal{F}_k\right] \tag{116}$$

$$\leq \frac{1}{\gamma}\lambda_{\max}(M_k)^{1+\alpha}\mathcal{L}_\epsilon(\theta_k)G(\theta_k)\mathbf{1}\left[\nu_j > k - 1\right]\mathbf{1}\left[\left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\nu_j > k - 1\right] \leq \delta\right] \tag{117}$$

$$\leq \frac{1}{\gamma}\lambda_{\max}(M_k)^{1+\alpha}\left(C_1 + C_2(F(\theta_k) - F_{l.b.})\mathbf{1}\left[\nu_j > k - 1\right] + C_3\delta^2\right), \tag{118}$$

where we have made use of Assumption 4, Markov's inequality, and Assumption 5 in the last two lines. Taking expectations, using $\sup_{k>j}\mathbb{E}\left[(F(\theta_k) - F_{l.b.})\mathbf{1}\left[\nu_j > k\right]\right] \leq N_j$ (Lemma 13), and applying Borel-Cantelli with Property 2, we conclude

$$\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\nu_j > k\right] > \delta + \mathcal{L}_\epsilon(\theta_k)^{\frac{1}{1+\alpha}}\gamma^{\frac{\alpha}{1+\alpha}}, \left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\nu_j > k - 1\right] \leq \delta\ i.o.\right] = 0. \tag{119}$$

Since γ is arbitrary, the result holds for a collection of $\gamma_n \downarrow 0$. As the countable union of measure zero events has measure zero,

$$\mathbb{P}\left[\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\nu_j > k\right] > \delta, \left\|\dot{F}(\theta_k)\right\|_2 \mathbf{1}\left[\nu_j > k - 1\right] \leq \delta\ i.o.\right] = 0. \tag{120}$$

Using this result in conjunction with Lemma 14, $\mathbb{P}\left[\lim_{k\to\infty}\left\|\dot{F}(\theta_{k+1})\right\|_2 \mathbf{1}\left[\nu_j > k\right] = 0\right] = 1$.

Hence, on the event $\{\nu_j = \infty\}$, $\lim_{k\to\infty}\|\dot{F}(\theta_k)\| = 0$. Thus, on $\cup_{j=0}^\infty\{\nu_j = \infty\}$, we conclude that $\lim_{k\to\infty}\|\dot{F}(\theta_k)\| = 0$. $\square$

D.4 Stopping Time Analysis

By Lemma 1, $\{\tau_j = k\} \subset \{\nu_j = k\}$. Therefore, $\{\tau_j \leq k\} \subset \{\nu_j \leq k\}$, and $\{\tau_j > k\} \supset \{\nu_j > k\}$. Using these relationships, we can compute the probabilities of $\{\tau_j = k\}$ and $\{\nu_j = k\}$.

Theorem 5. *Let $\{\tau_j : j + 1 \in \mathbb{N}\}$ be defined as in (10), and $\{\nu_j : j + 1 \in \mathbb{N}\}$ be defined as in (11). If Assumptions 1, 2 and 4 and Property 1 hold, and $\{\theta_k\}$ satisfy (4), then, for any $j + 1 \in \mathbb{N}$ and any $k + 1 \in \mathbb{N}$,*

$$\mathbb{P}\left[\tau_j = k\,|\,\mathcal{F}_k\right] \leq \mathbb{P}\left[\nu_j = k\,|\,\mathcal{F}_k\right] \leq \begin{cases} 0 & k \leq j, \\ \lambda_{\max}(M_k)^{1+\alpha} & k > j. \end{cases} \tag{121}$$

Moreover, if Property 2 also holds, then $\mathbb{P}\left[\cup_{j=0}^\infty\{\tau_j = \infty\}\right] = \mathbb{P}\left[\cup_{j=0}^\infty\{\nu_j = \infty\}\right] = 1$.

Proof. The case of $k \leq j$ is trivial. So consider only $k > j$. By the construction of $L(\cdot, \cdot)$ and $\mathcal{L}_\epsilon(\cdot)$, $\omega \in \{L(\theta_k, \theta_{k+1}) > \mathcal{L}_\epsilon(\theta_k)\}$ implies $\omega \in \{\|\theta_{k+1} - \theta_k\|_2 > (G(\theta_k) \vee \epsilon)^{\frac{1}{1+\alpha}}\}$. Using (4), Markov's inequality, Property 1, we conclude

$$\mathbb{P}\left[\tau_j = k \mid \mathcal{F}_k\right] \leq \mathbb{P}\left[\nu_j = k \mid \mathcal{F}_k\right] \leq \mathbb{P}\left[\left\|M_k \dot{f}(\theta_k, X_{k+1})\right\|_2^{1+\alpha} > G(\theta_k) \vee \epsilon \,\middle|\, \mathcal{F}_k\right] \tag{122}$$

$$\leq \frac{\lambda_{\max}(M_k)^{1+\alpha} \mathbb{E}\left[\left\|\dot{f}(\theta_k, X_{k+1})\right\|_2^{1+\alpha} \,\middle|\, \mathcal{F}_k\right]}{G(\theta_k) \vee \epsilon}. \tag{123}$$

Applying Assumption 4 supplies the bound on $\mathbb{P}\left[\tau_j = k \mid \mathcal{F}_k\right]$ and $\mathbb{P}\left[\nu_j = k \mid \mathcal{F}_k\right]$. For the second part, note

$$\mathbb{P}\left[\tau_j = \infty\right] \geq \mathbb{P}\left[\nu_j = \infty\right] \geq 1 - \mathbb{P}\left[\nu_j < \infty\right] \geq 1 - \sum_{k=j+1}^{\infty} \lambda_{\max}(M_k)^{1+\alpha}. \tag{124}$$

Moreover, $\{\nu_j = \infty\} \subset \{\nu_{j+1} = \infty\}$. Therefore,

$$\mathbb{P}\left[\bigcup_{j=0}^{\infty} \{\tau_j = \infty\}\right] \geq \mathbb{P}\left[\bigcup_{j=0}^{\infty} \{\nu_j = \infty\}\right] = \lim_{j \to \infty} \mathbb{P}\left[\nu_j = \infty\right]. \tag{125}$$

Since $\lim_j \mathbb{P}[\nu_j = \infty] \geq 1 - \lim_j \sum_{k=j+1}^{\infty} \lambda_{\max}(M_k)^{1+\alpha}$, applying Property 2 supplies the final result. $\qquad\square$