Online Frank-Wolfe with Arbitrary Delays

Abstract

The online Frank-Wolfe (OFW) method has gained much popularity for online convex optimization due to its projection-free property. Previous studies showed that OFW can attain an $O(T^{3/4})$ regret bound for convex losses and an $O(T^{2/3})$ regret bound for strongly convex losses. However, they assumed that each gradient queried by OFW is revealed immediately, which may not hold in practice and limits the application of OFW. To address this limitation, we proposed a delayed variant of OFW, which allows gradients to be delayed by arbitrary rounds. The main idea is to perform an update similar to OFW after receiving any delayed gradient, and play the latest decision for each round. Despite its simplicity, we prove that our delayed variant of OFW is able to achieve an $O(T^{3/4} + dT^{1/4})$ regret bound for convex losses and an $O(T^{2/3} + d \log T)$ regret bound for strongly convex losses, where d is the maximum delay. This is somewhat surprising since under a relatively large amount of delay (e.g., $d = O(\sqrt{T})$ for convex losses and $d = O(T^{2/3}/\log T)$ for strongly convex losses), the delayed variant of OFW enjoys the same regret bound like that of the original OFW.

1 Introduction

Online convex optimization (OCO) has become a leading paradigm for online learning due to its capability to model various problems from diverse domains such as online routing, online collaborative filtering, and online advertisement [Hazan, 2016]. In general, it is formulated as a structured repeated game between a player and an adversary. In each round t, the player first chooses a decision $\mathbf{x}_t$ from a convex decision set $\mathcal{K} \subseteq \mathbb{R}^n$, where n is the dimensionality. Then, the adversary selects a convex function $f_t(\mathbf{x}) : \mathcal{K} \mapsto \mathbb{R}$, and the player suffers a loss $f_t(\mathbf{x}_t)$. The player aims to choose decisions such that the regret

$$R(T) = \sum_{t=1}^{T} f_t(\mathbf{x}_t) - \min_{\mathbf{x} \in \mathcal{K}} \sum_{t=1}^{T} f_t(\mathbf{x})$$

is sublinear in the number of total rounds T. Online gradient descent (OGD) is a standard method for OCO, which enjoys an $O(\sqrt{T})$ regret bound for convex losses [Zinkevich, 2003] and an $O(\log T)$ regret bound for strongly convex losses [Hazan et al., 2007]. However, it needs to compute a projection onto the decision set to ensure the feasibility of each decision, which is computationally expensive for complex decision sets [Hazan and Kale, 2012].

To tackle this computational issue, Hazan and Kale [2012] proposed the online Frank-Wolfe (OFW) method, which has become one of the most commonly used algorithms for OCO over complex decision sets. The main advantage of OFW is its projection-free property: instead of performing the projection operation, it utilizes a linear optimization step to select a feasible decision, which could be much more efficient. For example, in the problem of online collaborative filtering, the decision set consists of matrices with a bounded trace norm, and the linear optimization step is at least an

order of magnitude faster than the projection operation [Hazan and Kale, 2012]. Moreover, it has been showed that OFW achieves an $O(T^{3/4})$ regret bound for convex losses [Hazan and Kale, 2012, Hazan, 2016] and an $O(T^{2/3})$ regret bound for strongly convex losses [Wan and Zhang, 2021, Garber and Kretzu, 2021], which are the best known regret bounds of projection-free methods without further assumptions.

However, OFW requires that the gradient $\nabla f_t(\mathbf{x}_t)$ is revealed immediately after making the decision $\mathbf{x}_t$, which is not necessarily satisfied in reality. For example, in the previously mentioned online collaborative filtering [Hazan and Kale, 2012], the decision is a prediction of a user-item rating matrix, and the corresponding gradient depends on the true rating of a user on a item, which may not be decided by the user immediately. Therefore, it is natural to consider a more practical setting, where the gradient $\nabla f_t(\mathbf{x}_t)$ arrives at the end of round $t + d_t - 1$, and $d_t \geq 1$ denotes an arbitrary delay. To handle this setting, one potential way is to combine OFW with an existing black-box technique for converting any traditional OCO algorithm into this delayed setting [Joulani et al., 2013]. To be precise, this black-box technique is to pool independent instances of OFW, each of which acts as a learner in the non-delayed setting over a subsequence of rounds. In each round, a single instance will be taken out from the pool, which makes a decision and then waits for its feedback before rejoining the pool. If the pool is empty, a new instance of OFW will be added to it. Moreover, according to Joulani et al. [2013], their black-box technique is able to attain a regret bound of $dR(T/d)$ by combining with a traditional OCO algorithm with $R(T)$ regret, where d is the maximum delay. As a result, combining it with OFW will attain an $O(d^{1/4}T^{3/4})$ regret bound for convex losses and an $O(d^{1/3}T^{2/3})$ regret bound for strongly convex losses, which magnify the regret bounds of OFW in the non-delayed setting by a coefficient about the delay. Thus, it is natural to ask whether the effect about delay can be further reduced.

In this paper, we give an affirmative answer by developing a simple method called delayed OFW, which is able to be robust to a relatively large amount of delay for both convex and strongly convex losses. Different from the black-box technique that needs to maintain multiple instances of OFW [Joulani et al., 2013], our delayed OFW is a natural extension of OFW in the delayed setting, which updates the decision similar to OFW after receiving any delayed gradient, and play the latest decision for each round. Our theoretical contributions are summarized as follows.

- First, we prove that our delayed OFW attains an $O(T^{3/4} + dT^{1/4})$ regret bound for convex losses, where d is the maximum delay, which matches the $O(T^{3/4})$ regret bound in the non-delayed setting as long as d does not exceed $O(\sqrt{T})$.

- Second, we prove that our delayed OFW attains an $O(T^{2/3} + d\log T)$ regret bound for strongly convex losses, which matches the $O(T^{2/3})$ regret bound in the non-delayed setting as long as d does not exceed $O(T^{2/3}/\log T)$.

Therefore, our regret bounds are strictly better than those achieved by combining the black-box technique [Joulani et al., 2013] with OFW, when the term about d in them is not dominant.

2 Related work

In this section, we briefly review related work on projection-free algorithms for OCO, and OCO under delayed feedback.

2.1 Projection-free algorithms for OCO

The OFW method [Hazan and Kale, 2012, Hazan, 2016] is the first projection-free algorithm for OCO, which is an online extension of the classical Frank-Wolfe method [Frank and Wolfe, 1956, Jaggi, 2013]. For convex losses, OFW first chooses an arbitrary $\mathbf{x}_1 \in \mathcal{K}$, and then iteratively updates its decision by the following linear optimization step

$$
\begin{aligned}
\mathbf{v}_t &\in \underset{\mathbf{x} \in \mathcal{K}}{\operatorname{argmin}} \langle \nabla F_t(\mathbf{x}_t), \mathbf{x} \rangle \\
\mathbf{x}_{t+1} &= \mathbf{x}_t + \sigma_t(\mathbf{v}_t - \mathbf{x}_t)
\end{aligned}
\tag{1}
$$

where $F_t(\mathbf{x})$ is a surrogate loss function defined as

$$F_t(\mathbf{x}) = \eta \sum_{i=1}^{t} \langle \nabla f_i(\mathbf{x}_i), \mathbf{x} \rangle + \|\mathbf{x} - \mathbf{x}_1\|_2^2 \tag{2}$$

and η, σ_t are two parameters. With appropriate parameters, it can attain an $O(T^{3/4})$ regret bound for convex losses.

If losses are convex and smooth, Hazan and Minasyan [2020] proposed a randomized projection-free method, which is based on a classical OCO method called follow the perturbed leader [Kalai and Vempala, 2005], and achieved a regret bound of $O(T^{2/3})$. Recently, Wan and Zhang [2021] proved that OFW can achieve an $O(T^{2/3})$ regret bound for strongly convex losses. Specifically, to utilize the strong convexity of losses, Wan and Zhang [2021] redefined $F_t(\mathbf{x})$ in (2) to

$$F_t(\mathbf{x}) = \sum_{i=1}^{t} \left(\langle \nabla f_i(\mathbf{x}_i), \mathbf{x} \rangle + \frac{\beta}{2} \|\mathbf{x} - \mathbf{x}_i\|_2^2 \right) \tag{3}$$

where β is the modulus of the strong convexity. The same regret bound was concurrently established by Garber and Kretzu [2021].

Moreover, projection-free OCO algorithms, which are able to achieve an $O(\sqrt{T})$ regret bound for convex losses and an $O(\log T)$ regret bound for strongly convex losses, were proposed for polyhedral sets [Garber and Hazan, 2016], smooth sets [Levy and Krause, 2019], and special sets that are accessible through a membership oracle [Mhammedi, 2021] or a separation oracle [Garber and Kretzu, 2022], respectively. Although the regret bounds of these algorithms are better than those of OFW, they can only be efficiently implemented in some special sets, and thus are still not able to replace OFW. Different from these algorithms limited to special sets, Wan and Zhang [2021] showed that by setting σ_t with a line search technique, OFW can adaptively utilize the strong convexity of decision sets to attain an $O(T^{2/3})$ regret bound for convex losses and an $O(\sqrt{T})$ regret bound for strongly convex losses.

Despite great progresses about projection-free OCO algorithms, these studies did not consider the practical problem of delayed feedback.

2.2 OCO under delayed feedback

The first work about OCO under delayed feedback was due to Weinberger and Ordentlich [2002], which focused on a special case with a fixed delay d', i.e., the feedback for each decision $\mathbf{x}_t$ is received at the end of round $t + d' - 1$. They proposed a technique that can convert any traditional OCO algorithm for the non-delayed setting into this delayed setting. Specifically, their technique is to run d' instances of a traditional OCO algorithm, where each instance is used every d' rounds, which is feasible because the feedback for any decision $\mathbf{x}_t$ must be available for selecting the decision $\mathbf{x}_{t+d'}$. If the traditional OCO algorithm enjoys a regret bound of $R(T)$ in the non-delayed setting, they proved that this technique achieves a regret bound of $d'R(T/d')$. As a result, by combining with OGD, the regret bound of this technique is $O(\sqrt{d'T})$ for convex losses and $O(d' \log T)$ for strongly convex losses. However, it needs to run d' instances of OGD, which requires more storage and computational resources than the original OGD. To address this problem, Langford et al. [2009] studied the same delayed setting, and showed that simply performing each gradient descent step in the original OGD with a delayed gradient can also achieve the $O(\sqrt{d'T})$ regret bound for convex losses and the $O(d' \log T)$ regret bound for strongly convex losses.

Furthermore, Joulani et al. [2013] extended the technique in Weinberger and Ordentlich [2002] to handle a more general setting, where each feedback is delayed by arbitrary rounds. For this setting, their technique is able to attain a regret bound of $dR(T/d)$ by combining with a traditional OCO algorithm with $R(T)$ regret, where d is the maximum delay. Note that although this technique can also convert OFW to the delayed setting, it can only attain an $O(d^{1/4}T^{3/4})$ regret bound for convex losses and an $O(d^{1/3}T^{2/3})$ regret bound for strongly convex losses, which cannot match regret bounds of OFW in the non-delayed setting as long as d is larger than $O(1)$. Additionally, similar to the technique in Weinberger and Ordentlich [2002], it also needs to run multiple instances of a traditional OCO algorithm, which could be prohibitively resource-intensive. Many studies

126 [Quanrud and Khashabi, 2015, Joulani et al., 2016, Wan et al., 2022, Flaspohler et al., 2021] proposed
127 delayed OCO algorithms, which only require the same storage and computational resources as in the
128 non-delayed setting, but did not consider the projection-free algorithms.

3 Main results

130 In this section, we first introduce necessary preliminaries including the problem setting, definitions,
131 and assumptions. Then, we present our delayed OFW and the corresponding theoretical guarantees
132 for convex and strongly convex losses, respectively.

3.1 Preliminaries

134 We consider the problem of OCO with arbitrary delays [Joulani et al., 2013, Quanrud and Khashabi,
135 2015]. Similar to the standard OCO, in each round $t = 1, \ldots, T$, the player first chooses a decision
136 $\mathbf{x}_t$ from the decision set $\mathcal{K}$, and then the adversary selects a convex function $f_t(\mathbf{x})$. However, different
137 from the standard OCO, the gradient $\mathbf{g}_t = \nabla f_t(\mathbf{x}_t)$ is revealed at the end of round $t + d_t - 1$, where
138 $d_t \geq 1$ denotes an arbitrary delay. As a result, the player actually receives gradients $\{\mathbf{g}_k | k \in \mathcal{F}_t\}$ at
139 the end of round t, where

$$\mathcal{F}_t = \{k | k + d_k - 1 = t\}.$$

140 Then, we recall the standard definition for strongly convex functions [Boyd and Vandenberghe, 2004].

141 **Definition 1** *A function $f(\mathbf{x}) : \mathcal{K} \to \mathbb{R}$ is called β-strongly convex over $\mathcal{K}$ if it holds that*

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{\beta}{2} \|\mathbf{y} - \mathbf{x}\|_2^2$$

142 *for all $\mathbf{x}, \mathbf{y} \in \mathcal{K}$.*

143 Finally, we introduce two assumptions, which are commonly used in studies about OCO [Shalev-
144 Shwartz, 2011, Hazan, 2016].

145 **Assumption 1** *The gradients of all loss functions are bounded by G, i.e., it holds that $\|\nabla f_t(\mathbf{x})\|_2 \leq$*
146 *G for any $\mathbf{x} \in \mathcal{K}$ and $t \in [T]$.*

147 **Assumption 2** *The diameter of the decision set $\mathcal{K}$ is bounded by D, i.e., it holds that $\|\mathbf{x} - \mathbf{y}\|_2 \leq D$*
148 *for any $\mathbf{x}, \mathbf{y} \in \mathcal{K}$.*

3.2 Delayed OFW for convex losses

150 As presented in (1) and (2), the original OFW for convex losses requires the gradient $\mathbf{g}_t$ before
151 making the decision $\mathbf{x}_{t+1}$. However, in the problem of OCO with arbitrary delays, this requirement
152 is not necessarily satisfied, because the player actually receives gradients $\{\mathbf{g}_k | k \in \mathcal{F}_t\}$ at each round
153 t, which may not contain $\mathbf{g}_t$. To address this limitation, we propose a natural generalization of OFW
154 to this delayed problem, which is described as follows.

155 The main idea is to update the decision similar to OFW for each received gradient, and play the
156 latest decision for each round. In this way, there exist some intermediate decisions that are not
157 really played. To facilitate presentations, we introduce an additional notation $\mathbf{y}_\tau$ to denote the τ-th
158 intermediate decision. Moreover, we denote the sum of τ received gradients by $\bar{\mathbf{g}}_\tau$. Initially, we
159 choose an arbitrary vector $\mathbf{y}_1 \in \mathcal{K}$ and set $\tau = 1, \bar{\mathbf{g}}_0 = \mathbf{0}$. At each round $t = 1, \ldots, T$, we play the
160 latest decision $\mathbf{x}_t = \mathbf{y}_\tau$ and query the gradient $\mathbf{g}_t = \nabla f_t(\mathbf{x}_t)$. Then, we receive delayed gradients
161 that are queried in a set of rounds $\mathcal{F}_t$, and perform the following steps for each received gradient.

162 Specifically, for any $k \in \mathcal{F}_t$, inspired by (2) of the original OFW, we first compute $\bar{\mathbf{g}}_\tau = \bar{\mathbf{g}}_{\tau-1} + \mathbf{g}_k$
163 and define

$$F_\tau(\mathbf{y}) = \eta \langle \bar{\mathbf{g}}_\tau, \mathbf{y} \rangle + \|\mathbf{y} - \mathbf{y}_1\|_2^2.$$

164 Then, similar to (1) of the original OFW, we perform the following update

$$\mathbf{v}_\tau \in \underset{\mathbf{y} \in \mathcal{K}}{\operatorname{argmin}} \langle \nabla F_\tau(\mathbf{y}_\tau), \mathbf{y} \rangle$$

$$\mathbf{y}_{\tau+1} = \mathbf{y}_\tau + \sigma_\tau (\mathbf{v}_\tau - \mathbf{y}_\tau)$$

Algorithm 1 Delayed OFW for Convex Losses

1: **Input:** η
2: **Initialization:** choose an arbitrary vector $\mathbf{y}_1 \in \mathcal{K}$ and set $\tau = 1, \bar{\mathbf{g}}_0 = \mathbf{0}$
3: **for** $t = 1, 2, \ldots, T$ **do**
4:     Play $\mathbf{x}_t = \mathbf{y}_\tau$ and query $\mathbf{g}_t = \nabla f_t(\mathbf{x}_t)$
5:     Receive a set of delayed gradients $\{\mathbf{g}_k | k \in \mathcal{F}_t\}$
6:     **for** $k \in \mathcal{F}_t$ **do**
7:         $\bar{\mathbf{g}}_\tau = \bar{\mathbf{g}}_{\tau-1} + \mathbf{g}_k$
8:         Define $F_\tau(\mathbf{y}) = \eta\langle\bar{\mathbf{g}}_\tau, \mathbf{y}\rangle + \|\mathbf{y} - \mathbf{y}_1\|_2^2$
9:         $\mathbf{v}_\tau \in \operatorname{argmin}_{\mathbf{y}\in\mathcal{K}}\langle\nabla F_\tau(\mathbf{y}_\tau), \mathbf{y}\rangle$
10:        $\mathbf{y}_{\tau+1} = \mathbf{y}_\tau + \sigma_\tau(\mathbf{v}_\tau - \mathbf{y}_\tau)$ with σ_τ in (4)
11:        $\tau = \tau + 1$
12:    **end for**
13: **end for**

---

where σ_τ is a parameter. Following Wan and Zhang [2021], it is set by a line search rule

$$\sigma_\tau = \operatorname*{argmin}_{\sigma\in[0,1]}\langle\sigma(\mathbf{v}_\tau - \mathbf{y}_\tau), \nabla F_\tau(\mathbf{y}_\tau)\rangle + \sigma^2\|\mathbf{v}_\tau - \mathbf{y}_\tau\|_2^2. \tag{4}$$

Finally, we update $\tau = \tau + 1$ so that τ still indexes the latest intermediate decision.

The detailed procedures are summarized in Algorithm 1, which is named as delayed OFW for convex losses. Define

$$d = \max\{d_1, \ldots, d_T\}.$$

We first establish the following theorem with respect to the regret of Algorithm 1.

Theorem 1 *For any $\mathbf{x}^* \in \mathcal{K}$, under Assumptions 1 and 2, Algorithm 1 with $\eta = \frac{D}{\sqrt{2}G(T+2)^{3/4}}$ has*

$$\sum_{t=1}^{T}[f_t(\mathbf{x}_t) - f_t(\mathbf{x}^*)] = O(T^{3/4} + dT^{1/4}).$$

Theorem 1 shows that our Algorithm 1 can achieve an $O(T^{3/4} + dT^{1/4})$ regret bound for convex losses with arbitrary delays. This bound matches the $O(T^{3/4})$ regret bound of OFW in the non-delayed setting [Hazan and Kale, 2012, Hazan, 2016], as long as d does not exceed $O(\sqrt{T})$. Moreover, it is better than the $O(d^{1/4}T^{3/4})$ regret bound achieved by combining the technique of Joulani et al. [2013] and the $O(T^{3/4})$ regret bound of OFW for convex losses, as long as d does not exceed $O(T^{2/3})$.

3.3 Delayed OFW for strongly convex losses

We proceed to handle β-strongly convex losses by slightly modifying Algorithm 1. Recall that in the standard OCO without delays, the critical idea of utilizing the strong convexity of losses is to replace the surrogate loss function in (1) by that in (3) [Wan and Zhang, 2021]. The main difference is that the regularization term in (3) is about all historical decisions, instead of only the initial decision.

Inspired by (3), we first redefine $F_\tau(\mathbf{y})$ in Algorithm 1 to

$$F_\tau(\mathbf{y}) = \langle\bar{\mathbf{g}}_\tau, \mathbf{y}\rangle + \sum_{i=1}^{\tau}\frac{\beta}{2}\|\mathbf{y} - \mathbf{y}_i\|_2^2.$$

Second, since $F_\tau(\mathbf{y})$ is modified, we adjust the line search rule to

$$\sigma_\tau = \operatorname*{argmin}_{\sigma\in[0,1]}\langle\sigma(\mathbf{v}_\tau - \mathbf{y}_\tau), \nabla F_\tau(\mathbf{y}_\tau)\rangle + \frac{\beta\tau\sigma^2}{2}\|\mathbf{v}_\tau - \mathbf{y}_\tau\|_2^2. \tag{5}$$

The detailed procedures are summarized in Algorithm 2, which is named as delayed OFW for strongly convex losses.

Then, we establish the following theorem about the regret of Algorithm 2.

Algorithm 2 Delayed OFW for Strongly Convex Losses

1: **Input:** β
2: **Initialization:** choose an arbitrary vector $\mathbf{y}_1 \in \mathcal{K}$ and set $\tau = 1, \bar{\mathbf{g}}_0 = \mathbf{0}$
3: **for** $t = 1, 2, \ldots, T$ **do**
4:      Play $\mathbf{x}_t = \mathbf{y}_\tau$ and query $\mathbf{g}_t = \nabla f_t(\mathbf{x}_t)$
5:      Receive a set of delayed gradients $\{\mathbf{g}_k | k \in \mathcal{F}_t\}$
6:      **for** $k \in \mathcal{F}_t$ **do**
7:          $\bar{\mathbf{g}}_\tau = \bar{\mathbf{g}}_{\tau-1} + \mathbf{g}_k$
8:          Define $F_\tau(\mathbf{y}) = \langle \bar{\mathbf{g}}_\tau, \mathbf{y} \rangle + \sum_{i=1}^{\tau} \frac{\beta}{2} \|\mathbf{y} - \mathbf{y}_i\|_2^2$
9:          $\mathbf{v}_\tau \in \arg\min_{\mathbf{y} \in \mathcal{K}} \langle \nabla F_\tau(\mathbf{y}_\tau), \mathbf{y} \rangle$
10:         $\mathbf{y}_{\tau+1} = \mathbf{y}_\tau + \sigma_\tau(\mathbf{v}_\tau - \mathbf{y}_\tau)$ with σ_τ in (5)
11:         $\tau = \tau + 1$
12:      **end for**
13: **end for**

Theorem 2 *For any $\mathbf{x}^* \in \mathcal{K}$, suppose Assumptions 1 and 2 hold, and all losses are β-strongly convex, Algorithm 2 has*

$$\sum_{t=1}^{T} [f_t(\mathbf{x}_t) - f_t(\mathbf{x}^*)] = O(T^{2/3} + d \log T).$$

Theorem 2 shows that our Algorithm 2 can achieve an $O(T^{2/3} + d \log T)$ regret bound for strongly convex losses with arbitrary delays. First, this bound is better than the $O(T^{3/4} + dT^{1/4})$ regret bound in Theorem 1, which is established by only using the convexity condition. Second, it matches the $O(T^{2/3})$ regret bound of OFW for strongly convex losses in the non-delayed setting [Garber and Kretzu, 2021, Wan and Zhang, 2021], as long as d does not exceed $O(T^{2/3} / \log T)$. Third, it is better than the $O(d^{1/3}T^{2/3})$ regret bound achieved by combining the technique of Joulani et al. [2013] and the $O(T^{2/3})$ regret bound of OFW for strongly convex losses, as long as d does not exceed $O(T^{2/3})$.

4 Theoretical analysis

In this section, we first introduce necessary preliminaries for our analysis, and then prove Theorems 1 and 2 by introducing some lemmas. Due to the limitation of space, the proofs for those lemmas are provided in the supplementary material.

4.1 Preliminaries

Because of the effect of delays, there may exist some gradients that arrive after round T. Although our Algorithms 1 and 2 do not need to use these gradients, they are useful for the analysis. As a result, in our analysis, we further set $\mathbf{x}_t = \mathbf{y}_\tau$ and perform steps 5 to 12 of Algorithms 1 and 2 for any $t = T+1, \ldots, T+d-1$. In this way, all gradients $\mathbf{g}_1, \mathbf{g}_2, \ldots, \mathbf{g}_T$ queried by our algorithms are utilized, which produces decisions $\mathbf{y}_1, \mathbf{y}_2, \ldots, \mathbf{y}_{T+1}$.

Then, let $\tau_t = 1 + \sum_{i=1}^{t-1} |\mathcal{F}_i|$ for any $t \in [T+d]$. It is not hard to verify that our Algorithms 1 and 2 ensure that

$$\mathbf{x}_t = \mathbf{y}_{\tau_t} \tag{6}$$

for any $t \in [T+d-1]$. Next, we define

$$\mathcal{I}_t = \begin{cases} \emptyset, & \text{if } |\mathcal{F}_t| = 0, \\ \{\tau_t, \tau_t + 1, \ldots, \tau_{t+1} - 1\}, & \text{otherwise.} \end{cases} \tag{7}$$

Let $s = \min\{t | t \in [T+d-1], |\mathcal{F}_t| > 0\}$. It is not hard to verify that

$$\cup_{t=s}^{T+d-1} \mathcal{I}_t = [T] \text{ and } \mathcal{I}_i \cap \mathcal{I}_j = \emptyset, \forall i \neq j \tag{8}$$

and

$$\cup_{t=s}^{T+d-1} \mathcal{F}_t = [T] \text{ and } \mathcal{F}_i \cap \mathcal{F}_j = \emptyset, \forall i \neq j. \tag{9}$$

To facilitate the analysis, we further denote the time-stamp of the τ-th gradient used in the update of our Algorithms 1 and 2 by c_τ. To help understanding, one can imagine that our Algorithms 1 and 2 also implement $c_\tau = k$ in their step 7. If $|\mathcal{F}_t| \neq 0$, we have

$$\{c_{\tau_t}, \ldots, c_{\tau_{t+1}-1}\} = \mathcal{F}_t. \tag{10}$$

Moreover, by using this notation, $F_\tau(\mathbf{y})$ defined in Algorithms 1 and 2 is respectively equivalent to

$$F_\tau(\mathbf{y}) = \eta \sum_{i=1}^{\tau} \langle \mathbf{g}_{c_i}, \mathbf{y} \rangle + \|\mathbf{y} - \mathbf{y}_1\|_2^2, \tag{11}$$

$$F_\tau(\mathbf{y}) = \sum_{i=1}^{\tau} \langle \mathbf{g}_{c_i}, \mathbf{y} \rangle + \sum_{i=1}^{\tau} \frac{\beta}{2} \|\mathbf{y} - \mathbf{y}_i\|_2^2. \tag{12}$$

4.2   Proof of Theorem 1

Let $t' = t + d_t - 1$ for any $t \in [T]$. According to the convexity of $f_t(\mathbf{x})$, we have

$$f_t(\mathbf{x}_t) - f_t(\mathbf{x}^*) \leq \langle \mathbf{g}_t, \mathbf{x}_t - \mathbf{x}^* \rangle = \langle \mathbf{g}_t, \mathbf{x}_{t'} - \mathbf{x}^* \rangle + \langle \mathbf{g}_t, \mathbf{x}_t - \mathbf{x}_{t'} \rangle$$
$$\leq \langle \mathbf{g}_t, \mathbf{x}_{t'} - \mathbf{x}^* \rangle + G\|\mathbf{x}_t - \mathbf{x}_{t'}\|_2 = \langle \mathbf{g}_t, \mathbf{y}_{\tau_{t'}} - \mathbf{x}^* \rangle + G\|\mathbf{y}_{\tau_t} - \mathbf{y}_{\tau_{t'}}\|_2$$

where the last equality is due to (6).

By summing over $t = 1, \ldots, T$, we have

$$\sum_{t=1}^{T} [f_t(\mathbf{x}_t) - f_t(\mathbf{x}^*)] \leq \sum_{t=1}^{T} \langle \mathbf{g}_t, \mathbf{x}_t - \mathbf{x}^* \rangle \leq \sum_{t=1}^{T} \langle \mathbf{g}_t, \mathbf{y}_{\tau_{t'}} - \mathbf{x}^* \rangle + \underbrace{\sum_{t=1}^{T} G\|\mathbf{y}_{\tau_t} - \mathbf{y}_{\tau_{t'}}\|_2}_{:=A}. \tag{13}$$

Then, we bound the first term in the right side of (13) as follows

$$\sum_{t=1}^{T} \langle \mathbf{g}_t, \mathbf{y}_{\tau_{t'}} - \mathbf{x}^* \rangle = \sum_{t=s}^{T+d-1} \sum_{i \in \mathcal{F}_t} \langle \mathbf{g}_i, \mathbf{y}_{\tau_{i+d_i-1}} - \mathbf{x}^* \rangle$$

$$= \sum_{t=s}^{T+d-1} \sum_{i \in \mathcal{F}_t} \langle \mathbf{g}_i, \mathbf{y}_{\tau_t} - \mathbf{x}^* \rangle = \sum_{t=s}^{T+d-1} \sum_{i=\tau_t}^{\tau_{t+1}-1} \langle \mathbf{g}_{c_i}, \mathbf{y}_{\tau_t} - \mathbf{x}^* \rangle$$

$$= \sum_{t=s}^{T+d-1} \sum_{i=\tau_t}^{\tau_{t+1}-1} \left(\langle \mathbf{g}_{c_i}, \mathbf{y}_i - \mathbf{x}^* \rangle + \langle \mathbf{g}_{c_i}, \mathbf{y}_{\tau_t} - \mathbf{y}_i \rangle \right) \tag{14}$$

$$= \sum_{t=1}^{T} \langle \mathbf{g}_{c_t}, \mathbf{y}_t - \mathbf{x}^* \rangle + \sum_{t=s}^{T+d-1} \sum_{i=\tau_t}^{\tau_{t+1}-1} \langle \mathbf{g}_{c_i}, \mathbf{y}_{\tau_t} - \mathbf{y}_i \rangle$$

$$\leq \underbrace{\sum_{t=1}^{T} \langle \mathbf{g}_{c_t}, \mathbf{y}_t - \mathbf{x}^* \rangle}_{:=B} + \underbrace{\sum_{t=s}^{T+d-1} \sum_{i=\tau_t}^{\tau_{t+1}-1} G\|\mathbf{y}_{\tau_t} - \mathbf{y}_i\|_2}_{:=C}$$

where the first equality is due to (9), the second equality is due to $i + d_i - 1 = t$ for any $i \in \mathcal{F}_t$, the third equality is due to (10), and the last equality is due to (7) and (8).

By combining (13) and (14), we have

$$\sum_{t=1}^{T} [f_t(\mathbf{x}_t) - f_t(\mathbf{x}^*)] \leq A + B + C. \tag{15}$$

Next, we proceed to bound terms A, B, and C defined above. Specifically, we first establish the following bound for the sum of terms A and C by carefully analyzing the distance $\|\mathbf{y}_{\tau_t} - \mathbf{y}_{\tau_{t'}}\|_2$ in the term A and the distance $\|\mathbf{y}_{\tau_t} - \mathbf{y}_i\|_2$ in the term C.

Lemma 1 *Let $\mathbf{y}_t^* = \operatorname{argmin}_{\mathbf{y} \in \mathcal{K}} F_{t-1}(\mathbf{y})$ for any $t \in [T+1]$, where $F_t(\mathbf{y})$ is defined in (11). Suppose Assumption 1 and 2 hold, and there exist some constants $\gamma > 0$ and $0 < \alpha \le 1$ such that $F_{t-1}(\mathbf{y}_t) - F_{t-1}(\mathbf{y}_t^*) \le \gamma(t+2)^{-\alpha}$ for any $t \in [T+1]$, Algorithm 1 ensures*

$$A + C \le 3dGD + 4Gd\sqrt{\gamma} + \frac{8G\sqrt{\gamma}}{2-\alpha}T^{1-\alpha/2} + \frac{3\eta G^2 dT}{2}$$

where $A = \sum_{t=1}^{T} G\|\mathbf{y}_{\tau_t} - \mathbf{y}_{\tau_{t'}}\|_2$ and $C = \sum_{t=s}^{T+d-1}\sum_{i=\tau_t}^{\tau_{t+1}-1} G\|\mathbf{y}_{\tau_t} - \mathbf{y}_i\|_2$.

Note that Lemma 1 introduces an assumption about $\mathbf{y}_t$ and $F_{t-1}(\mathbf{y})$. According to our Algorithm 1, $\mathbf{y}_t$ is actually generated by approximately minimizing $F_{t-1}(\mathbf{y})$ with a linear optimization step. Therefore, by following the analysis of the original OFW [Hazan, 2016], we show that this assumption can be satisfied with $\gamma = 8D^2$ and $\alpha = 1/2$.

Lemma 2 *Let $\mathbf{y}_t^* = \operatorname{argmin}_{\mathbf{y} \in \mathcal{K}} F_{t-1}(\mathbf{y})$ for any $t \in [T+1]$, where $F_t(\mathbf{y})$ is defined in (11). Under Assumptions 1 and 2, for any $t \in [T+1]$, Algorithm 1 with $\eta = \frac{D}{\sqrt{2}G(T+2)^{3/4}}$ has*

$$F_{t-1}(\mathbf{y}_t) - F_{t-1}(\mathbf{y}_t^*) \le \frac{8D^2}{\sqrt{t+2}}.$$

Then, by combining $\eta = \frac{D}{\sqrt{2}G(T+2)^{3/4}}$ with Lemmas 1 and 2, we have

$$A + C \le (3 + 8\sqrt{2})GDd + \frac{32\sqrt{2}GD}{3}T^{3/4} + \frac{3GDdT^{1/4}}{2\sqrt{2}} = O(T^{3/4} + dT^{1/4}). \quad (16)$$

Furthermore, by following the analysis of the original OFW [Hazan, 2016], we establish an upper bound for the term B.

Lemma 3 *Under Assumptions 1 and 2, for any $\mathbf{x}^* \in \mathcal{K}$, Algorithm 1 with $\eta = \frac{D}{\sqrt{2}G(T+2)^{3/4}}$ ensures*

$$\sum_{t=1}^{T} \langle \mathbf{g}_{c_t}, \mathbf{y}_t - \mathbf{x}^* \rangle \le \frac{11\sqrt{2}GD(T+2)^{3/4}}{3} + \frac{GDT^{1/4}}{\sqrt{2}}.$$

Finally, by combining (15), (16), and Lemma 3, we complete this proof.

4.3 Proof of Theorem 2

Since $f_t(\mathbf{x})$ is β-strongly convex, we have

$$\sum_{t=1}^{T} [f_t(\mathbf{x}_t) - f_t(\mathbf{x}^*)] \le \sum_{t=1}^{T} \langle \mathbf{g}_t, \mathbf{x}_t - \mathbf{x}^* \rangle - \sum_{t=1}^{T} \frac{\beta}{2}\|\mathbf{x}_t - \mathbf{x}^*\|_2^2. \quad (17)$$

Then, we note that the first term in the right side of (17) can be bounded by reusing (13) and (14). Specifically, we have

$$\sum_{t=1}^{T} [f_t(\mathbf{x}_t) - f_t(\mathbf{x}^*)] \le A + C + \sum_{t=1}^{T} \langle \mathbf{g}_{c_t}, \mathbf{y}_t - \mathbf{x}^* \rangle - \sum_{t=1}^{T} \frac{\beta}{2}\|\mathbf{x}_t - \mathbf{x}^*\|_2^2 \quad (18)$$

where terms A and C are defined in (13) and (14).

Next, we consider the last term in the right side of (18). For any $\mathbf{y}_t, \mathbf{x}_t, \mathbf{x}^* \in \mathcal{K}$, we have

$$\begin{aligned}
\|\mathbf{y}_t - \mathbf{x}^*\|_2^2 &= \|\mathbf{y}_t - \mathbf{x}_t\|_2^2 + \|\mathbf{x}_t - \mathbf{x}^*\|_2^2 + 2\langle \mathbf{y}_t - \mathbf{x}_t, \mathbf{x}_t - \mathbf{x}^* \rangle \\
&\le \|\mathbf{y}_t - \mathbf{x}_t\|_2^2 + \|\mathbf{x}_t - \mathbf{x}^*\|_2^2 + 2\|\mathbf{y}_t - \mathbf{x}_t\|_2\|\mathbf{x}_t - \mathbf{x}^*\|_2 \\
&\le 3D\|\mathbf{y}_t - \mathbf{x}_t\|_2 + \|\mathbf{x}_t - \mathbf{x}^*\|_2^2
\end{aligned}$$

where the last inequality is due to Assumption 2.

By combining the above inequality and (6) with (18), we have

$$\sum_{t=1}^{T}[f_t(\mathbf{x}_t) - f_t(\mathbf{x}^*)] \leq A + C + \sum_{t=1}^{T}\left(\langle \mathbf{g}_{c_t}, \mathbf{y}_t - \mathbf{x}^*\rangle - \frac{\beta}{2}\|\mathbf{y}_t - \mathbf{x}^*\|_2^2\right) + \underbrace{\sum_{t=1}^{T}\frac{3\beta D}{2}\|\mathbf{y}_t - \mathbf{y}_{\tau_t}\|_2}_{:=E}.$$

(19)

Then, we proceed to establish upper bounds for terms A, C, and E by carefully analyzing the distance $\|\mathbf{y}_{\tau_t} - \mathbf{y}_{\tau_{t'}}\|_2$ in the term A, the distance $\|\mathbf{y}_{\tau_t} - \mathbf{y}_i\|_2$ in the term C, and the distance $\|\mathbf{y}_t - \mathbf{y}_{\tau_t}\|_2$ in the term E.

Lemma 4 *Let $\mathbf{y}_t^* = \operatorname{argmin}_{\mathbf{y}\in\mathcal{K}} F_{t-1}(\mathbf{y})$ for any $t = 2, \ldots, T+1$, where $F_t(\mathbf{y})$ is defined in (12). Suppose Assumption 1 and 2 hold, all losses are β-strongly convex, and there exist some constants $\gamma > 0$ and $0 \leq \alpha < 1$ such that $F_{t-1}(\mathbf{y}_t) - F_{t-1}(\mathbf{y}_t^*) \leq \gamma(t-1)^\alpha$ for any $t = 2, \ldots, T+1$, Algorithm 1 ensures*

$$E \leq 3dD\sqrt{2\beta\gamma} + \frac{6D\sqrt{2\beta\gamma}}{1+\alpha}T^{(1+\alpha)/2} + 3\beta dD^2 + 3D(G+\beta D)d\ln T,$$

$$A + C \leq 3dGD + \frac{4G(G+\beta D)d(1+\ln T)}{\beta} + 4dG\sqrt{\frac{2\gamma}{\beta}} + \sqrt{\frac{2\gamma}{\beta}}\frac{8G}{1+\alpha}T^{(1+\alpha)/2},$$

where $A = \sum_{t=1}^{T} G\|\mathbf{y}_{\tau_t} - \mathbf{y}_{\tau_{t'}}\|_2$, $C = \sum_{t=s}^{T+d-1}\sum_{i=\tau_t}^{\tau_{t+1}-1} G\|\mathbf{y}_{\tau_t} - \mathbf{y}_i\|_2$, and $E = \sum_{t=1}^{T}\frac{3\beta D}{2}\|\mathbf{y}_t - \mathbf{y}_{\tau_t}\|_2$.

Note that Lemma 4 also introduces an assumption about $\mathbf{y}_t$ and $F_{t-1}(\mathbf{y})$. According to our Algorithm 2, $\mathbf{y}_t$ is actually generated by approximately minimizing $F_{t-1}(\mathbf{y})$ with a linear optimization step. Therefore, by following the analysis of OFW for strongly convex losses [Wan and Zhang, 2021], we show that this assumption is satisfied with $\gamma = 16(G+2\beta D)^2/\beta$ and $\alpha = 1/3$.

Lemma 5 *Let $\mathbf{y}_t^* = \operatorname{argmin}_{\mathbf{y}\in\mathcal{K}} F_{t-1}(\mathbf{y})$ for any $t = 2, \ldots, T+1$, where $F_t(\mathbf{y})$ is defined in (12). Suppose Assumption 1 and 2 hold, and all losses are β-strongly convex, for any $t = 2, \ldots, T+1$, Algorithm 2 has*

$$F_{t-1}(\mathbf{y}_t) - F_{t-1}(\mathbf{y}_t^*) \leq \frac{16(G+2\beta D)^2(t-1)^{1/3}}{\beta}.$$

By combining Lemmas 4 and 5, we have

$$A + C + E = O\left(T^{(1+\alpha)/2} + d\log T\right) = O(T^{2/3} + d\log T).$$

(20)

Furthermore, by following the analysis of OFW for strongly convex losses [Wan and Zhang, 2021], we establish an upper bound for the third term in (19).

Lemma 6 *Suppose Assumption 1 and 2 hold, and all losses are β-strongly convex, for any $\mathbf{x}^* \in \mathcal{K}$, Algorithm 2 ensures*

$$\sum_{t=1}^{T}\left(\langle \mathbf{g}_{c_t}, \mathbf{y}_t - \mathbf{x}^*\rangle - \frac{\beta}{2}\|\mathbf{y}_t - \mathbf{x}^*\|_2^2\right) \leq \frac{6\sqrt{2}(G+2\beta D)^2 T^{2/3}}{\beta} + \frac{2(G+2\beta D)^2\ln T}{\beta}$$
$$+ (G+\beta D)D.$$

Finally, by combining (19), (20), and Lemma 6, we complete this proof.

5 Conclusion and future work

In this paper, we propose delayed OFW for OCO with arbitrary delays. For convex losses, we show that it attains an $O(T^{3/4} + dT^{1/4})$ regret bound, which matches the $O(T^{3/4})$ regret bound of OFW in the non-delayed setting, as long as d does not exceed $O(\sqrt{T})$. When losses are strongly convex, we further prove that it can attain an $O(T^{2/3} + d\log T)$ regret bound, which matches the $O(T^{2/3})$ regret bound of OFW in the non-delayed setting, as long as d does not exceed $O(T^{2/3}/\log T)$. These results demonstrate that delayed OFW is robust to a relatively large amount of delay.

This paper only extends the classical OFW to the delayed setting. In the future, we will investigate how to develop delayed variants for other projection-free online algorithms.

References

Stephen Boyd and Lieven Vandenberghe. *Convex Optimization*. Cambridge University Press, 2004.

Genevieve E Flaspohler, Francesco Orabona, Judah Cohen, Soukayna Mouatadid, Miruna Oprescu, Paulo Orenstein, and Lester Mackey. Online learning with optimism and delay. In *Proceedings of the 38th International Conference on Machine Learning*, pages 3363–3373, 2021.

Marguerite Frank and Philip Wolfe. An algorithm for quadratic programming. *Naval Research Logistics Quarterly*, 3(1–2):95–110, 1956.

Dan Garber and Elad Hazan. A linearly convergent conditional gradient algorithm with applications to online and stochastic optimization. *SIAM Journal on Optimization*, 26(3):1493–1528, 2016.

Dan Garber and Ben Kretzu. Revisiting projection-free online learning: the strongly convex case. In *Proceedings of the 24th International Conference on Artificial Intelligence and Statistics*, pages 3592–3600, 2021.

Dan Garber and Ben Kretzu. New projection-free algorithms for online convex optimization with adaptive regret guarantees. *arXiv:2202.04721*, 2022.

Elad Hazan. Introduction to online convex optimization. *Foundations and Trends in Optimization*, 2 (3–4):157–325, 2016.

Elad Hazan and Satyen Kale. Projection-free online learning. In *Proceedings of the 29th International Conference on Machine Learning*, pages 1843–1850, 2012.

Elad Hazan and Edgar Minasyan. Faster projection-free online learning. In *Proceedings of the 33rd Annual Conference on Learning Theory*, pages 1877–1893, 2020.

Elad Hazan, Amit Agarwal, and Satyen Kale. Logarithmic regret algorithms for online convex optimization. *Machine Learning*, 69(2):169–192, 2007.

Martin Jaggi. Revisiting frank-wolfe: Projection-free sparse convex optimization. In *Proceedings of the 30th International Conference on Machine Learning*, pages 427–435, 2013.

Pooria Joulani, András György, and Csaba Szepesvári. Online learning under delayed feedback. In *Proceedings of the 30th International Conference on Machine Learning*, pages 1453–1461, 2013.

Pooria Joulani, András György, and Csaba Szepesvári. Delay-tolerant online convex optimization: Unified analysis and adaptive-gradient algorithms. *Proceedings of the 30th AAAI Conference on Artificial Intelligence*, pages 1744–1750, 2016.

Adam Kalai and Santosh Vempala. Efficient algorithms for online decision problems. *Journal of Computer and System Sciences*, 71(3):291–307, 2005.

John Langford, Alexander J. Smola, and Martin Zinkevich. Slow learners are fast. In *Advances in Neural Information Processing Systems 22*, pages 2331–2339, 2009.

Kfir Y. Levy and Andreas Krause. Projection free online learning over smooth sets. In *Proceedings of the 22nd International Conference on Artificial Intelligence and Statistics*, pages 1458–1466, 2019.

Zakaria Mhammedi. Efficient projection-free online convex optimization with membership oracle. *arXiv:2111.05818*, 2021.

Kent Quanrud and Daniel Khashabi. Online learning with adversarial delays. In *Advances in Neural Information Processing Systems 28*, pages 1270–1278, 2015.

Shai Shalev-Shwartz. Online learning and online convex optimization. *Foundations and Trends in Machine Learning*, 4(2):107–194, 2011.

Yuanyu Wan and Lijun Zhang. Projection-free online learning over strongly convex sets. In *Proceedings of the 35th AAAI Conference on Artificial Intelligence*, pages 10076–10084, 2021.

Yuanyu Wan, Wei-Wei Tu, and Lijun Zhang. Online strongly convex optimization with unknown delays. *Machine Learning*, 111(3):871–893, 2022.

Marcelo J. Weinberger and Erik Ordentlich. On delayed prediction of individual sequences. *IEEE Transactions on Information Theory*, 48(7):1959–1976, 2002.

Martin Zinkevich. Online convex programming and generalized infinitesimal gradient ascent. In *Proceedings of the 20th International Conference on Machine Learning*, pages 928–936, 2003.