A general approximation lower bound in L^p norm, with applications to feedforward neural networks

Abstract

We study the fundamental limits to the expressive power of neural networks. Given two sets F, G of real-valued functions, we first prove a general lower bound on how well functions in F can be approximated in $L^p(\mu)$ norm by functions in G, for any $p \geq 1$ and any probability measure μ. The lower bound depends on the packing number of F, the range of F, and the fat-shattering dimension of G. We then instantiate this bound to the case where G corresponds to a piecewise-polynomial feedforward neural network, and describe in details the application to two sets F: Hölder balls and multivariate monotonic functions. Beside matching (known or new) upper bounds up to log factors, our lower bounds shed some light on the similarities or differences between approximation in L^p norm or in sup norm, solving an open question by DeVore et al. [DHP21]. Our proof strategy differs from the sup norm case and uses a key probability result of Mendelson [Men02].

1 Introduction

Neural networks are known for their great expressive power: in classification, they can interpolate arbitrary labels [ZBH+21], while in regression they have universal approximation properties [Cyb89, Hor91, LLPS93, KL20], with approximation rates that can outperform those of linear approximation methods [Yar18, DHP21]. Though the approximation problem is often only one part of the underlying learning problem (where generalization and optimization properties are also at stake), understanding the fundamental limits to the approximation properties of neural networks is key, both conceptually and for practical issues such as designing the right network architecture for the right problem.

Setting and related works. One way to quantify the expressive power of neural networks is through the following problem (some informal statements will be made more precise in the next sections). Let G be the set of all functions $g_{\mathbf{w}} : \mathcal{X} \subset \mathbb{R}^d \to \mathbb{R}$ that can be represented by tuning the weights $\mathbf{w} \in \mathbb{R}^W$ of a feedforward neural network with a fixed architecture, and let F be any set of real-valued functions on $\mathcal{X}$. A natural question is: how well functions $f \in F$ can be approximated by functions $g_{\mathbf{w}} \in G$? More precisely, given a norm $\| \cdot \|$ on functions, what is the order of magnitude of the (worst-case) *approximation error of F by G* defined by

$$\sup_{f \in F} \inf_{g_{\mathbf{w}} \in G} \| f - g_{\mathbf{w}} \| , \tag{1}$$

and how small can it be given the numbers W, L of weights and layers, and some properties of F?

The case when $\| \cdot \|$ is the sup norm (defined by $\|f\|_\infty = \sup_{x \in \mathcal{X}} |f(x)|$) is rather well understood at least in some special cases. For example, when F is a Hölder ball of smoothness $s > 0$ and the

network uses the ReLU activation function, Yarotsky [Yar17] derived a lower bound on (1) of the order of $W^{-2s/d}$, later refined to $(LW)^{-s/d}$ (up to log factors) by [Yar18, YZ20] when the depth of the network varies from $L = 1$ to $L \approx W$. Using the bit extraction technique, these authors showed that these lower bounds are achievable (up to log factors) with a carefully designed ReLU network architecture. Refined results in terms of width and depth were obtained by [SYZ22] when $s \leq 1$, while some other activation functions were also studied in [YZ20].

In this paper, we study (1) with the $L^p(\mu)$ norm, defined by $\|f\|_{L^p(\mu)} = \left(\int_X |f(x)|^p d\mu(x) \right)^{1/p}$, for $p \geq 1$ and some probability measure μ on $\mathcal{X}$. Since this corresponds to approximating functions in F in a more "average" sense than in sup norm, a natural question is whether the same accuracy can be achieved with a smaller network or not. Unfortunately, however, the proof strategies behind the lower bounds of [Yar17, Yar18, YZ20, SYZ22] are specific to the sup norm (see Remark 1 in Section 3 for details). DeVore et al. [DHP21] indeed commented: "When we move to the case $p < \infty$, the situation is even less clear [...] we cannot use the VC dimension theory for $L^p(\Omega)$ approximation. [...] What is missing vis-à-vis Problem 8.13 is what the best bounds are and how we prove lower bounds for approximation rates in $L^p(\Omega)$, $p \neq \infty$."

Existing lower bounds in $L^p(\mu)$ norm. Several papers provided lower bounds in some special cases, under some restrictions on the set to approximate F, the neural network, the approximation metric, or the encoding map $f \in F \mapsto \mathbf{w}(f) \in \mathbb{R}^W$.

When F is a space of smoothness s, a first result which is based on [DHM89] states that when imposing the weights to depend continuously on the function to be approximated, one can not achieve a better approximation rate than $W^{-\frac{s}{d}}$.

For the same F, another result for $p = 2$ and for activation functions which are continuous ([Mai99, MMR99]) prove a lower bound on the approximation of functions of smoothness s on a compact of $\mathbb{R}^d$, by one hidden-layer neural networks, of order $W^{-\frac{s}{d-1}}$. A matching upper bound is proven for a particular activation function, which is sigmoidal but pathological ([MP99]). For this same activation function, they prove that contrary to the one-hidden-layer case, there is no lower bound in the case of two-hidden-layer networks. The result is based on the Kolmogorov-Arnold superposition theorem.

In [SX21], the authors study approximation by shallow neural networks with bounded weights and activations of the form ReLU^k for an integer k. They approximate the closure of the convex hull of shallow ReLU^k-neural networks with constrained weights. They obtain optimal lower bounds of order $W^{-\frac{1}{2} - \frac{2k+1}{2d}}$ in any norm $\| \cdot \|_X$ where X is a Banach space to which the approximation functions belong and such that these functions are uniformly bounded w.r.t. $\| \cdot \|_X$. Although we only consider approximation in $L^p(\mu)$ norm, our results complement the latter by addressing neural networks with unbounded weights and arbitrary depth, and general sets F.

Approximation lower bounds in $L^p(\mu)$ norm, $p \geq 1$, have also been studied in the quantized neural networks setting (networks with weights encoded with a fixed number of bits). In [PV18], under weak assumptions on the activation function, the authors prove a lower bound on the minimal number of nonzero weights W that are required for a network to approximate a class of binary classifiers with L^p error at most ε. They show that W is at least of the order $\varepsilon^{-\frac{p(d-1)}{\beta}} \log_2^{-1} (1/\varepsilon)$, where β is a smoothness parameter. Later works including [VP19, GR21] derive lower bounds for approximation by quantized networks in various norms.

Main contributions and outline of the paper. We prove lower bounds on the approximation error (1) in any $L^p(\mu)$ norm, for non-quantized networks of arbitrary depth, and general sets F. Our main contributions are the following.

In Section 2, we first prove a general lower bound for any two sets F, G of real-valued functions on a set $\mathcal{X}$ (Theorem 1). The lower bound depends on the packing number of F, the range of F, and the fat-shattering dimension of G. We then derive a versatile corollary when G corresponds to a piecewise-polynomial feedforward neural network (Corollary 1), solving the question by DeVore et al. [DHP21]. Importantly, our proof strategy still relies on VC dimension theory, but differs from the

sup norm case in using a key probability result of Mendelson [Men02], to relate approximation in $L^p(\mu)$ norm with the fat-shattering dimension of G.

In Sections 3–4 we apply this corollary to the approximation of two sets: Hölder balls and multivariate monotonic functions. Beside matching (known or new) upper bounds up to log factors, our lower bounds shed some light on the similarities or differences between approximation in L^p norm or in sup norm. In particular, with ReLU networks, Hölder balls are not easier to approximate in L^p norm than in sup norm. On the contrary, the approximation rate for multivariate monotonic functions depends on p. In Section 5, we outline several other examples of function sets F and G for which the general lower bound (Theorem 1) can also be easily applied. Finally, some proofs are postponed to the supplement, while some details on other existing lower bound proof strategies are provided in the supplement, in Appendix C.

Main definitions and notation. We provide below some definitions and notation that will be used throughout the paper. We denote the set of positive integers $\{1, 2, \ldots\}$ by $\mathbb{N}^*$ and let $\mathbb{N} := \mathbb{N}^* \cup \{0\}$. All sets considered in this paper will be assumed to be nonempty, and *measurable set* will be used to denote a set $\mathcal{X}$ (implicitly) endowed with a σ-algebra.

Let $p \in [1, +\infty]$ and $\mathcal{X}$ be any measurable set endowed with a probability measure μ. For any measurable function $f : \mathcal{X} \to \mathbb{R}$, the $L^p(\mu)$ norm of f is defined by $\|f\|_{L^p(\mu)} = \left(\int_{\mathcal{X}} |f(x)|^p d\mu(x) \right)^{1/p}$ (possibly infinite) if $p < +\infty$, and $\|f\|_{L^\infty(\mu)} = \operatorname{ess\,sup}_{x \in \mathcal{X}} |f(x)|$. We will write λ for the Lebesgue measure on $[0, 1]^d$.

For any $\varepsilon > 0$, two functions f_1, f_2 are said to be ε-*distant* in $\| \cdot \|$ if $\|f_1 - f_2\| > \varepsilon$. Let F be a set of functions from $\mathcal{X}$ to $\mathbb{R}$. A set $\{f_1, \ldots, f_N\} \subset F$ is said to be an ε-packing of F in $\| \cdot \|$ (or just an ε-packing for short) if for any $i \neq j \in \{1, \ldots, N\}$, f_i and f_j are ε-distant in $\| \cdot \|$. The ε-packing number $M(\varepsilon, F, \| \cdot \|)$ is the largest cardinality of ε-packings (possibly infinite).

For $\gamma \geq 0$, we say that a set $S = \{x_1 \ldots, x_N\} \subset \mathcal{X}$ is γ-*shattered* by F if there exists $r : S \to \mathbb{R}$ such that for any $E \subset S$, there exists $f \in F$ satisfying for all $i = 1, \ldots, N$, $f(x_i) > r(x_i) + \gamma$ if $x_i \in E$, and $f(x_i) < r(x_i) - \gamma$ if $x_i \notin E$. If this statement is true when we replace γ by 0, we say that F *pseudo shatters* S. The γ-*fat-shattering dimension*, denoted by $\mathrm{fat}_\gamma(F)$, is defined as the largest number $N \leq +\infty$ such that there exists $S \subset \mathcal{X}$ of cardinality N which is γ-shattered by F. The *pseudo dimension* $\mathrm{Pdim}(G)$ is defined similarly but for sets that are pseudo shattered.

A formal definition of feedforward neural networks is recalled in Appendix A. In short, in this paper, a *feedforward neural network architecture* $\mathcal{A}$ of depth $L \geq 1$ is a directed acyclic graph with $d \geq 1$ input neurons, $L - 1$ hidden layers (if $L \geq 2$), and an output layer with only one neuron. Skip connections are allowed, i.e., there can be connections between non-consecutive layers. Given an activation function $\sigma : \mathbb{R} \to \mathbb{R}$, a feedforward neural network architecture $\mathcal{A}$, and a vector $\mathbf{w} \in \mathbb{R}^W$ of weights assigned to all edges and non-input neurons (linear coefficients and biases), the network computes a function $g_{\mathbf{w}} : \mathbb{R}^d \to \mathbb{R}$ defined by recursively computing affine transformations for each hidden or output neuron, and then applying the activation function σ for hidden neurons only (see Appendix A for more details). Finally, we define $H_{\mathcal{A}} := \{g_{\mathbf{w}} : \mathbf{w} \in \mathbb{R}^W\}$ to be the set of all functions that can be represented by tuning all the weights assigned to the network.

A function $\sigma : \mathbb{R} \to \mathbb{R}$ is *piecewise-polynomial* on $K \geq 2$ pieces, with maximal degree $\nu \in \mathbb{N}$, if there exists a partition $I_1, \ldots, I_K$ of $\mathbb{R}$ into K nonempty intervals, such that σ restricted on each I_j is polynomial with degree at most ν.

2 A general approximation lower bound in $L^p(\mu)$ norm

In this section, we provide our two main results: a general lower bound on the $L^p(\mu)$ approximation error of F by G, i.e., $\sup_{f \in F} \inf_{g \in G} \|f - g\|_{L^p(\mu)}$, and a corollary when G corresponds to a feedforward neural network with a piecewise-polynomial activation function. The weak assumptions on F make the last result applicable to a wide range of cases of interest, as shown in Sections 3–5.

2.1 Main results

Our generic lower bound reads as follows, and is proved in Section 2.2. We follow the conventions $0 \times \log^2(0) = 0$ and $P^{-\frac{1}{\alpha}} \log^{-\frac{2}{\alpha}}(P) = +\infty$ when $P = 1$.

Theorem 1. *Let $1 \le p < +\infty$ and $\mathcal{X}$ be a measurable set endowed with a probability measure μ. Let F, G be two sets of real-valued functions defined on $\mathcal{X}$, such that all functions in F have the same finite range $[a, b]$, and $\mathrm{fat}_\gamma(G) < +\infty$ for all $\gamma > 0$. Then, there exists a constant $c > 0$ depending only on p such that*

$$\sup_{f \in F} \inf_{g \in G} \|f - g\|_{L^p(\mu)} \ge \inf \left\{ \varepsilon > 0 : \log M\big(3\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big) \le c \, \mathrm{fat}_{\frac{\varepsilon}{32}}(G) \log^2\left(\frac{2\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G)}{\varepsilon/(b-a)}\right) \right\}.$$
(2)

In particular, if $\log M\big(\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big) \ge c_0 \varepsilon^{-\alpha}$ for some $c_0, \varepsilon_0, \alpha > 0$ and all $\varepsilon \le \varepsilon_0$, and if $\mathrm{Pdim}(G) < +\infty$, then there exist constants $c_1, \varepsilon_1 > 0$ depending only on $b - a$, p, c_0, ε_0 and α such that

$$\sup_{f \in F} \inf_{g \in G} \|f - g\|_{L^p(\mu)} \ge \min \left\{ \varepsilon_1, \ c_1 \,\mathrm{Pdim}(G)^{-\frac{1}{\alpha}} \log^{-\frac{2}{\alpha}}\big(\mathrm{Pdim}(G)\big) \right\}.$$
(3)

The first lower bound (2) is generic but requires solving an inequation.[1] In (3) we solve this inequation when $\log M\big(\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big)$ grows at least polynomially in $1/\varepsilon$ (which is typical of nonparametric sets) and when G has finite pseudodimension $\mathrm{Pdim}(G)$. Though we will restrict our attention to such cases in all subsequent sections, we stress that the first bound should have broader applications. A first example is when $\mathrm{Pdim}(G) = +\infty$ but $\mathrm{fat}_\gamma(G) < +\infty$ is finite for all $\gamma > 0$ (e.g., for RKHS [Bel18]). The first bound should also be useful to prove (slightly) tighter lower bounds when $\log M\big(\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big)$ has a (slightly) different dependency on $1/\varepsilon$ (e.g., of the order of $\varepsilon^{-\alpha} \log^\beta(1/\varepsilon)$ as when F is the set of all multivariate cumulative distribution functions [BGL07]).

In the rest of the paper, we focus on the important special case when the approximation set G is the set of all real-valued functions that can be represented by tuning the weights of a feed-forward neural network with fixed architecture $\mathcal{A}$ and a piecewise-polynomial activation function. By combining Theorem 1 with known bounds on the pseudo-dimension [BHLM19], we obtain the following corollary, which bounds the approximation error in terms of the number W of weights and the depth L (i.e., the number of hidden and output layers). The proof is postponed to Appendix B.4.

Corollary 1. *Let $1 \le p < +\infty$, $d \ge 1$ and $\mathcal{X}$ be a measurable subset of $\mathbb{R}^d$ endowed with a probability measure μ. Let F be a set of functions from $\mathcal{X}$ to $[a, b]$, $a, b \in \mathbb{R}$, such that $\log M\big(\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big) \ge c_0 \varepsilon^{-\alpha}$ for some $c_0, \varepsilon_0, \alpha > 0$ and all $\varepsilon \le \varepsilon_0$.*

Let $H_\mathcal{A}$ be the set of all real-valued functions on $\mathcal{X}$ that can be represented by a feed-forward network with a fixed architecture $\mathcal{A}$ of depth $L \ge 1$, $W \ge 1$ variable weights and a piecewise-polynomial activation function of maximal degree $\nu \in \mathbb{N}$ on $K \ge 2$ pieces. Then, for $W \ge W_{\min}$,

$$\sup_{f \in F} \inf_{g \in H_\mathcal{A}} \|f - g\|_{L^p(\mu)} \ge \begin{cases} c_1 W^{-\frac{2}{\alpha}} \log^{-\frac{2}{\alpha}}(W) & \text{if } \nu \ge 2\,, \\ c_2 (LW)^{-\frac{1}{\alpha}} \log^{-\frac{3}{\alpha}}(W) & \text{if } \nu = 1\,, \\ c_3 W^{-\frac{1}{\alpha}} \log^{-\frac{3}{\alpha}}(W) & \text{if } \nu = 0\,, \end{cases}$$
(4)

where the constants $W_{\min}, c_1, c_2, c_3 > 0$ are independent from W.

There are equivalent ways to write the above corollary. For example, given a target accuracy $\varepsilon > 0$ and a depth $L \ge 1$, (4) yields a lower bound on the minimum number W of weights that are needed to get $\sup_{f \in F} \inf_{g \in H_\mathcal{A}} \|f - g\|_{L^p(\mu)} \le \varepsilon$. Some earlier approximation results were written this way (e.g., [Yar17, PV18]).

---

[1]Note that any $\varepsilon \ge (b-a)/3$ is a solution to this inequation, since $\log M\big(3\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big) = \log(1) = 0$ (because all functions in F are $[a, b]$-valued). Therefore, the right-hand side of (2) is at most $(b-a)/3$.

2.2 Proof of Theorem 1

In order to prove Theorem 1, we need two inequalities. The first one is straightforward (and appeared within proofs, e.g., in [YZ20]), but formalizes the key idea that if G approximates F with error ε, then G has to be at least as large as F. We use the conventions $\log(+\infty) = +\infty$ and $+\infty \leq +\infty$.

Lemma 1. *Let $p \geq 1$ and $\mathcal{X}$ be a measurable space endowed with a probability measure μ. Let F, G be two sets of real-valued functions defined on $\mathcal{X}$. If $\sup_{f \in F} \inf_{g \in G} \|f - g\|_{L^p(\mu)} < \varepsilon$, then*

$$\log M\big(3\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big) \leq \log M\big(\varepsilon, G, \|\cdot\|_{L^p(\mu)}\big) \ .$$

Proof. Let $P_F = \{f_1, \ldots, f_N\}$ be a 3ε-packing of F, with $N \geq 1$. Let $P_G = \{g_1, \ldots, g_N\}$ be a subset of G such that $\|f_i - g_i\|_{L^p(\mu)} \leq \varepsilon$ for all i. Note that the existence of such a P_G is guaranteed by the assumption $\sup_{f \in F} \inf_{g \in G} \|f - g\|_{L^p(\mu)} < \varepsilon$. Since the f_i's are pairwise 3ε-distant in $L^p(\mu)$, the triangle inequality entails that the g_i's are also at least pairwise ε-distant in $L^p(\mu)$. Therefore, P_G is an ε-packing of G, and the result follows. $\square$

The second inequality is a fundamental probability result due to Mendelson [Men02]. It bounds from above the ε-packing number in $L^p(\mu)$ norm of any uniformly bounded function set in terms of its fat-shattering dimension. Crucially, the inequality holds for finite $p \geq 1$, as opposed to the lower bound strategy of Yarotsky [Yar17, Yar18] (see also [DHP21]), that relates the VC-dimension with the approximation error in sup norm. The next statement is a straightforward generalization of a result of [Men02] initially stated for $[a, b] = [0, 1]$ and for Glivenko-Cantelli classes G (see Appendix B.1 for details).

Proposition 1 ([Men02], Corollary 3.12). *Let G be a set of functions from a measurable set $\mathcal{X}$ to $[a, b]$, $a, b \in \mathbb{R}$, and such that $\mathrm{fat}_\gamma(G) < +\infty$ for all $\gamma > 0$. Then for any $1 \leq p < +\infty$, there exists $c > 0$ depending only on p such that for every probability measure μ on $\mathcal{X}$ and every $\varepsilon > 0$,*

$$\log M\big(\varepsilon, G, \|\cdot\|_{L^p(\mu)}\big) \leq c\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G) \log^2\left(\frac{2(b-a)\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G)}{\varepsilon}\right) \ . \tag{5}$$

Proof (of Theorem 1). **Part 1.** We start by proving (2), using Proposition 1 as a key argument. Since functions in G are not necessarily uniformly bounded, we will apply Proposition 1 to the "clipped version of G". More precisely, for any function $g \in G$, we define its clipping (truncature) to $[a, b]$ as the function $\tilde{g} : \mathcal{X} \to \mathbb{R}$ given by $\tilde{g}(x) = \min(\max(a, g(x)), b)$ for all $x \in \mathcal{X}$. We then set $G_{[a,b]} = \{\tilde{g} : g \in G\}$, which by construction consists of functions that are all $[a, b]$-valued.

Noting that clipping can only help since elements of F are $[a, b]$-valued (see Lemma 4 in the supplement, Appendix B.2), we have

$$\sup_{f \in F} \inf_{g \in G} \|f - g\|_{L^p(\mu)} \geq \sup_{f \in F} \inf_{\tilde{g} \in G_{[a,b]}} \|f - \tilde{g}\|_{L^p(\mu)} \ . \tag{6}$$

Setting $\Delta := \sup_{f \in F} \inf_{\tilde{g} \in G_{[a,b]}} \|f - \tilde{g}\|_{L^p(\mu)}$, we now show that Δ is bounded from below by the right-hand side of (2). To that end, it suffices to show that every $\varepsilon > \Delta$ is a solution to the inequation

$$\log M\big(3\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big) \leq c\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G) \log^2\left(\frac{2(b-a)\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G)}{\varepsilon}\right) \ . \tag{7}$$

The last inequality is true whenever $\varepsilon \geq (b-a)/3$ (see Footnote 1 and note $c\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G) \geq 0$). We only need to prove (7) when $\Delta < \varepsilon < (b-a)/3$. In this case, by definition of Δ and by Lemma 1 applied to $G_{[a,b]}$, we have

$$\begin{aligned}
\log M\big(3\varepsilon, F, \|\cdot\|_{L^p(\mu)}\big) &\leq \log M\big(\varepsilon, G_{[a,b]}, \|\cdot\|_{L^p(\mu)}\big) \\
&\leq c\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G_{[a,b]}) \log^2\left(\frac{2(b-a)\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G_{[a,b]})}{\varepsilon}\right) \\
&\leq c\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G) \log^2\left(\frac{2(b-a)\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G)}{\varepsilon}\right) \ ,
\end{aligned} \tag{8}$$

where the second inequality follows from Proposition 1 (note from Lemma 3 in the supplement, Appendix B.2 that $\mathrm{fat}_\gamma(G_{[a,b]}) \le \mathrm{fat}_{\frac{\varepsilon}{32}}(G)$ for all $\gamma > 0$, which is finite by assumption), and where (8) follows from the next remark. Either $\mathrm{fat}_{\frac{\varepsilon}{32}}(G_{[a,b]}) = 0$, and (8) is true by the convention $0 \times \log^2(0) = 0$ and $c\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G) \ge 0$. Either $\mathrm{fat}_{\frac{\varepsilon}{32}}(G_{[a,b]}) \ge 1$, and (8) follows from $t \mapsto ct\log^2\left(\frac{2(b-a)t}{\varepsilon}\right)$ being non-decreasing on $[\varepsilon/(2(b-a)), +\infty)$ and $\varepsilon/(2(b-a)) \le 1/6 \le 1 \le \mathrm{fat}_{\frac{\varepsilon}{32}}(G_{[a,b]}) \le \mathrm{fat}_{\frac{\varepsilon}{32}}(G)$. To conclude, every $\varepsilon > \Delta$ satisfies (7), which implies that Δ is bounded from below by the right-hand side of (2). Combining with (6) concludes the proof of (2).

Part 2. Set $\varepsilon_1' = \min\left\{\frac{\varepsilon_0}{3}, 2(b-a)\right\}$. We now derive (3) from (2). To that end, setting $P = \mathrm{Pdim}(G)$, we show that every $\varepsilon > 0$ satisfying (7) is such that $\varepsilon \ge \min\{\varepsilon_1, c_1 P^{-\frac{1}{\alpha}} \log^{-\frac{2}{\alpha}}(P)\}$, where $\varepsilon_1 \in (0, \varepsilon_1']$ and $c_1 > 0$ will be defined later. Since the claimed lower bound on ε is true when $\varepsilon \ge \varepsilon_1'$, in the sequel we consider any solution ε to (7) such that $0 < \varepsilon < \varepsilon_1'$ (if such a solution exists).

By the assumption on $\log M\left(u, F, \|\cdot\|_{L^p(\mu)}\right)$ for $u = 3\varepsilon \le \varepsilon_0$, and then using (7), we have, setting $r = 2(b-a)$,

$$c_0(3\varepsilon)^{-\alpha} \le \log M\left(3\varepsilon, F, \|\cdot\|_{L^p(\mu)}\right) \le c\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G)\log^2\left(\frac{r\,\mathrm{fat}_{\frac{\varepsilon}{32}}(G)}{\varepsilon}\right) \le cP\log^2\left(\frac{rP}{\varepsilon}\right) ,$$

where the last inequality is because $t \mapsto ct\log^2\left(\frac{rt}{\varepsilon}\right)$ is non-decreasing on $[\varepsilon/r, +\infty)$, with $\varepsilon/r \le 1$, and $1 \le \mathrm{fat}_{\frac{\varepsilon}{32}}(G) \le \mathrm{Pdim}(G) = P$ (the lower bound of 1 follows from $c_0(3\varepsilon)^{-\alpha} > 0$).

Solving the inequation $c_0(3\varepsilon)^{-\alpha} \le cP\log^2(rP/\varepsilon)$ for ε (see Appendix B.3 for details), we get

$$\varepsilon \ge \min\left\{\varepsilon_1'', c_1 P^{-\frac{1}{\alpha}} \log^{-\frac{2}{\alpha}} P\right\} , \tag{9}$$

for some constants $\varepsilon_1'', c_1 > 0$ depending only on $p, c_0, b - a$ and α. Setting $\varepsilon_1 = \min\{\varepsilon_1'', \varepsilon_1'\}$ and noting that ε_1' only depends on ε_0 and $b - a$, we conclude the proof. $\qquad\square$

3 Approximation of Hölder balls by feed-forward neural networks

In this section, we apply Corollary 1 to establish nearly-tight lower bounds for the approximation of unit Hölder balls by feed-forward neural networks. Our main result is Proposition 3, which solves an open question by [DHP21].

Throughout the section, for any $s > 0$, we denote by n and α the unique members of the decomposition $s = n + \alpha$ such that $n \in \mathbb{N}$ and $0 < \alpha \le 1$.

For a set $\mathcal{X} \subset \mathbb{R}^d$, we follow [YZ20] and define the Hölder space $\mathcal{C}^{n,\alpha}(\mathcal{X})$ as the space of n times continuously differentiable functions with finite norm

$$\|f\|_{\mathcal{C}^{n,\alpha}} = \max\left\{\max_{\mathbf{n}:|\mathbf{n}|\le n}\|D^{\mathbf{n}}f\|_\infty, \max_{\mathbf{n}:|\mathbf{n}|=n}\sup_{x\ne y}\frac{|D^{\mathbf{n}}f(x) - D^{\mathbf{n}}f(y)|}{\|x-y\|_2^\alpha}\right\} ,$$

where, for $\mathbf{n} = (n_1, \cdots, n_d) \in \mathbb{N}^d$, $D^{\mathbf{n}}f = \left(\frac{\partial}{\partial x_1}\right)^{n_1}\cdots\left(\frac{\partial}{\partial x_d}\right)^{n_d}f$ denotes the $|\mathbf{n}|$-order partial derivative of f. We denote

$$F_{s,d} = \{f \in \mathcal{C}^{n,\alpha}([0,1]^d) : \|f\|_{\mathcal{C}^{n,\alpha}} \le 1\}.$$

Let λ denote the Lebesgue measure over $[0,1]^d$. In this section, we provide nearly matching upper and lower bounds for the $L^p(\lambda)$ approximation error of elements of $F_{s,d}$ by feed-forward ReLU neural networks. The bounds are expressed in terms of the number of weights of the network.

3.1 Known bounds on the sup norm approximation error

[YZ20] gives matching (up to a certain constant) lower and upper bounds of the sup norm approximation error of the elements of $F_{s,d}$ by feed-forward ReLU neural networks.

Proposition 2 ([YZ20]). *Let $d \in \mathbb{N}^*$, $s > 0$, $\gamma \in \left(\frac{s}{d}, \frac{2s}{d}\right]$. Consider $n \in \mathbb{N}$ and $\alpha \in (0, 1]$ such that $s = n + \alpha$.*

There exist positive constants $W_{\min}$ and c_1, depending only on d and n, such that for any integer $W \geq W_{\min}$, there exists a feed-forward ReLU neural network architecture $\mathcal{A}$ with $L = O(W^{\gamma \frac{d}{s} - 1})$ layers and W weights such that

$$\sup_{f \in F_{s,d}} \inf_{g \in H_{\mathcal{A}}} \|f - g\|_\infty \leq c_1 W^{-\gamma}. \tag{10}$$

In the meantime, there exists a constant c_2 depending only on d and n such that for any feed-forward ReLU neural network of architecture $\mathcal{A}$ with W weights and $L = o(W^{\gamma \frac{d}{s} - 1} / \log W)$ layers

$$\sup_{f \in F_{s,d}} \inf_{g \in H_{\mathcal{A}}} \|f - g\|_\infty \geq c_2 W^{-\gamma}. \tag{11}$$

It is worth stressing that, for any probability measure μ on $[0, 1]^d$, the upper bound (10) is automatically generalized to any smaller $L^p(\mu)$ norm, when $1 \leq p < +\infty$. However, the lower bound (11) does not necessarily apply when $\| \cdot \|_\infty$ is replaced with $\| \cdot \|_{L^p(\mu)}$, $1 \leq p < +\infty$. The lower bound of the next subsection shows that, in this setting, approximation in $L^p(\lambda)$ norm is not easier than in sup norm, solving an open question of DeVore et al. [DHP21].

3.2 Nearly-matching lower bounds of the $L^p(\lambda)$ approximation error

We first state a lower bound on the packing number of $F_{s,d}$, which is rather classical though hard to find in this specific form (see [BS67] for the L^∞ norm, or [ET96] for other Sobolev-type norms). For the sake of completeness, we give a proof of Lemma 2 in the supplement, Appendix D.1.

Lemma 2. *Let $s > 0$, $d \in \mathbb{N}^*$ and $1 \leq p < \infty$. There exist constants $\varepsilon_0, c_0 > 0$ such that for any $0 < \varepsilon \leq \varepsilon_0$,*

$$\log M\left(\varepsilon, F_{s,d}, \| \cdot \|_{L^p(\lambda)}\right) \geq c_0 \varepsilon^{-\frac{d}{s}}. \tag{12}$$

Given Lemma 2, we can use Corollary 1 to establish the next proposition and obtain the lower bound of the $L^p(\lambda)$ approximation error.

Proposition 3. *Let $d \in \mathbb{N}^*$, $s > 0$, $\gamma \in \left(\frac{s}{d}, \frac{2s}{d}\right]$ and $1 \leq p < +\infty$. Let ε_0, c_0 be defined as in Lemma 2. Consider $n \in \mathbb{N}$ and $\alpha \in (0, 1]$ such that $s = n + \alpha$.*

Let $\sigma : \mathbb{R} \to \mathbb{R}$ be a piecewise-affine function. There exist positive constants $c_1, W_{\min}$ depending only on d, p, c_0, ε_0 and σ such that for any architecture $\mathcal{A}$ of depth $1 \leq L \leq cW^{\gamma \frac{d}{s} - 1}$ (where c is a constant), $W \geq W_{\min}$ variable weights and activation σ, the set $H_{\mathcal{A}}$ satisfies

$$\sup_{f \in F_{s,d}} \inf_{g \in H_{\mathcal{A}}} \|f - g\|_{L^p(\lambda)} \geq c_1 W^{-\gamma} \log^{-\frac{3s}{d}}(W). \tag{13}$$

Note that, since ReLU is a piecewise-affine function, we obtain a lower bound which matches, up to logarithmic factors, the upper bound presented in the previous subsection.

Proof. From Lemma 2, there exists a constant $c_0 > 0$ such that $\log M\left(\varepsilon, \mathcal{M}_d, \| \cdot \|_{L^p(\lambda)}\right) \geq c_0 \varepsilon^{-\frac{d}{s}}$. Therefore, using Corollary 1 and $L \leq c(W^{\gamma \frac{d}{s} - 1})$, we obtain the result. $\square$

Remark 1 (Comparison with existing proof strategies in sup norm.)**.** *We would like to highlight a key difference between the proof of Proposition 3 and the lower bound proof strategies of [Yar17, Yar18, YZ20, SYZ22] that are specific to the sup norm. Their overall argument is roughly the following: if G can approximate any $f \in F$ in sup norm at accuracy $\varepsilon > 0$, since F contains many "oscillating" functions with oscillation amplitude roughly ε, then so must be the case for G (the sup norm is key here: **all** oscillations of any $f \in F$ are well approximated). Therefore, a small ε implies a large $\mathrm{VCdim}(G)$, which by contrapositive enables to lower bound the approximation error (1) with a*

decreasing function of $\mathrm{VCdim}(G)$, *and therefore as a function of L and W. In contrast, in the proof of Theorem 1, the key probability result of Mendelson (Proposition 1) enables us to show that, even if the oscillations of any $f \in F$ are only well approximated* **on average** *(in $L^p(\mu)$ norm) by G, then* $\mathrm{Pdim}(G)$ *must be large when ε is small. The conclusion is then the same: the approximation error in $L^p(\mu)$ norm can be lower bounded as a function of $\mathrm{Pdim}(G)$, and therefore in terms of L, W. This solves the question of DeVore et al. [DHP21] mentioned in the introduction, showing in particular that VC dimension theory can (surprisingly) be useful to prove L^p approximation lower bounds.*

4 Approximation of monotonic functions by feed-forward neural networks

In this section, we consider the problem of approximating the set $\mathcal{M}^d$ of all non-decreasing functions from $[0,1]^d$ to $[0,1]$. These are functions $f : [0,1]^d \to [0,1]$ that are non-decreasing along any line parallel to an axis, i.e., such that, for all $x, y \in [0,1]^d$,

$$x_i \leq y_i, \;\; i = 1, \ldots, d \implies f(x) \leq f(y) \,.$$

Next we focus on the approximation of $\mathcal{M}^d$ with Heaviside feed-forward neural networks. After proving an impossibility result for the sup norm, we show that the weaker goal of approximating $\mathcal{M}^d$ in L^p norm is feasible, and derive nearly matching lower and upper bounds. Interestingly, the approximation rates depend on $p \geq 1$, which is in sharp contrast with the case of Hölder balls, that are not easier to approximate in L^p norm than in sup norm (see Section 3).

4.1 Warmup: an impossibility result in sup norm

We start this section by showing that approximating monotonic functions of $d \geq 2$ variables in the sup norm is impossible with Heaviside neural networks.

Proposition 4. *For any neural network architecture $\mathcal{A}$ with the Heaviside activation, the set $H_{\mathcal{A}}$ satisfies*

$$\sup_{f \in \mathcal{M}^d} \inf_{g \in H_{\mathcal{A}}} \|f - g\|_{\infty} \geq \frac{1}{2}.$$

The proof of Proposition 4 is postponed to the supplement, Appendix E.2. We show a slightly stronger result, by exhibiting a single function $f \in \mathcal{M}^d$ such that the lower bound of $\frac{1}{2}$ holds simultaneously for all network architectures.

Next we study the approximation of $\mathcal{M}^d$ in $L^p(\lambda)$ norm.

4.2 Lower bound in $L^p(\lambda)$ norm

We start by proving a lower bound, as a direct consequence of Corollary 1 and a lower bound on the packing number due to [GW07].

Proposition 5. *Let $1 \leq p < +\infty$, $d \geq 1$, and let $\alpha = \max\{d, (d-1)p\}$. Let $\sigma : \mathbb{R} \to \mathbb{R}$ be a piecewise-polynomial function having maximal degree $\nu \in \mathbb{N}$. There exist positive constants $c_1, c_2, c_3, W_{\min}$ depending only on d, p, and σ such that for any architecture $\mathcal{A}$ of depth $L \geq 1$, $W \geq W_{\min}$ variable weights and activation σ, the set $H_{\mathcal{A}}$ satisfies*

$$\sup_{f \in \mathcal{M}^d} \inf_{g \in H_{\mathcal{A}}} \|f - g\|_{L^p(\lambda)} \geq \begin{cases} c_1 W^{-\frac{2}{\alpha}} \log^{-\frac{2}{\alpha}}(W) & \text{if } \nu \geq 2 \,, \\ c_2 (LW)^{-\frac{1}{\alpha}} \log^{-\frac{3}{\alpha}}(W) & \text{if } \nu = 1 \,, \\ c_3 W^{-\frac{1}{\alpha}} \log^{-\frac{3}{\alpha}}(W) & \text{if } \nu = 0 \,. \end{cases} \tag{14}$$

Proof. From [GW07], there exist constants $\varepsilon_0, c_0 > 0$ such that for $\varepsilon \leq \varepsilon_0$, $\log M(\varepsilon, \mathcal{M}_d, \|\cdot\|_{L^p}) \geq c_0 \varepsilon^{-\alpha}$. Using Corollary 1, we obtain the result. $\qquad\square$

4.3 Nearly-matching upper bound in $L^p(\lambda)$ norm

To the best of our knowledge, there does not exist any upper-bound of the $L^p(\lambda)$ approximation error of $\mathcal{M}^d$ with feed-forward neural networks. Checking that all the lower-bounds of Proposition 5 are tight is out of the scope of this paper and we leave it for future research[2]. However, we establish in the next proposition upper-bounds of the $L^p(\lambda)$ approximation error of $\mathcal{M}^d$ with feed-forward neural networks with the Heaviside activation function. This shows that, for the $L^p(\lambda)$ approximation error, the lower-bound obtained in (14), for $\nu = 0$, is tight up to logarithmic factors. The next proposition follows by reinterpreting a metric entropy upper bound of [GW07] in terms of Heaviside neural networks. The proof is postponed to Appendix E.1 in the supplement.

Proposition 6. *Let* $1 \le p < +\infty$, $d \in \mathbb{N} \setminus \{0, 1\}$ *and let* $\alpha = \max\{d, (d-1)p\}$. *There exist positive constants* $W_{\min}$ *and* c, *depending only on* d *and* p, *such that for any integer* $W \ge W_{\min}$, *there exists a feed-forward architecture* $\mathcal{A}$ *with two hidden layers,* W *weights and the Heaviside activation function such that the set* $H_\mathcal{A}$ *satisfies*

$$\sup_{f \in \mathcal{M}^d} \inf_{g \in H_\mathcal{A}} \|f - g\|_{L^p(\lambda)} \le \left\{ \begin{array}{ll} cW^{-\frac{1}{\alpha}} & \text{if } p(d-1) \ne d, \\ cW^{-\frac{1}{d}} \log(W) & \text{if } p(d-1) = d, \end{array} \right. \tag{15}$$

5 Conclusion and other possible applications

We proved a general lower bound on the approximation error of F by G in $L^p(\mu)$ norm (Theorem 1), in terms of generic properties of F and G (packing number of F, range of F, fat-shattering dimension of G). The proof relies on VC dimension theory as in the sup norm case, but uses an additional key probabilistic argument due to Mendelson ([Men02], see Proposition 1), solving a question raised by DeVore et al. [DHP21].

In Sections 3 and 4 we detailed two applications, where Corollary 1 yields nearly optimal approximation lower bounds in L^p norm, and which correspond to two examples where the approximation rate may depend or not depend on p.

Theorem 1 and Corollary 1 can be used to derive approximation lower bounds for many other cases. Corollary 1 only requires a (tight) lower bound on the packing number of F, for which approximation theory provides several examples. For instance, for the *Barron space* introduced in [Bar93], Petersen and Voigtlaender [PV21] showed a tight lower bound on the log packing number in $L^p(\lambda, [0,1]^d)$ norm, of order $\varepsilon^{-2d/(d+2)}$. Applying Corollary 1, this yields an approximation lower bound of $(LW)^{-\left(\frac{1}{2}+\frac{1}{d}\right)} \log^{-3\left(\frac{1}{2}+\frac{1}{d}\right)}(W)$ for ReLU networks (see Appendix F in the supplement for details). Other examples of sets F for which tight lower bounds on the packing number (or metric entropy) are available include: multivariate cumulative distribution functions [BGL07], multivariate convex functions [GS13], and functions with other shape constraints [GJ14].

Theorem 1 can also be applied to other approximating sets G, beyond classical feedforward neural networks, as soon as a (tight) upper bound on the fat-shattering dimension of G is available. For example, upper bounds were derived by [WS22] on the VC dimension of partially quantized networks, while [Bel18] derived bounds on the fat-shattering dimension of some RKHS. Investigating such applications and whether the obtained approximation lower bounds are rate-optimal is a natural research direction for the future.

References

[AB99]   Martin Anthony and Peter L. Bartlett. *Neural network learning: theoretical foundations*. Cambridge University Press, Cambridge, 1999.

---

[2]Obtaining an upper-bound for ReLU networks seems challenging. For example, the bit extraction technique used in [Yar18] to find a sharp upper bound heavily relies on the local smoothness assumption of the function to approximate, which is not satisfied in general for monotonic functions.

[Bar93]   Andrew Barron. Universal approximation bounds for superpositions of a sigmoidal function. *Information Theory, IEEE Transactions on*, 39:930 – 945, 06 1993.

[Bel18]   Mikhail Belkin. Approximation beats concentration? an approximation view on inference with smooth radial kernels. In Sébastien Bubeck, Vianney Perchet, and Philippe Rigollet, editors, *Proceedings of the 31st Conference On Learning Theory*, volume 75 of *Proceedings of Machine Learning Research*, pages 1348–1361. PMLR, 06–09 Jul 2018.

[BGL07]   Ron Blei, Fuchang Gao, and Wenbo V. Li. Metric entropy of high dimensional distributions. *Proceedings of the American Mathematical Society*, 135(12):4009–4018, 2007.

[BHLM19] Peter L. Bartlett, Nick Harvey, Christopher Liaw, and Abbas Mehrabian. Nearly-tight vc-dimension and pseudodimension bounds for piecewise linear neural networks. *Journal of Machine Learning Research*, 20(63):1–17, 2019.

[BS67]   M Š Birman and M Z Solomjak. Piecewise-polynomial approximation of functions of the classes w_p^α. 2(3):295–317, apr 1967.

[Cyb89]   G. Cybenko. Approximation by superpositions of a sigmoidal function. *Math. Control Signal Systems*, 2:303–314, 1989.

[DHM89]   Ronald A DeVore, Ralph Howard, and Charles Micchelli. Optimal nonlinear approximation. *Manuscripta mathematica*, 63(4):469–478, 1989.

[DHP21]   Ronald DeVore, Boris Hanin, and Guergana Petrova. Neural network approximation. *Acta Numerica*, 30:327–444, 2021.

[ET96]   D. E. Edmunds and H. Triebel. *Function Spaces, Entropy Numbers, Differential Operators*. Cambridge Tracts in Mathematics. Cambridge University Press, 1996.

[GJ04]   Paul W. Goldberg and Mark Jerrum. Bounding the vapnik-chervonenkis dimension of concept classes parameterized by real numbers. *Machine Learning*, 18:131–148, 2004.

[GJ14]   Piet Groeneboom and Geurt Jongbloed. *Nonparametric Estimation under Shape Constraints: Estimators, Algorithms and Asymptotics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2014.

[GR21]   Ingo Gühring and Mones Raslan. Approximation rates for neural networks with encodable weights in smoothness spaces. *Neural Networks*, 134:107–130, 2021.

[GS13]   Adityanand Guntuboyina and Bodhisattva Sen. Covering numbers for convex functions. *IEEE Transactions on Information Theory*, 59(4):1957–1965, 2013.

[GW07]   Fuchang Gao and Jon A. Wellner. Entropy estimate for high-dimensional monotonic functions. *Journal of Multivariate Analysis*, 98(9):1751–1764, 2007.

[Hor91]   Kurt Hornik. Approximation capabilities of multilayer feedforward networks. *Neural Networks*, 4(2):251–257, 1991.

[KB17]   Jason M. Klusowski and Andrew R. Barron. Minimax lower bounds for ridge combinations including neural nets. In *2017 IEEE International Symposium on Information Theory (ISIT)*, pages 1376–1380, 2017.

[KL20]   Patrick Kidger and Terry Lyons. Universal Approximation with Deep Narrow Networks. In Jacob Abernethy and Shivani Agarwal, editors, *Proceedings of Thirty Third Conference on Learning Theory*, volume 125 of *Proceedings of Machine Learning Research*, pages 2306–2327. PMLR, 09–12 Jul 2020.

[LLPS93]   Moshe Leshno, Vladimir Ya. Lin, Allan Pinkus, and Shimon Schocken. Multilayer feedforward networks with a nonpolynomial activation function can approximate any function. *Neural Networks*, 6(6):861–867, 1993.

[Mai99]   V.E Maiorov. On best approximation by ridge functions. *Journal of Approximation Theory*, 99(1):68–94, 1999.

[Men02]   S. Mendelson. Rademacher averages and phase transitions in glivenko-cantelli classes. *IEEE Transactions on Information Theory*, 48, 2002.

[MMR99]   V. Maiorov, R. Meir, and Joel Ratsaby. On the approximation of functional classes equipped with a uniform measure using ridge functions. *Journal of Approximation Theory*, 99:95–111, 1999.

[MP99]   Vitaly Maiorov and Allan Pinkus. Lower bounds for approximation by mlp neural networks. *Neurocomputing*, 25(1):81–91, 1999.

[PV18]   Philipp Petersen and Felix Voigtlaender. Optimal approximation of piecewise smooth functions using deep relu neural networks. *Neural Networks*, 108:296–330, 2018.

[PV21]   Philipp Petersen and Felix Voigtlaender. Optimal learning of high-dimensional classification problems using deep neural networks, 2021.

[SX21]   Jonathan W. Siegel and Jinchao Xu. Sharp bounds on the approximation rates, metric entropy, and n-widths of shallow neural networks. 2021.

[SYZ22]   Zuowei Shen, Haizhao Yang, and Shijun Zhang. Optimal approximation rate of relu networks in terms of width and depth. *Journal de Mathématiques Pures et Appliquées*, 157:101–135, 2022.

[VP19]   Felix Voigtländer and Philipp Petersen. Approximation in $l^p(\mu)$ with deep relu neural networks, 04 2019.

[WS22]   Yutong Wang and Clayton D. Scott. Vc dimension of partially quantized neural networks in the overparametrized regime. In *Proceedings of ICLR 2022*, 2022.

[Yar17]   Dmitry Yarotsky. Error bounds for approximations with deep relu networks. *Neural Networks*, 94:103–114, 2017.

[Yar18]   Dmitry Yarotsky. Optimal approximation of continuous functions by very deep relu networks. *arXiv preprint arXiv:1802.03620*, 2018.

[Yu97]   Bin Yu. Assouad, fano, and le cam. 1997.

[YZ20]   Dmitry Yarotsky and Anton Zhevnerchuk. The phase diagram of approximation rates for deep neural networks. *Advances in neural information processing systems*, 33:13005–13015, 2020.

[ZBH⁺21] Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning (still) requires rethinking generalization. *Communications of the ACM*, 64(3):107–115, 2021.