So3krates - Self-attention for higher-order geometric interactions on arbitrary length-scales

Abstract

The application of machine learning methods in quantum chemistry has enabled the study of numerous chemical phenomena, which are computationally intractable with traditional *ab-initio* methods. However, some quantum mechanical properties of molecules and materials depend on non-local electronic effects, which are often neglected due to the difficulty of modeling them efficiently. This work proposes a modified attention mechanism adapted to the underlying physics, which allows to recover the relevant non-local effects. Namely, we introduce *spherical harmonic coordinates* (SPHCs) to reflect higher-order geometric information for each atom in a molecule, enabling a non-local formulation of attention in the SPHC space. Our proposed model SO3KRATES – a self-attention based message passing neural network – uncouples geometric information from atomic features, making them independently amenable to attention mechanisms. We show that in contrast to other published methods, SO3KRATES is able to describe non-local quantum mechanical effects over arbitrary length scales. Further, we find evidence that the inclusion of higher-order geometric correlations increases data efficiency and improves generalization. SO3KRATES matches or exceeds state-of-the-art performance on popular benchmarks, notably, requiring a significantly lower number of parameters (0.25–0.4x) while at the same time giving a substantial speedup (6–14x for training and 2–11x for inference) compared to other models.

1   Introduction

Atomistic simulations use long time-scale molecular dynamics (MD) trajectories to predict macroscopic properties that arise from interactions on the microscopic scale [1–3]. Their predictive reliability is determined by the accuracy of the underlying *force field* (FF), which needs to be queried at every time step. This quickly becomes a computational bottleneck if the forces are determined from first principles, which may be required for accurate results. To that end, machine learning FFs (MLFFs) offer a computationally more efficient, yet accurate empirical alternative to expensive *ab-initio* methods [2, 4–24].

In recent years, *Geometric Deep Learning* has become a popular design paradigm, which exploits relevant symmetry groups of the underlying learning problem by incorporating a *geometric prior* [12, 25, 26]. This effectively restricts the learnable functions of the model to a subspace with a meaningful inductive bias. Prominent examples for such models are e.g. convolutional neural networks (CNNs) [27], which are equivariant w. r. t. the group of translations, or graph neural networks (GNNs) [28], which are invariant w. r. t. node permutation.

For molecular property prediction, it has been shown that equivariance w. r. t. the 3D rotation group SO(3) greatly improves data efficiency and accuracy of the learned FFs [29–32]. To achieve equivariance, architectures either rely on feature expansions in terms of spherical harmonics (SH) [33]

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

Figure 1: **(a)** Design principle of our proposed augmented message-passing scheme, where features **f** and spherical harmonics coordinates (SPHCs) χ are updated separately using information from both **f** and χ. **(b)** Overview of the SO3KRATES architecture. **(c)** Comparison of prediction accuracy and model size of different architectures. **(d)** Comparison of training and inference times of different architectures. **(e)** Illustration of learned SPHCs at different layers of the SO3KRATES architecture. **(f)** Low-dimensional projection of atomic SPHCs, showing that atoms far apart in Euclidean space can be mapped close together in SPHC space. **(g)** In contrast to local models, SO3KRATES is able to learn the long-range correlations between hydrogen rotors in cumulene and reproduces the *ab-initio* ground truth faithfully.

37 or explicitly include (dihedral) angles [29, 34]. While the latter scales quadratically (cubically) in
38 the number of neighboring atoms and has been shown to be geometrically incomplete [35], the
39 calculation of spherical harmonics scales only linear in the number of neighboring atoms, which
40 makes them a fast and accurate alternative [30, 32, 36].

41 However, current higher-order geometric representations based on SHs usually result in expensive
42 transformations, since an individual feature channel per SH degree (and order) is required. As a result,
43 going to higher degrees is computationally expensive and comes at the price of increasing complexity,
44 resulting in state-of-the-art (SOTA) models with millions of parameters [30, 32, 34]. However, in
45 order to be applicable to large molecular structures, models are required to be both efficient and
46 accurate on *all* length scales.

47 Non-local electronic effects have been outlined as one of the major challenges for a new generation of
48 MLFFs [21]. They result in non-local, higher-order geometric relations between atoms. Most current
49 architectures implicitly assume locality of interactions (expressed through a local neighborhood),
50 which prohibits an efficient description of all relevant atomic interactions at larger scales. Simply
51 increasing the cutoff radius used to determine local neighborhoods is not an adequate solution, since
52 it only shifts the problem to larger length scales [30].

---

* Reference times were taken from [34]. As our own timings were measured on a different GPU, we decreased the reported times according to speedup-factors reported in [37]. For full details, see appendix A.8.

In this work, we propose *spherical harmonic coordinates* (SPHCs), which encode higher-order geometric information for each node in a molecular graph (Fig. 1e). This is in stark contrast to current approaches, which consider molecules as three-dimensional point clouds with learned features and fixed atomic coordinates: we propose to make the SPHCs themselves a *learned* quantity. Through localization in the space of SPHCs (Fig. 1f), models are able to efficiently describe electronic effects that are non-local in three-dimensional Euclidean space (Fig. 1g).

We then present SO3KRATES (Fig. 1b), a self-attention based message-passing neural network (MPNN), which decouples atomic features and SPHCs and updates them individually (Fig. 1a). This resembles ideas from equivariant graph neural networks [25], but allows to go to arbitrarily high geometric orders. The separation of higher-order geometric and feature information allows to overcome the parametric and computational complexity usually encountered in models with higher-order geometric representations, since we only require a single feature channel (instead of one per SH degree and order). Thus, SO3KRATES resembles some early architectures like SCHNET [10] or PHYSNET [19] in parametric simplicity. We further show that SO3KRATES outperforms the popular sGDML [38] kernel model by a large margin in the low-data regime, a domain which has so far been considered to be dominated by kernel machines [21]. Numerical evidence suggests that the data efficiency of SO3KRATES is directly related to the maximal degree of geometric information encoded in the SPHCs. We then apply SO3KRATES to the well-established MD17 benchmark and show that our model achieves SOTA results, despite is light-weight structure and having only 0.25–0.4x the number of parameters of competitive architectures (Fig. 1c), while achieving speedups of 6–14x and 2–11x for training and inference, respectively (Fig. 1d).

Although we focus on quantum chemistry applications in this work, the developed methods are also applicable to other fields where long-ranged correlations in three-dimensional data are relevant. For example, models based on SPHCs may also be applicable to tasks like 3D shape classification or computer vision.

2 Preliminaries and Related Work

In the following, we review the most important concepts our method is based upon and relate it to prior work.

Message Passing Neural Networks MPNNs [14] carry over many of the benefits of convolutions to unstructured domains and have thus become one of the most promising approaches for the description of molecular properties. Their general working principle relies on the repeated iteration of message passing (MP) steps, which can be phrased as follows [14, 25]

$$\mathbf{m}_{ij} = m(\mathbf{f}_i, \mathbf{f}_j, \boldsymbol{r}_{ij}) \tag{1}$$

$$\mathbf{m}_i = \sum_{j \in \mathcal{N}(i)} \mathbf{m}_{ij} \tag{2}$$

$$\mathbf{f}_i' = u(\mathbf{f}_i, \mathbf{m}_i) . \tag{3}$$

Here, $\mathbf{m}_{ij}$ is the message between atoms i and j computed with the message function $m(\cdot)$, $\mathbf{m}_i$ is the aggregation of all messages in the neighborhood $\mathcal{N}(i)$ of atom i, and $u(\cdot)$ is an update function returning updated features $\mathbf{f}_i'$ based on the current features $\mathbf{f}_i$ and message $\mathbf{m}_i$. The neighborhood $\mathcal{N}(i)$ consists of all atoms which lie within a given cutoff radius around the atomic position $\boldsymbol{r}_i$, which ensures linear scaling in the number of atoms n. While earlier variants parametrized messages only in terms of inter-atomic distances [13, 19], more recent approaches also take higher-order geometric information into account [25, 29, 31, 32, 39, 40].

Molecules as Point Clouds A molecule can be considered as a point cloud of n atoms $\mathcal{P}_{3D}(\mathcal{R}, \mathcal{F})$, where $\mathcal{R} = (\boldsymbol{r}_1, ... \boldsymbol{r}_n)$ denotes the set of atomic positions $\boldsymbol{r}_i \in \mathbb{R}^3$ and $\mathcal{F} = (\mathbf{f}_1, \ldots, \mathbf{f}_n)$ is the set of rotationally invariant atomic descriptors, or features, $\mathbf{f}_i \in \mathbb{R}^F$. We write the distance vector pointing from the position of atom i to the position of atom j as $\boldsymbol{r}_{ij} = \boldsymbol{r}_j - \boldsymbol{r}_i$, the distance as $r_{ij} = \|\boldsymbol{r}_{ij}\|_2$ and the normalized distance vector as $\hat{\boldsymbol{r}}_{ij} = \boldsymbol{r}_{ij}/r_{ij}$. Given the point cloud, a density over Euclidean space assigning a vector value to each point $\boldsymbol{r}$ can be constructed as

$$\boldsymbol{\rho}(\boldsymbol{r}) = \sum_{i=1}^{n} \delta(\|\boldsymbol{r}_i - \boldsymbol{r}\|_2) \cdot \mathbf{f}_i , \tag{4}$$

where δ is the Dirac delta function. It can be shown that applying a convolutional filter on $\boldsymbol{\rho}(\boldsymbol{r})$ resembles the update steps used in MPNNs [41].

Equivariance Given a set of transformations that act on a vector space $\mathbb{A}$ as $S_g : \mathbb{A} \mapsto \mathbb{A}$ to which we associate an abstract group G, a function $f : \mathbb{A} \mapsto \mathbb{B}$ is said to be equivariant w.r.t. G if

$$f(S_g x) = T_g f(x) \,, \tag{5}$$

where $T_g : \mathbb{B} \mapsto \mathbb{B}$ is an equivalent transformation on the output space [25]. Thus, in order to say that f is equivariant, it must hold that under transformation of the input, the output transforms "in the same way". While equivariance has been a popular concept in signal processing for decades (cf. e.g. [42]), recent years have seen efforts to design group equivariant NNs and kernel methods, since respecting relevant symmetries builds an important inductive bias [12, 43, 44]. Examples are CNNs [27] which are equivariant w.r.t. translation, GNNs [14, 28] which are invariant ($T_g = \mathbb{I}$) w.r.t. permutation, or architectures which are equivariant w.r.t. the $\mathrm{SO}(3)$ group [25, 31, 33, 36, 45]. In this work, we consider the $\mathrm{SO}(3)$ group of rotations, such that $\mathbb{A}$ is the Euclidean space $\mathbb{R}^3$, where the corresponding group actions are given by rotation matrices $R_\theta \in \mathbb{R}^{3\times3}$.

Spherical Harmonics The spherical harmonics are special functions defined on the surface of the sphere $S^2 = \{\hat{\boldsymbol{r}} \in \mathbb{R}^3 : \|\hat{\boldsymbol{r}}\|_2 = 1\}$ and form an orthonormal basis for the irreducible representations (irreps) of $\mathrm{SO}(3)$. In the context of *tensor field networks* [33], they have been introduced as elementary building blocks for $\mathrm{SO}(3)$-equivariant neural networks. The spherical harmonics are commonly denoted as $Y_l^m(\hat{\boldsymbol{r}}) : S^2 \mapsto \mathbb{R}$, where the *degree* l determines all possible values of the *order* $m \in \{-l, \dots, +l\}$. They transform under rotation as

$$Y_l^m(R_\theta \, \hat{\boldsymbol{r}}) = \sum_{m'} D_{mm'}^l(R_\theta) \, Y_l^{m'}(\hat{\boldsymbol{r}}) \,, \tag{6}$$

where $D_{mm'}^l(R_\theta)$ are the entries of the *Wigner-D* matrix $\mathbf{D}^l(R_\theta) \in \mathbb{R}^{(2l+1)\times(2l+1)}$ [46]. Based on the spherical harmonics, we define a vector-valued function $\boldsymbol{Y}^{(l)} : S^2 \mapsto \mathbb{R}^{2l+1}$ for each degree l, with entries Y_l^m for all valid orders m of a given degree l. Since $\boldsymbol{Y}^{(l)}(R_\theta \, \hat{\boldsymbol{r}}) = \mathbf{D}^l(R_\theta)\boldsymbol{Y}^{(l)}(\hat{\boldsymbol{r}})$ (cf. eq. (6)), $\boldsymbol{Y}^{(l)}$ is equivariant w.r.t. $\mathrm{SO}(3)$.

Tensor Product Contractions The irreps $\boldsymbol{Y}^{(l_1)}$ and $\boldsymbol{Y}^{(l_2)}$ can be coupled by computing their tensor product $\boldsymbol{Y}^{(l_1)} \otimes \boldsymbol{Y}^{(l_2)}$, which can equivalently be expressed as a direct sum [33, 47]

$$\boldsymbol{Y}^{(l_1)} \otimes \boldsymbol{Y}^{(l_2)} = \bigoplus_{l_3=|l_1-l_2|}^{l_1+l_2} \overbrace{\widetilde{\boldsymbol{Y}}^{(l_3)}}^{:=\left(\boldsymbol{Y}^{(l_1)} \otimes_{l_3} \boldsymbol{Y}^{(l_2)}\right)} \,, \tag{7}$$

where the entry of order m_3 for the coupled irreps $\widetilde{\boldsymbol{Y}}^{(l_3)}$ is given by

$$\tilde{Y}_{m_3}^{l_3} = \sum_{m_1=-l_1}^{l_1} \sum_{m_2=-l_2}^{l_2} C_{m_1,m_2,m_3}^{l_1,l_2,l_3} Y_{m_1}^{l_1} Y_{m_2}^{l_2} \,, \tag{8}$$

and $C_{m_1,m_2,m_3}^{l_1,l_2,l_3}$ are the so-called *Clebsch-Gordon coefficients*. In the following, we will denote the tensor product of degrees l_1 and l_2 followed by "contraction" to l_3 (meaning the irreps of degree l_3 in the direct sum representation of their tensor product) as $\left(\boldsymbol{Y}^{(l_1)} \otimes_{l_2} \boldsymbol{Y}^{(l_2)}\right)$, which is a mapping of the form $\mathbb{R}^{(2l_1+1)\times(2l_2+1)} \mapsto \mathbb{R}^{2l_3+1}$, since $m_3 \in \{-l_3, \dots l_3\}$.

3 Methods

In the following, we describe the main methodological contributions of this work. We introduce the concept of an adapted point cloud $\mathcal{P}_{3D}(\mathcal{R}, \mathcal{X}, \mathcal{F})$, which incorporates the set of spherical harmonics coordinates (SPHCs) $\mathcal{X} = (\boldsymbol{\chi}_1, \dots, \boldsymbol{\chi}_n)$ (see below) in addition to features $\mathcal{F}$ and Euclidean coordinates $\mathcal{R}$. However, contrary to $\mathcal{R}$, SPHCs $\mathcal{X}$ are refined during the message passing updates. Having SPHCs as part of the molecular point cloud extends the idea of current MPNNs, which learn message functions on $\mathcal{R}$, only. Instead, we learn a message function m (cf. eq. (1)) on both, the (fixed) atomic coordinates $\mathcal{R}$ as well as on the SPHCs $\mathcal{X}$. This adapted message-passing scheme allows to learn non-local geometric corrections. Based on these design principles, we propose the SO3KRATES architecture.

Initialization Feature vectors are initialized from the atomic numbers $z_i \in \mathbb{N}$ (denoting which chemical element an atom belongs to) by an embedding map

$$\mathbf{f}_i = f_{\text{emb}}(z_i), \tag{9}$$

where $f_{\text{emb}} : \mathbb{N} \mapsto \mathbb{R}^F$. We define SPHCs $\boldsymbol{\chi}$ as the concatenation of degrees $\mathcal{L} \coloneqq \{l_{\min}, \ldots, l_{\max}\}$

$$\boldsymbol{\chi} = [\ \underbrace{\boldsymbol{\chi}^{(l_{\min})}}_{\in \mathbb{R}^{2l_{\min}+1}}, \ldots, \underbrace{\boldsymbol{\chi}^{(l_{\max})}}_{\in \mathbb{R}^{2l_{\max}+1}}\ ] \in \mathbb{R}^{(l_{\max}-l_{\min}+1)^2}, \tag{10}$$

such that their transformation under rotation can be expressed in terms of concatenated Wigner-D matrices (see appendix A.1). The short-hand $\boldsymbol{\chi}^{(l)} \in \mathbb{R}^{2l+1}$ refers to the subset of SPHCs with degree l. They are initialized as

$$\boldsymbol{\chi}_i^{(l)} = \frac{1}{\mathcal{C}_i} \sum_{j \in \mathcal{N}(i)} \phi_{r_{\text{cut}}}(r_{ij}) \cdot \mathbf{Y}^{(l)}(\hat{\boldsymbol{r}}_{ij}), \tag{11}$$

where $\mathcal{C}_i = \sum_{j \in \mathcal{N}(i)} \phi_{r_{\text{cut}}}(r_{ij})$, $\phi_{r_{\text{cut}}} : \mathbb{R} \mapsto \mathbb{R}$ is the cosine cutoff function [6], and the sum runs over the neighborhood $\mathcal{N}(i)$ of atom i.

Message Passing Update Two branches of attention-weighted MP steps are defined for the feature vectors $\mathbf{f}$ and SPHCs $\boldsymbol{\chi}$ (see Fig. 1a). After initialization (eqs. (9) and (11)), the features are updated as

$$\mathbf{f}_i' = \mathbf{f}_i + \sum_{j \in \mathcal{N}(i)} \phi_{r_{\text{cut}}}(r_{ij}) \cdot \alpha_{ij} \cdot \mathbf{f}_j, \tag{12}$$

where $\alpha_{ij} \in \mathbb{R}$ are self-attention [48, 49] coefficients (see below). In analogy to the feature vectors, it is possible to define an MP update for the SPHCs as

$$\boldsymbol{\chi}_i'^{(l)} = \boldsymbol{\chi}_i^{(l)} + \sum_{j \in \mathcal{N}(i)} \phi_{r_{\text{cut}}}(r_{ij}) \cdot \alpha_{ij}^{(l)} \cdot \mathbf{Y}^{(l)}(\hat{\boldsymbol{r}}_{ij}), \tag{13}$$

where individual attention coefficients $\alpha_{ij}^{(l)} \in \mathbb{R}$ for each degree of the SPHCs are computed using multi-head attention [48]. However, with this definition, both MP updates are limited to local neighborhoods $\mathcal{N}(i)$. To be able to model non-local effects, we introduce the SPHC distance matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ with entries $\chi_{ij} = \|\chi_i - \chi_j\|_2$, i.e. distances between two atoms i and j in SPHC space for all possible pair-wise combinations of n atoms. To have uniform scales, we further apply the softmax along each row of $\mathbf{X}$ to generate a rescaled matrix $\tilde{\mathbf{X}} = \text{softmax}(\mathbf{X})$ with entries $\tilde{\chi}_{ij}$. A polynomial cutoff function $\phi_{\tilde{\chi}_{\text{cut}}}$ [29] is then applied to $\tilde{\mathbf{X}}$ to define spherical neighborhoods $\mathcal{N}_{\boldsymbol{\chi}}(i)$ (see A.2), which may include atoms that are far away in Euclidean space (see Fig. 1f). The spherical cutoff distance is chosen as $\tilde{\chi}_{\text{cut}} = 1/n$ to ensure that spherical neighborhoods remain small, even when going to larger molecules. We then incorporate non-local geometric corrections into the MP update of the SPHCs as

$$\boldsymbol{\chi}_i'^{(l)} = \underbrace{\sum_{j \in \mathcal{N}(i)} \phi_{r_{\text{cut}}}(r_{ij}) \cdot \alpha_{ij}^{(l)} \cdot \mathbf{Y}^{(l)}(\hat{\boldsymbol{r}}_{ij})}_{\text{local in } \mathbb{R}^3} + \underbrace{\sum_{j \in \mathcal{N}_{\boldsymbol{\chi}}(i)} \phi_{\tilde{\chi}_{\text{cut}}}(\tilde{\chi}_{ij}) \cdot \alpha_{ij}^{(l)} \cdot \mathbf{Y}^{(l)}(\hat{\boldsymbol{r}}_{ij})}_{\text{local in } \boldsymbol{\chi}, \text{ but non-local in } \mathbb{R}^3}. \tag{14}$$

Self-Attention The self-attention coefficients in eqs. (12)–(14) are calculated as

$$\alpha_{ij} = \mathbf{f}_i^T(\mathbf{w}_{ij} \odot \mathbf{f}_j)/\sqrt{F}, \tag{15}$$

where $\mathbf{w}_{ij} \in \mathbb{R}^F$ is the output of a filter generating function and '$\odot$' denotes the element-wise product. The filter maps the Euclidean distance r_{ij} and per-degree SPHC distances $\chi_{ij}^{(l)} = \|\boldsymbol{\chi}_j^{(l)} - \boldsymbol{\chi}_i^{(l)}\|_2$ between the current SPHCs of atoms i and j into the feature space $\mathbb{R}^F$ (as a short-hand, we write the vector containing all per-degree SPHC distances as $[\chi_{ij}^{(l)}]_{l \in \mathcal{L}}$). It is built as the linear combination of two filter-generating functions

$$\mathbf{w}_{ij} = \underbrace{\phi_r(r_{ij})}_{\text{radial filter}} + \underbrace{\phi_s\left([\chi_{ij}^{(l)}]_{l \in \mathcal{L}}\right)}_{\text{spherical filter}}, \tag{16}$$

which separately act on the Euclidean and SPHC distances. We call $\phi_r : \mathbb{R} \mapsto \mathbb{R}^F$ the *radial filter* function and $\phi_s : \mathbb{R}^{|\mathcal{L}|} \mapsto \mathbb{R}^F$ the *spherical filter* function (an ablation study for ϕ_s can be found in appendix A.4). Since per-atom features $\mathbf{f}_i$, interatomic distances r_{ij}, and per-degree distances $\chi_{ij}^{(l)}$ are invariant under rotations (proof in appendix A.1), so are the self-attention coefficients α_{ij}.

Atomwise Interaction After each MP update, features and SPHCs are coupled with each other according to

$$\mathbf{f}_i' = \mathbf{f}_i + \phi_1 \left(\mathbf{f}_i, [\chi_i^{(l)}]_{l \in \mathcal{L}}, [\tilde{\chi}_i^{(l)}]_{l \in \mathcal{L}} \right) , \tag{17}$$

$$\boldsymbol{\chi}_i'^{(l)} = \boldsymbol{\chi}_i^{(l)} + \phi_2^{(l)} \left(\mathbf{f}_i, [\chi_i^{(l)}]_{l \in \mathcal{L}}, [\tilde{\chi}_i^{(l)}]_{l \in \mathcal{L}} \right) \boldsymbol{\chi}_i^{(l)} + \phi_3^{(l)} \left([\tilde{\chi}_i^{(l)}]_{l \in \mathcal{L}} \right) \tilde{\boldsymbol{\chi}}_i^{(l)} , \tag{18}$$

where $\phi_1 : \mathbb{R}^{F+2|\mathcal{L}|} \mapsto \mathbb{R}^F$, $\phi_2^{(l)} : \mathbb{R}^{F+2|\mathcal{L}|} \mapsto \mathbb{R}$, and $\phi_3^{(l)} : \mathbb{R}^{|\mathcal{L}|} \mapsto \mathbb{R}$. In the inputs to $\phi_{1,2,3}$, degree-wise scalars $\chi^{(l)} = \|\boldsymbol{\chi}^{(l)}\|_2$ are used to preserve equivariance. The coupling step additionally includes cross-degree coupled SPHCs $\tilde{\chi}_i^{(l)}$ for each degree l. Following [47] they are constructed as

$$\tilde{\boldsymbol{\chi}}_i^{(l)} = \sum_{l_1=l_{\min}}^{l_{\max}} \sum_{l_2=l_1+1}^{l_{\max}} k_{l_1,l_2,l} \left(\boldsymbol{\chi}_i^{(l_1)} \otimes_l \boldsymbol{\chi}_i^{(l_2)} \right) , \tag{19}$$

where $k_{l_1,l_2,l} \in \mathbb{R}$ are learnable coefficients for all valid combinations of l_1, l_2 given l and the term in brackets is the contraction of degrees l_1 and l_2 into degree l (eq. (8)).

SO3KRATES architecture Using the design paradigm above, we build the transformer network SO3KRATES, which consists of a self-attention block on $\mathcal{F}$ and $\mathcal{X}$ (eqs. (12) and (13)), respectively, as well as an interaction block (eqs. (17) and (18)) per layer. After initialization of the features and the SPHCs according to eqs. (9) and (11), they are updated iteratively by passing through n_l layers. Atomic energy contributions $E_i \in \mathbb{R}$ are predicted from the features of the final layer using a two-layered output block. The individual contributions are summed to the total energy prediction $E = \sum_i^n E_i$. See Fig. 1b for an overview. More details on the implementation, training details and network hyperparameters are given in appendix A.3.

4 Experiments

4.1 Non-Local Geometric Interactions

For efficiency reasons, MPNNs only consider interactions between atoms in local neighborhoods, i.e. within a cutoff radius r_{cut}. Thus, information can only be propagated over a distance of r_{cut} within a single MP step. Although multiple MP updates increase the effective cutoff distance, because information can "hop" between different neighborhoods as long as they share at least one atom, each MP step is accompanied by an undesirable loss of information, which limits the accuracy that can be obtained. Consequently, MPNNs are unable to describe non-local effects on length-scales that exceed the effective cutoff distance. To illustrate this problem, we consider the challenging open task [21] of learning the potential energy of cumulene molecules with different sizes (see Fig. 2a). Here, the relative orientation of the hydrogen rotors at the far ends of the molecule strongly influences its energy due to non-local electronic effects [21]. In order to be able to successfully learn the energy profile with a local model, the effective cutoff has to be large enough to allow information to propagate from one hydrogen rotor to the other.

As a representative example for MPNNs, we consider the recently proposed NEQUIP model [32], which achieves SOTA performance on several benchmarks. We find that even when the effective cutoff radius is large enough in principle, an MPNN with $n_l = 4$, $r_{\text{cut}} = 2.5$ Å, and $l_{\max} \leq 1$ fails to learn the correct energy profile. This is due to the fact that the relevant geometric information "cancels out" (similar to addition of vectors oriented in opposite directions) within each neighborhood, underlining the limited expressiveness of mean-field interactions in MPNNs. Only by including higher-order geometric correlations, e.g. going to $l_{\max} = 3$, the correct energy profile can be recovered (at the cost of computational efficiency). When going to even larger cumulene structures, however, the effective cutoff becomes too small and it is necessary to increase the number of MP layers to solve the task (again, at the cost of lower computational efficiency), which is illustrated

Figure 2: **(a)** Energy predictions for different cumulene structures with NEQUIP ($l_{\max} = 1$ and $l_{\max} = 3$) and SO3KRATES ($l_{\max} = 1$) models with and without geometric corrections (eqs. (13) and (14)). Cutoff radius and number of layers are kept constant at $r_{\mathrm{cut}} = 2.5\,\text{Å}$ and $n_l = 4$, respectively. **(b)** Visualization of attention coefficients α_{ij} from the geometric corrections (eq. (14)) after training for different angles Θ between hydrogen rotors.

in appendix A.5. Neither increasing the maximum degree of interactions $l_{\max}$, nor the number of layers n_l, is a satisfactory workaround: Instead of offering a general solution to describe non-local interactions, both options decrease computational efficiency, while only shifting the problem to larger length-scales or higher-order geometric correlations.

In contrast, our proposed SO3KRATES architecture is able to reproduce the energy profile for cumulene molecules of all sizes independent of the effective cutoff radius. Crucially, even with $l_{\max} = 1$, the predicted energy matches the *ab-initio* reference faithfully. We find that geometric corrections in the MP update of the SPHCs (cf. eq. (14)) are responsible for the increased capability of describing higher-order geometric correlations, as a SO3KRATES model with a naive MP update (cf. eq. (13)) fails to solve this task with $l_{\max} = 1$ (see Fig. 2a). We further confirm that the model picks up on the physically relevant interaction between the hydrogen rotors by analysing the attention values after training (Fig. 2b). To illustrate how SO3KRATES is able to describe non-local effects, we show a low-dimensional projection of the atomic SPHCs before and after training for the largest of the cumulene molecules (Fig. 3). After training, the SPHCs for hydrogen atoms at opposite ends of the molecule are embedded close together in SPHC space, allowing SO3KRATES to efficiently model the non-local geometric dependence between the hydrogen rotors.

4.2 Data Efficiency and Generalization

As pointed out in [31] and [32], equivariant features not only increase performance, but also improve data efficiency. The latter is particularly important, as *ab-initio* methods for reference data generation can become exceedingly expensive when high accuracy is required. Here, we use a subset of the recently introduced QM7-X data set [50], which we call QM7-X250. It contains 250 different molecular structures, each with 80 data points for training, 10 data points for validation and 11-3748 data points for testing (for details, see appendix A.6). The small number of training/validation samples per molecule makes it particularly suited for evaluating model behavior in the low data regime. In the following, we train (1) one model per structure in QM7-X250 and (2) one model for all structures in QM7-X250 ($250 \times 80 = 20k$ training points), which we refer to as *individual* and *joint* models, respectively.

We start by investigating the performance as a function of the maximal degree $l_{\max}$ and find that the error strongly decreases with higher $l_{\max}$ (Fig. 4a). As kernel methods are known to perform well in

Figure 3: **(a)** Cumulene structure in Euclidean space. **(b)** Low-dimensional projections of the SPHCs ($l_{\max} = 1$) in SO3KRATES before and after training. Carbon atoms are grey and hydrogen atoms on different ends of the molecule are green and red, respectively. During training, the model maps hydrogen atoms at opposite ends close together in SPHC space (which is not the case at initialization).

Figure 4: **(a)** Average force MAE as a function of $l_{\max}$ on 250 structures from QM7-X [50]. The blue solid line corresponds to SO3KRATES models trained on each of the 250 structures (à 80 training points), individually. The jointly trained SO3KRATES model (red solid line) is a single model trained on all 250×80 training points. **(b)** Accuracy of force predictions for unknown structures as function of $l_{\max}$. Generalization is only investigated for the jointly trained model. Due to the way the molecular descriptor is designed, SGDML [38] models can only be trained on individual structures.

the low-data regime [21], we compare our results to SGDML [38] kernel models, which only use distances as a molecular descriptor (corresponding to $l_{\max} = 0$). For $l_{\max} = 0$, we find SGDML gives competitive results, whereas for $l_{\max} = 1$, SO3KRATES starts to outperform SGDML. As soon as $l_{\max} \geq 2$, however, the prediction accuracy of SO3KRATES is greater than that of SGDML by a large margin. Thus, increasing the order of geometric information in the SPHCs leads to strong improvements in the low-data regime. For jointly trained models, we find that SO3KRATES outperforms SGDML even for $l_{\max} = 0$, with continuous improvement for increasing $l_{\max}$. In appendix A.7 we report energy *and* force errors across degrees and further experimental details.

The generalization capability of SO3KRATES is tested, by applying a jointly trained model to 25 completely unknown molecules from the QM7-X dataset (see, Fig. 4b, details in appendix A.7). Again, we find that force MAEs decrease with increasing $l_{\max}$. For reference, we compare SO3KRATES to individually trained SGDML models and find that SO3KRATES performs on par, or even slightly better, for $l_{\max} \geq 2$. Going beyond $l_{\max} = 2$ is found to only marginally improve generalization.

For completeness, we also apply SO3KRATES to the popular MD17 benchmark (see Table 1). We find, that SO3KRATES outperforms networks that have the same parameter complexity by a large margin (PAINN and NEWTONNET). Notably, it requires significantly less parameters than other SH based architectures (NEQUIP and SPOOKYNET), while performing only slightly worse or even on par with them. Compared to current SH based approaches, GEMNETQ needs less parameters (still ~ 2.5x more than SO3KRATES) to achieve competitive results. However, it requires the explicit calculation of dihedral angles which scales cubically in the number of neighboring atoms (compared to the linear scaling of all other listed architectures). Due to its scaling and lightweight structure, SO3KRATES can significantly reduce the time for training and inference (see Fig. 1d and A.8).

Table 1: MAE for energy (in kcal mol^{-1}) and forces (in $\text{kcal mol}^{-1}\,\text{Å}^{-1}$) of current state-of-the-art machine learning models on the MD17 benchmark for 1k training points. For each model, the number of paramaeters, as well as the scaling in the number of neighbors m is shown.

		NEQUIP [32] 3M | $\mathcal{O}(m)$	SPOOKYNET [30] 3M | $\mathcal{O}(m)$	GEMNETQ [34] 2.2M | $\mathcal{O}(m^3)$	PAINN [31] 600k | $\mathcal{O}(m)$	NEWTONNET [51] 500k | $\mathcal{O}(m)$	SO3KRATES 700k | $\mathcal{O}(m)$
Aspirin	*energy*	0.13	0.151	–	0.159	0.168	0.139
	forces	0.19	0.258	0.217	0.371	0.348	0.236
Ethanol	*energy*	0.05	0.052	–	0.063	0.061	0.052
	forces	0.09	0.094	0.088	0.230	0.211	0.096
Malondialdehyde	*energy*	0.08	0.079	–	0.091	0.096	0.077
	forces	0.13	0.167	0.159	0.319	0.323	0.147
Naphthalene	*energy*	0.11	0.116	–	0.117	0.118	0.115
	forces	0.04	0.089	0.051	0.083	0.084	0.074
Salicylic Acid	*energy*	0.11	0.114	–	0.114	0.115	0.106
	forces	0.09	0.180	0.124	0.209	0.197	0.145
Toluene	*energy*	0.09	0.094	–	0.097	0.094	0.095
	forces	0.05	0.087	0.060	0.102	0.088	0.073
Uracil	*energy*	0.10	0.105	–	0.104	0.107	0.103
	forces	0.08	0.119	0.104	0.140	0.149	0.111

5 Discussion and Conclusion

Due to the locality assumption used in most MPNNs, they are unable to model non-local electronic effects, which result in global geometric dependencies between different parts of a molecule. The length-scales of such interactions often greatly exceed the cutoff radius used in the MP step, and even though stacking multiple MP layers increases the effective cutoff, ultimately, MPNNs are not capable of efficiently modeling geometric dependencies on arbitrary length scales.

In this work, we contribute conceptually by proposing an efficient and scalable solution to this problem. We suggest a set of refinable, equivariant coordinates for point clouds in Euclidean space, called spherical harmonic coordinates (SPHCs). Non-local geometric effects can then be efficiently modeled by including geometric corrections, which are localized in the space of SPHCs, but non-local in Euclidean space. Further, we show that introducing spherical filter functions acting on the SPHCs increases geometric resolution and predictive accuracy.

We then propose the SO3KRATES architecture, a self-attention based MPNN, which decouples atomic features from higher-order geometric information. This allows to drastically decrease the parametric complexity while still achieving SOTA prediction accuracy. We show evidence that increasing the geometric order of SPHCs greatly improves model performance in the low-data regime, as well as generalization to unknown molecules.

A limitation of the current implementation of SO3KRATES is that spherical neighborhoods $\mathcal{N}_\chi$ in eq. (14) are computed from all pairwise distances in SPHC space. An alternative implementation could use a space partitioning scheme to find neighborhoods more efficiently. In a broader context, our work falls into the category of approaches that can help to reduce the vast computational complexity of molecular and material simulations. This can accelerate novel drug and material designs, which holds the promise of tackling societal challenges, such as climate change and sustainable energy supply [52]. Of course, our method could also be used for nefarious applications, e.g. design of chemical warfare, but this is true for all quantum chemistry methods.

Future research will focus on applications of SO3KRATES to materials and bio-molecules, which are typical examples of chemical systems where the accurate description of non-local effects is necessary to produce novel insights. Beyond atomic interactions, efficient treatment of non-local effects in point cloud data goes beyond the domain of quantum chemistry. Large-scale geometric dependencies could also be relevant in the domain of 3d computer vision, where the relative orientation of certain objects far apart from each other may play an important role. For example, consider LIDAR sensor data used in the development of self-driving cars. Here, non-local correlations, for example between two people on opposite sides of a road, might be relevant to determine whether to break or accelerate.

References

[1] Mark E Tuckerman. Ab initio molecular dynamics: basic concepts, current trends and novel applications. *J. Phys. Condens. Matter*, 14(50):R1297, 2002.

[2] Frank Noé, Alexandre Tkatchenko, Klaus-Robert Müller, and Cecilia Clementi. Machine learning for molecular simulation. *Annu. Rev. Phys. Chem.*, 71:361–390, 2020.

[3] John A Keith, Valentin Vassilev-Galindo, Bingqing Cheng, Stefan Chmiela, Michael Gastegger, Klaus-Robert Müller, and Alexandre Tkatchenko. Combining machine learning and computational chemistry for predictive insights into chemical systems. *Chemical Reviews*, 121 (16):9816–9872, 2021. URL https://pubs.acs.org/doi/abs/10.1021/acs.chemrev.1c00107.

[4] Jörg Behler and Michele Parrinello. Generalized neural-network representation of high-dimensional potential-energy surfaces. *Phys. Rev. Lett.*, 98(14):146401, 2007.

[5] Albert P Bartók, Mike C Payne, Risi Kondor, and Gábor Csányi. Gaussian Approximation Potentials: the accuracy of quantum mechanics, without the electrons. *Phys. Rev. Lett.*, 104(13): 136403, 2010.

[6] Jörg Behler. Atom-centered symmetry functions for constructing high-dimensional neural network potentials. *J. Chem. Phys.*, 134(7):074106, 2011.

[7] Albert P Bartók, Risi Kondor, and Gábor Csányi. On representing chemical environments. *Phys. Rev. B*, 87(18):184115, 2013.

[8] Zhenwei Li, James R Kermode, and Alessandro De Vita. Molecular dynamics with on-the-fly machine learning of quantum-mechanical forces. *Phys. Rev. Lett.*, 114(9):096405, 2015.

[9] Stefan Chmiela, Alexandre Tkatchenko, Huziel E Sauceda, Igor Poltavsky, Kristof T Schütt, and Klaus-Robert Müller. Machine learning of accurate energy-conserving molecular force fields. *Sci. Adv.*, 3(5):e1603015, 2017.

[10] Kristof T Schütt, Farhad Arbabzadah, Stefan Chmiela, Klaus-Robert Müller, and Alexandre Tkatchenko. Quantum-chemical insights from deep tensor neural networks. *Nat. Commun.*, 8: 13890, 2017.

[11] Michael Gastegger, Jörg Behler, and Philipp Marquetand. Machine learning molecular dynamics for the simulation of infrared spectra. *Chem. Sci.*, 8(10):6924–6935, 2017.

[12] Stefan Chmiela, Huziel E. Sauceda, Klaus-Robert Müller, and Alexandre Tkatchenko. Towards exact molecular dynamics simulations with machine-learned force fields. *Nat. Commun.*, 9(1): 3887, 2018. doi: 10.1038/s41467-018-06169-2.

[13] Kristof T Schütt, Huziel E Sauceda, Pieter-Jan Kindermans, Alexandre Tkatchenko, and Klaus-Robert Müller. SchNet – a deep learning architecture for molecules and materials. *J. Chem. Phys.*, 148(24):241722, 2018.

[14] Justin Gilmer, Samuel S Schoenholz, Patrick F Riley, Oriol Vinyals, and George E Dahl. Neural message passing for quantum chemistry. In *International Conference on Machine Learning*, pages 1263–1272. Pmlr, 2017.

[15] Justin S Smith, Olexandr Isayev, and Adrian E Roitberg. ANI-1: an extensible neural network potential with DFT accuracy at force field computational cost. *Chem. Sci.*, 8(4):3192–3203, 2017.

[16] Nicholas Lubbers, Justin S Smith, and Kipton Barros. Hierarchical modeling of molecular energies using a deep neural network. *J. Chem. Phys.*, 148(24):241715, 2018.

[17] Martin Stöhr, Leonardo Medrano Sandonas, and Alexandre Tkatchenko. Accurate many-body repulsive potentials for density-functional tight-binding from deep tensor neural networks. *J. Phys. Chem. Lett.*, 11:6835–6843, 2020.

[18] Felix A Faber, Anders S Christensen, Bing Huang, and O Anatole von Lilienfeld. Alchemical and structural distribution based representation for universal quantum machine learning. *J. Chem. Phys.*, 148(24):241717, 2018.

[19] Oliver T Unke and Markus Meuwly. PhysNet: A neural network for predicting energies, forces, dipole moments, and partial charges. *J. Chem. Theory Comput.*, 15(6):3678–3693, 2019.

[20] Anders S Christensen, Lars A Bratholm, Felix A Faber, and O Anatole von Lilienfeld. FCHL revisited: faster and more accurate quantum machine learning. *J. Chem. Phys.*, 152(4):044107, 2020.

[21] Oliver T Unke, Stefan Chmiela, Huziel E Sauceda, Michael Gastegger, Igor Poltavsky, Kristof T Schütt, Alexandre Tkatchenko, and Klaus-Robert Müller. Machine learning force fields. *Chemical Reviews*, 121(16):10142–10186, 2021.

[22] Yaolong Zhang, Ce Hu, and Bin Jiang. Embedded Atom Neural Network Potentials: efficient and accurate machine learning with a physically inspired representation. *J. Phys. Chem. Lett.*, 10(17):4962–4967, 2019.

[23] Silvan Käser, Oliver Unke, and Markus Meuwly. Reactive dynamics and spectroscopy of hydrogen transfer from neural network-based reactive potential energy surfaces. *New J. Phys.*, 22:55002, 2020.

[24] O Anatole von Lilienfeld, Klaus-Robert Müller, and Alexandre Tkatchenko. Exploring chemical compound space with quantum-based machine learning. *Nat. Rev. Chem.*, 4(7):347–358, 2020.

[25] Victor Garcia Satorras, Emiel Hoogeboom, and Max Welling. E (n) equivariant graph neural networks. In *International Conference on Machine Learning*, pages 9323–9332. PMLR, 2021.

[26] Michael M Bronstein, Joan Bruna, Taco Cohen, and Petar Veličković. Geometric deep learning: Grids, groups, graphs, geodesics, and gauges. *arXiv preprint arXiv:2104.13478*, 2021.

[27] Yann LeCun, Yoshua Bengio, et al. Convolutional networks for images, speech, and time series. *The handbook of brain theory and neural networks*, 3361(10):1995, 1995.

[28] Thomas N Kipf and Max Welling. Semi-supervised classification with graph convolutional networks. *arXiv preprint arXiv:1609.02907*, 2016.

[29] Johannes Klicpera, Janek Groß, and Stephan Günnemann. Directional message passing for molecular graphs. *arXiv preprint arXiv:2003.03123*, 2020.

[30] Oliver T Unke, Stefan Chmiela, Michael Gastegger, Kristof T Schütt, Huziel E Sauceda, and Klaus-Robert Müller. SpookyNet: Learning force fields with electronic degrees of freedom and nonlocal effects. *Nat. Commun.*, 12:7273, 2021.

[31] Kristof T Schütt, Oliver T Unke, and Michael Gastegger. Equivariant message passing for the prediction of tensorial properties and molecular spectra. *arXiv preprint arXiv:2102.03150*, 2021.

[32] Simon Batzner, Tess E Smidt, Lixin Sun, Jonathan P Mailoa, Mordechai Kornbluth, Nicola Molinari, and Boris Kozinsky. SE (3)-equivariant graph neural networks for data-efficient and accurate interatomic potentials. *arXiv preprint arXiv:2101.03164*, 2021.

[33] Nathaniel Thomas, Tess Smidt, Steven Kearnes, Lusann Yang, Li Li, Kai Kohlhoff, and Patrick Riley. Tensor field networks: rotation-and translation-equivariant neural networks for 3D point clouds. *arXiv preprint arXiv:1802.08219*, 2018.

[34] Johannes Klicpera, Florian Becker, and Stephan Günnemann. Gemnet: Universal directional graph neural networks for molecules. *arXiv preprint arXiv:2106.08903*, 2021.

[35] Sergey N Pozdnyakov, Michael J Willatt, Albert P Bartók, Christoph Ortner, Gábor Csányi, and Michele Ceriotti. Incompleteness of atomic structure representations. *Physical Review Letters*, 125(16):166001, 2020.

[36] Fabian B Fuchs, Daniel E Worrall, Volker Fischer, and Max Welling. Se (3)-transformers: 3d roto-translation equivariant attention networks. *arXiv preprint arXiv:2006.10503*, 2020.

[37] Aneja Alisha. Performance comparison between nvidia's geforce gtx 1080 and tesla p100 for deep learning. `https://github.com/alisha17/benchmarks`, 2017.

[38] Stefan Chmiela, Huziel E Sauceda, Igor Poltavsky, Klaus-Robert Müller, and Alexandre Tkatchenko. sgdml: Constructing accurate and data efficient molecular force fields using machine learning. *Computer Physics Communications*, 240:38–45, 2019.

[39] Brandon Anderson, Truong-Son Hy, and Risi Kondor. Cormorant: covariant molecular neural networks. *arXiv preprint arXiv:1906.04015*, 2019.

[40] Philipp Thölke and Gianni De Fabritiis. Equivariant transformers for neural network based molecular potentials. In *International Conference on Learning Representations*, 2021.

[41] Matan Atzmon, Haggai Maron, and Yaron Lipman. Point convolutional neural networks by extension operators. *arXiv preprint arXiv:1803.10091*, 2018.

[42] J-F Cardoso and Beate H Laheld. Equivariant adaptive source separation. *IEEE Transactions on signal processing*, 44(12):3017–3030, 1996.

[43] Taco S Cohen and Max Welling. Group equivariant convolutional networks. In *International conference on machine learning*, pages 2990–2999. PMLR, 2016.

[44] Geoffrey E Hinton, Alex Krizhevsky, and Sida D Wang. Transforming auto-encoders. In *International conference on artificial neural networks*, pages 44–51. Springer, 2011.

[45] Jonas Köhler, Leon Klein, and Frank Noé. Equivariant flows: exact likelihood generative learning for symmetric densities. In *International Conference on Machine Learning*, pages 5361–5370. PMLR, 2020.

[46] Eugene Paul Wigner. Gruppentheorie und ihre anwendung auf die quantenmechanik der atomspektren. 1931.

[47] Oliver Unke, Mihail Bogojeski, Michael Gastegger, Mario Geiger, Tess Smidt, and Klaus-Robert Müller. Se(3)-equivariant prediction of molecular wavefunctions and electronic densities. *Advances in Neural Information Processing Systems*, 34, 2021.

[48] Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Lukasz Kaiser, and Illia Polosukhin. Attention is all you need. *arXiv preprint arXiv:1706.03762*, 2017.

[49] Petar Velickovic, Guillem Cucurull, Arantxa Casanova, Adriana Romero, Pietro Lio, and Yoshua Bengio. Graph attention networks. *arXiv preprint arXiv:1710.10903*, 2017.

[50] Johannes Hoja, Leonardo Medrano Sandonas, Brian G Ernst, Alvaro Vazquez-Mayagoitia, Robert A DiStasio Jr, and Alexandre Tkatchenko. Qm7-x, a comprehensive dataset of quantum-mechanical properties spanning the chemical space of small organic molecules. *Scientific data*, 8(1):1–11, 2021.

[51] Mojtaba Haghighatlari, Jie Li, Xingyi Guan, Oufan Zhang, Akshaya Das, Christopher J Stein, Farnaz Heidar-Zadeh, Meili Liu, Martin Head-Gordon, Luke Bertels, et al. Newtonnet: A newtonian message passing network for deep learning of interatomic potentials and forces. *arXiv preprint arXiv:2108.02913*, 2021.

[52] Anubhav Jain, Shyue Ping Ong, Geoffroy Hautier, Wei Chen, William Davidson Richards, Stephen Dacek, Shreyas Cholia, Dan Gunter, David Skinner, Gerbrand Ceder, et al. Commentary: The materials project: A materials genome approach to accelerating materials innovation. *APL materials*, 1(1):011002, 2013.