Revisit last-iterate convergence of mSGD under milder requirement on step size

Abstract

Understanding convergence of SGD-based optimization algorithms can help deal with enormous machine learning problems. To ensure last-iterate convergence of SGD and momentum-based SGD (mSGD), the existing studies usually constrain the step size ϵ_n to decay as $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$, which however is rather conservative and may lead to slow convergence in the early stage of the iteration. In this paper, we relax this requirement by studying an alternate step size for the mSGD. First, we relax the requirement of the decay on step size to $\sum_{n=1}^{+\infty} \epsilon_n^{2+\eta_0} < +\infty$ $(0 \leq \eta_0 < 1/2)$. This implies that a larger step size, such as $\epsilon_n = \frac{1}{\sqrt{n}}$ can be utilized for accelerating the mSGD in the early stage. Under this new step size and some common conditions, we prove that the last iterate of mSGD for non-convex loss functions asymptotically converges to a connected set of stationary points. In addition, we show that this step size can indeed help make the algorithm converge to a neighbor of the stationary points quicker in the early stage. In addition, we establish the convergence of mSGD under a constant step size $\epsilon_n \equiv \epsilon > 0$ by removing the common requirement in the literature on the strong convexity of the loss function. Some experiments are given to illustrate the developed results.

1 Introduction

The booming development of machine learning over the past decade relies on the employment of effective optimization algorithms. A large number of such optimization algorithms are based on gradient descent (GD). The problem in learning can be cast as minimizing a loss function $g(\theta) \in \mathbb{R}$ over the choice of an N-dimensional real-valued parameter θ, i.e., by solving the problem

$$\theta^* = \arg \min_{\theta \in \mathbb{R}^N} g(\theta). \tag{1}$$

This problem can be solved with a typical GD algorithm through an iteration of the form

$$\theta_{n+1} = \theta_n - \epsilon_n \nabla_{\theta_n} g(\theta_n), \tag{2}$$

where θ_n is the estimate of θ^* at step n, ϵ_n is a positive step size (learning rate) to be designed, and $\nabla_{\theta_n} g(\theta_n)$ stands for the gradient of $g(\theta_n)$ at step n. Under certain technical conditions, θ_n in (2) can asymptotically (as $n \to \infty$) converge to the optimal solution θ^*. However, (2) is not efficient in machine leaning applications especially with enormous data. To accelerate (2), one attempt is stochastic gradient descent (SGD) originating from [1]. Instead of calculating $\nabla_{\theta_n} g(\theta_n)$ over all the data, SGD computes gradient estimate $\nabla_{\theta_n} g(\theta_n, \xi_n)$ by sampling a subset of data (using the distribution of the sampling noise $\{\xi_n\}$ and then updates the iterate as follows

$$\theta_{n+1} = \theta_n - \epsilon_n \nabla_{\theta_n} g(\theta_n, \xi_n). \tag{3}$$

In addition to the sampling noise, ξ_n can be used to model any external noises arising from the gradient computation.

However the price to pay for this data efficiency is that the employment of stochastic gradients (3) displaying a relatively slow convergence rate during the learning process. In order to improve the convergence rate of SGD, momentum-based stochastic gradient descend (mSGD), which reduces the update variance by averaging the past gradients ([2]), has been proposed. An iteration in mSGD is as follows ([3, 4]):

$$v_n = \alpha v_{n-1} + \epsilon_n \nabla_{\theta_n} g(\theta_n, \xi_n), \quad \theta_{n+1} = \theta_n - v_n, \tag{4}$$

where $\alpha \in [0, 1)$ and $\epsilon_n > 0$ are momentum coefficient and step size (learning rate), respectively. An alternative formulation of mSGD, named stochastic heavy ball (SHB) [5, 6], has also been proposed as follows

$$v_n = \beta_n v_{n-1} + (1 - \beta_n) \nabla_{\theta_n} g(\theta_n, \xi_n), \quad \theta_{n+1} = \theta_n - \gamma_n v_n, \tag{5}$$

where $\beta_n \in (0, 1)$ and γ_n are momentum coefficient and step size. It has been shown that mSGD and SHB are essentially equivalent [7]. In recent years, mSGD has been widely employed in many applications of deep learning such as image classification [8], fault diagnosis [9], statistical image reconstruction [10], among others. Moreover, a number of variants on momentum have been emerging, see, e.g., synthesized Nesterov variants [11], robust momentum [12], and PID-control based methods [13]. The importance of momentum in deep learning has been illustrated in [4] through experiments.

There has been a long line of literature analyzing the convergence of SGD and mSGD algorithms. Regarding SGD, authors in [14] studied the last-iterate convergence when the step size is chosen to decay as $O(\frac{1}{n})$. Authors in [15, 16] established the last-iterate convergence of SGD for strongly convex loss functions without the bounded gradients assumption. For the normalized mSGD (SHB), results from Polyak [17, 2] and Kaniovski [18] established the convergence (subsequence convergence and convergence of time averages) properties for convex loss functions. Igor Gitman [5] provided convergence results for mSGD (SHB) for non-convex loss functions under the somewhat restrictive requirement for uniform boundedness of a noise term $\mathbb{E}(\|\nabla_{\theta_n} g(\theta_n, \xi_n) - \nabla_{\theta_n} g(\theta_n)\|) \leq \delta$. It has been pointed out [19, 4] that the designs of momentum coefficients in [17, 2, 18, 5] may not be consistent with the requirements of some practical applications. The last-iterate convergence of mSGD for non-convex loss functions has recently been established in [20, 7] with the step size condition, i.e., $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$ that has been traditionally utilized in stochastic optimization [1, 21, 16, 15]. Under the strong convexity condition, since this condition requires the step size to decay relatively fast, it may lead to slow convergence especially in the early stage of the iteration. With the requirement that the loss function is strongly convex, [22] showed that the last-iterate of mSGD can converge to a neighbor of the stationary point when the step size is a constant.

2   Problem of interest and contributions

In this paper, we consider the problem of last-iterate convergence of mSGD under relaxed requirements on step sizes. Specifically, we seek to relax the condition $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$ [1, 21, 16, 15, 7] that has been required in the literature for proving last-iterate convergence of mSGD. We consider two possible step sizes: step size that decay as $\epsilon_n = \frac{1}{\sqrt{n}}$ and constant step size $\epsilon_n \equiv \epsilon > 0$. Using such larger step size is expected to lead to the iterate of mSGD converging faster. We note that since the SGD (3) is a special case of mSGD (4), the developed results in this paper for mSGD also work for SGD.

The contributions of this paper are summarized as follows:

1) Under the setting of step size decaying as $\epsilon_n = \frac{1}{\sqrt{n}}$, we prove that the last iterate of mSGD for non-convex loss functions can still converge to a connected set of stationary points almost surely. This result is more general than mean-square gradient convergence established in [20]. In fact, we generalize the result for any step sizes of the form $\sum_{n=1}^{+\infty} \epsilon_n^{2+\eta_0} < +\infty$ ($0 \leq \eta_0 < 1/2$), which is a superset of the widely required condition $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$ [1, 21, 16, 15, 7]. Given this condition, one can run mSGD by employing a larger step size like $\epsilon_n = \frac{1}{\sqrt{n}}$ for last-iterate convergence.

2) Under some mild conditions, we provide an estimate of the convergence rate of mSGD. Furthermore, we reveal that the new step size can indeed help to improve the convergence speed in the early stage of the iteration.

3) Under the setting of constant step size, first we prove that given any small neighborhood of stationary points, it is feasible to design a constant step size, such that there is a convergent iterate subsequence of mSGD staying within the neighborhood almost surely. In addition, we prove that the mean-square gradient can be arbitrary small by tunning the step size. Comparing with [22], we remove the requirement of strong convexity on loss function.

Regarding the definitions on sequence convergence, the following ones from literature are typical. Last-iterate convergence VS time-average convergence: For a stochastic variables sequence $\{\zeta_n\} \in \mathbb{R}^N$ with 0 as the only limit point. Then the sequence is said to satisfy last-iterate convergence if $\lim_{n \to +\infty} \|\zeta_n\| = 0$; and time-average mean-square convergence if $\lim_{n \to +\infty} \frac{1}{n} \sum_{k=1}^{n} \mathbb{E}(\|\zeta_k\|^2) = 0$. It can be proved that last-iterate convergence implies time-average convergence under the general setting of this problem.

Paper outline. The rest of the paper is organized as follows. In Section 3, we provide the main results of the paper on the convergence of mSGD under decaying step size and constant step size respectively. In Section 4, two experiments are given. Conclusion is made in Section 5. The main proofs are given in Appendix.

3 Main results

In this section, we provide the main results of this paper on the last-iterate convergence for mSGD in (4) under two possible step sizes: step sizes that decay as $\epsilon_n = \frac{1}{\sqrt{n}}$ and constant step size $\epsilon_n \equiv \epsilon > 0$.

3.1 Convergence of mSGD with decaying step size

The following assumptions are needed in this paper.

Assumption 3.1. *Loss function $g(\theta)$ in (1) satisfies the following conditions:*

1. *Noise sequence $\{\xi_n\}$ are mutually independent and independent of θ_1 and v_0, such that $g(\theta) = \mathbb{E}_{\xi_n}\left(g(\theta, \xi_n)\right)$ for any $\theta \in \mathbb{R}^N$.*

2. *$g(\theta)$ is a non-negative and continuously differentiable function. The set of its stationary points $J = \{\theta | \|\nabla_\theta g(\theta)\| = 0\}$ is a bounded set which has only finite connected components $J_1, ..., J_n$. In addition, there is $\epsilon_1 > 0$, such that for any i and $0 < d(\theta, J_i) < \epsilon_1$, it holds that $\left| g(\theta) - g_i \right| \neq 0$, where $g_i = \{g(\theta) | \theta \in J_i\}$ is a constant.*

3. *There are two constants $M' > 0$ and $a' > 0$ such that for any $\theta \in \mathbb{R}^N$ and $n \in \mathbb{N}_+$,*

$$E_{\xi_n}\left(\left\|\nabla_\theta g(\theta, \xi_n)\right\|^2\right) \leq M'\left\|\nabla_\theta g(\theta)\right\|^2 + a'. \tag{6}$$

4. *The gradient $\nabla_\theta g(\theta)$ satisfies the Lipschitz condition, i.e., there is a constant $c > 0$, such that for any $x, y \in \mathbb{R}^N$,*

$$\left\|\nabla_x g(x) - \nabla_y g(y)\right\| \leq c\|x - y\|.$$

Conditions 1, 3 and 4 in Assumption 3.1 are common in the literature [1, 21, 16, 15, 7]. Condition 2 is also a mild condition for the reasons as follows. In some works, the non-negative condition may be replaced by a finite low bound condition, i.e., $g(\theta) > \hat{l}_0 > -\infty$. These two conditions are essentially equivalent, since one can construct a new loss function $\overline{g} = g - \hat{l}_0$ for the finite low bound condition, such that the new loss function is non-negative. The rest of Condition 2 is quite general, since it allows the loss function to have multiple stationary points and to be non-convex.

Assumption 3.2. *In the mSGD (4), momentum coefficient $\alpha \in [0, 1)$ and the sequence of step size ϵ_n is positive and monotonically decreasing to zero, such that $\sum_{n=1}^{+\infty} \epsilon_n = +\infty$ and $\sum_{n=1}^{+\infty} \epsilon_n^{2+\eta_0} < +\infty$, where $0 \leq \eta_0 < 1/2$ is a constant.*

In Assumption 3.2, the momentum coefficient $\alpha \in [0, 1)$ is a constant. Compared with the setting in [6, 17, 18, 5], where $\alpha^{(n)}$ tends to 1 or 0, constant momentum coefficient is more general in practice [7, 19, 4]. Regarding the step size condition, it is more general than the one in many existing works on last-iterate convergence of SGD and mSGD [1, 21, 16, 15, 7, 13], i.e., $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$, which is obtained from Assumption 3.2 if $\eta_0 = 0$. Under the step size condition in Assumption 3.2, one can choose larger step size like $\epsilon_n = \frac{1}{\sqrt{n}}$, which however is not feasible in the existing condition $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$. Since the new step size can decay more slowly than the existing one, it provides more space for step size fine tinning, such that the algorithm can quickly converge to a neighborhood of the stationary point [23, 24, 22].

Then we attain the first main result in this paper on last-iterate convergence of mSGD with the new step size satisfying Assumption 3.2.

Theorem 3.1. *Consider the mSGD in* (4) *with any* $v_0 \in \mathbb{R}^N$ *and* $\theta_1 \in \mathbb{R}^N$. *Under Assumptions 3.1 and 3.2, there exists a connected set* $J^* \subseteq J$ *such that the iterate* θ_n *is convergent to the set* J^* *almost surely, i.e.,*

$$\lim_{n \to \infty} d(\theta_n, J^*) = 0, \qquad a.s.$$

Technically, there are several big challenges when we try to extend the condition $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$ to $\sum_{n=1}^{+\infty} \epsilon_n^{2+\delta_0} < +\infty$ with $0 \le \eta_0 < 1/2$. One of the challenges is: Usually when we try to prove $\sum_{n=1}^{+\infty} \zeta_n = \sum_{n=1}^{+\infty} \epsilon_n \nabla_{\theta_n} g(\theta_n)^T (\nabla_{\theta_n} g(\theta_n, \xi_n) - \nabla_{\theta_n} g(\theta_n))$ is convergent a.s., which is the vital step to get the last-iterate convergence, we use the $Martingale\ convergence\ theorem$, through which we just need to prove $\sum_{n=1}^{+\infty} \zeta_n^2$ is convergent. If the condition $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$ holds, we can use the following inequality to get $\sum_{n=1}^{+\infty} \zeta_n^2 < \sum_{n=1}^{+\infty} \hat{K} \epsilon_n^2 < +\infty$ ($\hat{K}$ is a constant) directly. However, under the new condition $\sum_{n=1}^{+\infty} \epsilon_n^{2+\delta_0}$, we can not use this approach to prove it. Instead, we introduce a more technical approach to handle this problem.

The result in Theorem 3.1 is quite general, since J^* is a connected set of stationary points. If there is only one stationary point (i.e., $J = J^* = \theta^*$), it follows from Theorem 3.1 that $\theta_n \to \theta^*$. Let $\alpha = 0$ in mSGD, then we can get a similar result for SGD

Corollary 3.1. *Consider the SGD in* (3) *with any* $\theta_1 \in \mathbb{R}^N$. *Under Assumptions 3.1 and 3.2, there exists a connected set* $J^* \subseteq J$ *such that the iterate* θ_n *is convergent to the set* J^* *almost surely, i.e.,*

$$\lim_{n \to \infty} d(\theta_n, J^*) = 0, \qquad a.s.$$

Comparing with the corresponding result in [7] (Theorem 1 in [7]), we relax the condition on the decay of step size. With this new condition, one can utilize step size with slow decay in the iteration of mSGD.

In the following, we aim to show the last-iterate convergence rate under the new step size condition. Since $\|\nabla_{\theta_n} g(\theta_n)\|^2$ is a stochastic variable, it is not favorable to estimate the almost-sure convergence rate of $\nabla_{\theta_n} g(\theta_n)$. Instead, the convergence rate of $\mathbb{E}\left(\|\nabla_\theta g(\theta_n)\|^2\right)$ is quite meaningful. In general, if we need to estimate the convergence rate of the last iterate, we usually need some extra assumptions to establish a quantitative relationship between g and ∇g. Existing works were established usually under the strong-convex assumption [25, 26, 13], local P-L condition [7] or convex loss function with some stability conditions, i.e., $\mathbb{E}_{\xi_n} \|\nabla_{\theta_n} g(\theta_n)\|^2 < \delta < +\infty$. These assumptions are all relatively strong to some extent, especially the stability condition which is very difficult to verify. In our paper, we need a milder assumption than the above conditions as follow.

Assumption 3.3. *Regarding the loss function* $g(\theta)$ *defined in* (1), *there exists* $\delta_0 > 0$, $\hat{T} > 0$, *such that for any* $\theta \in \{\theta | \|\nabla_\theta g(\theta)\|^2 / (g(\theta) - g^*) < \delta_0\}$, *it holds that* $g(\theta) < \hat{T}$, *where* $g^* = \inf_{\theta \in \mathbb{R}^N} g(\theta)$.

This assumption requires the value of the loss function $g(\theta)$ is bounded in the set $S = \{\theta | \|\nabla_\theta g(\theta)\|^2 / (g(\theta) - g^*) < \delta_0\}$. The motivation of this assumption is: For a stochastic algorithm, there is always a probability at which the iterates are far away from the true value. Since second-derivative information will have more positive effect on convergence (such as strongly convex loss functions), this condition restricts iterates which are far away due to extreme noise, so that it does not have significant effect on global updates. This requirement is milder than the strong convexity condition, which is commonly used in former works [25, 26, 13] for the analysis of the last-iterate

168 convergence. In addition, this assumption allows the loss function to be non-convex, and does not
169 need a local P-L condition used in [7] on the second derivative information near the stationary point.
170 At the same time, it can be satisfied by many loss functions commonly encountered in machine
171 learning. First of all, many loss functions usually used are bounded [27–30], and thus satisfy As-
172 sumption 3.3. For unbounded loss functions, there are many instances satisfying Assumption 3.3.
173 For linear regression functions, since they are strongly convex, it is easy to verify the assumption. In
174 the following, we explain (not in a strict manner but an illustrating one) this assumption can also be
175 satisfied by a two layer neural network with $Relu$ active function and square loss function: Given the
176 loss function

$$g(\theta, W) = \frac{1}{N} \sum_{i=1}^{N} \left| y_i - \theta^T \sigma(W x_i) \right|^2.$$

177 we can calculate its partial derivatives with respect to θ and W

$$\frac{\partial g(\theta, W)}{\partial \theta} = \frac{2}{N} \sum_{i=1}^{N} \left(y_i - \theta^T \sigma(W x_i) \right) \sigma(W x_i),$$

$$\frac{\partial g(\theta, W)}{\partial W} = \frac{2}{N} \sum_{i=1}^{N} \left(y_i - \theta^T \sigma(W x_i) \right) (\theta \odot \mathbf{1}_{W x_i \geq 0}) x_i^T.$$

178 For illustration, we just consider the situation which only have one date, i.e. $N = 1$. It holds that

$$\left\| \frac{\partial g(\theta, W)}{\partial(\theta, W)} \right\|^2 = \left\| \frac{\partial g(\theta, W)}{\partial \theta} \right\|^2 + \left\| \frac{\partial g(\theta, W)}{\partial W} \right\|^2 = 4 \left| y - \theta^T \sigma(W x) \right|^2 \|\sigma(W x)\|^2$$

$$+ 4 \left| y - \theta^T \sigma(W x) \right|^2 \|\theta \odot \mathbf{1}_{W x_i \geq 0}\|^2 \|x\|^2 = 4 \left(\|\sigma(W x)\|^2 + \|\theta \odot \mathbf{1}_{W x_i \geq 0}\|^2 \|x\|^2 \right) \left| y - \theta^T \sigma(W x) \right|^2.$$

179 As a result, $\|\nabla_\theta g(\theta)\|^2 / g(\theta) = 4 \left(\|\sigma(W x)\|^2 + \|\theta \odot \mathbf{1}_{W x_i \geq 0}\|^2 \|x\|^2 \right)$. When $4 \left(\|\sigma(W x)\|^2 + \|\theta \odot \right.$
180 $\left. \mathbf{1}_{W x_i \geq 0}\|^2 \|x\|^2 \right) < \delta_0$, we can get $\|\theta^T \sigma(W x)\|^2 \leq 1/2 \left(\|\sigma(W x)\|^2 + \|\theta \odot \mathbf{1}_{W x_i \geq 0}\|^2 \|x\|^2 \right) \leq 2\delta_0$.
181 This ensures that $g(\theta, W)$ bounded.

182 In the following theorem, we provide an estimate of the convergence rate for the mSGD in (4).

183 **Theorem 3.2.** *Consider the mSGD in* (4) *with any* $v_0 \in \mathbb{R}^N$ *under Assumptions 3.1–3.3. Then for*
184 *any* $\|\nabla_{\theta_1} g(\theta_1)\|^2$, *there is*

$$\mathbb{E} \left(\|\nabla_\theta g(\theta_n)\|^2 \right) = O \left(\left(e^{-\frac{2\hat{c}}{(1-\alpha)^2} \sum_{i=1}^{n} \epsilon_i} \right) \left(\sum_{i=1}^{n} \epsilon_i^2 e^{\frac{2\hat{c}}{(1-\alpha)^2} \sum_{k=1}^{i} \epsilon_k} \right) \right), \qquad (7)$$

185 *where* $\hat{c} = \min\{a/2T_0, 2\delta_0\}$.

186 Comparing with the result of the last-iterate convergence rate in [7], we remove the strong
187 growth condition, i.e., $\mathbb{E}_{\xi_n} \|\nabla_\theta g(\theta, \xi_n)\|^2 \leq M \|\nabla_\theta g(\theta)\|^2$, and relax the step size condition to
188 be $\sum_{n=1}^{+\infty} \epsilon_n^{2+\delta_0}$ $(0 < \delta_0 \leq 1/2)$. In the comparison with [20], we do not require the convexity of
189 loss function or $\mathbb{E} \|\nabla_{\theta_n} g(\theta_n)\|^2 < G$. In addition, we relax the step size condition in [20].

190 From (7), we see that larger step size may not lead to larger convergence rate. For example, given
191 two step sizes $\frac{(1-\alpha)^2}{2\hat{c}} \frac{1}{n}$ and $\frac{(1-\alpha)^2}{2\hat{c}} \frac{1}{n^t}$ (*some* $1 > t > 1/2$), they lead to the same convergence rate
192 $O(1/n)$. This is reasonable, since the convergence rate reflects how quickly when $n \to +\infty$ and a
193 larger step size usually makes the algorithm converge to a neighborhood of stationary point quicker.
194 We have the following theorem on this point.

195 **Theorem 3.3.** *Consider the mSGD in* (4) *with any* $v_0 \in \mathbb{R}^N$. *Under Assumptions 3.1–3.3, given any*
196 $a > 0$, *if* $\|\nabla_{\theta_1} g(\theta_1)\|^2 > a$, *then it holds that*

$$P(\tau^{(a)} \geq n) = O \left(e^{-\frac{2\hat{c}}{(1-\alpha)^2} \sum_{i=1}^{n} \epsilon_i} \right),$$

197 *where* $\tau^{(a)} = \min_{n>0} \{ \|\nabla_{\theta_n} g(\theta_n)\|^2 < a \}$ *and* $\hat{c} = \min\{a/2T_0, 2\delta_0\}$.

198 Theorem 3.3 provides a probability description of how fast the iterates of mSGD converge into the
199 preset neighborhood of stationary points. It can be illustrated as follows. It follows from Theorem 3.3
200 that

$$P(\tau_i^{(a_0)} < n) > 1 - \theta_0 e^{-\frac{\delta_0}{(1-\alpha)^2} \sum_{i=1}^{n} \epsilon_i},$$

where $\theta_0 > 0$ is a constant. From the inequality, as n increases, the event $\|\theta_n - \theta^{(i)}\| < a_0$ occurs with higher and higher probability tending to probability 1. The influence of momentum coefficient α and step size ϵ_n is also quantified. We note that with the measure in Theorem 3.3, the iterate θ_n under the step size condition in Assumption 3.2 can converge to a neighbor of the stationary point faster than under the traditional condition $\sum_{n=1}^{+\infty} \epsilon_n^2 < +\infty$ by setting the step size with a relatively slow decay speed as $\sum_{n=1}^{+\infty} \epsilon_n^{2+\delta_0}$ $(0 < \delta_0 \leq 1/2)$, such as $\epsilon_n = \frac{1}{\sqrt{n}}$ and $\eta_0 = 1/4$. Similar results (the large step size can make the iteration quickly converge to a neighbor of stationary point, but may converge slowly in the late stage of the algorithm) appear in [23, 24, 22] through experiments.

3.2 Convergence of mSGD with constant step size

In the previous subsection, we establish the last-iterate convergence of mSGD under a relaxed decay condition on step size. In this subsection, we study the convergence of mSGD under a constant step size $\epsilon_n \equiv \epsilon > 0$. We start with the following result on subsequence convergence.

Theorem 3.4. *Consider the mSGD in* (4) *with any $v_0 \in \mathbb{R}^N$ and $\theta_1 \in \mathbb{R}^N$. Under Assumption 3.1, for any $\varphi > 0$, there exists $\mu_0^{(\varphi)} > 0$, such that for any $\epsilon_n \equiv \epsilon < \mu_0^{(\varphi)}$, it holds that*

$$\left\| \nabla_{\theta_{k_n}^{(\varphi)}} g(\theta_{k_n}^{(\varphi)}) \right\|^2 \leq \varphi, \ a.s.$$

where $\{\theta_{k_n}^{(\varphi)}\}$ is a subsequence $\{\theta_n\}$.

Proof. Due to page constraint, the complete proof is given Appendix. Here, we provide a proof outline to present the main proof ideas.

To study the mSGD with constant step size, we notice the fact that the impact of the noise cannot be totally eliminated through the step size. But when $\|\nabla_{\theta_n} g(\theta_n)\|$ is relatively large (meaning the distance between θ_n and the stationary point θ^* is relatively large), it holds that $\mathbb{E}_{\xi_n} \left(\|\nabla_{\theta_n} g(\theta_n, \xi_n)\|^2 \right) \leq M' \|\nabla_{\theta_n} g(\theta_n)\|^2 + a' \leq (M' + \hat{k}) \|\nabla_{\theta_n} g(\theta_n)\|^2$. This implies that with a proper step size ϵ, the noise can be controlled. As a result, the algorithm can be stabilized around a stationary point. In the following, several key steps in the proof are provided.

Step 1: We define an event by $A_n^{(\varphi)} = \left\{ \|\nabla_{\theta_1} g(\theta_1)\|^2 > \varphi, \ \|\nabla_{\theta_2} g(\theta_2)\|^2 > \varphi, ..., \|\nabla_{\theta_n} g(\theta_n)\|^2 > \varphi \right\}$ and a characteristic function of this event by $I_n^{(\varphi)}$.

Step 2: We prove that the probability of $A_n^{(\varphi)}$ is tending to 0 as $n \to \infty$. First, we attain a recursion formula:

$$\mathbb{E} \left(I_{t+1}^{(\varphi)} g(\theta_{t+1}) \right) - \mathbb{E} \left(I_t^{(\varphi)} g(\theta_t) \right)$$

$$\leq - \alpha^{t-1} I_1^{(\varphi)} \mathbb{E} \left(\nabla_{\theta_1} g(\theta_1)^\mathsf{T} v_1 \right) - \sum_{i=2}^t \alpha^{t-i} (\epsilon - \hat{k} M' \epsilon^2) \mathbb{E} \left(I_{i-1}^{(\varphi)} \|\nabla_{\theta_i} g(\theta_i)\|^2 \right).$$

By taking the sum of multiple inequalities as above, we have

$$\sum_{i=2}^t \mathbb{E} \left(I_{i-1}^{(\varphi)} \|\nabla_{\theta_i} g(\theta_i)\|^2 \right) < +\infty.$$

This leads to

$$\mathbb{E} \left(I_{t-1}^{(\varphi)} \|\nabla_{\theta_t} g(\theta_t)\|^2 \right) \to 0.$$

From the *Chebyshev inequality*, we attain $P(A_n^{(\varphi)}) \to 0$.

Step 3: Step 2 implies that with probability 1 that there exists at least one instant n at which $\|\nabla_{\theta_n} g(\theta_n)\|^2 \leq \varphi$. We note that for the time instant m at which $\|\nabla_{\theta_m} g(\theta_m)\|^2 > \varphi$, then we can use a new start point. By repeating the process of Step 2, we attain the conclusion of this theorem.

$\square$

Theorem 3.4 shows that under the common conditions in the literature, the constant step size can guarantee the subsequence convergence of mSGD (4) in any small neighborhood of stationary point. In addition, we have a similar result on $\mathbb{E} \left(\|\nabla_{\theta_n} g(\theta_n)\|^2 \right)$ as follows.

(a) Three-layer neural network   (b) Four-layer neural network

Figure 1: Training loss-iteration chart on the Boston house-price data.

Proposition 3.1. *Suppose $\{\theta_n\}$ is a sequence generated by* (4) *with any $\theta_1 \in \mathbb{R}^N$ and $v_0 \in \mathbb{R}^N$. Under Assumptions 3.1 and 3.3, if $g(\theta)$ has only one optimal point $g^* = g(\theta^*)$ and $\inf_{d(\theta,\theta^*)>\delta_0} \|\nabla_\theta g(\theta)\|^2 > 0 \ \ (\forall \delta > 0)$, then for any $\varphi_0 > 0$, there exist scalars $\mu_0^{(\varphi_0)} > 0$ and $n_0 > 0$, such that for any $\epsilon_n \equiv \epsilon < \mu_0^{(\varphi_0)}$ and $n > n_0$, it holds that*

$$\mathbb{E}\left(\|\nabla_{\theta_n} g(\theta_n)\|^2\right) \leq \varphi_0.$$

This result is a substantial extension of the existing results on SGD with constant step size to mSGD. Comparing with the results in [22], we remove the requirement of μ-strong convexity on loss function.

4 Experiments

Since this paper focuses on convergence analysis on the well-known mSGD, two simple experiments on regression and classification tasks are given respectively. In the following, we mainly show the results with training datasets. More results with testing datasets are given in Appendix.

4.1 Regression task

In this subsection, we study a house price prediction problem with neural networks trained by mSGD.

Network Architectures. We respectively employ a 3-layer and a 4-layer fully-connected neural network with ReLu active function and squared loss function.

Implementation. We implement two neural networks to train on this dataset using keras. The first neural network consists of three fully connected layers with 13, 8, and 1 neurons. The second neural network consists of four fully connected layers with 13, 32, 16, 1 neurons respectively. We initialize the weights use glorot uniform algorithm. We use mSGD with a mini-batch size of 16, and use mean square error loss function to train the model. The learning rate starts from 0.1 and decay through different algorithms, momentum term coefficient is 0.9, and the models are trained for up to 1000 epochs, training takes about two hours at a time in 3080GPU. We do not use dropout.

Dataset. The training dataset we use is from Boston House Price Dataset. This dataset belongs to the regression prediction dataset, which consists of 506 sets of data with dimension 13. During the training process, we randomly divide 30% as the test set and normalize the data before training.

Results. By using the mSGD in (4) with three different step sizes respectively, i.e., $\epsilon_n = \frac{1}{n}, \epsilon_n = \frac{1}{n^{0.7}}$, and $\epsilon_n = \frac{1}{n^{0.5}}$. The experiment results are given in Figure 1. The figure shows that 1) the loss decays to zero under the three step size setting; 2) $\epsilon_n = \frac{1}{n^{0.5}}$ can make the loss tend to a small neighborhood of zero fastest among the three step size settings. This conforms to the theoretical analysis in Theorems 3.1–3.3.

(a) The training loss   (b) Accuracy of ResNet20 on CIFAR-10

Figure 2: Training and prediction performance on CIFAR-10.

4.2 Classification task

In this subsection, we study a image classification problem with neural networks trained by mSGD.

Network Architectures. We employ a 20-layer ResNet network. The convolutional layers mostly have 3×3 filters and follow two simple design rules: (i) for the same output feature map size, the layers have the same number of filters; (ii) if the feature map size is halved, the number of filters is doubled so as to preserve the time complexity per layer. We perform downsampling directly by convolutional layers that have a stride of 2 and padding with valid. The network ends with an average pooling layer and a 10-way fully-connected layer with softmax.

Implementation. We implement the ResNet20 network using Keras. We initialize the weights use glorot uniform algorithm. We use mSGD with a mini-batch size of 64, momentum coefficient 0.9 and use categorical crossentropy loss function to train the model. The learning rate starts from 0.1 and decays through different algorithms. The models are trained for up to 1000 epochs, taking about two hours at a time in 3080GPU. We do not use dropout.

Dataset. We use two different datasets CIFAR-10 and CIFAR-100. CIFAR-10 consists of 50k training images and 10k testing images in 10 classes. CIFAR-10 is a color image dataset closer to ubiquitous objects, and the image size is 32x32x3. CIFAR-100 consists of 50k training images and 10k testing images in 100 classes. Each category contains 500 training images and 100 testing images.CIFAR-100 is a color image dataset closer to ubiquitous objects,and the image size is 32x32x3. Normalize the dataset to between 0-1 before training.

Results. By using the mSGD in (4) with three different step sizes respectively, i.e., $\epsilon_n = \frac{1}{n}, \epsilon_n = \frac{1}{n^{0.7}}$, and $\epsilon_n = \frac{1}{n^{0.5}}$. The experiment results are given in Figure 3. The figure shows that 1) the loss decays to zero and the accuracy goes up to 1 under the three step size setting; 2) $\epsilon_n = \frac{1}{n^{0.5}}$ can make the loss tend to a small neighborhood of zero and make the accuracy tend to a small neighborhood of 1 fastest among the three step size settings, followed by the setting of $\epsilon_n = \frac{1}{n^{0.7}}$. These results conform to the theoretical analysis in Theorems 3.1–3.3.

5 Conclusion

In this paper, we studied the last-iterate convergence of momentum-based stochastic gradient descent (mSGD) under relaxed conditions on step sizes. First, under the relaxed condition $\sum_{n=1}^{+\infty} \epsilon_n^{2+\eta_0} < +\infty$ $(0 \leq \eta_0 < 1/2)$, we proved the last-iterate convergence of mSGD for non-convex loss functions. In addition, we showed that this step size can indeed help to improve the convergence speed in the early stage of the algorithm by quantifying the influence of the step size and momentum coefficient. This implies that a larger step size, such as $\epsilon_n = \frac{1}{\sqrt{n}}$ can be utilized for accelerating the mSGD. We also proved that the algorithm with a constant step size (i.e., $\epsilon_n \equiv \epsilon > 0$) can ensure the last-iterate convergence of mSGD without requiring the strong convexity assumption on loss function.

(a) The training loss   (b) Accuracy of ResNet20 on CIFAR-100

Figure 3: Training and prediction performance on CIFAR-100.

References

[1] H. Robbins and S. Monro, "A stochastic approximation method," *Annals of Mathematical Statistics*, vol. 22, no. 3, pp. 400–407, 1951.

[2] B. T. Polyak, "Some methods of speeding up the convergence of iteration methods," *USSR Computational Mathematics & Mathematical Physics*, vol. 4, no. 5, pp. 1–17, 1964.

[3] S. Zareba, A. Gonczarek, J. M. Tomczak, and J. Swiatek, "Accelerated learning for restricted Boltzmann machine with momentum term," in *Progress in Systems Engineering*, pp. 187–192, 2015.

[4] I. Sutskever, J. Martens, G. Dahl, and G. E. Hinton, "On the importance of initialization and momentum in deep learning," in *International Conference on Machine Learning*, pp. 1139–1147, 2013.

[5] I. Gitman, H. Lang, P. Zhang, and L. Xiao, "Understanding the role of momentum in stochastic gradient methods," *Advances in Neural Information Processing Systems*, vol. 32, 2019.

[6] A. M. Gupal and L. G. Bazhenov, "Stochastic analog of the conjugate-gradient method," *Cybernetics*, vol. 8, no. 1, pp. 138–140, 1972.

[7] R. Jin, Y. Xing, and X. He, "On the convergence of mSGD and AdaGrad for stochastic optimization," in *arXiv:2201.11204*, 2022.

[8] A. Krizhevsky, I. Sutskever, and G. E. Hinton, "Imagenet classification with deep convolutional neural networks," *Advances in Neural Information Processing Systems*, vol. 25, pp. 1097–1105, 2012.

[9] S. Tang, C. Shen, D. Wang, S. Li, W. Huang, and Z. Zhu, "Adaptive deep feature learning network with Nesterov momentum and its application to rotating machinery fault diagnosis," *Neurocomputing*, vol. 305, pp. 1–14, 2018.

[10] D. Kim, S. Ramani, and J. A. Fessler, "Combining ordered subsets and momentum for accelerated X-ray CT image reconstruction," *IEEE Transactions on Medical Imaging*, vol. 34, no. 1, pp. 167–178, 2014.

[11] M. O. Williams and T. E. Lovett, "Numerical diagnostics for systems of differential algebraic equations," *arXiv preprint arXiv:1602.07550*, 2016.

[12] S. Cyrus, B. Hu, B. Van Scoy, and L. Lessard, "A robust accelerated optimization algorithm for strongly convex functions," in *American Control Conference*, pp. 1376–1381, 2018.

[13] W. An, H. Wang, Q. Sun, J. Xu, Q. Dai, and L. Zhang, "A PID controller approach for stochastic optimization of deep networks," in *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, pp. 8522–8531, 2018.

[14] O. Shamir, "Making gradient descent optimal for strongly convex stochastic optimization," *Omnipress*, 2011.

[15] L. M. Nguyen, P. H. Nguyen, P. Richtárik, K. Scheinberg, M. Taká, and M. V. Dijk, "New convergence aspects of stochastic gradient algorithms," 2018.

[16] L. Nguyen, P. H. Nguyen, M. Dijk, P. Richtárik, K. Scheinberg, and M. Takác, "SGD and Hogwild! convergence without the bounded gradients assumption," in *International Conference on Machine Learning*, pp. 3750–3758, 2018.

[17] B. T. Polyak, "Comparison of convergence rate of one-step and multistep optimization algorithms in the presence of noise," *Izv. Akad. Nauk SSSR, Tekh. Kibern.*, vol. 1, pp. 9–12, 1977.

[18] Y. M. Kaniovskii, "Behaviour in the limit of iterations of the stochastic two-step method," *Zhurnal Vychislitel'noi Matematiki i Matematicheskoi Fiziki*, vol. 23, no. 1, pp. 13–20, 1983.

[19] S. L. Smith, P.-J. Kindermans, C. Ying, and Q. V. Le, "Don't decay the learning rate, increase the batch size," 2018.

[20] X. Li, M. Liu, and F. Orabona, "On the last iterate convergence of momentum methods," 2021.

[21] H.-F. Chen, *Stochastic Approximation and Its Applications*. Springer Science & Business Media, 2006.

[22] Y. Liu, Y. Gao, and W. Yin, "An improved analysis of stochastic gradient descent with momentum," *arXiv preprint arXiv:2007.07989*, 2020.

[23] N. Iyer, V. Thejas, N. Kwatra, R. Ramjee, and M. Sivathanu, "Lrtuner: A learning rate tuner for deep neural networks," 2021.

[24] K. N. S. S. Sastry, "Step size matters in deep learning," 2018.

[25] T. Zhang, "Solving large scale linear prediction problems using stochastic gradient descent algorithms," in *Proceedings of the twenty-first international conference on Machine learning*, p. 116, 2004.

[26] O. Shamir and T. Zhang, "Stochastic gradient descent for non-smooth optimization: Convergence results and optimal averaging schemes," *JMLR.org*, 2012.

[27] L. Yang, Z. Hui, and S. Yuan, "Bayes estimation of parameter of exponential distribution under a bounded loss function," *Research Journal of Mathematics Statistics*, vol. 5, no. 4, pp. 28–31, 2013.

[28] R. Leon and C. Wu, "A theory of performance measures in parameter design," *Statistica Sinica*, pp. 335–358, 1992.

[29] Berger and O. James, "Statistical decision theory and bayesian analysis," , vol. 10.1007/978-1-4757-4286-2, 2004.

[30] M. Towhidi and J. Behboodian, "Minimax estimation of a bounded parameter under some bounded loss functions," *Far East Journal of Theoretical Statistics*, vol. 6, no. 1, pp. 39–48, 2002.