A Non-Asymptotic Moreau Envelope Theory for High-Dimensional Generalized Linear Models

Abstract

We prove a new generalization bound that shows for any class of linear predictors in Gaussian space, the Rademacher complexity of the class and the training error under any continuous loss ℓ can control the test error under all *Moreau envelopes* of the loss ℓ. We use our finite-sample bound to directly recover the "optimistic rate" of Zhou et al. (2021) for linear regression with the square loss, which is known to be tight for minimal ℓ_2-norm interpolation, but we also handle more general settings where the label is generated by a potentially misspecified multi-index model. The same argument can analyze noisy interpolation of max-margin classifiers through the squared hinge loss, and establishes consistency results in spiked-covariance settings. More generally, when the loss is only assumed to be Lipschitz, our bound effectively improves Talagrand's well-known contraction lemma by a factor of two, and we prove uniform convergence of interpolators (Koehler et al. 2021) for all smooth, non-negative losses. Finally, we show that application of our generalization bound using localized Gaussian width will generally be sharp for empirical risk minimizers, establishing a non-asymptotic Moreau envelope theory for generalization that applies outside of proportional scaling regimes, handles model misspecification, and complements existing asymptotic Moreau envelope theories for M-estimation.

1 Introduction

Modern machine learning models often contain more parameters than the number of training samples. Despite the capacity to overfit, training these models without regularization has been empirically shown to achieve good generalization performance (Neyshabur et al. 2015; C. Zhang et al. 2017; Belkin et al. 2019). On the theoretical side, the study of minimal-norm interpolation has revealed fascinating phenomena that challenge traditional understandings of machine learning.

We now have a better understanding of how properties of the data distribution and algorithmic bias can affect generalization in high-dimensional linear regression. For example, data with a spiked covariance structure can ensure that the test error of ridge regression will be approximately constant once the regularization strength is small enough for the model to fit the signal (Zhou et al. 2021; Tsigler and Bartlett 2020), contradicting the classical U-shaped curve expected from arguments about the bias-variance tradeoff. Surprisingly, even when the signal is sparse, the risk of the minimal-ℓ_1 norm interpolator can be shown to converge much slower to the Bayes risk than the minimal-ℓ_2 norm interpolator in the junk feature setting (Chatterji and Long 2021; Koehler et al. 2021). In contrast, the minimal-ℓ_2 norm interpolator fails to achieve consistency in the isotropic setting, while the minimal-ℓ_1 norm interpolator is consistent with sparsity but suffers from an exponentially slow rate in the number of parameters d (G. Wang et al. 2021; Muthukumar et al. 2020). However, we can still achieve the minimax rate with minimal-ℓ_p norm interpolators with p extremely close to 1 (Donhauser et al. 2022).

In fact, many of the intriguing phenomena from the work above may be understood using the norm of a predictor; localized notions of uniform convergence have emerged as essential tools for doing so. Compared to other techniques, uniform convergence analyses can have the benefit of requiring neither particular proportional scaling regimes nor closed-form expressions for the learned model, since only an approximate estimate of its complexity is needed. Despite uniform convergence's potential for wider applicability, work in this area has mostly focused on linear regression settings with the strong assumptions that the conditional expectation of the label is linear with respect to the features, and that the residual has constant variance. In contrast, classical agnostic learning guarantees established by uniform convergence usually need only much weaker assumptions on the data distribution, and apply to a broader range of losses and function classes. For example, Srebro et al. (2010) show that bounds with "optimistic rate" hold generally for any smooth, nonnegative loss, though the hidden logarithmic factor in their result is too loose for explaining noisy interpolation; this was recently addressed by Zhou et al. (2021) in the special case of well-specified linear regression.

In this work, we take a step further towards agnostic interpolation learning and consider a high-dimensional generalized linear model (GLM) setting where the label is generated by a potentially misspecified model. We show a new generalization bound that allows us to use the Moreau envelopes of any continuous loss function as an intermediate tool. By optimizing over the smoothing parameter to balance the approximation and generalization errors, our general Moreau envelope theory yields sharp non-asymptotic generalization bounds in a wide variety of settings. Applying to linear regression with the square loss, we recover the optimistic rate of Zhou et al. (2021) and show that it can more generally be extended to handle model misspecification, such as nonlinear trends and heteroskedasticity. The generality of our result comes from the fact that taking the Moreau envelope of the square loss only scales by a constant; this property alone suffices to obtain a generalization guarantee in terms of the original square loss. The squared hinge loss enjoys the same property, and hence a completely analogous argument shows an optimistic rate in that setting. Combined with an analysis of the margin, we show a novel consistency result for max-margin classifiers.

More generally, we apply the Moreau envelope theory to obtain a generic bound for any Lipschitz loss and smooth, nonnegative loss with sharp constants. Looking specifically at the test error of an Empirical Risk Minimizer (ERM), we show our generalization bound with localized Gaussian width will be asymptotically sharp even when overfitting is not necessarily benign, yielding a version of the asymptotic Moreau envelope framework for analyzing M-estimators (El Karoui et al. 2013; Bean et al. 2013; Donoho and Montanari 2016; Thrampoulidis et al. 2018; Sur and Candès 2019) but for the problem of generalization. Numerical simulations on a variety of feature distributions and label generating processes confirm the wide applicability of our theory.

2 Related Work

The Moreau envelope has been useful in characterizing asymptotic properties of M-estimators in linear models (Bean et al. 2013; El Karoui et al. 2013; Donoho and Montanari 2016; El Karoui 2018; Thrampoulidis et al. 2018) and logistic regression (Sur and Candès 2019; Sur et al. 2019; Candès and Sur 2020; Salehi et al. 2019; Zhao et al. 2022). This theory focuses on estimation and inference under proportional asymptotics, rather than generalization, and does not provide any non-asymptotic result.

For linear regression, Bartlett et al. (2020) identify nearly-matching necessary and sufficient conditions for the consistency of minimal-ℓ_2 norm interpolation; their subsequent work (Tsigler and Bartlett 2020) shows generalization bounds for overparametrized ridge regression. Following their work, Negrea et al. (2020) and Zhou et al. (2020) explore the role of uniform convergence, including showing that uniformly bounding the difference between training error and test error fails to explain interpolation learning. Zhou et al. (2020) argue, however, that uniform convergence *of interpolators* is sufficient to establish consistency in a toy example. Koehler et al. (2021) extend their result to arbitrary data covariance and norm, recovering the benign overfitting conditions of Bartlett et al. (2020) as well as proving novel consistency results for the minimal-ℓ_1 norm interpolator. Based on this uniform convergence framework, G. Wang et al. (2021) establish tight bounds for the minimal-ℓ_1 norm interpolator under a sparse signal with isotropic data. Earlier work (Ju et al. 2020; Chinot et al. 2020; Li and Wei 2021) also studied the minimal-ℓ_1 norm interpolator, without showing consistency. Though the minimal-ℓ_1 norm interpolator suffers from an exponentially slow rate, Donhauser et al. (2022) show the minimal-ℓ_p norm interpolator can achieve faster rates with p close to 1. Zhou et al. (2021) show a risk-dependent ("localized") bound that extends the uniform convergence of

interpolators guarantee to predictors with arbitrary training loss, and used it to establish generalization for regularized estimators such as Ridge and LASSO. Our Moreau envelope theory builds on the techniques developed in this line of work to apply uniform convergence in the interpolation regime.

In terms of requirements on the data distribution, Bartlett et al. (2020) and Tsigler and Bartlett (2020) only require the feature vector to be sub-Gaussian, but assume a well-specified linear model for the conditional distribution of the label. The uniform convergence-based works also assume a well-specified linear model, but the assumptions are more restrictive in the sense that the marginal distribution of the feature needs to be *exactly* Gaussian because their proof techniques rely on the Gaussian Minimax Theorem (GMT). Our Moreau envelope theory's application to linear regression significantly relaxes the assumption on the label generating process, though it is still constrained by the Gaussian data assumption. Shamir (2022) also studies model misspecification in linear regression, but allows non-Gaussian features; and shows that benign overfitting does not necessarily occur in the most general setting, even with a spiked-covariance structure (see their Example 1).

For linear classification, Muthukumar et al. (2021) analyze ℓ_2 max-margin classifier by connecting to minimal-norm interpolation in regression. Similarly, our analysis in the classification case depends on the fact the squared hinge loss goes through the same transformation as the square loss under smoothing by Moreau envelope. Donhauser et al. (2022) prove generalization bounds for ℓ_p max-margin classifiers in the isotropic setting and do not consider the spiked-covariance case. Deng et al. (2021), Montanari et al. (2019) and Liang and Sur (2020) derive exact expressions for the asymptotic prediction risk of the ℓ_2 and $\ell_p, (p \in [1, 2))$ max-margin classifiers respectively. Though their proof techniques rely on GMT, our approaches are drastically different. We use GMT in order to show uniform convergence for a class of predictors and establish a non-asymptotic bound, whereas their results are asymptotic and assume a proportional scaling limit. This is a key distinction, because overfitting usually cannot be benign with proportional scaling (e.g. Proposition 1 of Donhauser et al. 2022). Similar lower bounds have also been shown in the context of linear regression (Muthukumar et al. 2020; G. Wang et al. 2021; Zhou et al. 2020).

Some concurrent works have obtained consistency results for max-margin classification in the spiked covariance setting. In particular, the May 2022 version of the work by Shamir (2022) also studies convergence to the minimum of the squared hinge loss, and obtains consistency under conditions similar to the benign covariance condition of Bartlett et al. (2020). During preparation of this manuscript we learned of concurrent work by Montanari et al., not yet publicly available, which also studies consistency results for classification. Comparing our Corollary 3 to Shamir (2022), their result applies to some non-Gaussian settings, but in the Gaussian setting their result is not as general as ours. (Combining Assumptions 1 and 2 of Theorem 7 there, they require that the norm of the data stays bounded, whereas our Corollary 3 applies when the first $o(n)$ eigenvalues of Σ diverge with n arbitrarily fast.) More conceptually, our result follows from a norm-based generalization bound that applies to all predictors and outside of the "benign overfitting" conditions, generalizing the result in (Koehler et al. 2021) and unlike the analysis in prior works.

3 Problem Formulation

GLM fomulation. Given a continuous loss function $f : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ and i.i.d. sample pairs (x_i, y_i) from some data distribution $\mathcal{D}$, we can learn a linear model $(\hat{w}, \hat{b})$ by minimizing the empirical loss $\hat{L}_f$ with the goal of achieving small population loss L_f:

$$\hat{L}_f(w, b) = \frac{1}{n} \sum_{i=1}^{n} f(\langle w, x_i \rangle + b, y_i), \quad L_f(w, b) = \mathbb{E}_{(x,y) \sim \mathcal{D}} f(\langle w, x \rangle + b, y). \quad (1)$$

Multi-index model. We assume that the data distribution $\mathcal{D}$ over (x, y) is given by

1. $x \sim \mathcal{N}(0, \Sigma)$ is a centered Gaussian with unknown covariance matrix Σ.

2. There are unknown weight vectors $w_1^*, ..., w_k^* \in \mathbb{R}^d$, a function $g : \mathbb{R}^{k+1} \to \mathbb{R}$, and a random variable ξ independent of x (not necessarily Gaussian) such that

$$\eta_i = \langle w_i^*, x \rangle, \quad y = g(\eta_1, ..., \eta_k, \xi). \quad (2)$$

We can assume that the distribution of x is centered without loss of generality since presence of a mean term simply corresponds to changing the bias term b: $\langle w, x \rangle + b = \langle w, x - \mu \rangle + (b - \langle w, \mu \rangle)$.

We can also assume that $\Sigma^{1/2}w_1^*, ..., \Sigma^{1/2}w_k^*$ are orthonormal without loss of generality since we have not imposed any assumption on the link function g. The multi-index model includes well-specified linear regression, by setting $k = 1$ and $g(\eta, \xi) = \eta + \xi$. It also allows nonlinear trends and heteroskedasticity (such as the model in Figure 1) by changing the definition of g. Since g need not be continuous, the label y can be binary, as in linear classification.

4 Moreau Envelope Generalization Theory

Our theory vitally depends on the Moreau envelope, defined as follows.

Definition 1. The *Moreau envelope* of $f : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ with parameter $\lambda \in \mathbb{R}^+$ is defined as the function $f_\lambda : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ given by

$$f_\lambda(\hat{y}, y) = \inf_u \, f(u, y) + \lambda(u - \hat{y})^2. \tag{3}$$

The Moreau envelope can be viewed as a smooth approximation to the original function f: in our parameterization, smaller λ corresponds to more smoothing. The map that outputs the minimizer u, known as the *proximal operator*, plays an important role in convex analysis (Parikh and Boyd 2014; Bauschke, Combettes, et al. 2011).

Our general theory, as stated in Theorem 1 below, essentially upper bounds the generalization gap between the population Moreau envelope L_{f_λ} and the original training loss $\hat{L}_f$ by the sum of two parts: a parametric component that can be controlled by the dimension, and a non-parametric component that can be controlled by a dimension-free complexity measure such as the Euclidean norm of the predictor. Typically, the first term is negligible since the dimension is small, and the complexity of fitting all the noise is absorbed into the second term. More precisely, we introduce the following definitions to formalize separating out a low dimensional component:

Definition 2. Under the model assumptions (2), define a projection matrix Q onto the space orthogonal to $w_1^*, ..., w_k^*$ and a mapping ϕ from $\mathbb{R}^d$ to $\mathbb{R}^{k+1}$ by

$$Q = I_d - \sum_{i=1}^k w_i^*(w_i^*)^T\Sigma, \quad \phi(w) = (\langle w, \Sigma w_1^*\rangle, ..., \langle w, \Sigma w_k^*\rangle, \|\Sigma^{1/2}Qw\|_2)^T. \tag{4}$$

We let $\Sigma^\perp = Q^T\Sigma Q$ denote the covariance matrix of $Q^T x$. We also define a low-dimensional *surrogate distribution* $\tilde{\mathcal{D}}$ over $\mathbb{R}^{k+1} \times \mathbb{R}$ by

$$\tilde{x} \sim \mathcal{N}(0, I_{k+1}) \quad \text{and} \quad \tilde{y} = g(x_1, ..., x_k, \xi). \tag{5}$$

Our main result is as follows:

Theorem 1. *Suppose $\lambda \in \mathbb{R}^+$ satisfies that for any $\delta \in (0, 1)$, there exists a continuous function $\epsilon_{\lambda,\delta} : \mathbb{R}^{k+1} \to \mathbb{R}$ such that with probability at least $1 - \delta/4$ over independent draws $(\tilde{x}_i, \tilde{y}_i)$ from the surrogate distribution $\tilde{\mathcal{D}}$ defined in (5), we have uniformly over all $(\tilde{w}, \tilde{b}) \in \mathbb{R}^{k+2}$ that*

$$\frac{1}{n}\sum_{i=1}^n f_\lambda(\langle \tilde{w}, \tilde{x}_i\rangle + \tilde{b}, \tilde{y}_i) \geq \mathop{\mathbb{E}}_{(\tilde{x}, \tilde{y})\sim\tilde{D}} [f_\lambda(\langle \tilde{w}, \tilde{x}\rangle + \tilde{b}, \tilde{y})] - \epsilon_{\lambda,\delta}(\tilde{w}, \tilde{b}). \tag{6}$$

Further, assume that for any $\delta \in (0, 1)$, there exists a continuous function $C_\delta : \mathbb{R}^d \to [0, \infty]$ such that with probability at least $1 - \delta/4$ over $x \sim \mathcal{N}(0, \Sigma)$, uniformly over all $w \in \mathbb{R}^d$,

$$\langle Qw, x\rangle \leq C_\delta(w). \tag{7}$$

Then it holds with probability at least $1 - \delta$ that uniformly over all $(w, b) \in \mathbb{R}^{d+1}$, we have

$$L_{f_\lambda}(w, b) \leq \hat{L}_f(w, b) + \epsilon_{\lambda,\delta}(\phi(w), b) + \frac{\lambda C_\delta(w)^2}{n}. \tag{8}$$

If we additionally assume that (6) holds uniformly for all $\lambda \in \mathbb{R}^+$, then (8) does as well.

As we will see in the application sections later, we can generally bound the difference between L_{f_λ} and L_f when the loss is assumed to be Lipschitz. If f is not Lipschitz but smooth, we can always write it as the Moreau envelope of another function $\tilde{f}$. In the special case of square loss or squared hinge loss, the Moreau envelope f_λ is proportional to f, meaning that (8) becomes a generalization guarantee in terms of L_f. Optimizing over λ will establish optimal bounds that recover the result of Koehler et al. (2021) and Zhou et al. (2021), and lead to many other novel contributions.

Remark 1. The complexity functional $C_\delta(w)$ should be thought of as a localized, high-probability version of Rademacher complexity. This is because the Gaussian width of a convex set $\mathcal{K}$, $\mathbb{E} \sup_{w \in \mathcal{K}} \langle w, x \rangle$, is the same as the Rademacher complexity of the class of linear functions $\{x \mapsto \langle w, x \rangle : w \in \mathcal{K}\}$ (Zhou et al. 2021, Proposition 1). A somewhat similar complexity functional appears in Panchenko (2003). Also, note (6) requires only *one-sided concentration* — see Remark 3.

4.1 VC Theory for Low-dimensional Concentration

To apply our generalization result (Theorem 1), we should check the low-dimensional concentration assumption (6). The quantitative bounds in the low-dimensional concentration (i.e. the precise form of error term $\epsilon_{\lambda, \delta}$) will inevitably depend on the exact setting we consider (see e.g. Vapnik 1982; Koltchinskii and Mendelson 2015; Lugosi and Mendelson 2019 for discussion).

First, we recall the following result from VC theory.

Theorem 2 (Special Case of Assertion 4 of Vapnik 1982, Chapter 7.8; see also Theorem 7.6)**.** *Suppose that $\mathcal{K} \subset \mathbb{R}^d$ and $\mathcal{B} \subset \mathbb{R}$, and $(x, y) \sim \mathcal{D}$ are random variables with x valued in $\mathbb{R}^d$ such that for some $\tau > 0$, for all $w \in \mathcal{K}, b \in \mathcal{B}$*

$$\frac{\left(\mathbb{E} f(\langle w, x \rangle + b, y)^4]\right)^{1/4}}{\mathbb{E} f(\langle w, x \rangle + b, y)} \leq \tau. \tag{9}$$

Suppose that the class of functions $\{(x, y) \mapsto \mathbb{1}\{f(\langle w, x \rangle + b, y) > t\} : w \in \mathcal{K}, b \in \mathcal{B}, t \in \mathbb{R}\}$ has VC dimension at most h. Then with probability at least $1 - \delta$, for $(x_1, y_1), \ldots, (x_n, y_n) \sim \mathcal{D}$ i.i.d.,

$$\frac{1}{n} \sum_{i=1}^n f(\langle w, x_i \rangle + b, y_i) \geq \left(1 - 8\tau \sqrt{\frac{h(\log(2n/h) + 1) + \log(12/\delta)}{n}}\right) \mathbb{E} f(\langle w, x \rangle + b, y)$$

uniformly over $w \in \mathcal{K}, b \in \mathcal{B}$.

The assumption (9) is standard (indeed, this is the setting primarily focused on in Vapnik 1982) and is sometimes referred to as *hypercontractivity* or *norm equivalence* in the literature; a variant of the result holds with 4 replaced by $1 + \epsilon$. In many settings of interest, this can be directly checked using the fact that x is Gaussian. Of course, our general result can be applied without this assumption, by using low-dimensional concentration under an alternative assumption: Vapnik (1982), Panchenko (2002), Panchenko (2003), and Mendelson (2017) have further discussion and alternative results. Combining the above result with Theorem 1 yields the following.

Corollary 1. *Let $\mathcal{K} \subset \mathbb{R}^d$ and $\mathcal{B} \subset \mathbb{R}$ be arbitrary. In the setting of Theorem 1, suppose that the surrogate distribution $\tilde{\mathcal{D}}$ satisfies assumption (9) uniformly over $(\phi(w), b)$ for $w \in \mathcal{K}, b \in \mathcal{B}$ and for fixed $\lambda \geq 0$ the class $\{(x, y) \mapsto \mathbb{1}\{f_\lambda(\langle \tilde{w}, \phi(x) \rangle + \tilde{b}, y) > t\} : (\tilde{w}, \tilde{b}) \in \mathbb{R}^{k+2}, t \in \mathbb{R}\}$ has VC dimension at most h. Then with probability at least $1 - \delta$,*

$$\left(1 - 8\tau \sqrt{\frac{h(\log(2n/h) + 1) + \log(48/\delta)}{n}}\right) L_{f_\lambda}(w, b) \leq \hat{L}_f(w, b) + \frac{\lambda C_\delta(w)^2}{n}.$$

Furthermore, if the class $\{(x, y) \mapsto \mathbb{1}\{f_\lambda(\langle \tilde{w}, \phi(x) \rangle + \tilde{b}, y) > t\} : (\tilde{w}, \tilde{b}) \in \mathbb{R}^{k+2}, t \in \mathbb{R}, \lambda \in \mathbb{R}_{\geq 0}\}$ has VC dimension at most h, then the same conclusion holds uniformly over λ.

The last conclusion (uniformity over λ) follows by going through the proof of Theorem 2, since it is based on reduction to uniform control of indicators. In every situation we will consider, it is easy to check that the VC dimension h in the theorem statement is $O(k)$, generally by reducing to the fact that halfspaces in $\mathbb{R}^k$ have VC dimension $k + 1$.

5 Applications

5.1 Linear Regression with Square Loss

In this section, we show how to recover optimistic rates (Zhou et al. 2021) for linear regression without assuming the model is well-specified. We will consider the square loss, $f(\hat{y}, y) = (\hat{y} - y)^2$. A key property of the square loss is that the Moreau envelope is proportional to itself:

$$f_\lambda(\hat{y}, y) = \inf_u (u - y)^2 + \lambda(u - \hat{y})^2 = \frac{\lambda}{1 + \lambda} f(\hat{y}, y). \tag{10}$$

218 Thus we can multiply by $(1 + \lambda)/\lambda$ in our generalization bound and solve for the optimal choice of λ.

219 **Corollary 2.** *Suppose f is the square loss and the surrogate distribution $\tilde{\mathcal{D}}$ satisfies assumption* (9)
220 *uniformly over $(w, b) \in \mathbb{R}^{k+1}$, then with probability at least $1 - \delta$, uniformly over all w, b we have*

$$\left(1 - 8\tau\sqrt{\frac{k(\log(2n/k) + 1) + \log(48/\delta)}{n}}\right) L_f(w, b) \leq \left(\sqrt{\hat{L}_f(w, b)} + C_\delta(w)/\sqrt{n}\right)^2.$$

221 As mentioned earlier, assumption (9) usually holds under mild conditions on y. For example,
222 τ can be chosen to be a constant when y is a bounded-degree polynomial of a Gaussian due to
223 Gaussian hypercontractivity (O'Donnell 2014, Section 11.1). Specializing Corollary 2 to interpolators
224 ($\hat{L}_f = 0$) recovers the uniform convergence of interpolators guarantee from Koehler et al. (2021).
225 Combined with a more general norm analysis in Section 6, we establish ℓ_2 benign overfitting with
226 misspecification. In the well-specified case, see Zhou et al. (2021) for detailed examples on ordinary
227 least squares, ridge regression, and LASSO.

5.2 Classification with Squared Hinge Loss

229 In this section, we show a novel optimistic rate bound for max-margin linear classification with the
230 squared hinge loss, $f(\hat{y}, y) = \max(0, 1 - y\hat{y})^2$. Its Moreau envelope is given by

$$f_\lambda(\hat{y}, y) = \inf_u \max(0, 1 - yu)^2 + \lambda(u - \hat{y})^2 = \begin{cases} 0 & \text{if } 1 - y\hat{y} \leq 0 \\ \frac{\lambda}{1+\lambda}(1 - y\hat{y})^2 & \text{if } 1 - y\hat{y} > 0 \end{cases} = \frac{\lambda}{1 + \lambda} f(\hat{y}, y).$$

231 We consider the case of a general binary response y valued in $\{\pm 1\}$ satisfying the model assumptions
232 in equation (2). In this case as well, f is proportional to its Moreau envelope; thus, the same proof as
233 for the squared loss shows that Corollary 2 continues to hold when square loss is replaced by squared
234 hinge loss! In Appendix E.3, we discuss certain settings (including noisy settings) where minimizing
235 the squared hinge loss also minimizes the zero-one loss, i.e. the misclassification rate.

5.3 Further applications

237 In this section, we discuss further interesting examples where our general theory applies. As before,
238 we can obtain a generalization bound by appealing to the general Corollary 1: we omit stating the
239 formal corollary for each case, and simply show the Moreau envelope and its consequences.

240 L_1 **loss (LAD) and Hinge.** For the L_1 loss $f(\hat{y}, y) = |\hat{y} - y|$ its Moreau envelope is given by

$$f_\lambda(\hat{y}, y) = \inf_u |u - y| + \lambda(u - \hat{y})^2 = \begin{cases} \lambda(\hat{y} - y)^2 & \text{if } |\hat{y} - y| \leq \frac{1}{2\lambda} \\ |\hat{y} - y| - \frac{1}{4\lambda} & \text{if } |\hat{y} - y| > \frac{1}{2\lambda} \end{cases},$$

241 which is 2λ times the Huber loss with parameter $\delta = \frac{1}{2\lambda}$. Therefore, the population Huber loss is
242 controlled by the empirical L_1 loss. Clearly, interpolators have zero training error under both L_1 and
243 L_2 training losses. We already see that Corollary 2 implies $(1 - o(1)) \mathbb{E}(\langle w, x \rangle + b - y)^2 \leq C_\delta(w)^2/n$.
244 Now, considering Corollary 1 with f the L_1 loss and using the above Moreau envelope calculation,
245 we can check that when $\lambda \to 0$ we reproduce the exact same bound, since the Huber loss becomes
246 the squared loss in the limit. Further insight into this phenomenon appears later in Theorem 4: the
247 Huber loss naturally shows up when computing the training error of the LAD estimator. An entirely
248 analogous situation occurs when f is the hinge loss $f(\hat{y}, y) := \max(0, 1 - \hat{y}y)$: its Moreau envelope
249 f_λ will be a rescaling of the Huber hinge loss (c.f. T. Zhang 2004).

250 **Lipschitz loss: improved contraction.** If f is M-Lipschitz in $\hat{y}$, Proposition 3.4 of Planiden and
251 X. Wang (2019) gives that $0 \leq f - f_\lambda \leq \frac{M^2}{4\lambda}$. Thus, assuming $k/n = o(1)$, Corollary 1 implies
252 $(1 - o(1))(L_f(w) - \frac{M^2}{4\lambda}) \leq \hat{L}_f(w) + \frac{\lambda C_\delta(w)^2}{n}$; optimizing over λ yields

$$(1 - o(1))L_f(w) \leq \hat{L}_f(w) + M\sqrt{\frac{C_\delta(w)^2}{n}}. \tag{11}$$

253 For $w \in \mathcal{K}$, a high-probability upper bound on the Rademacher complexity of the function class
254 $x \mapsto \langle w, x \rangle$ upper bounds the second term (Remark 1). In comparison, the standard symmetrization

Figure 1: Top: ridge regression with model misspecification; bottom: ℓ_2 margin classification with a logistic model. Here the features are Gaussian; see Appendix B for additional experiments on ℓ_1 regularization and non-Gaussian features. The covariance in the first column (isotropic) is $\Sigma = I_d$, in the second column (junk features) is $\Sigma = \mathrm{diag}(1, ..., 1, 0.05^2, ..., 0.05^2)$, and in the third (harmful features) is $\Sigma = \mathrm{diag}(1, ..., 1, \frac{1}{(k+1)^2}, ..., \frac{1}{d^2})$. For regression, the number of leading eigenvalues is $k = 3$, and the label is generated according to $y = 1.5\, x_1 + |x_1| \cos x_2 + x_3 \cdot \mathcal{N}(0, 0.5)$. For classification, the number of leading eigenvalues is $k = 1$ and $\mathrm{Pr}(y = 1 \mid x) = \mathrm{sigmoid}(5x_1 + 3)$. The risk bound is calculated using the expression $\left(\sqrt{\hat{L}} + \sqrt{\|w\|_2^2\, \mathrm{Tr}(\Sigma^\perp)/n}\right)^2$, which corresponds to the choice of $C(w)$ from Lemma 1, and is expected to be tight in the junk features setting. In the isotropic case, we use an easy improvement of the bound where w is projected to the orthogonal complement of the Bayes predictor w^* (Zhou et al. 2021). In the other cases, we use covariance splitting without projection of w. Each point on the curve is the average from repeated experiments, and shaded areas correspond to 95% bootstrap confidence intervals.

and contraction argument (Bartlett and Mendelson 2002) loses a factor of two. Note that if f is the L_1 loss, this applies with $M = 1$, and it can further be shown that the constant factor cannot be improved further (see Appendix E.4), but this bound is also not as tight as the version with Huber test loss.

Uniform Convergence of Interpolators for Smooth Losses. Suppose that f is an H-smooth (in the sense that $|\frac{\partial^2 f}{\partial \hat{y}^2}| \leq H$) and convex function of $\hat{y}$. In addition, assume that the minimum of $f(\hat{y}, y)$ is zero for any fixed y. Since f is H-smooth, there exists a function $\tilde{f}$ such that $f = \tilde{f}_{H/2}$ (Bauschke, Combettes, et al. 2011, Corollary 18.18). If w satisfies $\hat{L}_f(w) = 0$, then $\hat{L}_{\tilde{f}}(w) = 0$ as well, because the Moreau envelope preserves minima (Parikh and Boyd 2014, Chapter 3.1). Finally, by Corollary 2, if $k/n = o(1)$ we have uniformly over all w such that $\hat{L}_f(w) = 0$ that

$$(1 - o(1))L_f(w) \leq \frac{H}{2} \cdot \frac{C_\delta(w)^2}{n}, \tag{12}$$

which generalizes the main result of Koehler et al. (2021) to arbitrary smooth losses.

6 ℓ_2 Benign Overfitting

We can obtain conditions for consistency, like those of Bartlett et al. (2020), by simply combining a norm-based uniform generalization bound with an upper bound on the norm of interpolators. Koehler et al. (2021) used the same strategy, but our more powerful and general tools give better results:

1. For squared loss regression, we show that the assumption that the ground truth is generated by a linear model (i.e. well-specified), made in previous work, is not required. The same result holds under the much more general model[1] assumption of Section 3.

2. We show an analogous result in the *classification* setting, replacing the squared loss with the squared hinge loss. In fact, the argument is exactly the same in the two cases: all we need to use is that $f_\lambda = \frac{\lambda}{1+\lambda} f$ and that f is the square of a Lipschitz function.

First, the following lemma (essentially just the standard Rademacher complexity bound for the ℓ_2 ball) combines with Corollary 2 and its squared hinge loss version to give generalization bounds. These bounds are demonstrated in Figure 1 and Appendix B.

Lemma 1. *In the setting of Theorem 1, letting $\Sigma^\perp = Q^T \Sigma Q$, the following $C_\delta(w)$ will satisfy (7):*

$$C_\delta(w) = \|w\|_2 \left[\sqrt{\mathrm{Tr}(\Sigma^\perp)} + 2\sqrt{\|\Sigma^\perp\|_{op} \log(8/\delta)} \right].$$

Next, we provide a sufficient condition for a zero-training error predictor w to exist in an ℓ_2 ball. In the case of classification, this allows us to lower-bound the margin of the max-margin halfspace.

Definition 3 (Bartlett et al. 2020). The *effective ranks* of a covariance matrix Σ are

$$r(\Sigma) = \frac{\mathrm{Tr}(\Sigma)}{\|\Sigma\|_{op}} \quad \text{and} \quad R(\Sigma) = \frac{\mathrm{Tr}(\Sigma)^2}{\mathrm{Tr}(\Sigma^2)}.$$

Lemma 2. *Suppose that $f(\hat{y}, y)$ is either squared loss or squared hinge loss. Let $(w^\sharp, b^\sharp) \in \mathbb{R}^{d+1}$ be an arbitrary vector satisfying $Qw^\sharp = 0$ and with probability at least $1 - \delta/4$,*

$$\hat{L}_f(w^\sharp, b^\sharp) \le L_f(w^\sharp, b^\sharp) + \rho_1(w^\sharp, b^\sharp) \tag{13}$$

for some $\rho_1(w^\sharp, b^\sharp) > 0$. Then for any $\rho_2 \in (0, 1)$, provided $\Sigma^\perp = Q^T \Sigma Q$ satisfies

$$R(\Sigma^\perp) = \Omega\left(\frac{n \log^2(4/\delta)}{\rho_2} \right), \tag{14}$$

we have that with probability at least $1 - \delta$ that $\min_{\|w\| \le B} L_f(w, b^\sharp) = 0$ for $B > 0$ defined by
$B^2 = \|w^\sharp\|_2^2 + (1 + \rho_2)\frac{n}{\mathrm{Tr}(\Sigma_2^\perp)}(L_f(w^\sharp, b^\sharp) + \rho_1)$.

We note that for any vector $w^\sharp$, we have $L_f((I-Q)w^\sharp, b^\sharp) < L_f(w^\sharp, b^\sharp)$ by Jensen's inequality over $Q^T x$, so the assumption $Qw^\sharp = 0$ in the lemma is always satisfied for the minimizer of $L_f(w, b)$. Combining the norm bound Lemma 2 and the generalization bound Lemma 1 yields the following.

Theorem 3. *Let $(\hat{w}, \hat{b}) = \arg\min_{w \in \mathbb{R}^d, b \in \mathbb{R} : \hat{L}_f(w,b)=0} \|\hat{w}\|_2$ be the minimum-ℓ_2 norm predictor with zero training error. In the setting of Lemma 2, we have*

$$L_f(\hat{w}, \hat{b}) - \epsilon_\delta(\phi(\hat{w}), \hat{b}) \le (1 + \rho_3) \inf_{w^\sharp \in \mathbb{R}^d, b^\sharp \in \mathcal{B}} \left(L_f(w^\sharp, b^\sharp) + \rho_1(w^\sharp, b^\sharp) + \frac{\|w^\sharp\|_2^2 \, \mathrm{Tr}(\Sigma^\perp)}{n} \right),$$

where $\rho_3 > 0$ is defined by $1 + \rho_3 = (1 + \rho_2)\left[1 + 2\sqrt{\frac{\log(2/\delta)}{r(\Sigma^\perp)}}\right]^2$ and we recall $\rho_1(w^\sharp, b^\sharp)$ from (13).

We now show that this formally implies convergence to the optimal test loss (i.e. consistency) under the ℓ_2 benign overfitting conditions (15) from Bartlett et al. (2020) and Tsigler and Bartlett (2020):

Corollary 3. *Suppose that $\mathcal{D}_n$ is a sequence of data distributions following our model assumptions (2), with k_n such that $y = g(\eta_1, \ldots, \eta_{k_n}, \xi)$, and projection operator Q_n defined as in (4). Suppose f is either the squared loss or the squared hinge loss, and define $(w_n^\sharp, b_n^\sharp) = \arg\min_{w,b} L_{f,n}(w, b)$ where $L_{f,n}(w, b)$ is the population loss over distribution $\mathcal{D}_n$ with loss f. Suppose that the hypercontractivity assumption (9) holds with some fixed $\tau > 0$ for all $\mathcal{D}_n$. Define $\Sigma_n := \mathbb{E}_{\mathcal{D}_n}[xx^T]$ and $\Sigma_n^\perp = Q_n^T \Sigma_n Q_n$. Suppose that as $n \to \infty$, we have*

$$\frac{n}{R(\Sigma_n^\perp)} \to 0, \quad \frac{\|w_n^\sharp\|_2^2 \, \mathrm{Tr}(\Sigma_n^\perp)}{n} \to 0, \quad \frac{k_n}{n} \to 0. \tag{15}$$

---

[1]Theorem 3 of concurrent work (Shamir 2022) also proves benign overfitting for some misspecified models, but requires the very strong assumption $n^2 \|\Sigma^\perp\|_{OP} \to 0$ that fails to hold in several examples from Bartlett et al. 2020.

Then we have the following convergence in probability, as $n \to \infty$:

$$\frac{L_{f,n}(\hat{w}_n, \hat{b}_n)}{L_{f,n}(w_n^\sharp, b_n^\sharp)} \to 1, \tag{16}$$

where $(\hat{w}_n, \hat{b}_n) = \arg\min_{w \in \mathbb{R}^d, b \in \mathbb{R}: \hat{L}_f(w,b)=0} \|w\|_2$ is the minimum-norm interpolator, and $\hat{L}_{f,n}$ is the training error based on n i.i.d. samples from the distribution $\mathcal{D}_n$.

Note when applying Corollary 3, we have the flexibility to increase k_n and shrink $\Sigma_n^\perp$ by choosing additional weights w_i^* and letting the link function g ignore the extra components.

Remark 2 (Flatness of the test loss along regularization path). Our method can easily show a slightly stronger statement: let $(\hat{w}_n, \hat{b}_n) \in \arg\min_{\|w\| \leq B_n, b \in \mathbb{R}} \hat{L}_{f,n}(w_n^\sharp, b_n^\sharp)$ such that, if there are multiple minima, we pick the one with smallest $\|w\|$. As long as $B_n \geq \|w_n^\sharp\|$, we still have (16), and this is established uniformly over all sequences B_n satisfying the constraint. Therefore, under the benign overfitting conditions we get consistency as long as we do not over-regularize the predictor. See Figure 1 for an experimental demonstration of the flatness.

7 Training Error and Local Gaussian Width

Theorem 1 shows how to upper-bound the test error of a predictor (under the Moreau envelope loss) by its training error and an upper bound on the class complexity. The following theorem is the dual result, which upper-bounds the training error of the constrained ERM (Empirical Risk Minimizer) by the Moreau envelope and a complexity term. In particular, this general result is used to derive the norm bound for interpolators in Lemma 2 above.

Theorem 4. Let $\mathcal{K}, \mathcal{B}$ be bounded convex sets, and let $f(\hat{y}, y)$ be convex in $\hat{y}$. Suppose that τ is such that with probability at least $1 - \delta$, for $(\tilde{x}, \tilde{y})_{i=1}^n$ sampled i.i.d. from $\tilde{\mathcal{D}}$ we have

$$\min_{\tilde{w} \in \phi(\mathcal{K}), b_0 \in \mathcal{B}} \max_{\lambda \geq 0} \left[\frac{1}{n} \sum_{i=1}^n f_\lambda(\langle \tilde{w}, \tilde{x} \rangle + b_0, y_i) - \frac{\lambda}{n} \max_{w_0 \in \phi^{-1}(\tilde{w}) \cap \mathcal{K}} \langle x, Q w_0 \rangle^2 \right] \leq \tau. \tag{17}$$

Then with probability at least $1 - 2\delta$, $\min_{w \in \mathcal{K}, b \in \mathcal{B}} \hat{L}_f(w,b) \leq \tau$.

Note that the assumption (17) implicitly suggests a low-dimensional concentration assumption: we expect $\frac{1}{n} \sum_{i=1}^n f_\lambda(\langle \tilde{w}, \tilde{x} \rangle + b_0, y_i)$ to be approximately the test loss of $(\tilde{w}, b_0)$ under the surrogate distribution $\tilde{\mathcal{D}}$. As we discuss more in Appendix F, combining this training error bound with the correct choice of $C_\delta(w)$ in Theorem 1, which is essentially $C_\delta(w) = \mathbb{E} \max_{w_0 \in \phi^{-1}(\tilde{w}) \cap \mathcal{K}} \langle x, Q w_0 \rangle^2$), yields a matching lower bound to (8) on the Moreau envelope test loss and so our generalization bound is asymptotically sharp. This establishes a non-asymptotic analogue of the existing asymptotic Moreau envelope theory (see Section 2), and recovers the special case of well-specified linear models (Zhou et al. 2021).

8 Discussion

In this work, we significantly extend the localized uniform convergence technique developed in the study of noisy interpolation to any loss function and label generating process under mild conditions. Though the application of Moreau envelope to study GLMs is not new in the statistical literature, our general theory establishes novel non-asymptotic generalization bounds in a wide variety of overparameterized settings. We believe the generality of our framework may allow further applications in other areas of statistics, such as robust statistics and high-dimensional inference.

As mentioned in Section 2, the applicability of our theory is still considerably limited by the Gaussian data assumption, although it appears experimentally that it may hold much more broadly (Appendix B). Relaxing the Gaussian data assumption could allow us to study kernel methods and bring us closer to a theoretical understanding of deep neural networks. The GMT formulation also does not allow for multi-class classification or two-layer networks, because of their vector-valued outputs. Overcoming these two challenges seems to be crucial avenues for future work.

References

Bartlett, Peter L., Philip M. Long, Gábor Lugosi, and Alexander Tsigler (2020). "Benign overfitting in linear regression." *Proceedings of the National Academy of Sciences* 117.48, pp. 30063–30070. arXiv: 1906.11300.

Bartlett, Peter L. and Shahar Mendelson (2002). "Rademacher and Gaussian complexities: Risk bounds and structural results." *Journal of Machine Learning Research* 3.Nov, pp. 463–482.

Bauschke, Heinz H, Patrick L Combettes, et al. (2011). *Convex analysis and monotone operator theory in Hilbert spaces*. Vol. 408. Springer.

Bean, Derek, Peter J Bickel, Noureddine El Karoui, and Bin Yu (2013). "Optimal M-estimation in high-dimensional regression." *Proceedings of the National Academy of Sciences* 110.36, pp. 14563–14568.

Belkin, Mikhail, Daniel Hsu, Siyuan Ma, and Soumik Mandal (2019). "Reconciling modern machine learning practice and the bias-variance trade-off." *Proceedings of the National Academy of Sciences* 116.32, pp. 15849–15854. arXiv: 1812.11118.

Candès, Emmanuel J and Pragya Sur (2020). "The phase transition for the existence of the maximum likelihood estimate in high-dimensional logistic regression." *The Annals of Statistics* 48.1, pp. 27–42.

Chatterji, Niladri S. and Philip M. Long (2021). "Foolish Crowds Support Benign Overfitting." arXiv: 2110.02914.

Chinot, Geoffrey, Matthias Löffler, and Sara van de Geer (2020). "On the robustness of minimum-norm interpolators." arXiv: 2012.00807.

Deng, Zeyu, Abla Kammoun, and Christos Thrampoulidis (2021). "A model of double descent for high-dimensional binary linear classification." *Information and Inference: A Journal of the IMA*. arXiv: 1911.05822.

Donhauser, Konstantin, Nicolo Ruggeri, Stefan Stojanovic, and Fanny Yang (2022). "Fast rates for noisy interpolation require rethinking the effects of inductive bias." arXiv: 2203.03597.

Donoho, David and Andrea Montanari (2016). "High dimensional robust m-estimation: Asymptotic variance via approximate message passing." *Probability Theory and Related Fields* 166.3, pp. 935–969.

El Karoui, Noureddine (2018). "On the impact of predictor geometry on the performance on high-dimensional ridge-regularized generalized robust regression estimators." *Probability Theory and Related Fields* 170.1, pp. 95–175.

El Karoui, Noureddine, Derek Bean, Peter J Bickel, Chinghway Lim, and Bin Yu (2013). "On robust regression with high-dimensional predictors." *Proceedings of the National Academy of Sciences* 110.36, pp. 14557–14562.

Gordon, Yehoram (1985). "Some inequalities for Gaussian processes and applications." *Israel Journal of Mathematics* 50.4, pp. 265–289.

Ju, Peizhong, Xiaojun Lin, and Jia Liu (2020). "Overfitting Can Be Harmless for Basis Pursuit: Only to a Degree." *Advances in Neural Information Processing Systems*. arXiv: 2002.00492.

Koehler, Frederic, Lijia Zhou, Danica J. Sutherland, and Nathan Srebro (2021). "Uniform Convergence of Interpolators: Gaussian Width, Norm Bounds and Benign Overfitting." *Advances in Neural Information Processing Systems*. arXiv: 2106.09276.

Koltchinskii, Vladimir and Shahar Mendelson (2015). "Bounding the smallest singular value of a random matrix without concentration." *International Mathematics Research Notices* 2015.23, pp. 12991–13008.

Lecué, Guillaume and Shahar Mendelson (2013). "Learning subgaussian classes: Upper and minimax bounds." arXiv: 1305.4825.

Li, Yue and Yuting Wei (2021). "Minimum ℓ_1-norm interpolators: Precise asymptotics and multiple descent." arXiv: 2110.09502.

Liang, Tengyuan and Pragya Sur (2020). "A Precise High-Dimensional Asymptotic Theory for Boosting and Minimum-L1-Norm Interpolated Classifiers." arXiv: 2002.01586.

Lugosi, Gábor and Shahar Mendelson (2019). "Mean estimation and regression under heavy-tailed distributions: A survey." *Foundations of Computational Mathematics* 19.5, pp. 1145–1190.

Mendelson, Shahar (2014). "Learning without concentration." *Conference on Learning Theory*. PMLR, pp. 25–39.

Mendelson, Shahar (2017). "Extending the scope of the small-ball method." arXiv: 1709.00843.

Montanari, Andrea, Feng Ruan, Youngtak Sohn, and Jun Yan (2019). "The generalization error of max-margin linear classifiers: High-dimensional asymptotics in the overparametrized regime." arXiv: 1911.01544.

Muthukumar, Vidya, Adhyyan Narang, Vignesh Subramanian, Mikhail Belkin, Daniel Hsu, and Anant Sahai (2021). "Classification vs regression in overparameterized regimes: Does the loss function matter?" *Journal of Machine Learning Research* 22, pp. 1–69. arXiv: 2005.08054.

Muthukumar, Vidya, Kailas Vodrahalli, Vignesh Subramanian, and Anant Sahai (2020). "Harmless interpolation of noisy data in regression." *IEEE Journal on Selected Areas in Information Theory*. arXiv: 1903.09139.

Negrea, Jeffrey, Gintare Karolina Dziugaite, and Daniel M. Roy (2020). "In Defense of Uniform Convergence: Generalization via derandomization with an application to interpolating predictors." *International Conference on Machine Learning*. arXiv: 1912.04265.

Neyshabur, Behnam, Ryota Tomioka, and Nathan Srebro (2015). "In Search of the Real Inductive Bias: On the Role of Implicit Regularization in Deep Learning." *International Conference on Learning Representations – Workshop*. arXiv: 1412.6614.

O'Donnell, Ryan (2014). *Analysis of boolean functions*. Cambridge University Press.

Panchenko, Dmitry (2002). "Some Extensions of an Inequality of Vapnik and Chervonenkis." *Electronic Communications in Probability* 7, pp. 55–65. arXiv: 0405342.

Panchenko, Dmitry (2003). "Symmetrization approach to concentration inequalities for empirical processes." *The Annals of Probability* 31.4, pp. 2068–2081.

Parikh, Neal and Stephen Boyd (2014). "Proximal algorithms." *Foundations and Trends in optimization* 1.3, pp. 127–239.

Planiden, Chayne and Xianfu Wang (2019). "Proximal mappings and Moreau envelopes of single-variable convex piecewise cubic functions and multivariable gauge functions." *Nonsmooth optimization and its applications*. Springer, pp. 89–130.

Salehi, Fariborz, Ehsan Abbasi, and Babak Hassibi (2019). "The impact of regularization on high-dimensional logistic regression." *Advances in Neural Information Processing Systems* 32.

Shamir, Ohad (2022). "The Implicit Bias of Benign Overfitting." arXiv: 2201.11489.

Srebro, Nathan, Karthik Sridharan, and Ambuj Tewari (2010). "Optimistic Rates for Learning with a Smooth Loss." arXiv: 1009.3896.

Sur, Pragya and Emmanuel J Candès (2019). "A modern maximum-likelihood theory for high-dimensional logistic regression." *Proceedings of the National Academy of Sciences* 116.29, pp. 14516–14525.

Sur, Pragya, Yuxin Chen, and Emmanuel J Candès (2019). "The likelihood ratio test in high-dimensional logistic regression is asymptotically a rescaled chi-square." *Probability theory and related fields* 175.1, pp. 487–558.

Thrampoulidis, Christos, Ehsan Abbasi, and Babak Hassibi (2018). "Precise error analysis of regularized M-estimators in high dimensions." *IEEE Transactions on Information Theory* 64.8, pp. 5592–5628.

Thrampoulidis, Christos, Samet Oymak, and Babak Hassibi (2015). "Regularized linear regression: A precise analysis of the estimation error." *Conference on Learning Theory*.

Tsigler, Alexander and Peter L. Bartlett (2020). "Benign overfitting in ridge regression." arXiv: 2009.14286.

Van Handel, Ramon (2014). "Probability in High Dimension." Lecture notes, Princeton University. URL: https://web.math.princeton.edu/~rvan/APC550.pdf.

Vapnik, Vladimir (1982). *Estimation of dependences based on empirical data*. Springer Science & Business Media.

Wang, Guillaume, Konstantin Donhauser, and Fanny Yang (2021). "Tight bounds for minimum l1-norm interpolation of noisy data." arXiv: 2111.05987.

Zhang, Chiyuan, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals (2017). "Understanding deep learning requires rethinking generalization." *International Conference on Learning Representations*. arXiv: 1611.03530.

Zhang, Tong (2004). "Solving large scale linear prediction problems using stochastic gradient descent algorithms." *Proceedings of the twenty-first international conference on Machine learning*, p. 116.

Zhao, Qian, Pragya Sur, and Emmanuel J Candes (2022). "The asymptotic distribution of the MLE in high-dimensional logistic models: Arbitrary covariance." *Bernoulli* 28.3, pp. 1835–1861.

Zhou, Lijia, Frederic Koehler, Danica J Sutherland, and Nathan Srebro (2021). "Optimistic Rates: A Unifying Theory for Interpolation Learning and Regularization in Linear Regression." arXiv: 2112.04470.

Zhou, Lijia, Danica J. Sutherland, and Nathan Srebro (2020). "On Uniform Convergence and Low-Norm Interpolation Learning." *Advances in Neural Information Processing Systems*. arXiv: 2006. 05942.