Maximizing the robust margin provably overfits on noiseless data

Abstract

Numerous recent works show that overparameterization implicitly reduces variance, suggesting vanishing benefits for explicit regularization in high dimensions. However, this narrative has been challenged by empirical observations indicating that adversarially trained deep neural networks suffer from robust overfitting. While existing explanations attribute this phenomenon to noise or problematic samples in the training data set, we prove that even on entirely noiseless data, achieving vanishing adversarial logistic training loss is suboptimal compared to regularized counterparts.

1. Introduction

A modern narrative in machine learning is that regularization is not required for good performance for large overparameterized models. This narrative is based on theoretical and experimental findings that analyze the population risk in a setup where the train and test data follows the same distribution (see (Hastie et al., 2019; Bartlett et al., 2020; Yang et al., 2020; Nakkiran et al., 2020) and references therein). However, the successful adoption of machine learning models in real world applications crucially hinges on their robustness to adversarial attacks or distribution shifts. A popular method to achieve low *robust risk* is to minimize a robust training loss for example using adversarial training (Goodfellow et al., 2015). Mounting empirical evidence suggests that when minimizing the robust loss the narrative that regularization is not required is false: additional regularization, for example via early stopping the adversarial training, often leads to better generalization (Rice et al., 2020; Sagawa et al., 2020a;b). For example, Rice et al. (2020) show that overparameterized deep neural networks that are adversarially trained with ℓ_∞ perturbations benefit from early stopping on image data sets. This phenomenon is sometimes referred to as *robust overfitting*.

[1]Anonymous Institution, Anonymous City, Anonymous Region, Anonymous Country. Correspondence to: Anonymous Author <anon.email@domain.com>.

Preliminary work. Under review by the ICML 2021 workshop on A Blessing in Disguise: The Prospects and Perils of Adversarial Machine Learning. Do not distribute.

Figure 1. The robust risks for the robust max-margin solution ($\lambda \to 0$) are strictly higher than the regularized minimizer ($\lambda_{\mathrm{opt}} > 0$) even in the overparameterized regime $d \gg n$ and when the training data is noiseless (see detailed setting in Section 2 and experimental details in Appendix C). This illustrates how adversarial training benefits from regularization.

For noisy settings, the following explanation for robust overfitting has been proposed in previous work: since the robust risk amplifies estimation errors, its variance is larger and hence regularization can be beneficial for generalization (Sanyal et al., 2021). However, Figure 4 in Appendix B shows that even when estimating entirely noiseless signals, robust overfitting persists! This observation does not only apply to image data but also to simple linear models: Figure 1 shows the gap in performance between ridge regularized ℓ_∞-robust logistic regression and the robust max-margin solution.

Existing theoretical results do not predict nor explain the observations in Figure 1. In particular, it is perhaps unexpected that for noiseless data, strictly sacrificing robust data fit by increasing the weight of the ridge penalty leads to better generalization. In this paper, we show for robust logistic regression and a sparse ground truth that in the high-dimensional regime, where $d > n$, maximizing the robust margin leads to suboptimal generalization. Specifically,

- We prove that a strictly positive ridge penalty leads to a systematic improvement in generalization compared to the robust max-margin solution. Our results provide the first rigorous proof for robust overfitting even in the absence of noise.

- We experimentally show a perhaps unexpected regularizing effect of adding a moderate amount of wrong label information: because the noisy data cannot be robustly linearly separated, it prevents the unregularized minimizer from converging to the max-margin solution and thus results in better robust generalization.

2. Risk minimization framework

In this section, we describe the setup for our theoretical analysis of adversarial training with the logistic loss. We define the data generating process, the standard and robust population risks, and formally introduce the estimators that we analyze.

2.1. Problem setting

We consider a discriminative data-generating model for classification with covariates (or features) $x \in \mathbb{R}^d$ drawn from an isotropic Gaussian distribution $x \sim \mathcal{N}(0, I_d)$. In this paper we focus primarily on training with noiseless data, namely we observe deterministic labels given by $y = \text{sgn}\langle\theta^\star, x\rangle \in \{-1, +1\}$ for all covariates $x \in \mathbb{R}^d$, where the fixed vector $\theta^\star$ denotes the ground truth. We consider the scenario where the ground truth has inherent low-dimensional structure in the form of sparsity $\theta^\star = (1, 0, \ldots, 0)^\top$. As we discuss later in this section, ℓ_∞ robustness and sparsity via its convex surrogate, the ℓ_1 norm, are related objectives.

The results in this paper are of asymptotic nature and hold when $d/n \to \gamma$ as both the dimensionality d and the number of samples n tend to infinity. This high-dimensional regime is widely studied in the literature (Bühlmann & Van De Geer, 2011; Wainwright, 2019) as it yields precise predictions for many real world problems where both the input dimension and the data set size are large. It is also the predominant setting considered in previous theoretical papers that discuss overparameterized linear models (Dobriban & Wager, 2018; Hastie et al., 2019; Ali et al., 2020; Deng et al., 2021; Javanmard et al., 2020; Javanmard & Soltanolkotabi, 2020; Sur & Candès, 2019).

2.2. Robust risk for evaluation

The broad application of ML models in real-world decision-making processes increases requirements on their robustness. For example, for the image domain, robust classifiers should yield the same prediction when an image is attacked via an additive imperceptible ℓ_∞-perturbation that do not change the ground truth label. In this case, the estimator which has zero standard population risk also achieves zero robust population risk. Transferred to linear classification, we require such additive *consistent* perturbations to be orthogonal to $\theta^\star$, that is $\delta \in \mathcal{U}_c(\epsilon) := \{\delta \in \mathbb{R}^d : \|\delta\|_\infty \leq$

ϵ and $\langle\theta^\star, \delta\rangle = 0\}$. We hence evaluate the adversarially robust risk of a parameter θ with respect to consistent ℓ_∞-perturbations

$$\mathbf{R}_\epsilon(\theta) := \mathbb{E}_{X \sim \mathbb{P}} \min_{\delta \in \mathcal{U}_c(\epsilon)} \mathbb{1}_{\text{sgn}(\langle\theta, X+\delta\rangle) \neq \text{sgn}(\langle\theta^\star, X\rangle)}, \quad (1)$$

where the expectation is taken over the marginal feature distribution $\mathbb{P}$ and $\mathbb{1}$ is the indicator function.

For $\epsilon = 0$ we obtain the population $0 - 1$ risk, also called standard risk, denoted by $\mathbf{R}(\theta)$. For the 1-sparse ground truth $\theta^\star$ the risks have a closed form expression given in Lemma A.1 in Appendix A.1.

2.3. Interpolating and regularized estimator

We say that the data is *robustly separable* when the robust max-min margin solution exists and reads

$$\hat{\theta}_0 := \arg\min_\theta \|\theta\|_2 \text{ such that for all } i, \quad (2)$$

$$\max_{\delta \in \mathcal{U}(\epsilon)} y_i\langle\theta, x_i + \delta\rangle \geq 1.$$

Robust separability is guaranteed for noiseless data and for consistent perturbations $\mathcal{U}(\epsilon) = \mathcal{U}_c(\epsilon)$ that are orthogonal to the ground truth $\theta^\star$. However, minimizing with respect to consistent perturbations in Equation (2) requires the full knowledge of the ground truth during training, thus leaking information about $\theta^\star$. Instead, many papers to date (e.g. (Javanmard et al., 2020; Javanmard & Soltanolkotabi, 2020)) consider the unrestricted (and hence inconsistent) ℓ_∞-perturbation set $\mathcal{U}_{ic}(\epsilon) := \{\delta \in \mathbb{R}^d : \|\delta\|_\infty \leq \epsilon\}$. When training with inconsistent perturbations however, it is possible that the perturbed data crosses the true decision boundary and as a result, a minimizing solution fits wrong labels even when the actual training samples are noiseless.

We want to study the purely noiseless setting and hence focus primarily on adversarial training with consistent perturbations. This setting is also the one in which we expect regularization to help the least, according to classical wisdom. Having said that, we discuss inconsistent adversarial training in more detail in Section 4.2.

In this paper, we analyze the minimizer of the regularized ℓ_∞ robust logistic loss

$$\hat{\theta}_\lambda := \arg\min_\theta \frac{1}{n} \sum_{i=1}^n \max_{\delta \in \mathcal{U}(\epsilon)} \log(1 + e^{-\langle\theta, x_i+\delta\rangle y_i}) + \lambda\|\theta\|_2^2.$$

$$(3)$$

We study ridge regularized logistic regression since for robustly sepable data, the max-margin solution directionally aligns with minimizer $\hat{\theta}_\lambda$ for $\lambda \to 0$, i.e. $\hat{\theta}_0 = \lim_{\lambda \to 0} \hat{\theta}_\lambda$. This fact follows from the results in (Rosset et al., 2004), when using the closed form expression for the consistent robust risk (1). Since the robust risk is independent of the norm of the estimator, we simply refer to the max-margin solution as the normalized vector $\frac{\hat{\theta}_0}{\|\hat{\theta}_0\|_2}$.

(a) Fixed n, d and asymptotic predictions
(b) Benefit of ridge regularization
(c) Benefit of early stopping

Figure 2. (a) Robust and standard risks of the robust max-margin ($\lambda \to 0$) and robust regularized minimizer ($\lambda = 1$) using $\epsilon = 0.05$ as a function of the overparameterization ratio d/n for simulations (markers) and asymptotic theoretical predictions from Theorem 3.1 (lines). (b),(c) Normalized robust margins and risks of empirical simulations using $\epsilon = 0.1$ and $d/n = 8$, with respect to (b) increasing $1/\lambda$ and (c) gradient descent iterations when minimizing Equation (3) using $\lambda = 0$. The normalized robust margin is defined as $\min_i \min_{\delta \in \mathcal{U}(\epsilon)} y_i \langle \theta, x_i + \delta \rangle / \|\theta\|_2$. Both ridge regularization and early stopping yield superior robust and standard risks. All experiments use $n = 1000$ and consistent ℓ_∞ perturbations. See Appendix C for more details.

Finally, we remark that for standard logistic regression, gradient descent converges to the direction of the max margin solution (Ji & Telgarsky, 2019; Soudry et al., 2018). In fact, as argued in (Javanmard & Soltanolkotabi, 2020) the results can be straight forwardly extended to the robust loss (3).

3. Regularization for robustly separable data

In this section, we prove that the robust max-margin solution (2) overfits as it yields worse generalization than the regularized esimator (3) with optimally chosen $\lambda > 0$ for entirely noiseless training data. We derive precise asymptotic predictions that show a systematic benefit of ℓ_2-regularization for robust logistic regression when $d, n \to \infty$ and $d/n \to \gamma$. We further show that an early stopping gradient descent on the unregularized robust logistic loss in Equation (3) yields much smaller population risks than the robust max-margin solution at convergence.

3.1. Benefits of ridge regularization

We now present an informal statement describing the asymptotic standard and robust risk for linear classification with the logistic loss as $d, n \to \infty$. We refer to Appendix D for the precise statement and the proof. The latter is inspired by the works (Javanmard & Soltanolkotabi, 2020; Salehi et al., 2019) and uses the Convex Gaussian Minimax Theorem (CGMT) (Gordon, 1988; Thrampoulidis et al., 2015).

Theorem 3.1 (Informal)**.** *Assume that $\epsilon = \epsilon_0/\sqrt{d}$ for some constant ϵ_0 independent of n, d and $\theta^\star = (1, 0, \cdots, 0)$. Then, the robust and standard risks of the estimator $\hat{\theta}_\lambda$ from*

Equation (2),(3) with respect to consistent attacks and $\lambda \geq 0$ converge in probability as $d, n \to \infty$, $d/n \to \gamma$

$$\boldsymbol{R}(\hat{\theta}_\lambda) \to \frac{1}{\pi} \arccos\left(\frac{\nu_\parallel^\star}{\nu^\star}\right)$$

$$\boldsymbol{R}_\epsilon(\hat{\theta}_\lambda) \to \boldsymbol{R}(\hat{\theta}_\lambda) + \frac{1}{2}\mathrm{erf}\left(\frac{\epsilon_0\delta^\star}{\sqrt{2}\nu^\star}\right) + I\left(\frac{\epsilon_0\delta^\star}{\nu^\star}, \frac{\nu_\parallel^\star}{\nu^\star}\right)$$

with erf *the error function,*

$$I(t, u) := \int_0^t \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) \mathrm{erf}\left(\frac{xu}{\sqrt{2(1-u^2)}}\right) dx$$

and where $\nu^\star = \sqrt{(\nu_\perp^\star)^2 + (\nu_\parallel^\star)^2}$, and $\nu_\perp^\star, \nu_\parallel^\star, \delta^\star$ are the unique solution of a scalar optimization problem specified in Appendix D that depends on $\theta^\star, \gamma, \epsilon_0$ and λ.

We illustrate the theorem in Figure 2a. Figure 2a depicts the standard and robust risk obtained by the asymptotic theoretical predictions for $d, n \to \infty$ and simulations for $n = 1000$ and $\epsilon = 0.1$. The theoretical curves are obtained by solving the scalar optimization problem specified in Appendix D; the empirical setup is described in Appendix C. We observe that the precise asymptotics indeed predict benefits of regularization for robust logistic regression on noiseless training data. This trend is further supported by our simulation results closely following the theoretical curves. Furthermore, in Figure 2b we observe for a fixed overparameterization ratio $d/n = 8$, how optimizing the robust logistic loss well beyond 100% robust training accuracy (i.e. when the robust margin becomes positive) substantially hurts generalization.

(a) Consistent AT with noisy training data   (b) Consistent and inconsistent AT with noiseless data

Figure 3. (a) Training loss and robust risk with respect to increasing training label noise for $\epsilon = 0.1$ and $d = 8000$. We observe that, counterintuitively, moderate amounts of label noise yield estimators with lower robust risk since the noisy labels prevent interpolation. (b) Training loss and risks with respect to increasing train and test ϵ for $d = 500$. In contrast to training with consistent attacks, unregularized inconsistent adversarial training does not achieve vanishing training loss for large enough ϵ, and hence leads to a smaller robust risk. Both experiments use $n = 1000$ and perform unregularized ($\lambda \to 0$) training.

3.2. Benefits of early stopping

In addition to varying the ridge coefficient λ, we show that the same trends as for $1/\lambda \to \infty$ can be observed on the optimization path when training with gradient descent. Figure 2c indicates that early stopped gradient descent can improve the robust risk for logistic regression, similar to ℓ_2-regularization. Both early stopping and ℓ_2-regularization prevent interpolation and yield an estimator with significantly lower standard and robust risk compared to the interpolator ($\lambda \to 0$) respectively the converged estimator (gradient steps $\to \infty$).

4. The benefits of non-separable data

In the previous section we focused on settings with noiseless observations and studied the generalization performance of regularized estimators that do not maximize the robust margin. In this section, we show how, surprisingly, as a consequence of non-robust separability, adding noisy labels in the training loss also leads to an estimator with better generalization than the max-margin solution of the corresponding noiseless problem.

4.1. Avoiding the max-margin solution via label noise

We first consider the case where we introduce label noise to the training data. Figure 3a shows the robust and standard risk together with the training loss of estimator $\hat{\theta}_\lambda$ (3) for $\lambda \to 0$ with varying percentages of flipped labels. For low noise levels, the data is robustly separable and the training loss vanishes), yielding the max-margin solution (2). For large enough noise the constraints in (2) are not feasible anymore and the training loss of the estimator at convergence

starts to increase. As we can see, this effectively leads to an implicit regularization as the risk at this point starts to decrease again. In particular, training on the data with 20% label flips results in a robust risk that is $\sim 25\%$ lower than training on the clean dataset without any mislabeled samples.

4.2. Avoiding the max-margin solution via inconsistent training

For large ϵ and inconsistent perturbations $\mathcal{U}_{ic}(\epsilon)$, even noiseless training data might not be robustly separable and hence the limit of $\lambda \to 0$ does not result in the max margin solution. Consequently, this leads to a similar effect as adding label noise, which was discussed in Section 4.1. Indeed, Figure 3b reveals that the resulting estimator trained with the inconsistent adversarial loss achieves a better performance than the max-margin estimator (2).

5. Conclusions

In this paper we prove that on high-dimensional input data, ridge regularization of the robust logistic loss leads to a solution that generalizes better than the unregularized robust max-margin estimator that is obtained when training with gradient descent until convergence. We further show experimentally that early stopping gradient descent yields a similar benefit. While it is well known that covariate noise can act as an implicit regularization (Bishop, 1995), in contrast to common intuition, we demonstrate that the performance can also improve when introducing wrong labels in the training loss as it prevents the estimator from converging to the max-margin solution as $\lambda \to 0$.

References

Ali, A., Dobriban, E., and Tibshirani, R. The implicit regularization of stochastic gradient flow for least squares. In *Proceedings of the International Conference on Machine Learning (ICML)*, volume 119, pp. 233–244, 13–18 Jul 2020.

Bartlett, P. L., Long, P. M., Lugosi, G., and Tsigler, A. Benign overfitting in linear regression. *Proceedings of the National Academy of Sciences*, 117(48):30063–30070, 2020.

Bishop, C. Training with noise is equivalent to tikhonov regularization. *Neural Computation*, 7(1):108–116, 1995. ISSN 0899-7667.

Bühlmann, P. and Van De Geer, S. *Statistics for high-dimensional data: methods, theory and applications*. Springer Science & Business Media, 2011.

Bühlmann, P. et al. Invariance, causality and robustness. *Statistical Science*, 35(3):404–426, 2020.

Chen, Y. and Bühlmann, P. Domain adaptation under structural causal models. *arXiv preprint arXiv:2010.15764*, 2020.

Deng, Z., Kammoun, A., and Thrampoulidis, C. A model of double descent for high-dimensional binary linear classification. *Information and Inference: A Journal of the IMA*, 04 2021.

Dobriban, E. and Wager, S. High-dimensional asymptotics of prediction: Ridge regression and classification. *The Annals of Statistics*, 46(1):247 – 279, 2018.

Dong, C., Liu, L., and Shang, J. Data profiling for adversarial training: On the ruin of problematic data. *arXiv preprint arXiv:2102.07437*, 2021.

Goodfellow, I., Shlens, J., and Szegedy, C. Explaining and harnessing adversarial examples. In *Proceedings of the International Conference on Learning Representations (ICLR)*, 2015.

Gordon, Y. On milman's inequality and random subspaces which escape through a mesh in rn. In *Geometric aspects of functional analysis*, pp. 84–106. 1988.

Hastie, T., Montanari, A., Rosset, S., and Tibshirani, R. J. Surprises in high-dimensional ridgeless least squares interpolation. *arXiv preprint arXiv:1903.08560*, 2019.

Javanmard, A. and Soltanolkotabi, M. Precise statistical analysis of classification accuracies for adversarial training. *arXiv preprint arXiv:2010.11213*, 2020.

Javanmard, A., Soltanolkotabi, M., and Hassani, H. Precise tradeoffs in adversarial training for linear regression. In *Proceedings of the Conference on Learning Theory (COLT)*, pp. 2034–2078, 2020.

Ji, Z. and Telgarsky, M. The implicit bias of gradient descent on nonseparable data. In *Proceedings of the Conference on Learning Theory (COLT)*, pp. 1772–1798, 2019.

Jourani, A., Thibault, L., and Zagrodny, D. Differential properties of the moreau envelope. *Journal of Functional Analysis*, 266(3):1185–1237, 2014.

Nakkiran, P., Kaplun, G., Bansal, Y., Yang, T., Barak, B., and Sutskever, I. Deep double descent: Where bigger models and more data hurt. In *Proceedings of the International Conference on Learning Representations (ICLR)*, 2020.

Rice, L., Wong, E., and Kolter, Z. Overfitting in adversarially robust deep learning. In *Proceedings of the International Conference on Machine Learning (ICML)*, pp. 8093–8104, 2020.

Rosset, S., Zhu, J., and Hastie, T. Margin maximizing loss functions. In *Advances in Neural Information Processing Systems*, volume 16, 2004.

Sagawa, S., Koh, P. W., Hashimoto, T. B., and Liang, P. Distributionally robust neural networks. In *International Conference on Learning Representations*, 2020a.

Sagawa, S., Raghunathan, A., Koh, P. W., and Liang, P. An investigation of why overparameterization exacerbates spurious correlations. In *Proceedings of the International Conference on Machine Learning (ICML)*, volume 119, pp. 8346–8356, 13–18 Jul 2020b.

Salehi, F., Abbasi, E., and Hassibi, B. The impact of regularization on high-dimensional logistic regression. In *Advances in Neural Information Processing Systems*, volume 32, 2019.

Sanyal, A., Dokania, P. K., Kanade, V., and Torr, P. How benign is benign overfitting? In *Proceedings of the International Conference on Learning Representations (ICLR)*, 2021.

Sinha, A., Namkoong, H., and Duchi, J. Certifying some distributional robustness with principled adversarial training. In *International Conference on Learning Representations*, 2018.

Soudry, D., Hoffer, E., Nacson, M. S., Gunasekar, S., and Srebro, N. The implicit bias of gradient descent on separable data. *Journal of Machine Learning Research*, 19 (70):1–57, 2018.

Sur, P. and Candès, E. J. A modern maximum-likelihood theory for high-dimensional logistic regression. *Proceedings of the National Academy of Sciences*, 116(29):14516–14525, 2019.

Thrampoulidis, C., Oymak, S., and Hassibi, B. Regularized linear regression: A precise analysis of the estimation error. In *Proceedings of the Conference on Learning Theory (COLT)*, pp. 1683–1709, 2015.

Wainwright, M. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge University Press, 2019. ISBN 9781108498029.

Yang, Z., Yu, Y., You, C., Steinhardt, J., and Ma, Y. Rethinking bias-variance trade-off for generalization of neural networks. In *Proceedings of the International Conference on Machine Learning (ICML)*, pp. 10767–10777, 2020.

A. Setting

In Section A.1 we derive closed-form expressions of the standard and robust risks (1). Furthermore, Section A.2 discusses that the robust risk (1) upper-bounds the worst case risk under distributional mean shifts.

A.1. Closed-form of robust risk for classification

Similarly to linear regression, we can express the robust and standard risk for the linear classification model in Section 3 as stated in the following lemma.

Lemma A.1. *Assume that $\mathbb{P}_X$ is the isotropic Gaussian distribution and $\theta^\star = (1, 0, \cdots, 0)^\top$. Then,*

1. *For any non-decreasing loss $\ell : \mathbb{R} \to \mathbb{R}$ we have*

$$\max_{\delta_i \in \mathcal{U}_\infty(\epsilon)} \ell(y_i \langle x_i + \delta_i, \theta \rangle) = \ell(y_i \langle x_i, \theta \rangle - \epsilon \|\Pi_\perp \theta\|_1). \tag{4}$$

2. *For the 0-1 loss the robust risk (1) with respect to ℓ_∞ perturbations is given by*

$$\boldsymbol{R}_\epsilon(\theta) = \frac{1}{\pi} \arccos\left(\frac{\langle \theta^\star, \theta \rangle}{\|\theta\|_2}\right) + \frac{1}{2}\mathrm{erf}\left(\frac{\epsilon\|\Pi_\perp \theta\|_1}{\sqrt{2}\|\theta\|_2}\right) + I\left(\frac{\epsilon\|\Pi_\perp \theta\|_1}{\|\theta\|_2}, \frac{\langle \theta^\star, \theta \rangle}{\|\theta\|_2}\right), \tag{5}$$

   with

$$I(t, u) := \int_0^t \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) \Phi\left(\frac{xu}{\sqrt{2}\sqrt{1-u^2}}\right) dx. \tag{6}$$

Proof. We first prove Eq (4). Because ℓ is non-increasing, we have

$$\max_{\delta_i \in \mathcal{U}_\infty(\epsilon)} \ell(y_i \langle x_i + \delta_i, \theta \rangle) = \ell(\min_{\delta_i \in \mathcal{U}_\infty(\epsilon)} y_i \langle x_i + \delta_i, \theta \rangle)$$

$$= \ell(y_i \langle x_i, \theta \rangle + \min_{\|\delta_i\|_\infty \leq \epsilon, \delta_i \perp \theta^\star} \langle \delta_i, \theta \rangle).$$

While minimization over δ has no closed-form solution in general, we note that for $\theta^\star = (1, 0, \cdots, 0)$, we get $\min_{\|\delta_i\|_\infty \leq \epsilon, \delta_i \perp \theta^\star} \langle \delta_i, \theta \rangle = -\epsilon\|\Pi_\perp \theta\|_1$ and Equation (4).

Let $\mathbb{1}\{E\}$ be the indicator function, 1 if the event E holds. Since $\ell(\cdot) = \mathbb{1}_{\cdot \leq 0}$ is non-increasing we can use (4) and write

$$\boldsymbol{R}_\epsilon(\theta) = \mathbb{E}_X \max_{\delta \in \mathcal{U}_\infty(\epsilon)} \mathbb{1}\{\mathrm{sgn}(\langle X, \theta^\star \rangle)\langle X + \delta, \theta \rangle \leq 0\}$$

$$= \mathbb{E}_X \mathbb{1}\{\mathrm{sgn}(\langle X, \theta^\star \rangle)\langle X, \theta \rangle - \epsilon\|\Pi_\perp \theta\|_1 \leq 0\}.$$

Let $\widehat{\Pi}_\| := \frac{1}{\|\theta\|_2^2}\theta\theta^\top$ be the projection onto the subspace spanned by θ and $\widehat{\Pi}_\perp := I_d - \widehat{\Pi}_\|$ the projection onto the orthogonal subspace. Since X is a vector with i.i.d. standard normal distributed entries, we equivalently have

$$\boldsymbol{R}_\epsilon(\theta) = \mathbb{E}_{Z_1, Z_2} \mathbb{1}\{Z_1 \,\mathrm{sgn}\left(Z_1\|\widehat{\Pi}_\|\theta^\star\|_2 + Z_2\|\widehat{\Pi}_\perp\theta^\star\|_2\right) - \epsilon\frac{\|\Pi_\perp \theta\|_1}{\|\theta\|_2} \leq 0\}, \tag{7}$$

with Z_1, Z_2 two independent standard normal random variables. For brevity of notation, define $\nu = \epsilon\frac{\|\Pi_\perp \theta\|_1}{\|\theta\|_2}$ and $b(Z_1, Z_2) = \mathrm{sgn}\left(Z_1\|\widehat{\Pi}_\|\theta^\star\|_2 + Z_2\|\widehat{\Pi}_\perp\theta^\star\|_2\right) =: \mathrm{sgn}(\beta^\top Z)$ with $\beta^\top = (\|\widehat{\Pi}_\|\theta^\star\|_2, \|\widehat{\Pi}_\perp\theta^\star\|_2)$.

Define the event $A = \{\mathrm{sgn}\left(Z_1\|\widehat{\Pi}_\|\theta^\star\|_2 + Z_2\|\widehat{\Pi}_\perp\theta^\star\|_2\right) - \epsilon\frac{\|\Pi_\perp \theta\|_1}{\|\theta\|_2} \leq 0\}$. Because Z_2 is symmetric, the distribution of $Z_1 b(Z_1, Z_2)$ is symmetric, hence we can rewrite the risk

$$\boldsymbol{R}_\epsilon(\theta) = \underbrace{\mathbb{P}(b(Z_1, Z_2) \leq 0 | Z_1 \geq 0)}_{T_1} + \underbrace{\mathbb{P}(Z_1 \leq \nu, b(Z_1, Z_2) \geq 0 | Z_1 \geq 0)}_{T_2} \tag{8}$$

and derive expression for T_1, T_2 separately.

Step 1: Proof for T_1   Note that due to $\|\theta^\star\|_2 = 1$ we have $\|\beta\|_2 = 1$ and recall that $T_1 = \mathbb{P}(\beta^\top Z \le 0 | Z_1 \ge 0)$. Using the fact that both Z_1 and Z_2 are independent standard normal distributed random variables, a simple geometric argument then yields that $T_1 = \frac{\alpha}{\pi}$ with $\alpha = \arccos\left(\frac{\beta_1}{\sqrt{\beta_1^2 + \beta_2^2}}\right) = \arccos(\beta_1)$. Noting that $\beta_1 = \|\widehat{\Pi}_\| \theta^\star\|_2 = \frac{\langle \theta^\star, \theta \rangle}{\|\theta\|_2}$ then yields $T_1 = \frac{1}{\pi}\arccos\left(\frac{\langle\theta^\star,\theta\rangle}{\|\theta\|_2}\right)$.

Step 2: Proof for T_2   First, assume that $\langle\theta^\star,\theta\rangle \ge 0$. We separate the event $\mathcal{V} = \{Z_1 \le \nu, b(Z_1, Z_2) \ge 0\}$ into two events $\mathcal{V} = \mathcal{V}_1 \cup \mathcal{V}_2$

$$\mathcal{V}_1 = \{Z_1 \le \nu, Z_2 \ge 0\} \text{ and } \mathcal{V}_2 = \{Z_1 \le \nu, b(Z_1, Z_2) \ge 0, Z_2 \le 0\}.$$

The conditional probability of the first event is directly given

$$\mathbb{P}(\mathcal{V}_1 | Z_1 \ge 0) = \mathbb{P}(Z_2 \ge 0)\mathbb{P}(Z_1 \le \nu | Z_1 \ge 0) = \frac{1}{2}\mathrm{erf}\left(\frac{\nu}{\sqrt{2}}\right).$$

Hence it only remains to find an expression for $\mathbb{P}(\mathcal{V}_2 | Z_1 \ge 0)$. Letting μ denote the standard normal distribution, we can write

$$\mathbb{P}(Z_1 \le \nu, Z_2 \le 0, b(Z_1, Z_2) \ge 0 | Z_1 \ge 0) = 2 \int_0^\nu \int_0^{\frac{\beta_1 x}{\beta_2}} d\mu(y) d\mu(x) = \int_0^\nu \frac{1}{2}\mathrm{erf}\left(\frac{\beta_1 x}{\beta_2}\right) d\mu(x).$$

Together with Step 1, Equation (5) follows by noting that $\beta_1^2 + \beta_2^2 = 1$. Finally, the proof for the case where $\langle\theta^\star,\theta\rangle \le 0$ follows exactly from the same argument and the proof is complete. □

A.2. Distribution shift robustness and consistent adversarial robustness

In this section we rigorously introduce distribution shift robustness and show the relation to consistent ℓ_p adversarial robustness for certain types of distribution shifts.

When learned models are deployed in the wild, the i.i.d. assumption does not always hold. That is, the test loss might be evaluated on samples from a slightly different distribution. Shifts in the mean of the covariate distribution is a standard intervention studied in the invariant causal prediction literature (Bühlmann et al., 2020; Chen & Bühlmann, 2020). For mean shifts in the null space of the ground truth $\theta^\star$ we define an alternative evaluation metric that we refer to as the *distributionally robust risk* defined as follows:

$$\tilde{\mathbf{R}}_\epsilon(\theta) := \max_{\mathbb{Q} \in \mathcal{V}_q(\epsilon; \mathbb{P})} \mathbb{E}_{X \sim \mathbb{Q}} \ell_{\text{test}}(\langle\theta, X + \delta\rangle, \langle\theta^\star, X\rangle), \text{ with}$$

$$\mathcal{V}_p(\epsilon; \mathbb{P}) := \{\mathbb{Q} \in \mathcal{P} : \|\mu_\mathbb{P} - \mu_\mathbb{Q}\|_p \le \epsilon \text{ and } \langle\mu_\mathbb{P} - \mu_\mathbb{Q}, \theta^\star\rangle = 0\}$$

where $\mathcal{V}_p$ is the neighborhood of mean shifted probability distributions.

A duality between distribution shift robustness and adversarial robustness has been established in earlier work such as (Sinha et al., 2018) for general convex, continuous losses ℓ_{test}. For our setting, the following lemma holds.

Lemma A.2. *For any $\epsilon \ge 0$ and θ, we have $\tilde{\mathbf{R}}_\epsilon(\theta) \le \mathbf{R}_\epsilon(\theta)$.*

Proof. The proof follows directly from the definition and consistency of the perturbations $\mathcal{U}_p(\epsilon)$ and orthogonality of the mean shifts for the neighborhood $\mathcal{V}_p$. By defining the random variable $w = x - \mu_\mathbb{P}$ for $x \sim \mathbb{P}$ we have the distributional equivalence

$$x' = \mu_\mathbb{P} + \delta + w \overset{d}{=} x + \delta$$

for $x' \sim \mathbb{Q}$ and $x \sim \mathbb{P}$ with $\mu_\mathbb{Q} - \mu_\mathbb{P} = \delta$ and hence

$$\tilde{\mathbf{R}}_\epsilon(\theta) = \max_{\mathbb{Q} \in \mathcal{V}_p(\epsilon)} \mathbb{E}_{x \sim \mathbb{Q}} \ell_{\text{test}}(\langle\theta, x\rangle, \langle\theta^\star, x\rangle) = \max_{\|\delta\|_p \le \epsilon, \delta \perp \theta^\star} \mathbb{E}_{x \sim \mathbb{P}} \ell_{\text{test}}(\langle\theta, x + \delta\rangle, \langle\theta^\star, x\rangle)$$

$$\le \mathbb{E}_{x \sim \mathbb{P}} \max_{\|\delta\|_p \le \epsilon, \delta \perp \theta^\star} \ell_{\text{test}}(\langle\theta, x + \delta\rangle, \langle\theta^\star, x\rangle) = \mathbf{R}_\epsilon(\theta)$$

where the first line follows from orthogonality of the mean-shift to $\theta^\star$. □

B. Neural networks on sanitized binary MNIST

Figure 4. Increase in robust test accuracy when using early stopping for single hidden layer neural networks with different widths on a subset of MNIST classes 1 vs 3. The curves depict the differences of the test robust accuracy between the best and last iteration of training. Even if we discard training samples that are difficult to fit (pruned data), early stopping yields strictly positive benefits for the robust accuracy.

Figure 4 shows that robust overfitting in the overparameterized regime also occurs for single hidden layer neural networks on an image classification problem that we chose to be arguably devoid of noise. We consider binary classification on MNIST classes 1 vs 3 and further reduce variance by removing "difficult" samples: We train networks of width $p \in \{10, 100, 10k\}$ on the full MNIST training dataset and discard all images that take at least one of the models more than 100 epochs of training to learn. While some recent work argues that such sanitation procedures can effectively mitigate robust overfitting (Dong et al., 2021), we still observe a significant increase from the final test robust accuracies to the best (early stopped) ones in Figure 4.

We train all networks on a subset of $n = 2000$ training samples using vanilla mini-batch stochastic gradient descent with learning rate $\nu_p = \sqrt{0.1/p}$ that we multiply by 0.1 after 300 epochs. This learning rate schedule minimizes the training loss efficiently; we did not perform tuning using test or validation data. For the robust test error, we approximate worst-case ℓ_∞ perturbations using 10-step SGD attacks on each test sample.

C. Experimental details

In this section we provide additional details on our experiments. All our code including instructions and hyperparameters can be found here: `We_will_add_the_link_here_in_the_camera_ready_version.`

C.1. Empirical predictions

If not mentioned otherwise, all empirical simulations use noiseless i.i.d. samples from our synthetic data model as described in Section 2.1. Whenever we simulate label noise, we flip a fixed percentage of all training labels chosen uniformly at random. However, we calculate all risks in closed-form without noise and, in the robust case, with consistent perturbations. We approximate the integral for the robust 0-1 risk in Theorem 3.1 using a numerical integral solver since we cannot obtain a solution analytically.

We fit all logistic regression models except in Figure 2c by minimizing the (regularized) robust logistic loss (3) using the *CVXPY* library in combination with the *Mosek* convex programming solver. To obtain the max-margin solution whenever feasible for $\lambda \to 0$, we optimize the constrained problem (2) directly since (3) with $\lambda = 0$ has many optimal solutions.

In Figure 1, we use $\epsilon = 0.1$ for the robust risk and consistent adversarial training on $n = 1000$ samples. In the noisy case, we flip 2% of all training labels. For Figure 2c, we run zero-initialized gradient descent on the unregularized loss ($\lambda = 0$) for $500k$ iterations. We start with a small initial step size of 0.01 that we double every $30k$ steps until iteration $300k$.

C.2. Theoretical predictions

In order to obtain the asymptotic theoretical predictions for logistic regression in Figure 2a corresponding to the empirical simulations with $n = 1000$ and $\epsilon = 0.05$, we obtain the solution of the optimization problems in Theorem D.1,D.2 with $\epsilon_0 = 0.05\sqrt{1000\gamma}$ by solving the system of equations in Corollary D.4,D.5 using *MATLAB*'s optimization toolbox where we approximate expectations via numerical integration.

D. Details on Theorem 3.1

In this section we give a formal statement for Theorem 3.1. The results are based on the Convex Gaussian Minimax Theorem (CGMT)(Gordon, 1988; Thrampoulidis et al., 2015). The results presented in this section have similarities with the ones in (Javanmard & Soltanolkotabi, 2020), however, we study the discriminative data model with features drawn from a single Gaussian and 1-sparse ground truth. In contrast, in the paper (Javanmard & Soltanolkotabi, 2020), the authors study a generative data model with features drawn from two Gaussians. Furthermore, while not for adversarial training, several papers have already studied standard logistic regression for isotropic Gaussian features in high dimensions (Salehi et al., 2019; Sur & Candès, 2019).

An immediate consequence of the proof of Lemma A.1 is that the adversarial loss from Equation (3) writes as

$$\mathcal{L}_{\epsilon,\lambda}(\theta) = \frac{1}{n}\sum_{i=1}^{n}\ell_{\text{train}}(y_i\langle\theta, x_i\rangle - \epsilon\|\Pi_\perp\hat{\theta}\|_1) + \lambda\|\theta\|_2^2. \tag{9}$$

Note that the equation holds for consistent ℓ_∞ perturbations with logistic loss $\ell_{\text{train}}(x) = \log(1 + \exp(-x))$ and 1-sparse ground truth.

Let $\mathcal{M}_f(x, t) = \min_y \frac{1}{2t}(x - y)^2 + f(y)$ be the Moreau envelope and let $Z_\parallel, Z_\perp$ be two independent standard normal distributed random variables. We can now state Theorem D.1 which describes the asymptotic risk of $\hat{\theta}_\lambda$, for $\lambda > 0$, and for the asymptotic regime where $d, n \to \infty$. The proof of the theorem can be found in Appendix D.4.

Theorem D.1. *Assume that the we have i.i.d.random features x_i drawn form an isotropic Gaussian with noiseless observations $y_i = \text{sgn}(\langle x_i, \theta^\star\rangle)$ and ground truth $\theta^\star = (1, 0, \ldots, 0)^\top$. Further, assume that $\lambda > 0$ and $\epsilon = \epsilon_0/\sqrt{d}$, where ϵ_0 is a numerical constant. Let $(\nu_\perp^\star, \nu_\parallel^\star, r^\star, \delta^\star, \mu^\star, \tau^\star)$ be the unique solution of*

$$\min_{\substack{\nu_\perp \geq 0, \tau \geq 0, \\ \nu_\parallel, \delta \geq 0}} \max_{\substack{r \geq 0, \\ \mu \geq 0}} \mathbb{E}_{Z_\parallel, Z_\perp}\left[\mathcal{M}_\ell\left(|Z_\parallel|\nu_\parallel + Z_\perp\nu_\perp - \epsilon_0\delta, \frac{\tau}{r}\right)\right] - \delta\mu + \frac{r\tau}{2} + \lambda(\nu_\perp^2 + \nu_\parallel^2)$$

$$-\nu_\perp\sqrt{\left[(\mu^2 + \gamma r^2) - (\mu^2 + \gamma r^2)\text{erf}(\mu/(\sqrt{\gamma}r\sqrt{2})) - \sqrt{\frac{2}{\pi}}\sqrt{\gamma}r\mu\exp(-\mu^2/(\gamma r^2 2))\right]}. \tag{10}$$

Then, for $\lambda > 0$ the estimator $\hat{\theta}_\lambda$ from Equation (3) with logistic loss and consistent ℓ_∞ attacks satisfies asymptotically as $d, n \to \infty$, $d/n \to \gamma$

$$\frac{1}{\sqrt{d}}\|\Pi_\perp\hat{\theta}_\lambda\|_1 \to \delta \text{ and } \langle\hat{\theta}_0(\epsilon), \theta^\star\rangle \to \nu_\parallel \text{ and } \|\hat{\theta}_\lambda\|_2^2 \to \nu_\parallel^2 + \nu_\perp^2. \tag{11}$$

The convergences hold true in probability.

For $\lambda > 0$, the loss in Equation (9) has a unique minimizer. In contrast, for $\lambda = 0$ the minimizer of the loss in Equation (9) is not unique since the observations are noiseless and the adversarial attacks are consistent. As discussed in Section 3, we study instead the penalized max-margin solution from Equation (2). The asymptotic behavior of the corresponding solution is characterized in Theorem D.2 with proof in Appendix D.5.

Theorem D.2. *Assume that the we have i.i.d.random features x_i drawn form an isotropic Gaussian with noiseless observations $y_i = \text{sgn}(\langle x_i, \theta^\star\rangle)$ and $\theta^\star = (1, 0, \cdots, 0)^\top$ as describes in Section 3. Further, assume that $\epsilon = \epsilon_0/\sqrt{d}$. Let*

$(\nu_{\perp}^{\star}, \nu_{\parallel}^{\star}, r^{\star}, \delta^{\star}, \zeta^{\star}, \kappa^{\star})$ *be the unique solution of*

$$\min_{\substack{\nu_{\perp} \geq 0, \\ \delta \geq 0, \kappa, \nu_{\parallel}}} \max_{\zeta, r \geq 0} \nu_{\parallel}^2 - \kappa \nu_{\perp}^2 - \delta\zeta - \frac{\gamma r^2}{4(1+\kappa)} + r\sqrt{\mathbb{E}_{Z_{\parallel}, Z_{\perp}} \left[\max\left(0, 1 + \epsilon_0 \delta - |Z_{\parallel}|\nu_{\parallel} + Z_{\perp}\nu_{\perp}\right)^2\right]}$$

$$+ \frac{1}{2(1+\kappa)}\left(\frac{\gamma r^2 + \zeta^2}{2}\mathrm{erf}\left(\frac{\zeta}{\sqrt{2}\sqrt{\gamma}r}\right) - \frac{\zeta^2}{2} + \frac{\sqrt{\gamma}r\zeta}{\sqrt{2\pi}}\exp\left(-\frac{\zeta^2}{2\gamma r^2}\right)\right). \quad (12)$$

Then, the estimator $\hat{\theta}_0(\epsilon)$ from Equation (2) with logistic loss and consistent ℓ_{∞} attacks satisfies asymptotically as $d, n \to \infty$, $d/n \to \gamma$

$$\frac{1}{\sqrt{d}}\|\Pi_{\perp}\hat{\theta}_0(\epsilon)\|_1 \to \delta \text{ and } \langle\hat{\theta}_0(\epsilon), \theta^{\star}\rangle \to \nu_{\parallel} \text{ and } \|\hat{\theta}_0(\epsilon)\|_2^2 \to \nu_{\parallel}^2 + \nu_{\perp}^2. \quad (13)$$

The convergences hold true in probability.

Remark D.3. *Theorem 3.1 is obtained form Theorem D.1,D.2 when inserting the expression from Equation (11),(13) into the expression of the risk in Lemma A.1.*

D.1. Solving the optimization problems

To obtain concrete predictions from Theorems D.1 and D.2 we need to solve the corresponding optimization problems. Using the fact that the objective is concave in r, μ respectively r, τ_r and convex in all other variables, we find the unique solution by solving the systems of equations stated in Corollary D.4,D.5 presented below — assuming that the solution is not obtained on the boundary. For numerical predictions, we solve the systems of equations using a least squares solver in *MATLAB* where we approximated the expectations using a standard numerical integrator.

Regularization $\lambda > 0$: We make use of the following well known facts on the derivative of the Moreau envelope (e.g. see (Jourani et al., 2014)):

$$\frac{\partial \mathcal{M}_f(x,t)}{\partial x} = \frac{1}{t}(x - \mathrm{Prox}_f(x,t)) \text{ and } \frac{\partial \mathcal{M}_f(x,t)}{\partial t} = -\frac{1}{2t^2}(x - \mathrm{Prox}_f(x,t))^2$$

where $\mathrm{Prox}_f(x,\mu)$ is the proximal operator

$$\mathrm{Prox}_f(x,\mu) = \arg\min_t f(t) + \frac{1}{2\mu}(t-x)^2.$$

Let $M = \mathbb{E}_{Z_{\parallel}, Z_{\perp}}\left[\mathcal{M}_{\ell}(Z_{\parallel}\nu_{\parallel} + Z_{\perp}\nu_{\perp} - \epsilon\delta, \frac{\tau}{r})\right]$ and denote with $M_{\nu_{\parallel}} = \mathbb{E}_{Z_{\parallel}, Z_{\perp}}\left[\frac{\partial_x \mathcal{M}_{\ell}}{\partial x}(Z_{\parallel}\nu_{\parallel} + Z_{\perp}\nu_{\perp} - \epsilon\delta, \frac{\tau}{r})\frac{\partial(Z_{\parallel}\nu_{\parallel} + Z_{\perp}\nu_{\perp} - \epsilon\delta)}{\partial \nu_{\perp}}\right]$ the derivative of the expected Moreau envelope with respect to $\nu_{\parallel}$ and similar for the other variables.

Further, define $g = (\mu^2 + \gamma r^2) - (\mu^2 + \gamma r^2)\mathrm{erf}(\mu/(\sqrt{\gamma}r\sqrt{2})) - \sqrt{\frac{2}{\pi}}\sqrt{\gamma}r\mu\exp(-\mu^2/(\gamma r^2 2))$ and denote with g_{μ} respectively g_r the corresponding partial derivatives. This allows us to express the optimization objective form Theorem D.1 as

$$C = M - \nu_{\perp}\sqrt{g(\mu,r)} - \delta\mu + \frac{r\tau}{2} + \lambda(\nu_{\perp}^2 + \nu_{\parallel}^2).$$

Since the problem is concave in r, μ and convex in the other variables, we can equivalently solve the the system of equations $\nabla C = 0$, which gives us the following corollary.

Corollary D.4. *Assume that the solution of Equation (10) form Theorem D.1 is not obtained on the boundary. Then, the*

solution satisfies the following system of Equations.

$$0 = \frac{\partial}{\partial \nu_\|}C = M_{\nu_\|} + 2\lambda\nu_\|$$

$$0 = \frac{\partial}{\partial \nu_\perp}C = M_{\nu_\perp} + 2\lambda\nu_\perp - \sqrt{g}$$

$$0 = \frac{\partial}{\partial r}C = M_r - \frac{\nu_\perp}{2\sqrt{g}}g_r + \frac{\tau}{2}$$

$$0 = \frac{\partial}{\partial \delta}C = M_\delta - \mu$$

$$0 = \frac{\partial}{\partial \mu}C = -\frac{\nu_\perp}{2\sqrt{g}}g_\mu - \delta$$

$$0 = \frac{\partial}{\partial \tau}C = M_\tau + \frac{r}{2}$$

D.2. Interpolation $\lambda = 0$:

Similarly, we can also reformulate the optimization problem in Theorem D.2. Again, for brevity of notation, let $T = \mathbb{E}_{Z,Z_\sigma}\left(1 + \epsilon_0\delta - Z_\sigma\nu_\| + Z\nu_\perp\right)_+^2$ and $H = \left(\frac{\gamma r^2 + \zeta^2}{2}\mathrm{erf}\left(\frac{\zeta}{\sqrt{2}\sqrt{\gamma}r}\right) - \frac{\zeta^2}{2} + \frac{\sqrt{\gamma}r\zeta}{\sqrt{2\pi}}\exp\left(-\frac{\zeta^2}{2\gamma r^2}\right)\right)$. Further, denote with $T_{\nu_\perp}$ the derivative with respect to the corresponding variable and similarly for H_r. As a result, we can write the optimization objective from Theorem D.2 as

$$C = \nu_\|^2 - \kappa\nu_\perp^2 - \delta\zeta - \frac{\gamma r^2}{4s} + 2sH + r\sqrt{T} + \frac{\tau_r\kappa}{2}.$$

Since the solution of the optimization problem satisfies again $\nabla C = 0$, we analogously obtain the following corollary.

Corollary D.5. *Assume that the solution of Equation* (12) *form Theorem D.2 is not obtained on the boundary. Then, the solution satisfies the following system of Equations.*

$$0 = \frac{\partial}{\partial \nu_\|}C = 2\nu_\| + \frac{1}{2\sqrt{T}}rT_{\nu_\|}$$

$$0 = \frac{\partial}{\partial \nu_\perp}C = -2\kappa\nu_\perp + \frac{1}{2\sqrt{T}}rT_{\nu_\perp}$$

$$0 = \frac{\partial}{\partial r}C = \sqrt{T} + H_r - \frac{\gamma r}{2(1+\kappa)}$$

$$0 = \frac{\partial}{\partial \delta}C = -\zeta + \frac{1}{2\sqrt{T}}rT_\delta$$

$$0 = \frac{\partial}{\partial \zeta}C = -\delta + H_y$$

$$0 = \frac{\partial}{\partial \kappa}C = -\nu_\perp^2 + H_\kappa + \frac{\gamma r^2}{4(1+\kappa)^2}$$

D.3. Label noise

While our results assume noiseless observations $y_i = \mathrm{sgn}(\langle x_i, \theta^\star\rangle)$, Theorem D.1,D.2 can be extended to the case where additional label noise is added to the observations. That is, it can be extended to the case where $y_i = \mathrm{sgn}(\langle x_i, \theta^\star\rangle)\xi$ with $\mathbb{P}(\xi = 1) = 1 - \sigma$ and $\mathbb{P}(\xi = -1) = \sigma$. Note that the only modification need for both theorems is to replace $|Z_\||$ with $\xi|Z_\||$.

However, as discussed in Section 4.1, label noise in the train data may prevent interpolation even when $\lambda = 0$ and hence Equation (9) may also have a unique bounded solution for $\lambda = 0$. In fact, we note that the asymptotic threshold $\gamma^\star$ such that the train loss does not vanish with high probability for all $\gamma < \gamma^\star$ and vanishes with high probability for all $\gamma \geq \gamma^\star$ as $d/n \to \gamma$ can be found when solving a low dimensional scalar optimization problem similar to Theorem 6.1 (Javanmard & Soltanolkotabi, 2020).

D.4. Proof of Theorem D.1

The proof is similar to the proof of Theorem 6.4 (Javanmard & Soltanolkotabi, 2020). Denote with $X \in \mathbb{R}^{n \times d}$ the input data matrx and with $y \in \mathbb{R}^n$ the vector containing the observations. Recall that the estimator $\hat{\theta}$ is given by

$$\hat{\theta} = \arg\min_{\theta} \frac{1}{n} \sum_{i=1}^{n} \ell(y_i \langle x_i, \theta \rangle - \epsilon \|\Pi_{\perp}\theta\|_1) + \lambda\|\theta\|_2^2$$

$$= \arg\min_{\theta} \frac{1}{n} \sum_{i=1}^{n} \ell(v_i - \epsilon\|\Pi_{\perp}\theta\|_1) + \lambda\|\theta\|_2^2 \text{ such that } v = D_y X\theta, \tag{14}$$

where $\ell(x) = \log(1 + \exp(-x))$ is the logistic loss, $X \in \mathbb{R}^{n \times d}$ is the data matrix and D_y the diagonal matrix with entries $(D_y)_{i,i} = y_i$. We can then introduce the Lagrange multiplier u to obtain:

$$\min_{\theta,v} \max_{u} \frac{1}{n} \sum_{i=1}^{n} \ell(v_i - \epsilon\|\Pi_{\perp}\theta\|_1) + \frac{1}{n}u^{\top}D_y X\theta - \frac{1}{n}u^{\top}v + \lambda\|\theta\|_2^2.$$

Furthermore, we can separate $X = X\Pi_{\perp} + X\Pi_{\|}$, which gives us:

$$\min_{\theta,v} \max_{u} \frac{1}{n} \sum_{i=1}^{n} \ell(v_i - \epsilon\|\Pi_{\perp}\theta\|_1) + \frac{1}{n}u^{\top}D_y X\Pi_{\|}\theta + \frac{1}{n}u^{\top}D_y X\Pi_{\perp}\theta - \frac{1}{n}u^{\top}v + \lambda\|\theta\|_2^2.$$

Convex Gaussian Minimax Theorem We can now make use of the CGMT, which states that

$$\min_{\theta \in U_{\theta}} \max_{u \in U_u} u^{\top}X\theta + \psi(u,\theta), \tag{15}$$

with ψ convex in θ and concave in u has asymptotically point wise the same solution when $d, n \to \infty$, $d/n \to \gamma$ as

$$\min_{\theta \in U_{\theta}} \max_{u \in U_u} \|u\|_2 g^{\top}\theta + u^{\top}h\|\theta\|_2 + \psi(u,\theta), \tag{16}$$

with $g \in \mathbb{R}^n$, $u \in \mathbb{R}^d$ random vectors with i.i.d. standard normal distributed entries. Here, the sets U_{θ} and U_u are compact. As commonly referred to in the literature, we call Equation (20) the primal optimization problem and Equation (16) the auxiliary optimization problem. We refer to (Thrampoulidis et al., 2015) for the precise statement.

Since $X\Pi_{\perp}$ is independent of the observations y and of $X\Pi_{\|}$, and because Equation (14) has a unique bounded solution as $\lambda > 0$, we can apply the CGMT, which gives:

$$\min_{\theta,v} \max_{u} \frac{1}{n} \sum_{i=1}^{n} \ell(v_i - \epsilon\|\Pi_{\perp}\theta\|_1) + \frac{1}{n}u^{\top}D_y X\Pi_{\|}\theta + \frac{1}{n}\|u^{\top}D_y\|_2 g^{\top}\Pi_{\perp}\theta$$

$$+ \frac{1}{n}u^{\top}D_y h\|\Pi_{\perp}\theta\|_2 - \frac{1}{n}u^{\top}v + \lambda\|\theta\|_2^2,$$

where $g \in \mathbb{R}^d$ and $h \in \mathbb{R}^n$ are vectors with i.i.d. standard normal distributed entries.

Scalarization of the optimization problem We now aim to simplify the optimization problem. In a first step, we maximize over u. For this, define $r = \|u\|_2/\sqrt{n}$, which allows us to equivalently write:

$$\min_{\theta,v} \max_{r \geq 0} \frac{1}{n} \sum_{i=1}^{n} \ell(v_i - \epsilon\|\Pi_{\perp}\theta\|_1) + \frac{r}{\sqrt{n}}\|D_y X\Pi_{\|}\theta + D_y h\|\Pi_{\perp}\theta\|_2 - v\|_2 + \frac{1}{\sqrt{n}}rg^{\top}\Pi_{\perp}\theta + \lambda\|\theta\|_2^2,$$

where we have used the fact that $\|u^{\top}D_y\|_2 = \|u\|_2$. Next, we note that because the problem is convex in r and concave in θ, v, we can swap maximization with minimization. In order to proceed, we want to separate $\Pi_{\perp}\theta$ form the loss $\ell(v, \Pi_{\perp}\theta) := \frac{1}{n} \sum_{i=1}^{n} \ell(v_i - \epsilon\|\Pi_{\perp}\theta\|_1)$. We can write $\ell(v, \Pi_{\perp}\theta)$ in terms of its conjugate with respect to $\Pi_{\perp}\theta$:

$$\ell(v, \Pi_{\perp}\theta) = \sup_{w} \frac{1}{\sqrt{d}}w^{\top}\Pi_{\perp}\theta - \tilde{\ell}(v, w),$$

where $\tilde{\ell}$ is the conjugate of ℓ. In particular, we can use Lemma A.2 in the paper (Javanmard & Soltanolkotabi, 2020) which shows that

$$\tilde{\ell}(v, w) = \sup_{\delta \geq 0} \frac{\sqrt{d}}{\sqrt{d}} \delta \|w\|_\infty - \frac{1}{n} \sum_{i=1}^n \ell(v_i - \sqrt{d}\epsilon\delta) = \sup_{\delta \geq 0} \delta \|w\|_\infty - \frac{1}{n} \sum_{i=1}^n \ell(v_i - \epsilon_0\delta).$$

Interchanging the order of maximization over w and minimization over δ, we get

$$\max_{r \geq 0} \min_{\substack{\theta, v, \\ \delta \geq 0}} \frac{1}{n} \sum_{i=1}^n \ell(v_i - \epsilon_0\delta) + \frac{r}{\sqrt{n}} \left\| D_y X \Pi_\| \theta + D_y h \|\Pi_\perp \theta\|_2 - v \right\|_2 + \lambda \|\theta\|_2^2 \tag{17}$$

$$+ \max_w \left[\frac{1}{\sqrt{d}} w^\top \Pi_\perp \theta - \delta \|w\|_\infty + \frac{1}{\sqrt{n}} r g^\top \Pi_\perp \theta \right]$$

where we used again the fact that we can interchange the order maximization and minimization. Next, we simplify the optimization over θ. Write $\Pi_\| \theta = \Pi_\| 1 \nu_\|$ where $\nu_\| \in \mathbb{R}$ and let $\nu_\perp = \|\Pi_\perp \theta\|_2$. We can simplify

$$\max_{r \geq 0} \min_{\substack{\nu_\perp \geq 0, \\ \delta \geq 0, \\ \nu_\|, v}} \frac{1}{n} \sum_{i=1}^n \ell(v_i - \epsilon_0\delta) + \frac{r}{\sqrt{n}} \|D_y X \Pi_\| 1 \nu_\| + D_y h \nu_\perp - v\|_2 + \lambda(\nu_\|^2 + \nu_\perp^2) \tag{18}$$

$$+ \max_w \left[-\frac{1}{\sqrt{d}} \nu_\perp \|\Pi_\perp (w - \sqrt{\gamma} r g)\|_2 - \delta \|w\|_\infty \right]$$

In order to obtain a low dimensional scalar optimization problem, we still need to scalarize the optimization over w and v. For this, we replace the term $\|D_y X \Pi_\| 1 \nu_\| + D_y h \nu_\perp - v\|_2$ with its square, which is achieved by using the following identity $\min_{\tau \geq 0} \frac{x^2}{2\tau} + \frac{\tau}{2} = x$. Hence,

$$\max_{r \geq 0} \min_{\substack{\nu_\perp \geq 0, \tau \geq 0, \\ \delta \geq 0, \\ \nu_\|, v}} \frac{1}{n} \sum_{i=1}^n \ell(v_i - \epsilon_0\delta) + \frac{r}{2\tau n} \|D_y X \Pi_\| 1 \nu_\| + D_y h \nu_\perp - v\|_2^2 + \frac{\tau r}{2} + \lambda(\nu_\|^2 + \nu_\perp^2)$$

$$+ \max_w \left[-\frac{1}{\sqrt{d}} \nu_\perp \|\Pi_\perp (w - \sqrt{\gamma} r g)\|_2 - \delta \|w\|_\infty \right]$$

interchanging again the order of maximization and minimization, we can now separately solve the following two inner optimization problems:

$$\max_w \; -\nu_\perp \frac{1}{d} \|\Pi_\perp (w - \sqrt{\gamma} r g)\|_2 - \delta \|w\|_\infty$$

$$\min_v \; \frac{r}{2\tau n} \|D_y X \Pi_\| 1 \nu_\| + D_y h \nu_\perp - v\|_2^2 + \sum_{i=1}^n \ell(v_i - \epsilon_0\delta)$$

First term   Let $\mathrm{ST}_t(x) = \begin{cases} 0 & |x| \leq t \\ \mathrm{sgn}(x)(|x| - t) & \text{else} \end{cases}$ be the soft threshold function. We have

$$\max_w \; -\nu_\perp \frac{1}{\sqrt{d}} \|\Pi_\perp (w - \sqrt{\gamma} r g)\|_2 - \delta \|w\|_\infty$$

$$= -\min_w \nu_\perp \frac{1}{\sqrt{d}} \|\Pi_\perp (w - \sqrt{\gamma} r g)\|_2 - \delta \|w\|_\infty$$

$$\overset{\mu = \|w\|_\infty}{=} -\min_{\mu \geq 0} \nu_\perp \sqrt{\frac{1}{d} \sum_{i=2}^d \left(\mathrm{ST}_\mu \left(\sqrt{\gamma} r g_i \right) \right)^2} + \delta\mu$$

$$\overset{\text{LLN as } d \to \infty}{\to} -\min_{\mu \geq 0} \nu_\perp \sqrt{\mathbb{E}_Z \left(\mathrm{ST}_\mu \left(\sqrt{\gamma} r Z \right) \right)^2} + \delta\mu,$$

where we used in the third line that the ground truth $\theta^\star$ is 1-sparse and in the last line that the expectation exists. Finally, we can further simplify

$$
\begin{aligned}
\mathbb{E}_Z\left(\mathrm{ST}_\mu\left(\sqrt{\gamma}rZ\right)\right)^2 &= \gamma r^2 \mathbb{E}_Z\left(\mathrm{ST}_{\mu/(\sqrt{\gamma}r)}(Z)\right)^2 \\
&= \gamma r^2 \mathbb{E}_Z(Z - \mu/(\sqrt{\gamma}r))^2 - \mathbb{E}_Z \mathbb{1}_{|Z|\leq\mu/(\sqrt{\gamma}r)}(Z - \mu/(\sqrt{\gamma}r))^2 \\
&= (\mu^2 + \gamma r^2)\left(1 - \mathrm{erf}(\mu/(\sqrt{2\gamma}r))\right) - \sqrt{\gamma}r\mu\sqrt{\frac{2}{\pi}}\exp(-\mu/(2\gamma r^2)).
\end{aligned}
$$

Hence, we can conclude the first term.

Second term For the second term we also aim to apply the law of large numbers. We have

$$
\min_v \frac{r}{2\tau n}\|D_y X\Pi_\| 1\nu_\| + D_y h\nu_\perp - v\|_2^2 + \frac{1}{n}\sum_{i=1}^n \ell(v_i - \epsilon_0\delta)
$$

$$
\overset{\tilde{v}=v-\epsilon\delta}{=} \min_{\tilde{v}} \frac{r}{2\tau n}\|D_y X\Pi_\| 1\nu_\| + D_y h\nu_\perp - \tilde{v} - \epsilon_0\delta\|_2^2 + \frac{1}{n}\sum_{i=1}^n \ell(\tilde{v}_i)
$$

$$
\overset{\tilde{v}=v-\epsilon\delta}{=} \min_{\tilde{v}} \frac{1}{n}\sum_{i=1}^n \frac{r}{2\tau n}\left((D_y X\Pi_\| 1\nu_\|)_i + (D_y h\nu_\perp)_i - \tilde{v}_i - \epsilon_0\delta\right)^2 + \ell(\tilde{v}_i)
$$

$$
\overset{\mathrm{LLN}}{\to} \mathbb{E}_{Z_\|, Z_\perp}\left[\mathcal{M}_\ell(|Z_\|| \nu_\| + Z_\perp\nu_\perp - \epsilon_0\delta, \frac{\tau}{r})\right],
$$

where in the last line we used that $(D_y X\Pi_\| \Pi_\perp 1)_i = y_i x_i^\top \theta^\star = \xi_\sigma \mathrm{sgn}(x_i^\top \theta^\star) x_i^\top \theta^\star$ which has the same distribution as $|Z_\||$. Further, to apply the law of large numbers, we need to show that the Moreau envelope exists. Similarly to Theorem 1 (Salehi et al., 2019), this follows immediately when noting that $\mathcal{M}_\ell(x, \mu) \leq \ell(x) = \log(1 + \exp(-x)) \leq \log(2) + |x|$. Finally, we obtain the desired optimization problem when combining these results.

Uniform convergences So far, we only proved point wise convergence to the same solution of the primal 14 and auxiliary 10 optimization problem. However, since we also optimize over v, we need to show uniform convergence to guarantee that the solutions are asymptotically the same. This follows when using similar arguments as in Lemma A.5 (Thrampoulidis et al., 2015). Finally, note that the optima $\delta^\star$ in Equation (17) satisfies $\delta^\star = \frac{1}{\sqrt{d}}\|\Pi_\perp\theta\|$, and similarly the optima $\nu_\perp^\star$ and $\nu_\|^\star$ in Equation (18) satisfy $\nu_\perp^\star = \|\Pi_\perp\theta\|_2$ and $\nu_\|^\star = \langle\theta, \theta^\star\rangle$. Hence we can conclude the proof as the solutions 10, (14) converge asymptotically to the same optima.

D.5. Proof of Theorem D.2

The proof is similar to the proof of Theorem 6.3 (Javanmard & Soltanolkotabi, 2020). Recall the penalized max margin solution form Equation (2):

$$
\min_\theta \|\theta\|_2^2 \text{ sucht that forall } i : \langle\theta, x_i\rangle - \epsilon\|\Pi_\perp\theta\|_1 \geq 1 \tag{19}
$$

When introducing the Lagrange multiplier ζ and u we can write equivalently write:

$$
\min_{\theta, \delta} \max_{\substack{u:u_i\geq 0, \\ \zeta\geq 0}} \|\theta\|_2^2 + \frac{1}{n}u^\top\left(1 + 1\epsilon_0\delta - D_y X\theta\right) + \zeta\left(\frac{\|\Pi_\perp\theta\|_1}{\sqrt{d}} - \delta\right).
$$

and separating X again into $X\Pi_\perp + X\Pi_\|$, we get

$$
\min_{\theta, \delta} \max_{\substack{u:u_i\geq 0, \\ \zeta\geq 0}} \|\theta\|_2^2 + \frac{1}{n}u^\top\left(1 + 1\epsilon_0\delta - D_y X\Pi_\|\theta - D_y X\Pi_\perp\theta\right) + \zeta\left(\frac{\|\Pi_\perp\theta\|_1}{\sqrt{d}} - \delta\right) \tag{20}
$$

Convex Gaussian Minimax Theorem Since the adversarial attacks are consistent and the observations are noiseless, we know the solution in Equation (19) exists for all d, n. Yet, in order to apply the CGMT, we have to show that we can restrict u and θ to compact sets. This follows for a simple trick as explained in Section D.3.1 (Javanmard & Soltanolkotabi, 2020).

Hence, the primal optimization problem from Equation (20) can be asymptotically replaced with the following auxiliary optimization:

$$\min_{\theta,\delta} \max_{\substack{u:u_i \geq 0, \\ \zeta \geq 0}} \|\theta\|_2^2 + \frac{1}{n}u^\top \left(1 + 1\epsilon_0\delta - D_y X\Pi_\| \theta + D_y h\|\Pi_\perp \theta\|_2\right) + \frac{1}{n}\|u\|_2 g^\top \Pi_\perp \theta + \zeta\left(\frac{\|\Pi_\perp \theta\|_1}{\sqrt{d}} - \delta\right)$$

Scalarization of the optimization problem The goal is again to scalarize the optimization problem. As a first step, we can solve the optimization over u when defining $r = \frac{\|u\|_2}{\sqrt{n}}$:

$$\min_{\theta,\delta} \max_{\substack{r \geq 0, \\ \zeta \geq 0}} \|\theta\|_2^2 + \frac{r}{\sqrt{n}}\| \max\left(0, 1 + 1\epsilon_0\delta - D_y X\Pi_\| \theta + D_y h\|\Pi_\perp \theta\|_2\right) \|_2$$

$$+ \frac{r\sqrt{\gamma}}{\sqrt{d}}\|u\|_2 g^\top \Pi_\perp \theta + \zeta\left(\frac{1}{\sqrt{d}}\|\Pi_\perp \theta\|_1 - \delta\right),$$

where $\max$ is taken elementwise over the vector. We now want to separate $\|\Pi_\perp \theta\|_2$ form the term in $\max(.)$. This is achieved by introducing the variables $\nu_\perp \geq 0$ and the Lagrange multiplier κ. Further, we set $\nu_\| = \langle \theta^\star, \Pi_\| \theta\rangle$, which allows us to equivalently write

$$\min_{\substack{\nu_\perp \geq 0, \\ \nu_\|,\delta}} \max_{\substack{r \geq 0, \\ \zeta \geq 0,\kappa}} \nu_\|^2 + \|\Pi_\perp \theta\|_2^2 + \kappa(\|\Pi_\perp \theta\|_2^2 - \nu_\perp^2) + \frac{r}{\sqrt{n}}\| \max\left(0, 1 + 1\epsilon_0\delta - D_y X\Pi_\| \theta_0\nu_\| + D_y h\nu_\perp\right) \|_2$$

$$+ \frac{r\sqrt{\gamma}}{\sqrt{d}}\|u\|_2 g^\top \Pi_\perp \theta + \zeta\left(\frac{1}{\sqrt{d}}\|\Pi_\perp \theta\|_1 - \delta\right)$$

We further note that

$$\frac{r}{\sqrt{n}}\| \max\left(0, 1 + 1\epsilon_0\delta - D_y X\Pi_\| \theta_0\nu_\| + D_y h\nu_\perp\right) \|_2$$

$$\overset{\text{LLN}}{\to} r\sqrt{\mathbb{E}_{Z_\|,Z_\perp}\left[\max\left(0, 1 + \epsilon_0\delta - |Z_\||\nu_\| + Z_\perp\nu_\perp\right)^2\right]} =: \sqrt{T}.$$

Next, by completion of the squares we get

$$\min_{\substack{\nu_\perp \geq 0, \\ \nu_\|,\delta}} \max_{\substack{r \geq 0, \\ \zeta \geq 0,\kappa}} \nu_\|^2 + (1+\kappa)\|\Pi_\perp \theta + \frac{r\sqrt{\gamma}}{\sqrt{d}2(1+\kappa)}g\|_2^2 - \frac{r^2\gamma}{4(1+\kappa)}\|g/\sqrt{d}\|_2^2$$

$$- \kappa\nu_\perp^2 + \sqrt{T} + \zeta\left(\frac{1}{\sqrt{d}}\|\Pi_\perp \theta\|_1 - \delta\right),$$

with $\|g/\sqrt{d}\|_2^2 \to 1$. Hence, it only remains to solve the inner optimization over $\Pi_\perp \theta$. We can write:

$$\min_{\Pi_\perp \theta} (1+\kappa)\|\Pi_\perp \theta + \frac{r\sqrt{\gamma}}{\sqrt{d}2(1+\kappa)}g\|_2^2 + \zeta\frac{\|\Pi_\perp \theta\|_1}{\sqrt{d}}$$

$$\overset{\tilde{\theta}_\perp = \frac{\Pi_\perp \theta}{\sqrt{d}}}{=} \min_{\tilde{\theta}_\perp} \frac{1}{d}(1+\kappa)\|\tilde{\theta}_\perp + \frac{r\sqrt{\gamma}}{2(1+\kappa)}g\| + \zeta\frac{\|\Pi_\perp \theta\|_1}{d}$$

$$= \frac{1}{d}\sum_{i=2}^{d} \min_{(\tilde{\theta}_\perp)_i} (1+\kappa)((\tilde{\theta}_\perp)_i + \frac{r\sqrt{\gamma}}{2(1+\kappa)}g_i)^2 + \zeta|(\tilde{\theta}_\perp)_i|$$

$$= \frac{1}{d}2(1+\kappa)\sum_{i=2}^{d} \min_{(\tilde{\theta}_\perp)_i} \frac{1}{2}((\tilde{\theta}_\perp)_i + \frac{r\sqrt{\gamma}}{2(1+\kappa)}g_i)^2 + \frac{\zeta}{2(1+\kappa)}|(\tilde{\theta}_\perp)_i|$$

$$= \frac{1}{d}2(1+\kappa)\sum_{i=2}^{d} \ell_H(-\frac{r\sqrt{\gamma}}{2(1+\kappa)}g_i, \frac{\zeta}{2(1+\kappa)})$$

$$\to 2(1+\kappa)\mathbb{E}_Z \ell_H\left(\frac{r\sqrt{\gamma}}{2(1+\kappa)}Z, \frac{\zeta}{2(1+\kappa)}\right)$$

where ℓ_{H} is the Huber loss, given by $\ell_{\mathrm{H}}(x, y) = \begin{cases} 0.5x^2 & |x| \leq y \\ y(|x| - 0.5y) \end{cases}$. Finally, we can conclude the proof from

$$\mathbb{E}_Z \, \ell_{\mathrm{H}}(aZ, b) = \frac{a^2 + b^2}{2} \mathrm{erf}\left(\frac{b}{\sqrt{2}a}\right) - \frac{b^2}{2} + \frac{ab}{\sqrt{2\pi}} \exp\left(-\frac{b^2}{2a^2}\right)$$