Nonuniform Negative Sampling and Log Odds Correction with Rare Events Data

Abstract

We investigate the issue of parameter estimation with nonuniform negative sampling for imbalanced data. We first prove that, with imbalanced data, the available information about unknown parameters is only tied to the relatively small number of positive instances, which justifies the usage of negative sampling. However, if the negative instances are subsampled to the same level of the positive cases, there is information loss. To maintain more information, we derive the asymptotic distribution of a general inverse probability weighted (IPW) estimator and obtain the optimal sampling probability that minimizes its variance. To further improve the estimation efficiency over the IPW method, we propose a likelihood-based estimator by correcting log odds for the sampled data and prove that the improved estimator has the smallest asymptotic variance among a large class of estimators. It is also more robust to pilot misspecification. We validate our approach on simulated data as well as a real click-through rate dataset with more than 0.3 trillion instances, collected over a period of a month. Both theoretical and empirical results demonstrate the effectiveness of our method.

1 Introduction

Imbalanced data are ubiquitous in scientific fields and applications with binary response outputs, where the number of instances in the positive class is much smaller than that in the negative class. For example, in online search or recommendation systems where billions of impressions appear each day, non-click impressions usually dominate. Using these non-click impressions as negative instances is prevalent and proved to be useful especially in modern machine learning systems [20, 12, 4, 9, 14]. A common approach to address imbalanced data is to balance it by subsampling the negative class [6, 18] and/or up-sampling the positive class [2, 10, 19, 5], and a great deal of attentions have been drawn to this problem, see [15, 16, 3, 7, 22, 25, 1, 24, 8, 17, 28], and the references therein.

In this paper, we focus on subsampling the negative class, i.e., negative sampling, while keeping all rare positive instances. This is beneficial for modern online learning systems, where negative sampling significantly reduces data volume in distributed storage while saving training time for multiple downstream models. Rare events data and negative sampling are studied in [28], but it focused on linear logistic regression and only considered uniform sampling. We approach this problem under general binary response models from the perspective of optimal subsampling, which aims to minimize the asymptotic variance of the IPW estimator [30, 26, 29, 31]. This topic was not well investigated neither for imbalanced data nor for negative sampling. In addition, existing optimal probability formulation minimizes the conditional asymptotic variance and the variations due to the data randomness are ignored. We fill these gaps by deriving the asymptotic distributions for both full data and subsample estimators under general binary response models, and we find the unconditional optimal probability for negative sampling. In addition, since IPW may inflate the variance [13, 21],

we develop a more efficient likelihood-based estimator through nonuniform log odds correction to avoid IPW. Our main contributions are summarized as follows.

- Under a general binary response model with rare events data, we find that the difference between the full data estimator and the true parameter converges to a normal distribution at a rate that is tied to the number of rare positive instances only. This indicates the possibility to throw away the majority of negative instances without information loss, i.e, it justifies the usage of negative sampling.

- We show that there is no asymptotic information loss with aggressive negative sampling if the negative instances kept dominates the positive instances. However there is information loss expressed as a variance inflation if the negative instances are brought down to the same level of the positive instances. For this case, we obtain optimal subsampling probabilities by minimizing the unconditional asymptotic variance of a general IPW estimator.

- We develop a likelihood estimator through nonuniform log odds correction for sampled data, and prove that it has the smallest asymptotic variance among a large class of estimators.

- We apply the proposed method to a real online streaming click-through rate (CTR) dataset with more than 0.3 trillion instances and demonstrate its effectiveness.

The rest of paper is organized as follows. Section 2 defines the problem this paper focuses on and shows the full data results. Section 3 presents the optimal probability for the IPW estimator. Section 4 proposes the likelihood-based estimator and establishes its asymptotic optimality. Section 5 discusses practical implementation and the theoretical properties of the resulting estimators. Section 6 presents numerical experiments with simulated data, and applies the proposed method to a real online streaming CTR dataset with more than 0.3 trillion instances. Section 7 concludes the paper and points out limitations of our investigation. Proofs and additional experiments with misspecifications are in the supplementary materials.

2   Problem setup and full data results

Let $\{(\mathbf{x}_i, y_i)\}_{i=1}^N$ be training data of size N that satisfies the binary response model,

$$\Pr(y = 1 \mid \mathbf{x}) = p(\mathbf{x}; \boldsymbol{\theta}), \tag{1}$$

where $y \in \{0, 1\}$ is the binary class label ($y = 1$ for the positive and $y = 0$ for the negative), $\mathbf{x}$ is the vector of features, and $\boldsymbol{\theta}$ is the unknown d-dimensional parameter.

Let N_1 be the number of positive instances ($N_1 = \sum_{i=1}^N y_i$) and N_0 be the number of negative instances ($N_0 = N - N_1$). We consider the scenario of massive imbalanced data, i.e., $N_1 \ll N_0$. For asymptotic investigations when N_1 is much smaller than N_0, it is more appropriate to assume that N_1 increases in a slower rate compared with N_0, i.e., $N_1/N_0 \to 0$ in probability as $N \to \infty$ [28]. This requires $\Pr(y = 1) = \mathbb{E}\{p(\mathbf{x}; \boldsymbol{\theta})\} \to 0$ as $N \to \infty$ on the model side. We assume that the parameter $\boldsymbol{\theta}$ contains two components $\boldsymbol{\theta} = (\alpha, \boldsymbol{\beta}^{\mathrm{T}})^{\mathrm{T}}$ and the log odds can be written as

$$g(\mathbf{x}; \boldsymbol{\theta}) := \log\left\{\frac{p(\mathbf{x}; \boldsymbol{\theta})}{1 - p(\mathbf{x}; \boldsymbol{\theta})}\right\} = \alpha + f(\mathbf{x}; \boldsymbol{\beta}),$$

where the true parameter, denoted as $\boldsymbol{\theta}^* = (\alpha^*, \boldsymbol{\beta}^{*\mathrm{T}})^{\mathrm{T}}$, satisfies that $\alpha^* \to -\infty$ as $N \to \infty$ and $\boldsymbol{\beta}^*$ is fixed. Here $f(\mathbf{x}; \boldsymbol{\beta})$ is a smooth function of $\boldsymbol{\beta}$, such as a linear function or a nonlinear neural network model. If it is linear, the model reduces to the logistic regression model. A diverging α^* and a fixed $\boldsymbol{\beta}^*$ indicate that both the marginal and conditional probabilities for a positive instance are small. Heuristically, this implies that a change in the feature value does make a rare event become a large probability event. We can also allow $\boldsymbol{\beta}^*$ to change with N, but as long as $\boldsymbol{\beta}^*$ has a finite limit, the situation is essentially the same as a fixed $\boldsymbol{\beta}^*$. So here we assume $\boldsymbol{\beta}^*$ to be fixed to simplify the presentation.

Throughout this paper, we use $\dot{g}(\mathbf{x}; \boldsymbol{\theta})$ and $\ddot{g}(\mathbf{x}; \boldsymbol{\theta})$ to denote the gradient and Hessian of $g(\mathbf{x}; \boldsymbol{\theta})$ with respect to (w.r.t.) $\boldsymbol{\theta}$, respectively. Let $\|\mathbf{v}\|$ be the Frobenius norm and $\mathbf{v}^{\otimes 2} = \mathbf{v}\mathbf{v}^{\mathrm{T}}$ for a vector or a matrix $\mathbf{v}$. We denote $\pi(\mathbf{x}, y)$ as the sampling probability for an instance $(\mathbf{x}, y)$. For negative cases, we sometimes use a shorthand notation $\pi(\mathbf{x})$ to denote its sampling probability. We use $o_P(1)$ to represent a random quantity that converges to 0 in probability as $N \to \infty$.

2.1 It is the number of positive instances that matters

We show that for rare events data, the estimation error rate for $\boldsymbol{\theta}$ is related to N_1 instead of N. Using the full data, the most efficient estimator of $\boldsymbol{\theta}$ is the maximum likelihood estimator (MLE)

$$\hat{\boldsymbol{\theta}}_{\mathrm{f}} := \arg\max_{\boldsymbol{\theta}} \sum_{i=1}^{N} \left[y_i g(\mathbf{x}_i; \boldsymbol{\theta}) - \log\left\{ 1 + e^{g(\mathbf{x}_i; \boldsymbol{\theta})} \right\} \right].$$

To investigate the theoretical properties of the estimator $\hat{\boldsymbol{\theta}}_{\mathrm{f}}$, we make the following assumptions.

Assumption 1. The first, second, and third derivatives of $f(\mathbf{x}; \boldsymbol{\beta})$ and $e^{f(\mathbf{x}; \boldsymbol{\beta})} f(\mathbf{x}; \boldsymbol{\beta})$ w.r.t. any components of $\boldsymbol{\beta}$ are bounded by a square intergrable random variable $B(\mathbf{x})$.

Assumption 2. The matrix $\mathbb{E}\{\dot{g}^{\otimes 2}(\mathbf{x}; \boldsymbol{\theta}^*)\}$ is finite and positive definite.

Assumption 1 imposes constraints on the smoothness of $f(\mathbf{x}; \boldsymbol{\theta})$ w.r.t. $\boldsymbol{\theta}$ and on the tightness of the distribution of $\mathbf{x}$. Assumption 2 ensures that all components of $\boldsymbol{\theta}$ are estimable.

Theorem 1. *Let* $\mathbf{V}_{\mathrm{f}} = \mathbb{E}\{e^{f(\mathbf{x}; \boldsymbol{\beta}^*)}\}\mathbf{M}_{\mathrm{f}}^{-1}$ *where* $\mathbf{M}_{\mathrm{f}} = \mathbb{E}\{e^{f(\mathbf{x}; \boldsymbol{\beta}^*)}\dot{g}^{\otimes 2}(\mathbf{x}; \boldsymbol{\theta}^*)\}$. *Under Assumptions 1 and 2, as* $N \to \infty$,

$$\sqrt{N_1}(\hat{\boldsymbol{\theta}}_{\mathrm{f}} - \boldsymbol{\theta}^*) \longrightarrow \mathbb{N}(\mathbf{0}, \ \mathbf{V}_{\mathrm{f}}), \quad \text{in distribution.}$$

This result shows that even if all the N instances are used for model training, the resulting estimator $\hat{\boldsymbol{\theta}}_{\mathrm{f}}$ converges to the true parameter $\boldsymbol{\theta}^*$ at the rate of $N_1^{-1/2}$, which is much slower than the usual rate of $N^{-1/2}$ for regular cases. This indicates that with rare events data, the available information about unknown parameters is at the scale of N_1, and it is the number of positive instances that matters.

2.2 Negative sampling

Since the available information ties to the number of positive instances instead of the full data size, one can keep all the positive instances and significantly subsmaple the negative instances to reduce the computational cost. For better estimation efficiency, we consider nonuniform sampling. Let $\varphi(\mathbf{x}) > 0$ be an integrable function and ρ be the sampling rate on the negative class. Without loss of generality, assume $\mathbb{E}\{\varphi(\mathbf{x})\} = 1$ so that $\pi(\mathbf{x}_i) := \rho\varphi(\mathbf{x}_i)$ is the sampling probability for the i-th data point if $y_i = 1$. We present the nonuniform negative sampling procedure in Algorithm 1.

Algorithm 1 Negative sampling

For $i = 1, ..., N$:   if $y_i = 1$,   include $\{\mathbf{x}_i, y_i, \pi(\mathbf{x}_i, y_i) = 1\}$ in the sample;
                 if $y_i = 0$,   calculate $\varphi(\mathbf{x}_i)$ and generate $u_i \sim \mathbb{U}(0, 1)$;
                                if $u_i \leq \rho\varphi(\mathbf{x}_i)$, include $\{\mathbf{x}_i, y_i, \pi(\mathbf{x}_i, y_i) = \rho\varphi(\mathbf{x}_i)\}$ in the sample.

---

3 Weighted estimation and its optimal negative sampling probability

Let $\delta_i = 1$ if the i-th data point is selected and $\delta_i = 0$ otherwise. Given a subsample taken according to $\pi(\mathbf{x}_i, y_i) = y_i + (1 - y_i)\rho\varphi(\mathbf{x}_i)$, $i = 1, ..., N$, the IPW estimator of $\boldsymbol{\theta}$ is

$$\hat{\boldsymbol{\theta}}_w = \arg\max_{\boldsymbol{\theta}} \sum_{i=1}^{N} \delta_i \pi^{-1}(\mathbf{x}_i, y_i) \left[y_i g(\mathbf{x}_i; \boldsymbol{\theta}) - \log\left\{ 1 + e^{g(\mathbf{x}_i; \boldsymbol{\theta})} \right\} \right]. \tag{2}$$

We need an assumption on the subsampling rate to investigate the subsample estimators.

Assumption 3. The subsampling rate ρ satisfies that $c_N := e^{\alpha^*}/\rho \to c$ for a constant $c \in [0, \infty)$.

It can be shown that $c_N \mathbb{E}\{e^{f(\mathbf{x}; \boldsymbol{\beta}^*)}\} = N_1(N_0\rho)^{-1}\{1 + o_P(1)\}$, so $c\mathbb{E}\{e^{f(\mathbf{x}; \boldsymbol{\beta}^*)}\}$ is the asymptotic positive/negative ratio for the sample. The theorem below shows the asymptotic distribution of $\hat{\boldsymbol{\theta}}_w$.

Theorem 2. *Under Assumptions 1-3, if* $\mathbb{E}[\{\varphi(\mathbf{x}) + \varphi^{-1}(\mathbf{x})\}B^2(\mathbf{x})] < \infty$ *then as* $N \to \infty$,

$$\sqrt{N_1}(\hat{\boldsymbol{\theta}}_w - \boldsymbol{\theta}^*) \longrightarrow \mathbb{N}(\mathbf{0}, \ \mathbf{V}_w), \quad \text{in distribution,}$$

where $\mathbf{V}_w = \mathbf{V}_{\mathrm{f}} + \mathbf{V}_{\mathrm{sub}}$ *and* $\mathbf{V}_{\mathrm{sub}} = c\mathbb{E}\{e^{f(\mathbf{x}; \boldsymbol{\beta}^*)}\}\mathbf{M}_{\mathrm{f}}^{-1}\mathbb{E}\{\varphi^{-1}(\mathbf{x})e^{2f(\mathbf{x}; \boldsymbol{\beta}^*)}\dot{g}^{\otimes 2}(\mathbf{x}; \boldsymbol{\theta}^*)\}\mathbf{M}_{\mathrm{f}}^{-1}$.

The subsampled estimator $\hat{\boldsymbol{\theta}}_w$ have the same convergence rate as the full data estimator. However, the asymptotic variance $\mathbf{V}_w$ may be inflated by the term $\mathbf{V}_{\mathrm{sub}}$ from subsampling the negative cases. Here $\mathbf{V}_{\mathrm{sub}}$ is zero if $c = 0$. If we keep much more negative instances than the positive instances in the subsample ($c = 0$), then there is no asymptotic information loss due to subsampling ($\mathbf{V}_w = \mathbf{V}_{\mathrm{f}}$). In this scenario, the number of negative instances can still be aggressively reduced ($\rho \to 0$) so that the computational efficiency is significantly improved, and the sampling function $\varphi(\mathbf{x})$ does not play an significant role. If we have to reduce the negative instances to the same level of the positive instances ($0 < c < \infty$), then the variance inflation $\mathbf{V}_{\mathrm{sub}}$ is not negligible. In this scenario, a well designed sampling function $\varphi(\mathbf{x})$ it is more relevant and critical.

Theorem 2 shows that the distribution of the estimation error $err := \sqrt{N_1}(\hat{\boldsymbol{\theta}}_w - \boldsymbol{\theta}^*)$ is approximated by a normal distribution $\mathbb{N}(\mathbf{0}, \mathbf{V}_w)$. Thus for any $\epsilon > 0$ the probability of excess error $\Pr(err > \epsilon)$ is approximated by $\Pr\{\|\mathbb{N}(\mathbf{0}, \mathbf{V}_w)\| > \epsilon\} = \Pr\left(\sum_{j=1}^d \lambda_j z_j^2 > \epsilon\right)$, where λ_j's are eigenvalues of $\mathbf{V}_w$ and z_j's are independent standard normal random variables. Therefore, a smaller trace of $\mathbf{V}_w$ means a smaller probability of excess error of the same level since $\mathrm{tr}(\mathbf{V}_w) = \sum_{j=1}^d \lambda_j$.

The following theorem gives the optimal negative sampling function for the IPW estimator.

Theorem 3. *For a given sampling rate ρ, the asymptotically optimal $\varphi(\mathbf{x})$ that minimizes $\mathrm{tr}(\mathbf{V}_w)$ is*

$$\varphi_{\mathrm{os}}(\mathbf{x}) = \frac{\min\{t(\mathbf{x}; \boldsymbol{\theta}^*), T\}}{\mathbb{E}[\min\{t(\mathbf{x}; \boldsymbol{\theta}^*), T\}]}, \tag{3}$$

where $t(\mathbf{x}; \boldsymbol{\theta}) = p(\mathbf{x}; \boldsymbol{\theta})\|\mathbf{M}_{\mathrm{f}}^{-1}\dot{g}(\mathbf{x}; \boldsymbol{\theta})\|$ and T is the maximum number so that $\rho[\min\{t(\mathbf{x}; \boldsymbol{\theta}^*), T\}] \leq \mathbb{E}[\min\{t(\mathbf{x}; \boldsymbol{\theta}^*), T\}]$ with probability one.

Remark 1. If $\rho \to 0$ then $\rho t(\mathbf{x}; \boldsymbol{\theta}^*) \leq \mathbb{E}\{t(\mathbf{x}; \boldsymbol{\theta}^*)\}$ almost surely, so T can be dropped (i.e., $T = \infty$). If $\lim_{N \to \infty} \rho > 0$ then $c = 0$, so the variance inflation due to subsampling is negligible. Thus, the truncation term T can be ignored in practice with imbalanced data; it only plays an role for not very imbalanced data when the sampling ratio is not very small.

In $\varphi_{\mathrm{os}}(\mathbf{x})$, $t(\mathbf{x}; \boldsymbol{\theta}^*)$ consists of two components, $p(\mathbf{x}; \boldsymbol{\theta}^*)$ and $\|\mathbf{M}_{\mathrm{f}}^{-1}\dot{g}(\mathbf{x}; \boldsymbol{\theta}^*)\|$. The term $p(\mathbf{x}; \boldsymbol{\theta}^*)$ is from the binary response model structure, and it gives a higher preference for a data point with larger $p(\mathbf{x}; \boldsymbol{\theta}^*)$ in the negative class. The term $\|\mathbf{M}_{\mathrm{f}}^{-1}\dot{g}(\mathbf{x}; \boldsymbol{\theta}^*)\|$ corresponds to the A optimality criterion [23]. For computational benefit in optimal sampling, the A optimality is often replaced by the L optimality [e.g., 30, 26]. Under the L optimality, $\|\mathbf{M}_{\mathrm{f}}^{-1}\dot{g}(\mathbf{x}; \boldsymbol{\theta}^*)\|$ is replaced by $\|\dot{g}(\mathbf{x}; \boldsymbol{\theta}^*)\|$, and this minimizes the asymptotic mean squared error (MSE) of $\mathbf{M}_{\mathrm{f}}\hat{\boldsymbol{\theta}}_w$. In case the gradient is difficult to obtain, one could ignore the gradient term and simply use $p(\mathbf{x}; \boldsymbol{\theta}^*)$ to replace $t(\mathbf{x}; \boldsymbol{\theta}^*)$. Although this does not give an optimal sampling probability, it outperforms the uniform subsampling because it takes into account the information of the model structure like the local case-control (LCC) sampling.

4 More efficient estimation based on the likelihood with log odds correction

4.1 Nonuniform log odds correction

The optimal function $\varphi_{\mathrm{os}}(\mathbf{x})$ assigns larger probabilities to more informative instances. However, the IPW estimator in (2) assigns smaller weights to more informative instances in estimation, so the resulting estimator can be improved to have higher estimation efficiency. A naive unweighted estimator is biased, and the bias correction approach in [8, 27] does not work for the sampling procedure in Algorithm 1 even when the underlying model is the logistic regression. We adopted the idea of [11] and seek to define a more efficient estimator through finding the corrected likelihood of the sampled data based on any negative sampling probability $\pi(\mathbf{x})$. For data included in the subsample (where $\delta = 1$), the conditional probability of $y = 1$ is

$$p_\pi(\mathbf{x}; \boldsymbol{\theta}) := \Pr(y = 1 \mid \mathbf{x}, \delta = 1) = \frac{1}{1 + e^{-g(\mathbf{x}; \boldsymbol{\theta}) - l}}, \tag{4}$$

where $l = -\log\{\pi(\mathbf{x})\}$. Please see the detailed derivation of (4) in the supplement. To avoid the IPW, we propose the estimator based on the log odds corrected likelihood in (4), namely,

$$\hat{\boldsymbol{\theta}}_{\mathrm{lik}} = \arg\max_{\boldsymbol{\theta}} \ell_{\mathrm{lik}}(\boldsymbol{\theta}), \quad \text{where} \quad \ell_{\mathrm{lik}}(\boldsymbol{\theta}) = \sum_{i=1}^N \delta_i \left[y_i g(\mathbf{x}_i; \boldsymbol{\theta}) - \log\left\{ 1 + e^{g(\mathbf{x}_i; \boldsymbol{\theta}) + l_i} \right\} \right]. \tag{5}$$

158 With $\hat{\boldsymbol{\theta}}_w$ in (2), $\pi(\mathbf{x}_i)$ is in the inverse. If an instance with much smaller $\pi(\mathbf{x}_i)$ is selected in the
159 sample, it dominates the objective function, making the resulting estimator unstable. With $\hat{\boldsymbol{\theta}}_{\text{lik}}$, this
160 problem is ameliorated because $\pi(\mathbf{x}_i)$ is in the logarithm in the log-likelihood of (5) and $\log(v)$ goes
161 to infinity much slower than v^{-1} as $v \downarrow 0$.

4.2 Theoretical analysis of the likelihood-based estimator

163 The following shows the asymptotic distribution of $\hat{\boldsymbol{\theta}}_{\text{lik}}$.

164 **Theorem 4.** *Under Assumptions 1-3, if* $\mathbb{E}\{e^{f(\mathbf{x};\boldsymbol{\beta}^*)}\varphi^{-1}(\mathbf{x})B(\mathbf{x})\} < \infty$, *then as* $N \to \infty$,

$$\sqrt{N_1}(\hat{\boldsymbol{\theta}}_{\text{lik}} - \boldsymbol{\theta}) \longrightarrow \mathbb{N}(\mathbf{0}, \mathbf{V}_{\text{lik}}) \quad \text{in distribution},$$

165 *where* $\mathbf{V}_{\text{lik}} = \mathbb{E}\{e^{f(\mathbf{x};\boldsymbol{\beta}^*)}\}\boldsymbol{\Lambda}_{\text{lik}}^{-1}$ *and* $\boldsymbol{\Lambda}_{\text{lik}} = \mathbb{E}\left[\frac{e^{f(\mathbf{x};\boldsymbol{\beta}^*)}\dot{g}^{\otimes 2}(\mathbf{x};\boldsymbol{\theta}^*)}{1+c\varphi^{-1}(\mathbf{x})e^{f(\mathbf{x};\boldsymbol{\beta}^*)}}\right]$.

166 Theorem 3 shows that the proposed estimator $\hat{\boldsymbol{\theta}}_{\text{lik}}$ is asymptotic normal with variance $\mathbf{V}_{\text{lik}}$. We
167 compare $\mathbf{V}_{\text{lik}}$ with $\mathbf{V}_w$ to show that $\hat{\boldsymbol{\theta}}_{\text{lik}}$ has a higher estimation efficiency than $\hat{\boldsymbol{\theta}}_w$.

168 **Theorem 5.** *If* $\mathbf{V}_w$ *and* $\mathbf{V}_{\text{lik}}$ *are finite and positive definite, then* $\mathbf{V}_{\text{lik}} \leq \mathbf{V}_w$, *i.e.,* $\mathbf{V}_w - \mathbf{V}_{\text{lik}}$ *is*
169 *non-negative definite. The equality holds when* $c = 0$ *and in this case* $\mathbf{V}_{\text{lik}} = \mathbf{V}_w = \mathbf{V}_{\text{f}}$.

170 Since the estimator $\hat{\boldsymbol{\theta}}_{\text{lik}}$ in Eg. (5) is based on the conditional log-likelihood of the sampled data, it
171 actually has the highest estimation efficiency among a class of asymptotically unbiased estimators.

172 **Theorem 6.** *Let* $\mathbf{X} = (\mathbf{x}_1, ..., \mathbf{x}_N)^{\text{T}}$ *be the feature matrix and* $\mathcal{D}_\delta$ *denote the sampled data. Let* $\hat{\boldsymbol{\theta}}_U$
173 *be a subsample estimator with the following asymptotic representation:*

$$\hat{\boldsymbol{\theta}}_U = \mathbf{U}(\boldsymbol{\theta}^*; \mathcal{D}_\delta) + N_1^{-1/2}o_P(1), \tag{6}$$

174 *where the variance* $\mathbb{V}\{\mathbf{U}(\boldsymbol{\theta}^*; \mathcal{D}_\delta)\}$ *exists, and* $\mathbf{U}(\boldsymbol{\theta}^*; \mathcal{D}_\delta)$ *satisfies that* $\mathbb{E}\{\mathbf{U}(\boldsymbol{\theta}^*; \mathcal{D}_\delta) \mid \mathbf{X}\} = \boldsymbol{\theta}^*$
175 *and* $\mathbb{E}\{\dot{\mathbf{U}}(\boldsymbol{\theta}^*; \mathcal{D}_\delta) \mid \mathbf{X}\} = \mathbf{0}$ *with* $\dot{\mathbf{U}}(\boldsymbol{\theta}^*; \mathcal{D}_\delta) = \partial\mathbf{U}(\boldsymbol{\theta}^*; \mathcal{D}_\delta)/\partial\boldsymbol{\theta}^{*\text{T}}$. *If* $N_1\mathbb{V}\{\mathbf{U}(\boldsymbol{\theta}^*; \mathcal{D}_\delta)\} \to \mathbf{V}_U$
176 *in probability, then* $\mathbf{V}_U \geq \mathbf{V}_{\text{lik}}$.

177 **Remark 2.** Theorem 6 tells us that $\hat{\boldsymbol{\theta}}_{\text{lik}}$ is statistically the most efficient among a class of estimators,
178 which includes both $\hat{\boldsymbol{\theta}}_w$ and $\hat{\boldsymbol{\theta}}_{\text{lik}}$ as special cases. The condition $\mathbb{E}\{\mathbf{U}(\boldsymbol{\theta}; \mathcal{D}_\delta) \mid \mathbf{X}\} = \mathbf{0}$ ensures that
179 the estimator is asymptotically unbiased. The condition $\mathbb{E}\{\dot{\mathbf{U}}(\boldsymbol{\theta}^*; \mathcal{D}_\delta) \mid \mathbf{X}\} = \mathbf{0}$ can be intuitively
180 interpreted as the requirement that the derivative of the $o_P(1)$ term in (6) is also negligible. Clearly,
181 this is satisfied by all unbiased estimators for which the $o_P(1)$ term in (6) is $\mathbf{0}$.

5 Practical considerations

183 In this section, we first show how we design a more practical estimator based on the previous results
184 and then present the theoretical analysis behind these improvements.

5.1 Making estimators more practical

186 Like the LCC sampling [8] and the A-optimal sampling [27], the $\varphi_{\text{os}}(\mathbf{x})$ in (3) depends on $\boldsymbol{\theta}$, and thus
187 a pilot value of $\boldsymbol{\theta}$, say $\tilde{\boldsymbol{\theta}}$, is required in practice. Here, $\tilde{\boldsymbol{\theta}}$ can be constructed from a pilot sample, say
188 $(\tilde{\mathbf{x}}_i, \tilde{y}_i)$, $i = 1, ..., \tilde{n}$, taken by uniform sampling from both classes with equal expected sample sizes.
189 With $\tilde{\boldsymbol{\theta}}$ obtained, calculate $\tilde{t}_i = p(\tilde{\mathbf{x}}_i; \tilde{\boldsymbol{\theta}})\|\tilde{\mathbf{M}}_{\text{f}}^{-1}\dot{g}(\tilde{\mathbf{x}}_i; \tilde{\boldsymbol{\theta}})\|$ for $i = 1, ..., \tilde{n}$, where $\tilde{\mathbf{M}}_{\text{f}}$ is the Hessian
190 matrix of the pilot sample objective function. In case that the Hessian matrix and gradients are hard
191 to record, the gradient term can be dropped, i.e., use $\tilde{t}_i = p(\tilde{\mathbf{x}}; \tilde{\boldsymbol{\theta}})$. As mentioned in Remark 1, T can
192 be ignored for very imbalanced data. The expectation in the denominator of (3) can be approximated
193 by mimicking the method of moment estimator, namely by

$$\tilde{\omega} = 2N_1(\tilde{n}N)^{-1}\sum_{\tilde{y}_i=1}\tilde{t}_i + 2N_0(\tilde{n}N)^{-1}\sum_{\tilde{y}_i=0}\tilde{t}_i. \tag{7}$$

194 In practice, it is common to set a lower threshold, say ϱ, on sampling probabilities, to ensure that all
195 instances have positive probabilities to be selected. This is more critical for the IPW estimator while
196 the likelihood estimator is not very sensitive to very small sampling probabilities.

197 Denote the pilot value as $\tilde{\boldsymbol{\vartheta}} = (\tilde{\boldsymbol{\theta}}, \tilde{\omega})$. The following probabilities can be practically implemented

$$\pi_\varrho^{\mathrm{os}}(\mathbf{x}; \tilde{\boldsymbol{\vartheta}}) = \min[\max\{\rho \tilde{\varphi}_{\mathrm{os}}(\mathbf{x}), \varrho\}, 1], \quad \text{where} \quad \tilde{\varphi}_{\mathrm{os}}(\mathbf{x}) = \tilde{\omega}^{-1} t(\mathbf{x}; \tilde{\boldsymbol{\theta}}), \tag{8}$$

198 and we use this in our experiments in Section 6. The $\tilde{\omega}$ in (7) is essentially a normalizing term so
199 that the expected subsample size is ρ. For some practical systems such as online models, $\tilde{\omega}$ is often
200 treated as a tuning parameter. We will illustrated this in our real data example in Section 6.2.

5.2 Theoretical analysis of practical estimators

202 Denote $\hat{\boldsymbol{\theta}}_w^{\tilde{\boldsymbol{\vartheta}}}$ and $\hat{\boldsymbol{\theta}}_{\mathrm{lik}}^{\tilde{\boldsymbol{\vartheta}}}$ the IPW estimator and the likelihood estimator, respectively, with practically
203 estimated optimal probability in (8). We have the following results.

204 **Theorem 7.** *Assume that the pilot estimator $\tilde{\boldsymbol{\vartheta}}$ is independent of the data, $\tilde{\varphi}_{\mathrm{os}}(\mathbf{x}) \to \varphi_{\mathrm{plt}}(\mathbf{x})$ in*
205 *probability, and $\rho^{-1}\varrho \to c_l > 0$. Under Assumptions 1-3, as $N \to \infty$, the $\hat{\boldsymbol{\theta}}_w^{\tilde{\boldsymbol{\vartheta}}}$ satisfies that,*

$$\sqrt{N_1}(\hat{\boldsymbol{\theta}}_w^{\tilde{\boldsymbol{\vartheta}}} - \boldsymbol{\theta}) \to \mathbb{N}(\mathbf{0}, \mathbf{V}_w^{\mathrm{plt}}) \quad \text{in distribution,}$$

206 *where $\mathbf{V}_w^{\mathrm{plt}}$ has the same expression of $\mathbf{V}_w$ except that $\varphi(\mathbf{x})$ is replaced by $\max\{\varphi_{\mathrm{plt}}(\mathbf{x}), c_l\}$. If*
207 $\Pr\{\varphi_{\mathrm{plt}}(\mathbf{x}) \geq c_l\} = 1$ *and the pilot estimator is consistent, then $\mathbf{V}_w^{\mathrm{plt}}$ achieves the optimal variance.*

208 **Theorem 8.** *Assume that the pilot estimator $\tilde{\boldsymbol{\vartheta}}$ is independent of the data, $\tilde{\varphi}_{\mathrm{os}}(\mathbf{x}) \to \varphi_{\mathrm{plt}}(\mathbf{x})$ in*
209 *probability, and $\varrho = o(\rho)$. Under Assumptions 1-3, as $N \to \infty$, $\hat{\boldsymbol{\theta}}_{\mathrm{lik}}^{\tilde{\boldsymbol{\vartheta}}}$ satisfies that,*

$$\sqrt{N_1}(\hat{\boldsymbol{\theta}}_{\mathrm{lik}}^{\tilde{\boldsymbol{\vartheta}}} - \boldsymbol{\theta}) \to \mathbb{N}(\mathbf{0}, \mathbf{V}_{\mathrm{lik}}^{\mathrm{plt}}) \quad \text{in distribution,}$$

210 *where $\mathbf{V}_{\mathrm{lik}}^{\mathrm{plt}}$ has the same expression of $\mathbf{V}_{\mathrm{lik}}$ except that $\varphi(\mathbf{x})$ is replaced by $\varphi_{\mathrm{plt}}(\mathbf{x})$. If the pilot*
211 *estimator is consistent, then $\mathbf{V}_{\mathrm{lik}}^{\mathrm{plt}}$ achieves the variance with the optimal probability.*

212 **Remark 3.** Theorems 7 and 8 do not require $\tilde{\boldsymbol{\vartheta}}$ to be consistent, i.e., the pilot estimator can be
213 misspecified. However $\tilde{\boldsymbol{\vartheta}}$ has to be consistent in order to achieve the asymptotic variances with the
214 optimal probability. Compared with the likelihood estimator $\hat{\boldsymbol{\theta}}_{\mathrm{lik}}^{\tilde{\boldsymbol{\vartheta}}}$, the IPW estimator $\hat{\boldsymbol{\theta}}_w^{\tilde{\boldsymbol{\vartheta}}}$ requires a
215 stronger condition on ϱ to have asymptotic normality, because $\pi_\varrho^{\mathrm{os}}(\mathbf{x}; \tilde{\boldsymbol{\vartheta}})$ is in the denominator of the
216 objective function for $\hat{\boldsymbol{\theta}}_w^{\tilde{\boldsymbol{\vartheta}}}$ so the turbulence of $\tilde{\boldsymbol{\vartheta}}$ is amplified. The requirement $c_l > 0$ means that ϱ
217 cannot be too small in practice. The likelihood estimator does not have this constraint.

6 Numerical experiments

219 We present simulated results and report the performance on a CTR dataset with more than 0.3 trillion
220 instances. More experiments with model and pilot misspecifications are available in the supplement.

6.1 Experiments on simulated data

222 We implemented the following negative sampling methods for logistic regression. 1) uniW: uniform
223 sampling with the IPW estimator. 2) uniLik: uniform sampling with the likelihood estimator. 3)
224 optW: optimal sampling with the IPW estimator. 4) optLik: optimal sampling with the likelihood
225 estimator. We also implement the full data MLE (Full) and the LCC for comparisons.

226 In each repetition of the simulation, we generate full data of size $N = 5 \times 10^5$ from the logistic
227 regression with $g(\mathbf{x}; \boldsymbol{\theta}) = \alpha + \mathbf{x}^{\mathrm{T}}\boldsymbol{\beta}$. We set the true $\boldsymbol{\beta}^*$ to be a 6×1 vector of ones and set different
228 values for α for different feature distributions so that the positive/negative ratio is close to 1:400. We
229 consider the following feature distributions: (a) Normal distribution which is symmetric with light
230 tails; the true intercept is $\alpha = -7.65$. (b) Log-normal distribution which is asymmetric and positively
231 skewed; the true intercept is $\alpha = -0.5$. (c) t_3 distribution which is symmetric with heavier tails; the
232 true intercept is $\alpha = -7$.

233 We set the sampling rate as $\rho = 0.002, 0.004, 0.006, 0.01$, and 0.02 for all sampling methods. In each
234 repetition, uniform samples of average size 100 are selected from each class to calculate the pilot
235 estimates, so the uncertainty due to the pilot estimates are taken into account. We also consider pilot
236 misspecification by adding a uniform random number from $\mathbb{U}(0, 1.5)$ to the pilot so that the pilot is

(a). **x**'s are normal   (b) **x**'s are lognormal   (c) **x**'s are t_3

Figure 1: Log(MSEs) for different estimators (the smaller the better). The top row uses consistent pilot estimators; the bottom uses misspecified pilot estimators.

systematically different from the true parameter. We repeat the simulation for $R = 1000$ times to calculate the MSE as $R^{-1} \sum_{r=1}^{R} \|\hat{\boldsymbol{\theta}}^{(r)} - \boldsymbol{\theta}\|^2$, where $\boldsymbol{\theta}^{(r)}$ is the estimate at the r-th repetition.

Results are presented in Figure 1. The MSEs of all estimators decrease as the sampling rate ρ increases. The optLik outperforms other sampling methods in general, and its performance is close to the full data MLE when $\rho = 0.02$, especially if the pilot is consistent. The advantage of optLik over optW is more evident for smaller ρ. If the pilot is misspecified, optLik is quite robust but optLik and LCC significantly deteriorate. Since uniLik and uniW do not require a pilot, they are not affected by pilot misspecification, but they have much larger MSE than optLik and optW when the pilot is consistent, especially uniW. Different feature distributions also affect the performances of different sampling methods.

6.2 Experiments on real data

We conduct experiments on a real online streaming CTR dataset[1] with more than 0.3 trillion instances. There are 299 input features, including user and ad profiles together with rich context features. We concatenate all features as a single high-dimensional vector **x**, whose length is greater than $5,000$. We use a 3-layer fully connected deep neural network as a feature extractor that maps **x** into a 256-dimensional dense features $h(\mathbf{x})$. The downstream model is a linear logistic regression model taking $h(\mathbf{x})$ as input to predict the binary output whether the user clicks the ad or not.

The positive/negative ratio is around 1:80. Due to limited storage, we first uniformly drop 80% negative instances and try various negative sampling algorithms on the rest 60 billion instances. On this pre-processed dataset, we then split out 1 percent of the data for testing, and do negative sampling on the rest data. The entire training procedure scans the subsampled data set from begin to end in one-pass according to the timestamp. The testing is done all the way through training. We calculate the area under the ROC curve (AUC) for testing instances. The entire model is trained on a distributed machine learning platform using 1,200 CPU cores and can finish within two days.

We adopt the sampling probability $\pi_\varrho^{os}(\mathbf{x}; \tilde{\boldsymbol{\vartheta}})$ as demonstrated in (8) and use the empirical sampling rate to calibrate the normalizing term $\tilde{\omega}$. As mentioned before, we remove the gradient term for implementation tractability so $t(\mathbf{x}; \tilde{\boldsymbol{\theta}}) = p(\mathbf{x}; \tilde{\boldsymbol{\theta}})$. As the result will show, this approximate score produces consistent results with the simulation experiments. We compare four methods: 1) uniW and 2) uniLik are the same as in simulated experiments; and 3) optW and 4) optLik represent nonuniform sampling probability using $\pi_\varrho^{os}(\mathbf{x}; \tilde{\boldsymbol{\vartheta}})$ with the IPW estimator and with the likelihood estimator. We use "opt" to refer to both optW and optLik when focusing on the sampling.

---

[1]Since more detailed descriptions of the data source will expose authors' identity, we will uncover this information after the double-blind review process.

We study two sets of negative sample rates (w.r.t. the original 0.3 trillion data): (i) $[0.01, 0.05]$. In this case, negative instances still outnumber positive instances. (ii) $[0.001, 0.008]$. In this case, negative subsampling is too aggressive that positive instances dominates negative instances. We demonstrate the result in Figure 2. When the sample rate is moderate (Figure 2.(b)), opt is nearly optimal and can significantly outperform uniform subsampling. There is still a small gap between the IPW estimator and the likelihood estimator. When the sample rate is extremely small (Figure 2.(a)), opt becomes sub-optimal and its gap w.r.t. uniform sampling is closer. Still we see a clear difference between the IPW estimator and the likelihood estimator. This is due to a larger asymptotic variance for uniform sampling when the sampling rate goes to zero. Note that on this huge data set, a small relative AUC gain (e.g., 0.0005) usually could bring about significant revenue gain in products.

Figure 2: Empirical testing AUC of subsample estimators for different sample sizes (the larger the better).

(a). Extremely small sampling rates. (b) Moderate sampling rates.

6.3 Sensitivity analysis

In practice, we find it necessary to use cross-validation to find the optimal ϱ in (8) given a fixed sample rate. As demonstrated in Figure 3, for each sampling rate we tune ϱ from a very small value (in this case, 10^{-6}) all the way to its largest value. When ϱ achieves its largest value, $\rho\tilde{\omega}^{-1} = 0$ and $\pi_{\varrho}^{\mathrm{os}}(\mathbf{x}; \tilde{\boldsymbol{\vartheta}})$ reduces to uniform negative sampling. We observe that $\pi_{\varrho}^{\mathrm{os}}(\mathbf{x}; \tilde{\boldsymbol{\vartheta}})$ with a moderate ϱ achieves the best results for various sampling rates. Note that ϱ is the lower truncation level for negative sampling probabilities. Thus, a larger ϱ means more "simple" negative instances ($\rho\tilde{\omega}^{-1}p(\mathbf{x}_i; \tilde{\boldsymbol{\theta}})$ is smaller than ϱ) will have better chances to be selected. This coincides with recent empirical results for negative subsampling in large search systems [14], where they mix "simple" (smaller $p(\mathbf{x}_i; \tilde{\boldsymbol{\theta}})$) and "hard" (larger $p(\mathbf{x}_i; \tilde{\boldsymbol{\theta}})$) negative instances. We also observe an empirical variance of around 0.0001 for each setup, demonstrating that the relative improvement is consistent.

Figure 3: AUC for IPW and likelihood estimators when tuning the truncation lower bound ϱ.

7 Conclusions and limitations

In this paper, we have derived asymptotic distributions for full data and subsample estimators with rare events data. We have also found the optimal sampling probability that minimizes the unconditional asymptotic variance of the IPW estimator, and proposed the likelihood estimator

through nonuniform log odds correction to further improve the estimation efficiency. The optimal probability depends on unknown parameters so we have discussed practical implementations and investigated the convergences of the resultant estimators. Experiments on both simulated and real big online streaming data confirm our theoretical findings.

This paper has the following limitations: 1) We assume that the model is correctly specified. Theoretical properties with model misspecification is not considered, and this important question requires future studies. 2) The assumption of a fixed β^* means that both the marginal and conditional probabilities for a positive instance to occur are small. If the proportion of the positive instances for a given feature in a dataset is large, this assumption may not be appropriate. Further investigation is need to see if our results is still applicable. 3) Oversampling the rare positive instances is another common practice to deal with imbalanced data. Its application together with negative sampling is not considered.

References

[1] N. V. Chawla. Data mining for imbalanced datasets: An overview. In *Data Mining and Knowledge Discovery Handbook*, pages 875–886. Springer, 2009.

[2] N. V. Chawla, K. W. Bowyer, L. O. Hall, and W. P. Kegelmeyer. Smote: synthetic minority over-sampling technique. *Journal of Artificial Intelligence Research*, 16:321–357, 2002.

[3] N. V. Chawla, N. Japkowicz, and A. Kotcz. Editorial: special issue on learning from imbalanced data sets. *ACM SIGKDD Explorations Newsletter*, 6(1):1–6, 2004.

[4] J. Chen, B. Sun, H. Li, H. Lu, and X.-S. Hua. Deep ctr prediction in display advertising. In *Proceedings of the 24th ACM International Conference on Multimedia (MM)*, pages 811–820, 2016.

[5] G. Douzas and F. Bacao. Self-organizing map oversampling (somo) for imbalanced data set learning. *Expert Systems with Applications*, 82:40–52, 2017.

[6] C. Drummond, R. C. Holte, et al. C4. 5, class imbalance, and cost sensitivity: why under-sampling beats over-sampling. In *Workshop on Learning from Imbalanced Datasets II*, volume 11, pages 1–8. Citeseer, 2003.

[7] A. Estabrooks, T. Jo, and N. Japkowicz. A multiple resampling method for learning from imbalanced data sets. *Computational intelligence*, 20(1):18–36, 2004.

[8] W. Fithian and T. Hastie. Local case-control sampling: Efficient subsampling in imbalanced data sets. *Annals of statistics*, 42(5):1693, 2014.

[9] H. Guo, R. Tang, Y. Ye, Z. Li, and X. He. Deepfm: a factorization-machine based neural network for ctr prediction. *arXiv preprint arXiv:1703.04247*, 2017.

[10] H. Han, W.-Y. Wang, and B.-H. Mao. Borderline-smote: A new over-sampling method in imbalanced data sets learning. In D.-S. Huang, X.-P. Zhang, and G.-B. Huang, editors, *Advances in Intelligent Computing*, pages 878–887, Berlin, Heidelberg, 2005. Springer Berlin Heidelberg.

[11] L. Han, K. M. Tan, T. Yang, T. Zhang, et al. Local uncertainty sampling for large-scale multiclass logistic regression. *Annals of Statistics*, 48(3):1770–1788, 2020.

[12] X. He, J. Pan, O. Jin, T. Xu, B. Liu, T. Xu, Y. Shi, A. Atallah, R. Herbrich, S. Bowers, et al. Practical lessons from predicting clicks on ads at facebook. In *Proceedings of the Eighth International Workshop on Data Mining for Online Advertising*, pages 1–9, 2014.

[13] T. Hesterberg. Weighted average importance sampling and defensive mixture distributions. *Technometrics*, 37(2):185–194, 1995.

[14] J.-T. Huang, A. Sharma, S. Sun, L. Xia, D. Zhang, P. Pronin, J. Padmanabhan, G. Ottaviano, and L. Yang. Embedding-based retrieval in facebook search. In *Proceedings of the 26th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*, pages 2553–2561, 2020.

[15] N. Japkowicz. Learning from imbalanced data sets: Papers from the AAAI workshop, AAAI, 2000. Technical Report WS-00-05, 2000.

[16] G. King and L. Zeng. Logistic regression in rare events data. *Political analysis*, 9(2):137–163, 2001.

[17] G. Lemaître, F. Nogueira, and C. K. Aridas. Imbalanced-learn: A python toolbox to tackle the curse of imbalanced datasets in machine learning. *The Journal of Machine Learning Research*, 18(1):559–563, 2017.

[18] X. Liu, J. Wu, and Z. Zhou. Exploratory undersampling for class-imbalance learning. *IEEE Transactions on Systems, Man, and Cybernetics, Part B (Cybernetics)*, 39(2):539–550, 2009.

[19] J. Mathew, C. K. Pang, M. Luo, and W. H. Leong. Classification of imbalanced data by oversampling in kernel space of support vector machines. *IEEE Transactions on Neural Networks and Learning Systems*, 29(9):4065–4076, 2017.

[20] H. B. McMahan, G. Holt, D. Sculley, M. Young, D. Ebner, J. Grady, L. Nie, T. Phillips, E. Davydov, D. Golovin, et al. Ad click prediction: a view from the trenches. In *Proceedings of the 19th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pages 1222–1230, 2013.

[21] A. Owen and Y. Zhou. Safe and effective importance sampling. *Journal of the American Statistical Association*, 95(449):135–143, 2000.

[22] A. B. Owen. Infinitely imbalanced logistic regression. *The Journal of Machine Learning Research*, 8:761–773, 2007.

[23] F. Pukelsheim. *Optimal design of experiments*. SIAM, 2006.

[24] M. M. Rahman and D. Davis. Addressing the class imbalance problem in medical datasets. *International Journal of Machine Learning and Computing*, 3(2):224, 2013.

[25] Y. Sun, M. S. Kamel, A. K. Wong, and Y. Wang. Cost-sensitive boosting for classification of imbalanced data. *Pattern Recognition*, 40(12):3358–3378, 2007.

[26] D. Ting and E. Brochu. Optimal subsampling with influence functions. In *Advances in neural information processing systems*, pages 3650–3659, 2018.

[27] H. Wang. More efficient estimation for logistic regression with optimal subsamples. *Journal of Machine Learning Research*, 20(132):1–59, 2019.

[28] H. Wang. Logistic regression for massive data with rare events. In H. D. III and A. Singh, editors, *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 9829–9836. PMLR, 13–18 Jul 2020.

[29] H. Wang and Y. Ma. Optimal subsampling for quantile regression in big data. *Biometrika*, 108(1):99–112, 2021.

[30] H. Wang, R. Zhu, and P. Ma. Optimal subsampling for large sample logistic regression. *Journal of the American Statistical Association*, 113(522):829–844, 2018.

[31] J. Yu, H. Wang, M. Ai, and H. Zhang. Optimal distributed subsampling for maximum quasi-likelihood estimators with massive data. *Journal of the American Statistical Association*, 0(0):1–12, 2020.