On the Sample Complexity of Learning with Geometric Stability

Abstract

Many supervised learning problems involve high-dimensional data such as images, text, or graphs. In order to make efficient use of data, it is often useful to leverage certain geometric priors in the problem at hand, such as invariance to translations, permutation subgroups, or stability to small deformations. We study the sample complexity of learning problems where the target function presents such invariance and stability properties, by considering spherical harmonic decompositions of such functions on the sphere. We provide non-parametric rates of convergence for kernel methods, and show improvements in sample complexity by a factor equal to the size of the group when using an invariant kernel over the group, compared to the corresponding non-invariant kernel. These improvements are valid when the sample size is large enough, with an asymptotic behavior that depends on spectral properties of the group. Finally, these gains are extended beyond invariance groups to also cover geometric stability to small deformations, modeled here as subsets (not necessarily subgroups) of permutations.

1 Introduction

Learning from high-dimensional data is known to be statistically intractable without strong assumptions on the problem. A canonical example is learning Lipschitz functions, which generally requires a number of samples exponential in the dimension due to the curse of dimensionality (*e.g.*, [27]). Many high-dimensional machine learning problems involve highly structured data such as images, text, or graphs, and may exhibit invariance to certain transformations of the input data, such as permutations, translations or rotations, and near invariance to small deformations. More precisely, if $\mathcal{X}$ is the high-dimensional data domain, and G is a set of transformations $\sigma : \mathcal{X} \to \mathcal{X}$, the learning task can be alleviated if one knows in advance that the target function f varies smoothly under transformations in G: $|f(\sigma \cdot x) - f(x)|$ is uniformly small over $x \in \mathcal{X}$ for $\sigma \in G$.

To further motivate this property, it is useful to view the data domain $\mathcal{X}$ as a space of signals $\mathcal{X} = L^2(\Omega; \mathbb{R})$ defined over a geometric domain Ω, such as a 2d grid. The set of transformations G can then be articulated in terms of Ω rather than $\mathcal{X}$, a much simpler geometric object, and then *lifted* into $\mathcal{X}$ by composition: if $\sigma : \Omega \to \Omega$, and $x \in \mathcal{X}$ then $(\sigma \cdot x)(u) := x(\sigma^{-1}(u))$ for every $u \in \Omega$. The smoothness to transformations can thus be interpreted as a form of *geometric stability*.

In this paper, we quantify the sample complexity gains brought by geometric stability. Concretely, we consider target functions f defined on the sphere $\mathcal{X} = \mathbb{S}^{d-1}$ in d dimensions with finite $L^2(\mathbb{S}^{d-1})$ norm. In this case, we view the geometric domain as the discrete 1d grid $\Omega = [1, \ldots, d]$, and consider geometric transformations G as subsets of the symmetric group of permutations of d elements. Given a set G (not necessarily a group), we consider the *smoothing* operator given by $S_G f(x) = \frac{1}{|G|} \sum_{\sigma \in G} f(\sigma \cdot x)$ for $f \in L^2(\mathbb{S}^{d-1})$, and assume that our target function f is geometrically stable, in the sense that $f = S_G g$ for some $g \in L^2(\mathbb{S}^{d-1})$. In words, the smoothing

operator S_G replaces the prediction $f(x)$ by the average over transformations of x. In particular, functions which are invariant under the action of $\sigma \in G$, namely

$$f(\sigma \cdot x) = f(x), \quad \sigma \in G, x \in \mathbb{S}^{d-1}, \tag{1}$$

are also stable, with $f = S_G f$.

Building on the recent work [20], we proceed by studying harmonic decompositions of such functions using spherical harmonics [13], which generalize Fourier series on the circle to higher dimensions. This allows us to obtain rates of approximation for invariant and geometrically stable functions with varying levels of smoothness, and to study the generalization properties of invariant kernel methods using kernels defined on the sphere. Specifically, our main contributions are:

- By comparing spectral properties of usual kernels on the sphere with invariant ones, we find that the latter provide improvements in sample complexity by a factor of the order of the size of the group when the sample size is large enough (Section 3).

- We study how this improvement factor varies with sample size, in terms of the structure of the group and on spectral properties of the permutation matrices it contains (Section 4).

- We extend the invariance analysis to geometrically stable functions, establishing similar gains in sample complexity that depend on the size of the transformation subset (Section 5).

Our proofs rely on comparing the dimension of invariant and non-invariant spherical harmonics at a given degree, and showing that their ratio decays to the inverse group size as the degree tends to infinity. In contrast to [20], we consider the dimension to be fixed and study non-parametric rates of convergence for potentially non-smooth target functions and general groups of permutations, while they consider high-dimensional limits where only polynomials can be learned, and focus on invariance to translation groups.

Related work. Invariance and deformation stability have been analysed using convolutional neural network-type architectures such as the scattering transform [19, 6], or convolutional kernels [5, 17]. While these works characterise the stability in terms of the dyadic structure of convolutional filters (such as wavelets), they do not cover a statistical analysis of sample complexity. Similarly, models of compositional functions such as those in [10, 21, 24] study the benefit of hierarchical representations with local connectivity for approximation, while [18] studies computational benefits of local connectivity; yet these works do not consider invariance or stability. [20] studies similar benefits but focus on polynomial target functions, and consider high-dimensional limits, while we consider a fixed dimension and obtain non-parametric rates of convergence. While their framework is potentially applicable to general groups, their results only cover the case of translation groups, leading to improvements in sample complexity of at most a factor equal to the dimension. [14] also sudies benefits of group invariance, but focuses on linear models, and only considers interpolating estimators. [9] study benefits of equivariant kernels in structured prediction problems. [12] studies generalisation advantages of CNNs over fully-connected models, while our focus is on non-parametrics.

2 Preliminaries

In this section, we describe our setup and provide some background on harmonic decompositions on the sphere, and how these are affected by invariance.

Statistical learning setup. We consider a supervised learning problem where the data distribution ρ on input-label pairs (x, y) is such that $x \in \mathbb{S}^{d-1}$ and $\mathbb{E}[y|x] = f^*(x)$ for some target function f^* in $L^2(\mathbb{S}^{d-1})$. For simplicity, we will assume that x is uniformly distributed on the sphere, and denote the uniform measure on $\mathbb{S}^{d-1}$ by $d\tau$. We consider a regression setup with L^2 risk given by

$$R(f) = \mathbb{E}_{(x,y)\sim\rho}\left[(f(x) - y)^2\right].$$

For a given estimator $\hat{f}_n$ based on n samples from ρ, the goal is then to obtain generalization bounds as a function of n on the excess risk

$$\mathbb{E}[R(\hat{f}_n)] - R(f^*) = \mathbb{E}[\|\hat{f}_n - f^*\|^2_{L^2(d\tau)}], \tag{2}$$

where the expectation is over the n samples. Such bounds are well-studied for various classes of target functions f^* such as smoothness classes, and estimators such as kernel ridge regression. These are

typically studied through harmonic decompositions of f^* and of a kernel function in appropriate L^2 bases, which then relate function regularity and decays of Fourier coefficients.

Harmonic analysis on the sphere. When considering functions in $L^2(d\tau)$, an appropriate choice of orthonormal basis is that of spherical harmonic polynomials [1, 13]. More precisely, denote $\{Y_{k,j}\}_{j=1}^{N(d,k)}$ denote an orthonormal basis of the space $V_{d,k}$ of spherical harmonics of degree k, *i.e.*, homogeneous harmonic polynomials of degree k, where $N(d,k) = \frac{2k+d-2}{k}\binom{k+d-3}{d-2}$. Then, the collection $\{Y_{k,j} : k \geq 0, j = 1, \ldots, N(d,k)\}$ forms an orthonormal basis of $L^2(d\tau)$, so that any function $f \in L^2(d\tau)$ may be written

$$f(x) = \sum_{k \geq 0} \sum_{j=1}^{N(d,k)} a_{k,j} Y_{k,j}(x), \tag{3}$$

with $\sum_k \sum_{j=1}^{N(d,k)} a_{k,j}^2 < \infty$. Similarly, any dot-product kernel $K(x,x') = \kappa(\langle x, x' \rangle)$ on the sphere may be written

$$\kappa(\langle x, x' \rangle) = \sum_{k \geq 0} \mu_k \sum_{j=1}^{N(d,k)} Y_{k,j}(x) Y_{k,j}(x'), \tag{4}$$

where μ_k is given by

$$\mu_k = \frac{\omega_{d-2}}{\omega_{d-1}} \int_{-1}^{1} \kappa(t) P_{d,k}(t)(1-t^2)^{(d-3)/2} dt. \tag{5}$$

Here, ω_{p-1} is the surface measure of the sphere in p dimensions, and $P_{d,k}$ are Legendre or Gegenbauer polynomials of degree k in d dimensions (normalized with $P_{d,k}(1) = 1$),which form an orthogonal basis of $L^2([-1,1], dw)$, with $dw(t) = (1-t^2)^{(d-3)/2} dt$. When the kernel K is positive definite and is used in the context of kernel ridge regression with data uniformly distributed on the sphere, then the μ_k also correspond to the eigenvalues of the covariance operator. These eigenvalues and their decay then control the statistical properties of the kernel ridge regression estimator [7].

Spherical harmonics and group-invariant functions. In order to describe harmonic decompositions of functions satisfying the group invariance property (1) for a discrete group G, we follow [20] and define the symmetrization operator

$$S_G f(x) = \frac{1}{|G|} \sum_{\sigma \in G} f(\sigma \cdot x). \tag{6}$$

This operator acts as a projection from $L^2(d\tau)$ to a subset thereof which contains invariant functions. It can be shown that the spaces $V_{d,k}$ of spherical harmonics of degree k are stable by S [20], and we may then define an orthonormal basis of $\overline{V}_{d,k} := S_G V_{d,k}$ consisting of *invariant* spherical harmonics $\{\overline{Y}_{k,j}\}_{j=1}^{\overline{N}(d,k)}$. We then have the following lemma.

Lemma 1 (Representation of projection [20]). *For any $k \geq 0$, we have*

$$\gamma_d(k) := \frac{\overline{N}(d,k)}{N(d,k)} = \frac{1}{|G|} \sum_{\sigma \in G} \mathbb{E}_x \left[P_{d,k}(\langle \sigma \cdot x, x \rangle) \right]. \tag{7}$$

The quantity $\gamma_d(k)$ will play an important role in determining the gains in sample complexity brought by invariance. We will show in Section 4 that $\gamma_d(k)$ converges to $1/|G|$ for large k, with an asymptotic behavior that is governed by spectral properties of the elements of the group.

3   Sample Complexity of Invariant Kernels

We begin our study by focusing on the invariant case. In this section, we study the sample complexity of learning invariant functions, by considering kernel ridge regression estimators and providing non-parametric rates of convergence that illustrate the gains achievable with invariant kernels compared to non-invariant ones.

Kernel ridge regression (KRR) and invariant kernels. For a positive definite kernel K with RKHS $\mathcal{H}_K$, we consider the KRR estimator $\hat{f}_\lambda$ given by

$$\hat{f}_\lambda := \arg \min_{f \in \mathcal{H}_K} \frac{1}{n} \sum_{i=1}^n (f(x_i) - y_i)^2 + \lambda \|f\|_{\mathcal{H}_K}^2. \tag{8}$$

We consider the following kernels, which we assume positive definite, given for $x, x' \in \mathbb{S}^{d-1}$ by

$$K(x, x') = \kappa(\langle x, x' \rangle), \qquad K_G(x, x') = \frac{1}{|G|} \sum_{\sigma \in G} \kappa(\langle \sigma \cdot x, x' \rangle), \tag{9}$$

with $\kappa(u) \leq 1$. A common example for κ is the arc-cosine kernel [8], which arises from infinite-width shallow neural networks with ReLU activations. The following integral operator defined on $L^2(d\tau)$ and its eigen decomposition play an important role for the statistical and approximation properties of kernel methods:

$$T_K f(x) = \int K(x, x') f(x') d\tau(x'). \tag{10}$$

We now show that its spectral properties are closely related for K and K_G.

Lemma 2 (Spectral properties of K and K_G.). *There exists a basis of spherical harmonics in which the operators T_K and T_{K_G} are jointly diagonalized. They admit the same eigenvalues μ_k given by* (5)*, with multiplicity $N(d, k)$ for T_K and $\overline{N}(d, k)$ for T_{K_G}.*

The decay of the eigenvalues μ_k controls the smoothness of functions in the RKHS, for instance when μ_k decays polynomially, T_K behaves similarly to powers of the Laplacian on the sphere, leading to functional spaces similar to Sobolev spaces. For the example of the arc-cosine kernel, μ_k decays as k^{-d-2}. leading to an RKHS containing functions with $d/2 + 1$ bounded derivatives [2].

Approximation error. The approximation error of kernel methods is often controlled by the following quantity (*e.g.*, [3, 11]):

$$A_\mathcal{H}(\lambda, f^*) = \inf_{f \in \mathcal{H}} \|f - f^*\|_{L^2(d\tau)}^2 + \lambda \|f\|_\mathcal{H}^2, \tag{11}$$

where f^* is a target function in $L^2(d\tau)$, and $\mathcal{H}$ is a given RKHS. In particular, if f^* is smooth enough so that $f^* \in \mathcal{H}$, then we have $A_\mathcal{H}(\lambda, f^*) \leq \lambda \|f^*\|_\mathcal{H}^2$, while if $f^* \notin \mathcal{H}$, *e.g.*, if f^* is only Lipschitz, then $A_\mathcal{H}(\lambda, f^*)$ typically grows much faster with λ. We now show a useful result for invariant targets, showing that in this case the approximation error is the same for the kernels K and K_G.

Lemma 3 (Approximation error for invariant functions.). *If f^* is invariant to the group G, so that $f^* = Sf^*$, then we have*

$$A_{\mathcal{H}_K}(\lambda, f^*) = A_{\mathcal{H}_{K_G}}(\lambda, f^*). \tag{12}$$

Degrees of freedom. The above result suggests that any gains of using K_G instead of K for learning invariant functions should come from estimation rather than approximation error. The estimation error of ridge rigression estimators is typically controlled with the following quantity, often called *degrees of freedom* or *effective dimension* (*e.g.*, [3, 16]):

$$\mathcal{N}_K(\lambda) = \text{Tr}(\Sigma_K(\Sigma_K + \lambda I)^{-1}) = \sum_{m \geq 0} \frac{\lambda_m}{\lambda_m + \lambda}, \tag{13}$$

where $\Sigma_K = \mathbb{E}_x [K(x, \cdot) \otimes_{\mathcal{H}_K} K(x, \cdot)]$ is the covariance operator and $(\lambda_m)_{m \geq 0}$ its eigenvalues, taking multiplicity into account, which are the same as those of T_K when data is distributed according to $d\tau$ [7]. We then obtain the following simple result relating $\mathcal{N}_{K_G}$ to $\mathcal{N}_K$.

Lemma 4 (Degrees of freedom for K and K_G.). *For any $\ell \geq 0$, we have*

$$\mathcal{N}_{K_G}(\lambda) \leq D(\ell) + \nu_d(\ell) \mathcal{N}_K(\lambda),$$

where $D(\ell) := \sum_{k < \ell} \overline{N}(d, k)$ and $\nu_d(\ell) := \sup_{k \geq \ell} \gamma_d(k)$, with γ_d given in (7)*.*

This suggests that for a fixed ℓ, the effective dimension of K_G is controlled by a factor $\nu_d(\ell)$ times that of K, up to a finite fixed dimension $D(\ell)$. For difficult non-parametric problems which require small λ at large sample sizes, the second term will tend to dominate, so that having a small $\nu_d(\ell)$ may help reduce sample complexity compared to using the vanilla kernel K, an observation which we make rigorous below.

Generalization bound for KRR. Armed with the above lemmas on approximation error and degrees of freedom, we now study generalization of KRR under the following assumptions:

(A1) *capacity condition*: $\mathcal{N}_K(\lambda) \leq C_K \lambda^{-1/\alpha}$ with $\alpha > 1$.

(A2) *source condition*: there exists $r > \frac{\alpha-1}{2\alpha}$ and $g \in L^2(d\tau)$ with $\|g\|_{L^2(d\tau)} \leq C_{f^*}$ such that $f^* = T_K^r g$.

(A3) *invariance*: f^* is G-invariant.

(A4) *problem noise*: ρ is such that $\mathbb{E}_\rho[(y - f^*(x))^2 | x] \leq \sigma_\rho^2$.

The first, second, and fourth conditions are commonly used in the kernel methods literature [7]. (A1) characterizes the "size" of the RKHS, and is satisfied when the eigenvalues λ_m of T_K decay as $k^{-\alpha}$. On the sphere, $\alpha = \frac{2s}{d-1}$ corresponds to having s bounded derivatives, *e.g.*, we have $s = d/2 + 1$ for the arc-cosine kernel. The parameter r in (A2) defines the regularity of f^* relative to that of the kernel: $r = 1/2$ corresponds to $f^* \in \mathcal{H}_K$, while larger (resp. smaller) r implies f^* is more (resp. less) smooth. The condition on r is needed for our specific bound, which is based on [3, Proposition 7.2], but may be bypassed using different algorithms or analyses [15, 23]. We now present our bound on the excess risk.

Theorem 5 (Generalization of invariant kernel.). *Assume (A1-4). Let $\nu_d(\ell)$ be as in Lemma 4, or an upper bound thereof, and assume $\nu_0 := \inf_{\ell \geq 0} \nu_d(\ell) > 0$. Let*

$$n \geq \max\left\{ \|f^*\|_\infty^2 / \sigma_\rho^2, (C_1/\nu_0)^{\frac{\alpha}{2\alpha r + 1 - \alpha}} \right\}, \tag{14}$$

and define

$$\ell_n := \sup\{\ell : D(\ell) \leq C_2 \nu_d(\ell)^{\frac{2\alpha r}{2\alpha r + 1}} n^{\frac{1}{2\alpha r + 1}}\}. \tag{15}$$

We then have, for $\lambda = C_3(\nu_d(\ell_n)/n)^{\alpha/(2\alpha r + 1)}$,

$$\mathbb{E}[R(\hat{f}_\lambda) - R(f^*)] \leq C_4 \left(\frac{\nu_d(\ell_n)}{n} \right)^{\frac{2\alpha r}{2\alpha r + 1}}. \tag{16}$$

In the same setting, KRR with kernel K and $\lambda = C_3 n^{\frac{-\alpha}{2\alpha r + 1}}$ achieves $\mathbb{E}[R(\hat{f}_\lambda) - R(f^*)] \leq C_4 n^{\frac{-2\alpha r}{2\alpha r + 1}}$, where C_3, C_4 are the same constants as for the invariant kernel. Here, the constants $C_{1:4}$ only depend on the parameters of assumptions (A1-4).

The theorem shows that the generalization error for the invariant kernel behaves as if it effectively had access to $n/\nu_d(\ell_n)$ samples. Note that $\nu_d(\ell_n) \leq 1$ and ν_d is decreasing, so that we always have some improvement in sample complexity, and this gets better when ℓ_n is large. In particular, we show in Section 4 that $\nu_d(\ell)$ converges to $1/|G|$, so that asymptotically the gain in sample complexity can be as large as the size of the group, which in some cases may grow exponentially in d.

Estimating ℓ_n and $\nu_d(\ell_n)$. Since $D(\ell)$ increases with ℓ, Eq. (15) suggests that ℓ_n increases with n. We now provide intuition on how we might expect ℓ_n and $\nu_d(\ell_n)$ to behave in a situation of interest where we know an asymptotic equivalent of ν_d. Namely, assume that we have

$$\nu_d(\ell) \approx \nu_0 + c\ell^{-\beta}.$$

We provide such asymptotic equivalents in Section 4, where $\nu_0 = 1/|G|$, and β depends on spectral properties of the elements of G. For some large groups, β may be small, in which case we may consider other approximations with larger β, at the cost of a larger ν_0. When ℓ_n is large, using the approximation $N(d,k) \approx k^{d-2}$, we have $D(\ell) \approx \sum_{k=0}^{\ell-1} k^{d-2} \approx \ell^{d-1}$. Hiding constants other than ν_0, we may then consider ℓ_n to be solution of

$$\ell^{\frac{(d-1)(2\alpha r + 1)}{2\alpha r}} = n^{\frac{1}{2\alpha r}}(\nu_0 + \ell^{-\beta}).$$

Since the l.h.s. increases, while the r.h.s. decreases with ℓ, we must have $\ell_n \geq \max(\ell_{n,1}, \ell_{n,2})$, with

$$\ell_{n,1}^{\frac{(d-1)(2\alpha r + 1)}{2\alpha r}} = n^{\frac{1}{2\alpha r}} \nu_0, \quad \text{and} \quad \ell_{n,2}^{\frac{(d-1)(2\alpha r + 1)}{2\alpha r}} = n^{\frac{1}{2\alpha r}} \ell_{n,2}^{-\beta}.$$

This yields

$$\nu_d(\ell_n) \leq \nu_0 + C \min\left\{ (\nu_0^{2\alpha r} n)^{\frac{-\beta}{(d-1)(2\alpha r + 1)}}, n^{\frac{-\beta}{(d-1)(2\alpha r + 1) + 2\beta\alpha r}} \right\}. \tag{17}$$

Notice that when $\beta \ll d$, both exponents of n display a curse of dimensionality, but this curse goes away as β grows. Note also that the first exponent yields a faster rate, but one that is only achieved for large n due to the factor $\nu_0^{2\alpha r}$, which may be small for large groups.

Curse of dimensionality and optimality. Note that the obtained bound is still cursed for an invariant target f^*, in the sense that the exponent in the rate is of order $1/d$ when f^* is only assumed to be Lipschitz,[1] raising the question of whether this can be improved. We note that since $\gamma_d(k) = \Omega(1/|G|)$ (as we show in Section 4), the asymptotic decays (as a function of k) of the coefficients of f^* and of the eigenvalues of T_{K_G} are similar to those for the non-invariant case, which implies that these rates cannot be improved (see, *e.g.*, [7]), and that constant-factor improvements in the sample complexity may be the best we can hope for. Nevertheless, this may still lead to significant gains in practice.

4 Counting Invariant Polynomials

In this section, we study the asymptotic behavior of $\gamma_d(k)$, given in (7), when $k \to \infty$ and the dimension d and the group G are fixed. This quantity can be seen as capturing the fraction of orthogonal spherical harmonics of degree k that are invariant to G, and helps us control the possible gains in sample complexity for learning invariant functions, as described in Section 3. Denoting $\gamma_{d,\sigma}(k) := \mathbb{E}_x[P_{d,k}(\langle \sigma \cdot x, x\rangle)]$, we will show that $\gamma_{d,\sigma}(k)$ vanishes for large k for any σ that is not the identity. This implies that $\gamma_d(k)$ converges to $1/|G|$, since we trivially have $\gamma_{d,Id}(k) = P_{d,k}(1) = 1$. We further characterize the asymptotic behavior of $\gamma_d(k)$ in terms of properties of the group elements. In the following we consider the case of G being a subgroup of S_d, the groups of permutations on d elements.

Decay of $\gamma_{d,\sigma}(k)$. Our main insight is to leverage the fact that when σ is not the identity, then the random variable $z_\sigma = \langle \sigma \cdot x, x\rangle$ when $x \sim \tau$ admits a density on $[-1, 1]$, which we denote q_σ. This by itself will prove sufficient to show that $\gamma_{d,\sigma}(k)$ decays for large k, thanks to the oscillatory behavior of $P_{d,k}$. We can then further characterize its asymptotic behavior by studying the singularities of q_σ, leveraging the seminal work of Saldanha and Tomei [25]. In particular, these depend on spectral properties of the matrix associated to σ. We summarize this in the next proposition.

Proposition 6 (Asymptotic behavior of $\gamma_{d,\sigma}(k)$.)**.** *Let A_σ be the matrix associated to $\sigma \neq \mathrm{Id}$, that is such that $\sigma \cdot x = A_\sigma x$. Denote by Λ_σ the set of (complex) eigenvalues of A_σ, and by m_λ the multiplicity of $\lambda \in \Lambda_\sigma$. When $k \to \infty$, we have the asymptotic equivalent $\gamma_{d,\sigma}(k) = \sum_{\lambda \in \Lambda_\sigma} \gamma_{d,\sigma,\lambda}(k)$, where*

$$\gamma_{d,\sigma,\lambda}(k) \lesssim \begin{cases} k^{-d+m_\lambda} + o(k^{-d+m_\lambda}), & \text{if } \lambda \in \{\pm 1\}, \\ k^{-d+m_\lambda+5} + o(k^{-d+m_\lambda+5}), & \text{otherwise}, \end{cases} \tag{18}$$

where $\lesssim$ hides constants that may depend on d, σ and λ.

Every permutation $\sigma \in \mathcal{S}_d$ (where $\mathcal{S}_d$ is the symmetric group of permutations) can be decomposed into cycles on disjoint orbits; the eigenvalues λ (and their multiplicities m_λ) of a matrix A_σ admit an interpretation based on such decomposition. Indeed, since A_σ is unitary, its eigenvalues are of the form $\lambda = e^{2\pi i \theta}$, and one can verify that necessarily $\theta = \frac{p}{q} \in \mathbb{Q}$. Furthermore, assuming w.l.o.g. that q is prime, such eigenvalue appears whenever σ contains a cycle of length a multiple of q. In particular, the multiplicity of the eigenvalue 1, m_1, corresponds to the total number of cycles in such a decomposition, which we will denote by $c(\sigma)$. Then $\gamma_{d,\sigma}(k)$ can be controlled as follows.

Corollary 7 (Decay of $\gamma_{d,\sigma}(k)$)**.** *Let $\sigma \neq \mathrm{Id}$, and let $c(\sigma)$ denote the number of cycles in σ. Then,*

$$\gamma_{d,\sigma}(k) \lesssim \begin{cases} k^{-d+c(\sigma)}, & \text{if } c(\sigma) > \frac{d}{2} + 2, \\ k^{-d/2+7}, & \text{otherwise}. \end{cases}$$

Decay on specific subgroups. We may now use Corollary 7 to study the asymptotic behaviour of $\gamma_d(k)$ as $k \to \infty$ for various choices of subgroups of $\mathcal{S}_d$, using the following result.

Corollary 8 (Upper bounds with permutation statistics)**.** *Let G be a subset of $\mathcal{S}_d$ and define*

$$\zeta(G, n) := |\{\sigma \in G : c(\sigma) > n\}|$$

for $n \in [d-1]$. Then we have

$$\gamma_d(k) \leq \frac{\zeta(G, n)}{|G|} + O\left(k^{-d+\max\{n,\, d/2+7\}}\right),$$

with equality if $n = d-1$.

---

[1]This corresponds to taking r and α such that $2\alpha r \approx 2/(d-1)$, leading to a rate $n^{-2/(2+d-1)}$.

Example 9 (Translations). *Let $G = C_d$ be the cyclic group on d elements. Then it holds*

$$\gamma_d(k) = \frac{1}{d} + O\left(k^{-d/2+7}\right) .$$

This follows by noticing that every translation σ (but the identity) satisfies $c(\sigma) \leq d/2$.

Example 10 (Local translations). *Let $d = s \cdot r$ (with $r, s \geq 5$), and consider the group composed of traslations over r blocks of coordinates of size s; i.e., the block-cyclic group*

$$G = \{\sigma \; : \; \sigma = \sigma^{(1)} \circ \cdots \circ \sigma^{(r)}\}$$

where each $\sigma^{(i)}$ is a translation over the set $\{(i-1)s+1, \ldots, is\}$, for $i \in [r]$. Then it holds

$$\gamma_d(k) = \frac{1}{s^r} + O\left(k^{-s/2+1}\right) . \tag{19}$$

This follows by noticing that every local traslation σ (but the identity) satisfies $c(\sigma) \leq (d-s) + s/2$.

Example 11 (Full permutation group). *For the case of $G = \mathcal{S}_d$, we can split the group based on the value of $\mathrm{Fix}(\sigma)$, the number of elements fixed by a permutation σ. Denote*

$$\xi(G, n) := |\sigma \in G \; : \; \mathrm{Fix}(\sigma) > n| = \sum_{j=n+1}^{d} \binom{d}{j} !(d-j) .$$

for $n \in [d-1]$, where $!k$ denotes the k-th subfactorial. Then we have

$$\gamma_d(k) \leq \frac{\xi(G, n)}{d!} + O\left(k^{-d/2+\max\{n/2,\, 7\}}\right) ,$$

with equality for $n = d - 1$. This follows from the fact that $c(\sigma) \leq \mathrm{Fix}(\sigma) + (d - \mathrm{Fix}(\sigma))/2$. In particular, it follows

$$\gamma_d(k) \leq \frac{2}{(n+1)!} + O\left(k^{-d/2+\max\{n/2,\, 7\}}\right) .$$

Using different values of n may yield different upper bounds with faster decays, but larger constants.

5 Beyond Group Invariance: Geometric Stability

In this section, we study gains in sample complexity when the target function f^* is not fully invariant to a group, but may be stable under small geometric changes on the input. We formalize this by considering a similar averaging operator S_G, but we allow G to be a generic set of permutations instead of a group, and allow for a weighted average:

$$S_G f(x) := \sum_{\sigma \in G} h(\sigma) f(\sigma \cdot x), \tag{20}$$

where $h(\sigma) \geq 0$ for all $\sigma \in G$ and $\sum_{\sigma \in G} h(\sigma) = 1$. We assume that G is "symmetric", *i.e.*, $\sigma^{-1} \in G$ when $\sigma \in G$ and $h(\sigma^{-1}) = h(\sigma)$, so that S_G is self-adjoint. In this case, images of S_G are not invariant functions, but may nevertheless exhibit a form of "local" stability to small perturbations of the input data. For instance, if G consists of local translations by at most a few pixels, or if G consists of all translations but h is localized around the identity, then applying S_G yields functions that are stable to local translations. We may also consider a more structured set G of permutations that resemble local deformations, consisting of both a global translation as well as different local translations at different scales, as we describe below.

Spectral properties of S_G. Note that in this setup, we no longer have that S_G is a projection, however we may still view it as a *smoothing* operator, which attenuates certain harmonics that are "less" invariant than others. The next lemma shows related spectral properties to the invariant case.

Lemma 12 (Spectral properties of S_G). *There exists a basis of spherical harmonics $\overline{Y}_{k,j}$, for $k \geq 0$, and $j = 1, \ldots, N(d, k)$, in which the operator S_G is diagonal, with eigenvalues $\lambda_{k,j} \geq 0$. In analogy to Lemma 1, we have*

$$\gamma_d(k) := N(d, k)^{-1} \sum_{j=1}^{N(d,k)} \lambda_{k,j} = \sum_{\sigma \in G} h(\sigma) \mathbb{E}_x \left[P_{d,k}(\langle \sigma \cdot x, x \rangle) \right]. \tag{21}$$

We also define $\nu_d(\ell) := \sup_{k \geq \ell} \gamma_d(k)$.

Sample complexity of stable kernel. In analogy to Section 3, we may consider a stable kernel

$$K_G(x, x') = \sum_{\sigma \in G} h(\sigma)\kappa(\langle \sigma \cdot x, x' \rangle). \tag{22}$$

Then, it is easy to check that the integral operator of K_G is given by $T_{K_G} = S_G T_K$. In contrast to Section 3, we no longer have that the approximation errors of K and K_G are the same in general on "geometrically stable" functions, since the notion is not precisely defined. Nevertheless, we may represent favorable targets f^* as those whose coefficients decay similarly at each frequency k to those of S_G, by viewing it as a smoothing of some L^2 function g^*, *i.e.*, $f^* = S_G^r g^*$ for some exponent r. With this in mind, we make the following assumptions, replacing assumptions (A1-3) of Section 3.

(A5) *capacity*: the eigenvalues $(\xi_m)_{m \geq 0}$ of T_K satisfy $\xi_m \leq C(m+1)^{-\alpha}$.

(A6) *source condition*: there exists $r > \frac{\alpha-1}{2\alpha}$ and $g \in L^2(d\tau)$ with $\|g\|_{L^2(d\tau)} \leq C_{f^*}$ such that $f^* = S_G^r T_K^r g$.

Note that (A6) corresponds to a standard source condition with the kernel K_G (since $T_{K_G}^r = S_G^r T_K^r$), yet it reveals how K_G jointly performs smoothing on the sphere, through T_K, as well as on the permutation group through S_G. While these two forms of smoothing appear "entangled" in this assumption, one may balance them by choosing different levels of smoothing in the kernel function κ, or by averaging multiple times in (22). Assumption (A5) is needed for obtaining a variant of Lemma 4, and implies (A1) with $C_K \propto C^{1/\alpha}$. We then obtain the following generalization bound.

Theorem 13 (Generalization with geometric stability.). *Assume (A4-6), and assume* $\nu_0 := \inf_{\ell \geq 0} \nu_d(\ell) > 0$. *Let* $n \geq \max(\|f^*\|_\infty^2 / \sigma_\rho^2, (C_1/\nu_0)^{1/(2\alpha r+1-\alpha)})$, *and define*

$$\ell_n := \sup\{\ell : D(\ell) \leq C_2 \nu_d(\ell)^{\frac{2r}{2\alpha r+1}} n^{\frac{1}{2\alpha r+1}}\}. \tag{23}$$

We then have, for $\lambda = C_3(\nu_d(\ell_n)^{1/\alpha}/n)^{\alpha/(2\alpha r+1)}$,

$$\mathbb{E}[R(\hat{f}_\lambda) - R(f^*)] \leq C_4 \left(\frac{\nu_d(\ell_n)^{1/\alpha}}{n}\right)^{\frac{2\alpha r}{2\alpha r+1}}. \tag{24}$$

In the same setting, KRR with kernel K obtains the same bound with $\nu_d(\ell_n)^{1/\alpha}$ replaced by 1. Here, the constants $C_{1:4}$ only depend on the parameters of assumptions (A4-6).

Note that the theorem is very similar to Theorem 5, albeit with a gain in sample complexity dictated by $\nu_d(\ell_n)^{1/\alpha}$ instead of $\nu_d(\ell_n)$. This is due to the fact that in contrast to the invariant case, where $\gamma_d(k)$ in (7) can help precisely control the number of invariant spherical harmonics, in this case $\gamma_d(k)$ as computed in (21) can only give information about the sum of the eigenvalues $\lambda_{k,j}$ at frequency k, which may be insufficient to precisely estimate the gains in effective dimension. For smooth kernels with fast decays (large α), this discrepancy can weaken the sample complexity gains shown by our bounds, but for slow decays ($\alpha \approx 1$), the gains are similar to the invariant case.

Deformation-like stability. For inputs x defined as signals $x \in L^2(\Omega)$ over a continuous domain $\Omega \subseteq \mathbb{R}^s$, $s = 1, 2$, the action of 'small' diffeomorphisms $\varphi : \Omega \to \Omega$ as $(\varphi \cdot x)(u) = x(\varphi^{-1}(u))$ is a powerful diagnostic of performance of trainable CNNs [22], and a key guiding principle for scattering representations [19, 6]. In these works, the basic deformation cost is measured as $\|\varphi\| := \sup_u \|\nabla\varphi(u) - \mathbf{I}\|$. We instantiate an equivalent of small deformations in our finite-dimensional setting as follows.

$$\Phi_\varepsilon := \{\sigma \in \mathcal{S}_d : |\sigma(u) - \sigma(u') - (u - u')| \leq \varepsilon|u - u'|\} \qquad G_\epsilon := \{\sigma\sigma'^{-1} : \sigma, \sigma' \in \Phi_\varepsilon\},$$

where the differences are taken modulo d, and G_ε symmetrizes Φ_ε so that S_{G_ε} (considered here with a constant h) is self-adjoint. For $\varepsilon = 0$, we recover the translation group described in Example 9. We can verify that $\varepsilon = 1$ also corresponds to the translation group (due to the constraint that $\sigma(u) \neq \sigma(u')$ whenever $u \neq u'$), thus the first non-trivial model corresponds to $\varepsilon = 2$.

Proposition 14 (Upper bound on $\gamma_d(k)$ for deformations.). *Let $\Phi = \Phi_2$ and $G = G_2$ as above. Then it holds $|\Phi| \geq 2.16^d$, and as a consequence $|G| \geq |\Phi| \geq 2.16^d$. We then have*

$$\gamma_d(k) \leq 2.16^{-d} + O(k^{-1}).$$

We thus verify that small deformations, already with $\varepsilon = 2$, provide a substantial gain relative to rigid translations, since $|G|$ now grows exponentially with the dimension, rather than linearly. We note that faster decays may be obtained using a finer analysis of the cycle statistics of G, as in Example 11.

Figure 1: Comparison of KRR with invariant and non-invariant kernels. (left) permutation-invariant target with $d = 6$, comparison between various invariant kernels (cyclic, block-cyclic, and permutation groups). (center/right) invariant vs non-invariant kernels on invariant target functions with $d = 12$, for block-cyclic groups G of two different sizes.

6 Numerical Experiments

In this section, we provide simple numerical experiments on synthetic data which illustrate our theoretical results. In Figure 1, we consider KRR on 5 000 training samples with inputs uniformly distributed on $\mathbb{S}^{d-1}$, and outputs generated according to a target non-smooth function $f^* = S_G g^*$, with $g^*(x) = \mathbb{1}\{w_*^\top x \geq 0.7\}$, where the averaging operator S_G is over different groups in each plot. The regularization parameter λ is optimized on 5 000 test samples. We use the dot-product kernel function $\kappa(u) = (u + 1)\kappa_1(u)$, where κ_1 is the arc-cosine kernel of degree 1, which corresponds to an infinite-width shallow ReLU network [8].

When the target is permutation-invariant, we can see in Figure 1(left) that the kernel based on permutation invariance leads to the largest gain in sample complexity compared to those which use cyclic or block-cyclic groups. Since the permutation group has the largest cardinality, this is consistent with our finding that the gains may be of the order of the size of the group. Figures 1(center/right) consider the example of cyclic translations on local blocks of size 2 or 6 (Example 10 with $s = 2$ or 6), and illustrate that the improvement in sample complexity happens later for $s = 2$ than $s = 6$, which is consistent with the slower decays of γ_d obtained in (19) due to the larger number of cycles.

7 Discussion and Conclusion

We have studied how geometric invariance or stability assumptions on target functions enable more efficient learning, with improvements in sample complexity which may be as large as the number of permutations considered in the group or set of elements to which the target is invariant or stable. In particular, this gain can be exponential in the dimension if we consider, *e.g.*, all permutations, local translations on small blocks, or permutations that resemble small deformations. This last example provides a strong justification for seeking models and architectures that are stable to deformations, a natural prior when learning functions on images [22]. In that respect, our results provide a theoretical baseline to assess learning guarantees under geometric priors: by designing appropriate geometrically stable kernels, we simultaneously address approximation and generalisation errors within a framework of convex optimization.

That said, while these gains may be large in practice, the obtained rates may still be cursed by dimension if the target is non-smooth. This suggests that further assumptions may be needed to learn efficiently on high-dimensional geometric data, which may be exploited perhaps through richer architectures that involve local connectivity, hierarchy, or feature learning to exploit regularities in the target function beyond what our invariant/stable kernels can capture. Another interesting question is to study whether it is possible to adapt to general symmetries present in the target, instead of encoding them in the model with an appropriate kernel as we do here.

References

[1] K. Atkinson and W. Han. *Spherical harmonics and approximations on the unit sphere: an introduction*, volume 2044. Springer Science & Business Media, 2012.

[2] F. Bach. Breaking the curse of dimensionality with convex neural networks. *Journal of Machine Learning Research (JMLR)*, 18(1):629–681, 2017.

[3] F. Bach. *Learning Theory from First Principles (draft)*. 2021.

[4] A. Bietti and F. Bach. Deep equals shallow for ReLU networks in kernel regimes. In *Proceedings of the International Conference on Learning Representations (ICLR)*, 2021.

[5] A. Bietti and J. Mairal. Group invariance, stability to deformations, and complexity of deep convolutional representations. *Journal of Machine Learning Research (JMLR)*, 20(25):1–49, 2019.

[6] J. Bruna and S. Mallat. Invariant scattering convolution networks. *IEEE Transactions on Pattern Analysis and Machine Intelligence (PAMI)*, 35(8):1872–1886, 2013.

[7] A. Caponnetto and E. De Vito. Optimal rates for the regularized least-squares algorithm. *Foundations of Computational Mathematics*, 7(3):331–368, 2007.

[8] Y. Cho and L. K. Saul. Kernel methods for deep learning. In *Advances in Neural Information Processing Systems (NIPS)*, 2009.

[9] C. Ciliberto, F. Bach, and A. Rudi. Localized structured prediction. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2019.

[10] N. Cohen and A. Shashua. Inductive bias of deep convolutional networks through pooling geometry. *arXiv preprint arXiv:1605.06743*, 2016.

[11] F. Cucker and S. Smale. On the mathematical foundations of learning. *Bulletin of the American mathematical society*, 39(1):1–49, 2002.

[12] S. S. Du, Y. Wang, X. Zhai, S. Balakrishnan, R. Salakhutdinov, and A. Singh. How many samples are needed to estimate a convolutional neural network? In *Advances in Neural Information Processing Systems (NeurIPS)*, 2018.

[13] C. Efthimiou and C. Frye. *Spherical harmonics in p dimensions*. World Scientific, 2014.

[14] B. Elesedy and S. Zaidi. Provably strict generalisation benefit for equivariant models. *arXiv preprint arXiv:2102.10333*, 2021.

[15] S. Fischer and I. Steinwart. Sobolev norm learning rates for regularized least-squares algorithms. *Journal of Machine Learning Research (JMLR)*, 21(205):1–38, 2020.

[16] T. Hastie, R. Tibshirani, and J. Friedman. *The elements of statistical learning: data mining, inference, and prediction*. Springer Science & Business Media, 2009.

[17] J. Mairal, P. Koniusz, Z. Harchaoui, and C. Schmid. Convolutional kernel networks. In *Advances in Neural Information Processing Systems (NIPS)*, 2014.

[18] E. Malach and S. Shalev-Shwartz. Computational separation between convolutional and fully-connected networks. In *Proceedings of the International Conference on Learning Representations (ICLR)*, 2021.

[19] S. Mallat. Group invariant scattering. *Communications on Pure and Applied Mathematics*, 65(10):1331–1398, 2012.

[20] S. Mei, T. Misiakiewicz, and A. Montanari. Learning with invariances in random features and kernel models. In *Conference on Learning Theory (COLT)*, 2021.

[21] H. N. Mhaskar and T. Poggio. Deep vs. shallow networks: An approximation theory perspective. *Analysis and Applications*, 14(06):829–848, 2016.

[22] L. Petrini, A. Favero, M. Geiger, and M. Wyart. Relative stability toward diffeomorphisms in deep nets indicates performance. *arXiv preprint arXiv:2105.02468*, 2021.

[23] L. Pillaud-Vivien, A. Rudi, and F. Bach. Statistical optimality of stochastic gradient descent on hard learning problems through multiple passes. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2018.

[24] T. Poggio, H. Mhaskar, L. Rosasco, B. Miranda, and Q. Liao. Why and when can deep-but-not shallow-networks avoid the curse of dimensionality: a review. *International Journal of Automation and Computing*, 14(5):503–519, 2017.

[25] N. C. Saldanha and C. Tomei. The accumulated distribution of quadratic forms on the sphere. *Linear algebra and its applications*, 245:335–351, 1996.

[26] A. J. Smola, Z. L. Ovari, and R. C. Williamson. Regularization with dot-product kernels. In *Advances in Neural Information Processing Systems (NIPS)*, 2001.

[27] M. J. Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.