FAST CONVERGENCE OF STOCHASTIC SUBGRADIENT METHOD UNDER INTERPOLATION

ABSTRACT

This paper studies the behaviour of the stochastic subgradient descent (SSGD) method applied to over-parameterized nonsmooth optimization problems that satisfy an interpolation condition. By leveraging the composite structure of the empirical risk minimization problems. We prove that SSGD converges, respectively, with rates $\mathcal{O}(1/\epsilon)$ and $\mathcal{O}(\log(1/\epsilon))$ for convex and strongly-convex objectives when interpolation holds. These rates coincide with established rates for the stochastic gradient descent (SGD) method applied to smooth problems that also satisfy an interpolation condition. Our analysis provides a partial explanation for the empirical observation that sometimes SGD and SSGD behave similarly for training smooth and nonsmooth machine learning models. We also prove that the rate $\mathcal{O}(1/\epsilon)$ is optimal for the subgradient method in the convex and interpolation setting.

1 INTRODUCTION

Gradient descent (GD) and subgradient descent (subGD) methods are simple and effective first-order optimization algorithms for training machine learning models. The convergence-rate analyses for these methods depend crucially on the smoothness of the objective function. It is well understood that there is a fundamental gap between the convergence rates of gradient-like methods for smooth and nonsmooth problems (Shor, 1984; Nemirovski & Yudin, 1983; Nesterov, 2005; Bubeck, 2015; Beck, 2017). Table 1 summarizes the rates for the main problem classes.

However, in the practice of training machine learning models, the nonsmootheness from the model is not causing much trouble (Glorot et al., 2011; Goodfellow et al., 2016). Neural networks with nonsmooth activation function such as ReLU activation can usually be trained as fast as the one with smooth activation function such as softplus, see Figure 1.1 as a motivating experiment from us to compare the convergence of gradient and subgradient based methods for smooth and nonsmooth neural networks on the MNIST dataset[1] to classify digits 0 and 1. We can see that the gradient and subgradient based methods, either batch or stochastic version, has similar convergence behaviour for smooth and nonsmooth neural networks. Therefore there is a discrepancy between theory and practice.

The success of overparameterized models—such as deep and wide neural networks—has instigated a trend in the analysis of stochastic variants of gradient descent in the *interpolation* setting, in which the model achieves zero training loss, and therefore fits all of the training data (Schmidt & Le Roux, 2013; Bassily et al., 2018; Ma et al., 2018b; Jain et al., 2018; Vaswani et al., 2019a;b; Wu et al., 2019; Liu & Belkin, 2020). This recent series of papers offer insight into the fast convergence of SGD and new approaches for algorithm design. This line of analysis, however, focus exclusively on smooth objective function, and cannot explain the effectiveness of SSGD for training nonsmooth neural networks.

We present a formal analysis showing that SSGD for nonsmooth objectives could converge as fast as smooth objectives in the interpolation setting. Our contributions include:

- A description of semi-smoothness properties of a function useful for the iteration complexity analysis of convex objectives in the interpolation context. Under mild conditions, semi-

---

[1] http://yann.lecun.com/exdb/mnist/

objective $f(\mathbf{x})$	smooth	nonsmooth
convex	$\mathcal{O}(1/\epsilon)$	$\mathcal{O}(1/\epsilon^2)$
strongly convex	$\mathcal{O}(\log(1/\epsilon))$	$\mathcal{O}(1/\epsilon)$

Table 1: Worst-case iteration complexity of batch gradient and subgradient methods.

Figure 1.1: The convergence of gradient-based (GD and SGD) and subgradient-based (subGD and SSGD) methods for smooth and nonsmooth neural networks

smoothness under interpolation allows us to prove that SSGD has iteration complexity $\mathcal{O}(1/\epsilon)$ for convex objectives, and $\mathcal{O}(\log(1/\epsilon))$ for strongly-convex objectives. These rates match the convergence rates of SGD for convex and smooth objective under interpolation.

- Proof that the iteration bound $\mathcal{O}(1/\epsilon)$ is optimal in the convex and interpolation setting. In contrast to the case with a smooth objective function, subgradient-based methods cannot be further accelerated for nonsmooth model—even with the interpolation assumption.

2 RELATED WORK

2.1 FIRST-ORDER METHODS FOR NONSMOOTH OPTIMIZATION

The subgradient method and its convergence analysis for general convex and nonsmooth problems was first described by Shor in the late 1960s and 70s; see Shor (1984). Nemirovski & Yudin (1983) subsequently established that the iteration complexity $\mathcal{O}(1/\epsilon^2)$ described by Shor (1984) is optimal for methods that can only access a subgradient oracle. Subsequent works for minimizing general convex and nonsmooth problems include stochastic subgradient methods (Polyak, 1987; Shamir & Zhang, 2013), dual averaging (Nesterov, 2009), and acceleration via smoothing (Nesterov, 2005; Beck & Teboulle, 2012). More recently, Zhang et al. (2020) and Shamir (2020), among others, described the tractability and complexity of get approximate stationary point for general nonsmooth and nonconvex problems.

A related line of work involves the analysis of optimization algorithms for partially nonsmooth objectives that take the form $f(\mathbf{x}) + g(\mathbf{x})$, where the function f is smooth, and the function g, which usually represents a regularizer, is convex and nonsmooth. Many models in feature selection and compressed sensing fall into this category of problems, which are usually solved by variations of the proximal-gradient method (Nesterov, 2007; Beck & Teboulle, 2009; Xiao, 2010; Parikh & Boyd, 2014; Defazio et al., 2014; Allen-Zhu, 2017). These approaches typically use special properties of the regularization function g, which generally do not apply in the context we consider in this paper.

2.2 INTERPOLATION HELPS OPTIMIZATION

The interpolation condition implies that the residual between the prediction of the model and data vanishes. In the context of nonlinear least-squares, for example, it is known that interpolation allows the Gauss-Newton method to converge at a local superlinear rate, although in general it can only achieve a local linear rate (Bertsekas, 1997). Conditional gradient method can also enjoy a linear convergence rate for least square objective when interpolation holds (Beck & Teboulle, 2004). Recent interest in over-parameterized models for machine learning problems makes interpolation (or near interpolation) a relevant assumption for the analysis and development of new algorithms. Under interpolation, stochastic gradient descent enjoys the same convergence rate as its deterministic counterpart, gradient descent (Schmidt & Le Roux, 2013; Ma et al., 2018a; Bassily et al., 2018). Some techniques that originally developed for gradient descent, such as line-search and Nesterov acceleration, are proven to also hold for SGD when interpolation is satisfied (Vaswani et al., 2019b;a;

Algorithm 1 Stochastic subgradient descent. The learning rate function $\alpha_t : \mathbb{N} \to \mathbb{R}_+$ returns the learning rate at iteration t.

1: **Initialize:** $w^{(1)} \in \mathbb{R}^d$
2: **for** $t = 1, 2, \ldots$ **do**
3:     select $i \in \{1, 2, \ldots, n\}$ uniformly at random
4:     compute $g^{(t)} \in \partial f_i(w^{(t)})$
5:     $w^{(t+1)} = w^{(t)} - \alpha_t g^{(t)}$
6: **end for**

Liu & Belkin, 2020; Jain et al., 2018) Related work in this vein includes improved analysis of stochastic second-order optimization algorithms (Meng et al., 2020) and adaptive stochastic gradient descent (Loizou et al., 2020; Vaswani et al., 2020) under interpolation. Another line of work (Qian et al., 2019; Khaled & Richtárik, 2020) in this direction aims to explore assumptions other than the classic bounded noise assumption for the optimization theory of training over-parameterized model, such as the expected growth condition (Schmidt & Le Roux, 2013), expected smoothness (Qian et al., 2019; Gower et al., 2018), etc. All of the work just mentioned involves differentiable and smooth objective functions, and cannot explain the fast convergence of SSGD for nonsmooth objectives that is often observed in practice.

3   PROBLEM SETTING

We consider the unconstrained empirical risk-minimization problem

$$\underset{w \in \mathbb{R}^d}{\text{minimize}} \ \ f(w) := \frac{1}{n} \sum_{i=1}^{n} f_i(w) \qquad \text{where each} \qquad f_i(w) := \ell(h_i(w)) \tag{3.1}$$

and n is the number of data points. Throughout the paper, we use w^* to denote any solution of eq. (3.1), and thus $f^* := f(w^*)$ is the optimal objective value. We assume that the 1-dimensional loss function $\ell : \mathbb{R} \to \mathbb{R}_+$ is nonnegative, convex, and is 1-smooth, i.e.,

$$|\ell'(\alpha) - \ell'(\beta)| \le |\alpha - \beta| \quad \forall \alpha, \beta \in \mathbb{R}.$$

Without loss of generality, we also assume $\ell(0) = 0$. Common examples for the loss function include the 2-norm, negative entropy, and the 2-norm hinge loss function. The n functions h_i are Lipschitz continuous with respect to the fixed parameter L:

$$|h_i(w_1) - h_i(w_2)| \le L\|w_1 - w_2\| \quad \forall w_1, w_2 \in \mathbb{R}^d, \ \forall i \in [n].$$

We make no assumption on their smoothness properties. Many key machine learning tasks can be formulated as (3.1), including training deep neural networks with nonsmooth activations, such as the ReLU function. Here and throughout, the function $\|\cdot\|$ is the 2-norm a vector, unless otherwise specified.

Our analysis relies on the Clark *generalized* gradient (Clarke, 1990) of the nonconvex and nonsmooth function h, defined as the convex hull of all valid limiting gradients:

$$\partial h(w) := \text{conv}\left\{ u \mid u = \lim_{k \to \infty} \nabla h(w_k), \ w_k \to w \right\},$$

This definition additionally requires h to be almost everywhere differentiable by Rademacher's theorem. We refer readers to Clarke (1990), and more recently, to Zhang et al. (2020), who use this generalized gradient in a related analysis. These properties of the generalized gradient are needed for our analysis:

(chain rule) $\partial f_i(w) = \ell'(h(w)) \cdot \partial h(w),$

(gradient bound) $\|\partial h_i(w)\| \le L, \ \forall w \in \mathbb{R}^d, \ \forall i \in [n].$

The second property follows from the L-Lipschitz continuity of each function h_i. We define the norm of the generalized gradient at a vector w as

$$\|\partial h_i(w)\| = \sup\left\{ \|z\| \mid z \in \partial h_i(w) \right\}.$$

Algorithm 1 describes the stochastic subgradient descent (SSGD) method.

4 MAIN RESULTS

4.1 BOUNDS AND LIPSCHITZ PROPERTIES OF THE GENERALIZED GRADIENT

Our analysis hinges on establishing that the objective function f is differentiable at all of its minimizers, i.e., at all vectors w such that $f(w) = 0$. This is implied by the following proposition, which holds even without the interpolation condition.

Proposition 4.1 (Generalized gradient bound). *For all d-vectors w,*

$$\|\partial f_i(w)\|^2 \leq 2L^2 f_i(w) \qquad \forall i \in [n]. \tag{4.1}$$

Consequently,

$$\|\partial f(w)\|^2 \leq \frac{1}{n}\sum_{i=1}^n \|\partial f_i(w)\|^2 \leq 2L^2 f(w). \tag{4.2}$$

Equation (4.2) implies that if the objective value $f(w) = 0$, then the generalized gradient contains only the origin: $\partial f(w) = \{0\}$. It thus follows from Clarke (1981, Property 10) that f is differentiable at any solution with zero objective value. This means that any solution with a zero value is a fixed point of the subgradient method, which stands in contrast to a solution where the function is nonsmooth. This property of f makes it possible for the subgradient method to converge to solution with constant learning rate.

Proof of Proposition 4.1. Fix an arbitrary d-vector w. The subdifferential $\partial f_i(w) = \ell'(h_i(w)) \cdot \partial h_i(w)$. Because each function h_i is L-Lipschitz continuous, we can then deduce that

$$
\begin{aligned}
\|\partial f_i(w)\|^2 &\leq L^2 \left[\ell'(h_i(w))\right]^2 \\
&\overset{\text{(i)}}{=} L^2 (\ell'(h_i(w)) - \ell'(0))^2 \\
&\overset{\text{(ii)}}{\leq} 2L^2 (\ell(h_i(w)) - \ell(0)) \\
&= 2L^2 f_i(w).
\end{aligned} \tag{4.3}
$$

Step (i) follows from the assumption that $\ell(0) = \min_{\lambda \in \mathbb{R}} \ell(\lambda) = 0$, which implies $\ell'(0) = 0$. Step (ii) follows from the fact that any convex L-smooth function $u : \mathbb{R}^n \to \mathbb{R}$ satisfies the bound

$$u(y) - u(x) - \langle \nabla u(x), y - x \rangle \geq \frac{1}{2L}\|\nabla u(y) - \nabla u(x)\|^2;$$

see Nesterov (2014, Theorem 2.1.5). Make the identifications

$$u = \ell, \quad y = h_i(w), \quad x = 0$$

to immediately obtain eq. (4.3) and thus the proof for eq. (4.1).

Equation (4.2) can be obtained directly from Jensen's inequality:

$$\|\partial f(w)\|^2 = \left\|\frac{1}{n}\sum_{i=1}^n \partial f_i(w)\right\|^2 \leq \frac{1}{n}\sum_{i=1}^n \|\partial f_i(w)\|^2 \leq \frac{1}{n}\sum_{i=1}^n 2L^2 f_i(w) = 2L^2 f(w).$$

$\square$

The composite structure of the functions f_i allows us develop a semi-Lipschitz bound on their generalized gradients. Moreover, we show that it is possible to provide a global convex majorant for each function f_i, which holds without assuming convexity. Interestingly, eq. (4.5) overlaps with the "semi-smoothness" property of over-parameterized neural networks derived by Allen-Zhu et al. (2019, Theorem 4). In contrast to that result, however, we do not use any special properties of over-parameterized neural networks. Proposition 4.2 may thus be of more general interest.

> **Proposition 4.2** (Semi-smoothness)**.** *For all vectors w_1 and w_2, and each $i \in [n]$,*
>
> $$\|\partial f_i(w_2) - \partial f_i(w_1)\| \le L^2 \|w_2 - w_1\| + 2L\sqrt{2\min\{f_i(w_1), f_i(w_2)\}}, \qquad (4.4)$$
>
> *and*
>
> $$f_i(w_2) \le f_i(w_1) + \langle \partial f_i(w_1), w_2 - w_1 \rangle + \frac{L^2}{2}\|w_2 - w_1\|^2 + 2L\|w_2 - w_1\|\sqrt{2f_i(w_1)}. \tag{4.5}$$

See Appendix A for the proof of this result.

4.2 Convergence rate of stochastic subgradient descent

We now present a global convergence analysis for the SSGD algorithm under the additional assumption that the objective f of eq. (3.1) is convex. We develop a bound on the expected progress of the objective value that depends on the minimizing value f^*, rather than on the Lipschitz bound on the function itself, which is the usual bound in the literature. Our proof is based on a simple modification of the classical proof of subgradient descent method.

> **Theorem 4.3** (Global convergence rate of SSGD)**.** *Assume f is convex. Then for any positive integer T and any learning rate function α_t that satisfies $\sum_{t=1}^{T}(\alpha_t - L^2\alpha_t^2) > 0$,*
>
> $$\min_{t \in [T]} \mathbb{E}[f(w^t) - f^*] \le \frac{\|w^{(1)} - w^*\|^2 + 2L^2 f^* \sum_{t=1}^{T} \alpha_t^2}{2\sum_{t=1}^{T}(\alpha_t - L^2\alpha_t^2)}. \tag{4.6}$$

Proof. Each iterate $w^{(t)}$ satisfies the bound

$$\mathbb{E}[\|w^{(t+1)} - w^*\|^2 \mid w^{(t)}] = \frac{1}{n}\sum_{i=1}^{n}\|w^{(t)} - \alpha_t g_i^{(t)} - w^*\|^2$$

$$= \|w^{(t)} - w^*\|^2 - 2\alpha_t \left\langle \frac{1}{n}\sum_{i=1}^{n} g_i^{(t)}, w^{(t)} - w^* \right\rangle + \frac{1}{n}\sum_{i=1}^{n}\alpha_t^2 \|g_i^{(t)}\|^2$$

$$\overset{(i)}{\le} \|w^{(t)} - w^*\|^2 - 2\alpha_t(f(w^{(t)}) - f(w^*)) + \frac{1}{n}\sum_{i=1}^{n}\alpha_t^2 \|g_i^{(t)}\|^2 \qquad (4.7)$$

$$\overset{(ii)}{\le} \|w^{(t)} - w^*\|^2 - 2\alpha_t(f(w^{(t)}) - f(w^*)) + 2\alpha_t^2 L^2 f(w^{(t)}), \qquad (4.8)$$

where (i) follows from the convexity of f, and (ii) follows from Proposition 4.1. Take expectations on both sides of the inequality (4.8) and rearrange to obtain

$$(2\alpha_t - 2L^2\alpha_t^2) \cdot \mathbb{E}[(f(w^{(t)}) - f^*)] \le \mathbb{E}[\|w^{(t)} - w^*\|^2] - \mathbb{E}[\|w^{(t+1)} - w^*\|^2] + 2L^2\alpha_t^2 f^*. \tag{4.9}$$

Summing inequality (4.9) over $t \in \{1, 2, \dots, T\}$ yields

$$\sum_{t=1}^{T}(2\alpha_t - 2L^2\alpha_t^2)\mathbb{E}[f(w^t) - f^*] \le \|w^{(1)} - w^*\|^2 + 2L^2 f^* \sum_{t=1}^{T}\alpha_t^2.$$

Divide both sides by $2\sum_{t=1}^{T}(\alpha_t - L^2\alpha_t^2) > 0$ to obtain the required result. $\qquad \square$

This proof mirrors closely the classical proof of SSGD, which assumes that the subdifferential of each f_i is bounded, i.e., $\|\partial f_i(w)\| \le G$ for some constant G. We use Proposition 4.1 in eq. (4.7) to avoid the bounded subgradient assumption and to express the convergence rate using the minimal value f^*. This modification allows us to leverage the interpolation assumption that $f^* = 0$. We use Theorem 4.3 to immediately deduce the following convergence rate result.

Corollary 4.4 (Global convergence rate of SSGD with constant learning rate). *Assume that f is convex and that the learning rate $\alpha_t = 1/(2L^2)$ is constant for all $t > 0$. Then for any positive integer T, the SSGD iterates w^t satisfy*

$$\min_{t \in [T]} \mathbb{E}[f(w^t) - f^*] \leq (2L^2/T)\|w^{(1)} - w^*\|^2 + f^*.$$

Furthermore, when $f^* = 0$ (interpolation holds),

$$\min_{t \in [T]} \mathbb{E}[f(w^t)] - f^*] \leq (2L^2/T)\|w^{(1)} - w^*\|^2.$$

Corollary 4.4 indicates that the SSGD method converges at rate $\mathcal{O}(1/\epsilon)$ when interpolation holds. This rate matches the convergence rate of SGD for smooth objective functions under interpolation (Schmidt & Le Roux, 2013). When interpolation does not hold, Corollary 4.4 implies that SSGD, on expectation, could obtain objective value lower than $2f^* + \epsilon$ in $\mathcal{O}(1/\epsilon)$ time, which could be close to f^* when f^* is close to 0 (interpolation nearly holds).

Next, we derive convergence rate under the stronger assumption that f is strongly convex, which means that

$$f(w_1) \geq f(w_2) + \langle g, w_1 - w_2 \rangle + \frac{\mu}{2}\|w_1 - w_2\|^2 \quad \forall w_1, w_2 \in \mathbb{R}^d, \ \forall g \in \partial f(w_2) \tag{4.10}$$

for some constant $\mu > 0$. Note that recent works indicated that in order to prove linear convergence rate, the strong convexity assumption can be relaxed to other weaker assumptions that could hold even for some nonconvex functions (Karimi et al., 2016; Qian et al., 2019). For simplicity, we assume strong convexity in our analysis.

Theorem 4.5 (Global convergence rate of SSGD under strong convexity). *Assume that f is μ-strongly convex and that the learning rate $\alpha_t = 1/L^2$ is constant for all $t > 0$. Then for any positive integer T, the SSGD iterates $w^{(t)}$ satisfy*

$$\mathbb{E}[\|w^{(T)} - w^*\|^2] \leq \left(1 - \frac{\mu}{2L^2}\right)^{T-1} \|w^{(1)} - w^*\|^2 + \frac{4}{\mu}f^*. \tag{4.11}$$

Proof. Use the definition of the SSGD iteration to obtain

$$\mathbb{E}[\|w^{(t+1)} - w^*\|^2 \mid w^{(t)}]$$

$$\stackrel{(i)}{=} \|w^{(t)} - w^*\|^2 - 2\alpha_t \left\langle \frac{1}{n}\sum_{i=1}^n g_i^{(t)}, w^{(t)} - w^* \right\rangle + \frac{1}{n}\sum_{i=1}^n \alpha_t^2 \|g_i^{(t)}\|^2$$

$$\stackrel{(ii)}{\leq} \|w^{(t)} - w^*\|^2 - 2\alpha_t(f(w^{(t)}) - f^*) - \frac{\mu\alpha_t}{2}\|w^* - w^{(t)}\|^2 + \frac{1}{n}\sum_{i=1}^n \alpha_t^2 \|g_i^{(t)}\|^2$$

$$\stackrel{(iii)}{\leq} \left(1 - \frac{\mu\alpha_t}{2}\right)\|w^{(t)} - w^*\|^2 - 2\alpha_t(f(w^{(t)}) - f^*) + 2\alpha_t^2 L^2 f(w^{(t)})$$

$$\stackrel{(iv)}{=} \left(1 - \frac{\mu}{2L^2}\right)\|w^{(t)} - w^*\|^2 + \frac{2}{L^2}f^*, \tag{4.12}$$

where (i) follows from the same argument of the proof of Theorem 4.3; (ii) follows from the μ-strong convexity of f (see eq. (4.11)); (iii) follows from eq. (4.1); and (iv) follows from the definition of the learning rate $\alpha_t = \frac{1}{L^2}$.

Taking expectation to both sides of eq. (4.12) and recursively apply it to $t \in \{1, 2, \ldots, T\}$ to deduce

$$\mathbb{E}[\|w^{(T)} - w^*\|^2] \leq \left(1 - \frac{\mu}{2L^2}\right)^{T-1} \|w^{(1)} - w^*\|^2 + \sum_{t=0}^{T-2} \left(1 - \frac{\mu}{2L^2}\right)^t \frac{2}{L^2}f^*$$

$$\stackrel{(i)}{\leq} \left(1 - \frac{\mu}{2L^2}\right)^{T-1} \|w^{(1)} - w^*\|^2 + \frac{4}{\mu}f^*, \tag{4.13}$$

where (i) follows from the fact that $\sum_{t=0}^{\infty}(1 - \frac{\mu}{2L^2})^t = 2L^2/\mu$.

$\square$

Theorem 4.5 indicates that the SSGD can converges to the ball centered at w^* with radius $2\sqrt{f^*/\mu}$ at a linear rate. If additionally interpolation holds, SSGD converges to the solution linearly. Again, this rate matches the convergence rate of SGD for smooth and strongly convex objectives in the interpolation setting (Schmidt & Le Roux, 2013; Ma et al., 2018a). Similar to Corollary 4.4, when interpolation is nearly satisfied, SSGD converges linearly to an $2\sqrt{f^*/\mu}$-approximate solution.

Corollary 4.4 and Theorem 4.5 also provide insight into the effect of learning rate schedules on the performance of SSGD. A learning rate schedule α_t that decays as $t^{-1/2}$ is optimal because it causes the algorithm to exhibit a complexity bound of $\mathcal{O}(1/\epsilon^2)$, which is the theoretical lower bound (Nemirovski & Yudin, 1983). However, the learning rate schedule $\mathcal{O}(t^{-1/2})$ is slow in practice. Corollary 4.4 and Theorem 4.5 partially explain the discrepancy between the theory and practice: many machine learning models exhibit interpolation or near interpolation, and an aggressive constant learning-rate schedule works better than the conservative worst-case optimal learning rate $\alpha_t = \mathcal{O}(t^{-1/2})$.

4.3 LOWER BOUNDS

We have proven that, under interpolation, SGD for smooth problems and SSGD for nonsmooth problems exhibit the same convergence rates. This causes us to consider the following questions:

- Is it possible to induce momentum-type acceleration for SSGD under interpolation? Vaswani et al. (2019a) and Liu & Belkin (2020) showed recently that acceleration is possible for SGD under interpolation. It seems plausible that similar techniques could be used for nonsmooth problems.

- Can the composite structure $f_i = \ell \circ h_i$, which is central to our analysis, be used to establish an improved convergence rate for SSGD without interpolation? In other words, we know the lower bound on the iteration complexity for any subgradient method for nonsmooth functions is $\Omega(1/\epsilon^2)$. What, then, is the lower bound for minimizing the structured nonsmooth function $\ell \circ h$?

Unfortunately, the answer to these questions is "no", as we show below. First, we derive the lower iteration bound for any algorithm with access only to a subgradient oracle in the interpolation setting.

Theorem 4.6 (Lower bound with interpolation). *Given $t < d$ and positive constants L and R, and an initial vector $w^{(1)}$. Let $\ell = (\cdot)^2$. Then there exists an L-Lipschitz function h such that $f := l \circ h$ is convex, $f^* = \min_w f(w) = 0$, $\|w^{(1)} - w^*\| \le R$ such that*

$$\min_{1 \le s \le t} f(w^{(s)}) - f^* \ge \frac{L^2 R^2}{t+1}. \tag{4.14}$$

Proof. With out loss of generality, we assume the initial point $w^{(1)} = 0$. Let $h(w) = L\|w - w^*\|_\infty$, where

$$w^* = \Big[\underbrace{\frac{R}{\sqrt{t+1}}, \ldots, \frac{R}{\sqrt{t+1}}}_{t+1 \text{ entries}}, \underbrace{0, \ldots, 0}_{(d-t-1) \text{ entries}} \Big].$$

It is easy to check that ℓ is 1-smooth, h is L-Lipschitz, $f = \ell \circ h$ is convex, and $f^* = 0$.

We follow Nemirovski & Yudin (1983) and assume without loss of generality that any algorithm that uses only subgradient information generates iterates $w^{(s)} \in \text{span}\{e_1, e_2, \ldots, e_t\}$ for all $s \le t$. Therefore, for all $s \le t$, $f(w^{(s)}) - f^* = L^2 R^2/(t+1)$, as required. $\square$

Theorem 4.6 established $\Omega(1/\epsilon)$ iteration complexity for subgradient method under interpolation. Combining this result with the $\mathcal{O}(1/\epsilon)$ iteration complexity in Corollary 4.4, we can conclude that the rate $\mathcal{O}(1/\epsilon)$ is optimal in the interpolation setting and no acceleration is possible for subgradient-based methods. Then we proceed to the lower bound with assuming interpolation.

Theorem 4.7 (Lower bound without interpolation). *Given $t < d, L, R > 0$ and an initial point $w^{(1)}$. Let $\ell(\cdot) = (\cdot)^2$, and there exist L-Lipschitz function h satisfying $f := l \circ h$ is convex, $\|w^{(1)} - w^*\| \le R$*

such that

$$\min_{1 \leq s \leq t} f(w^{(s)}) - f^* \geq \frac{2L^2 R^2}{\sqrt{t+1}}. \tag{4.15}$$

Proof. Similar to the proof of Theorem 4.6, we set $w^{(1)} = 0$. But we change the construction of $h(w)$ to $h(w) = L(\|w - w^*\|_\infty + R)$, and w^* is defined in the same way as in the proof of Theorem 4.6:

$$w^* = \Big[\underbrace{\frac{R}{\sqrt{t+1}}, \ldots, \frac{R}{\sqrt{t+1}}}_{t+1 \text{ entries}}, \underbrace{0, \ldots, 0}_{(d-t-1) \text{ entries}} \Big].$$

Following the same analysis of the proof of Theorem 4.6. $w^{(s)} \in \text{span}\{e_1, e_2, \ldots, e_t\} \ \forall s \leq t$ and

$$f(w^{(s)}) - f^* = L^2 \left(\frac{R}{\sqrt{t+1}} + R \right)^2 - L^2 R^2 \geq \frac{2L^2 R^2}{\sqrt{t+1}} \qquad \forall s \leq t,$$

which completes the proof. $\square$

Theorem 4.7 provides an $\Omega(1/\epsilon^2)$ iteration complexity for subgradient-based methods for solving the structured function $\ell \circ h$. This matches the lower bound for solving general nonsmooth objective function with subgradient-based method, and implies that the structure $\ell \circ h$ itself without interpolation cannot give us an improved iteration complexity of subgradient-based methods.

5 CONCLUSION AND DISCUSSION

An empirical minimization problem based on composite functions has sufficient structure to allow for a tight convergence analysis that explains the effectiveness of stochastic subgradient descent methods on nonsmooth problems with interpolation. Surprisingly, the complexity bounds $\mathcal{O}(1/\epsilon)$ and $\mathcal{O}(\log(1/\epsilon))$ that we prove under interpolation match those of stochastic gradient descent for smooth functions.

Our convergence analysis is based on the the convexity properties of f. As we mention in connection with Theorem 4.5, the strong convexity assumption can be relaxed to weaker conditions, but even these exclude important nonconvex models that appear in neural networks. It is still an open problem as to whether a linear convergence rate for subgradient methods under a weaker assumptions, such as the restricted secant inequality (RSI). In Section 4.3 we proved that the rate $\mathcal{O}(1/\epsilon)$ is optimal for subgradient-based methods under interpolation. However, this lower bound holds only for subgradient-based algorithms. Smoothing techniques based on Moreau envelopes (Nesterov, 2005) can sometimes lead to acceleration for nonsmooth optimization, which may be further avenue to explore for obtaining an accelerated SSGD method.

REFERENCES

Zeyuan Allen-Zhu. Katyusha: The first direct acceleration of stochastic gradient methods. *Journal of Machine Learning Research*, 18:221:1–221:51, 2017.

Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. A convergence theory for deep learning via over-parameterization. In *Proceedings of ICML*, pp. 242–252, 2019.

Raef Bassily, Mikhail Belkin, and Siyuan Ma. On exponential convergence of sgd in non-convex over-parametrized learning. *arXiv e-prints*, art. arXiv:1811.02564, Nov 2018.

Amir Beck. *First-order methods in optimization*, volume 25. SIAM, 2017.

Amir Beck and Marc Teboulle. A conditional gradient method with linear rate of convergence for solving convex linear systems. *Mathematical Methods of Operations Research*, 59(2):235–247, 2004.

Amir Beck and Marc Teboulle. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM J. Imaging Sci.*, 2(1):183–202, 2009.

Amir Beck and Marc Teboulle. Smoothing and first order methods: A unified framework. *SIAM Journal on Optimization*, 22(2):557–580, 2012.

Dimitri P Bertsekas. Nonlinear programming. *Journal of the Operation Research Society*, 48(3), 1997.

Sébastien Bubeck. Convex optimization: Algorithms and complexity. *Foundation and Trends in Machine Learning*, 8(3-4):231–357, 2015.

Frank H Clarke. Generalized gradients of lipschitz functionals. *Advances in Mathematics*, 40(1): 52–67, 1981.

Frank H. Clarke. *Optimization and nonsmooth analysis*, volume 5. SIAM, 1990.

Aaron Defazio, Francis R. Bach, and Simon Lacoste-Julien. SAGA: A fast incremental gradient method with support for non-strongly convex composite objectives. In *Proceedings of NeurIPS*, pp. 1646–1654, 2014.

Xavier Glorot, Antoine Bordes, and Yoshua Bengio. Deep sparse rectifier neural networks. In *Proceedings of AISTATS*, pp. 315–323, 2011.

Ian Goodfellow, Yoshua Bengio, and Aaron Courville. *Deep Learning*. The MIT Press, 2016. ISBN 0262035618.

Robert M. Gower, Peter Richtárik, and Francis Bach. Stochastic quasi-gradient methods: Variance reduction via jacobian sketching. *arXiv e-prints*, art. arXiv:1805.02632, May 2018.

Prateek Jain, Sham M. Kakade, Rahul Kidambi, Praneeth Netrapalli, and Aaron Sidford. Accelerating stochastic gradient descent for least squares regression. In *Proceedings of COLT*, pp. 545–604, 2018.

Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition. In *Proceedings of ECML PKDD*, pp. 795–811, 2016.

Ahmed Khaled and Peter Richtárik. Better theory for sgd in the nonconvex world. *arXiv e-prints*, art. arXiv:2002.03329, Feb 2020.

Chaoyue Liu and Mikhail Belkin. Accelerating SGD with momentum for over-parameterized learning. In *Proceedings of ICLR*, 2020.

Nicolas Loizou, Sharan Vaswani, Issam Laradji, and Simon Lacoste-Julien. Stochastic polyak step-size for sgd: An adaptive learning rate for fast convergence. *arXiv e-prints*, art. arXiv:2002.10542, Feb 2020.

Siyuan Ma, Raef Bassily, and Mikhail Belkin. The power of interpolation: Understanding the effectiveness of SGD in modern over-parametrized learning. In *Proceedings of ICML*, pp. 3325–3334, 2018a.

Siyuan Ma, Raef Bassily, and Mikhail Belkin. The power of interpolation: Understanding the effectiveness of SGD in modern over-parametrized learning. In *Proceedings of ICML*, pp. 3331–3340, 2018b.

Si Yi Meng, Sharan Vaswani, Issam Hadj Laradji, Mark Schmidt, and Simon Lacoste-Julien. Fast and furious convergence: Stochastic second order methods under interpolation. In *Proceedings of AISTATS*, pp. 1375–1386, 2020.

Arkadi Nemirovski and David Yudin. *Problem Complexity and Method Efficiency in Optimization*. Wiley Interscience, 1983.

Yu. Nesterov. Gradient methods for minimizing composite objective function. CORE Discussion Papers 2007076, Université catholique de Louvain, Center for Operations Research and Econometrics (CORE), 2007.

Yurii Nesterov. Primal-dual subgradient methods for convex problems. *Mathematical programming*, 120(1):221–259, 2009.

Yurii Nesterov. *Introductory Lectures on Convex Optimization: A Basic Course*. Springer Publishing Company, Incorporated, 1 edition, 2014. ISBN 1461346916.

Yurii E. Nesterov. Smooth minimization of non-smooth functions. *Mathematical Programming*, 103 (1):127–152, 2005.

Neal Parikh and Stephen Boyd. Proximal algorithms. *Foundation and Trends in Optimization*, 1(3): 127–239, 2014.

Boris T Polyak. Introduction to optimization. translations series in mathematics and engineering. *Optimization Software*, 1987.

Xun Qian, Peter Richtárik, Robert M. Gower, Alibek Sailanbayev, Nicolas Loizou, and Egor Shulgin. SGD with arbitrary sampling: General analysis and improved rates. In *Proceedings of ICML*, pp. 5200–5209, 2019.

Mark Schmidt and Nicolas Le Roux. Fast convergence of stochastic gradient descent under a strong growth condition. *arXiv e-prints*, art. arXiv:1308.6370, Aug 2013.

Ohad Shamir. Can we find near-approximately-stationary points of nonsmooth nonconvex functions? *arXiv e-prints*, art. arXiv:2002.11962, Feb 2020.

Ohad Shamir and Tong Zhang. Stochastic gradient descent for non-smooth optimization: convergence results and optimal averaging schemes. In *Proceedings of ICML*, 2013.

Naum Z Shor. *Minimization methods for non-differentiable functions*. Springer Series in Computational Mathematics, Springer, 1984.

Sharan Vaswani, Francis Bach, and Mark Schmidt. Fast and faster convergence of SGD for over-parameterized models and an accelerated perceptron. In *Proceedings of AISTATS*, pp. 1195–1204, 2019a.

Sharan Vaswani, Aaron Mishkin, Issam H. Laradji, Mark Schmidt, Gauthier Gidel, and Simon Lacoste-Julien. Painless stochastic gradient: Interpolation, line-search, and convergence rates. In *Proceedings of NeurIPS*, 2019b.

Sharan Vaswani, Frederik Kunstner, Issam Laradji, Si Yi Meng, Mark Schmidt, and Simon Lacoste-Julien. Adaptive gradient methods converge faster with over-parameterization (and you can do a line-search). *arXiv e-prints*, art. arXiv:2006.06835, Jun 2020.

Xiaoxia Wu, Simon S. Du, and Rachel Ward. Global convergence of adaptive gradient methods for an over-parameterized neural network. *arXiv e-prints*, art. arXiv:1902.07111, Feb 2019.

Lin Xiao. Dual averaging methods for regularized stochastic learning and online optimization. *Journal of Machine Learning Research*, 11:2543–2596, 2010.

Jingzhao Zhang, Hongzhou Lin, Suvrit Sra, and Ali Jadbabaie. On complexity of finding stationary points of nonsmooth nonconvex functions. In *Proceedings of ICML*, 2020.

APPENDIX

A THE PROOF OF PROPOSITION 4.2

Proof of Proposition 4.2. Given $w_1, w_2 \in \mathbb{R}^d$, $\forall i \in [n]$,

$$
\begin{aligned}
&\|\partial f_i(w_2) - \partial f_i(w_1)\| \\
\leq\ & \|\ell'(h_i(w_2))\partial h_i(w_2) - \ell'(h_i(w_1))\partial h_i(w_1)\| \\
\leq\ & \|\ell'(h_i(w_2))\partial h_i(w_2) - \ell'(h_i(w_2))\partial h_i(w_1) + \ell'(h_i(w_2))\partial h_i(w_1) - \ell'(h_i(w_1))\partial h_i(w_1)\| \\
\leq\ & \|\ell'(h_i(w_2))\partial h_i(w_2) - \ell'(h_i(w_2))\partial h_i(w_1)\| + \|\ell'(h_i(w_2))\partial h_i(w_1) - \ell'(h_i(w_1))\partial h_i(w_1)\| \\
\leq\ & \|\ell'(h_i(w_2))(\partial h_i(w_2) - \partial h_i(w_1))\| + \|(\ell'(h_i(w_2)) - \ell'(h_i(w_1)))\partial h_i(w_1)\| \\
\leq\ & |\ell'(h_i(w_2))|\|\partial h_i(w_2) - \partial h_i(w_1)\| + |\ell'(h_i(w_2)) - \ell'(h_i(w_1))|\|\partial h_i(w_1)\| \\
\leq\ & 2L|\ell'(h_i(w_2))| + |h_i(w_2) - h_i(w_1)| \times L \\
\overset{(i)}{\leq}\ & 2L\sqrt{2f_i(w_2)} + L^2\|w_2 - w_1\|,
\end{aligned}
\tag{A.1}
$$

where (i) follows the same argument as the proof of Proposition 4.1:

$$
|\ell'(h_i(w_2))| = |\ell'(h_i(w_2)) - \ell'(0)| \leq \sqrt{2(\ell(h_i(w_2) - \ell(0))} = \sqrt{2f_i(w_2)}.
$$

Exchange w_1 and w_2, we can also get

$$
\|\partial f_i(w_2) - \partial f_i(w_1)\| \leq 2L\sqrt{2f_i(w_1)} + L^2\|w_2 - w_1\|.
\tag{A.2}
$$

Combining Equation (A.2) and Equation (A.1), we finished the proof for Equation (4.4).

Given $w_1, w_2 \in \mathbb{R}^d$, note that $f_i(w)$ is almost every differentiable by Rademacher's Theorem, thus

$$
\begin{aligned}
f_i(w_2) &= f_i(w_1) + \int_0^1 \langle \partial f_i(w_1 + \tau(w_2 - w_1)), w_2 - w_1\rangle d\tau \\
&= f_i(w_1) + \langle \partial f_i(w_1), w_2 - w_1\rangle + \int_0^1 \langle \partial f_i(w_1 + \tau(w_2 - w_1)) - \partial f(w_1), w_2 - w_1\rangle d\tau
\end{aligned}
\tag{A.3}
$$

Note that

$$
\begin{aligned}
&\int_0^1 \langle \partial f_i(w_1 + \tau(w_2 - w_1)) - \partial f_i(w_1), w_2 - w_1\rangle d\tau \\
\leq\ & \int_0^1 \|\partial f_i(w_1 + \tau(w_2 - w_1)) - \partial f_i(w_1)\|\|w_2 - w_1\|d\tau \\
\leq\ & \int_0^1 \left(2L\sqrt{2f_i(w_1)} + L^2\|\tau(w_2 - w_1)\|\right)\|w_2 - w_1\|d\tau \\
\leq\ & 2L\|w_2 - w_1\|\sqrt{2f_i(w_1)} + \frac{L^2}{2}\|w_2 - w_1\|^2.
\end{aligned}
\tag{A.4}
$$

Plug Equation (A.4) into Equation (A.3), we finished the proof for Equation (4.5). □