A Theoretical Analysis on Feature Learning in Neural Networks: Emergence from Inputs and Advantage over Fixed Features

Abstract

An important characteristic of neural networks is their ability to learn representations of the input data with effective features for prediction, which is believed to be a key factor to their superior empirical performance. To better understand the source and benefit of feature learning in neural networks, we consider learning problems motivated by practical data, where the labels are determined by a set of class relevant patterns and the inputs are generated from these along with some background patterns. We prove that neural networks trained by gradient descent can succeed on these problems. The success relies on the emergence and improvement of effective features, which are learned among exponentially many candidates efficiently by exploiting the data (in particular, the structure of the input distribution). In contrast, no linear models on data-independent features of polynomial sizes can learn to as good errors. Furthermore, if the specific input structure is removed, then no polynomial algorithm in the Statistical Query model can learn even weakly. These results provide theoretical evidence showing that feature learning in neural networks depends strongly on the input structure and leads to the superior performance. Our preliminary experimental results on synthetic and real data also provide positive support.

1 Introduction

Various empirical studies have shown that an important characteristic of neural networks is their feature learning ability, i.e., to learn a feature mapping for the inputs which allow accurate prediction (e.g., Zeiler & Fergus (2014); Girshick et al. (2014); Zhang et al. (2019); Manning et al. (2020)). This is widely believed to be a key factor to their remarkable success in many applications, in particular, an advantage over traditional machine learning methods. To understand their success, it is then crucial to understand the source and benefit of feature learning in neural networks. Empirical observations show that networks can learn neurons that correspond to different semantic patterns in the inputs (e.g., eyes, bird shapes, tires, etc. in images (Zeiler & Fergus, 2014; Girshick et al., 2014)). Moreover, recent progress (e.g., Caron et al. (2018); Chen et al. (2020b); He et al. (2020); Jing & Tian (2020)) shows that one can even learn a feature mapping using only unlabeled inputs and then learn an accurate predictor (usually a linear function) on it using labeled data. This further demonstrates the feature learning ability of neural networks and that these input distributions contain important information for learning useful features. These empirical observations strongly suggest that the structure of the input distribution is crucial for feature learning and feature learning is crucial for the strong performance. However, it is largely unclear how practical training methods (gradient descent or its variants) learn important patterns from the inputs and whether this is necessary for obtaining the superior performance, since the empirical studies do not exclude the possibility that some other training methods can achieve similar performance without feature learning or with feature learning that does not exploit the input structure. Rigorous theoretical investigations are thus needed for answering these fundamental questions: *How can effective features emerge from inputs in the training dynamics of gradient descent? Is learning features from inputs necessary for the superior performance?*

Compared to the abundant empirical evidence, the theoretical understanding still remains largely open. One line of work (e.g. Jacot et al. (2018); Li & Liang (2018); Du et al. (2019); Allen-Zhu et al.

(2019); Chizat et al. (2019); Cao et al. (2020) and many others) shows in certain regime, sufficiently overparameterized networks are approximately linear models, i.e., a linear function on the Neural Tangent Kernel (NTK). This falls into the traditional approach of linear models on fixed features, which also includes random features (Rahimi & Recht, 2008) and other kernel methods (Kamath et al., 2020). The kernel viewpoint thus does not explain feature learning in networks nor the advantage over fixed features. A recent line of work (e.g. Daniely & Malach (2020); Bai & Lee (2019); Ghorbani et al. (2020); Yehudai & Shamir (2019); Allen-Zhu & Li (2019; 2020a); Li et al. (2020); Malach et al. (2021) and others) shows examples where neural networks provably enjoy the advantage over fixed features, under different settings and assumptions. While providing insightful results on the separation of the two approaches, they have not investigated whether the input structure is a crucial factor for feature learning and thus the advantage. Also, most studies have not analyzed how gradient descent can learn important input patterns as effective features, or rely on strong assumptions like models or data atypical in practice (e.g., special networks, Gaussian data, etc).

Towards a more thorough understanding, we propose to analyze learning problems motivated by practical data, where the labels are determined by a set of class-relevant patterns and the inputs are generated from these along with some background patterns. We use comparison for our study: (1) by comparing network learning with fixed feature approaches on these problems, we analyze the emergence of effective features and demonstrate feature learning leads to the advantage over fixed features; (2) by comparing these problems to those with the input structure removed, we demonstrate that the input structure is crucial for feature learning and prediction performance.

More precisely, we obtain the following results. We first prove that two-layer networks trained by gradient descent can efficiently learn to small errors on these problems, and then prove that no linear models on fixed features of polynomial sizes can learn to as good errors. These two results thus establish the provable advantage of networks and implies that feature learning leads to this advantage. More importantly, our analysis reveals the dynamics of feature learning: the network first learns a rough approximation of the effective features, then improves them to get a set of good features, and finally learns an accurate classifier on these features. Notably, the improvement of the effective features in the second phase is needed for obtaining the provable advantage. The analysis also reveals the emergence and improvement of the effective features are by exploiting the data, and in particular, they rely on the input structure. To formalize this, we further prove the third result: if the specific input structure is removed and replaced by a uniform distribution, then no polynomial algorithm can even weakly learn in the Statistical Query (SQ) learning model, not to mention the advantage over fixed features. Since SQ learning includes essentially all known algorithms (in particular, mini-batch stochastic gradient descent used in practice), this implies that feature learning depends strongly on the input structure. Finally, we perform simulations on synthetic data to verify our results. We also perform experiments on real data and observe similar phenomena, which show that our analysis provides useful insights for the practical network learning.

Our analysis then provides theoretical support for the following principle: *feature learning in neural networks depends strongly on the input structure and leads to the superior performance.* In particular, our results make it explicit that learning features from the input structure is crucial for the superior performance. This suggests that input-distribution-free analysis (e.g., traditional PAC learning) may not be able to explain the practical success, and advocates an emphasis of the input structure in the analysis. While these results are for our proposed problem setting and network learning in practice can be more complicated, the insights obtained match existing empirical observations and are supported by our experiments. The compelling evidence hopefully can attract more attention to further studies on modeling the input structure and analyzing feature learning.

2 RELATED WORK

Neural Tangent Kernel (NTK) and Linearization of Neural Networks. One line of work explains the success of sufficiently over-parameterized neural network by connecting them to linear methods like NTK (e.g. Jacot et al. (2018); Li & Liang (2018); Matthews et al. (2018); Lee et al. (2019); Novak et al. (2019); Yang (2019); Du et al. (2019); Allen-Zhu et al. (2019); Ji & Telgarsky (2019); Cao et al. (2020); Geiger et al. (2020); Chizat et al. (2019) and more). Though their approaches are different, they all base on the observation that when the neural network is sufficiently large, the weight of the neurons stays closed to the initialization during the training. In this situation, training with gradient

descent appears no different from solving a kernel method problem. This view is typically referred to as the NTK regime, or lazy training, or linearization. However, neural networks used in practice are usually not large enough to enter this regime, and the neurons are frequently observed to traverse instead of staying close to their initialization during the training process. Furthermore, in this regime, network learning is essentially the traditional approach of linear methods over fixed features, which cannot establish or explain feature learning and the advantage of network learning.

Advantage of Neural Networks over Linear Models on Fixed Features. Since the benign network learning results via gradient descent are not well explained by the NTK view, a recent line of work has turned to show learning settings where neural networks provably have advantage over linear models on fixed features (e.g. Daniely & Malach (2020); Refinetti et al. (2021); Malach et al. (2021); Dou & Liang (2020); Bai & Lee (2019); Ghorbani et al. (2020); Allen-Zhu & Li (2019); see the nice summary in Malach et al. (2021)). While formally establishing the advantages of network learning over fixed feature approaches, they have not answered the two fundamental questions this work focuses on; in particular, they have not studied whether the input structure is a crucial factor for feature learning and thus the advantage. For example, Ghorbani et al. (2020) show the advantage of networks in approximation power and Dou & Liang (2020) show their statistical advantage, but they do not consider the learning dynamics (i.e., how the training method obtains the good network). Allen-Zhu & Li (2019) prove the advantage of the networks for PAC learning with labels given by a depth-2 ResNet and Allen-Zhu & Li (2020a) prove for Gaussian inputs with labels given by a multiple-layer network, while neither considers the influence of the input structure on feature learning or the advantage. Daniely & Malach (2020) prove the advantage of the networks for the task of learning sparse parities on specifically designed input distributions that help the gradient descent learn effective features for prediction, and Malach et al. (2021) consider similar learning problems but with specifically designed differentiable models. On the other hand, neither study explores if the input structure is needed for the learning, and the data distributions and models are also atypical in practice, while our setting better connects to the practice. More technical discussions can be found in Appendix A.

There are also other theoretical studies on feature learning in networks (e.g. Yehudai & Ohad (2020); Zhou et al. (2021); Diakonikolas et al. (2020); Frei et al. (2020)), which however do not directly relate feature learning to the input structure or the advantage.

3   PROBLEM SETUP

To motivate our setup, consider images with various kinds of patterns like lines and circles. Some patterns are relevant for the labels while others are not; if the image contains a sufficient number of the former, then we are confident that the image belongs to a certain class. Dictionary learning, or sparse coding (e.g. Olshausen & Field (1997); Vinje & Gallant (2000); Blei et al. (2003); Yang et al. (2009)) is a classic model of such data. We thus model the patterns as a dictionary, generate a hidden vector indicating the presence of the patterns, and generate the input and label from this vector.

Let $\mathcal{X} = \mathbb{R}^d$ be the input space, and $\mathcal{Y} = \{\pm 1\}$ be the label space. Suppose $M \in \mathbb{R}^{d \times D}$ is an unknown dictionary with D columns that can be regarded as patterns. For simplicity, assume M is orthonormal. Let $\tilde{\phi} \in \{0, 1\}^D$ be a hidden vector that indicates the presence of each pattern. Let $A \subseteq [D]$ be a subset of size k corresponding to the class relevant patterns. Then the input is generated by $M\tilde{\phi}$, and the label can be any binary function on the number of class relevant patterns. More precisely, let $P \subseteq [k]$. We first sample $\tilde{\phi}$ from a distribution $\mathcal{D}_{\tilde{\phi}}$, and then generate the input $\tilde{x}$ and the class label y from $\tilde{\phi}, A, P$:

$$\tilde{\phi} \sim \mathcal{D}_{\tilde{\phi}}, \quad \tilde{x} = M\tilde{\phi}, \quad y = \begin{cases} +1, & \text{if } \sum_{i \in A} \tilde{\phi}_i \in P, \\ -1, & \text{otherwise.} \end{cases} \quad (1)$$

Learning with Input Structure. We allow quite general $\mathcal{D}_{\tilde{\phi}}$ with the following assumptions:

(A0) The class probabilities are balanced: $\Pr[\sum_{i \in A} \tilde{\phi}_i \in P] = 1/2$.

(A1) The patterns in A are correlated with the labels: for any $i \in A$, $\gamma = \mathbb{E}[y\tilde{\phi}_i] - \mathbb{E}[y]\mathbb{E}[\tilde{\phi}_i] > 0$.

(A2) Each pattern outside A is independent of all other patterns and identically distributed. Let $p_{\mathrm{o}} := \Pr[\tilde{\phi}_i = 1] \leq 1/2$ denote the probability they appear.

Let $\mathcal{D}(A, P, \mathcal{D}_{\tilde{\phi}})$ denote the distribution on $(\tilde{x}, y)$ corresponding to some A, P, and $\mathcal{D}_{\tilde{\phi}}$. Given parameters $\Xi = (d, D, k, \gamma, p_{\text{o}})$, the family $\mathcal{F}_{\Xi}$ of distributions for learning is the set of all $\mathcal{D}(A, P, \mathcal{D}_{\tilde{\phi}})$ with $A \subseteq [D]$, $P \subseteq [k]$, and $\mathcal{D}_{\tilde{\phi}}$ satisfying the above assumptions. Appendix F presents results for more general settings (e.g., incoherent dictionary, unbalanced classes, etc.). Our learning problems here already include some interesting special cases:

Example 1. Suppose k is odd, and $P = \{i \in [k] : i > k/2\}$ for some threshold, i.e., we will set the label $y = +1$ when more than a half of the relevant patterns are presented in the input.

Example 2. Suppose k is odd, and let $P = \{i \in [k] : i \text{ is odd}\}$, i.e., the labels are given by the parity function on $\tilde{\phi}_j (j \in A)$. This is useful to prove our lower bounds via the properties of parities.

Learning Without Input Structure. For comparison, we also consider learning problems without input structure. The data is generated as above but with different distributions $\mathcal{D}_{\tilde{\phi}}$:

(**A1'**) The patterns are independent, and for any $i \in [D], \Pr[\tilde{\phi}_i = 1] = 1/2$.

Given parameters $\Xi_0 = (d, D, k)$, the family $\mathcal{F}_{\Xi_0}$ of distributions without input structure is the set of all the distributions with $A \subseteq [D]$, $P \subseteq [k]$ and $\mathcal{D}_{\tilde{\phi}}$ satisfying the above assumptions.

3.1 Neural Network Learning

Networks. We consider training a two-layer network via gradient descent on the data distribution:[1]

$$g(x) = \sum_{i=1}^{2m} a_i \sigma(\langle w_i, x \rangle + b_i). \tag{2}$$

where $w_i \in \mathbb{R}^d, b_i, a_i \in \mathbb{R}$, and $\sigma(z) = \min(1, \max(z, 0))$ is the truncated rectified linear unit (ReLU) activation function. Let $\theta = \{w_i, b_i, a_i\}_{i=1}^{2m}$ denote all the parameters, and let superscript (t) denote the time step, e.g., $g^{(t)}$ denote the network at time step t with $\theta^{(t)} = \{w_i^{(t)}, b_i^{(t)}, a_i^{(t)}\}$.

Loss Function. Similar to typical practice, we will normalize the data for learning: first compute $x = (\tilde{x} - \mathbb{E}[\tilde{x}])/\tilde{\sigma}$ where $\tilde{\sigma}^2 = \sum_{i=1}^{d}(\tilde{x}_i - \mathbb{E}[\tilde{x}_i])^2$ is the variance of the data, and then train on (x, y). This is equivalent to setting $\phi = (\tilde{\phi} - \mathbb{E}[\tilde{\phi}])/\tilde{\sigma}$ and generating $x = M\phi$. For $(\tilde{x}, y)$ from $\mathcal{D}$ and the normalized (x, y), we will simply say $(x, y) \sim \mathcal{D}$.

For the training, we consider the hinge-loss $\ell(y, \hat{y}) = \max\{1 - y\hat{y}, 0\}$. We will inject some noise ξ to the neurons for the convenience of the analysis. (This can be viewed as using a smoothed version of the activation $\tilde{\sigma}(z) = \mathbb{E}_{\xi}\sigma(z + \xi)$ similar to those in existing studies like Allen-Zhu & Li (2020b); Malach et al. (2021). See Section 5 for more explanations.) Formally, the loss is:

$$L_{\mathcal{D}}(g; \sigma_{\xi}) = \mathbb{E}_{(x,y)}[\ell(y, g(x; \xi))], \text{ where } g(x; \xi) = \sum_{i=1}^{2m} a_i \mathbb{E}_{\xi}[\sigma(\langle w_i, x \rangle + b_i + \xi_i)] \tag{3}$$

where $\xi \sim \mathcal{N}(0, \sigma_{\xi}^2 I_{m \times m})$ are independent Gaussian noise. Let $L_{\mathcal{D}}(g)$ denote the typical hinge-loss without noise. We also consider ℓ_2 regularization: $R(g; \lambda_a, \lambda_w) = \sum_{i=1}^{2m} \lambda_a |a_i|^2 + \lambda_w \|w_i\|_2^2$ with regularization coefficients λ_a, λ_w.

Training Process. We first perform an unbiased initialization: for every $i \in [m]$, initialize $w_i^{(0)} \sim \mathcal{N}(0, \sigma_w^2 I_{d \times d})$ with $\sigma_w = 1/k$, $b_i^{(0)} \sim \mathcal{N}(0, \sigma_b^2)$ with $\sigma_b = 1/k^2$, $a_i^{(0)} \sim \mathcal{N}(0, \sigma_a^2)$ with $\sigma_a = \tilde{\sigma}^2/(\gamma k^2)$, and then set $w_{m+i}^{(0)} = w_i^{(0)}, b_{m+i}^{(0)} = b_i^{(0)}, a_{m+i}^{(0)} = -a_i^{(0)}$. We then do gradient updates:

$$\theta^{(t)} = \theta^{(t-1)} - \eta^{(t)} \nabla_{\theta} \left(L_{\mathcal{D}}(g^{(t-1)}; \sigma_{\xi}^{(t)}) + R(g^{(t-1)}; \lambda_a^{(t)}, \lambda_w^{(t)}) \right), \text{ for } t = 1, 2, \ldots, T, \tag{4}$$

for some choice of the hyperparameters $\eta^{(t)}, \lambda_a^{(t)}, \lambda_w^{(t)}, \sigma_{\xi}^{(t)}$, and T.

---

[1] The analysis can be easily generalized to τ-approximate gradients, e.g., by the typical mini-batch SGD.

4 MAIN RESULTS

Provable Guarantee for Neural Networks. The network learning has the following guarantee:

Theorem 1. *For any $\delta, \epsilon \in (0, 1)$, if $p_o = \Omega(k^2/D)$, $k = \Omega\left(\log^2\left(Dm/(\delta\gamma)\right)\right)$, and $m \geq \max\{\Omega(k^{12}/\epsilon^{3/2}), D\}$, then with properly set hyperparameters, we have for any $\mathcal{D} \in \mathcal{F}_\Xi$, with probability at least $1 - \delta$, there exists $t \in [T]$ such that $\Pr[\text{sign}(g^{(t)}(x)) \neq y] \leq L_\mathcal{D}(g^{(t)}) \leq \epsilon$.*

The theorem shows that for a wide range of the background pattern probability p_o and the number of class relevant patterns k, the network trained by gradient descent can obtain a small 0-1 classification error with high probability. More importantly, the analysis shows the success comes from feature learning. In the early stage, the network learns and improves the neuron weights such that on the corresponding features (i.e., the neurons' outputs) there exists an accurate classifier; afterwards it can learn such an accurate classifier on these features. Next section will provide a detailed discussion on the feature learning and also the choice of the hyperparameters.

Lower Bound for Fixed Features. Empirical observations and Theorem 1 do not exclude the possibility that some learning methods without feature learning can achieve similar performance. We thus prove a lower bound for the fixed feature approach, i.e., linear models on on a set of data-independent features. Such models include random feature approaches (Rahimi & Recht, 2008) and approximate various kernel methods (see the discussion in Kamath et al. (2020)).

Theorem 2. *Suppose Ψ is a data-independent feature mapping of dimension N with bounded features, i.e., $\Psi : \mathcal{X} \to [-1, 1]^N$. For $B > 0$, the family of linear models on Ψ with bounded norm B is $\mathcal{H}_B = \{h(\tilde{x}) : h(\tilde{x}) = \langle \Psi(\tilde{x}), w \rangle, \|w\|_2 \leq B\}$. If $3 < k \leq D/16$ and k is odd, then there exists $\mathcal{D} \in \mathcal{F}_\Xi$ such that all $h \in \mathcal{H}_B$ have hinge-loss at least $p_o \left(1 - \frac{\sqrt{2N}B}{2^k}\right)$.*

The theorem shows that using *fixed* features, one cannot get loss nontrivially smaller than p_o unless with exponentially large models. In contrast, viewing the hidden neurons as *learned* features with $\Psi_i = \sigma(\langle w_i, x \rangle + b_i)$, network learning can achieve loss ϵ for any $\epsilon \in (0, 1)$ with models of polynomial sizes. This comparison then shows the data-dependent feature learning ability of the networks can help overcome the exponential requirement.

Lower Bound for Without Input Structure. Existing results do not exclude the possibility that some learning methods without exploiting the input structure can achieve strong performance. To show the necessity of the input structure, we consider learning $\mathcal{F}_{\Xi_0}$ with input structure removed.

We obtain a lower bound for such learning problems in the classic Statistical Query (SQ) model (Kearns, 1998). In this model, the algorithm can only receive information about the data through statistical queries. A statistical query is specified by some polynomially-computable property predicate Q of labeled instances and a tolerance parameter $\tau \in [0, 1]$. For a query (Q, τ), the algorithm receives a response $\hat{P}_Q \in [P_Q - \tau, P_Q + \tau]$, where $P_Q = \Pr[Q(x, y) \text{ is true}]$. Notice that a query can be simulated using the average of roughly $O(1/\tau^2)$ random data samples with high probability. The SQ model captures almost all common learning algorithms (except Gaussian elimination) including the commonly used mini-batch SGD, and thus is suitable for our purpose.

Theorem 3. *For any algorithm in the Statistical Query model that can learn over $\mathcal{F}_{\Xi_0}$ to classification error less than $\frac{1}{2} - \frac{1}{\binom{D}{k}^3}$, either the number of queries or $1/\tau$ must be at least $\frac{1}{2}\left(\frac{D}{k}\right)^{1/3}$.*

The theorem shows that without the input structure, polynomial algorithms in the SQ model cannot get a classification error nontrivially smaller than random guessing. In contrast, with the input structure, network learning can achieve small classification errors with a polynomial algorithm. This comparison between the two problem settings then shows that the input structure is crucial for network learning, in particular, for achieving the advantage over fixed feature models.

5 PROOF SKETCHES

Here we provide the sketch of our analysis, focusing on the key intuition and discussing some interesting implications. The complete proofs are included in Appendix B-D.

5.1 PROVABLE GUARANTEES OF NEURAL NETWORKS

Overall Intuition. We first show that there is a two-layer network that can represent the target labeling function, whose neurons can be viewed as the "ground-truth" features to be learned. We then show that after the first gradient step, the hidden neurons of the trained network become close to the ground-truth: their weights contain large components along the class relevant patterns but small along the background patterns. We further show that in the second gradient step, these features get improved: the "signal-noise" ratio between the components for class relevant patterns and those for the background ones becomes larger, giving a set of good features. Finally, we show that the remaining steps learn an accurate classifier on these features.

Existence of A Good Network. We show that there is a two-layer network that can fit the labels.

Lemma 4. *For any $\mathcal{D} \in \mathcal{F}_\Xi$, there exists a network $g^*(x) = \sum_{i=1}^n a_i^* \sigma(\langle w_i^*, x \rangle + b_i^*)$ with $y = g^*(x)$ for any $(x, y) \sim \mathcal{D}$. Furthermore, the number of neurons $n = 3(k+1)$, $|a_i^*| \leq 32k, 1/(32k) \leq |b_i^*| \leq 1/2$, $w_i^* = \tilde{\sigma} \sum_{j \in A} M_j/(4k)$, and $|\langle w_i^*, x \rangle + b_i^*| \leq 1$ for any $i \in [n]$ and $(x, y) \sim \mathcal{D}$.*

In particular, the weights of the neurons are proportional to $\sum_{j \in A} M_j$, the sum of the class relevant patterns. We thus focus on analyzing how the network learns such neuron weights.

Feature Emergence in the First Gradient Step. The gradient for w_i (ignoring the noise) is:

$$\frac{\partial L_{\mathcal{D}}(g)}{\partial w_i} = -a_i \mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ y \mathbb{I}[yg(x) \leq 1] \sigma'[\langle w_i, x \rangle + b_i] x \right\} = -a_i \mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ yx \sigma'[\langle w_i, x \rangle + b_i] \right\}$$

where the last step is due to $g(x) = 0$ by the unbiased initialization. Let $q_j = \langle M_j, w_i \rangle$ denote the component along the direction of the pattern M_j. Then the component of the gradient on M_j is:

$$\left\langle M_j, \frac{\partial}{\partial w_i} L_{\mathcal{D}}(g) \right\rangle = -a_i \mathbb{E} \left\{ y \phi_j \sigma'[\langle w_i, x \rangle + b_i] \right\} = -a_i \mathbb{E} \left\{ y \phi_j \sigma' \left[\sum_{\ell \in [D]} \phi_\ell q_\ell + b_i \right] \right\}.$$

The key intuition is that with the randomness of ϕ_ℓ (and potentially that of the injected noise ξ), the random variable under σ' is not significantly affected by a small subset of $\phi_\ell q_\ell$. For example, for class relevant patterns $j \in A$, let $\mathbb{I}_{[D]} := \sigma' \left[\sum_{\ell \in [D]} \phi_\ell q_\ell + b_i \right]$ and $\mathbb{I}_{-A} := \sigma' \left[\sum_{\ell \notin A} \phi_\ell q_\ell + b_i \right]$. We have $\mathbb{I}_{[D]} \approx \mathbb{I}_{-A}$ and thus:

$$\left\langle M_j, \frac{\partial}{\partial w_i} L_{\mathcal{D}}(g) \right\rangle \propto \mathbb{E} \left\{ y \phi_j \mathbb{I}_{[D]} \right\} \approx \mathbb{E} \left\{ y \phi_j \mathbb{I}_{-A} \right\} = \mathbb{E} \left\{ y \phi_j \right\} \mathbb{E} [\mathbb{I}_{-A}] = \frac{\gamma}{\tilde{\sigma}} \mathbb{E} [\mathbb{I}_{-A}]$$

since y only depends on $\phi_j (j \in A)$. Then the gradient has a nontrivial component along the pattern. Similarly, for background patterns $j \notin A$, the component of the gradient along M_j is close to 0.

Lemma 5 (Informal). *Assume p_o, k as in Theorem 1 and $\sigma_\xi^{(1)} < 1/k$, then with high probability $\frac{\partial}{\partial w_i} L_{\mathcal{D}}(g^{(0)}; \sigma_\xi^{(1)}) = -a_i^{(0)} \sum_{j=1}^D M_j T_j$ where for a small ϵ_e:*

- *if $j \in A$, then $|T_j - \beta \gamma / \tilde{\sigma}| \leq O(\epsilon_e / \tilde{\sigma})$ with $\beta \in [\Omega(1), 1]$;*

- *if $j \notin A$, then $|T_j| \leq O(\sigma_\phi^2 \epsilon_e \tilde{\sigma})$.*

By setting $\lambda_w^{(1)} = 1/(2\eta^{(1)})$, we have $w_i^{(1)} = \eta^{(1)} a_i^{(0)} \sum_{j=1}^D M_j T_j \approx \eta^{(1)} a_i^{(0)} \frac{\beta \gamma}{\tilde{\sigma}} \sum_{j \in [D]} M_j$. For small p_o, e.g., $p_o = \tilde{O}(k^2/D)$, these neurons can already allow accurate prediction. However, for such small p_o, we cannot show a provable advantage of networks over fixed features. Furthermore, for larger p_o meaning a significant number of background patterns in the input, the approximation error terms $T_j (j \notin A)$ together can overwhelm the signals $T_j (j \in A)$ and lead to bad prediction, even though each term is small. Fortunately, we will show that the second gradient step can improve the weights by decreasing the ratio between $T_j (j \notin A)$ and $T_j (j \in A)$.

Feature Improvement in the Second Gradient Step. We note that by setting a small $\eta^{(1)}$, after the update we still have $yg(x; \xi) < 1$ for most $(x, y) \sim \mathcal{D}$ and thus the gradient in the second step is:

$$\frac{\partial}{\partial w_i} L_{\mathcal{D}}(g; \sigma_\xi) \approx -a_i \mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ yx \mathbb{E}_\xi \sigma'[\langle w_i, x \rangle + b_i + \xi_i] \right\}.$$

We can then follow the intuition for the first step again. For $j \in A$, the component $\langle M_j, \frac{\partial}{\partial w_i} L_{\mathcal{D}}(g) \rangle$ is roughly proportional to $\frac{\gamma}{\tilde{\sigma}} \mathbb{E}[\mathbb{I}_{-A,\xi}]$ where $\mathbb{I}_{-A,\xi} := \sigma' \left[\sum_{\ell \notin A} \phi_\ell q_\ell + b_i + \xi_i \right]$. While $\phi_\ell q_\ell$ may not have large enough variance, the injected noise ξ_i makes sure that a nontrivial amount of data activate the neuron.[2] Then $\mathbb{I}_{-A,\xi} \neq 0$, leading to a nontrivial component along M_j, similar to the first step. On the other hand, for $j \notin A$, the approximation error term T_j depends on how well $\sigma' \left[\sum_{\ell \notin A, \ell \neq j} \phi_\ell q_\ell + b_i + \xi_i \right]$ approximates $\sigma' \left[\sum_{\ell \in [D]} \phi_\ell q_\ell + b_i + \xi_i \right]$. Since the q_ℓ's (the weight's component along M_ℓ) in the second step are small compared to those in the first step, we can then get a small error term T_j.

Lemma 6 (Informal). *Assume p_o, k as in Theorem 1, $\eta^{(1)}$ is sufficiently small, and $\sigma_\xi^{(2)} = 1/k^{3/2}$, then with high probability $\frac{\partial}{\partial w_i} L_{\mathcal{D}}(g^{(1)}; \sigma_\xi^{(2)}) = -a_i^{(1)} \sum_{j=1}^{D} M_j T_j$ where for a small $\epsilon_{e2} \ll \epsilon_e$:*

- *if $j \in A$, then $T_j \approx \beta \gamma / \tilde{\sigma}$ with $\beta \in [\Omega(1), 1]$;*

- *if $j \notin A$, then $|T_j| \leq O(\sigma_\phi^2 \epsilon_{e2} \tilde{\sigma}) + \frac{1}{\tilde{\sigma}} \exp(-\Theta(p_o D))$.*

So roughly the ratio between $T_j (j \notin A)$ over $T_j (j \in A)$ improves from $O(\sigma_\phi^2 \tilde{\sigma}^2 \epsilon_e / \gamma)$ after the first step to $O(\sigma_\phi^2 \tilde{\sigma}^2 \epsilon_{e2} / \gamma)$ after the second step, giving good features allowing accurate prediction.

Remark. The analysis can be carried out for more gradient steps following similar intuition, while we analyze two steps for simplicity.

Remark. Readers may notice that the network can be overparameterized. With sufficient overparameterization and proper initialization and step sizes, network learning becomes approximately NTK. However, here our learning scheme allows going beyond this kernel regime: we use aggressive gradient updates $\lambda_w^{(t)} = 1/(2\eta^{(t)})$ in the first two steps, completely forgetting the old weights to learn effective features. Using proper initialization and aggressive updates early to escape the kernel regime has been studied in existing work (e.g., Woodworth et al. (2020); Li et al. (2019)). Our result thus adds another concrete example.

Classifier Learning Stage. Given the learned features, we are then ready to show the remaining gradient steps can learn accurate classifiers. Intuitively, with small hyperparameter values ($\eta^{(t)} = \frac{k^2}{T m^{1/3}}, \lambda_a^{(t)} = \lambda_w^{(t)} \leq \frac{k^3}{\tilde{\sigma} m^{1/3}}, \sigma_\xi^{(t)} = 0$ for $2 < t \leq T = m^{4/3}$), the first layer's weights do not change too much and thus the learning is similar to convex learning using the learned features. Formally, our proof uses the online convex optimization technique in Daniely & Malach (2020).

5.2 Lower Bounds

The lower bounds are based on the following observation: our problem setup is general enough to include learning sparse parity functions. Consider an odd k, and let $P = \{i \in [k] : i \text{ is odd}\}$. Then y is given by $\Pi_A(z) := \prod_{j \in A} z_j$ for $z_j = 2\tilde{\phi}_j - 1$, i.e., the parity function on $z_j (j \in A)$. Then known results for learning parity functions can be applied to prove our lower bounds.

Lower Bound for Fixed Features. We show that $\mathcal{F}_{\boxminus}$ contains learning problems that consist of a mixture of two distributions with weights p_o and $1 - p_o$ respectively, where in the first distribution $\mathcal{D}_A^{(1)}$, $\tilde{x}$ is given by the uniform distribution over $\tilde{\phi}$ and the label y is given by the parity function on A. On such $\mathcal{D}_A^{(1)}$, Daniely & Malach (2020) shows that exponentially large models over fixed features is needed to get nontrivial loss. Intuitively, there are exponentially many labeling functions Π_A that are uncorrelated (i.e., "orthogonal" to each other): $\mathbb{E}[\Pi_{A_1} \Pi_{A_2}] = 0$ for any A_1 and A_2. Note that the best approximation of Π_A by a fixed set of features Ψ_i's is its projection on the linear span of the features. Then with polynomial-size models, there always exists some Π_A far from the linear span.

---

[2]Equivalently, the network uses $\tilde{\sigma}(z) = \mathbb{E}_\xi \sigma(z + \xi)$, a Gaussian smoothed version of σ, and the smoothing allows z slightly outside the activated region of σ to generate gradient for the learning. Empirically it is not needed since typically sufficient data can activate the neurons. One potential reason is that the data have their own noise to achieve a similar effect (a remote analog being noisy gradients can help the optimization). Further analysis on such an effect is left for future work.

Remark. It is instructive to compare to network learning, which finds the effective weights $\sum_{j \in A} M_j$ among the exponentially many candidates corresponding to different A's. This can be done efficiently by exploiting the data since the gradient is roughly proportional to $\mathbb{E}\{yx\} = \sum_{j \in A} M_j$. The network then learns *data-dependent* features on which polynomial size linear models can achieve small loss.

Lower Bound for Learning without Input Structure. Clearly, $\mathcal{F}_{\Xi_0}$ contains the distributions $\mathcal{D}_A^{(1)}$ described above. The lower bound then follows from classic SQ learning results (Blum et al., 1994).

Remark. The SQ lower bound analysis does not apply to $\mathcal{F}_\Xi$, because in $\mathcal{F}_\Xi$ the input distribution is related the labeling function. This allows networks to learn with polynomial time/sample. While both the labeling function and the input distribution affect the learning, few existing studies explicitly point out the importance of the input structure. We thus emphasize the input structure is crucial for networks to learn effective features and achieve superior performance.

6 EXPERIMENTS

Our experiments mainly focus on feature learning and the effect of the input structure. We first perform simulations on our learning problems to (1) verify our main theorems on the benefit of feature learning and the effect of input structure; (2) verify our analysis of feature learning in networks. We then check if our insights carry over to real data: (3) whether similar feature learning is presented in real network/data; (4) whether damaging the input structure lowers the performance. The results are consistent with our analysis and provide positive support for the theory. Below we present part of the results and include the complete experimental details and results in Appendix E.

Simulation: Verification of the Main Results. We generate data according to our problem setup, with $d = 500, D = 100, k = 5, p_{\mathrm{o}} = 1/2$, a randomly sampled A, and labels given by the parity function. We then train a two-layer network with $m = 300$ following our learning process, and for comparison, we also use two fixed feature methods (the NTK and random feature methods based on the same network). Finally, we also use these three methods on the data distribution with the input structure removed (i.e., $\mathcal{F}_{\Xi_0}$ in Theorem 3).

Figure 1 shows that the results are consistent with our results. Network learning gets high test accuracy while the two fixed feature methods get significantly lower accuracy. Furthermore, when the input structure is removed, all three methods get test accuracy similar to random guessing.

Figure 1: Test accuracy on simulated data with or without input structure.

Simulation: Feature Learning in Networks. We compute the cosine similarities between the weights w_i's and visualize them by Multidimensional Scaling. (Recall that our analysis is on the *directions* of the weights without considering their *scaling*, and thus it is important to choose cosine similarity rather than say the typical Euclidean distance.) Figure 2 shows that the results are as predicted by our analysis. After the first gradient step, some weights begin to cluster around the ground-truth $\sum_{j \in A} M_j$ (or $-\sum_{j \in A} M_j$ due to the a_i in the gradient update which can be positive or negative). After the second step, the weights get improved and well-aligned with the ground-truth (with cosine similarities > 0.99). Furthermore, if a classifier is trained on the features after the first step, the test accuracy is about 52%; if the same is done after the second step, the test accuracy is about 100%. This demonstrates while some effective features emerge in the first step, they need to be improved in the second step to get accurate prediction.

Real Data: Feature Learning in Networks. We perform experiments on MNIST (LeCun et al., 1998; Deng, 2012), CIFAR10 (Krizhevsky, 2012), and SVHN (Netzer et al., 2011). On MNIST, we train a two-layer network with $m = 50$ on the subset with labels 0/1 and visualize the neurons' weights as in the simulation. Figure 3 shows a similar feature learning phenomenon: effective features emerge after a few steps and then get improved to form two clusters. Similar results are observed on other datasets. These suggest the insights obtained in our analysis are also applicable to the real data.

Figure 2: Visualization of the weights w_i's after initialization/one gradient step/two steps in network learning on the synthetic data. The red star denotes the ground-truth $\sum_{j \in A} M_j$; the orange star is $-\sum_{j \in A} M_j$. The red/orange dots are the weights closest to the red/orange star, respectively.

Figure 3: Visualization of the neurons' weights in a two-layer network trained on the subset of MNIST data with label 0/1. The weights gradually form two clusters.

Figure 4: Test accuracy at different steps for an equal mixture of Gaussian inputs with data: (a) MNIST, (b) CIFAR10, (c) SVHN.

Real Data: The Effect of Input Structure. Since we cannot directly manipulate the input distribution of real data, we perform controlled experiments by injecting different inputs. For labeled dataset $\mathcal{L}$ and injected input $\mathcal{U}$, we first train a teacher network fitting $\mathcal{L}$, then use the teacher network to give labels on a mixture of inputs from $\mathcal{L}$ and $\mathcal{U}$, and finally train a student network on this new dataset $\mathcal{M}$ consisting of the mixed inputs and the teacher network's labels. Checking the student' performance on different parts of $\mathcal{M}$ and comparing to those by directly training the student on the original data $\mathcal{L}$ can reveal the impact of chaning the input structure. We use MNIST, CIFAR10, or SVHN as $\mathcal{L}$, and use Gaussian or images in Tiny ImageNet (Le & Yang, 2015) as $\mathcal{U}$. The networks for MNIST are two-layer with $m = 9$, and those for CIFAR10/SVHN are ResNet-18 convolutional neural networks (He et al., 2016).

Figure 4 shows the results on an equal mixture of data and Gaussian. It presents the test accuracy of the student on the original data part, the Gaussian part, and the whole mixture. For example, on CIFAR10, the network learns well over the CIFAR10 part (with accuracy similar to directly training on the original data) but learns slower with worse accuracy on the Gaussian part. Furthermore, the accuracy on the whole mixture is lower than that of training on the original CIFAR10. This shows that the input structure indeed has a significant impact on the learning. While MNIST+Gaussian shows a less significant trend (possibly because the tasks are simpler), the other datasets show similar significant trends as CIFAR10+Gaussian (the results using Tiny ImageNet are in the appendix).

7 ETHICS STATEMENT

Our paper is mostly theoretical in nature and thus we foresee no immediate negative ethical impact. We are of the opinion that our theoretical framework may lead to better understanding and inspire development of improved network learning methods, which may have a positive impact in practice. In addition to the theoretical machine learning community, we perceive that our conceptual message that the input structure is crucial for the network learning's performance can be beneficial to engineering-inclined machine learning researchers.

8 REPRODUCIBILITY STATEMENT

For theoretical results in the Section 4, a complete proof is provided in the Appendix Section B-D. The theoretical results and complete proofs for a setting more general than that in the main text are provided in the Appendix Section F. For experiments in the Section 6, complete details and experimental results are provided in the Appendix Section E. The source code with explanations and comments is provided in the supplementary material.

REFERENCES

Zeyuan Allen-Zhu and Yuanzhi Li. What can resnet learn efficiently, going beyond kernels? In *Advances in Neural Information Processing Systems*, 2019.

Zeyuan Allen-Zhu and Yuanzhi Li. Backward feature correction: How deep learning performs deep learning. *arXiv preprint arXiv:2001.04413*, 2020a.

Zeyuan Allen-Zhu and Yuanzhi Li. Feature purification: How adversarial training performs robust deep learning. *arXiv preprint arXiv:2005.10190*, 2020b.

Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. A convergence theory for deep learning via over-parameterization. In *International Conference on Machine Learning*, 2019.

Yu Bai and Jason D Lee. Beyond linearization: On quadratic and higher-order approximation of wide neural networks. In *International Conference on Learning Representations*, 2019.

Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine-learning practice and the classical bias–variance trade-off. *Proceedings of the National Academy of Sciences*, 116(32):15849–15854, 2019.

David M Blei, Andrew Y Ng, and Michael I Jordan. Latent dirichlet allocation. *the Journal of machine Learning research*, 3:993–1022, 2003.

Avrim Blum, Merrick Furst, Jeffrey Jackson, Michael Kearns, Yishay Mansour, and Steven Rudich. Weakly learning dnf and characterizing statistical query learning using fourier analysis. In *Proceedings of the twenty-sixth annual ACM symposium on Theory of computing*, pp. 253–262, 1994.

Yuan Cao, Zhiying Fang, Yue Wu, Ding-Xuan Zhou, and Quanquan Gu. Towards understanding the spectral bias of deep learning, 2020.

Mathilde Caron, Piotr Bojanowski, Armand Joulin, and Matthijs Douze. Deep clustering for unsupervised learning of visual features. In *European Conference on Computer Vision*, 2018.

Minshuo Chen, Yu Bai, Jason D Lee, Tuo Zhao, Huan Wang, Caiming Xiong, and Richard Socher. Towards understanding hierarchical learning: Benefits of neural representations. *arXiv preprint arXiv:2006.13436*, 2020a.

Ting Chen, Simon Kornblith, Mohammad Norouzi, and Geoffrey Hinton. A simple framework for contrastive learning of visual representations. In *International Conference on Machine Learning*, 2020b.

Lenaic Chizat, Edouard Oyallon, and Francis Bach. On lazy training in differentiable programming. In *Advances in Neural Information Processing Systems*, 2019.

Amit Daniely and Eran Malach. Learning parities with neural networks. *Advances in Neural Information Processing Systems*, 2020.

Li Deng. The mnist database of handwritten digit images for machine learning research. *IEEE Signal Processing Magazine*, 29(6):141–142, 2012.

Ilias Diakonikolas, Surbhi Goel, Sushrut Karmalkar, Adam R Klivans, and Mahdi Soltanolkotabi. Approximation schemes for relu regression. In *Conference on Learning Theory*, 2020.

Xialiang Dou and Tengyuan Liang. Training neural networks as learning data-adaptive kernels: Provable representation and approximation benefits. *Journal of the American Statistical Association*, 2020.

Simon Du, Jason Lee, Haochuan Li, Liwei Wang, and Xiyu Zhai. Gradient descent finds global minima of deep neural networks. In *International Conference on Machine Learning*, 2019.

Cong Fang, Hanze Dong, and Tong Zhang. Over parameterized two-level neural networks can learn near optimal feature representations, 2019.

Cong Fang, Hanze Dong, and Tong Zhang. Mathematical models of overparameterized neural networks. *Proceedings of the IEEE*, 109(5):683–703, 2021.

Spencer Frei, Yuan Cao, and Quanquan Gu. Agnostic learning of a single neuron with gradient descent. In *Advances in Neural Information Processing Systems*, 2020.

Mario Geiger, Stefano Spigler, Arthur Jacot, and Matthieu Wyart. Disentangling feature and lazy training in deep neural networks. *Journal of Statistical Mechanics: Theory and Experiment*, 2020 (11):113301, 2020.

Behrooz Ghorbani, Song Mei, Theodor Misiakiewicz, and Andrea Montanari. When do neural networks outperform kernel methods? In *Advances in Neural Information Processing Systems*, 2020.

Ross Girshick, Jeff Donahue, Trevor Darrell, and Jitendra Malik. Rich feature hierarchies for accurate object detection and semantic segmentation. In *Computer Vision and Pattern Recognition*, 2014.

Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep residual learning for image recognition. In *Computer Vision and Pattern Recognition*, 2016.

Kaiming He, Haoqi Fan, Yuxin Wu, Saining Xie, and Ross Girshick. Momentum contrast for unsupervised visual representation learning. In *Computer Vision and Pattern Recognition*, 2020.

Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. In *Advances in neural information processing systems*, 2018.

Ziwei Ji and Matus Telgarsky. Polylogarithmic width suffices for gradient descent to achieve arbitrarily small test error with shallow relu networks. In *International Conference on Learning Representations*, 2019.

Longlong Jing and Yingli Tian. Self-supervised visual feature learning with deep neural networks: A survey. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 2020.

Pritish Kamath, Omar Montasser, and Nathan Srebro. Approximate is good enough: Probabilistic variants of dimensional and margin complexity. In *Conference on Learning Theory*, 2020.

Michael Kearns. Efficient noise-tolerant learning from statistical queries. *Journal of the ACM*, 1998.

Frederic Koehler and Andrej Risteski. The comparative power of relu networks and polynomial kernels in the presence of sparse latent structure. In *International Conference on Learning Representations*, 2018.

Alex Krizhevsky. Learning multiple layers of features from tiny images. *University of Toronto*, 2012.

Ya Le and Xuan Yang. Tiny imagenet visual recognition challenge. *CS 231N*, 2015.

Yann LeCun, Léon Bottou, Yoshua Bengio, and Patrick Haffner. Gradient-based learning applied to document recognition. *Proceedings of the IEEE*, 86(11):2278–2324, 1998.

Jaehoon Lee, Lechao Xiao, Samuel Schoenholz, Yasaman Bahri, Roman Novak, Jascha Sohl-Dickstein, and Jeffrey Pennington. Wide neural networks of any depth evolve as linear models under gradient descent. *Advances in neural information processing systems*, 2019.

Yuanzhi Li and Yingyu Liang. Learning overparameterized neural networks via stochastic gradient descent on structured data. In *Advances in Neural Information Processing Systems*, 2018.

Yuanzhi Li, Colin Wei, and Tengyu Ma. Towards explaining the regularization effect of initial large learning rate in training neural networks. *Advances in Neural Information Processing Systems*, 2019.

Yuanzhi Li, Tengyu Ma, and Hongyang R Zhang. Learning over-parametrized two-layer neural networks beyond ntk. In *Conference on Learning Theory*, 2020.

Eran Malach, Pritish Kamath, Emmanuel Abbe, and Nathan Srebro. Quantifying the benefit of using differentiable learning over tangent kernels. *arXiv preprint arXiv:2103.01210*, 2021.

Christopher D Manning, Kevin Clark, John Hewitt, Urvashi Khandelwal, and Omer Levy. Emergent linguistic structure in artificial neural networks trained by self-supervision. *Proceedings of the National Academy of Sciences*, 117(48):30046–30054, 2020.

Alexander G de G Matthews, Mark Rowland, Jiri Hron, Richard E Turner, and Zoubin Ghahramani. Gaussian process behaviour in wide deep neural networks. In *International Conference on Learning Representations*, 2018.

Preetum Nakkiran, Gal Kaplun, Yamini Bansal, Tristan Yang, Boaz Barak, and Ilya Sutskever. Deep double descent: Where bigger models and more data hurt. In *International Conference on Learning Representations*, 2020.

Yuval Netzer, Tao Wang, Adam Coates, Alessandro Bissacco, Bo Wu, and Andrew Y Ng. Reading digits in natural images with unsupervised feature learning. 2011.

Roman Novak, Lechao Xiao, Jaehoon Lee, Yasaman Bahri, Daniel A Abolafia, Jeffrey Pennington, and Jascha Sohl-Dickstein. Bayesian convolutional neural networks with many channels are gaussian processes. In *International Conference on Learning Representations*, 2019.

B. Olshausen and D. Field. Sparse coding with an overcomplete basis set: A strategy employed by v1? *Vision Research*, 37:3311–3325, 1997.

Ali Rahimi and Benjamin Recht. Random features for large-scale kernel machines. In *Advances in Neural Information Processing Systems*, 2008.

Maria Refinetti, Sebastian Goldt, Florent Krzakala, and Lenka Zdeborová. Classifying high-dimensional gaussian mixtures: Where kernel methods fail and neural networks succeed, 2021.

William E Vinje and Jack L Gallant. Sparse coding and decorrelation in primary visual cortex during natural vision. *Science*, 287(5456):1273–1276, 2000.

Blake Woodworth, Suriya Gunasekar, Jason D Lee, Edward Moroshko, Pedro Savarese, Itay Golan, Daniel Soudry, and Nathan Srebro. Kernel and rich regimes in overparametrized models. In *Conference on Learning Theory*, 2020.

Greg Yang. Scaling limits of wide neural networks with weight sharing: Gaussian process behavior, gradient independence, and neural tangent kernel derivation. *arXiv preprint arXiv:1902.04760*, 2019.

Jianchao Yang, Kai Yu, Yihong Gong, and Thomas Huang. Linear spatial pyramid matching using sparse coding for image classification. In *Computer Vision and Pattern Recognition*, 2009.

Gilad Yehudai and Shamir Ohad. Learning a single neuron with gradient methods. In *Conference on Learning Theory*, 2020.

Gilad Yehudai and Ohad Shamir. On the power and limitations of random features for understanding neural networks. *Advances in Neural Information Processing Systems*, 2019.

Matthew D Zeiler and Rob Fergus. Visualizing and understanding convolutional networks. In *European Conference on Computer Vision*, 2014.

Chiyuan Zhang, Samy Bengio, and Yoram Singer. Are all layers created equal? *arXiv preprint arXiv:1902.01996*, 2019.

Mo Zhou, Rong Ge, and Chi Jin. A local convergence theory for mildly over-parameterized two-layer neural network. In *Conference on Learning Theory*, 2021.

Appendix

Section A presents more technical discussion on related work. Section B-D provides the complete proofs for our results in the main text. Section E provides the complete details and experimental results for our experiments.

Finally, Section F provides the theoretical results and complete proofs for a setting more general than that in the main text, allowing incoherent dictionaries, unbalanced classes, and Gaussian noise in the data.

A   MORE TECHNICAL DISCUSSION ON RELATED WORK

Advantage of Neural Networks over Linear Models on Fixed Features. A recent line of work has turned to show learning settings where network learning provably has advantage over linear models on fixed features; see the nice summary in Malach et al. (2021). Here we highlight the results and focuses of the existing related studies and discuss the differences from ours.

Yehudai & Shamir (2019) shows that the random feature method fails to learn even a single ReLU neuron on Gaussian inputs unless its size is exponentially large in dimension. This points out the limitation of the random feature method (belonging to the fixed feature approach) but does not consider feature learning in networks.

Some studies show that a single ReLU neuron can be learnt by gradient descent (Yehudai & Ohad, 2020; Diakonikolas et al., 2020; Frei et al., 2020). The analysis typically involves feature learning. However, their focus is different: they do not show the advantage over fixed feature methods and do not consider the effect of the input structures.

Zhou et al. (2021) shows that in a special teacher-student setting, the student network will do exact local convergence in a surprising way that all student neurons will converge to one of the teacher neurons. The work does not consider the effect of the input structure nor the advantage over fixed features.

Dou & Liang (2020) explains the advantage of network learning by constructing adaptive Reproducing Kernel Hilbert Space (RKHS) indexed by the training process of the neural network, and shows that adaptive RKHS benefits from a smaller function space containing the residue comparing to RKHS. The work shows the statistical advantage of networks over data-independent kernels, but does not consider the optimization for learning the network.

Ghorbani et al. (2020) considers data generated from a hidden vector with two subsets of variables, each uniformly distributed in a high-dimensional sphere (with a different radius), while the label is determined by only the first subset of variables. It shows the existence of good neural networks that can overcome the curse of dimensionality by representing the best low-dimensional hidden structure. However, it studies the approximation power of neural networks rather than the learning, i.e., it does not show how to learn the good network.

Fang et al. (2019) argues that in the infinite width limit, a two-layer neural network will learn a nearly optimal feature representation in the distribution sense, thanks to the convexity of the limit problem. It is unclear how this result helps to understand the feature learning procedure for practical networks, which is usually a non-convex process.

Chen et al. (2020a) considers a fixed, randomly initialized neural network as a representation function fed into another trainable network which is the quadratic Taylor model of a wide two-layer network. It shows that learning over the random representation can achieve improved sample complexities compared to learning over the raw data. However, the representation considered is not learned, which is different from our focus on feature learning.

Allen-Zhu & Li (2020a) considers Gaussian inputs with labels given by a multiple-layer network with quadratic activations and skip connections (with the assumption of information gap on the weights), and studies training a deep network with quadratic activation. It shows that the trained network can learn proper representations and obtain small errors while no polynomial fixed feature methods can. On the other hand, it does not focus on the influence of input structure on feature learning: note that its input distribution contains no information about the "ground-truth" features in the target network.

It also points out that the learned features get improved during training: higher-level layers will help lower-level layers to improve by backpropagating correction signals. Our analysis also shows feature improvement which however is by signals from the input distribution.

Allen-Zhu & Li (2019) considers PAC learning with labels given by a depth-2 ResNet, and studies training an overparameterized depth-2 ResNet (using uniform inputs over Boolean hypercube as an example). It shows the trained network can obtain small errors while no polynomial kernel methods can obtain as good errors. Similar to [1], it doesn't focus on the influence of input structure on feature learning or the advantage of networks.

Daniely & Malach (2020) considers the task of learning sparse parities with two-layer networks, and the analysis suggests that the ability to learn the label-correlated features also seems to be critical towards the success of neural networks, although the authors did not explore much in this direction. Malach et al. (2021) also considers similar learning problems but with specifically designed models for the problems. The learning problems considered in Daniely & Malach (2020); Malach et al. (2021) have input distributions that leak information about the target labeling function, which is similar to our setting, and their analysis also shows that the first gradient descent can learn a set of good features and later steps can learn an accurate classifier on top. However, they have not explicitly considered if the input structure is crucial for the feature learning and consequently the advantage over fixed features. Our work focuses on studying if the input structures can crucially affect the advantage of networks using a learning problem motivated by practical data, while previous studies use the input structure in the analysis but do not explicitly focus on the effect. Furthermore, there are a few more important differences. First, their focuses are different from ours. Daniely & Malach (2020) focuses on showing neural networks can learn targets (i.e., k-parity functions) that are inherently non-linear. Our analysis generalizes to more general distributions, including practically motivated ones. Malach et al. (2021) focuses on strong separations between learning with gradient descent on differentiable models (including typical neural networks) and learning using the corresponding tangent kernels. The analysis is on specific differentiable models, while our work is on two-layer neural networks similar to practical ones. Second, our analysis relies on the feature improvement in the second gradient step. This is not an artifact of the analysis but comes from our problem setup. While in Daniely & Malach (2020) the data distribution allows some neurons to be sufficiently good after the first gradient step and needs no feature improvement, our setup is more general where the data distribution may not have a similar strong benign effect and thus needs feature improvement in the second gradient step.

Sparse Coding and Subspace Data Models. To analyze neural networks' performance, various data models have been considered. A practical way to model the underlying structure of data is by assuming that a set of hidden variables exists and the input data is a high dimensional projection of the hidden vector (possibly with noise). Along this line, the classic sparse coding model has been used in existing works for analyzing networks. Koehler & Risteski (2018) considers such a data distribution where the label is given by a linear function on the hidden sparse vector, but studies the approximation power of networks and classic polynomial methods rather than the learning. Allen-Zhu & Li (2020b) considers similar data distributions, but studies the performance of networks under adversarial perturbations. Another type of related data models assumes that the label is determined by a subset of hidden variables. Ghorbani et al. (2020) considers a hidden vector with two subsets of variables, each uniformly distributed in a high-dimensional sphere (with a different radius), while the label is determined by only the first subset of variables. However, Ghorbani et al. (2020) studies the approximation power of neural networks rather than the learning. Compared to these studies, our work assumes the input is given by a dictionary multiplied with a hidden vector (not necessarily sparse) while the label is determined by a subset of the hidden vector, as motivated by pattern recognition applications in practice. Furthermore, we focus on the learning ability of networks instead of approximation.

B COMPLETE PROOFS FOR PROVABLE GUARANTEES OF NEURAL NETWORKS

Notations.   For a vector v and an index set I, let v_I denote the vector containing the entries of v indexed by I, and v_{-I} denote the vector containing the entries of v with indices outside I.

By initialization, $w_i^{(0)}$ for $i \in [m]$ are i.i.d. copies of the same random variable $w^{(0)} \sim \mathcal{N}(0, \sigma_w^2 I_{d \times d})$; similar for $a^{(0)}$ and $b^{(0)}$. Let $q_\ell := \langle w^{(0)}, M_\ell \rangle$, then $\langle w^{(0)}, x \rangle = \langle \phi, q \rangle$. Similarly, define $q_{i,\ell}^{(t)} := \langle w_i^{(t)}, M_\ell \rangle$. Let $\sigma_\phi^2 := p_o(1 - p_o)/\tilde{\sigma}^2$ denote the variance of ϕ_ℓ for $\ell \notin A$.

We also define the following sets to denote typical initialization. For a fixed $\delta \in (0, 1)$, define

$$\mathcal{G}_w(\delta) := \left\{ w \in \mathbb{R}^d : q_\ell = \langle w, M_\ell \rangle, \frac{\sigma_w^2(D-k)}{2} \le \sum_{\ell \notin A} q_\ell^2 \le \frac{3\sigma_w^2(D-k)}{2}, \right.$$

$$\left. \max_\ell |q_\ell| \le \sigma_w \sqrt{2\log(Dm/\delta)} \right\}, \tag{5}$$

$$\mathcal{G}_a(\delta) := \{ a \in \mathbb{R} : |a| \le \sigma_a \sqrt{2\log(m/\delta)} \}. \tag{6}$$

$$\mathcal{G}_b(\delta) := \{ b \in \mathbb{R} : |b| \le \sigma_b \sqrt{2\log(m/\delta)} \}. \tag{7}$$

B.1 EXISTENCE OF A GOOD NETWORK

we first show that there exists a network that can fit the data distribution.

Lemma 7. *For some $s, a, b \in \mathbb{R}$ with $a, b \ge 0$, define a function $\delta_{s,a,b} : \mathbb{R} \to \mathbb{R}$ as*

$$\delta_{s,a,b}(z) = a\sigma_r(z - s + b) - 2a\sigma_r(z - s) + a\sigma_r(z - s - b). \tag{8}$$

where $\sigma_r(z) = \max\{z, 0\}$ is the ReLU activation function. Then

$$\delta_{s,a,b}(z) = \begin{cases} 0 & \text{when } z \le s - b, \\ a(z - s) + ab & \text{when } s - b \le z \le s, \\ -a(z - s) + ab & \text{when } s \le z \le s + b, \\ 0 & \text{when } s + b \le z. \end{cases} \tag{9}$$

That is, $\delta_{s,a,b}(z)$ linearly interpolates between $(s - b, 0), (s, ab), (s + b, 0)$ when $z \in [s - b, s + b]$, and is 0 elsewhere.

Proof of Lemma 7. This can be simply verified for the four cases of the value of z. □

Lemma 8 (Restatement of Lemma 4). *For any $\mathcal{D} \in \mathcal{F}_\Xi$, there exists a network $g^*(x) = \sum_{i=1}^n a_i^* \sigma(\langle w_i^*, x \rangle + b_i^*)$ with $y = g^*(x)$ for any $(x, y) \sim \mathcal{D}$. Furthermore, the number of neurons $n = 3(k + 1)$, $|a_i^*| \le 32k$, $1/(32k) \le |b_i^*| \le 1/2$, $w_i^* = \tilde{\sigma} \sum_{j \in A} M_j/(4k)$, and $|\langle w_i^*, x \rangle + b_i^*| \le 1$ for any $i \in [n]$ and $(x, y) \sim \mathcal{D}$.*

Proof of Lemma 4. Let $w = \tilde{\sigma} \sum_{j \in A} M_j$ and let $\mu = \sum_{j \in A} \mathbb{E}[\tilde{\phi}_j]$. We have

$$\langle w, x \rangle = \tilde{\sigma} \sum_{j \in A} \langle M_j, M\phi \rangle = \tilde{\sigma} \sum_{j \in A} \phi_j = \sum_{j \in A} \tilde{\phi}_j - \mu. \tag{10}$$

Then by Lemma 7,

$$g_1^*(x) := \sum_{p \in P} \delta_{p - \mu, 2, 1/2}(\langle w, x \rangle) - \sum_{p \notin P, 0 \le p \le k} \delta_{p - \mu, 2, 1/2}(\langle w, x \rangle) \tag{11}$$

$$= \sum_{p \in P} \delta_{p, 2, 1/2}(\langle w, x \rangle + \mu) - \sum_{p \notin P, 0 \le p \le k} \delta_{p, 2, 1/2}(\langle w, x \rangle + \mu) \tag{12}$$

$$= \sum_{p \in P} \delta_{p, 2, 1/2}\left(\sum_{j \in A} \tilde{\phi}_j \right) - \sum_{p \notin P, 0 \le p \le k} \delta_{p, 2, 1/2}\left(\sum_{j \in A} \tilde{\phi}_j \right) \tag{13}$$

$$= y \tag{14}$$

for any $(x, y) \sim \mathcal{D}$. Similarly,

$$g_2^*(x) := \sum_{p \in P} \delta_{p-\mu+1/4,4,1/2}(\langle w, x \rangle) - \sum_{p \notin P, 0 \leq p \leq k} \delta_{p-\mu+1/4,4,1/2}(\langle w, x \rangle) \tag{15}$$

$$= \sum_{p \in P} \delta_{p+1/4,4,1/2}(\langle w, x \rangle + \mu) - \sum_{p \notin P, 0 \leq p \leq k} \delta_{p+1/4,4,1/2}(\langle w, x \rangle + \mu) \tag{16}$$

$$= \sum_{p \in P} \delta_{p+1/4,4,1/2} \left(\sum_{j \in A} \tilde{\phi}_j \right) - \sum_{p \notin P, 0 \leq p \leq k} \delta_{p+1/4,4,1/2} \left(\sum_{j \in A} \tilde{\phi}_j \right) \tag{17}$$

$$= y \tag{18}$$

for any $(x, y) \sim \mathcal{D}$. Note that the bias terms in g_1^* and g_2^* have distance at least $1/4$, then at least one of them satisfies that all its bias terms have absolute value $\geq 1/8$. Pick that one and denote it as $g(x) = \sum_{i=1}^n a_i \sigma_r(\langle w_i, x \rangle + b_i)$. By the positive homogeneity of σ_r, we have

$$g(x) = \sum_{i=1}^n 4k a_i \sigma_r(\langle w_i, x \rangle/(4k) + b_i/(4k)). \tag{19}$$

Since for any $(x, y) \sim \mathcal{D}$, $|\langle w_i, x \rangle/(4k) + b_i/(4k)| \leq 1$, then

$$g(x) = \sum_{i=1}^n 4k a_i \sigma(\langle w_i, x \rangle/(4k) + b_i/(4k)) \tag{20}$$

where σ is the truncated ReLU. Now we can set $a_i^* = 4k a_i, w_i^* = w_i/(4k), b_i^* = b_i/(4k)$, to get our final g^*. $\qquad\square$

B.2 Initialization

We first show that with high probability, the initial weights are in typical positions.

Lemma 9. *For any $\delta \in (0, 1)$, with probability at least $1 - \delta - 2\exp\left(-\Theta(D-k)\right)$ over $w^{(0)}$,*

$$\sigma_w^2(D-k)/2 \leq \sum_{\ell \notin A} q_\ell^2 \leq 3\sigma_w^2(D-k)/2,$$

$$\max_\ell |q_\ell| \leq \sigma_w \sqrt{2\log(D/\delta)}.$$

With probability at least $1 - \delta$ over $b^{(0)}$,

$$|b^{(0)}| \leq \sigma_b \sqrt{2\log(1/\delta)}.$$

With probability at least $1 - \delta$ over $a^{(0)}$,

$$|a^{(0)}| \leq \sigma_a \sqrt{2\log(1/\delta)}.$$

Proof of Lemma 9. From $q \sim \mathcal{N}(0, \sigma_w^2 I_{d \times d})$, we have:

- With probability $\geq 1 - \delta/2$, $\max_\ell |q_\ell| \leq \sqrt{2\sigma_w^2 \log \frac{D}{\delta}}$, and

- For any subset $S \subseteq [D]$, with probability $\geq 1 - 2\exp\left(-\Theta(|S|)\right)$, $\|q_S\|_2^2 \in \left(\frac{|S|\sigma_w^2}{2}, \frac{3|S|\sigma_w^2}{2}\right)$.

Similar for $b^{(0)}$ and $a^{(0)}$. The lemma then follows. $\qquad\square$

Lemma 10. *We have:*

- *With probability $\geq 1 - \delta - 2m\exp(-\Theta(D-k))$ over $w_i^{(0)}$'s, for all $i \in [2m]$, $w_i^{(0)} \in \mathcal{G}_w(\delta)$.*

- *With probability $\geq 1 - \delta$ over $b_i^{(0)}$'s, for all $i \in [2m]$, $b_i^{(0)} \in \mathcal{G}_b(\delta)$.*

- *With probability $\geq 1 - \delta$ over $a_i^{(0)}$'s, for all $i \in [2m]$, $a_i^{(0)} \in \mathcal{G}_a(\delta)$.*

Proof of Lemma 10. This follows from Lemma 9 by union bound. $\square$

The following lemma about the typical $w_i^{(0)}$'s will be useful for later analysis.

Lemma 11. *Fix $\delta \in (0, 1)$. For any $w_i^{(0)} \in \mathcal{G}_w(\delta)$, we have*

$$\Pr_\phi \left[\sum_{\ell \notin A} \phi_\ell q_{i,\ell}^{(0)} \geq \Theta \left(\sqrt{(D-k)\sigma_\phi^2 \sigma_w^2} \right) \right] = \Theta(1) - \frac{O(\log^{3/2}(Dm/\delta))}{\sqrt{(D-k)\sigma_\phi^2 \tilde{\sigma}^2}}. \tag{21}$$

Consequently, when $p_o = \Omega(k^2/D)$ and $k = \Omega(\log^2(Dm/\delta))$,

$$\Pr_\phi \left[\sum_{\ell \notin A} \phi_\ell q_{i,\ell}^{(0)} \geq \Theta(\sigma_w) \right] = \Theta(1) - \frac{O(1)}{k^{1/4}}. \tag{22}$$

Proof of Lemma 11. Note that for $\ell \notin A$, $\mathbb{E}[\phi_\ell] = 0$, $\mathbb{E}[\phi_\ell^2] = \sigma_\phi^2$, and $\mathbb{E}[|\phi_\ell|^3] = \Theta(\sigma_\phi^2/\tilde{\sigma})$. Then the statement follows from Berry-Esseen Theorem. $\square$

B.3 SOME AUXILIARY LEMMAS

The expression of the gradients will be used frequently.

Lemma 12.

$$\frac{\partial}{\partial w_i} L_\mathcal{D}(g; \sigma_\xi) = -a_i \mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ y \mathbb{I}[yg(x; \xi) \leq 1] \mathbb{E}_{\xi_i} \mathbb{I}[\langle w_i, x \rangle + b_i + \xi_i \in (0, 1)]x \right\}, \tag{23}$$

$$\frac{\partial}{\partial b_i} L_\mathcal{D}(g; \sigma_\xi) = -a_i \mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ y \mathbb{I}[yg(x; \xi) \leq 1] \mathbb{E}_{\xi_i} \mathbb{I}[\langle w_i, x \rangle + b_i \in (0, 1)] \right\}, \tag{24}$$

$$\frac{\partial}{\partial a_i} L_\mathcal{D}(g; \sigma_\xi) = -\mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ y \mathbb{I}[yg(x; \xi) \leq 1] \mathbb{E}_{\xi_i} \sigma(\langle w_i, x \rangle + b_i + \xi_i) \right\}. \tag{25}$$

Proof of Lemma 12. It follows from straightforward calculation. $\square$

We now show that a small subset of the entries in ϕ, q does not affect the probability distribution of $\langle \phi, q \rangle$ much.

Lemma 13. *Suppose $\nu \sim \mathcal{N}(0, \sigma^2)$. For any $B \supseteq A$ and any b:*

$$\left| \Pr_{\phi_{-B}, \nu} \left\{ \langle \phi, q \rangle + \nu \geq b \right\} - \Pr_{\phi_{-B}, \nu} \left\{ \langle \phi_{-B}, q_{-B} \rangle + \nu \geq b \right\} \right| \tag{26}$$

$$\leq O \left(\frac{|\langle \phi_B, q_B \rangle|}{(\sigma_\phi^2 \|q_{-B}\|_2^2 + \sigma^2)^{1/2}} + \frac{\sigma^3 + \sigma_\phi^2 \|q_{-B}\|_3^3/\tilde{\sigma}}{(\sigma^2 + \sigma_\phi^2 \|q_{-B}\|_2^2)^{3/2}} \right). \tag{27}$$

Similarly,

$$\left| \Pr_{\phi_{-B}, \nu} \left\{ \langle \phi, q \rangle \geq b \right\} - \Pr_{\phi_{-B}, \nu} \left\{ \langle \phi_{-B}, q_{-B} \rangle \geq b \right\} \right| \tag{28}$$

$$\leq O \left(\frac{|\langle \phi_B, q_B \rangle|}{\sigma_\phi \|q_{-B}\|_2} + \frac{\|q_{-B}\|_3^3}{\tilde{\sigma} \sigma_\phi \|q_{-B}\|_2^3} \right). \tag{29}$$

Proof of Lemma 13. Note that for $\ell \notin A$, $\mathbb{E}[\phi_\ell] = 0$, $\mathbb{E}[\phi_\ell^2] = \sigma_\phi^2$, and $\mathbb{E}[|\phi_\ell|^3] = \Theta(\sigma_\phi^2/\tilde{\sigma})$. Let $t = |\langle \phi_B, q_B \rangle|$. Then by the Berry-Esseen Theorem,

$$\left| \Pr_{\phi_{-B}} \left\{ \langle \phi, q \rangle + \nu \geq b \right\} - \Pr_{\phi_{-B}} \left\{ \langle \phi_{-B}, q_{-B} \rangle + \nu \geq b \right\} \right| \tag{30}$$

$$\leq \Pr_{\phi_{-B}} \left\{ \langle \phi_{-B}, q_{-B} \rangle + \nu \in [-t + b, t + b] \right\} \tag{31}$$

$$\leq \frac{2t}{(\sigma_\phi^2 \|q_{-B}\|_2^2 + \sigma^2)^{1/2}} + \frac{O(\sigma^3 + \sigma_\phi^2 \|q_{-B}\|_3^3/\tilde{\sigma})}{(\sigma^2 + \sigma_\phi^2 \|q_{-B}\|_2^2)^{3/2}}. \tag{32}$$

The second statement follows from a similar argument. $\square$

We also have the following auxiliary lemma for later calculations.

Lemma 14.

$$\mathbb{E}_{\phi_A} \{y\} = 0, \tag{33}$$

$$\mathbb{E}_{\phi_A} \{|y|\} = 1, \tag{34}$$

$$\mathbb{E}_{\phi_j} \{|\phi_j|\} = 2\sigma_\phi^2 \tilde{\sigma}, \text{ for } j \notin A, \tag{35}$$

$$\mathbb{E}_{\phi_A} \{y\phi_j\} = \frac{\gamma}{\tilde{\sigma}}, \tag{36}$$

$$\mathbb{E}_{\phi_A} \{|y\phi_j|\} \leq \frac{1}{\tilde{\sigma}}, \text{ for all } j \in [D]. \tag{37}$$

Proof of Lemma 14.

$$\mathbb{E}_{\phi_A} \{y\} = \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A} \{y | y = v\} \Pr[y = v] \tag{38}$$

$$= \frac{1}{2} \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A} \{y | y = v\} \tag{39}$$

$$= 0. \tag{40}$$

$$\mathbb{E}_{\phi_A} \{|y|\} = \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A} \{|y| \,|\, y = v\} \Pr[y = v] \tag{41}$$

$$= \frac{1}{2} \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A} \{|y| \,|\, y = v\} \tag{42}$$

$$= 1. \tag{43}$$

$$\mathbb{E}_{\phi_j} \{|\phi_j|\} = \frac{|-p_o|(1 - p_o) + |1 - p_o|p_o}{\tilde{\sigma}} = 2\sigma_\phi^2 \tilde{\sigma}. \tag{44}$$

$$\mathbb{E}_{\phi_A} \{y\phi_j\} = \mathbb{E}_{\phi_A} \left\{ y \frac{\tilde{\phi}_j - \mathbb{E}[\tilde{\phi}_j]}{\tilde{\sigma}} \right\} \tag{45}$$

$$= \frac{1}{\tilde{\sigma}} \mathbb{E}_{\phi_A} \left\{ y\tilde{\phi}_j - y\mathbb{E}[\tilde{\phi}_j] \right\} \tag{46}$$

$$= \frac{\gamma}{\tilde{\sigma}}. \tag{47}$$

$$\mathbb{E}_{\phi_A} \{|y\phi_j|\} = \mathbb{E}_{\phi_A} \{|\phi_j|\} \tag{48}$$

$$\leq \frac{1}{\tilde{\sigma}}. \tag{49}$$

$\square$

B.4 FEATURE EMERGENCE: FIRST GRADIENT STEP

We will show that w.h.p. over the initialization, after the first gradient step, there are neurons that represent good features.

We begin with analyzing the gradients.

Lemma 15 (Full version of Lemma 5). *Fix $\delta \in (0,1)$ and suppose $w_i^{(0)} \in \mathcal{G}_w(\delta)$, $b_i^{(0)} \in \mathcal{G}_b(\delta)$ for all $i \in [2m]$. Let*

$$\epsilon_e := \frac{k \log^{1/2}(Dm/\delta) + \log^{3/2}(Dm/\delta)}{\sqrt{\sigma_\phi^2 \tilde{\sigma}^2 (D-k)}}.$$

If $p_o = \Omega(k^2/D)$, $k = \Omega(\log^2(Dm/\delta))$, and $\sigma_\xi^{(1)} < 1/k$, then

$$\frac{\partial}{\partial w_i} L_\mathcal{D}(g^{(0)}; \sigma_\xi^{(1)}) = -a_i^{(0)} \sum_{j=1}^D M_j T_j \tag{50}$$

where T_j satisfies:

- *if $j \in A$, then $|T_j - \beta\gamma/\tilde{\sigma}| \leq O(\epsilon_e/\tilde{\sigma})$, where $\beta \in [\Omega(1), 1]$ and depends only on $w_i^{(0)}, b_i^{(0)}$;*

- *if $j \notin A$, then $|T_j| \leq O(\sigma_\phi^2 \epsilon_e \tilde{\sigma})$.*

Proof of Lemma 15. Consider one neuron index i and omit the subscript i in the parameters. Since the unbiased initialization leads to $g^{(0)}(x; \xi^{(1)}) = 0$, we have

$$\frac{\partial}{\partial w} L_\mathcal{D}(g^{(0)}; \sigma_\xi^{(1)}) \tag{51}$$

$$= -a^{(0)} \mathbb{E}_{(x,y)\sim\mathcal{D}} \left\{ y\mathbb{I}[yg^{(0)}(x;\xi^{(1)}) \leq 1]\mathbb{E}_{\xi^{(1)}}\mathbb{I}[\langle w^{(0)}, x\rangle + b^{(0)} + \xi^{(1)} \in (0,1)]x \right\} \tag{52}$$

$$= -a^{(0)} \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(1)}} \left\{ y\mathbb{I}[\langle w^{(0)}, x\rangle + b^{(0)} + \xi^{(1)} \in (0,1)]x \right\} \tag{53}$$

$$= -a^{(0)} \sum_{j=1}^D M_j \underbrace{\mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(1)}} \left\{ y\mathbb{I}[\langle w^{(0)}, x\rangle + b^{(0)} + \xi^{(1)} \in (0,1)]\phi_j \right\}}_{:=T_j}. \tag{54}$$

First, consider $j \in A$.

$$T_j = \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(1)}} \left\{ y\mathbb{I}[\langle w^{(0)}, x\rangle + b^{(0)} + \xi^{(1)} \in (0,1)]\phi_j \right\} \tag{55}$$

$$= \mathbb{E}_{\phi_A,\xi^{(1)}} \left\{ y\phi_j \Pr_{\phi_{-A}} \left[\langle \phi, q\rangle + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right\}. \tag{56}$$

Let

$$I_a := \Pr_{\phi_{-A}} \left[\langle \phi, q\rangle + b^{(0)} + \xi^{(1)} \in (0,1) \right], \tag{57}$$

$$I_a' := \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A}\rangle + b^{(0)} + \xi^{(1)} \in (0,1) \right]. \tag{58}$$

We have

$$|\mathbb{E}_{\xi^{(1)}}(I_a - I_a')| \tag{59}$$

$$\leq \mathbb{E}_{\xi^{(1)}} \left| \Pr_{\phi_{-A}} \left[\langle \phi, q\rangle + b^{(0)} + \xi^{(1)} \geq 0 \right] - \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A}\rangle + b^{(0)} + \xi^{(1)} \geq 0 \right] \right| \tag{60}$$

$$+ \Pr_{\phi_{-A},\xi^{(1)}} \left[\langle \phi, q\rangle + b^{(0)} + \xi^{(1)} \geq 1 \right] + \Pr_{\phi_{-A},\xi^{(1)}} \left[\langle \phi_{-A}, q_{-A}\rangle + b^{(0)} + \xi^{(1)} \geq 1 \right]. \tag{61}$$

Then by Lemma 13,

$$\left| \Pr_{\phi_{-A}} \left[\langle \phi, q\rangle + b^{(0)} + \xi^{(1)} \geq 0 \right] - \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A}\rangle + b^{(0)} + \xi^{(1)} \geq 0 \right] \right| = O(\epsilon_e). \tag{62}$$

Note that $\sum_{\ell \notin A} \mathrm{Var}(\phi_\ell q_\ell) = \Theta(\sigma_\phi^2 \sigma_w^2 (D - k)) = \Theta(\sigma_w^2)$, and $|\phi_\ell| \leq \frac{1}{\tilde{\sigma}}$. If $p_o = \Omega(k^3/D)$ then apply Bennett's inequality in the small-deviation regime; $p_o = O(k^3/D)$ and $p_o = \Omega(k^2/D)$ then apply Bennett's inequality in the large-deviation regime. In both cases we have:

$$\Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle \geq 1/4 \right] = \exp(-\Omega(k)) = O(\epsilon_e). \tag{63}$$

We also have:

$$\Pr_{\xi^{(1)}} \left[b^{(0)} + \xi^{(1)} \geq 1/4 \right] = \exp(-\Omega(k)) = O(\epsilon_e). \tag{64}$$

Therefore,

$$\Pr_{\phi_{-A}, \xi^{(1)}} \left[\langle \phi, q \rangle + b^{(0)} + \xi^{(1)} \geq 1 \right] = \exp(-\Omega(k)) = O(\epsilon_e) \tag{65}$$

where the last step follows from the assumption on σ_w and k. A similar argument gives:

$$\Pr_{\phi_{-A}, \xi^{(1)}} \left[\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} + \xi^{(1)} \geq 1 \right] = \exp(-\Omega(k)) = O(\epsilon_e). \tag{66}$$

Then we have

$$\left| T_j - \mathbb{E}_{\phi_A, \xi^{(1)}} \left\{ y \phi_j I'_a \right\} \right| \tag{67}$$
$$\leq \mathbb{E}_{\phi_A} \left\{ |y \phi_j| \left| \mathbb{E}_{\xi^{(1)}} (I_a - I'_a) \right| \right\} \tag{68}$$
$$\leq O(\epsilon_e) \mathbb{E}_{\phi_A} \left\{ |y \phi_j| \right\} \tag{69}$$
$$\leq O(\epsilon_e/\tilde{\sigma}) \tag{70}$$

where the last step is from Lemma 14. Furthermore,

$$\mathbb{E}_{\phi_A, \xi^{(1)}} \left\{ y \phi_j I'_a \right\} \tag{71}$$
$$= \mathbb{E}_{\phi_A} \left\{ y \phi_j \right\} \mathbb{E}_{\xi^{(1)}} [I'_a] \tag{72}$$
$$= \mathbb{E}_{\phi_A} \left\{ y \phi_j \right\} \Pr_{\phi_{-A}, \xi^{(1)}} \left[\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} + \xi^{(1)} \in (0, 1) \right] \tag{73}$$

By Lemma 11, the assumption on p_o, and (63), we have

$$\Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} \in (0, 1/2) \right] \geq \Omega(1) - O(1/k^{1/4}), \tag{74}$$

$$\Pr_{\xi^{(1)}} \left[\xi^{(1)} \in (0, 1/2) \right] = 1/2 - \exp(-\Omega(k)), \tag{75}$$

This leads to

$$\beta := \mathbb{E}_{\xi^{(1)}} [I'_a] = \Pr_{\phi_{-A}, \xi^{(1)}} \left[\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} + \xi^{(1)} \in (0, 1) \right] \geq \Omega(1). \tag{76}$$

By Lemma 14, $\mathbb{E}_{\phi_A} \left\{ y \phi_j \right\} = \gamma/\tilde{\sigma}$. Therefore,

$$|T_j - \beta \gamma/\tilde{\sigma}| \leq O(\epsilon_e/\tilde{\sigma}). \tag{77}$$

Now, consider $j \notin A$. Let B denote $A \cup \{j\}$.

$$T_j = \mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ y \phi_j \mathbb{I} \left[\langle \phi, q \rangle + b^{(0)} + \xi^{(1)} \in (0, 1) \right] \right\} \tag{78}$$

$$= \mathbb{E}_{\phi_B} \mathbb{E}_{\phi_{-B}, \xi^{(1)}} \left\{ y \phi_j \mathbb{I} \left[\langle \phi, q \rangle + b^{(0)} + \xi^{(1)} \in (0, 1) \right] \right\} \tag{79}$$

$$= \mathbb{E}_{\phi_B, \xi^{(1)}} \left\{ y \phi_j \Pr_{\phi_{-B}} \left[\langle \phi, q \rangle + b^{(0)} + \xi^{(1)} \in (0, 1) \right] \right\}. \tag{80}$$

Let

$$I_b := \Pr_{\phi_{-B}} \left[\langle \phi, q \rangle + b^{(0)} + \xi^{(1)} \in (0, 1) \right], \tag{81}$$

$$I'_b := \Pr_{\phi_{-B}} \left[\langle \phi_{-B}, q_{-B} \rangle + b^{(0)} + \xi^{(1)} \in (0, 1) \right]. \tag{82}$$

Similar as above, we have $|\mathbb{E}_{\xi^{(1)}}(I_b - I'_b)| \leq O(\epsilon_e)$ by Lemma 13. Then by Lemma 14,

$$\left| T_j - \mathbb{E}_{\phi_B, \xi^{(1)}} \left\{ y \phi_j I'_b \right\} \right| \tag{83}$$

$$\leq \mathbb{E}_{\phi_B} \left\{ |y\phi_j| |\mathbb{E}_{\xi^{(1)}}(I_b - I'_b)| \right\} \tag{84}$$

$$\leq O(\epsilon_e) \mathbb{E}_{\phi_A} \left\{ |y| \right\} \mathbb{E}_{\phi_j} \left\{ |\phi_j| \right\} \tag{85}$$

$$\leq O(\epsilon_e) \times O(\sigma_\phi^2 \tilde{\sigma}) \tag{86}$$

$$= O(\sigma_\phi^2 \epsilon_e \tilde{\sigma}). \tag{87}$$

Furthermore,

$$\mathbb{E}_{\phi_B, \xi^{(1)}} \left\{ y \phi_j I'_b \right\} = \mathbb{E}_{\phi_A} \left\{ y \right\} \mathbb{E}_{\phi_j} \left\{ \phi_j \right\} \mathbb{E}_{\xi^{(1)}} [I'_b] = 0. \tag{88}$$

Therefore,

$$|T_j| \leq O(\sigma_\phi^2 \epsilon_e \tilde{\sigma}). \tag{89}$$

$\square$

Lemma 16. *Under the same assumptions as in Lemma 15,*

$$\frac{\partial}{\partial b_i} L_{\mathcal{D}}(g^{(0)}; \sigma_\xi^{(1)}) = -a_i^{(0)} T_b \tag{90}$$

where $|T_b| \leq O(\epsilon_e)$.

Proof of Lemma 16. Consider one neuron index i and omit the subscript i in the parameters. Since the unbiased initialization leads to $g^{(0)}(x; \xi^{(1)}) = 0$, we have

$$\frac{\partial}{\partial b} L_{\mathcal{D}}(g^{(0)}; \sigma_\xi^{(1)}) \tag{91}$$

$$= -a^{(0)} \mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ y \mathbb{I}[y g^{(0)}(x; \xi) \leq 1] \mathbb{E}_{\xi^{(1)}} \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \tag{92}$$

$$= -a^{(0)} \underbrace{\mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ y \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\}}_{:=T_b}. \tag{93}$$

Similar to the proof in Lemma 5,

$$\left| \Pr_{\phi_{-A}} [\langle \phi, q \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] - \Pr_{\phi_{-A}} [\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right| = O(\epsilon_e). \tag{94}$$

Then

$$\left| T_b - \mathbb{E}_{\phi_A, \xi^{(1)}} \left\{ y \Pr_{\phi_{-A}} [\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \right| \tag{95}$$

$$= \mathbb{E}_{\phi_A, \xi^{(1)}} \left\{ |y| \left| \Pr_{\phi_{-A}} [\langle \phi, q \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] - \Pr_{\phi_{-A}} [\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right| \right\} \tag{96}$$

$$\leq O(\epsilon_e) \mathbb{E}_{\phi_A} \left\{ |y| \right\} \tag{97}$$

$$\leq O(\epsilon_e). \tag{98}$$

Also,

$$\mathbb{E}_{\phi_A, \xi^{(1)}} \left\{ y \Pr_{\phi_{-A}} [\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \tag{99}$$

$$= \mathbb{E}_{\phi_A} \left\{ y \right\} \Pr_{\phi_{-A}, \xi^{(1)}} [\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \tag{100}$$

$$= 0. \tag{101}$$

Therefore, $|T_b| \leq O(\epsilon_e)$. $\square$

Lemma 17. *We have*

$$\frac{\partial}{\partial a_i} L_{\mathcal{D}}(g^{(0)}; \sigma_\xi^{(1)}) = -T_a \tag{102}$$

where $|T_a| \leq O(\max_\ell q_{i,\ell}^{(0)})$. *So if* $w_i^{(0)} \in \mathcal{G}(\delta)$, $|T_a| \leq O(\sigma_w \sqrt{\log(Dm/\delta)})$.

Proof of Lemma 17. Consider one neuron index i and omit the subscript i in the parameters. Since the unbiased initialization leads to $g^{(0)}(x; \xi^{(1)}) = 0$, we have

$$\frac{\partial}{\partial a} L_{\mathcal{D}}(g^{(0)}; \sigma_\xi^{(1)}) \tag{103}$$

$$= -\mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ y\mathbb{I}[yg^{(0)}(x; \xi^{(1)}) \leq 1] \mathbb{E}_{\xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) \right\} \tag{104}$$

$$= -\underbrace{\mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ y\sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) \right\}}_{:=T_a}. \tag{105}$$

Let ϕ'_A be an independent copy of ϕ_A, ϕ' be the vector obtained by replacing in ϕ the entries ϕ_A with ϕ'_A, and let $x' = M\phi'$ and its label is y'. Then

$$|T_a| = \left| \mathbb{E}_{\phi_A} \left\{ y\mathbb{E}_{\phi_{-A}, \xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) \right\} \right| \tag{106}$$

$$\leq \frac{1}{2} \left| \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) | y = 1 \right\} \right. \tag{107}$$

$$\left. - \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) | y = -1 \right\} \right| \tag{108}$$

$$\leq \frac{1}{2} \left| \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) | y = 1 \right\} \right. \tag{109}$$

$$\left. - \mathbb{E}_{\phi'_A} \left\{ \mathbb{E}_{\phi_{-A}, \xi^{(1)}} \sigma(\langle w^{(0)}, x' \rangle + b^{(0)} + \xi^{(1)}) | y' = -1 \right\} \right|. \tag{110}$$

Since σ is 1-Lipschitz,

$$|T_a| \leq \frac{1}{2} \mathbb{E}_{\phi_A, \phi'_A} \left\{ \mathbb{E}_{\phi_{-A}} \left| \langle w^{(0)}, x \rangle - \langle w^{(0)}, x' \rangle \right| | y = 1, y' = -1 \right\} \tag{111}$$

$$\leq \frac{1}{2} \mathbb{E}_{\phi_{-A}} \left(\mathbb{E}_{\phi_A} \left\{ \left| \langle w^{(0)}, x \rangle \right| | y = 1 \right\} + \mathbb{E}_{\phi'_A} \left\{ \left| \langle w^{(0)}, x' \rangle \right| | y' = -1 \right\} \right) \tag{112}$$

$$= \mathbb{E}_{\phi_{-A}, \phi_A} \left| \langle w^{(0)}, x \rangle \right| \tag{113}$$

$$= \mathbb{E}_x \left| \langle w^{(0)}, x \rangle \right| \tag{114}$$

$$\leq \sqrt{\mathbb{E}_x \langle w^{(0)}, x \rangle^2} \tag{115}$$

$$\leq \max_\ell q_{i,\ell}^{(0)} \sqrt{\mathbb{E}_x \left(\sum_{\ell \in [D]} \phi_\ell^2 + \sum_{j \neq \ell: j, \ell \in A} |\phi_j \phi_\ell| \right)} \tag{116}$$

$$\leq \max_\ell q_{i,\ell}^{(0)} \sqrt{\mathbb{E}_x (1 + O(1))} \tag{117}$$

$$= \Theta(\max_\ell q_{i,\ell}^{(0)}). \tag{118}$$

$\square$

With the bounds on the gradient, we now summarize the results for the weights after the first gradient step.

Lemma 18. *Set*

$$\lambda_w^{(1)} = 1/(2\eta^{(1)}), \lambda_a^{(1)} = \lambda_b^{(1)} = 0, \sigma_\xi^{(1)} = 1/k^{3/2}.$$

Fix $\delta \in (0,1)$ *and suppose* $w_i^{(0)} \in \mathcal{G}_w(\delta), b_i^{(0)} \in \mathcal{G}_b(\delta)$ *for all* $i \in [2m]$. *If* $p_o = \Omega(k^2/D)$, $k = \Omega(\log^2(Dm/\delta))$, *then for all* $i \in [m]$, $w_i^{(1)} = \sum_{\ell=1}^D q_{i,\ell}^{(1)} M_\ell$ *satisfying*

- *if* $\ell \in A$, *then* $|q_{i,\ell}^{(1)} - \eta^{(1)} a_i^{(0)} \beta\gamma/\tilde{\sigma}| \leq O\left(\frac{|\eta^{(1)} a_i^{(0)}|\epsilon_e}{\tilde{\sigma}}\right)$, *where* $\beta \in [\Omega(1), 1]$ *and depends only on* $w_i^{(0)}, b_i^{(0)}$;

- *if* $\ell \notin A$, *then* $|q_{i,\ell}^{(1)}| \leq O\left(\sigma_\phi^2 |\eta^{(1)} a_i^{(0)}|\epsilon_e\tilde{\sigma}\right)$;

and

- $b_i^{(1)} = b_i^{(0)} + \eta^{(1)} a_i^{(0)} T_b$ *where* $|T_b| = O(\epsilon_e)$;

- $a_i^{(1)} = a_i^{(0)} + \eta^{(1)} T_a$ *where* $|T_a| = O(\sigma_w \sqrt{\log(Dm/\delta)})$.

Proof of Lemma 18. This follows from Lemma 10 and Lemma 15-17. □

B.5 FEATURE IMPROVEMENT: SECOND GRADIENT STEP

We first show that with properly set $\eta^{(1)}$, for most x, $|g^{(1)}(x; \sigma_\xi^{(2)})| < 1$ and thus $yg^{(1)}(x; \sigma_\xi^{(2)}) < 1$.

Lemma 19. *Fix* $\delta \in (0,1)$ *and suppose* $w_i^{(0)} \in \mathcal{G}_w(\delta), b_i^{(0)} \in \mathcal{G}_b(\delta), a_i^{(0)} \in \mathcal{G}_a(\delta)$ *for all* $i \in [2m]$. *If* $p_o = \Omega(k^2/D)$, $k = \Omega(\log^2(Dm/\delta))$, $\sigma_a \leq \tilde{\sigma}^2/(\gamma k^2)$, $\eta^{(1)} = O\left(\frac{\gamma}{km\sigma_a}\right)$, *and* $\sigma_\xi^{(2)} \leq 1/k$, *then with probability* $\geq 1 - \exp(-\Theta(k))$ *over* (x,y), *we have* $yg^{(1)}(x; \sigma_\xi^{(2)}) < 1$. *Furthermore, for any* $i \in [2m]$, $\left|\langle w_i^{(1)}, x \rangle\right| = \left|\langle q_i^{(1)}, \phi \rangle\right| = O(\eta^{(1)}\tilde{\sigma}/\gamma)$, $\left|\langle (q_i^{(1)})_{-A}, \phi_{-A} \rangle\right| = O(\eta^{(1)}\tilde{\sigma}/\gamma)$, *and* $\left|b_i^{(1)} - b_{m+i}^{(1)}\right| = O(|\eta^{(1)} a_i^{(0)}|\epsilon_e)$.

Proof of Lemma 19. Note that $w_i^{(0)} = w_{m+i}^{(0)}, b_i^{(0)} = b_{m+i}^{(0)}$, and $a_i^{(0)} = a_{m+i}^{(0)}$. Then the gradient for w_{m+i} is the negation of that for w_{m+i}, the gradient for b_{m+i} is the negation of that for b_{m+i}, and the gradient for a_{m+i} is the same as that for a_{m+i}. With probability $\geq 1 - \exp(-\Theta(\max\{2p_o(D - k), k\}))$, among all $j \notin A$, we have that at most $2p_o(D - k) + k$ of ϕ_j are $(1 - p_o)/\tilde{\sigma}$, while the others are $-p_o/\tilde{\sigma}$. For data points with ϕ satisfying this, we have:

$$\left|g^{(1)}(x; \sigma_\xi^{(2)})\right| \tag{119}$$

$$= \left|\sum_{i=1}^{2m} a_i^{(1)} \mathbb{E}_{\xi^{(2)}} \sigma(\langle w_i^{(1)}, x \rangle + b_i^{(1)} + \xi_i^{(2)})\right| \tag{120}$$

$$= \left|\sum_{i=1}^m \left(a_i^{(1)} \mathbb{E}_{\xi^{(2)}} \sigma(\langle w_i^{(1)}, x \rangle + b_i^{(1)} + \xi_i^{(2)}) + a_{m+i}^{(1)} \mathbb{E}_{\xi^{(2)}} \sigma(\langle w_{m+i}^{(1)}, x \rangle + b_{m+i}^{(1)} + \xi_{m+i}^{(2)})\right)\right| \tag{121}$$

$$\leq \left|\sum_{i=1}^m \left(a_i^{(1)} \mathbb{E}_{\xi^{(2)}} \sigma(\langle w_i^{(1)}, x \rangle + b_i^{(1)} + \xi_i^{(2)}) + a_{m+i}^{(1)} \mathbb{E}_{\xi^{(2)}} \sigma(\langle w_i^{(1)}, x \rangle + b_i^{(1)} + \xi_i^{(2)})\right)\right| \tag{122}$$

$$+ \left|\sum_{i=1}^m \left(-a_{m+i}^{(1)} \mathbb{E}_{\xi^{(2)}} \sigma(\langle w_i^{(1)}, x \rangle + b_i^{(1)} + \xi_i^{(2)}) + a_{m+i}^{(1)} \mathbb{E}_{\xi^{(2)}} \sigma(\langle w_{m+i}^{(1)}, x \rangle + b_{m+i}^{(1)} + \xi_i^{(2)})\right)\right|. \tag{123}$$

Then we have

$$\left|g^{(1)}(x;\sigma_\xi^{(2)})\right| \le \sum_{i=1}^m \left|2\eta^{(1)}T_a\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_i^{(1)},x\rangle + b_i^{(1)} + \xi_i^{(2)})\right| \tag{124}$$

$$+ \sum_{i=1}^m \left|a_{m+i}^{(1)}\right|\left(\left|\langle w_i^{(1)} - w_{m+i}^{(1)},x\rangle\right| + \left|b_i^{(1)} - b_{m+i}^{(1)}\right|\right) \tag{125}$$

$$\le \sum_{i=1}^m \left|2\eta^{(1)}T_a\right|\left(\left|\langle w_i^{(1)},x\rangle + b_i^{(1)}\right| + \mathbb{E}_{\xi^{(2)}}\left|\xi_i^{(2)}\right|\right) \tag{126}$$

$$+ \sum_{i=1}^m \left|a_{m+i}^{(1)}\right|\left(\left|\langle w_i^{(1)} - w_{m+i}^{(1)},x\rangle\right| + \left|b_i^{(1)} - b_{m+i}^{(1)}\right|\right). \tag{127}$$

We have $|T_a| = O(\sigma_w\sqrt{\log(Dm/\delta)})$, and

$$\left|\langle w_i^{(1)},x\rangle\right| \le O(|\eta^{(1)}a_i^{(0)}|)(\beta\gamma/\tilde{\sigma} + \epsilon_{\mathrm{e}}/\tilde{\sigma})\frac{k}{\tilde{\sigma}} \tag{128}$$

$$+ O(|\eta^{(1)}a_i^{(0)}|\sigma_\phi^2\epsilon_{\mathrm{e}}\tilde{\sigma})((2p_{\mathrm{o}}(D-k)+k)(1-p_{\mathrm{o}})/\tilde{\sigma} + p_{\mathrm{o}}D/\tilde{\sigma}) \tag{129}$$

$$\le O(|\eta^{(1)}a_i^{(0)}|)(k\gamma/\tilde{\sigma}^2 + \epsilon_{\mathrm{e}}k/\tilde{\sigma}^2 + (k+p_{\mathrm{o}}D)\sigma_\phi^2\epsilon_{\mathrm{e}}) \tag{130}$$

$$\le O(\eta^{(1)}(1+p_{\mathrm{o}}\tilde{\sigma})/\gamma). \tag{131}$$

$$\left|b_i^{(1)}\right| \le \left|b_i^{(0)}\right| + \left|\eta^{(1)}a_i^{(0)}T_b\right| \tag{132}$$

$$\le \frac{\sqrt{\log(m/\delta)}}{k^2} + \left|\eta^{(1)}a_i^{(0)}\frac{\epsilon_{\mathrm{e}}}{\tilde{\sigma}}\right|. \tag{133}$$

$$\mathbb{E}_{\xi^{(2)}}\left|\xi_i^{(2)}\right| \le O(\sigma_\xi^{(2)}). \tag{134}$$

$$|a_{m+i}^{(1)}| \le |a_i^{(0)}| + |\eta^{(1)}T_a| \le |a_i^{(0)}| + O(\eta^{(1)}\sigma_w\sqrt{\log(Dm/\delta)}). \tag{135}$$

$$\left|\langle w_i^{(1)} - w_{m+i}^{(1)},x\rangle\right| = 2\left|\langle w_i^{(1)},x\rangle\right| = O(\eta^{(1)}(1+p_{\mathrm{o}}\tilde{\sigma})/\gamma). \tag{136}$$

$$\left|b_i^{(1)} - b_{m+i}^{(1)}\right| = 2|\eta^{(1)}a_i^{(0)}T_b| = O(|\eta^{(1)}a_i^{(0)}|\epsilon_{\mathrm{e}}). \tag{137}$$

Then we have

$$\left|g^{(1)}(x;\sigma_\xi^{(2)})\right| \le O\left(m\eta^{(1)}\sigma_w\sqrt{\log(Dm/\delta)}\right)\left(\frac{\eta^{(1)}}{\gamma} + \frac{\sqrt{\log(m/\delta)}}{k^2} + \left|\eta^{(1)}a_i^{(0)}\frac{\epsilon_{\mathrm{e}}}{\tilde{\sigma}}\right| + \sigma_\xi^{(2)}\right) \tag{138}$$

$$+ O\left(m(|a_i^{(0)}| + \eta^{(1)}\sigma_w\sqrt{\log(Dm/\delta)})\right)\left(\frac{\eta^{(1)}}{\gamma} + \left|\eta^{(1)}a_i^{(0)}\frac{\epsilon_{\mathrm{e}}}{\tilde{\sigma}}\right|\right) \tag{139}$$

$$= O\left(m\eta^{(1)}\sigma_w\frac{\log(Dm/\delta)}{k} + m|a_i^{(0)}|\left(\frac{\eta^{(1)}}{\gamma} + \left|\eta^{(1)}a_i^{(0)}\frac{\epsilon_{\mathrm{e}}}{\tilde{\sigma}}\right|\right)\right) \tag{140}$$

$$= O\left(m\eta^{(1)}\sigma_w\frac{\log(Dm/\delta)}{k} + m|a_i^{(0)}|\frac{\eta^{(1)}}{\gamma} + m\sigma_a\eta^{(1)}\frac{k}{\gamma}\right) \tag{141}$$

$$< 1. \tag{142}$$

Then $\left|yg^{(1)}(x;\sigma_\xi^{(2)})\right| < 1$. Finally, the statement on $\left|\langle (q_i^{(1)})_{-A}, \phi_{-A}\rangle\right|$ follows from a similar calculation on $\left|\langle w_i^{(1)},x\rangle\right| = \left|\langle q_i^{(1)},\phi\rangle\right|$. $\qquad\square$

We are now ready to analyze the gradients in the second gradient step.

Lemma 20 (Full version of Lemma 6). *Fix* $\delta \in (0,1)$ *and suppose* $w_i^{(0)} \in \mathcal{G}_w(\delta), b_i^{(0)} \in \mathcal{G}_b(\delta), a_i^{(0)} \in \mathcal{G}_a(\delta)$ *for all* $i \in [2m]$*. Let* $\epsilon_{e2} := O\left(\frac{\eta^{(1)}|a_i^{(0)}|k(\gamma+\epsilon_e)}{\tilde{\sigma}^2\sigma_\xi^{(2)}}\right) + \exp(-\Theta(k))$*. If*

$k = \Omega(\log^2(Dm/\delta))$ *and* $k = O(D)$, $\sigma_a \leq \tilde{\sigma}^2/(\gamma k^2)$, $\eta^{(1)} = O\left(\frac{\gamma}{km\sigma_a}\right)$, *and* $\sigma_\xi^{(2)} = 1/k^{3/2}$, *then*

$$\frac{\partial}{\partial w_i} L_\mathcal{D}(g^{(1)}; \sigma_\xi^{(2)}) = -a_i^{(1)} \sum_{j=1}^{D} M_j T_j \tag{143}$$

where T_j satisfies:

- *if $j \in A$, then $|T_j - \beta\gamma/\tilde{\sigma}| \leq O(\epsilon_{e2}/\tilde{\sigma} + \eta^{(1)}/\sigma_\xi^{(2)} + \eta^{(1)}|a_i^{(0)}|\epsilon_e/(\tilde{\sigma}\sigma_\xi^{(2)}))$, where $\beta \in [\Omega(1), 1]$ and depends only on $w_i^{(0)}, b_i^{(0)}$;*

- *if $j \notin A$, then $|T_j| \leq \frac{1}{\tilde{\sigma}}\exp(-\Theta(k)) + O(\sigma_\phi^2\epsilon_{e2}\tilde{\sigma})$.*

Proof of Lemma 20. Consider one neuron index i and omit the subscript i in the parameters. By Lemma 19, $\Pr[yg^{(1)}(x; \xi^{(2)}) > 1] \leq \exp(-\Theta(k))$. Let $\mathbb{I}_x = \mathbb{I}[yg^{(1)}(x; \xi^{(2)}) \leq 1]$.

$$\frac{\partial}{\partial w} L_\mathcal{D}(g^{(1)}; \sigma_\xi^{(2)}) \tag{144}$$

$$= -a^{(1)}\mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{y\mathbb{I}_x\mathbb{E}_{\xi^{(2)}}\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0,1)]x\right\} \tag{145}$$

$$= -a^{(1)}\sum_{j=1}^{D} M_j \underbrace{\mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{y\mathbb{I}_x\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0,1)]\phi_j\right\}}_{:=T_j}. \tag{146}$$

Let $T_{j1} := \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{y\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0,1)]\phi_j\right\}$. We have

$$|T_j - T_{j1}| \tag{147}$$

$$= \left|\mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{y(1 - \mathbb{I}_x)\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0,1)]\phi_j\right\}\right| \tag{148}$$

$$\leq \frac{1}{\tilde{\sigma}}\mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}|1 - \mathbb{I}_x| \tag{149}$$

$$\leq \frac{1}{\tilde{\sigma}}\exp(-\Theta(k)). \tag{150}$$

So it is sufficient to bound T_{j1}. For simplicity, we use q as a shorthand for $q_i^{(1)}$.

First, consider $j \in A$.

$$T_{j1} = \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{y\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0,1)]\phi_j\right\} \tag{151}$$

$$= \mathbb{E}_{\phi_A}\left\{y\phi_j \Pr_{\phi_{-A},\xi^{(2)}}\left[\langle\phi, q\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right]\right\}. \tag{152}$$

Let

$$I_a := \Pr_{\xi^{(2)}}\left[\langle\phi, q\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right], \tag{153}$$

$$I_a' := \Pr_{\xi^{(2)}}\left[\langle\phi_{-A}, q_{-A}\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right]. \tag{154}$$

By the property of the Gaussian $\xi^{(2)}$, that $|\langle\phi_A, q_A\rangle| = O(\frac{\eta^{(1)}|a_i^{(0)}|k(\gamma+\epsilon_e)}{\tilde{\sigma}^2})$, and that $|\langle\phi, q\rangle| = |\langle w_i^{(1)}, x\rangle| = O(\eta^{(1)}/\gamma) < O(1/k)$ and $|\langle\phi_{-A}, q_{-A}\rangle| = O(\eta^{(1)}/\gamma) < O(1/k)$, we have

$$|I_a - I_a'| \leq \left|\Pr_{\xi^{(2)}}\left[\langle\phi, q\rangle + b^{(1)} + \xi^{(2)} \geq 0\right] - \Pr_{\xi^{(2)}}\left[\langle\phi_{-A}, q_{-A}\rangle + b^{(1)} + \xi^{(2)} \geq 0\right]\right| \tag{155}$$

$$+ \Pr_{\xi^{(2)}}\left[\langle\phi, q\rangle + b^{(1)} + \xi^{(2)} \geq 1\right] + \Pr_{\xi^{(2)}}\left[\langle\phi_{-A}, q_{-A}\rangle + b^{(1)} + \xi^{(2)} \geq 1\right] \tag{156}$$

$$= O\left(\frac{\eta^{(1)}|a_i^{(0)}|k(\gamma+\epsilon_e)}{\tilde{\sigma}^2\sigma_\xi^{(2)}}\right) + \exp(-\Theta(k)) = O(\epsilon_{e2}). \tag{157}$$

This leads to

$$\left|T_{j1} - \mathbb{E}_{\phi_A, \phi_{-A}}\left\{y\phi_j I_a'\right\}\right| \tag{158}$$

$$\leq \mathbb{E}_{\phi_A}\left\{|y\phi_j|\left|\mathbb{E}_{\phi_{-A}}(I_a - I_a')\right|\right\} \tag{159}$$

$$\leq O(\epsilon_{e2})\mathbb{E}_{\phi_A}\left\{|y\phi_j|\right\} \tag{160}$$

$$\leq O(\epsilon_{e2}/\tilde{\sigma}) \tag{161}$$

where the last step is from Lemma 14. Furthermore,

$$\mathbb{E}_{\phi_A, \phi_{-A}}\left\{y\phi_j I_a'\right\} \tag{162}$$

$$= \mathbb{E}_{\phi_A}\left\{y\phi_j\right\}\mathbb{E}_{\phi_{-A}}[I_a'] \tag{163}$$

$$= \mathbb{E}_{\phi_A}\left\{y\phi_j\right\}\Pr_{\phi_{-A}, \xi^{(2)}}\left[\langle\phi_{-A}, q_{-A}\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right]. \tag{164}$$

By Lemma 19, we have $|\langle\phi_{-A}, q_{-A}\rangle| \leq O(\eta^{(1)}\tilde{\sigma}/\gamma)$. Also, $|b^{(1)} - b^{(0)}| \leq O(\eta^{(1)}|a_i^{(0)}|\epsilon_e)$. By the property of $\xi^{(2)}$,

$$\left|\Pr_{\xi^{(2)}}\left[\langle\phi_{-A}, q_{-A}\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right] - \Pr_{\xi^{(2)}}\left[b^{(0)} + \xi^{(2)} \in (0,1)\right]\right| \tag{165}$$

$$\leq O(\eta^{(1)}\tilde{\sigma}/(\gamma\sigma_\xi^{(2)})) + O(\eta^{(1)}|a_i^{(0)}|\epsilon_e/\sigma_\xi^{(2)}). \tag{166}$$

On the other hand,

$$\beta := \Pr_{\phi_{-A}, \xi^{(2)}}\left[b^{(0)} + \xi^{(2)} \in (0,1)\right] = \Pr_{\xi^{(2)}}\left[\xi^{(2)} \in (-b^{(0)}, 1 - b^{(0)})\right] \tag{167}$$

$$= \Omega(1) \tag{168}$$

and β only depends on $b^{(0)}$. By Lemma 14, $\mathbb{E}_{\phi_A}\left\{y\phi_j\right\} = \gamma/\tilde{\sigma}$. Therefore,

$$|T_{j1} - \beta\gamma/\tilde{\sigma}| \leq O(\epsilon_{e2}/\tilde{\sigma}) + O(\eta^{(1)}/\sigma_\xi^{(2)}) + O(\eta^{(1)}|a_i^{(0)}|\epsilon_e/(\tilde{\sigma}\sigma_\xi^{(2)})). \tag{169}$$

Now, consider $j \notin A$. Let B denote $A \cup \{j\}$.

$$T_{j1} = \mathbb{E}_{(x,y)\sim\mathcal{D}, \xi^{(2)}}\left\{y\phi_j\mathbb{I}\left[\langle\phi, q\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right]\right\} \tag{170}$$

$$= \mathbb{E}_{\phi_B}\mathbb{E}_{\phi_{-B}, \xi^{(2)}}\left\{y\phi_j\mathbb{I}\left[\langle\phi, q\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right]\right\} \tag{171}$$

$$= \mathbb{E}_{\phi_B}\left\{y\phi_j\Pr_{\phi_{-B}, \xi^{(2)}}\left[\langle\phi, q\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right]\right\}. \tag{172}$$

Let

$$I_b := \Pr_{\xi^{(2)}}\left[\langle\phi, q\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right], \tag{173}$$

$$I_b' := \Pr_{\xi^{(2)}}\left[\langle\phi_{-B}, q_{-B}\rangle + b^{(1)} + \xi^{(2)} \in (0,1)\right]. \tag{174}$$

Similar as above, we have $|I_b - I_b'| \leq \epsilon_{e2}$. Then by Lemma 14,

$$\left|T_{j1} - \mathbb{E}_{\phi_B, \phi_{-B}}\left\{y\phi_j I_b'\right\}\right| \tag{175}$$

$$\leq \mathbb{E}_{\phi_B}\left\{|y\phi_j||\mathbb{E}_{\phi_{-B}}(I_b - I_b')|\right\} \tag{176}$$

$$\leq O(\epsilon_{e2})\mathbb{E}_{\phi_j}\left\{|\phi_j|\right\} \tag{177}$$

$$\leq O(\epsilon_e) \times O(\sigma_\phi^2\tilde{\sigma}) \tag{178}$$

$$= O(\sigma_\phi^2\epsilon_{e2}\tilde{\sigma}). \tag{179}$$

Furthermore,

$$\mathbb{E}_{\phi_B, \phi_{-B}}\left\{y\phi_j I_b'\right\} = \mathbb{E}_{\phi_A}\left\{y\right\}\mathbb{E}_{\phi_j}\left\{\phi_j\right\}\mathbb{E}_{\phi_{-B}}[I_b'] = 0. \tag{180}$$

Therefore,

$$|T_{j1}| \leq O(\sigma_\phi^2\epsilon_{e2}\tilde{\sigma}). \tag{181}$$

$$\square$$

Lemma 21. *Under the same assumptions as in Lemma 20,*

$$\frac{\partial}{\partial b} L_{\mathcal{D}}(g^{(1)}; \sigma_\xi^{(2)}) = -a_i^{(1)} T_b \tag{182}$$

where $|T_b| \le \exp(-\Omega(k)) + O(\epsilon_{e2})$.

Proof of Lemma 21. Consider one neuron index i and omit the subscript i in the parameters. By Lemma 19, $\Pr[yg^{(1)}(x; \xi^{(2)}) > 1] \le \exp(-\Omega(k))$. Let $\mathbb{I}_x = \mathbb{I}[yg^{(1)}(x; \xi^{(2)}) \le 1]$.

$$\frac{\partial}{\partial b} L_{\mathcal{D}}(g^{(1)}; \sigma_\xi^{(2)}) \tag{183}$$

$$= -a^{(1)} \underbrace{\mathbb{E}_{(x,y)\sim\mathcal{D}} \left\{ y \mathbb{I}_x \mathbb{E}_{\xi^{(2)}} \mathbb{I}[\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)} \in (0,1)] \right\}}_{:=T_b}. \tag{184}$$

Let $T_{b1} := \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}} \left\{ y \mathbb{I}[\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)} \in (0,1)] \right\}$. We have

$$|T_b - T_{b1}| \tag{185}$$

$$= \left| \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}} \left\{ y(1 - \mathbb{I}_x) \mathbb{I}[\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)} \in (0,1)] \right\} \right| \tag{186}$$

$$\le \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}} |1 - \mathbb{I}_x| \tag{187}$$

$$\le \exp(-\Omega(k)). \tag{188}$$

So it is sufficient to bound T_{b1}. For simplicity, we use q as a shorthand for $q_i^{(1)}$.

$$T_{b1} = \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}} \left\{ y \mathbb{I} \left[\langle \phi, q \rangle + b^{(1)} + \xi^{(2)} \in (0,1) \right] \right\} \tag{189}$$

$$= \mathbb{E}_{\phi_A} \mathbb{E}_{\phi_{-A},\xi^{(2)}} \left\{ y \mathbb{I} \left[\langle \phi, q \rangle + b^{(1)} + \xi^{(2)} \in (0,1) \right] \right\} \tag{190}$$

$$= \mathbb{E}_{\phi_A} \left\{ y \Pr_{\phi_{-A},\xi^{(2)}} \left[\langle \phi, q \rangle + b^{(1)} + \xi^{(2)} \in (0,1) \right] \right\}. \tag{191}$$

Let

$$I_b := \Pr_{\xi^{(2)}} \left[\langle \phi, q \rangle + b^{(1)} + \xi^{(2)} \in (0,1) \right], \tag{192}$$

$$I_b' := \Pr_{\xi^{(2)}} \left[\langle \phi_{-A}, q_{-A} \rangle + b^{(1)} + \xi^{(2)} \in (0,1) \right]. \tag{193}$$

Similar as in Lemma 6, we have $|I_b - I_b'| \le \epsilon_{e2}$. Then by Lemma 14,

$$\left| T_{b1} - \mathbb{E}_{\phi_A,\phi_{-A}} \{y I_b'\} \right| \tag{194}$$

$$\le \mathbb{E}_{\phi_A} \left\{ |\mathbb{E}_{\phi_{-A}}(I_b - I_b')| \right\} \tag{195}$$

$$\le O(\epsilon_{e2}). \tag{196}$$

Furthermore,

$$\mathbb{E}_{\phi_A,\phi_{-A}} \{y I_b'\} = \mathbb{E}_{\phi_A} \{y\} \mathbb{E}_{\phi_{-A}}[I_b'] = 0. \tag{197}$$

Therefore, $|T_{b1}| \le O(\epsilon_{e2})$ and the statement follows. $\square$

Lemma 22. *Under the same assumptions as in Lemma 20,*

$$\frac{\partial}{\partial a_i} L_{\mathcal{D}}(g^{(1)}; \sigma_\xi^{(2)}) = -T_a \tag{198}$$

where $|T_a| = O(\eta^{(1)} \tilde{\sigma}/\gamma) + \exp(-\Omega(k))\text{poly}(Dm)$.

Proof of Lemma 22. Consider one neuron index i and omit the subscript i in the parameters. By Lemma 19, $\Pr[yg^{(1)}(x; \xi^{(2)}) > 1] \leq \exp(-\Omega(k))$. Let $\mathbb{I}_x = \mathbb{I}[yg^{(1)}(x; \xi^{(2)}) \leq 1]$.

$$\frac{\partial}{\partial a} L_{\mathcal{D}}(g^{(1)}; \sigma_{\xi}^{(2)}) \tag{199}$$

$$= -\underbrace{\mathbb{E}_{(x,y)\sim\mathcal{D}} \left\{ y\mathbb{I}_x \mathbb{E}_{\xi^{(2)}} \sigma(\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)}) \right\}}_{:=T_a}. \tag{200}$$

Let $T_{a1} := \mathbb{E}_{(x,y)\sim\mathcal{D}} \left\{ y\mathbb{E}_{\xi^{(2)}} \sigma(\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)}) \right\}$. We have

$$|T_a - T_{a1}| \tag{201}$$

$$= \left| \mathbb{E}_{(x,y)\sim\mathcal{D}} \left\{ y(1 - \mathbb{I}_x)\mathbb{E}_{\xi^{(2)}} \sigma(\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)}) \right\} \right| \tag{202}$$

$$\leq \mathbb{E}_{(x,y)\sim\mathcal{D}, \xi^{(2)}} |1 - \mathbb{I}_x| \tag{203}$$

$$\leq \exp(-\Omega(k)). \tag{204}$$

So it is sufficient to bound T_{a1}. For simplicity, we use q as a shorthand for $q_i^{(1)}$.

Let ϕ'_A be an independent copy of ϕ_A, ϕ' be the vector obtained by replacing in ϕ the entries ϕ_A with ϕ'_A, and let $x' = M\phi'$ and its label is y'. Then

$$|T_{a1}| := \left| \mathbb{E}_{\phi_A} \left\{ y\mathbb{E}_{\phi_{-A}, \xi^{(2)}} \sigma(\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)}) \right\} \right| \tag{205}$$

$$\leq \frac{1}{2} \left| \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \xi^{(2)}} \sigma(\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(1)}) | y = 1 \right\} \right. \tag{206}$$

$$\left. - \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \xi^{(2)}} \sigma(\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)}) | y = -1 \right\} \right| \tag{207}$$

$$\leq \frac{1}{2} \left| \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \xi^{(2)}} \sigma(\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)}) | y = 1 \right\} \right. \tag{208}$$

$$\left. - \mathbb{E}_{\phi'_A} \left\{ \mathbb{E}_{\phi_{-A}, \xi^{(2)}} \sigma(\langle w^{(1)}, x' \rangle + b^{(1)} + \xi^{(2)}) | y' = -1 \right\} \right| \tag{209}$$

$$\leq \frac{1}{2} \mathbb{E}_{\phi_A, \phi'_A} \left\{ \mathbb{E}_{\phi_{-A}} \left| \langle w^{(1)}, x \rangle - \langle w^{(1)}, x' \rangle \right| | y = 1, y' = -1 \right\} \tag{210}$$

$$\leq \frac{1}{2} \mathbb{E}_{\phi_{-A}} \left(\mathbb{E}_{\phi_A} \left\{ \left| \langle w^{(1)}, x \rangle \right| | y = 1 \right\} + \mathbb{E}_{\phi'_A} \left\{ \left| \langle w^{(1)}, x' \rangle \right| | y' = -1 \right\} \right) \tag{211}$$

$$\leq \mathbb{E}_{\phi_{-A}, \phi_A} \left| \langle w^{(1)}, x \rangle \right| \tag{212}$$

$$= \mathbb{E}_x \left| \langle w^{(1)}, x \rangle \right| \tag{213}$$

$$= O(\eta^{(1)} \tilde{\sigma} / \gamma) + \exp(-\Omega(k)) \times D \times \|q^{(1)}\|_\infty \|\phi\|_\infty \tag{214}$$

$$= O(\eta^{(1)} \tilde{\sigma} / \gamma) + \exp(-\Omega(k)) \frac{D|\eta^{(1)} a^{(0)}|(\gamma + \epsilon_e)}{\tilde{\sigma}^2} \tag{215}$$

$$= O(\eta^{(1)} \tilde{\sigma} / \gamma) + \exp(-\Omega(k)) \text{poly}(Dm) \tag{216}$$

where the fourth step follows from that σ is 1-Lipschitz, the third to the last step from Lemma 19, and the second to the last step from Lemma 18. $\square$

With the above lemmas about the gradients, we are now ready to show that at the end of the second step, we get a good set of features for accurate prediction.

Lemma 23. *Set*

$$\eta^{(1)} = \frac{\gamma^2 \tilde{\sigma}}{km^3}, \lambda_a^{(1)} = 0, \lambda_w^{(1)} = 1/(2\eta^{(1)}), \sigma_\xi^{(1)} = 1/k^{3/2}, \tag{217}$$

$$\eta^{(2)} = 1, \lambda_a^{(2)} = \lambda_w^{(2)} = 1/(2\eta^{(2)}), \sigma_\xi^{(2)} = 1/k^{3/2}. \tag{218}$$

Fix $\delta \in (0, O(1/k^3))$. If $p_o = \Omega(k^2/D)$, $k = \Omega\left(\log^2\left(\frac{Dm}{\delta\gamma}\right)\right)$, and $m \geq \max\{\Omega(k^4), D\}$, then with probability at least $1 - \delta$ over the initialization, there exist $\tilde{a}_i$'s such that $\tilde{g}(x) := \sum_{i=1}^{2m} \tilde{a}_i \sigma(\langle w_i^{(2)}, x \rangle + b_i^{(2)})$ satisfies $L_{\mathcal{D}}(\tilde{g}) = 0$. Furthermore, $\|\tilde{a}\|_0 = O(m/k)$, $\|\tilde{a}\|_\infty = O(k^5/m)$, and $\|\tilde{a}\|_2^2 = O(k^9/m)$. Finally, $\|a^{(2)}\|_\infty = O\left(\frac{1}{km^2}\right)$, $\|w_i^{(2)}\|_2 = O(\tilde{\sigma}/k)$, and $|b_i^{(2)}| = O(1/k^2)$ for all $i \in [2m]$.

Proof of Lemma 23. By Lemma 4, there exists a network $g^*(x) = \sum_{\ell=1}^{3(k+1)} a_\ell^* \sigma(\langle w_\ell^*, x \rangle + b_\ell^*)$ satisfying $g^*(x)$ for all $(x, y) \sim \mathcal{D}$. Furthermore, $|a_i^*| \leq 32k$, $1/(32k) \leq |b_i^*| \leq 1/2$, $w_i^* = \tilde{\sigma} \sum_{j \in A} M_j/(4k)$, and $|\langle w_i^*, x \rangle + b_i^*| \leq 1$ for any $i \in [n]$ and $(x, y) \sim \mathcal{D}$. Now we fix an ℓ, and show that with high probability there is a neuron in $g^{(2)}$ that can approximate the ℓ-th neuron in g^*.

By Lemma 10, with probability $1 - 2\delta$ over $w_i^{(0)}$'s, they are all in $\mathcal{G}_w(\delta)$; with probability $1 - \delta$ over $a_i^{(0)}$'s, they are all in $\mathcal{G}_a(\delta)$; with probability $1 - \delta$ over $b_i^{(0)}$'s, they are all in $\mathcal{G}_b(\delta)$. Under these events, by Lemma 18, Lemma 20 and 21, for any neuron $i \in [2m]$, we have

$$w_i^{(2)} = a_i^{(1)} \sum_{j=1}^{D} M_j T_j, \tag{219}$$

$$b_i^{(2)} = b_i^{(1)} + a_i^{(1)} T_b. \tag{220}$$

where

- if $j \in A$, then $|T_j - \beta\gamma/\tilde{\sigma}| \leq \epsilon_{w1} := O(\epsilon_{e2}/\tilde{\sigma} + \eta^{(1)}/\sigma_\xi^{(2)} + \eta^{(1)}|a_i^{(0)}|\epsilon_e/(\tilde{\sigma}\sigma_\xi^{(2)}))$, where $\beta \in [\Omega(1), 1]$ and depends only on $w_i^{(0)}, b_i^{(0)}$;

- if $j \notin A$, then $|T_j| \leq \epsilon_{w2} := \frac{1}{\tilde{\sigma}} \exp(-\Theta(k)) + O(\sigma_\phi^2 \epsilon_{e2} \tilde{\sigma})$.

- $|T_b| \leq \epsilon_b := \frac{1}{\tilde{\sigma}} \exp(-\Theta(k)) + O(\epsilon_{e2})$.

Given the initialization, with probability $\Omega(1)$ over $b_i^{(0)}$, we have

$$|b_i^{(0)}| \in \left[\frac{1}{2k^2}, \frac{2}{k^2}\right], \text{sign}(b_i^{(0)}) = \text{sign}(b_\ell^*). \tag{221}$$

Finally, since $\frac{4k|b_\ell^*|\beta\gamma}{|b_i^{(0)}|\tilde{\sigma}^2} \in [\Omega(k^2\gamma/\tilde{\sigma}^2), O(k^3\gamma/\tilde{\sigma}^2)]$ and depends only on $w_i^{(0)}, b_i^{(0)}$, we have that for $\epsilon_a = \Theta(1/k^2)$, with probability $\Omega(\epsilon_a) > \delta$ over $a_i^{(0)}$,

$$\left|\frac{4k|b_\ell^*|\beta\gamma}{|b_i^{(0)}|\tilde{\sigma}^2} a_i^{(0)} - 1\right| \leq \epsilon_a, \quad |a_i^{(0)}| = O\left(\frac{\tilde{\sigma}^2}{k^2\gamma}\right). \tag{222}$$

Let $n_a = \epsilon_a m/4$. For the given value of m, by (219)-(222) we have with probability $\geq 1 - 5\delta$ over the initialization, for each ℓ there is a different set of neurons $I_\ell \subseteq [m]$ with $|I_\ell| = n_a$ and such that for each $i_\ell \in I_\ell$,

$$|b_{i_\ell}^{(0)}| \in \left[\frac{1}{2k^2}, \frac{2}{k^2}\right], \quad \text{sign}(b_{i_\ell}^{(0)}) = \text{sign}(b_\ell^*), \tag{223}$$

$$\left|\frac{4k|b_\ell^*|\beta\gamma}{|b_{i_\ell}^{(0)}|\tilde{\sigma}^2} a_{i_\ell}^{(0)} - 1\right| \leq \epsilon_a, \quad |a_{i_\ell}^{(0)}| = O\left(\frac{\tilde{\sigma}^2}{k^2\gamma}\right). \tag{224}$$

We also have

$$\left| \langle w_{i_\ell}^{(2)}, x \rangle - \frac{a_{i_\ell}^{(0)} \beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| \tag{225}$$

$$\leq \left| \langle w_{i_\ell}^{(2)}, x \rangle - \frac{a_{i_\ell}^{(1)} \beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| + \left| \frac{a_{i_\ell}^{(1)} \beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle - \frac{a_{i_\ell}^{(0)} \beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| \tag{226}$$

$$= \left| a_{i_\ell}^{(1)} \sum_{j=1}^{D} T_j \phi_j - \frac{a_{i_\ell}^{(1)} \beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \phi_j \right| + \left| a_{i_\ell}^{(1)} - a_{i_\ell}^{(0)} \right| \left| \frac{\beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \phi_j \right| \tag{227}$$

$$= \left| a_{i_\ell}^{(1)} \right| \left| \sum_{j=1}^{D} T_j \phi_j - \frac{\beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \phi_j \right| + \left| a_{i_\ell}^{(1)} - a_{i_\ell}^{(0)} \right| \left| \frac{\beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \phi_j \right|. \tag{228}$$

We have $\left| a_{i_\ell}^{(1)} - a_{i_\ell}^{(0)} \right| = O(\eta^{(1)} \sigma_w \sqrt{\log(Dm/\delta)})$, and

$$\left| \sum_{j=1}^{D} T_j \phi_j - \frac{\beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \phi_j \right| \leq \left| \sum_{j \in A} (T_j - \frac{\beta\gamma}{\tilde{\sigma}}) \phi_j \right| + \left| \sum_{j \notin A} T_j \phi_j \right| \tag{229}$$

$$\leq O(k\epsilon_{w1}/\tilde{\sigma}) + O(D\epsilon_{w2}/\tilde{\sigma}) =: \epsilon_\phi. \tag{230}$$

For the given values of parameters, we have

$$\epsilon_{e2} = O\left(\frac{\gamma}{m^2} \right), \tag{231}$$

$$\epsilon_{w1} = O\left(\frac{k\gamma}{m^2 \tilde{\sigma}} + \frac{\gamma \epsilon_e}{k m^2} \right), \tag{232}$$

$$\epsilon_{w2} = O\left(\frac{\gamma}{m^2 \tilde{\sigma}} \right), \tag{233}$$

$$\epsilon_b = O\left(\frac{\gamma}{m^2} \right), \tag{234}$$

$$\epsilon_\phi = O\left(\frac{k^2 \gamma}{m^2 \tilde{\sigma}^2} + \frac{\gamma \epsilon_e}{m^2 \tilde{\sigma}} + \frac{\gamma}{m \tilde{\sigma}^2} \right). \tag{235}$$

Therefore,

$$\left| \langle w_{i_\ell}^{(2)}, x \rangle - \frac{a_{i_\ell}^{(0)} \beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| \tag{236}$$

$$\leq \left| a_{i_\ell}^{(1)} \right| \epsilon_\phi + \left| a_{i_\ell}^{(1)} - a_{i_\ell}^{(0)} \right| \frac{k\gamma}{\tilde{\sigma}^2} \tag{237}$$

$$\leq O\left(\frac{\tilde{\sigma}^2}{k^2 \gamma} + \eta^{(1)} \sigma_w \sqrt{\log(Dm/\delta)} \right) \left(\frac{k^2 \gamma}{m^2 \tilde{\sigma}^2} + \frac{\gamma \epsilon_e}{m^2 \tilde{\sigma}} + \frac{\gamma}{m \tilde{\sigma}^2} \right) \tag{238}$$

$$+ O\left(\eta^{(1)} \sigma_w \sqrt{\log(Dm/\delta)} \right) \frac{k\gamma}{\tilde{\sigma}^2} \tag{239}$$

$$\leq O\left(\frac{1}{m} \right). \tag{240}$$

We also have by Lemma 18 and 21:

$$|b_{i_\ell}^{(2)} - b_{i_\ell}^{(0)}| \leq O\left(\eta^{(1)} |a_{i_\ell}^{(0)}| \epsilon_e + |a_{i_\ell}^{(1)}| \left(\frac{1}{\tilde{\sigma}} \exp(-\Theta(k)) + \epsilon_{e2} \right) \right) \leq O\left(\frac{1}{m} \right). \tag{241}$$

Now, construct $\tilde{a}$ such that $\tilde{a}_{i_\ell} = \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|n_a}$ for each ℓ and each $i_\ell \in I_\ell$, and $\tilde{a}_i = 0$ elsewhere. Then

$$|\tilde{g}(x) - 2g^*(x)| \tag{242}$$

$$= \left| \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \tilde{a}_{i_\ell} \sigma \left(\langle w_{i_\ell}^{(2)}, x \rangle + b_{i_\ell}^{(2)} \right) - \sum_{\ell=1}^{3(k+1)} 2a_\ell^* \sigma \left(\langle w_\ell^*, x \rangle + b_\ell^* \right) \right| \tag{243}$$

$$= \left| \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|n_a} \sigma \left(\langle w_{i_\ell}^{(2)}, x \rangle + b_{i_\ell}^{(2)} \right) - \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|n_a} \sigma \left(\frac{|b_{i_\ell}^{(0)}|}{|b_\ell^*|} \langle w_\ell^*, x \rangle + b_{i_\ell}^{(0)} \right) \right| \tag{244}$$

$$\leq \left| \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \frac{1}{n_a} \left(\frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|} \sigma \left(\langle w_{i_\ell}^{(2)}, x \rangle + b_{i_\ell}^{(2)} \right) - \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|} \sigma \left(\frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle + b_{i_\ell}^{(0)} \right) \right) \right| \tag{245}$$

$$+ \left| \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \frac{1}{n_a} \left(\frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|} \sigma \left(\frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle + b_{i_\ell}^{(0)} \right) - \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|} \sigma \left(\frac{|b_{i_\ell}^{(0)}|}{|b_\ell^*|} \langle w_\ell^*, x \rangle + b_{i_\ell}^{(0)} \right) \right) \right| \tag{246}$$

$$\leq 3(k+1) \max_\ell \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|} O\left(\frac{1}{m}\right) + \tag{247}$$

$$3(k+1) \max_\ell \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|} \frac{\tilde{\sigma}|b_{i_\ell}^{(0)}|}{4k|b_\ell^*|} \left| \frac{4ka_{i_\ell}^{(0)}\beta\gamma|b_\ell^*|}{\tilde{\sigma}^2|b_{i_\ell}^{(0)}|} - 1 \right| \frac{k}{\tilde{\sigma}} \tag{248}$$

$$= O\left(\frac{k^4}{m} + k^2\epsilon_a\right) \tag{249}$$

$$\leq 1. \tag{250}$$

Here the second equation follows from that σ is positive-homogeneous in $[0, 1]$, $|\langle w_\ell^*, x \rangle + b_\ell^*| \leq 1$, $|b_{i_\ell}^{(0)}|/|b_\ell^*| \leq 1$. This guarantees $y\tilde{g}(x) \geq 1$. Changing the scaling of δ leads to the statement.

Finally, the bounds on $\tilde{a}$ follow from the above calculation. The bound on $\|a^{(2)}\|_2$ follows from Lemma 22, and those on $\|w_i^{(2)}\|_2$ and $\|b_i^{(2)}\|_2$ follow from (219)(220) and the bounds on $a_i^{(1)}$ and $b_i^{(1)}$ in Lemma 18. $\qquad\square$

B.6 CLASSIFIER LEARNING STAGE

Once we have a set of good features, we are now ready to prove that the later steps will learn an accurate classifier. The intuition is that the first layer's weights do not change much and the second layer's weights get updated till achieving good accuracy. In particular, we will employ the online optimization technique from Daniely & Malach (2020).

We begin by showing that the first layer's weights do not change too much.

Lemma 24. *Assume the same conditions as in Lemma 23. Suppose for $t > 2$, $\lambda_a^{(t)} = \lambda_w^{(t)} = \lambda$, $\eta^{(t)} = \eta$ for some $\lambda, \eta \in (0, 1)$, and $\sigma_\xi^{(t)} = 0$. Then for any $t > 2$ and $i \in [2m]$,*

$$|a_i^{(t)}| \leq \eta t + O\left(\frac{1}{km^2}\right), \tag{251}$$

$$\|w_i^{(t)} - w_i^{(2)}\|_2 \leq O\left(\frac{t\eta\lambda\tilde{\sigma}}{k}\right) + \eta^2 t^2 + O\left(\frac{t}{km^2}\right), \tag{252}$$

$$|b_i^{(t)} - b_i^{(2)}| \leq O\left(\frac{t\eta\lambda}{k^2}\right) + \eta^2 t^2 + O\left(\frac{t}{km^2}\right). \tag{253}$$

Proof of Lemma 24. First, we bound the size of $|a_i^{(t)}|$:

$$|a_i^{(t)}| = \left| (1 - 2\eta\lambda)a_i^{(t-1)} - \eta\frac{\partial}{\partial a_i}L_{\mathcal{D}}(g^{(t-1)}) \right| \tag{254}$$

$$\leq \left| (1 - 2\eta\lambda)a_i^{(t-1)} - \eta\mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{ y\mathbb{I}[yg^{(t-1)}(x) \leq 1]\sigma(\langle w_i^{(t-1)}, x\rangle + b_i^{(t-1)}) \right\} \right| \tag{255}$$

$$\leq |a_i^{(t-1)}| + \eta \tag{256}$$

which leads to

$$|a_i^{(t)}| \leq \eta t + |a_i^{(2)}| \tag{257}$$

where $|a_i^{(2)}| = O\left(\frac{1}{km^2}\right)$. We are now to bound the change of $w_i^{(t)}$ and $b_i^{(t)}$.

$$\|w_i^{(t)} - w_i^{(2)}\|_2 \tag{258}$$

$$= \left\| (1 - 2\eta\lambda)w_i^{(t-1)} - \eta\frac{\partial}{\partial w_i}L_{\mathcal{D}}(g^{(t-1)}) - w_i^{(2)} \right\|_2 \tag{259}$$

$$\leq \left\| (1 - 2\eta\lambda)w_i^{(t-1)} \right. \tag{260}$$

$$\left. + \eta a_i^{(t-1)}\mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{ y\mathbb{I}[yg^{(t-1)}(x) \leq 1]\mathbb{I}[\langle w_i^{(t-1)}, x\rangle + b_i^{(t-1)} \in (0,1)]x \right\} - w_i^{(2)} \right\|_2 \tag{261}$$

$$\leq \left\| (1 - 2\eta\lambda)w_i^{(t-1)} - w_i^{(2)} \right\|_2 \tag{262}$$

$$+ \eta\left\| a_i^{(t-1)}\mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{ y\mathbb{I}[yg^{(t-1)}(x) \leq 1]\mathbb{I}[\langle w_i^{(t-1)}, x\rangle + b_i^{(t-1)} \in (0,1)]x \right\} \right\|_2 \tag{263}$$

$$\leq (1 - 2\eta\lambda)\left\| w_i^{(t-1)} - w_i^{(2)} \right\|_2 + 2\eta\lambda\left\| w_i^{(2)} \right\|_2 + \eta\left| a_i^{(t-1)} \right| \tag{264}$$

leading to

$$\|w_i^{(t)} - w_i^{(2)}\|_2 \leq 2t\eta\lambda\left\| w_i^{(2)} \right\|_2 + \eta^2 t^2 + t|a_i^{(2)}|. \tag{265}$$

Note that $\|w_i^{(2)}\|_2 = O(\tilde{\sigma}/k)$.

$$|b_i^{(t)} - b_i^{(2)}| = \left| b_i^{(t-1)} - \eta\frac{\partial}{\partial b_i}L_{\mathcal{D}}(g^{(t-1)}) - b_i^{(2)} \right| \tag{266}$$

$$\leq \left| b_i^{(t-1)} - b_i^{(2)} \right| \tag{267}$$

$$+ \eta\left| a_i^{(t-1)}\mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{ y\mathbb{I}[yg^{(t-1)}(x) \leq 1]\mathbb{I}[\langle w_i^{(t-1)}, x\rangle + b_i^{(t-1)} \in (0,1)] \right\} \right| \tag{268}$$

$$\leq \left| b_i^{(t-1)} - b_i^{(2)} \right| + \eta\left| a_i^{(t-1)} \right| \tag{269}$$

leading to

$$|b_i^{(t)} - b_i^{(2)}| \leq \eta^2 t^2 + t|a_i^{(2)}|. \tag{270}$$

Note that $\left| b_i^{(2)} \right| = O(1/k^2)$. $\qquad\square$

Lemma 25. *Assume the same conditions as in Lemma 24. Let $g_{\tilde{a}}^{(t)}(x) = \sum_{i=1}^m \tilde{a}_i\sigma(\langle w_i^{(t)}, x\rangle + b_i^{(t)})$. Then*

$$|\ell(g_{\tilde{a}}^{(t)}(x), y) - \ell(g_{\tilde{a}}^{(2)}(x), y)| \leq \|\tilde{a}\|_2\sqrt{\|\tilde{a}\|_0}\left(O\left(\frac{t\eta\lambda\tilde{\sigma}}{k}\right) + \eta^2 t^2 + O\left(\frac{t}{km^2}\right) \right). \tag{271}$$

Proof of Lemma 25. It follows from that

$$|\ell(g_{\tilde{a}}^{(t)}(x), y) - \ell(g_{\tilde{a}}^{(2)}(x))| \tag{272}$$

$$\leq |g_{\tilde{a}}^{(t)}(x) - g_{\tilde{a}}^{(2)}(x)| \tag{273}$$

$$\leq \|\tilde{a}\|_2 \sqrt{\|\tilde{a}\|_0} \max_{i \in [2m]} \left| \sigma(\langle w_i^{(t)}, x \rangle + b_i^{(t)}) - \sigma(\langle w_i^{(2)}, x \rangle + b_i^{(2)}) \right| \tag{274}$$

$$\leq \|\tilde{a}\|_2 \sqrt{\|\tilde{a}\|_0} \max_{i \in [2m]} \left(\left| \langle w_i^{(t)} - w_i^{(2)}, x \rangle \right| + \left| b_i^{(t)} - b_i^{(2)} \right| \right). \tag{275}$$

and Lemma 24. □

B.7 PROOF OF THEOREM 1

Based on the above lemmas, following the same argument as in the proof of Theorem 2 in Daniely & Malach (2020), we get our main theorem.

Theorem 26 (Full version of Theorem 1). *Set*

$$\eta^{(1)} = \frac{\gamma^2 \tilde{\sigma}^2}{km^3}, \lambda_a^{(1)} = 0, \lambda_w^{(1)} = 1/(2\eta^{(1)}), \sigma_\xi^{(1)} = 1/k^2, \tag{276}$$

$$\eta^{(2)} = 1, \lambda_a^{(2)} = \lambda_w^{(2)} = 1/(2\eta^{(2)}), \sigma_\xi^{(2)} = 1/k^2, \tag{277}$$

$$\eta^{(t)} = \eta = \frac{k^2}{Tm^{1/3}}, \lambda_a^{(t)} = \lambda_w^{(t)} = \lambda \leq \frac{k^3}{\tilde{\sigma}m^{1/3}}, \sigma_\xi^{(t)} = 0, \text{ for } 2 < t \leq T. \tag{278}$$

For any $\delta \in (0, 1)$, if $p_o = \Omega(k^2/D)$, $k = \Omega\left(\log^2\left(\frac{Dm}{\delta\gamma}\right)\right)$, $m \geq \max\{\Omega(k^4), D\}$, then we have for any $\mathcal{D} \in \mathcal{F}_\Xi$, with probability at least $1 - \delta$, there exists $t \in [T]$ such that

$$\Pr[\text{sign}(g^{(t)}(x)) \neq y] \leq L_\mathcal{D}(g^{(t)}) = O\left(\frac{k^8}{m^{2/3}} + \frac{k^3 T}{m^2} + \frac{k^2 m^{2/3}}{T}\right). \tag{279}$$

Consequently, for any $\epsilon \in (0, 1)$, if $T = m^{4/3}$, and $m \geq \max\{\Omega(k^{12}/\epsilon^{3/2}), D\}$, then

$$\Pr[\text{sign}(g^{(t)}(x)) \neq y] \leq L_\mathcal{D}(g^{(t)}) \leq \epsilon. \tag{280}$$

Proof of Theorem 1. Consider $\tilde{L}_\mathcal{D}(g^{(t)}) = \mathbb{E}[\ell(g^{(t)}, y)] + \lambda_a^{(t)}\|a^{(t)}\|_2^2$. Note that the gradient update using $\tilde{L}_\mathcal{D}(g^{(t)})$ is the same as the update in our learning algorithm. Then by Theorem 27, Lemma 23, and Lemma 25,

$$\frac{1}{T}\sum_{t=3}^{T} \tilde{L}_\mathcal{D}(g^{(t)}) \leq \frac{\|\tilde{a}\|_2^2}{2} + \|\tilde{a}\|_2 \sqrt{\|\tilde{a}\|_0} \left(O\left(\frac{T\eta\lambda\tilde{\sigma}}{k}\right) + \eta^2 T^2 + O\left(\frac{T}{km^2}\right) \right) \tag{281}$$

$$+ \frac{\|\tilde{a}\|_2^2}{2\eta T} + \|a^{(2)}\|_2 \sqrt{m} + \eta m \tag{282}$$

$$\leq O\left(\frac{k^9}{m} + k^4 \eta^2 T^2 + \frac{k^3 T}{m^2} + \frac{k^9}{\eta T m} + \eta m\right). \tag{283}$$

$$\leq O\left(\frac{k^8}{m^{2/3}} + \frac{k^3 T}{m^2} + \frac{k^2 m^{2/3}}{T}\right). \tag{284}$$

The statement follows from that 0-1 classification error is bounded by the hinge-loss. □

Theorem 27 (Theorem 13 in Daniely & Malach (2020)). *Fix some η, and let $f_1, \ldots, f_T$ be some sequence of convex functions. Fix some θ_1, and assume we update $\theta_{t+1} = \theta_t - \eta\nabla f_t(\theta_t)$. Then for every θ^* the following holds:*

$$\frac{1}{T}\sum_{t=1}^{T} f_t(\theta_t) \leq \frac{1}{T}\sum_{t=1}^{T} f_t(\theta^*) + \frac{1}{2\eta T}\|\theta^*\|_2^2 + \|\theta_1\|_2 \frac{1}{T}\sum_{t=1}^{T}\|\nabla f_t(\theta_t)\|_2 + \eta \frac{1}{T}\sum_{t=1}^{T}\|\nabla f_t(\theta_t)\|_2^2. \tag{285}$$

C   Lower Bound for Linear Models on Fixed Feature Mappings

Theorem 28 (Restatement of Theorem 2). *Suppose Ψ is a data-independent feature mapping of dimension N with bounded features, i.e., $\Psi : \mathcal{X} \to [-1, 1]^N$. Define for $B > 0$:*

$$\mathcal{H}_B = \{h(\tilde{x}) : h(\tilde{x}) = \langle \Psi(\tilde{x}), w \rangle, \|w\|_2 \le B\}. \tag{286}$$

Then, if $3 < k \le D/16$ and k is odd, then there exists $\mathcal{D} \in \mathcal{F}_\Xi$ such that all $h \in \mathcal{H}_B$ have hinge-loss at least $p_o \left(1 - \frac{\sqrt{2N}B}{2^k} \right)$.

Proof of Theorem 2. We first show that $\mathcal{F}_\Xi$ contains some distributions that are essentially sparse parity learning problems, and then we invoke the lower bound result from existing work for such problems.

Consider $\mathcal{D}$ defined as follows.

- Let $P = \{i \in [k] : i \text{ is odd}\}$. That is, if there are odd numbers of 1's in $\tilde{\phi}_A$, then $y = +1$.

- Let $\mathcal{D}_{\tilde{\phi}}^{(0)}$ be a distribution where all entries $\tilde{\phi}_j$ are i.i.d. with $\Pr[\tilde{\phi}_j = 0] = \Pr[\tilde{\phi}_j = 1] = 1/2$. Let $\mathcal{D}^{(0)}$ be the distribution over $(\tilde{x}, y)$ induced by $\mathcal{D}_{\tilde{\phi}}^{(0)}$ and the above P.

- Let $\mathcal{D}_{\tilde{\phi}}^{(1)}$ be a distribution where all entries $\tilde{\phi}_j$ for $j \notin A$ are i.i.d. with $\Pr[\tilde{\phi}_j = 1] = p_o/(2 - 2p_o)$, while $\Pr[\tilde{\phi}_A = (0, 0, \ldots, 0)] = \Pr[\tilde{\phi}_A = (1, 1, \ldots, 1)] = 1/2$. Let $\mathcal{D}^{(1)}$ be the distribution over $(\tilde{x}, y)$ induced by $\mathcal{D}_{\tilde{\phi}}^{(1)}$ and the above P.

- Let $\mathcal{D}_A^{\mathrm{mix}} = p_o \mathcal{D}^{(0)} + (1 - p_o) \mathcal{D}^{(1)}$.

It can be verified that such distributions are included in $\mathcal{F}_\Xi$ for $\gamma = \Theta(1)$.

Assume for contradiction that for all $\mathcal{D} \in \mathcal{F}_\Xi$, there exists $h^* \in \mathcal{H}_B$ such that $h = \langle \Psi, w^* \rangle$ loss smaller than $p_o \left(1 - \frac{\sqrt{2N}B}{2^k} \right)$. Then for all the distributions $\mathcal{D}_A^{\mathrm{mix}}$ defined above, we have

$$\mathbb{E}_{\mathcal{D}^{(0)}}[\ell(h^*(\tilde{x}), y)] < 1 - \frac{\sqrt{2N}B}{2^k}. \tag{287}$$

Now let $\mathcal{D}_z$ be a distribution over $z \in \{-1, +1\}^D$ with i.i.d. entries z_j and $\Pr[z_j = -1] = \Pr[z_j = +1] = 1/2$. Let $f_A(z) = \prod_{j \in A} z_j$ be the k-sparse parity functions. Let $\Psi'(z) = \Psi(M(z + 1)/2)$. Then we have $h'(z) = \langle \Psi'(z), w^* \rangle$ such that for all A,

$$\mathbb{E}_{\mathcal{D}_z}[\ell(h'(z), f_A(z))] < 1 - \frac{\sqrt{2N}B}{2^k}. \tag{288}$$

This is contradictory to Theorem 29. $\qquad\square$

The following theorem is implicit in the proof in Theorem 1 in Daniely & Malach (2020).

Theorem 29. *For a subset $A \subseteq [D]$ of size k, let the distribution $\mathcal{D}_A$ over (z, y) defined as follows: z is uniform over $\{\pm 1\}^D$ and $y = \prod_{i \in A} z_i$. Fix some $\Psi : \{\pm 1\}^D \to [-1, +1]^N$, and define:*

$$\mathcal{H}_\Psi^B = \{z \to \langle \Psi(z), w \rangle : \|w\|_2 \le B\}.$$

If k is odd and $k \le D/16$, then there exists some A such that

$$\min_{h \in \mathcal{H}_\Psi^B} \mathbb{E}_{\mathcal{D}_A}[\ell(h(z), y)] \ge 1 - \frac{\sqrt{2N}B}{2^k}.$$

D LOWER BOUND FOR LEARNING WITHOUT INPUT STRUCTURE

First recall the Statistical Query model (Kearns, 1998). In this model, the learning algorithm can only receive information about the data through statistical queries. A statistical query is specified by some property predicate Q of labeled instances, and a tolerance parameter $\tau \in [0, 1]$. When the algorithm asks a statistical query (Q, τ), it receives a response $\hat{P}_Q \in [P_Q - \tau, P_Q + \tau]$, where $P_Q = \Pr[Q(x, y) \text{ is true}]$. Q is also required to be polynomially computable, i.e., for any (x, y) $Q(x, y)$ can be computed in polynomial time. Notice that a statistical query can be simulated by empirical average of a large random sample of data of size roughly $O(1/\tau^2)$ to assure the tolerance τ with high probability.

Blum et al. (1994) introduces the notion of Statistical Query dimension, which is convenient for our purpose.

Definition 30 (Definition 2 in Blum et al. (1994)). *For concept class C and distribution $\mathcal{D}$, the statistical query dimension SQ-DIM$(C, \mathcal{D})$ is the largest number d such that C contains d concepts $c_1, \ldots, c_d$ that are nearly pairwise uncorrelated: specifically, for all $i \neq j$,*

$$| \Pr_{x \sim \mathcal{D}}[c_i(x) = c_j(x)] - \Pr_{x \sim \mathcal{D}}[c_i(x) \neq c_j(x)]| \leq 1/d^3. \tag{289}$$

Theorem 31 (Theorem 12 in Blum et al. (1994)). *In order to learn C to error less than $1/2 - 1/d^3$ in the Statistical Query model, where $d = SQ\text{-}DIM(C, \mathcal{D})$, either the number of queries or $1/\tau$ must be at least $\frac{1}{2}d^{1/3}$.*

We now use the above tools to prove our lower bound.

Theorem 32 (Restatement of Theorem 3). *For any algorithm in the Statistical Query model that can learn over $\mathcal{F}_{\Xi_0}$ to error less than $\frac{1}{2} - \frac{1}{\binom{D}{k}^3}$, either the number of queries or $1/\tau$ must be at least $\frac{1}{2}\binom{D}{k}^{1/3}$.*

Proof of Theorem 3. Consider the following concept class and marginal distribution:

- Let $\mathcal{D}$ be the distribution over $\tilde{x}$, given by $\tilde{x} = M\tilde{\phi}$ and $\tilde{\phi}_j$ are i.i.d. with $\Pr[\tilde{\phi}_j = 0] = \Pr[\tilde{\phi}_j = 1] = 1/2$.

- Let C be the class of functions $y = g_A(\tilde{\phi}) = \mathbb{I}[\sum_j(1 - \tilde{\phi}_j) \text{ is odd}]$ for different $A \subseteq [D]$.

The distributions over $(\tilde{x}, y)$ induced by $(C, \mathcal{D})$ are a subset of $\mathcal{F}_{\Xi_0}$. It is then sufficient to show that SQ-DIM$(C, \mathcal{D}) \geq \binom{D}{k}$.

It is easy to see that C are essentially the sparse parity functions: if $z_j = 2\tilde{\phi}_j - 1$, then $g_A(\tilde{\phi}) = \prod_{j \in A} z_j$. This then implies that the g_A's are uncorrelated, so SQ-DIM$(C, \mathcal{D}) \geq \binom{D}{k}$. $\square$

E COMPLETE EXPERIMENTAL RESULTS

Our experiments mainly focus on feature learning and the effect of the input structure. We first perform simulations on our learning problems to (1) verify our main theorems on the benefit of feature learning and the effect of input structure (2) verify our analysis of feature learning in networks. We then check if our insights carry over to real data: (3) whether similar feature learning is presented in real network/data; (4) whether damaging the input structure lowers the performance. The results are consistent with our analysis and provide positive support for the theory.

The experiments were ran 5 times with different random seeds, and the average results (accuracy) are reported. The standard deviations of the results are smaller than 0.5% and thus we do not present them for clarity. The hardware specifications are 4 Intel(R) Core(TM) i7-7700HQ CPU @ 2.80GHz, 16 GB RAM, and one NVIDIA GPU GTX1080.

E.1 SIMULATION

We train a two-layer network following our learning process. We use two fixed feature methods: the NTK (Fang et al., 2021) and random feature (RF) methods based on the same network and random initialization as the network learning. More precisely, in the NTK method, we randomly initialize the network and take its NTK and learn a classifier on it. In the RF method, we freeze the first layer of the network, and train the second layer (on the random features given by the frozen neurons). The training step number is the same as that in network learning. We also test these three methods on the data distribution with input structure removed (i.e., $\mathcal{F}_{\Xi_0}$ in Theorem 3). For comparison, we take the representation of our two-layer network at step one/step two, named One Step/Two Step (fix the weight of the 1st layer after the first step/second step to train the weight of the second layer), and train the best classifiers on top of them.

Recall that our analysis is on the *directions* of the weights without considering their *scaling*, and thus it is important to choose cosine similarity rather than the typical ℓ_2 distance. Thus, we use metric Cos Similarity $\max_{\{i \in [2m]\}} \cos(w_i, \sum_{j \in A} M_j)$ in our tables, and use Multidimensional Scaling to plot the weights distribution. The simulation dataset size is 20000. During training, the batch size is 1000, while for the first two steps we use the approximate full gradient (batch size is 20000). Each step is corresponding to one weights update.

E.1.1 PARITY LABELING

Setting. We generate data according to the parity function data distributions used in our proof of the lower bound for fixed features (Theorem 2), with $d = 500, D = 100, k = 5, p_o = 1/2$, with a randomly sampled A. More precisely, we consider $\mathcal{D}$ defined as follows.

- Let $P = \{i \in [k] : i \text{ is odd}\}$. That is, if there are odd numbers of 1's in $\tilde{\phi}_A$, then $y = +1$.

- Let $\mathcal{D}_{\tilde{\phi}}^{(0)}$ be a distribution where all entries $\tilde{\phi}_j$ are i.i.d. with $\Pr[\tilde{\phi}_j = 0] = \Pr[\tilde{\phi}_j = 1] = 1/2$. Let $\mathcal{D}^{(0)}$ be the distribution over $(\tilde{x}, y)$ induced by $\mathcal{D}_{\tilde{\phi}}^{(0)}$ and the above P.

- Let $\mathcal{D}_{\tilde{\phi}}^{(1)}$ be a distribution where all entries $\tilde{\phi}_j$ for $j \notin A$ are i.i.d. with $\Pr[\tilde{\phi}_j = 1] = p_o/(2 - 2p_o)$, while $\Pr[\tilde{\phi}_A = (0, 0, \ldots, 0)] = \Pr[\tilde{\phi}_A = (1, 1, \ldots, 1)] = 1/2$. Let $\mathcal{D}^{(1)}$ be the distribution over $(\tilde{x}, y)$ induced by $\mathcal{D}_{\tilde{\phi}}^{(1)}$ and the above P.

- Let $\mathcal{D}_A^{\text{mix}} = p_o \mathcal{D}^{(0)} + (1 - p_o)\mathcal{D}^{(1)}$.

The network and the training follow Section 3, where the network size is $m = 300$ and the training time $T = 600$ steps.

Verification of the Main Results. Figure 5 shows that the results are consistent with our analysis. Network learning gets high test accuracy while the two fixed feature methods get significantly lower accuracy. Furthermore, when the input structure is removed, all three methods get test accuracy similar to random guessing.

Figure 5: Test accuracy on simulated data under parity labeling with or without input structure.

Figure 6: Visualization of the weights w_i's after initialization/one gradient step/two gradient steps/few gradient steps in network learning under parity labeling. The red star denotes the ground-truth $\sum_{j \in A} M_j$; the orange star is $-\sum_{j \in A} M_j$. The red dots are the weights closest to the red star after two steps; the orange ones are for the orange star.

Model	Network	NTK	RF	One Step	Two Step	Network w/o structure
Train Acc (%)	100.0	84.0	74.7	51.3	100.0	100.0
Test Acc (%)	100.0	86.4	76.0	52.2	100.0	52.0
Cos Similarity	0.997	NA	0.114	0.848	0.997	0.253

Table 1: Parity labeling results in six methods.

Feature Learning in Networks Figure 6 shows that the results are as predicted by our analysis. After the first gradient step, some weights begin to cluster around the ground-truth $\sum_{j \in A} M_j$ (or $-\sum_{j \in A} M_j$ due to we have a_i in the gradient update which can be positive or negative). After the second step the weights get improved and well-aligned with the ground-truth (with cosine similarity > 0.99).

Table 1 shows the results for different methods. Recall that the Cos Similarity metric is $\max_{\{i \in [2m]\}} \cos(w_i, \sum_{j \in A} M_j)$, which reports the cosine value of the closest one. One Step refers to the method where we take the neurons after one gradient step, freeze their weights, and train a classifier on top; similar for Two Step. One Step gets test accuracy about 52%, while Two Step gets accuracy about 100%. This demonstrates that while some effective feature emerge in the first step, they need to be improved in the second step for accurate prediction. NTK, random feature, One Step all failed, while Network and Two Step can achieve 100% test accuracy. Network w/o structure refers to training the network on data without the input structure. It overfits the training dataset with 52% test accuracy.

Figure 7: Test accuracy on simulated data under interval labeling with or without input structure.

Figure 8: Visualization of the weights w_i's after initialization/one gradient step/two gradient steps/few gradient steps in network learning under interval labeling. The red star denotes the ground-truth $\sum_{j \in A} M_j$; the orange star is $-\sum_{j \in A} M_j$. The red dots are the weights closest to the red star after two steps; the orange ones are for the orange star.

Model	Network	NTK	RF	One Step	Two Step	Network w/o structure
Train Acc (%)	100.0	100.0	76.4	44.1	100.0	100.0
Test Acc (%)	100.0	100.0	73.2	41.0	100.0	100.0
Cos Similarity	1.00	NA	0.153	0.901	0.994	0.965

Table 2: Interval labeling results in six methods.

E.1.2 INTERVAL LABELING

Setting. We also tried interval function, where $y = 1$ if $\sum_{i \in A} \tilde{\phi}_i$ is in the range $[t_1, t_2]$ with $t_1 = 20$ and $t_2 = 30$, otherwise $y = -1$. We use $d = 500, D = 100, k = 30$. The $\tilde{\phi}_i$'s are independent, and $\Pr[\tilde{\phi}_i = 1] = 2/3$ for any $i \in A$, and $\Pr[\tilde{\phi}_i = 1] = 1/2$ otherwise. When the input structure is removed, we set $\Pr[\tilde{\phi}_i = 1] = 1/2$ for all i's.

The network and training again follows Section 3 with a network size $m = 100$ and the training time $T = 200$ steps.

Verification of the Main Results. Figure 7 shows that network learning learns the fastest, NTK learns slower but reaches similar test accuracy, while random feature can only reach a decent but lower accuracy. This is because for such simpler labeling functions, fixed feature methods can still achieve good performance (note that the lower bound does not hold for such a case), while the performance depends on what fixed features to use.

Furthermore, when the input structure is removed, the methods still get similar (or only slightly worse) performance as with input structure. This shows that when the labeling function is simple, the help of the input structure for learning may not be needed. In the experiments on real data, we will show that when the input structure is changed, it indeed leads to lower performance which suggests

Figure 9: Visualization of the neurons' weights in a two-layer network trained on the subset of MNIST data with label 0/1. The weights gradually form two clusters.

Figure 10: Visualization of the neurons' weights in a two-layer network trained on the subset of CIFAR10 data with label airplane/automobile. The weights gradually form two clusters.

Figure 11: Visualization of the neurons' weights in a two-layer network trained on the subset of SVHN data with label 0/1. The weights gradually form four clusters.

that the labeling function in practice is typically more complicated than this interval labeling setting, and the help of the input structure is significant for learning.

Feature Learning in Networks. Figure 8 shows the phenomenon of feature learning similar to that in the parity labeling setting. Table 2 shows the test accuracy of six different methods. Random feature and One Step failed, while Network, NTK and Two Step succeed showing that interval labeling setting is a simpler case than parity labeling setting.

E.2 REAL DATA: FEATURE LEARNING IN NETWORKS

We take the subset of MNIST (Deng, 2012) with labels 0/1, CIFAR10 (Krizhevsky, 2012) with labels airplane/automobile and SVHN (Netzer et al., 2011) with labels 0/1, and train a two-layer network with $m = 50$. We use traditional weight initialization method (random Gaussian) and training method (SGD with momentum $= 0.95$ without regularization) in this section, for our purpose of investigating the training dynamics in practice.

Then we visualize the neurons' weights following the same method in the simulation. Figure 9, Figure 10 and Figure 11 show a similar feature learning phenomenon: effective features emerge after a few steps, and then get improved to form clusters. This shows the insights obtained on our learning problems are also applicable to the real data.

E.3 REAL DATA: THE EFFECT OF INPUT STRUCTURE

To study the influence of the input structure, we propose to keep the labeling function unchanged, vary the input distributions, and exam the change of the loss surface and the training dynamics. We first describe the detailed experimental methodology, which allows us to generate data with similar labeling function but different input distributions. Then we perform experiments on the generated datasets to investigate the change of the learning due to the change in the input distributions, and present the experimental results. Finally, we also perform experiments to verify the intuition behind our experimental method.

E.3.1 EXPERIMENTAL METHODOLOGY

We consider the following experimental method. Given an original dataset $\mathcal{L} = \{(x_i, y_i)\}_{i=1}^n$ (e.g., CIFAR10) and an unlabeled dataset $\mathcal{U} = \{\tilde{x}_i\}_{i=1}^m$ from a proposed distribution $P_{\mathcal{U}}$ (e.g., Gaussians), first extend the labeling function of $\mathcal{L}$ to $\mathcal{U}$, giving synthetic labels $\tilde{y}_i$ to $\tilde{x}_i$. Then train a neural network on the union of $\mathcal{L}$ and the synthetic data $\mathcal{L}_{\mathcal{U}} = \{(\tilde{x}_i, \tilde{y}_i)\}_{i=1}^m$. By investigating the new training dynamics, in particular the difference on the original part $\mathcal{L}$ and the synthetic part $\mathcal{L}_{\mathcal{U}}$, we can see the effect of the input structure. The original dataset should be from real-world data, since one of our goals is to compare them with synthetic data, and identify the properties of real-world data important for the success of learning.

A natural idea is to first learn a powerful network $f(x)$ (called the teacher) on $\mathcal{L}$ to approximate the true labeling function, then apply f on $\mathcal{U}$ to generate synthetic labels, and finally train another network (called the student) on the synthetic data and original data. However, we found that naïvely implementation of this idea fails miserably: the support of $\mathcal{L}$ and $\mathcal{U}$ can be typically different, and the powerful network learned over $\mathcal{L}$ can have entirely different behavior on $\mathcal{U}$. Therefore, we need to control the size of the teacher f so that the labeling on $\mathcal{U}$ has similar complexity as that on $\mathcal{L}$. For our purpose, we can define the complexity of the labeling on $\mathcal{L}$ as the minimum size of the teacher achieving an approximation error ϵ for a chosen ϵ, if the ground-truth data distribution of $\mathcal{L}$ is known. However, given only limited data, we cannot faithfully estimate the needed size of the teacher, and need to take into account the variance introduced by the finite data.

Our key idea is to use the U-shaped curve of the bias-variance trade-off and select the size of the teacher at the minimum of the U-shaped curve. Since recent works (Belkin et al., 2019; Nakkiran et al., 2020) show that neural networks can have a double descent curve for the error v.s. model complexity, we thus plot the double descent curve, and find the minimum in the classical regime (corresponding to the traditional U-shape curve).

Our method is designed based on the following two reasons. First, on the U-shaped curve, the complexity of the network is still roughly controlled by that of the number of parameters. The local minimum of the U-shaped curve is a good measurement of the complexity of the data. If the ground-truth is much more complicated than the teacher, then increasing the teacher's size leads to a significant decrease in the approximation error (bias) compared to a small increase in the variance, that is, we will be on the left-hand side of the U-shaped. In contrast, on the right-hand side of the U-shaped, increasing the teacher's size leads to a small decrease in the bias compared to a significant increase in the variance. That is, the complexity of the ground-truth is comparable to or lower than the teacher. So the local minimum approximates the complexity of the ground-truth labeling function.

Second, the local minimum point is chosen to get the best approximation of the true labels. This helps to maintain the labeling from the real-world data and thus helps our investigation on the input, since too drastic change in the labeling can affect the training.

We note that the method is not perfect. First, the teacher at the local minimum of U-shape may not have very high accuracy, especially on more complicated data. To alleviate this, we also use the teacher to give synthetic labels y_i' to x_i in $\mathcal{L}$, and train the student network on $\mathcal{L}' = \{(x_i, y_i')\}_{i=1}^n$. Though this introduces some differences from the original labels, it is acceptable for our purpose of studying the inputs. Furthermore, ensuring the consistency of the labels on the original input in $\mathcal{L}$ and $\mathcal{U}$ is important in our experiments. Second, the measurement is an approximation due to variance. Since only limited labeled data is available, it's important and necessary to calibrate the measurement w.r.t. the level of variance on the given dataset.

Method Description. Algorithm 1 presents the details. For a fixed network architecture for the teacher f, it first varies the network size and plots the double descent curve. Then it selects the local minimum in the classic regime of U-shape and trains the teacher with the corresponding size. In practice, we observed that the teacher might have unbalanced probabilities for different classes on $\mathcal{U}$ if its training does not take into account $\mathcal{U}$. Therefore, we propose the following heuristic regularization using $x \in \mathcal{U}$, where λ is a regularization weight, and $f(x)$ is the probabilities over classes given by the teacher:

$$R(x) = R_1(x) + \lambda R_2(x) \tag{290}$$

$$R_1(x) = \sum_j \left(\frac{\sum_i f(x)_j}{m} \ln \frac{\sum_i f(x)_j}{m} \right) \tag{291}$$

$$R_2(x) = -\frac{1}{m} \sum_i \sum_j (f(x)_j \ln(f(x)_j)). \tag{292}$$

Here, $R_1(x)$ guarantees that each kind of label has the same average probability to be generated, and $R_2(x)$ pushes the probability away from uniform to avoid the case that the class probabilities for each data point converge to uniform.

Algorithm 1 Learning the teacher network to generate synthetic labels for studying the effect of the input structure

---

Input: teacher architecture f, labeled dataset $\mathcal{L} = \{(x_i, y_i)\}_{i=1}^n$, unlabeled dataset $\mathcal{U} = \{\tilde{x}_i\}_{i=1}^m$. Let i to be the size of f, f_i to be the teacher of size i.
for $i = 1$ to n **do**
    Train f_i on $\mathcal{L}$ and let l_i denote the test loss
end for
Plot l_i v.s. i, identify the classical regime, and the size i_t corresponding to the local minimum in classical regime.
Train f_{i_t} on $\mathcal{L}$ with a regularizer $R(x)$ on $\mathcal{U}$ defined in (290).
Output: f_{i_t}

E.3.2 EXPERIMENTAL RESULTS

Network models. Here we use one-hidden-layer fully-connected networks with m hidden units and quadratic activation functions. The network is denoted as $\mathrm{FC}(m)$. We use $\mathrm{ResNet}(m)$, which is a ResNet-18 convolutional neural network (He et al., 2016) with m filters in the first residual block. It is obtained by scaling the number of filters in each block proportionally from the standard ResNet-18 network which is $\mathrm{ResNet}(64)$.

Datasets. We use MNIST (Deng, 2012), CIFAR10 (Krizhevsky, 2012) and SVHN (Netzer et al., 2011) as $\mathcal{L}$, and use Gaussian and images in Tiny ImageNet (Le & Yang, 2015) as $\mathcal{U}$. We generate the mixture data, where the fraction of the unlabeled data is denoted as α.

Setup. We first use Algorithm 1 on the labeled data $\mathcal{L}$ and the unlabeled data $\mathcal{U}$ to get a synthetic labeling function (the teacher network) and then use it to give synthetic labels on a mixture of inputs from $\mathcal{L}$ and $\mathcal{U}$. For MNIST, the teacher network learned is $\mathrm{FC}(9)$, where the number of the hidden units is determined by Algorithm 1. See empirical verification in Figure 15. For CIFAR10 and SVHN, the teacher networks are $\mathrm{ResNet}(5)$ and $\mathrm{ResNet}(2)$, respectively, as determined by our method. The student network for MNIST is $\mathrm{FC}(9)$, and those for CIFAR10 and SVHN are $\mathrm{ResNet}(9)$ and $\mathrm{ResNet}(8)$, respectively. Finally, we train the student networks on these new datasets with perturbed input distributions.

Figure 12 shows the results on an equal mixture of data and Gaussian. It presents the test accuracy of the student on the original data part, the Gaussian part, and the whole mixture. For example, for CIFAR10, the test accuracy on the whole mixture is lower than that of training on the original CIFAR10, showing that the input structure indeed has a significant impact on the learning. Furthermore, the network learns well over the CIFAR10 part (with accuracy similar to that on the original data) but learns slower with worse accuracy on the Gaussian part. This suggests that the CIFAR10 input structure is still helping the network to learn effective features. While the results on

Figure 12: Test accuracy at different steps for an equal mixture $\alpha = 0.5$ of Gaussian inputs with data: (a) MNIST, (b) CIFAR10, (c) SVHN.

Figure 13: Test accuracy at different steps for an equal mixture $\alpha = 0.5$ of Tiny ImageNet inputs with data: (a) CIFAR10, (b) SVHN.

Figure 14: Test accuracy at different steps for varying mixture α of Gaussian inputs with CIFAR10: (a) $\alpha = 0.25$, (b) $\alpha = 0.50$, (c) $\alpha = 0.75$.

MNIST+Gaussian do not show a significant trend (possibly because the tasks there are simpler), the results on SVHN+Gaussian show similar significant trends as CIFAR10+Gaussian.

Figure 14 shows the results when we vary the fraction of the Gaussian data α. We observe that the test accuracy curve on the original part and that on the synthetic part have roughly the same trend for different α as before, further verifying our insights.

Figure 13 shows the results when mixed with Tiny ImageNet data instead of Gaussians. It shows a similar trend, while the performance on the Tiny ImageNet part is higher than that on the Gaussian part. This suggests that compared to Gaussians, the Tiny ImageNet data has helpful input structures, though not as helpful as that on the original data for learning the particular labeling.

(a) Teacher's double descent curve   (b) Student's curve when Teacher=FC(9)

(c) Student's curve when Teacher=FC(50)   (d) Student's curve when Teacher=FC(500)

Figure 15: Double descent curves of the students trained on data with synthetic labels (Loss v.s. Parameter number).

E.3.3 EMPIRICAL VERIFICATION OF OUR METHOD

We also perform experiments to verify the intuition behind our methodology, i.e., the method gives a synthetic labeling function with roughly the same complexity on the original inputs and the injected inputs. We first use our method on MNIST and samples (of the same size as MNIST) from a Gaussian to get the teacher FC(9); the double descent curve is in Figure 15(a). Then we train students on the Gaussian data with synthetic labels from the teacher, and plot the double descent curve for the students in Figure 15(b). The local minimums of the two U-shapes are roughly the same, matching our reasoning. Then we also train larger teachers and plot the double descent curve for students on Gaussian data. Figure 15(c) Teacher size 50. Figure 15(d) Teacher size 500. The local minimum of the U-shape becomes larger when the teacher gets larger, again matching our reasoning.

F PROVABLE GUARANTEES FOR NEURAL NETWORKS IN A MORE GENERAL SETTING

This section provides the analysis in a more general setting. We first describe the learning problems, and then provide the proofs following similar intuitions as for the simpler settings in the main text.

F.1 PROBLEM SETUP

Let $\mathcal{X} = \mathbb{R}^d$ be the input space, and $\mathcal{Y} = \{\pm 1\}$ be the label space. Suppose $M \in \mathbb{R}^{d \times D}$ is a dictionary with D elements, where each element M_j can be regarded as a pattern. We assume quite general incoherent dictionary:

(D) M is μ-incoherent, i.e., the columns of M are unit vectors, and for any $i \neq j, |\langle M_i, M_j \rangle| \leq \mu/\sqrt{d}$.

Note that the setting in the main text corresponds to $\mu = 0$.

Let $\tilde{\phi} \in \{0, 1\}^D$ be a hidden vector that indicates the presence of each pattern, and $\mathcal{D}_{\tilde{\phi}}$ a distribution for $\tilde{\phi}$. Let $A \subseteq [D]$ be a subset of size k corresponding to the class relevant patterns. Let $P \subseteq [k]$. We first sample $\tilde{\phi}$ from $\mathcal{D}_{\tilde{\phi}}$, and then generate the input $\tilde{x}$ and the class label y from $\tilde{\phi}, A, P$ by:

$$\tilde{x} = M\tilde{\phi} + \zeta, \quad y = \begin{cases} +1, & \text{if } \sum_{i \in A} \tilde{\phi}_i \in P, \\ -1, & \text{otherwise} \end{cases} \tag{293}$$

where the Gaussian noise $\zeta \sim \mathcal{N}(0, \sigma_\zeta^2 I_{d \times d})$ is independent from $\tilde{\phi}$. Note that the setting in the main text corresponds to $\sigma_\zeta = 0$.

We allow general $\mathcal{D}_{\tilde{\phi}}$ with the following assumptions:

(A1) The patterns in A are correlated with the labels: for any $i \in A$, for $v \in \{\pm 1\}$ let $\gamma_v = \mathbb{E}[y\tilde{\phi}_i | y = v]$, then $\gamma := (\gamma_{+1} + \gamma_{-1})/2 > 0$.

(A2) Each pattern outside A is independent of all other patterns and identically distributed. Let $p_o := \Pr[\tilde{\phi}_i = 1] \leq 1/2$ denote the probability they appear.

Note that we allow imbalanced classes. Let $p_{\min} := \min(\Pr[y = -1], \Pr[y = +1])$. If the classes are balanced, then the assumption **(A1)** implies the assumption **(A1)** in the main text, so the setting here is more general.

Let $\mathcal{D}(A, P, \mathcal{D}_{\tilde{\phi}})$ denote the distribution on $(\tilde{x}, y)$ corresponding to some A, P, and $\mathcal{D}_{\tilde{\phi}}$. Given parameters $\Xi = (d, D, k, \gamma, p_o, \mu, \sigma_\zeta)$, the family $\mathcal{F}_\Xi$ of distributions for learning is the set of all $\mathcal{D}(A, P, \mathcal{D}_{\tilde{\phi}})$ with $A \subseteq [D]$, $P \subseteq [k]$, and $\mathcal{D}_{\tilde{\phi}}$ satisfying the above assumptions.

One special case is the mixture of two Gaussians.

Example. Suppose M has one single column v, and $y = +1$ if $\tilde{\phi} = 1$ and $y = -1$ otherwise. Then the data distribution is simply a mixture of two Gaussians: $\tilde{x} \sim \frac{v}{2} + \mathcal{N}(y\frac{v}{2}, \sigma_\zeta^2 I_{d \times d})$.

F.1.1 NEURAL NETWORK LEARNING

Again, we will normalize the data for learning: we first compute $x = (\tilde{x} - \mathbb{E}[\tilde{x}])/\tilde{\sigma}$ where $\tilde{\sigma}^2 := \sum_{i=1}^d (\tilde{x}_i - \mathbb{E}[\tilde{x}_i])^2 = \sum_{j \in [D]} \text{Var}(\tilde{\phi}_j) + d\sigma_\zeta^2$ is the variance of the data, and then train on (x, y). This is equivalent to setting $\phi = (\tilde{\phi} - \mathbb{E}[\tilde{\phi}])/\tilde{\sigma}$ and generating $x = M\phi + \zeta/\sigma_\zeta$. For $(\tilde{x}, y)$ from $\mathcal{D}$ and the normalized (x, y), we will simply say $(x, y) \sim \mathcal{D}$.

The learning will be the same as that in the main text, except that we will use a weighted loss to handle the imbalanced classes in the first two steps for feature learning, and then use the unweighted loss in the remaining steps. Formally, the weighted loss is:

$$L_\mathcal{D}^\alpha(g; \sigma_\xi) = \mathbb{E}_{(x,y)}[\alpha_y \ell(y, g(x; \xi))], \tag{294}$$

where the class weights $\alpha_v = \frac{1}{2\Pr[y=v]}$ for $v \in \{\pm 1\}$.

F.2 MAIN RESULT

In this setting, we have the following theorem:

Theorem 33. *Set*

$$\eta^{(1)} = \frac{\gamma^2 p_{\min}\tilde{\sigma}}{km^3}, \lambda_a^{(1)} = 0, \lambda_w^{(1)} = 1/(2\eta^{(1)}), \sigma_\xi^{(1)} = 1/k^2, \tag{295}$$

$$\eta^{(2)} = 1, \lambda_a^{(2)} = \lambda_w^{(2)} = 1/(2\eta^{(2)}), \sigma_\xi^{(2)} = 1/k^2, \tag{296}$$

$$\eta^{(t)} = \eta = \frac{k^2}{Tm^{1/3}}, \lambda_a^{(t)} = \lambda_w^{(t)} = \lambda \leq \frac{k^3}{\tilde{\sigma}m^{1/3}}, \sigma_\xi^{(t)} = 0, \text{ for } 2 < t \leq T. \tag{297}$$

For any $\delta \in (0, O(1/k^3))$, if $\mu \leq O(\sqrt{d}/D)$, $\sigma_\zeta \leq O(\min\{1/\tilde{\sigma}, \tilde{\sigma}/\sqrt{d}\})$, $p_o = \Omega(k^2/D)$, $k = \Omega\left(\log^2\left(\frac{Dmd}{\delta\gamma p_{\min}}\right)\right)$, $m \geq \max\{\Omega(k^4), D, d\}$, then we have for any $\mathcal{D} \in \mathcal{F}_\Xi$, with probability at least $1 - \delta$, there exists $t \in [T]$ such that

$$\Pr[\text{sign}(g^{(t)}(x)) \neq y] \leq L_\mathcal{D}(g^{(t)}) = O\left(\frac{k^8}{m^{2/3}} + \frac{k^3 T}{m^2} + \frac{k^2 m^{2/3}}{T}\right). \tag{298}$$

Consequently, for any $\epsilon \in (0,1)$, if $T = m^{4/3}$, and $m \geq \max\{\Omega(k^{12}/\epsilon^{3/2}), D\}$, then

$$\Pr[\text{sign}(g^{(t)}(x)) \neq y] \leq L_\mathcal{D}(g^{(t)}) \leq \epsilon. \tag{299}$$

The rest of the section is devoted to the proof of this theorem.

F.3 NOTATIONS

Recall some notations that we will use throughout the analysis.

For a vector v and an index set I, let v_I denote the vector containing the entries of v indexed by I, and v_{-I} denote the vector containing the entries of v with indices outside I.

Let $\rho := M^\top M$. Then we have $\rho_{jj} = 1$ for any j, and $|\rho_{j\ell}| \leq \mu/\sqrt{d}$ for any $j \neq \ell$.

By initialization, $w_i^{(0)}$ for $i \in [m]$ are i.i.d. copies of the same random variable $w^{(0)} \sim \mathcal{N}(0, \sigma_w^2 I_{d\times d})$; similar for $a^{(0)}$ and $b^{(0)}$. Let $\sigma_\phi^2 := p_o(1 - p_o)/\tilde{\sigma}^2$ denote the variance of ϕ_ℓ for $\ell \notin A$.

Let $q_\ell := \langle w^{(0)}, M_\ell \rangle$. Similarly, define $q_{i,\ell}^{(t)} := \langle w_i^{(t)}, M_\ell \rangle$.

We also define the following sets to denote typical initialization. For a fixed $\delta \in (0,1)$, define

$$\mathcal{G}_w(\delta) := \left\{ w \in \mathbb{R}^d : q_\ell = \langle w, M_\ell \rangle, \frac{\sigma_w^2 d}{2} \leq \|w^{(0)}\|_2^2 \leq \frac{3\sigma_w^2 d}{2}, \tag{300} \right.$$

$$\frac{\sigma_w^2(D-k)}{2} \leq \sum_{\ell \notin A} q_\ell^2 \leq \frac{3\sigma_w^2(D-k)}{2},$$

$$\left. \max_\ell |q_\ell| \leq \sigma_w\sqrt{2\log(Dm/\delta)} \right\}, \tag{301}$$

$$\mathcal{G}_a(\delta) := \{a \in \mathbb{R} : |a| \leq \sigma_a\sqrt{2\log(m/\delta)}\}. \tag{302}$$

$$\mathcal{G}_b(\delta) := \{b \in \mathbb{R} : |b| \leq \sigma_b\sqrt{2\log(m/\delta)}\}. \tag{303}$$

F.4 EXISTENCE OF A GOOD NETWORK

We first show that there exists a network that can fit the data distribution.

Lemma 34. *Suppose $\frac{k\mu}{\sqrt{d}}\frac{p_oD+k}{\tilde{\sigma}} \leq \frac{1}{16}$. For any $\mathcal{D} \in \mathcal{F}_\Xi$, there exists a network $g^*(x) = \sum_{i=1}^{n} a_i^* \sigma(\langle w_i^*, x\rangle + b_i^*)$ which satisfies*

$$\Pr_{(x,y)\sim\mathcal{D}}[yg^*(x) \leq 1] \leq \exp(-\Omega(p_oD+k)) + \exp\left(-\Omega\left(\frac{1}{\sigma_\zeta^2(k+k^2\mu/\sqrt{d})}\right)\right).$$

Furthermore, the number of neurons $n = 3(k+1)$, $|a_i^*| \leq 64k$, $1/(64k) \leq |b_i^*| \leq 1/4$, $w_i^* = \tilde{\sigma}\sum_{j\in A} M_j/(8k)$, and $|\langle w_i^*, x\rangle + b_i^*| \leq 1$ for any $i \in [n]$ and $(x,y) \sim \mathcal{D}$.

Consequently, if furthermore we have $k\mu/\sqrt{d} < 1$ and $\sigma_\zeta < 1/k$, then

$$\Pr_{(x,y)\sim\mathcal{D}}[yg^*(x) \leq 1] \leq \exp(-\Omega(k)).$$

Proof of Lemma 34. Let $w = \tilde{\sigma}\sum_{j\in A} M_j$ and let $\mu = \sum_{j\in A} \mathbb{E}[\tilde{\phi}_j]$. We have

$$\langle w, x\rangle = \tilde{\sigma}\sum_{j\in A}\langle M_j, M\phi\rangle + \langle w, \zeta/\tilde{\sigma}\rangle \tag{304}$$

$$= \sum_{j\in A}\phi_j + \sum_{j\in A, \ell\neq j}\rho_{j\ell}\phi_\ell + \langle w, \zeta/\tilde{\sigma}\rangle \tag{305}$$

$$= \sum_{j\in A}\tilde{\phi}_j - \mu + \underbrace{\sum_{j\in A, \ell\neq j}\rho_{j\ell}\phi_\ell + \langle w, \zeta/\tilde{\sigma}\rangle}_{:=\epsilon_x}. \tag{306}$$

With probability $\geq 1 - \exp(-\Theta(\max\{2p_o(D-k), k\}))$, among all $j \notin A$, we have that at most $2p_o(D-k)+k$ of ϕ_j are $(1-p_o)/\tilde{\sigma}$, while the others are $-p_o/\tilde{\sigma}$, and thus

$$\left|\sum_{j\in A, \ell\neq j}\rho_{j\ell}\phi_\ell\right| \leq \frac{k\mu}{\sqrt{d}}\frac{p_oD+k}{\tilde{\sigma}} \leq \frac{1}{16}. \tag{307}$$

Furthermore, $\langle w, \zeta\rangle \sim \mathcal{N}(0, \sigma_\zeta^2\|w\|_2^2)$ and $\|w\|_2^2 \leq \tilde{\sigma}^2(k+k^2\mu/\sqrt{d})$, we have

$$\Pr[|\langle w, \zeta/\tilde{\sigma}\rangle| \leq 1/16] \geq 1 - \exp\left(-\Theta\left(\frac{1}{\sigma_\zeta^2\|w\|_2^2/\tilde{\sigma}^2}\right)\right) \tag{308}$$

$$\geq 1 - \exp\left(-\Theta\left(\frac{1}{\sigma_\zeta^2(k+k^2\mu/\sqrt{d})}\right)\right). \tag{309}$$

For good data points with ϕ and ζ satisfying the above, we have $|\epsilon_x| \leq 1/8$. By Lemma 7,

$$g_1^*(x) := \sum_{p\in P}\delta_{p-\mu,4,1/2}(\langle w, x\rangle) - \sum_{p\notin P, 0\leq p\leq k}\delta_{p-\mu,4,1/2}(\langle w, x\rangle) \tag{310}$$

$$= \sum_{p\in P}\delta_{p,4,1/2}(\langle w, x\rangle + \mu) - \sum_{p\notin P, 0\leq p\leq k}\delta_{p,4,1/2}(\langle w, x\rangle + \mu) \tag{311}$$

$$= \sum_{p\in P}\delta_{p,4,1/2}\left(\sum_{j\in A}\tilde{\phi}_j + \epsilon_x\right) - \sum_{p\notin P, 0\leq p\leq k}\delta_{p,4,1/2}\left(\sum_{j\in A}\tilde{\phi}_j + \epsilon_x\right). \tag{312}$$

Then for good data points, we have $yg_1^*(x) \geq 1$. Similarly,

$$g_2^*(x) := \sum_{p\in P}\delta_{p-\mu+1/4,8,1/2}(\langle w, x\rangle) - \sum_{p\notin P, 0\leq p\leq k}\delta_{p-\mu+1/4,8,1/2}(\langle w, x\rangle) \tag{313}$$

$$= \sum_{p\in P}\delta_{p+1/4,8,1/2}(\langle w, x\rangle + \mu) - \sum_{p\notin P, 0\leq p\leq k}\delta_{p+1/4,8,1/2}(\langle w, x\rangle + \mu) \tag{314}$$

$$= \sum_{p\in P}\delta_{p+1/4,8,1/2}\left(\sum_{j\in A}\tilde{\phi}_j + \epsilon_x\right) - \sum_{p\notin P, 0\leq p\leq k}\delta_{p+1/4,8,1/2}\left(\sum_{j\in A}\tilde{\phi}_j + \epsilon_x\right). \tag{315}$$

Then for good data points, we have $yg_2^*(x) \geq 1$.

Note that the bias terms in g_1^* and g_2^* have distance at least $1/4$, then at least one of them satisfies that all its bias terms have absolute value $\geq 1/8$. Pick that one and denote it as $g(x) = \sum_{i=1}^n a_i \sigma_{\mathrm{r}}(\langle w_i, x\rangle + b_i)$. By the positive homogeneity of σ_{r}, we have

$$g(x) = \sum_{i=1}^n 8ka_i\sigma_{\mathrm{r}}(\langle w_i, x\rangle/(8k) + b_i/(8k)). \tag{316}$$

Since for any good data points, $|\langle w_i, x\rangle/(8k) + b_i/(8k)| \leq 1$, then

$$g(x) = \sum_{i=1}^n 8ka_i\sigma(\langle w_i, x\rangle/(8k) + b_i/(8k)) \tag{317}$$

where σ is the truncated ReLU. Now we can set $a_i^* = 8ka_i, w_i^* = w_i/(8k), b_i^* = b_i/(8k)$, to get our final g^*. $\qquad\square$

F.5 INITIALIZATION

We first show that with high probability, the initial weights are in typical positions.

Lemma 35. *Suppose* $D\mu/\sqrt{d} \leq 1/16$. *For any* $\delta \in (0, 1)$, *with probability at least* $1 - \delta - 2\exp\left(-\Theta(D - k)\right)$ *over* $w^{(0)}$,

$$\sigma_w^2 d/2 \leq \|w^{(0)}\|_2^2 \leq 3\sigma_w^2 d/2,$$

$$\sigma_w^2(D - k)/2 \leq \sum_{\ell \notin A} q_\ell^2 \leq 3\sigma_w^2(D - k)/2,$$

$$\max_\ell |q_\ell| \leq \sigma_w\sqrt{2\log(D/\delta)}.$$

With probability at least $1 - \delta$ *over* $b^{(0)}$,

$$|b^{(0)}| \leq \sigma_b\sqrt{2\log(1/\delta)}.$$

With probability at least $1 - \delta$ *over* $a^{(0)}$,

$$|a^{(0)}| \leq \sigma_a\sqrt{2\log(1/\delta)}.$$

Proof of Lemma 35. The bound on $\|w^{(0)}\|_2^2$ follows from the property of Gaussians.

Note that $q = M^\top w^{(0)} \sim \mathcal{N}(0, \sigma_w^2\rho)$. We have with probability $\geq 1 - \delta/2$, $\max_\ell |q_\ell| \leq \sqrt{2\sigma_w^2\log\frac{D}{\delta}}$.

For any subset $S \subseteq [D]$, let ρ_S denote the submatrix of ρ containing the rows and columns indexed by S. Then $q_S = M^\top w^{(0)} \sim \mathcal{N}(0, \sigma_w^2\rho_S)$. By diagonalizing ρ_S and then applying Bernstein's inequality, we have With probability $\geq 1 - 2\exp\left(-\Theta(|S|/\|\rho\|_2)\right)$, $\|q_S\|_2^2 \in \left((\|\rho_S\|_F^2 - \frac{|S|}{4})\sigma_w^2, (\|\rho_S\|_F^2 + \frac{|S|}{4})\sigma_w^2\right)$. By Gershgorin circle theorem, we have

$$\|\rho\|_2 \leq 1 + (|S| - 1)\mu/\sqrt{d} \leq 17/16.$$

Similarly, we have

$$\frac{3}{4}|S| \leq \left(\frac{15}{16}\right)^2|S| \leq \|\rho_S\|_F^2 \leq \left(\frac{17}{16}\right)^2|S| \leq \frac{5}{4}|S|.$$

The bounds on q then follow.

The bounds on $b^{(0)}$ and $a^{(0)}$ follow from the property of Gaussians. $\qquad\square$

Lemma 36. *Suppose* $D\mu/\sqrt{d} \leq 1/16$. *We have:*

- *With probability $\geq 1-\delta-2m\exp(-\Theta(D-k))$ over $w_i^{(0)}$'s, for all $i \in [2m]$, $w_i^{(0)} \in \mathcal{G}_w(\delta)$.*

- *With probability $\geq 1-\delta$ over $b_i^{(0)}$'s, for all $i \in [2m]$, $b_i^{(0)} \in \mathcal{G}_b(\delta)$.*

- *With probability $\geq 1-\delta$ over $a_i^{(0)}$'s, for all $i \in [2m]$, $a_i^{(0)} \in \mathcal{G}_a(\delta)$.*

Proof of Lemma 36. This follows from Lemma 35 by union bound. $\qquad\square$

Then we get the same lemma as before about the typical $w_i^{(0)}$'s as before:

Lemma 37. *Fix $\delta \in (0,1)$. For any $w_i^{(0)} \in \mathcal{G}_w(\delta)$, we have*

$$\Pr_\phi \left[\sum_{\ell \notin A} \phi_\ell q_{i,\ell}^{(0)} \geq \Theta\left(\sqrt{(D-k)\sigma_\phi^2\sigma_w^2}\right) \right] = \Theta(1) - \frac{O(\log^{3/2}(Dm/\delta))}{\sqrt{(D-k)\sigma_\phi^2\tilde{\sigma}^2}}. \tag{318}$$

Consequently, when $p_o = \Omega(k^2/D)$ and $k = \Omega(\log^2(Dm/\delta))$,

$$\Pr_\phi \left[\sum_{\ell \notin A} \phi_\ell q_{i,\ell}^{(0)} \geq \Theta(\sigma_w) \right] = \Theta(1) - \frac{O(1)}{k^{1/4}}. \tag{319}$$

Proof of Lemma 37. Note that for $\ell \notin A$, $\mathbb{E}[\phi_\ell] = 0$, $\mathbb{E}[\phi_\ell^2] = \sigma_\phi^2$, and $\mathbb{E}[|\phi_\ell|^3] = \Theta(\sigma_\phi^2/\tilde{\sigma})$. Then the statement follows from Berry-Esseen Theorem. $\qquad\square$

F.6   SOME AUXILIARY LEMMAS

The expression of the gradients will be used frequently.

Lemma 38.

$$\frac{\partial}{\partial w_i} L_\mathcal{D}^\alpha(g;\sigma_\xi) = -a_i \mathbb{E}_{(x,y)\sim\mathcal{D}} \left\{ \alpha_y y \mathbb{I}[yg(x;\xi) \leq 1] \mathbb{E}_{\xi_i} \mathbb{I}[\langle w_i, x\rangle + b_i + \xi_i \in (0,1)]x \right\}, \tag{320}$$

$$\frac{\partial}{\partial b_i} L_\mathcal{D}^\alpha(g;\sigma_\xi) = -a_i \mathbb{E}_{(x,y)\sim\mathcal{D}} \left\{ \alpha_y y \mathbb{I}[yg(x;\xi) \leq 1] \mathbb{E}_{\xi_i} \mathbb{I}[\langle w_i, x\rangle + b_i \in (0,1)] \right\}, \tag{321}$$

$$\frac{\partial}{\partial a_i} L_\mathcal{D}^\alpha(g;\sigma_\xi) = -\mathbb{E}_{(x,y)\sim\mathcal{D}} \left\{ \alpha_y y \mathbb{I}[yg(x;\xi) \leq 1] \mathbb{E}_{\xi_i} \sigma(\langle w_i, x\rangle + b_i + \xi_i) \right\}. \tag{322}$$

Proof of Lemma 38. It follows from straightforward calculation. $\qquad\square$

We also have the following auxiliary lemma for later calculations.

Lemma 39.

$$\mathbb{E}_{\phi_A}\left\{ \alpha_y y \right\} = 0, \tag{323}$$

$$\mathbb{E}_{\phi_A}\left\{ |\alpha_y y| \right\} = 1, \tag{324}$$

$$\mathbb{E}_{\phi_j}\left\{ |\phi_j| \right\} = 2\sigma_\phi^2\tilde{\sigma}, \text{ for } j \notin A, \tag{325}$$

$$\mathbb{E}_{\phi_A}\left\{ \alpha_y y \phi_j \right\} = \frac{\gamma}{\tilde{\sigma}}, \tag{326}$$

$$\mathbb{E}_{\phi_A}\left\{ |\alpha_y y \phi_j| \right\} \leq \frac{1}{\tilde{\sigma}}, \text{ for all } j \in [D]. \tag{327}$$

Proof of Lemma 39.

$$\mathbb{E}_{\phi_A}\{\alpha_y y\} = \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{\alpha_y y | y = v\} \Pr[y = v] \tag{328}$$

$$= \frac{1}{2} \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{y | y = v\} \tag{329}$$

$$= 0. \tag{330}$$

$$\mathbb{E}_{\phi_A}\{|\alpha_y y|\} = \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{|\alpha_y y| \, | y = v\} \Pr[y = v] \tag{331}$$

$$= \frac{1}{2} \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{|y| \, | y = v\} \tag{332}$$

$$= 1. \tag{333}$$

$$\mathbb{E}_{\phi_j}\{|\phi_j|\} = \frac{|-p_o|(1 - p_o) + |1 - p_o|p_o}{\tilde{\sigma}} = 2\sigma_\phi^2 \tilde{\sigma}. \tag{334}$$

$$\mathbb{E}_{\phi_A}\{\alpha_y y \phi_j\} = \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{\alpha_y y \phi_j | y = v\} \Pr[y = v] \tag{335}$$

$$= \frac{1}{2} \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{y \phi_j | y = v\} \tag{336}$$

$$= \frac{1}{2} \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\left\{ y \frac{\tilde{\phi}_j - \mathbb{E}[\tilde{\phi}_j]}{\tilde{\sigma}} \bigg| y = v \right\} \tag{337}$$

$$= \frac{1}{2\tilde{\sigma}}(\gamma_{+1} + \gamma_{-1}) = \frac{\gamma}{\tilde{\sigma}}. \tag{338}$$

$$\mathbb{E}_{\phi_A}\{|\alpha_y y \phi_j|\} = \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{|\alpha_v y \phi_j| \, | y = v\} \Pr[y = v] \tag{339}$$

$$\leq \frac{1}{2} \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{|y \phi_j| \, | y = v\} \tag{340}$$

$$\leq \frac{1}{2} \sum_{v \in \{\pm 1\}} \mathbb{E}_{\phi_A}\{|y \phi_j| \, | y = v\} \tag{341}$$

$$\leq \frac{1}{\tilde{\sigma}}. \tag{342}$$

$\square$

F.7 FEATURE EMERGENCE: FIRST GRADIENT STEP

We will show that w.h.p. over the initialization, after the first gradient step, there are neurons that represent good features.

We begin with analyzing the gradients.

Lemma 40. *Fix $\delta \in (0, 1)$ and suppose $w_i^{(0)} \in \mathcal{G}_w(\delta), b_i^{(0)} \in \mathcal{G}_b(\delta)$ for all $i \in [2m]$. Let*

$$\epsilon_e := \frac{k \log^{1/2}(Dm/\delta) + \log^{3/2}(Dm/\delta)}{\sqrt{\sigma_\phi^2 \tilde{\sigma}^2 (D - k)}},$$

$$\epsilon_\nu := \frac{\sigma_\zeta \sigma_w \log(Dm/\delta)/\tilde{\sigma} + \log^{3/2}(Dm/\delta)}{\sqrt{\sigma_\phi^2 \tilde{\sigma}^2 (D - k)}},$$

If $\sigma_\zeta^2 \sigma_w^2 d / \tilde{\sigma}^2 = O(1/k)$, $p_o = \Omega(k^2/D)$, $k = \Omega(\log^2(Dmd/\delta))$, and $\sigma_\xi^{(1)} \leq 1/k$, then

$$\frac{\partial}{\partial w_i} L_\mathcal{D}^\alpha(g^{(0)}; \sigma_\xi^{(1)}) = -a_i^{(0)} \left(\sum_{j=1}^D M_j T_j + \nu \right) \tag{343}$$

where T_j satisfies:

- if $j \in A$, then $|T_j - \beta\gamma/\tilde{\sigma}| \leq O(\epsilon_e/\tilde{\sigma})$, where $\beta \in [\Omega(1), 1]$ and depends only on $w_i^{(0)}, b_i^{(0)}$;

- if $j \notin A$, then $|T_j| \leq O(\sigma_\phi^2 \epsilon_e \tilde{\sigma})$;

- $|\nu_j| \leq O\left(\frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} \epsilon_\nu \right) + \frac{\sigma_\zeta d}{\tilde{\sigma}} e^{-\Theta(k)}$.

Proof of Lemma 40. Consider one neuron index i and omit the subscript i in the parameters. Since the unbiased initialization leads to $g^{(0)}(x; \xi^{(1)}) = 0$, we have

$$\frac{\partial}{\partial w} L_\mathcal{D}^\alpha(g^{(0)}; \sigma_\xi^{(1)}) \tag{344}$$

$$= -a^{(0)} \mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ \alpha_y y \mathbb{I}[y g^{(0)}(x; \xi^{(1)}) \leq 1] \mathbb{E}_{\xi^{(1)}} \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] x \right\} \tag{345}$$

$$= -a^{(0)} \mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ \alpha_y y \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] x \right\} \tag{346}$$

$$= -a^{(0)} \sum_{j=1}^D M_j \underbrace{\mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ \alpha_y y \phi_j \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\}}_{:= T_j} \tag{347}$$

$$- a^{(0)} \underbrace{\mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta}{\tilde{\sigma}} \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\}}_{:= \nu} \tag{348}$$

First, consider $j \in A$.

$$T_j = \mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ \alpha_y y \phi_j \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \tag{349}$$

$$= \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \alpha_y y \phi_j \Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0, 1) \right] \right\}. \tag{350}$$

where $\iota := \langle w^{(0)}, \zeta/\tilde{\sigma} \rangle$. Let

$$I_a := \Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + b^{(0)} + \iota + \xi^{(1)} \in (0, 1) \right], \tag{351}$$

$$I_a' := \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0, 1) \right]. \tag{352}$$

We have

$$|\mathbb{E}_{\zeta, \xi^{(1)}}(I_a - I_a')| \tag{353}$$

$$\leq \mathbb{E}_{\zeta, \xi^{(1)}} \left| \Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \geq 0 \right] - \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \geq 0 \right] \right| \tag{354}$$

$$+ \Pr_{\phi_{-A}, \zeta, \xi^{(1)}} \left[\langle \phi, q \rangle + b^{(0)} + \iota + \xi^{(1)} \geq 1 \right] + \Pr_{\phi_{-A}, \zeta, \xi^{(1)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \geq 1 \right]. \tag{355}$$

Then by Lemma 13,

$$\left| \Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \geq 0 \right] - \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \geq 0 \right] \right| = O(\epsilon_e). \tag{356}$$

Note that $\sum_{\ell \notin A} \mathrm{Var}(\phi_\ell q_\ell) = \Theta(\sigma_\phi^2 \sigma_w^2(D-k)) = \Theta(\sigma_w^2)$, and $|\phi_\ell| \leq \frac{1}{\tilde{\sigma}}$. If $p_{\mathrm{o}} = \Omega(k^3/D)$ then apply Bennett's inequality in the small-deviation regime; $p_{\mathrm{o}} = O(k^3/D)$ and $p_{\mathrm{o}} = \Omega(k^2/D)$ then apply Bennett's inequality in the large-deviation regime. In both cases we have:

$$\Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle \geq 1/4 \right] = \exp(-\Omega(k)) = O(\epsilon_{\mathrm{e}}). \tag{357}$$

We also have:

$$\Pr_{\xi^{(1)}} \left[b^{(0)} + \xi^{(1)} \geq 1/4 \right] = \exp(-\Omega(k)) = O(\epsilon_{\mathrm{e}}). \tag{358}$$

Note that $\iota \sim \mathcal{N}(0, \Theta(\sigma_\zeta^2 \sigma_w^2 d/\tilde{\sigma}^2))$, and by assumption on σ_ζ we have $\sigma_\zeta^2 \sigma_w^2 d/\tilde{\sigma}^2 = O(1/k)$, so

$$\Pr_{\zeta} \left[\iota \geq 1/4 \right] = \exp(-\Omega(k)) = O(\epsilon_{\mathrm{e}}). \tag{359}$$

Therefore,

$$\Pr_{\phi_{-A}, \zeta, \xi^{(1)}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \geq 1 \right] = \exp(-\Omega(k)) = O(\epsilon_{\mathrm{e}}) \tag{360}$$

where the last step follows from the assumption on σ_w and k. A similar argument gives:

$$\Pr_{\phi_{-A}, \zeta, \xi^{(1)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \geq 1 \right] = \exp(-\Omega(k)) = O(\epsilon_{\mathrm{e}}). \tag{361}$$

In summary,

$$|\mathbb{E}_{\zeta, \xi^{(1)}}(I_a - I_a')| = O(\epsilon_{\mathrm{e}}). \tag{362}$$

Then we have

$$\left| T_j - \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \alpha_y y \phi_j I_a' \right\} \right| \tag{363}$$

$$\leq \mathbb{E}_{\phi_A} \left\{ |\alpha_y y \phi_j| \left| \mathbb{E}_{\zeta, \xi^{(1)}} (I_a - I_a') \right| \right\} \tag{364}$$

$$\leq O(\epsilon_{\mathrm{e}}) \mathbb{E}_{\phi_A} \left\{ |\alpha_y y \phi_j| \right\} \tag{365}$$

$$\leq O(\epsilon_{\mathrm{e}}/\tilde{\sigma}) \tag{366}$$

where the last step is from Lemma 39. Furthermore,

$$\mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \alpha_y y \phi_j I_a' \right\} \tag{367}$$

$$= \mathbb{E}_{\phi_A} \left\{ \alpha_y y \phi_j \right\} \mathbb{E}_{\zeta, \xi^{(1)}} [I_a'] \tag{368}$$

$$= \mathbb{E}_{\phi_A} \left\{ \alpha_y y \phi_j \right\} \Pr_{\phi_{-A}, \zeta, \xi^{(1)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \tag{369}$$

By Lemma 11, the assumption on p_{o}, and (357), we have

$$\Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + b^{(0)} \in (0, 1/2) \right] \geq \Omega(1) - O(1/k^{1/4}), \tag{370}$$

$$\Pr_{\zeta, \xi^{(1)}} \left[\iota + \xi^{(1)} \in (0, 1/2) \right] = 1/2 - \exp(-\Omega(k)), \tag{371}$$

This leads to

$$\beta := \mathbb{E}_{\zeta, \xi^{(1)}} [I_a'] = \Pr_{\phi_{-A}, \zeta, \xi^{(1)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \geq \Omega(1). \tag{372}$$

By Lemma 39, $\mathbb{E}_{\phi_A} \left\{ \alpha_y y \phi_j \right\} = \gamma/\tilde{\sigma}$. Therefore,

$$|T_j - \beta\gamma/\tilde{\sigma}| \leq O(\epsilon_{\mathrm{e}}/\tilde{\sigma}). \tag{373}$$

Now, consider $j \notin A$. Let B denote $A \cup \{j\}$.

$$T_j = \mathbb{E}_{(x,y) \sim \mathcal{D}, \zeta, \xi^{(1)}} \left\{ \alpha_y y \phi_j \mathbb{I} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right\} \tag{374}$$

$$= \mathbb{E}_{\phi_B} \mathbb{E}_{\phi_{-B}, \zeta, \xi^{(1)}} \left\{ \alpha_y y \phi_j \mathbb{I} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right\} \tag{375}$$

$$= \mathbb{E}_{\phi_B, \zeta, \xi^{(1)}} \left\{ \alpha_y y \phi_j \Pr_{\phi_{-B}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right\}. \tag{376}$$

Let

$$I_b := \Pr_{\phi_{-B}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0, 1) \right], \tag{377}$$

$$I_b' := \Pr_{\phi_{-B}} \left[\langle \phi_{-B}, q_{-B} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0, 1) \right]. \tag{378}$$

Similar as above, we have $|\mathbb{E}_{\zeta, \xi^{(1)}} (I_b - I_b')| \le O(\epsilon_e)$ by Lemma 13. Then by Lemma 39,

$$\left| T_j - \mathbb{E}_{\phi_B, \zeta, \xi^{(1)}} \left\{ \alpha_y y \phi_j I_b' \right\} \right| \tag{379}$$

$$\le \mathbb{E}_{\phi_B} \left\{ |\alpha_y y \phi_j| |\mathbb{E}_{\zeta, \xi^{(1)}} (I_b - I_b')| \right\} \tag{380}$$

$$\le O(\epsilon_e) \mathbb{E}_{\phi_A} \left\{ |\alpha_y y| \right\} \mathbb{E}_{\phi_j} \left\{ |\phi_j| \right\} \tag{381}$$

$$\le O(\epsilon_e) \times 1 \times O(\sigma_\phi^2 \tilde{\sigma}) \tag{382}$$

$$= O(\sigma_\phi^2 \epsilon_e \tilde{\sigma}). \tag{383}$$

Furthermore,

$$\mathbb{E}_{\phi_B, \zeta, \xi^{(1)}} \left\{ \alpha_y y \phi_j I_b' \right\} = \mathbb{E}_{\phi_A} \left\{ \alpha_y y \right\} \mathbb{E}_{\phi_j} \left\{ \phi_j \right\} \mathbb{E}_{\zeta, \xi^{(1)}} [I_b'] = 0. \tag{384}$$

Therefore,

$$|T_j| \le O(\sigma_\phi^2 \epsilon_e \tilde{\sigma}). \tag{385}$$

Finally, consider ν_j.

$$\nu_j = \mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \tag{386}$$

$$= \mathbb{E}_{\phi_A, \phi_{-A}, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} \mathbb{I}[\langle \phi, q \rangle + \iota_j + \iota_{-j} + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \tag{387}$$

$$= \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} \Pr_{\phi_{-A}} [\langle \phi, q \rangle + \iota_j + \iota_{-j} + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \tag{388}$$

where $\iota_j := w_j^{(0)} \zeta_j / \tilde{\sigma}$ and $\iota_{-j} := \langle w^{(0)}, \zeta/\tilde{\sigma} \rangle - \iota_j$.

With probability $\ge 1 - d \exp(-\Theta(k))$ over ζ, for any j, $|\zeta_j| \le O(\sigma_\zeta \sqrt{\log(k)})$. Let $\mathcal{G}_\zeta$ denote this event.

Let

$$I_j := \Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + \iota_j + \iota_{-j} + b^{(0)} + \xi^{(1)} \in (0, 1) \right], \tag{389}$$

$$I_j' := \Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + \iota_{-j} + b^{(0)} + \xi^{(1)} \in (0, 1) \right]. \tag{390}$$

Similar as in Lemma 13, we have $|\mathbb{E}_{\zeta, \xi^{(1)}}[I_j - I_j' | \mathcal{G}_\zeta]| \le O(\epsilon_\nu)$. Then

$$|\mathbb{E}_{\zeta, \xi^{(1)}} (I_b - I_b')| \le |\mathbb{E}_{\zeta, \xi^{(1)}} [(I_b - I_b') | \mathcal{G}_\zeta]| + \Pr[-\mathcal{G}_\zeta] \tag{391}$$

$$\le O(\epsilon_\nu + d \exp(-\Theta(k))). \tag{392}$$

$$\left| \nu_j - \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} I_j' \right\} \right| \tag{393}$$

$$= \left| \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} (I_j - I_j') \right\} \right| \tag{394}$$

$$\le \left| \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} (I_j - I_j') | \mathcal{G}_\zeta \right\} \right| + \left| \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} (I_j - I_j') | -\mathcal{G}_\zeta \right\} \right| \Pr[-\mathcal{G}_\zeta]. \tag{395}$$

The first term is bounded by

$$\left| \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} (I_j - I'_j) | \mathcal{G}_\zeta \right\} \right| \tag{396}$$

$$\leq \mathbb{E}_{\phi_A} \left\{ \frac{\alpha_y y \sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} |\mathbb{E}_{\zeta, \xi^{(1)}}[I_b - I'_b | \mathcal{G}_\zeta]| \right\} \tag{397}$$

$$\leq O(\epsilon_\nu) \mathbb{E}_{\phi_A} \left\{ |\alpha_y y| \right\} \frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} \tag{398}$$

$$\leq O(\epsilon_\nu) \times 1 \times \frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} \tag{399}$$

$$= O\left(\frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} \epsilon_\nu \right). \tag{400}$$

The second term is bounded by

$$\left| \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} (I_j - I'_j) | - \mathcal{G}_\zeta \right\} \right| \Pr[-\mathcal{G}_\zeta] \tag{401}$$

$$\leq \left| \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} (I_j - I'_j) | - \mathcal{G}_\zeta \right\} \right| \times d e^{-\Theta(k)} \tag{402}$$

$$\leq \mathbb{E}_{\phi_A} \left| \frac{\alpha_y y}{\tilde{\sigma}} \right| \times \mathbb{E}_\zeta \left\{ |\zeta_j| | - \mathcal{G}_\zeta \right\} \times d e^{-\Theta(k)} \tag{403}$$

$$\leq \frac{\sigma_\zeta}{\tilde{\sigma}} \times d e^{-\Theta(k)} \tag{404}$$

$$\leq \frac{\sigma_\zeta d}{\tilde{\sigma}} e^{-\Theta(k)}. \tag{405}$$

Furthermore,

$$\mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} I'_j \right\} = \mathbb{E}_{\phi_A} \left\{ \alpha_y y \right\} \mathbb{E}_{\zeta_j} \left\{ \frac{\zeta_j}{\tilde{\sigma}} \right\} \mathbb{E}_{\zeta_{-j}}[I'_j] = 0. \tag{406}$$

Therefore,

$$|\nu_j| \leq O\left(\frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} \epsilon_\nu \right) + \frac{\sigma_\zeta d}{\tilde{\sigma}} e^{-\Theta(k)}. \tag{407}$$

$\square$

Lemma 41. *Under the same assumptions as in Lemma 40,*

$$\frac{\partial}{\partial b_i} L_{\mathcal{D}}^\alpha(g^{(0)}; \sigma_\xi^{(1)}) = -a_i^{(0)} T_b \tag{408}$$

where $|T_b| \leq O(\epsilon_e)$.

Proof of Lemma 41. Consider one neuron index i and omit the subscript i in the parameters. Since the unbiased initialization leads to $g^{(0)}(x; \xi^{(1)}) = 0$, we have

$$\frac{\partial}{\partial b} L_{\mathcal{D}}^\alpha(g^{(0)}; \sigma_\xi^{(1)}) \tag{409}$$

$$= -a^{(0)} \mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ \alpha_y y \mathbb{I}[y g^{(0)}(x; \xi^{(1)}) \leq 1] \mathbb{E}_{\xi^{(1)}} \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \tag{410}$$

$$= -a^{(0)} \mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ \alpha_y y \mathbb{I}[\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)} \in (0, 1)] \right\} \tag{411}$$

$$= -a^{(0)} \underbrace{\mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \alpha_y y \Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0, 1) \right] \right\}}_{:= T_b}. \tag{412}$$

where $\iota := \langle w^{(0)}, \zeta/\tilde{\sigma}\rangle$. Similar to the proof in Lemma 40,

$$\left| \mathbb{E}_{\zeta, \xi^{(1)}} \left(\Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right. \right. \tag{413}$$

$$\left. \left. - \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right) \right| = O(\epsilon_e). \tag{414}$$

Then

$$\left| T_b - \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \alpha_y y \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right\} \right| \tag{415}$$

$$= \mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ |\alpha_y y| \left| \Pr_{\phi_{-A}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right. \right. \tag{416}$$

$$\left. \left. - \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right| \right\} \tag{417}$$

$$\leq O(\epsilon_e) \mathbb{E}_{\phi_A} \left\{ |\alpha_y y| \right\} \tag{418}$$

$$\leq O(\epsilon_e). \tag{419}$$

Also,

$$\mathbb{E}_{\phi_A, \zeta, \xi^{(1)}} \left\{ \alpha_y y \Pr_{\phi_{-A}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \right\} \tag{420}$$

$$= \mathbb{E}_{\phi_A} \left\{ \alpha_y y \right\} \Pr_{\phi_{-A}, \zeta, \xi^{(1)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right] \tag{421}$$

$$= 0. \tag{422}$$

Therefore, $|T_b| \leq O(\epsilon_e)$. $\qquad\square$

Lemma 42. *We have*

$$\frac{\partial}{\partial a_i} L_{\mathcal{D}}^{\alpha}(g^{(0)}; \sigma_{\xi}^{(1)}) = -T_a \tag{423}$$

where $|T_a| \leq O(\max_{\ell} q_{i,\ell}^{(0)})$. *So if* $w_i^{(0)} \in \mathcal{G}(\delta)$, $|T_a| \leq O(\sigma_w \sqrt{\log(Dm/\delta)})$.

Proof of Lemma 42. Consider one neuron index i and omit the subscript i in the parameters. Since the unbiased initialization leads to $g^{(0)}(x; \xi^{(1)}) = 0$, we have

$$\frac{\partial}{\partial a} L_{\mathcal{D}}^{\alpha}(g^{(0)}; \sigma_{\xi}^{(1)}) \tag{424}$$

$$= -\mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ \alpha_y y \mathbb{I}[y g^{(0)}(x; \xi^{(1)}) \leq 1] \mathbb{E}_{\xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) \right\} \tag{425}$$

$$= -\underbrace{\mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(1)}} \left\{ \alpha_y y \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) \right\}}_{:=T_a}. \tag{426}$$

Let ϕ'_A be an independent copy of ϕ_A, ϕ' be the vector obtained by replacing in ϕ the entries ϕ_A with ϕ'_A, and let $x' = M\phi' + \zeta/\tilde{\sigma}$ and its label is y'. Then

$$|T_a| = \left| \mathbb{E}_{\phi_A} \left\{ \alpha_y y \mathbb{E}_{\phi_{-A}, \zeta, \xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) \right\} \right| \tag{427}$$

$$\leq \frac{1}{2} \left| \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \zeta, \xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) | y = 1 \right\} \right. \tag{428}$$

$$\left. - \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \zeta, \xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) | y = -1 \right\} \right| \tag{429}$$

$$\leq \frac{1}{2} \left| \mathbb{E}_{\phi_A} \left\{ \mathbb{E}_{\phi_{-A}, \zeta, \xi^{(1)}} \sigma(\langle w^{(0)}, x \rangle + b^{(0)} + \xi^{(1)}) | y = 1 \right\} \right. \tag{430}$$

$$\left. - \mathbb{E}_{\phi'_A} \left\{ \mathbb{E}_{\phi_{-A}, \zeta, \xi^{(1)}} \sigma(\langle w^{(0)}, x' \rangle + b^{(0)} + \xi^{(1)}) | y' = -1 \right\} \right|. \tag{431}$$

Since σ is 1-Lipschitz,

$$|T_a| \leq \frac{1}{2} \mathbb{E}_{\phi_A, \phi'_A} \left\{ \mathbb{E}_{\phi_{-A}} \left| \langle w^{(0)}, M\phi \rangle - \langle w^{(0)}, M\phi' \rangle \right| | y = 1, y' = -1 \right\} \tag{432}$$

$$\leq \frac{1}{2} \mathbb{E}_{\phi_{-A}} \left(\mathbb{E}_{\phi_A} \left\{ \left| \langle w^{(0)}, M\phi \rangle \right| | y = 1 \right\} + \mathbb{E}_{\phi'_A} \left\{ \left| \langle w^{(0)}, M\phi' \rangle \right| | y' = -1 \right\} \right) \tag{433}$$

$$\leq \max_\ell q_{i,\ell}^{(0)} \sqrt{ \mathbb{E}_\phi \left(\sum_{\ell \in [D]} \phi_\ell^2 + \sum_{j \neq \ell : j, \ell \in A} |\phi_j \phi_\ell| \right) } \tag{434}$$

$$\leq \max_\ell q_{i,\ell}^{(0)} \sqrt{ \mathbb{E}_\phi (1 + O(1)) } \tag{435}$$

$$= \Theta(\max_\ell q_{i,\ell}^{(0)}). \tag{436}$$

$\square$

With the bounds on the gradient, we now summarize the results for the weights after the first gradient step.

Lemma 43. *Set*
$$\lambda_w^{(1)} = 1/(2\eta^{(1)}), \lambda_a^{(1)} = \lambda_b^{(1)} = 0, \sigma_\xi^{(1)} = 1/k^{3/2}.$$

Fix $\delta \in (0,1)$ and suppose $w_i^{(0)} \in \mathcal{G}_w(\delta), b_i^{(0)} \in \mathcal{G}_b(\delta)$ for all $i \in [2m]$. If $p_o = \Omega(k^2/D)$, $k = \Omega(\log^2(Dm/\delta))$, then for all $i \in [m]$, $w_i^{(1)} = \sum_{\ell=1}^D q_{i,\ell}^{(1)} M_\ell + v$ satisfying

- *if $\ell \in A$, then $|q_{i,\ell}^{(1)} - \eta^{(1)} a_i^{(0)} \beta \gamma / \tilde{\sigma}| \leq O\left(\frac{|\eta^{(1)} a_i^{(0)}| \epsilon_e}{\tilde{\sigma}} \right)$, where $\beta \in [\Omega(1), 1]$ and depends only on $w_i^{(0)}, b_i^{(0)}$;*

- *if $\ell \notin A$, then $|q_{i,\ell}^{(1)}| \leq O\left(|\eta^{(1)} a_i^{(0)}| \sigma_\phi^2 \epsilon_e \tilde{\sigma} \right)$;*

- *$|v_j| \leq O\left(|\eta^{(1)} a_i^{(0)}| \left(\frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} \epsilon_\nu + \frac{\sigma_\zeta d}{\tilde{\sigma}} e^{-\Theta(k)} \right) \right).$*

and

- *$b_i^{(1)} = b_i^{(0)} + \eta^{(1)} a_i^{(0)} T_b$ where $|T_b| = O(\epsilon_e)$;*

- *$a_i^{(1)} = a_i^{(0)} + \eta^{(1)} T_a$ where $|T_a| = O(\sigma_w \sqrt{\log(Dm/\delta)})$.*

Proof of Lemma 43. This follows from Lemma 36 and Lemma 40-42. $\square$

F.8 Feature Improvement: Second Gradient Step

We first show that with properly set $\eta^{(1)}$, for most x, $|g^{(1)}(x; \sigma_\xi^{(2)})| < 1$ and thus $yg^{(1)}(x; \sigma_\xi^{(2)}) < 1$.

Lemma 44. *Fix* $\delta \in (0,1)$ *and suppose* $w_i^{(0)} \in \mathcal{G}_w(\delta), b_i^{(0)} \in \mathcal{G}_b(\delta), a_i^{(0)} \in \mathcal{G}_a(\delta)$ *for all* $i \in [2m]$. *If* $D\mu/\sqrt{d} \leq 1/16$, $\sigma_\zeta \tilde{\sigma} = O(1)$, $\sigma_\zeta^2 d/\tilde{\sigma}^2 = O(1)$, $p_o = \Omega(k^2/D)$, $k = \Omega(\log^2(Dm/\delta))$, $\sigma_a \leq \tilde{\sigma}^2/(\gamma k^2)$, $\eta^{(1)} = O\left(\frac{\gamma}{km\sigma_a\tilde{\sigma}}\right)$, *and* $\sigma_\xi^{(2)} \leq 1/k$, *then with probability* $\geq 1 - (d+D)\exp(-\Omega(k))$ *over* (x,y), *we have* $yg^{(1)}(x; \sigma_\xi^{(2)}) < 1$. *Furthermore, for any* $i \in [2m]$, $\left|\langle w_i^{(1)}, \zeta/\tilde{\sigma}\rangle\right| = O(\eta^{(1)}\tilde{\sigma}/\gamma)$, $\left|\langle q_i^{(1)}, \phi\rangle\right| = O(\eta^{(1)}\tilde{\sigma}/\gamma)$, *and* $\left|\langle (q_i^{(1)})_{-A}, \phi_{-A}\rangle\right| = O(\eta^{(1)}\tilde{\sigma}/\gamma)$, *and for any* $j \in [d], \ell \in [D]$, $|\zeta_j| \leq O(\sigma_\zeta\sqrt{\log(k)})$ *and* $|\langle \zeta, D_\ell\rangle| \leq O(\sigma_\zeta\sqrt{\log(k)})$.

Proof of Lemma 44. Note that $w_i^{(0)} = w_{m+i}^{(0)}$, $b_i^{(0)} = b_{m+i}^{(0)}$, and $a_i^{(0)} = a_{m+i}^{(0)}$. Then the gradient for w_{m+i} is the negation of that for w_{m+i}, the gradient for b_{m+i} is the negation of that for b_{m+i}, and the gradient for a_{m+i} is the same as that for a_{m+i}.

$$\left|g^{(1)}(x; \sigma_\xi^{(2)})\right| \tag{437}$$

$$= \left|\sum_{i=1}^{2m} a_i^{(1)}\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_i^{(1)}, x\rangle + b_i^{(1)} + \xi_i^{(2)})\right| \tag{438}$$

$$= \left|\sum_{i=1}^{m}\left(a_i^{(1)}\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_i^{(1)}, x\rangle + b_i^{(1)} + \xi_i^{(2)}) + a_{m+i}^{(1)}\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_{m+i}^{(1)}, x\rangle + b_{m+i}^{(1)} + \xi_{m+i}^{(2)})\right)\right| \tag{439}$$

$$\leq \left|\sum_{i=1}^{m}\left(a_i^{(1)}\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_i^{(1)}, x\rangle + b_i^{(1)} + \xi_i^{(2)}) + a_{m+i}^{(1)}\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_i^{(1)}, x\rangle + b_i^{(1)} + \xi_i^{(2)})\right)\right| \tag{440}$$

$$+ \left|\sum_{i=1}^{m}\left(-a_{m+i}^{(1)}\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_i^{(1)}, x\rangle + b_i^{(1)} + \xi_i^{(2)}) + a_{m+i}^{(1)}\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_{m+i}^{(1)}, x\rangle + b_{m+i}^{(1)} + \xi_i^{(2)})\right)\right|. \tag{441}$$

Then we have

$$\left|g^{(1)}(x; \sigma_\xi^{(2)})\right| \leq \sum_{i=1}^{m}\left|2\eta^{(1)}T_a\mathbb{E}_{\xi^{(2)}}\sigma(\langle w_i^{(1)}, x\rangle + b_i^{(1)} + \xi_i^{(2)})\right| \tag{442}$$

$$+ \sum_{i=1}^{m}\left|a_{m+i}^{(1)}\right|\left(\left|\langle w_i^{(1)} - w_{m+i}^{(1)}, x\rangle\right| + \left|b_i^{(1)} - b_{m+i}^{(1)}\right|\right) \tag{443}$$

$$\leq \sum_{i=1}^{m}\left|2\eta^{(1)}T_a\right|\left(\left|\langle w_i^{(1)}, x\rangle + b_i^{(1)}\right| + \mathbb{E}_{\xi^{(2)}}\left|\xi_i^{(2)}\right|\right) \tag{444}$$

$$+ \sum_{i=1}^{m}\left|a_{m+i}^{(1)}\right|\left(\left|\langle w_i^{(1)} - w_{m+i}^{(1)}, x\rangle\right| + \left|b_i^{(1)} - b_{m+i}^{(1)}\right|\right). \tag{445}$$

With probability $\geq 1 - \exp(-\Omega(\max\{2p_o(D-k), k\}))$, among all $j \notin A$, we have that at most $2p_o(D-k) + k$ of ϕ_j are $(1-p_o)/\tilde{\sigma}$, while the others are $-p_o/\tilde{\sigma}$. With probability $\geq 1 - (d+D)\exp(-\Omega(k))$ over ζ, for any j, $|\zeta_j| \leq O(\sigma_\zeta\sqrt{\log(k)})$ and $|\langle\zeta, D_\ell\rangle| \leq O(\sigma_\zeta\sqrt{\log(k)})$. For data points with ϕ and ζ satisfying these, we have:

Claim 1. $\left|\langle w_i^{(1)}, x\rangle\right| \leq O(\eta^{(1)}/\gamma)(1 + p_o\tilde{\sigma} + \tilde{\sigma}/\sqrt{k})$.

Proof of Claim 1.

$$\left|\langle w_i^{(1)}, x\rangle\right| = \left|\langle \sum_{\ell=1}^{D} q_{i,\ell}^{(1)} M_\ell + \upsilon, \sum_{j=1}^{D} \phi_j M_j + \zeta/\tilde{\sigma}\rangle\right| \tag{446}$$

$$\leq \left|\langle \sum_{\ell=1}^{D} q_{i,\ell}^{(1)} M_\ell, \sum_{j=1}^{D} \phi_j M_j\rangle\right| + \left|\langle \sum_{\ell=1}^{D} q_{i,\ell}^{(1)} M_\ell, \zeta/\tilde{\sigma}\rangle\right| + \left|\langle \upsilon, \sum_{j=1}^{D} \phi_j M_j\rangle\right| + |\langle \upsilon, \zeta/\tilde{\sigma}\rangle|. \tag{447}$$

For each term above we bound as follows. Note that $\epsilon_{\mathrm{e}} = O(k \log^{1/2}(Dm/\delta)/\sqrt{p_{\mathrm{o}}D})$. Let

$$B_1 := \beta\gamma/\tilde{\sigma} + \epsilon_{\mathrm{e}}/\tilde{\sigma}, \tag{448}$$

$$B_2 := \sigma_\phi^2 \epsilon_{\mathrm{e}} \tilde{\sigma} = O(p_{\mathrm{o}}\epsilon_{\mathrm{e}}/\sqrt{p_{\mathrm{o}}D}), \tag{449}$$

$$C_1 = \frac{k}{\tilde{\sigma}}, \tag{450}$$

$$C_2 := (2p_{\mathrm{o}}(D-k)+k)(1-p_{\mathrm{o}})/\tilde{\sigma} + p_{\mathrm{o}}D/\tilde{\sigma} = O(p_{\mathrm{o}}D/\tilde{\sigma}). \tag{451}$$

Then

$$|a_i^{(0)}|B_1 C_1 = O(\log(Dm/\delta)/k + \log(Dm/\delta)\epsilon_{\mathrm{e}}/(\gamma k)) = O(1/\gamma), \tag{452}$$

$$|a_i^{(0)}|B_2 C_2 = O(p_{\mathrm{o}}\tilde{\sigma}/\gamma), \tag{453}$$

$$|a_i^{(0)}|B_1 C_2 = O(p_{\mathrm{o}}D/k + \sqrt{p_{\mathrm{o}}D}/\gamma), \tag{454}$$

$$|a_i^{(0)}|B_2 C_1 = O(p_{\mathrm{o}}\epsilon_{\mathrm{e}}/(\gamma\sqrt{k})) = O(1/\gamma), \tag{455}$$

Then by the assumption on μ,

$$\left|\langle \sum_{\ell=1}^{D} q_{i,\ell}^{(1)} M_\ell, \sum_{j=1}^{D} \phi_j M_j\rangle\right| \tag{456}$$

$$= \left|\sum_{\ell \in A} \langle q_{i,\ell}^{(1)} M_\ell, M_\ell \phi_\ell\rangle\right| + \left|\sum_{\ell \notin A} \langle q_{i,\ell}^{(1)} M_\ell, M_\ell \phi_\ell\rangle\right| + \left|\sum_{\ell \neq j} \langle q_{i,\ell}^{(1)} M_\ell, M_j \phi_j\rangle\right| \tag{457}$$

$$\leq O(|\eta^{(1)} a_i^{(0)}|) \left(B_1 C_1 + B_2 C_2 + \frac{\mu}{\sqrt{d}}(kB_1(C_1+C_2) + DB_2(C_1+C_2)) \right) \tag{458}$$

$$\leq O(|\eta^{(1)} a_i^{(0)}|) \left(B_1 C_1 + B_2 C_2 + \frac{k\mu}{\sqrt{d}}B_1 C_2 + B_2 C_1 \right) \tag{459}$$

$$\leq O(\eta^{(1)})(1/\gamma + p_{\mathrm{o}}\tilde{\sigma}/\gamma + p_{\mathrm{o}}/\gamma + 1/\gamma) \tag{460}$$

$$\leq O(\eta^{(1)}/\gamma)(1 + p_{\mathrm{o}}\tilde{\sigma}). \tag{461}$$

By the assumption on σ_ζ,

$$\left|\langle \sum_{\ell=1}^{D} q_{i,\ell}^{(1)} M_\ell, \zeta/\tilde{\sigma}\rangle\right| \tag{462}$$

$$\leq O(|\eta^{(1)} a_i^{(0)}|)(kB_1 + DB_2)\frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} \tag{463}$$

$$\leq O(\eta^{(1)})(kB_1 + DB_2)\frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}}\frac{\tilde{\sigma}^2 \sqrt{\log(Dm/\delta)}}{\gamma k^2} \tag{464}$$

$$\leq O(\eta^{(1)}) \left(\sigma_\zeta \log(Dm/\delta) \left(\frac{1}{k} + \frac{\epsilon_{\mathrm{e}}}{\gamma k} \right) + \sigma_\zeta \tilde{\sigma}\frac{\log(k)\log(Dm/\delta)}{\gamma k} \right) \tag{465}$$

$$\leq O(\eta^{(1)}/\gamma). \tag{466}$$

Also note that $|\nu_j| \le O\left(\frac{\sigma_\zeta \log^2(Dm/\delta)}{\tilde{\sigma}\sqrt{p_o D}}\right)$. Then by the assumption on σ_ζ,

$$\left|\langle v, \sum_{j=1}^{D} \phi_j M_j \rangle\right| \tag{467}$$

$$\le O(|\eta^{(1)} a_i^{(0)}|) \times \sqrt{d} \times O\left(\frac{\sigma_\zeta \log^2(Dm/\delta)}{\tilde{\sigma}\sqrt{p_o D}}\right) \times (C_1 + C_2) \tag{468}$$

$$\le O(|\eta^{(1)}/\gamma|). \tag{469}$$

Finally, we have

$$|\langle v, \zeta/\tilde{\sigma} \rangle| \le \sum_{j=1}^{d} |v_j||\zeta_j/\tilde{\sigma}| \tag{470}$$

$$\le O(|\eta^{(1)} a_i^{(0)}|) \times d \times \frac{\sigma_\zeta \log^2(Dm/\delta)}{\tilde{\sigma}\sqrt{p_o D}} \frac{\tilde{\sigma}\sqrt{\log(k)}}{\tilde{\sigma}} \tag{471}$$

$$\le O(\eta^{(1)}/\gamma)\frac{\tilde{\sigma}}{\sqrt{k}}. \tag{472}$$

$$\square$$

We also have:

$$|T_a| = O(\sigma_w \sqrt{\log(Dm/\delta)}) \tag{473}$$

$$\left|b_i^{(1)}\right| \le \left|b_i^{(0)}\right| + \left|\eta^{(1)} a_i^{(0)} T_b\right| \tag{474}$$

$$\le \frac{\sqrt{\log(m/\delta)}}{k^2} + \left|\eta^{(1)} a_i^{(0)} \epsilon_e\right|. \tag{475}$$

$$\mathbb{E}_{\xi^{(2)}} \left|\xi_i^{(2)}\right| \le O(\sigma_\xi^{(2)}). \tag{476}$$

$$|a_{m+i}^{(1)}| \le |a_i^{(0)}| + |\eta^{(1)} T_a| \le |a_i^{(0)}| + O(\eta^{(1)} \sigma_w \sqrt{\log(Dm/\delta)}). \tag{477}$$

$$\left|\langle w_i^{(1)} - w_{m+i}^{(1)}, x \rangle\right| = 2\left|\langle w_i^{(1)}, x \rangle\right| = O(\eta^{(1)} \tilde{\sigma}/\gamma). \tag{478}$$

$$\left|b_i^{(1)} - b_{m+i}^{(1)}\right| = 2|\eta^{(1)} a_i^{(0)} T_b| = O(|\eta^{(1)} a_i^{(0)}|\epsilon_e). \tag{479}$$

Then we have

$$\left|g^{(1)}(x; \sigma_\xi^{(2)})\right| \le O\left(m\eta^{(1)} \sigma_w \sqrt{\log(Dm/\delta)}\right) \left(\frac{\eta^{(1)} \tilde{\sigma}}{\gamma} + \frac{\sqrt{\log(m/\delta)}}{k^2} + \left|\eta^{(1)} a_i^{(0)} \epsilon_e\right| + \sigma_\xi^{(2)}\right) \tag{480}$$

$$+ O\left(m(|a_i^{(0)}| + \eta^{(1)} \sigma_w \sqrt{\log(Dm/\delta)})\right) \left(\frac{\eta^{(1)} \tilde{\sigma}}{\gamma} + \left|\eta^{(1)} a_i^{(0)} \epsilon_e\right|\right) \tag{481}$$

$$= O\left(m\eta^{(1)} \sigma_w \frac{\log(Dm/\delta)}{k} + m|a_i^{(0)}| \left(\frac{\eta^{(1)} \tilde{\sigma}}{\gamma} + \left|\eta^{(1)} a_i^{(0)} \epsilon_e\right|\right)\right) \tag{482}$$

$$= O\left(m\eta^{(1)} \sigma_w \frac{\log(Dm/\delta)}{k} + m|a_i^{(0)}|\frac{\eta^{(1)} \tilde{\sigma}}{\gamma} + m|a_i^{(0)}|\frac{\eta^{(1)} \tilde{\sigma}}{\gamma\sqrt{k}}\right) \tag{483}$$

$$< 1. \tag{484}$$

Then $\left|yg^{(1)}(x; \sigma_\xi^{(2)})\right| < 1$. Finally, the statement on $\left|\langle (q_i^{(1)})_{-A}, \phi_{-A} \rangle\right|$ follows from a similar calculation on $\left|\langle w_i^{(1)}, x \rangle\right| = \left|\langle q_i^{(1)}, \phi \rangle\right|$. $\square$

We are now ready to analyze the gradients in the second gradient step.

Lemma 45. *Fix $\delta \in (0, 1)$ and suppose $w_i^{(0)} \in \mathcal{G}_w(\delta), b_i^{(0)} \in \mathcal{G}_b(\delta), a_i^{(0)} \in \mathcal{G}_a(\delta)$ for all $i \in [2m]$.
Let $\epsilon_{e2} := O\left(\frac{\eta^{(1)}|a_i^{(0)}|k(\gamma+\epsilon_e)}{\tilde{\sigma}^2\sigma_\xi^{(2)}}\right) + \exp(-\Theta(k))$. If $D\mu/\sqrt{d} \leq 1/16$, $\sigma_\zeta\tilde{\sigma} = O(1)$, $\sigma_\zeta^2 d/\tilde{\sigma}^2 = O(1)$,
$p_o = \Omega(k^2/D)$, $k = \Omega(\log^2(Dm/\delta))$, $\sigma_a \leq \tilde{\sigma}^2/(\gamma k^2)$, $\eta^{(1)} = O\left(\frac{\gamma}{km\sigma_a\tilde{\sigma}}\right)$, and $\sigma_\xi^{(2)} = 1/k^{3/2}$,
then*

$$\frac{\partial}{\partial w_i}L_\mathcal{D}(g^{(1)}; \sigma_\xi^{(2)}) = -a_i^{(1)}\left(\sum_{j=1}^D M_j T_j + \nu\right) \tag{485}$$

where T_j satisfies:

- *if $j \in A$, then $|T_j - \beta\gamma/\tilde{\sigma}| \leq O(\epsilon_{e2}/\tilde{\sigma} + \eta^{(1)}/\sigma_\xi^{(2)} + \eta^{(1)}|a_i^{(0)}|\epsilon_e/(\tilde{\sigma}\sigma_\xi^{(2)}))$, where $\beta \in [\Omega(1), 1]$ and depends only on $w_i^{(0)}, b_i^{(0)}$;*

- *if $j \notin A$, then $|T_j| \leq \frac{1}{\tilde{\sigma}}\exp(-\Omega(k)) + O(\sigma_\phi^2\tilde{\sigma}\epsilon_{e2})$;*

- *$|\nu_j| \leq O\left(\frac{\eta^{(1)}\sigma_\zeta}{\gamma\sigma_\xi^{(2)}}\right) + \exp(-\Omega(k))$.*

Proof of Lemma 45. Consider one neuron index i and omit the subscript i in the parameters. By Lemma 44, with probability at least $1 - (d + D)\exp(-\Omega(k)) = 1 - \exp(-\Omega(k))$ over (x, y), $yg^{(1)}(x; \xi^{(2)}) > 1$ and furthermore, for any $i \in [2m]$, $\left|\langle w_i^{(1)}, \zeta/\tilde{\sigma}\rangle\right| = O(\eta^{(1)}\tilde{\sigma}/\gamma)$, $\left|\langle q_i^{(1)}, \phi\rangle\right| = O(\eta^{(1)}\tilde{\sigma}/\gamma)$, and $\left|\langle (q_i^{(1)})_{-A}, \phi_{-A}\rangle\right| = O(\eta^{(1)}\tilde{\sigma}/\gamma)$, and for any $j \in [d], \ell \in [D]$, $|\zeta_j| \leq O(\sigma_\zeta\sqrt{\log(k)})$ and $|\langle \zeta, D_\ell\rangle| \leq O(\sigma_\zeta\sqrt{\log(k)})$. Let $\mathbb{I}_x$ be the indicator of this event.

$$\frac{\partial}{\partial w}L_\mathcal{D}^\alpha(g^{(1)}; \sigma_\xi^{(2)}) \tag{486}$$

$$= -a^{(1)}\mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{\alpha_y y\mathbb{I}_x\mathbb{E}_{\xi^{(2)}}\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0, 1)]x\right\} \tag{487}$$

$$= -a^{(1)}\sum_{j=1}^D M_j \underbrace{\mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{\alpha_y y\mathbb{I}_x\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0, 1)]\phi_j\right\}}_{:=T_j} \tag{488}$$

$$- a^{(1)}\underbrace{\mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{\frac{\alpha_y y\zeta}{\tilde{\sigma}}\mathbb{I}_x\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0, 1)]\right\}}_{:=\nu}. \tag{489}$$

Let $T_{j1} := \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{\alpha_y y\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0, 1)]\phi_j\right\}$. We have

$$|T_j - T_{j1}| \tag{490}$$

$$= \left|\mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{\alpha_y y(1 - \mathbb{I}_x)\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0, 1)]\phi_j\right\}\right| \tag{491}$$

$$\leq \frac{1}{\tilde{\sigma}}\exp(-\Omega(k)). \tag{492}$$

Similarly, let $\nu' := \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{\frac{\alpha_y y\zeta}{\tilde{\sigma}}\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0, 1)]\right\}$. We have

$$|\nu - \nu'| \tag{493}$$

$$= \left|\mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}}\left\{\frac{\alpha_y y\zeta}{\tilde{\sigma}}(1 - \mathbb{I}_x)\mathbb{I}[\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)} \in (0, 1)]\right\}\right| \tag{494}$$

$$\leq \frac{\sigma_\zeta}{\tilde{\sigma}}\exp(-\Omega(k)). \tag{495}$$

So it is sufficient to bound T_{j1} and ν'. For simplicity, we use q as a shorthand for $q_i^{(1)}$.

First, consider $j \in A$.

$$T_{j1} = \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}} \left\{ \alpha_y y \mathbb{I}[\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)} \in (0,1)] \phi_j \right\} \tag{496}$$

$$= \mathbb{E}_{\phi_A} \left\{ \alpha_y y \phi_j \Pr_{\phi_{-A},\xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right] \right\} \tag{497}$$

where $\iota := \langle w^{(1)}, \zeta/\tilde{\sigma} \rangle$. Let

$$I_a := \Pr_{\xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right], \tag{498}$$

$$I_a' := \Pr_{\xi^{(2)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right]. \tag{499}$$

By the property of the Gaussian $\xi^{(2)}$, that $|\langle \phi_A, q_A \rangle| = O(\frac{\eta^{(1)}|a_i^{(0)}|k(\gamma+\epsilon_e)}{\tilde{\sigma}^2})$, and that $|\iota| = |\langle w_i^{(1)}, \zeta/\tilde{\sigma} \rangle|, |\langle \phi, q \rangle|, |\langle \phi_{-A}, q_{-A} \rangle|$ are all $O(\eta^{(1)}\tilde{\sigma}/\gamma) < O(1/k)$, we have

$$|I_a - I_a'| \tag{500}$$

$$\leq \left| \Pr_{\xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \geq 0 \right] - \Pr_{\xi^{(2)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(1)} + \xi^{(2)} \geq 0 \right] \right| \tag{501}$$

$$+ \Pr_{\xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \geq 1 \right] + \Pr_{\xi^{(2)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(1)} + \xi^{(2)} \geq 1 \right] \tag{502}$$

$$= O\left(\frac{\eta^{(1)}|a_i^{(0)}|k(\gamma+\epsilon_e)}{\tilde{\sigma}^2 \sigma_\xi^{(2)}} \right) + \exp(-\Theta(k)) = O(\epsilon_{e2}). \tag{503}$$

This leads to

$$\left| T_{j1} - \mathbb{E}_{\phi_A,\phi_{-A}} \left\{ \alpha_y y \phi_j I_a' \right\} \right| \tag{504}$$

$$\leq \mathbb{E}_{\phi_A} \left\{ |\alpha_y y \phi_j| \left| \mathbb{E}_{\phi_{-A}}(I_a - I_a') \right| \right\} \tag{505}$$

$$\leq O(\epsilon_{e2}) \mathbb{E}_{\phi_A} \left\{ |\alpha_y y \phi_j| \right\} \tag{506}$$

$$\leq O(\epsilon_{e2}/\tilde{\sigma}) \tag{507}$$

where the last step is from Lemma 39. Furthermore,

$$\mathbb{E}_{\phi_A,\phi_{-A}} \left\{ \alpha_y y \phi_j I_a' \right\} \tag{508}$$

$$= \mathbb{E}_{\phi_A} \left\{ \alpha_y y \phi_j \right\} \mathbb{E}_{\phi_{-A}}[I_a'] \tag{509}$$

$$= \mathbb{E}_{\phi_A} \left\{ \alpha_y y \phi_j \right\} \Pr_{\phi_{-A},\xi^{(2)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right]. \tag{510}$$

By Lemma 19, we have $|\langle \phi_{-A}, q_{-A} \rangle + \iota| \leq O(\eta^{(1)}\tilde{\sigma}/\gamma)$. Also, $|b^{(1)} - b^{(0)}| \leq O(\eta^{(1)}|a_i^{(0)}|\epsilon_e)$. By the property of $\xi^{(2)}$,

$$\left| \Pr_{\xi^{(2)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right] - \Pr_{\xi^{(2)}} \left[b^{(0)} + \xi^{(2)} \in (0,1) \right] \right| \tag{511}$$

$$\leq O(\eta^{(1)}\tilde{\sigma}/(\gamma\sigma_\xi^{(2)})) + O(\eta^{(1)}|a_i^{(0)}|\epsilon_e/\sigma_\xi^{(2)}). \tag{512}$$

On the other hand,

$$\beta := \Pr_{\phi_{-A},\xi^{(2)}} \left[b^{(0)} + \xi^{(2)} \in (0,1) \right] = \Pr_{\xi^{(2)}} \left[\xi^{(2)} \in (-b^{(0)}, 1-b^{(0)}) \right] \tag{513}$$

$$= \Omega(1) \tag{514}$$

and β only depends on $b^{(0)}$. By Lemma 39, $\mathbb{E}_{\phi_A} \left\{ \alpha_y y \phi_j \right\} = \gamma/\tilde{\sigma}$. Therefore,

$$|T_{j1} - \beta\gamma/\tilde{\sigma}| \leq O(\epsilon_{e2}/\tilde{\sigma}) + O(\eta^{(1)}/\sigma_\xi^{(2)}) + O(\eta^{(1)}|a_i^{(0)}|\epsilon_e/(\tilde{\sigma}\sigma_\xi^{(2)})). \tag{515}$$

Now, consider $j \notin A$. Let B denote $A \cup \{j\}$.

$$T_{j1} = \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}} \left\{ \alpha_y y \phi_j \mathbb{I} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right] \right\} \tag{516}$$

$$= \mathbb{E}_{\phi_B} \mathbb{E}_{\phi_{-B},\zeta,\xi^{(2)}} \left\{ \alpha_y y \phi_j \mathbb{I} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right] \right\} \tag{517}$$

$$= \mathbb{E}_{\phi_B} \left\{ \alpha_y y \phi_j \Pr_{\phi_{-B},\zeta,\xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right] \right\}. \tag{518}$$

Let

$$I_b := \Pr_{\xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right], \tag{519}$$

$$I_b' := \Pr_{\xi^{(2)}} \left[\langle \phi_{-B}, q_{-B} \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1) \right]. \tag{520}$$

Similar as above, we have $|I_b - I_b'| \le \epsilon_{e2}$. Then by Lemma 39,

$$\left| T_{j1} - \mathbb{E}_{\phi_B,\phi_{-B},\zeta} \left\{ \alpha_y y \phi_j I_b' \right\} \right| \tag{521}$$

$$\le \mathbb{E}_{\phi_B} \left\{ |\alpha_y y \phi_j| |\mathbb{E}_{\phi_{-B},\zeta}(I_b - I_b')| \right\} \tag{522}$$

$$\le O(\epsilon_{e2}) \mathbb{E}_{\phi_A} \left\{ |\alpha_y y| \right\} \mathbb{E}_{\phi_j} \left\{ |\phi_j| \right\} \tag{523}$$

$$\le O(\epsilon_{e2}) \times O(\sigma_\phi^2 \tilde{\sigma}) \tag{524}$$

$$= O(\sigma_\phi^2 \tilde{\sigma} \epsilon_{e2}). \tag{525}$$

Furthermore,

$$\mathbb{E}_{\phi_B,\phi_{-B},\zeta} \left\{ \alpha_y y \phi_j I_b' \right\} = \mathbb{E}_{\phi_A} \left\{ \alpha_y y \right\} \mathbb{E}_{\phi_j} \left\{ \phi_j \right\} \mathbb{E}_{\phi_{-B}} [I_b'] = 0. \tag{526}$$

Therefore,

$$|T_{j1}| \le O(\sigma_\phi^2 \tilde{\sigma} \epsilon_{e2}). \tag{527}$$

Finally, consider ν_j'.

$$\nu_j' = \mathbb{E}_{(x,y)\sim\mathcal{D},\xi^{(2)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} \mathbb{I}[\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)} \in (0,1)] \right\} \tag{528}$$

$$= \mathbb{E}_{\phi_A,\phi_{-A},\zeta,\xi^{(2)}} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} \mathbb{I}[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1)] \right\} \tag{529}$$

$$= \mathbb{E}_{\phi_A,\phi_{-A},\zeta} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} \Pr_{\xi^{(2)}}[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0,1)] \right\} \tag{530}$$

Let

$$I_j := \Pr_{\xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(0)} + \xi^{(1)} \in (0,1) \right], \tag{531}$$

$$I_j' := \Pr_{\xi^{(2)}} \left[\langle \phi, q \rangle + b^{(0)} + \xi^{(1)} \in (0,1) \right]. \tag{532}$$

Since $|\iota| \le O(\eta^{(1)} \tilde{\sigma}/\gamma)$, we have $|I_j - I_j'| \le O(\eta^{(1)} \tilde{\sigma}/(\gamma \sigma_\xi^{(2)}))$. Then

$$\left| \nu_j' - \mathbb{E}_{\phi_A,\phi_{-A},\zeta} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} I_j' \right\} \right| \tag{533}$$

$$= \left| \mathbb{E}_{\phi_A,\phi_{-A},\zeta} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} (I_j - I_j') \right\} \right| \tag{534}$$

$$\le O(\eta^{(1)} \tilde{\sigma}/(\gamma \sigma_\xi^{(2)})) \mathbb{E}_{\phi_A,\phi_{-A},\zeta} \left| \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} \right| \tag{535}$$

$$\le O(\eta^{(1)} \tilde{\sigma}/(\gamma \sigma_\xi^{(2)})) \mathbb{E}_{\phi_A} |\alpha_y y| \mathbb{E}_\zeta \left| \frac{\zeta_j}{\tilde{\sigma}} \right| \tag{536}$$

$$\le O(\eta^{(1)} \tilde{\sigma}/(\gamma \sigma_\xi^{(2)})) \times 1 \times \frac{\sigma_\zeta}{\tilde{\sigma}} \tag{537}$$

$$\le O(\eta^{(1)} \sigma_\zeta/(\gamma \sigma_\xi^{(2)})). \tag{538}$$

Furthermore,

$$\mathbb{E}_{\phi_A, \phi_{-A}, \zeta} \left\{ \frac{\alpha_y y \zeta_j}{\tilde{\sigma}} I'_j \right\} = \mathbb{E}_{\phi_A, \phi_{-A}} \left\{ \frac{\alpha_y y}{\tilde{\sigma}} I'_j \right\} \mathbb{E}_\zeta \left\{ \zeta_j \right\} = 0. \tag{539}$$

Therefore,

$$|\nu_j| \le O\left(\frac{\eta^{(1)} \sigma_\zeta}{\gamma \sigma_\xi^{(2)}} \right) + \exp(-\Omega(k)). \tag{540}$$

$\square$

Lemma 46. *Under the same assumptions as in Lemma 45,*

$$\frac{\partial}{\partial b} L_{\mathcal{D}}(g^{(1)}; \sigma_\xi^{(2)}) = -a_i^{(1)} T_b \tag{541}$$

where $|T_b| \le \exp(-\Omega(k)) + O(\epsilon_{e2})$.

Proof of Lemma 46. Consider one neuron index i and omit the subscript i in the parameters. By Lemma 44, $\Pr[y g^{(1)}(x; \xi^{(2)}) > 1] \le \exp(-\Omega(k))$. Let $\mathbb{I}_x = \mathbb{I}[y g^{(1)}(x; \xi^{(2)}) \le 1]$.

$$\frac{\partial}{\partial b} L_{\mathcal{D}}^\alpha(g^{(1)}; \sigma_\xi^{(2)}) \tag{542}$$

$$= -a^{(1)} \underbrace{\mathbb{E}_{(x,y) \sim \mathcal{D}} \left\{ \alpha_y y \mathbb{I}_x \mathbb{E}_{\xi^{(2)}} \mathbb{I}[\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)} \in (0, 1)] \right\}}_{:= T_b}. \tag{543}$$

Let $T_{b1} := \mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(2)}} \left\{ \alpha_y y \mathbb{I}[\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)} \in (0, 1)] \right\}$. We have

$$|T_b - T_{b1}| \tag{544}$$

$$= \left| \mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(2)}} \left\{ \alpha_y y (1 - \mathbb{I}_x) \mathbb{I}[\langle w^{(1)}, x \rangle + b^{(1)} + \xi^{(2)} \in (0, 1)] \right\} \right| \tag{545}$$

$$\le \exp(-\Omega(k)). \tag{546}$$

So it is sufficient to bound T_{b1}. For simplicity, we use q as a shorthand for $q_i^{(1)}$.

$$T_{b1} = \mathbb{E}_{(x,y) \sim \mathcal{D}, \xi^{(2)}} \left\{ \alpha_y y \mathbb{I} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0, 1) \right] \right\} \tag{547}$$

$$= \mathbb{E}_{\phi_A} \mathbb{E}_{\phi_{-A}, \xi^{(2)}} \left\{ \alpha_y y \mathbb{I} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0, 1) \right] \right\} \tag{548}$$

$$= \mathbb{E}_{\phi_A} \left\{ \alpha_y y \Pr_{\phi_{-A}, \xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0, 1) \right] \right\}. \tag{549}$$

Let

$$I_b := \Pr_{\xi^{(2)}} \left[\langle \phi, q \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0, 1) \right], \tag{550}$$

$$I'_b := \Pr_{\xi^{(2)}} \left[\langle \phi_{-A}, q_{-A} \rangle + \iota + b^{(1)} + \xi^{(2)} \in (0, 1) \right]. \tag{551}$$

Similar as in Lemma 6, we have $|I_b - I'_b| \le \epsilon_{e2}$. Then by Lemma 39,

$$\left| T_{b1} - \mathbb{E}_{\phi_A, \phi_{-A}} \left\{ \alpha_y y I'_b \right\} \right| \tag{552}$$

$$= \left| \mathbb{E}_{\phi_A, \phi_{-A}} \left\{ \alpha_y y (I_b - I'_b) \right\} \right| \tag{553}$$

$$= O(\epsilon_{e2}) \mathbb{E}_{\phi_A, \phi_{-A}} |\alpha_y y| \tag{554}$$

$$\le O(\epsilon_{e2}). \tag{555}$$

Furthermore,

$$\mathbb{E}_{\phi_A, \phi_{-A}} \left\{ \alpha_y y I'_b \right\} = \mathbb{E}_{\phi_A} \left\{ \alpha_y y \right\} \mathbb{E}_{\phi_{-A}} [I'_b] = 0. \tag{556}$$

Therefore, $|T_{b1}| \le O(\epsilon_{e2})$ and the statement follows. $\square$

Lemma 47. *Under the same assumptions as in Lemma 45,*

$$\frac{\partial}{\partial a_i} L_{\mathcal{D}}(g^{(1)}; \sigma_\xi^{(2)}) = -T_a \tag{557}$$

where $|T_a| = O\left(\frac{\eta^{(1)}\tilde{\sigma}}{\gamma p_{\min}}\right) + \exp(-\Omega(k))\text{poly}\left(\frac{dD}{p_{\min}}\right).$

Proof of Lemma 47. Consider one neuron index i and omit the subscript i in the parameters. By Lemma 44, $\Pr[yg^{(1)}(x; \xi^{(2)}) > 1] \leq \exp(-\Theta(k))$. Let $\mathbb{I}_x = \mathbb{I}[yg^{(1)}(x; \xi^{(2)}) \leq 1]$.

$$\frac{\partial}{\partial a} L_{\mathcal{D}}^\alpha(g^{(1)}; \sigma_\xi^{(2)}) \tag{558}$$

$$= -\underbrace{\mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{\alpha_y y \mathbb{I}_x \mathbb{E}_{\xi^{(2)}} \sigma(\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)})\right\}}_{:=T_a}. \tag{559}$$

Let $T_{a1} := \mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{\alpha_y y \mathbb{E}_{\xi^{(2)}} \sigma(\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)})\right\}$. We have

$$|T_a - T_{a1}| \tag{560}$$

$$= \left|\mathbb{E}_{(x,y)\sim\mathcal{D}}\left\{\alpha_y y(1 - \mathbb{I}_x)\mathbb{E}_{\xi^{(2)}} \sigma(\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)})\right\}\right| \tag{561}$$

$$\leq \exp(-\Omega(k)). \tag{562}$$

So it is sufficient to bound T_{a1}. For simplicity, we use q as a shorthand for $q_i^{(1)}$.

Let ϕ_A' be an independent copy of ϕ_A, ϕ' be the vector obtained by replacing in ϕ the entries ϕ_A with ϕ_A', and let $x' = M\phi' + \zeta$ and its label is y'. Then

$$|T_{a1}| := \left|\mathbb{E}_{\phi_A}\left\{\alpha_y y \mathbb{E}_{\phi_{-A}, \zeta, \xi^{(2)}} \sigma(\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)})\right\}\right| \tag{563}$$

$$\leq \frac{1}{2}\left|\mathbb{E}_{\phi_A}\left\{\mathbb{E}_{\phi_{-A}, \zeta, \xi^{(2)}} \sigma(\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(1)})|y = 1\right\}\right. \tag{564}$$

$$\left. - \mathbb{E}_{\phi_A}\left\{\mathbb{E}_{\phi_{-A}, \zeta, \xi^{(2)}} \sigma(\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)})|y = -1\right\}\right| \tag{565}$$

$$\leq \frac{1}{2}\left|\mathbb{E}_{\phi_A}\left\{\mathbb{E}_{\phi_{-A}, \zeta, \xi^{(2)}} \sigma(\langle w^{(1)}, x\rangle + b^{(1)} + \xi^{(2)})|y = 1\right\}\right. \tag{566}$$

$$\left. - \mathbb{E}_{\phi_A'}\left\{\mathbb{E}_{\phi_{-A}, \zeta, \xi^{(2)}} \sigma(\langle w^{(1)}, x'\rangle + b^{(1)} + \xi^{(2)})|y' = -1\right\}\right| \tag{567}$$

$$\leq \frac{1}{2}\mathbb{E}_{\phi_A, \phi_A'}\left\{\mathbb{E}_{\phi_{-A}}\left|\langle w^{(1)}, x\rangle - \langle w^{(1)}, x'\rangle\right| |y = 1, y' = -1\right\} \tag{568}$$

$$\leq \frac{1}{2}\mathbb{E}_{\phi_{-A}}\left(\mathbb{E}_{\phi_A}\left\{\left|\langle w^{(1)}, M\phi\rangle\right| |y = 1\right\} + \mathbb{E}_{\phi_A'}\left\{\left|\langle w^{(1)}, M\phi'\rangle\right| |y' = -1\right\}\right) \tag{569}$$

$$\leq \mathbb{E}_{\phi_{-A}, \phi_A}\left|\alpha_y\langle w^{(1)}, M\phi\rangle\right| \tag{570}$$

$$= \mathbb{E}_\phi\left|\alpha_y\langle w^{(1)}, M\phi\rangle\right| \tag{571}$$

$$= O(\eta^{(1)}\tilde{\sigma}/\gamma) + \exp(-\Omega(k)) \times \frac{\sqrt{dD}}{\tilde{\sigma}} \times \|w^{(1)}\|_\infty \tag{572}$$

$$= O\left(\frac{\eta^{(1)}\tilde{\sigma}}{\gamma p_{\min}}\right) + \exp(-\Omega(k))\text{poly}(dD/p_{\min}) \tag{573}$$

where the fourth step follows from that σ is 1-Lipschitz, and the second to the last line from Lemma 44 and that $\left|\langle w^{(1)}, M\phi\rangle\right| \leq \|w^{(1)}\|_\infty\sqrt{d}\|M\phi\|_2^2$. $\qquad\square$

With the above lemmas about the gradients, we are now ready to show that at the end of the second step, we get a good set of features for accurate prediction.

Lemma 48. *Set*

$$\eta^{(1)} = \frac{\gamma^2 p_{\min}\tilde{\sigma}}{km^3}, \lambda_a^{(1)} = 0, \lambda_w^{(1)} = 1/(2\eta^{(1)}), \sigma_\xi^{(1)} = 1/k^{3/2}, \tag{574}$$

$$\eta^{(2)} = 1, \lambda_a^{(2)} = \lambda_w^{(2)} = 1/(2\eta^{(2)}), \sigma_\xi^{(2)} = 1/k^{3/2}. \tag{575}$$

Fix $\delta \in (0, O(1/k^3))$. If $D\mu/\sqrt{d} \leq 1/16$, $\sigma_\zeta\tilde{\sigma} = O(1)$, $\sigma_\zeta^2 d/\tilde{\sigma}^2 = O(1)$, $p_o = \Omega(k^2/D)$, $k = \Omega\left(\log^2\left(\frac{Dmd}{\delta\gamma p_{\min}}\right)\right)$, and $m \geq \max\{\Omega(k^4), D, d\}$, then with probability at least $1 - \delta$ over the initialization, there exist $\tilde{a}_i$'s such that $\tilde{g}(x) := \sum_{i=1}^{2m} \tilde{a}_i\sigma(\langle w_i^{(2)}, x\rangle + b_i^{(2)})$ satisfies $L_\mathcal{D}(\tilde{g}) \leq \exp(-\Omega(k))$. Furthermore, $\|\tilde{a}\|_0 = O(m/k)$, $\|\tilde{a}\|_\infty = O(k^5/m)$, and $\|\tilde{a}\|_2^2 = O(k^9/m)$. Finally, $\|a^{(2)}\|_\infty = O\left(\frac{1}{km^2}\right)$, $\|w_i^{(2)}\|_2 = O(\tilde{\sigma}/k)$, and $|b_i^{(2)}| = O(1/k^2)$ for all $i \in [2m]$.

Proof of Lemma 48. By Lemma 34, there exists a network $g^*(x) = \sum_{\ell=1}^{3(k+1)} a_\ell^*\sigma(\langle w_\ell^*, x\rangle + b_\ell^*)$ satisfying

$$\Pr_{(x,y)\sim\mathcal{D}}[yg^*(x) \leq 1] \leq \exp(-\Omega(k)).$$

Furthermore, the number of neurons $n = 3(k+1)$, $|a_i^*| \leq 64k, 1/(64k) \leq |b_i^*| \leq 1/4$, $w_i^* = \tilde{\sigma}\sum_{j\in A} M_j/(8k)$, and $|\langle w_i^*, x\rangle + b_i^*| \leq 1$ for any $i \in [n]$ and $(x, y) \sim \mathcal{D}$. Now we fix an ℓ, and show that with high probability there is a neuron in $g^{(2)}$ that can approximate the ℓ-th neuron in g^*.

With probability $\geq 1 - \exp(-\Omega(\max\{2p_o(D-k), k\}))$, among all $j \notin A$, we have that at most $2p_o(D-k) + k$ of ϕ_j are $(1-p_o)/\tilde{\sigma}$, while the others are $-p_o/\tilde{\sigma}$. With probability $\geq 1 - (d + D)\exp(-\Omega(k))$ over ζ, for any j, $|\zeta_j| \leq O(\sigma_\zeta\sqrt{\log(k)})$ and $|\langle\zeta, D_\ell\rangle| \leq O(\sigma_\zeta\sqrt{\log(k)})$. Below we consider data points with ϕ and ζ satisfying these.

By Lemma 36, with probability $1 - 2\delta$ over $w_i^{(0)}$'s, they are all in $\mathcal{G}_w(\delta)$; with probability $1 - \delta$ over $a_i^{(0)}$'s, they are all in $\mathcal{G}_a(\delta)$; with probability $1 - \delta$ over $b_i^{(0)}$'s, they are all in $\mathcal{G}_b(\delta)$. Under these events, by Lemma 43, Lemma 45 and 46, for any neuron $i \in [2m]$, we have

$$w_i^{(2)} = a_i^{(1)}\left(\sum_{j=1}^D M_j T_j + \nu\right), \tag{576}$$

$$b_i^{(2)} = b_i^{(1)} + a_i^{(1)}T_b. \tag{577}$$

where

- if $j \in A$, then $|T_j - \beta\gamma/\tilde{\sigma}| \leq \epsilon_{w1} := O(\epsilon_{e2}/\tilde{\sigma} + \eta^{(1)}/\sigma_\xi^{(2)} + \eta^{(1)}|a_i^{(0)}|\epsilon_e/(\tilde{\sigma}\sigma_\xi^{(2)}))$, where $\beta \in [\Omega(1), 1]$ and depends only on $w_i^{(0)}, b_i^{(0)}$;

- if $j \notin A$, then $|T_j| \leq \epsilon_{w2} := \frac{1}{\tilde{\sigma}}\exp(-\Omega(k)) + O(\sigma_\phi^2\tilde{\sigma}\epsilon_{e2})$;

- $|\nu_j| \leq \epsilon_\nu := O\left(\frac{\eta^{(1)}\sigma_\zeta}{\gamma\sigma_\xi^{(2)}}\right) + \exp(-\Omega(k))$.

- $|T_b| \leq \epsilon_b := \exp(-\Omega(k)) + O(\epsilon_{e2})$.

Given the initialization, with probability $\Omega(1)$ over $b_i^{(0)}$, we have

$$|b_i^{(0)}| \in \left[\frac{1}{2k^2}, \frac{2}{k^2}\right], \text{sign}(b_i^{(0)}) = \text{sign}(b_\ell^*). \tag{578}$$

Finally, since $\frac{8k|b_\ell^*|\beta\gamma}{|b_i^{(0)}|\tilde{\sigma}^2} \in [\Omega(k^2\gamma/\tilde{\sigma}^2), O(k^3\gamma/\tilde{\sigma}^2)]$ and depends only on $w_i^{(0)}, b_i^{(0)}$, we have that for $\epsilon_a = \Theta(1/k^2)$, with probability $\Omega(\epsilon_a) > \delta$ over $a_i^{(0)}$,

$$\left|\frac{8k|b_\ell^*|\beta\gamma}{|b_i^{(0)}|\tilde{\sigma}^2}a_i^{(0)} - 1\right| \leq \epsilon_a, \quad |a_i^{(0)}| = O\left(\frac{\tilde{\sigma}^2}{k^2\gamma}\right). \tag{579}$$

Let $n_a = \epsilon_a m/4$. For the given value of m, by (576)-(579) we have with probability $\geq 1 - 5\delta$ over the initialization, for each ℓ there is a different set of neurons $I_\ell \subseteq [m]$ with $|I_\ell| = n_a$ and such that for each $i_\ell \in I_\ell$,

$$|b_{i_\ell}^{(0)}| \in \left[\frac{1}{2k^2}, \frac{2}{k^2}\right], \quad \text{sign}(b_{i_\ell}^{(0)}) = \text{sign}(b_\ell^*), \tag{580}$$

$$\left|\frac{8k|b_\ell^*|\beta\gamma}{|b_{i_\ell}^{(0)}|\tilde{\sigma}^2}a_{i_\ell}^{(0)} - 1\right| \leq \epsilon_a, \quad |a_{i_\ell}^{(0)}| = O\left(\frac{\tilde{\sigma}^2}{k^2\gamma}\right). \tag{581}$$

Now, construct $\tilde{a}$ such that $\tilde{a}_{i_\ell} = \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|n_a}$ for each ℓ and each $i_\ell \in I_\ell$, and $\tilde{a}_i = 0$ elsewhere. To show that it gives accurate predictions, we first consider bounding some error terms.

For the given values of parameters, we have

$$\epsilon_{e2} = O\left(\frac{\gamma}{m^2}\right), \tag{582}$$

$$\epsilon_{w1} = O\left(\frac{k\gamma}{m^2\tilde{\sigma}} + \frac{\gamma\epsilon_e}{km^2}\right), \tag{583}$$

$$\epsilon_{w2} = O\left(\frac{\gamma}{m^2\tilde{\sigma}}\right), \tag{584}$$

$$\epsilon_\nu = O\left(\frac{\gamma k}{m^3}\right), \tag{585}$$

$$\epsilon_b = O\left(\frac{\gamma}{m^2}\right). \tag{586}$$

We also have the following useful claims.

Claim 2. $\sum_{\ell \in A} |\langle M_\ell, x\rangle| \leq O\left(\frac{k}{\tilde{\sigma}}\right).$

Proof of Claim 2.

$$\sum_{\ell \in A} |\langle M_\ell, x\rangle| \tag{587}$$

$$\leq \sum_{\ell \in A}\left(|\phi_j| + \left|\sum_{j\neq\ell} M_\ell^\top M_j\phi_j\right| + |M_\ell^\top\zeta/\tilde{\sigma}|\right) \tag{588}$$

$$\leq O\left(\frac{k}{\tilde{\sigma}}\right) + O\left(kD\frac{\mu}{\sqrt{d}\tilde{\sigma}}\right) + O\left(k\frac{\sigma_\zeta\sqrt{\log(k)}}{\tilde{\sigma}}\right) \tag{589}$$

$$\leq O\left(\frac{k}{\tilde{\sigma}}\right). \tag{590}$$

$\square$

Claim 3.

$$\left|\langle w_{i_\ell}^{(2)}, x\rangle - \frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}}\sum_{j\in A}\langle M_j, x\rangle\right| \leq O\left(\frac{1}{m}\right). \tag{591}$$

Proof of Claim 3.

$$\left|\langle w_{i_\ell}^{(2)}, x\rangle\right| = \left|\langle\sum_{\ell=1}^D a_{i_\ell}^{(1)}T_\ell M_\ell + v, x\rangle\right| \tag{592}$$

$$\leq \left|\langle\sum_{\ell\in A} a_{i_\ell}^{(1)}T_\ell M_\ell, x\rangle\right| + \left|\langle\sum_{\ell\notin A} a_{i_\ell}^{(1)}T_\ell M_\ell, x\rangle\right| + |\langle v, x\rangle|. \tag{593}$$

Then

$$\left| \langle w_{i_\ell}^{(2)}, x \rangle - \frac{a_{i_\ell}^{(0)} \beta \gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| \tag{594}$$

$$\leq \left| \langle w_{i_\ell}^{(2)}, x \rangle - \frac{a_{i_\ell}^{(1)} \beta \gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| + \left| \frac{a_{i_\ell}^{(1)} \beta \gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle - \frac{a_{i_\ell}^{(0)} \beta \gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| \tag{595}$$

$$\leq \left| \langle w_{i_\ell}^{(2)}, x \rangle - \frac{a_{i_\ell}^{(1)} \beta \gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| + \left| a_{i_\ell}^{(1)} - a_{i_\ell}^{(0)} \right| \left| \frac{\beta \gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right|. \tag{596}$$

The first term is

$$\left| \langle w_{i_\ell}^{(2)}, x \rangle - \frac{a_{i_\ell}^{(1)} \beta \gamma}{\tilde{\sigma}} \sum_{j \in A} \langle M_j, x \rangle \right| \tag{597}$$

$$\leq \left| a_{i_\ell}^{(1)} \right| \left(\left| \langle \sum_{\ell \in A} \left(T_\ell - \frac{\beta \gamma}{\tilde{\sigma}} \right) M_\ell, x \rangle \right| + \left| \langle \sum_{\ell \notin A} T_\ell M_\ell, x \rangle \right| + |\langle \nu, x \rangle| \right). \tag{598}$$

By Claim 2,

$$\left| \langle \sum_{\ell \in A} \left(T_\ell - \frac{\beta \gamma}{\tilde{\sigma}} \right) M_\ell, x \rangle \right| \leq \sum_{\ell \in A} \left| T_\ell - \frac{\beta \gamma}{\tilde{\sigma}} \right| |\langle M_\ell, x \rangle| \tag{599}$$

$$\leq \epsilon_{w1} \sum_{\ell \in A} |\langle M_\ell, x \rangle| \tag{600}$$

$$\leq O\left(\frac{k \epsilon_{w1}}{\tilde{\sigma}} \right). \tag{601}$$

$$\left| \langle \sum_{\ell \notin A} T_\ell M_\ell, x \rangle \right| \leq \left| \sum_{\ell \notin A} T_\ell \phi_j \right| + \left| \sum_{\ell \notin A, j \neq \ell} T_\ell M_\ell^\top M_j \phi_j \right| + \left| \sum_{\ell \notin A} T_\ell M_\ell^\top \zeta / \tilde{\sigma} \right| \tag{602}$$

$$\leq O\left(\frac{D \epsilon_{w2}}{\tilde{\sigma}} \right) + O\left(D^2 \epsilon_{w2} \frac{\mu}{\sqrt{d} \tilde{\sigma}} \right) + O\left(D \epsilon_{w2} \frac{\sigma_\zeta \sqrt{\log(k)}}{\tilde{\sigma}} \right) \tag{603}$$

$$\leq O\left(\frac{D \epsilon_{w2}}{\tilde{\sigma}} \right). \tag{604}$$

$$|\langle \nu, x \rangle| \leq \left| \langle \nu, \sum_{\ell \in [D]} M_\ell \phi_\ell \rangle \right| + |\langle \nu, \zeta / \tilde{\sigma} \rangle| \tag{605}$$

$$\leq \sum_{\ell \in [D]} |\phi_\ell \langle \nu, M_\ell \rangle| + |\langle \nu, \zeta / \tilde{\sigma} \rangle| \tag{606}$$

$$\leq O\left(\frac{p_o D + k}{\tilde{\sigma}} \right) \epsilon_\nu \sqrt{d} + d \epsilon_\nu \frac{O(\sigma_\zeta \sqrt{\log(k)})}{\tilde{\sigma}} \tag{607}$$

$$\leq O\left(\frac{\epsilon_\nu \sqrt{d}}{\tilde{\sigma}} \right) \left(p_o D + \sqrt{d} \epsilon_\nu \sqrt{\log(k)} \right) \tag{608}$$

$$\leq O\left(\frac{\epsilon_\nu \sqrt{d}}{\tilde{\sigma}} \right) \left(p_o D + \tilde{\sigma} \sqrt{\log(k)} \right) \tag{609}$$

$$\leq O\left(\frac{p_o D \epsilon_\nu \sqrt{d}}{\tilde{\sigma}} \right). \tag{610}$$

Then by (582)-(586),

$$\epsilon_\phi := \left(\frac{k\epsilon_{w1}}{\tilde{\sigma}} + \frac{D\epsilon_{w2}}{\tilde{\sigma}} + \frac{p_o D\epsilon_\nu \sqrt{d}}{\tilde{\sigma}} \right) = O\left(\frac{k^2\gamma}{m^2\tilde{\sigma}^2} + \frac{\gamma\epsilon_e}{m^2\tilde{\sigma}} + \frac{\gamma}{m\tilde{\sigma}^2} + \frac{\gamma k}{m^{3/2}\tilde{\sigma}} \right). \tag{611}$$

We have $\left| a_{i_\ell}^{(1)} - a_{i_\ell}^{(0)} \right| = O(\eta^{(1)}\sigma_w\sqrt{\log(Dm/\delta)})$. So the first term is bounded by

$$\left| \langle w_{i_\ell}^{(2)}, x \rangle - \frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}} \sum_{j\in A} \langle M_j, x \rangle \right| \leq \left| a_{i_\ell}^{(1)} \right| \epsilon_\phi \tag{612}$$

$$\leq O\left(\frac{\tilde{\sigma}^2}{k^2\gamma} + \eta^{(1)}\sigma_w\sqrt{\log(Dm/\delta)} \right) \left(\frac{k^2\gamma}{m^2\tilde{\sigma}^2} + \frac{\gamma\epsilon_e}{m^2\tilde{\sigma}} + \frac{\gamma}{m\tilde{\sigma}^2} + \frac{\gamma k}{m^{3/2}\tilde{\sigma}} \right) \leq O\left(\frac{1}{m} \right). \tag{613}$$

By Claim 2, the second term is bounded by

$$\left| a_{i_\ell}^{(1)} - a_{i_\ell}^{(0)} \right| \left| \frac{\beta\gamma}{\tilde{\sigma}} \sum_{j\in A} \langle M_j, x \rangle \right| \leq O\left(\frac{k\eta^{(1)}\sigma_w\sqrt{\log(Dm/\delta)}\gamma}{\tilde{\sigma}^2} \right) \leq O\left(\frac{\gamma^3}{m^3} \right). \tag{614}$$

Combining the bounds on the two terms leads to the claim. $\qquad\square$

Claim 4.

$$|b_{i_\ell}^{(2)} - b_{i_\ell}^{(0)}| \leq O\left(\frac{1}{k^2 m} \right). \tag{615}$$

Proof of Claim 4. By Lemma 43 and 46:

$$|b_{i_\ell}^{(2)} - b_{i_\ell}^{(0)}| \leq |b_{i_\ell}^{(2)} - b_{i_\ell}^{(1)}| + |b_{i_\ell}^{(1)} - b_{i_\ell}^{(0)}| \tag{616}$$

$$\leq O\left(\eta^{(1)}|a_{i_\ell}^{(0)}|\epsilon_e + |a_{i_\ell}^{(1)}|\left(\exp(-\Omega(k)) + \epsilon_{e2}\right) \right) \tag{617}$$

$$\leq O\left(\frac{\gamma}{km^2} + \frac{1}{k^2 m} \right) \leq O\left(\frac{1}{k^2 m} \right). \tag{618}$$

$$\square$$

We are now ready to show $\tilde{g}$ is close to $2g^*$.

$$|\tilde{g}(x) - 2g^*(x)| \tag{619}$$

$$= \left| \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \tilde{a}_{i_\ell}\sigma\left(\langle w_{i_\ell}^{(2)}, x \rangle + b_{i_\ell}^{(2)} \right) - \sum_{\ell=1}^{3(k+1)} 2a_\ell^*\sigma\left(\langle w_\ell^*, x \rangle + b_\ell^* \right) \right| \tag{620}$$

$$= \left| \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|n_a}\sigma\left(\langle w_{i_\ell}^{(2)}, x \rangle + b_{i_\ell}^{(2)} \right) - \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|n_a}\sigma\left(\frac{|b_{i_\ell}^{(0)}|}{|b_\ell^*|}\langle w_\ell^*, x \rangle + b_{i_\ell}^{(0)} \right) \right| \tag{621}$$

$$\leq \left| \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \frac{1}{n_a}\left(\frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}\sigma\left(\langle w_{i_\ell}^{(2)}, x \rangle + b_{i_\ell}^{(2)} \right) - \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}\sigma\left(\frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}}\sum_{j\in A}\langle M_j, x \rangle + b_{i_\ell}^{(0)} \right) \right) \right| \tag{622}$$

$$+ \left| \sum_{\ell=1}^{3(k+1)} \sum_{i_\ell \in I_\ell} \frac{1}{n_a}\left(\frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}\sigma\left(\frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}}\sum_{j\in A}\langle M_j, x \rangle + b_{i_\ell}^{(0)} \right) - \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}\sigma\left(\frac{|b_{i_\ell}^{(0)}|}{|b_\ell^*|}\langle w_\ell^*, x \rangle + b_{i_\ell}^{(0)} \right) \right) \right|. \tag{623}$$

Here the second equation follows from that σ is positive-homogeneous in $[0, 1]$, $|\langle w_\ell^*, x \rangle + b_\ell^*| \leq 1$, $|b_{i_\ell}^{(0)}|/|b_\ell^*| \leq 1$.

By Claim 3 and 4, the first term is bounded by:

$$3(k+1)\max_\ell \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}\left(\left|\langle w_{i_\ell}^{(2)}, x\rangle - \frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}}\sum_{j\in A}\langle M_j, x\rangle\right| + |b_{i_\ell}^{(2)} - b_{i_\ell}^{(0)}|\right) \tag{624}$$

$$\leq 3(k+1)\max_\ell \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}O\left(\frac{1}{m}\right) \tag{625}$$

$$\leq O\left(\frac{k^4}{m}\right). \tag{626}$$

By Claim 2, the second term is bounded by:

$$3(k+1)\max_\ell \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}\left|\frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}}\sum_{j\in A}\langle M_j, x\rangle - \frac{|b_{i_\ell}^{(0)}|}{|b_\ell^*|}\langle w_\ell^*, x\rangle\right| \tag{627}$$

$$\leq 3(k+1)\max_\ell \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}\left|\frac{a_{i_\ell}^{(0)}\beta\gamma}{\tilde{\sigma}}\sum_{j\in A}\langle M_j, x\rangle - \frac{|b_{i_\ell}^{(0)}|\tilde{\sigma}}{8k|b_\ell^*|}\sum_{j\in A}\langle M_j, x\rangle\right| \tag{628}$$

$$\leq 3(k+1)\max_\ell \frac{2a_\ell^*|b_\ell^*|}{|b_{i_\ell}^{(0)}|}\frac{\tilde{\sigma}|b_{i_\ell}^{(0)}|}{8k|b_\ell^*|}\left|\frac{8ka_{i_\ell}\beta\gamma|b_\ell^*|}{\tilde{\sigma}^2|b_{i_\ell}^{(0)}|} - 1\right|\left|\sum_{j\in A}\langle M_j, x\rangle\right| \tag{629}$$

$$\leq 3(k+1)\max_\ell O(a_\ell^*\epsilon_a) \tag{630}$$

$$\leq O\left(k^2\epsilon_a\right). \tag{631}$$

Then

$$|\tilde{g}(x) - 2g^*(x)| = O\left(\frac{k^4}{m} + k^2\epsilon_a\right) \leq 1. \tag{632}$$

This guarantees $y\tilde{g}(x) \geq 1$. Changing the scaling of δ leads to the statement.

Finally, the bounds on $\tilde{a}$ follow from the above calculation. The bound on $\|a^{(2)}\|_2$ follows from Lemma 47, and those on $\|w_i^{(2)}\|_2$ and $\|b_i^{(2)}\|_2$ follow from (576)(577) and the bounds on $a_i^{(1)}$ and $b_i^{(1)}$ in Lemma 43. $\qquad\square$

F.9 CLASSIFIER LEARNING STAGE AND MAIN THEOREM

Once we have a good set of features in Lemma 48, we can follow exactly the same argument as in Section B.6 and B.7 for the simplified setting, and arrive at the main theorem for the general setting:

Theorem 49 (Restatement of Theorem 33). *Set*

$$\eta^{(1)} = \frac{\gamma^2 p_{\min}\tilde{\sigma}}{km^3}, \lambda_a^{(1)} = 0, \lambda_w^{(1)} = 1/(2\eta^{(1)}), \sigma_\xi^{(1)} = 1/k^2, \tag{633}$$

$$\eta^{(2)} = 1, \lambda_a^{(2)} = \lambda_w^{(2)} = 1/(2\eta^{(2)}), \sigma_\xi^{(2)} = 1/k^2, \tag{634}$$

$$\eta^{(t)} = \eta = \frac{k^2}{Tm^{1/3}}, \lambda_a^{(t)} = \lambda_w^{(t)} = \lambda \leq \frac{k^3}{\tilde{\sigma}m^{1/3}}, \sigma_\xi^{(t)} = 0, \text{ for } 2 < t \leq T. \tag{635}$$

For any $\delta \in (0, O(1/k^3))$, if $\mu \leq O(\sqrt{d}/D)$, $\sigma_\zeta \leq O(\min\{1/\tilde{\sigma}, \tilde{\sigma}/\sqrt{d}\})$, $p_o = \Omega(k^2/D)$, $k = \Omega\left(\log^2\left(\frac{Dmd}{\delta\gamma p_{\min}}\right)\right)$, $m \geq \max\{\Omega(k^4), D, d\}$, then we have for any $\mathcal{D} \in \mathcal{F}_{\boxminus}$, with probability at least $1 - \delta$, there exists $t \in [T]$ such that

$$\Pr[\text{sign}(g^{(t)}(x)) \neq y] \leq L_{\mathcal{D}}(g^{(t)}) = O\left(\frac{k^8}{m^{2/3}} + \frac{k^3 T}{m^2} + \frac{k^2 m^{2/3}}{T}\right). \tag{636}$$

Consequently, for any $\epsilon \in (0, 1)$, if $T = m^{4/3}$, and $m \geq \max\{\Omega(k^{12}/\epsilon^{3/2}), D\}$, then

$$\Pr[\text{sign}(g^{(t)}(x)) \neq y] \leq L_{\mathcal{D}}(g^{(t)}) \leq \epsilon. \tag{637}$$