Unlabelled Sample Compression Schemes for Intersection-Closed Classes and Extremal Classes

Abstract

The sample compressibility of concept classes plays an important role in learning theory, as a sufficient condition for PAC learnability, and more recently as an avenue for robust generalisation in adaptive data analysis. Whether compression schemes of size $O(d)$ must necessarily exist for all classes of VC dimension d is unknown, but conjectured to be true by Warmuth. Recently Chalopin, Chepoi, Moran, and Warmuth (2018) gave a beautiful unlabelled sample compression scheme of size VC dimension for all maximum classes: classes that meet the Sauer-Shelah-Perles Lemma with equality. They also offered a counterexample to compression schemes based on a promising approach known as corner peeling. In this paper we simplify and extend their proof technique to deal with so-called extremal classes of VC dimension d which contain maximum classes of VC dimension $d - 1$. A criterion is given which would imply that all extremal classes admit unlabelled compression schemes of size d. We also prove that all intersection-closed classes with VC dimension d admit unlabelled compression schemes of size at most $11d$.

1   Introduction

Littlestone and Warmuth (1986) initiated the study of sample compression schemes as an alternative characterisation of probably approximately correct (PAC) learnability, to finite Vapnik-Chervonenkis (VC) dimension. Informally a sample compression scheme for a concept class comprises a compressor and reconstructor: while the compressor represents samples labelled by the class as a labelled or unlabelled subsample, the reconstructor must be able to recover a concept consistent with the original labelled sample given only this representation. Littlestone and Warmuth (1986) demonstrated a remarkably simple proof of PAC learnability for any concept classes with labelled compression schemes provided compressed representations have bounded size, constant in the original sample size. They conjectured that bounded labelled compression schemes should also be necessary for PAC learnability—completing the characterisation—specifically that all concept classes of VC dimension d have labelled compression schemes of size d. This conjecture, for size $O(d)$, remains open now more than 35 years later, and stands as one of the oldest open problems in learning theory. In addition to proving PAC generalisation bounds (von Luxburg, Bousquet, and Schölkopf, 2004; Langford, 2005), compression has found important connections to generalisation in the adaptive data analysis setting (Cummings, Ligett, Nissim, Roth, and Wu, 2016).

While a characterisation of learnability by compression would be completed by a relaxation of the original Littlestone and Warmuth (1986) conjecture from labelled schemes of size d to size $poly(d)$, for all VC-d classes, Warmuth (2003) later conjectured $O(d)$. In introducing unlabelled compression schemes—where the compressor's representations omit labels, Kuzmin and Warmuth (2007) conjectured $O(d)$ unlabelled compression schemes. Clearly unlabelled schemes of any size imply labelled compression schemes of the same size.

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

Notable progress towards resolving the sample compression conjecture has involved compressing families of concept classes. Shortly after the original conjecture Floyd (1989) proved that all maximum concept classes $\mathcal{C}$ (those meeting the Sauer-Shelah-Perles Lemma with equality) can be label compressed to size VC $(\mathcal{C})$ (Floyd and Warmuth, 1995). Chalopin et al. (2018) recently proved the same bound but with unlabelled compression, while also providing an elegant counter example to the same result for so-called corner peeling schemes. This shows that the constructions of corner peeling schemes by Kuzmin and Warmuth (2007) and Rubinstein and Rubinstein (2012) are incorrect. Moran and Yehudayoff (2016) proved that all VC-d classes can be label compressed to size exponential in d. Recently Pálvölgyi and Tardos (2020) offered examples of VC classes which cannot admit unlabelled compression schemes of size d.

Moran and Warmuth (2016) discover labelled compression schemes for extremal classes, which contain cubes for any coordinates shattered by the class, and generalise maximum classes. It is unknown if all extremal classes admit unlabelled compression schemes of size $O(d)$. In this paper, we give a criterion on extremal classes to admit unlabelled compression schemes of size VC dimension

A promising approach to the general conjecture—and a strong motivation for compressing special families of concept classes–is via embedding. In this sense, there is interest in embedding general VC-classes into larger classes with good compression properties. Rubinstein, Rubinstein, and Bartlett (2015) proved that VC-d classes cannot in general be embedded in maximum classes of VC dimension less than $2d - 1$, leaving $O(d)$-maximum embeddings still open.

Intersection-closed classes—those closed under coordinate-wise multiplication—are another promising focus for compression, as any class can be embedded into an intersection-closed class. In this paper, we show that any intersection-closed class can be embedded into an extremal intersection-closed class with only a linear increase in VC dimension. These extremal intersection-closed classes have VC-sized unlabelled compression schemes by Chalopin et al. (2018). As such, we establish for the first time that intersection-closed classes can be unlabelled compressed to size linear in their VC dimension. We offer a short proof of the result by Chalopin et al. (2018) that maximum classes have VC dimension d-size unlabelled compression schemes and extend this to extremal classes which contain maximum classes of VC dimension $d - 1$.

Finally we give an efficient method to compute the VC dimension of the intersection closure of a class.

2 Preliminaries

Given $n \in \mathbb{N}$ we call a subset of the binary n-cube $\mathcal{C} \subseteq \{0, 1\}^n$, a *concept class* with elements $v \in \mathcal{C}$ *concepts*. These come from restrictions of classifier families, on samples of n instances.[1] We equate each instance in $[n] = \{1, \dots, n\}$ with *coordinates*, *axes* or *colours* of the n-cube. An important combinatorial parameter is the VC dimension, which is used to bound concept class cardinality.

Definition 2.1 (Vapnik and Chervonenkis, 1971). The *VC dimension* of concept class $\mathcal{C} \subseteq \{0, 1\}^n$ is defined as VC $(\mathcal{C}) = \max \{|I| : I \subseteq [n], \mathrm{proj}_I (\mathcal{C}) = \{0, 1\}^{|I|}\}$, where $\mathrm{proj}_I (\mathcal{C}) = \{(v_i)_{i \in I} : v \in \mathcal{C}\}$ is the set of *coordinate projections* of the concepts of $\mathcal{C}$ on coordinates $I \subseteq [n]$.

Lemma 2.2 (Sauer-Shelah-Perles Lemma, Vapnik and Chervonenkis, 1971; Shelah, 1972; Sauer, 1972). *The cardinality of any concept class $\mathcal{C} \subseteq \{0, 1\}^n$ is bounded by $|\mathcal{C}| \leq \sum_{i=0}^{\mathrm{VC}(\mathcal{C})} \binom{n}{i}$.*

Concept classes meeting Sauer's Lemma with equality are called *maximum* (Welzl, 1987). A cubical–complex structure can be put on concept classes $\mathcal{C} \subseteq \{0, 1\}^n$. To begin, each $v \in \mathcal{C}$ is a vertex, and pairs of Hamming-1 separated vertices are edges, of the *one-inclusion graph* of $\mathcal{C}$ (Haussler, Littlestone, and Warmuth, 1994). Each higher-order cube is composed of points varying over a set of coordinates, the cube's *colours*; a cube's common values on non-colour coordinates make up its *anchor*. (An equivalent view is that d-cubes arise as pre-images of points in the binary m-cube under a projection from the binary $m + d$-cube to the binary m-cube.) Any d-maximum class has a special structure as a complete collection of d-cubes:

Definition 2.3 (Rubinstein, Bartlett, and Rubinstein, 2009). A set of d-cubes $\mathcal{S}$ of cardinality $\binom{n}{d}$ is called *d-complete* if for all $I \subseteq [n]$ of cardinality d, there exists $c_I \in \mathcal{S}$ with $\mathrm{proj}_I (c_I) = \{0, 1\}^d$.

---

[1]Consider family of classifiers $\mathcal{F}$ mapping input domain $\mathcal{X}$ to labels $\{0, 1\}$. On sample $\mathbf{x} \in \mathcal{X}^n$, the family is represented by induced concept class $\mathcal{C} = \{(f(x_1), \dots, f(x_n) : f \in \mathcal{F}\}$.

87 Another property of a concept class $\mathcal{C}$ is of being *shortest-path closed*—also known as *isometric*
88 (Chalopin et al., 2018)—where every pair of points in $\mathcal{C}$ has a shortest path contained in $\mathcal{C}$ (Kuzmin
89 and Warmuth, 2007; Rubinstein and Rubinstein, 2012).

90 We will be especially interested in *extremal* classes—also known as *ample* classes (Chalopin et al.,
91 2018). A class $\mathcal{C}$ is extremal if whenever $\mathcal{C}$ shatters some $I \subseteq [n]$, there is a subcube $c \subseteq \mathcal{C}$ which
92 maps bijectively to the $|I|$-cube via $\mathrm{proj}_I(\cdot)$. All maximum classes are extremal and extremal classes
93 are shortest-path closed (Moran and Warmuth, 2016), while the complements of extremal classes
94 are themselves extremal (Bandelt, Chepoi, Dress, and Koolen, 2006; Bollobás and Radcliffe, 1995;
95 Lawrence, 1983). An extremal counter-part to Sauer's Lemma was independently discovered by
96 Pajor (1985), Bollobás and Radcliffe (1995), Dress (1997), and Anstee, Rónyai, and Sali (2002).

97 **Lemma 2.4** (Sandwich Lemma). *For any extremal class $\mathcal{C}$, its cardinality is equal to both the number*
98 *of shatterings of $\mathcal{C}$ and the number of cube types (an equivalence class of cubes which all have the*
99 *same set of colours) in $\mathcal{C}$.*

100 Extremal and maximum classes and the VC dimension, have an elegant recursive structure that is best
101 studied by inductive arguments. One such tool is the projection operator $\mathrm{proj}_I(C)$. The *reduction*
102 $\mathcal{C}^I \subseteq \mathrm{proj}_I(\mathcal{C})$ is the subset of points with non-unique pre-images under the projection, while the
103 *tail* is the subset of $\mathrm{proj}_I(\mathcal{C})$ with unique pre-image under the projection (Welzl, 1987; Kuzmin and
104 Warmuth, 2007).

105 Unlabelled compression schemes are defined by *representation mappings*. A party known as the
106 *compressor* uses a representation mapping to summarise a C-labelled sample of any size with a
107 small unlabelled subset of instances. A second party, the *reconstructor*, can invert this representation
108 thanks to condition 1, to a concept that is consistent with the original labelled sample, thanks to
109 condition 2, (Chalopin et al., 2018).

110 **Definition 2.5** (Kuzmin and Warmuth, 2007). Consider $\mathcal{C} \subseteq \{0,1\}^n$. Mapping r is a *representation*
111 *mapping of size $k < n$* of $\mathcal{C}$ if it satisfies

112   1. r is an injection between $\mathcal{C}$ and subsets of $[n]$ of size at most k; and

113   2. (non-clashing) $\mathrm{proj}_{r(v)\cup r(v')}(v) \neq \mathrm{proj}_{r(v)\cup r(v')}(v')$ for all $v \neq v' \in \mathcal{C}$.

3 The intersection closure operator

115 Examples of intersection-closed classes include axis-aligned hyperrectangles in $\mathbb{R}^m$, monomials,
116 which are unions of orthogonal subcubes, and any union of orthogonal intersection-closed sub-
117 classes (Moran and Warmuth, 2016). Closed-below classes (Rubinstein et al., 2009) are intersection
118 closed and extremal. All extremal classes of VC dimension 1 are intersection closed, so long as the
119 origin is located in the class (Moran and Warmuth, 2016).

120 We will use the notation $v \odot w$ to denote the intersection of the vertices v, w. This is just the
121 operation of bitwise Boolean multiplication $v \odot w = (v_1 w_1, \ldots, v_n w_n)$. We can likewise let
122 $u^1 \odot \ldots \odot u^k = (\prod_{i=1}^k u_1^i, \ldots, \prod_{i=1}^k u_n^i)$ denote the (iterated) intersection of vertices $u^1, \ldots, u^k$.

123 **Definition 3.1.** A class $\mathcal{C}$ in the binary n-cube is called *intersection closed* if $v \odot w \in \mathcal{C}$, whenever
124 $v, w \in \mathcal{C}$. Let $\overline{\mathcal{C}} = \odot(\mathcal{C}) = \{u^1 \odot \ldots \odot u^k\} : k \leq n, \{u^1, \ldots, u^k\} \subseteq \mathcal{C}\}$ denote the *intersection*
125 *closure* of any class $\mathcal{C}$ obtained via intersections of all finite subsets of $\mathcal{C}$'s concepts.

126 **Lemma 3.2.** *For any concept class $\mathcal{C} \subset \{0,1\}^n$, the intersection closure $\overline{\mathcal{C}}$ is the unique smallest*
127 *intersection-closed class containing $\mathcal{C}$, and it is contained in any intersection-closed class containing*
128 *$\mathcal{C}$. Moreover for any $J \subseteq [n]$, we must have $\overline{\mathrm{proj}_J(\mathcal{C})} = \mathrm{proj}_J(\overline{\mathcal{C}})$.*

129 We now give a useful result on computing the VC dimension of the intersection closure of any class.

130 **Definition 3.3.** Suppose $\mathcal{C}$ is a class in the binary n-cube with VC dimension d. We will say that
131 $\mathcal{C}$ satisfies the *k-close cube condition* if there is a vertex $v \in \{0,1\}^n$ and a complete union of
132 $(n-k-1)$-cubes in the complement of $\mathcal{C}$ so that each cube is distance at most one from v.

133 **Theorem 3.4.** *Suppose $\mathcal{C}$ is a class in the binary n-cube. If the VC dimension $\overline{d}$ of the intersection*
134 *closure $\overline{\mathcal{C}}$ of $\mathcal{C}$ is at most k then $\mathcal{C}$ satisfies the k-close cube condition. Conversely if $\mathcal{C}$ satisfies the*
135 *k-close cube condition, then there is a choice of origin so that the intersection closure $\overline{\mathcal{C}}$ of $\mathcal{C}$ has VC*
136 *dimension at most k relative to this origin.*

Proof. Suppose first that $\overline{d} \leq k$. Then any projection p of $\overline{C}$ to a $(k+1)$-cube is not onto. Assume $p(C)$ contained the vertex $w = (1, 1, \ldots, 1)$ and all vertices $w^i = (1, 1, \ldots, 1, 0, 1, \ldots, 1)$ with 0 in the ith position and all other entries 1. Then since $p(\overline{C})$ is intersection closed, the image is the whole $(k+1)$-cube contrary to assumption. The reason is that taking intersections of combinations of the vertices w^i gives all the vertices in the $(k+1)$-cube. So at least one of the vertices $w, w^1, \ldots, w^{k+1}$ must not be in $p(C)$. But then there is an $(n - k - 1)$-cube with anchor of this form in the complement of C by taking the inverse image under p of this vertex. By choosing v as the vertex with all entries 1 in the n-cube, we have shown the existence of a collection of cubes in the complement of C as claimed. For clearly each of our $(n - k - 1)$-cubes has distance at most one from v.

Conversely suppose that C has a collection of cubes in its complement as above. We claim that $\overline{d} \leq k$ for a suitable choice of origin. For suppose that any projection p of $\overline{C}$ to the $(k+1)$-cube is selected. By Lemma 3.2, $p(\overline{C}) = \overline{p(C)}$. But now by assumption, if we choose the origin so that v is the vertex with all entries 1, then there is an $(n - k - 1)$-cube in the complement of $\overline{C}$ which has anchor with entries all 1's except for at most one entry 0 and the anchor projects to a vertex of the $(k+1)$-cube. Hence the projection p of $\overline{C}$ is not onto. $\qquad\square$

We end with a useful property of the intersection closure operator. We will abuse notation by saying a class A is *shortest path closed inside a class B*, if $A \subset B$ and any two vertices of A are connected by a shortest path inside B.

Theorem 3.5. *Suppose C is shortest-path closed. Then the intersection closure $\overline{C}$ is also shortest-path closed.*

Proof. We first observe the easy fact that if λ is a shortest path in C consisting of vertices $v^1, v^2, \ldots, v^k$ and $u \in C$ then $u \odot \lambda := \{u \odot v^i : i \in [k]\}$ is a shortest path in $\overline{C}$ possibly with repetitions. For an edge between successive vertices v^i, v^{i+1} is transformed into $u \odot v^i, u \odot v^{i+1}$ which is either an edge of the same colour or a single vertex.

Our main claim that if C is shortest-path closed then $X = \{v : v = u \odot w, u, w \in C\}$ is shortest path closed inside $\overline{C}$. Once this claim is established, the proof follows easily by repeating this step. For we can apply the claim to the class X to prove that the class $\{v : v = u \odot w, u, w \in \mathcal{X}\}$ consisting of all elements of C together with intersections of up to 4 elements of C is shortest-path closed. inside $\overline{X} = \overline{C}$. Repeating at most $log_2 n$ times, $\overline{C}$ is obtained and is shortest-path closed. (Note that since $v = v \odot v$, elements of C can be written as intersections of pairs of elements of C).

The first step in proving the main claim is to observe that there is a shortest path in X connecting any element $u \in C$ to any element $u \odot v \in X$. To construct this path, start with a shortest path λ between u, v in C and then form $u \odot \lambda$. As above this is a shortest path, possibly with repetitions, between $u, u \odot v$ in X. We conclude that given two vertices $u, v \in C$, there is a shortest path in X from u to v passing through $u \odot v$, by concatenating shortest paths in X from u to $u \odot v$ and $u \odot v$ to v.

To prove the main claim, choose elements $v, v' \in X$. We want to find a shortest path in $\overline{C}$ connecting v, v'. Let $v = u \odot w, v' = u' \odot w'$ where $u, w, u', w' \in C$. By the first step, there are shortest paths λ, λ' in X connecting u, u' and w, w' and passing through $u \odot u', w \odot w'$ respectively.

Next, observe that $v \odot v' = u \odot w \odot u' \odot w' = u \odot u' \odot w \odot w'$. Split λ, λ' into λ_i, λ'_i for $i = 1, 2$ so that λ_1, λ'_1 join u, w and $u \odot u', w \odot w'$ respectively and λ_2, λ'_2 join $u \odot u', w \odot w'$ and u', w' respectively. To establish the main claim, it clearly suffices to construct shortest paths in $\overline{C}$ from v to $v \odot v'$ snd from $v \odot v'$ to v'. By symmetry, we will focus on the first of these.

We will prove a slightly stronger claim. Namely assume λ_1 consists of ordered vertices $u = z^1, z^2, \ldots, z^k = u \odot u'$ and λ'_1 has ordered vertices $w = y^1, y^2, \ldots, y^r = w \odot w'$. We will show that a shortest path in X connecting $v, v \odot v'$ can be chosen to have all vertices of the form $z^i \odot y^j$.

To complete the proof, consider the rectangular grid of intersections $z^i \odot y^j$, for $1 \leq i \leq k$ and $1 \leq j \leq r$ with $k, r \geq 1$. We claim that there is a shortest path between the diagonally opposite entries $v = z^1 \odot y^1, v \odot v' = z^k \odot y^r$ using vertices in this grid.

It suffices to use the boundary of the grid to do this. So for example, consider the first path $z^i \odot y^1$ for $1 \leq i \leq k$, followed by a second path $z^k \odot y^j$ for $1 \leq j \leq r$. By the first observation in this proof,

187 the first path is shortest with possible repetitions, as is the second path. To complete the argument,
188 we need to show that the concatenation of these two paths is also shortest with possible repetitions.

189 Notice that the path $\lambda_1 = z^1, z^2, \ldots, z^k$ runs between ends z^1, z^k where z^k is between the origin
190 of the binary n-cube and z_1. So the same follows for the path with possible repetitions $z^i \odot y^1$ for
191 $1 \leq i \leq k$. The second path $z^k \odot y^j$ for $1 \leq j \leq r$ is of the same form and hence the concatenation
192 of these two paths is also shortest with repetitions and the theorem proof is complete. $\qquad\square$

4 Extremal intersection-closed classes

194 **Theorem 4.1.** *Suppose that $\mathcal{C}$ is shortest-path and intersection closed. Then $\mathcal{C}$ is extremal.*

195 *Proof.* Suppose a projection $p : \{0,1\}^n \to \{0,1\}^k$ shatters $\mathcal{C}$. We will prove there is a k-subcube c
196 of $\mathcal{C}$ which witnesses the shattering.

197 Consider all the vertices of $\mathcal{C}$ which map to the vertex $w = (1, 1, \ldots, 1)$ in the k-cube. Choose a
198 vertex v in this set which has smallest distance to the origin. (Since $\mathcal{C}$ is shortest-path closed and
199 intersection closed, the origin is in $\mathcal{C}$ and distance in $\mathcal{C}$ is the same as distance in the n-cube.)

200 Our claim is that there are vertices $v^i \in \mathcal{C}, i \in [k]$, so that the distance from v to each v^i is one and
201 $p(v^i) = (1, 1, \ldots, 0, \ldots, 1) = w^i$ the vertex in the k-cube with all coordinates 1 except a single 0 at
202 the ith coordinate. Once this claim is established, it follows that since $\mathcal{C}$ is intersection closed, taking
203 the intersections of subcollections of the vertices $\{v^i : i \in [k]\}$ generates the required k-subcube c
204 of $\mathcal{C}$. Note c witnesses the shattering, since projection commutes with intersection and the vertices
205 w, w^i generate c under the operation of taking intersections.

206 To prove the claim, choose any vertex $u \in \mathcal{C}$ so that $p(u) = w^i$. Since $\mathcal{C}$ is intersection closed,
207 $u' = v \odot u \in \mathcal{C}$. Then $p(u') = p(v \odot u)) = p(v) \odot p(u) = w \odot w^i = w^i$. Note that u' is a vertex
208 between v and the origin, *i.e.*, for any coordinate where u' has value 1 then v must also. Since $\mathcal{C}$ is
209 shortest-path closed, a shortest path λ exists in $\mathcal{C}$ from v to u'. Moreover λ has vertices in order with
210 distances decreasing to the origin, since u' is between v and the origin.

211 Let v' be the first vertex on λ after v. Since $p(\lambda)$ is a shortest path in the k-cube from w to w^i, it
212 follows that $p(v')$ is either w or w^i. If the former, this contradicts our choice of v as a closest vertex
213 in $\mathcal{C}$ to the origin which projects to w. If the latter then we have found our vertex v^i with distance 1
214 to v and which projects to w^i. $\qquad\square$

215 *Remark* 4.2. Chalopin et al. (2018) discusses the significance of properties of extremal intersection-
216 closed classes, also known as *conditional antimatroids*.

217 *Remark* 4.3. There is a partial order on the binary n-cube given by $w \leq v$ when w is between v and
218 the origin or equivalently there is a shortest path from v to the origin passing through w. Choose an
219 arbitrary ordering on $[n]$. We will denote by $\lambda(v, w)$ the shortest path between v, w with vertices
220 $v = v_1, v_2, \ldots v_k = w$ where $v_i.v_{i+1}$ differ in the first coordinate x_j in the ordering of indices j
221 amongst all coordinates where v_i, w differ.

222 The main result of this section is the following:

223 **Theorem 4.4.** *Any intersection-closed class $\mathcal{C}$ embeds in a shortest-path closed and intersection-*
224 *closed class $\mathcal{C}^*$ so that $d^* \leq 11d$ where d, d^* are the VC dimensions of $\mathcal{C}, \mathcal{C}^*$ respectively. To*
225 *construct $\mathcal{C}^*$, for every pair of vertices $v, w \in \mathcal{C}$ with $w < v$, add all the vertices in $\lambda(v, w)$ to $\mathcal{C}$.*

226 **Lemma 4.5.** *If $\mathcal{C}$ is intersection closed then $\mathcal{C}^*$ as constructed by Theorem 4.4 is intersection closed.*

227 *Proof.* By definition, for any vertices $u, u' \in \mathcal{C}^*$, there are vertices $v, w, v', w' \in \mathcal{C}$ so that $u \in$
228 $\lambda(v, w), u' \in \lambda(v', w')$. Moreover, $w \leq v, w' \leq v'$.

229 Since $\mathcal{C}$ is intersection closed, both $v \odot v'$ and $w \odot w'$ are in $\mathcal{C}$. We claim that $u \odot u'$ is on the path
230 $\lambda(v \odot v', w \odot w')$—note that as part of this claim, we require that $w \odot w' \leq v \odot v'$.

231 We begin with the latter claim. This is easy—$w \odot w'$ has entry 1 at precisely the coordinates where
232 both w, w' have entry 1. But since $w \leq v, w' \leq v', v, v'$ has entry 1 at all the coordinates where
233 w, w' respectively has entry 1. So for any coordinate where $w \odot w'$ has entry 1, it follows immediately
234 that both v, v' have entry 1 and so does $v \odot v'$.

Next, the set of coordinates where v has entry 1 and u has entry 0 forms an initial segment of the coordinates where v has entries 1, using the ordering of the coordinates, since $u \in \lambda(v, w)$. A similar analysis applies to the set of coordinates of u' which are 0 but the entries for v' are 1.

To complete the proof that $\mathcal{C}^*$ is intersection closed, we show that the set of coordinates where the entries of $v \odot v'$ are 1 and the entries of $u \odot u'$ are 0, forms an initial segment of the ordering of the coordinates where the entries of $v \odot v'$ are 1. It will then follow that $u \odot u' \in \lambda(v \odot v', w \odot w')$.

Now the set of coordinates where $v \odot v'$ has entries 1 are exactly those coordinates where both v, v' have entries 1. Moreover $u \odot u'$ has entry 0 at a coordinate if either u or u' has entry 0. We complete the proof using an argument by contradiction. Suppose that there is a coordinate x_i such that $u \odot u'$ has entry 1 but this occurs before a coordinate x_j where $u \odot u'$ has entry 0 and moreover $v \odot v'$ has entry 1. Then at x_j, either u or u' has entry 0. Without loss of generality assume the former. Since $u \odot u'$ has entry 1 at x_i it follows that u has entry 1 at x_i. Note also that v has entry 1 at both x_i, x_j. But this is a contradiction since the coordinates where v has entry 1 but u has entry 0 should form an initial segment of the set of coordinates where v has entry 1. But we have found $j > i$ where u has entry 1 at x_i and 0 at x_j. This establishes that $\mathcal{C}^*$ is intersection closed. $\square$

Lemma 4.6. *If $\mathcal{C}$ is intersection closed then $\mathcal{C}^*$ as constructed by Theorem 4.4 is shortest-path closed.*

Proof. By Lemma 4.5, it suffices to show that if $v, w \in \mathcal{C}^*$ then there are shortest paths from v, w to $v \odot w$, since putting these together will give a shortest path from v to w. Next, it suffices to show there is a shortest path in $\mathcal{C}^*$ between any pair of vertices v, u in $\mathcal{C}^*$, where $u \leq v$, for we can then apply this to the pairs $v, v \odot w$ and $w, v \odot w$.

We claim that there is a vertex $v_1 \in \mathcal{C}$ so that $u \leq v_1, v \leq v_1$ and there are shortest paths in $\mathcal{C}^*$ joining both v_1, u and v_1, v. Firstly, since $v \in \mathcal{C}^*$, there are vertices $v_1, v_2 \in \mathcal{C}$ so that v is in $\lambda(v_1, v_2)$. Hence $v \leq v_1$ and there is a shortest path in $\mathcal{C}^*$ from v_1 to v, namely a segment of $\lambda(v_1, v_2)$.

Next, since $u \leq v$, then $u \leq v_1$. Moreover u lies on a path $\lambda(u_1, u_2)$ where $u_1, u_2 \in \mathcal{C}$, since $u \in \mathcal{C}^*$. Then $v_1 \odot u_1 \in \mathcal{C}$ and $v_1 \odot u_1 \leq u_1$. Since $u \leq v_1, u \leq u_1$, it follows that $u \leq v_1 \odot u_1$.

Notice that u must lie on $\lambda(v_1 \odot u_1, u_2)$. To verify this, observe all the coordinates where entries of $v_1 \odot u_1$ differ from entries of u are also coordinates where u_1 entries differ from u entries. Consequently, coordinates where entries of u and u_2 differ, are lower in the coordinate ordering than those where entries of $v_1 \odot u_1$ and u differ. So this confirms that $u \in \lambda(v_1 \odot u_1, u_2)$.

Combining $\lambda(v_1, v_1 \odot u_1)$ and $\lambda(v_1 \odot u_1, u)$ gives a shortest path from v_1 to u in $\mathcal{C}^*$. So this completes the proof of the claim about v_1.

To recapitulate, we have constructed shortest paths in $\mathcal{C}$ from v_1 to both u, v in $\mathcal{C}^*$, where $v \leq v_1, u \leq v_1, u \leq v$. To complete the proof that there is a shortest path from v to u in $\mathcal{C}$, we use induction on the Hamming distance from v_1 to v. To begin the induction, suppose this distance is 0. Then clearly we are done, since there is a shortest path in $\mathcal{C}^*$ from v to u.

Assume the distance from v_1 to v is k and the result is true for $v_1*, v*, u*$ as above, where the distance from v_1* to $v*$ is less than k. Let v' be the first vertex in the shortest path from v_1 to v. Our aim is to show that there is a shortest path in $\mathcal{C}^*$ from v' to u. By induction, it then follows there is a shortest path in $\mathcal{C}^*$ from v to u.

The argument is similar to that in Theorem 3.5. Let $v_1 = y^1, y^2, \ldots, y^r = u$ be a shortest path in $\mathcal{C}^*$. We can construct a shortest path $v' = v' \odot y^1, v' \odot y^2, \ldots, v' \odot y^r = u$ where there is a single repetition. Namely, since one of the edges joining y^j, y^{j+1} must have the same colour as the edge between v_1, v', then $v' \odot y^j = v' \odot y^{j+1}$. All other pairs of vertices are distance one apart and since $\mathcal{C}^*$ is intersection closed, this shortest path is in $\mathcal{C}^*$. So this completes the proof. $\square$

Proof. (Theorem 4.4) It remains to prove that the VC dimension of $\mathcal{C}^*$ is at most $11d$.

Choose any projection $p : \{0, 1\}^n \to \{0, 1\}^{11d}$. By Sauer's Lemma, since the VC dimension of $p(\mathcal{C})$ is at most d, the cardinality $|p(\mathcal{C})|$ is at most $F(d; 11d, \frac{1}{2}) \times 2^{11d}$ where $F(k; n, p)$ is the cumulative binomial probability function.

Algorithm 1 constructing $\mathcal{C}^*$ involves finding and adjoining shortest paths connecting suitable ordered pairs of points in $\mathcal{C}$. So $|\mathcal{C}^*| \leq n \times |\mathcal{C}|^2$. Similarly we see that $|p(\mathcal{C}^*)| \leq 11d \times |p(\mathcal{C})|^2$, since the process of adjoining shortest paths is preserved under projection. Hence $|p(\mathcal{C}^*| \leq 11d \times$

$F(d; 11d, \frac{1}{2})^2 \times 2^{22d} \times \frac{1}{2}$. Note that as pairs (v, w) are ordered by requiring that $w \leq v$, we get a factor of $\frac{1}{2}$. We claim that this upper bound is strictly less than 2^{11d}. Then p cannot shatter and the VC dimension of $\mathcal{C}^*$ is at most $11d - 1$.

The key idea is to use Chernoff's inequality to estimate $F(d; 11d, \frac{1}{2})$. This is:

$$F\left(d; 11d, \frac{1}{2}\right) \leq \exp\left((-11d) \times D\left(\frac{1}{11} \middle\| \frac{1}{2}\right)\right) ,$$

where $D(\frac{1}{11} \| \frac{1}{2}) = \frac{1}{11} \log \frac{2}{11} + (1 - \frac{1}{11}) \log 2(1 - \frac{1}{11})$. Substituting this into the previous upper bound and simplifying, we obtain $|p(\mathcal{C}^*)| \leq \frac{11d \times 2^{22d}}{2} \exp(22d \log 11 - 2d \log 2 - 20d \log 20)$. We can further simplify this to $|p(\mathcal{C}^*)| \leq 11d \times 11^{22d} \times 10^{-20d} \times \frac{1}{2}$. To complete the argument, we need to show the right side is strictly less than 2^{11d}. Hence it suffices to show that

$$(11d)^{\frac{1}{d}} \times 11^{22} \times 10^{-20} \times \frac{1}{2^{\frac{1}{d}}} < 2^{11}$$

Now if $d \geq 5$ then $11^{\frac{1}{d}} < 1.7$ and $d^{\frac{1}{d}} < 1.4$. So we can compute the left side is at most $1.7 \times 1.4 \times 121 \times \frac{11}{10}^{20} < 1937$ whereas the right side is 2048. (Here we leave out the factor $2^{-\frac{1}{d}}$.)

If $d = 4$ then $\frac{11}{2}^{\frac{1}{4}} < 1.6$ and $4^{\frac{1}{4}} < 1.42$. So we can compute the left side is as at most $1.6 \times 1.42 \times 121 \times \frac{11}{10}^{20} = 1810$ whereas the right side is 2048. So this completes the case when $d \geq 4$. For the other cases, it is convenient to work directly, as Chernoff's inequality is not sharp.

For $d = 1$, $|p(\mathcal{C})| \leq 12$, as it has VC dimension at most 1 in the 11-cube. Hence $|p(\mathcal{C}^*)| \leq 11 \times 12^2 \times \frac{1}{2} = 792$ which is less than 2048.

Next suppose $d = 2$. $|p(\mathcal{C})| \leq 254$, as it has VC dimension at most 2 in the 22-cube. Hence $|p(\mathcal{C}^*)|$ is bounded above by $22 \times 254^2 \times \frac{1}{2} = 709,676$ which is less than $2048^2 = 4,194,304$.

Finally assume $d = 3$. $|p(\mathcal{C})| \leq 6018$, as it has VC dimension at most 3 in the 33-cube. Hence $|p(\mathcal{C}^*)|$ is bounded above by $33 \times 6018^2 \times \frac{1}{2} = 597,569,346$ which is less than $2048^3 = 8,589,934592$. $\quad\square$

Corollary 4.7. *If $\mathcal{C}$ is an intersection-closed class, then $\mathcal{C}$ has a (corner peeling) unlabelled compression scheme of size at most $11d$.*

Proof. This follows immediately by combining Theorem 4.4 and (Chalopin et al., 2018, Theorem 4.9). $\quad\square$

5 Unlabelled compression schemes for extremal classes

The aim is to give a condition on extremal classes which implies that they have unlabelled compression schemes. We then give some situations where this criterion is satisfied. We conjecture that it is always valid.

5.1 High-level strategy

The idea is inspired by Chalopin et al. (2018). Start with a pair $(\mathcal{C}, \mathcal{D})$ consisting of an extremal class and a proper extremal subclass. The condition is that for every cube in $\mathcal{C}$ which is not maximal, there is a cube with the same colours in $\mathcal{D}$. We call this the *cubical colour condition* of the pair $\mathcal{C}, \mathcal{D}$ and abbreviate it by ccc.

We will construct a bijection ϕ between the set $\mathbb{S}$ of maximal cubes c of $\mathcal{C}$ not in $\mathcal{D}$ and $\mathcal{C} \setminus \mathcal{D}$, so that $v = \phi(c) \in c$. Note the cardinalities of these two sets are the same, by the Sandwich Lemma 2.4 applied to the two extremal classes $\mathcal{C}, \mathcal{D}$. Then $\phi^{-1} : \mathcal{C} \setminus \mathcal{D} \to \mathbb{S}$ gives a representation map for $\mathcal{C} \setminus \mathcal{D}$, which also separates the vertices of $\mathcal{C} \setminus \mathcal{D}$ from the vertices of $\mathcal{D}$. Therefore, any representation of $\mathcal{D}$ can be combined with the representation of $\mathcal{C} \setminus \mathcal{D}$ to give a representation of $\mathcal{C}$.

Suppose we can find a sequence of proper pairs of extremal subclasses $(\mathcal{C}, \mathcal{D}), (\mathcal{D}, \mathcal{E}), \ldots, (\mathcal{X}, \mathcal{Y})$, all satisfying ccc, ending with $\mathrm{VC}(\mathcal{Y}) \leq 2$. Then we can start with a representation map of the VC-2 class $\mathcal{Y}$ (Chalopin et al., 2018) and extend to a representation of $\mathcal{X}$, then extending the representation all the way to $\mathcal{C}$ using this strategy.

5.2 Detailed construction

We construct $\phi : \mathcal{C} \setminus \mathcal{D} \to \mathbb{S}$ and show it is a bijection. To begin with, denote the extremal class which is the complement of $\mathcal{D}$ by $\mathcal{D}^*$. Then every vertex of $\mathcal{C} \setminus \mathcal{D}$ is in at least one maximal cube of $\mathcal{D}^*$.

We say that two cubes c, c^* in $\{0, 1\}^n$ are *complementary* if the dimensions d, d^* of c, c^* satisfy $d + d^* = n$ and $c \cap c^*$ is a single vertex. Recall that $\mathbb{S}$ is the set of maximal cubes c of $\mathcal{C}$ not in $\mathcal{D}$. Denote the set of maximal cubes of D^* which intersect $\mathcal{C}$ by $\mathbb{S}^*$.

Lemma 5.1. *Assume that $(\mathcal{C}, \mathcal{D})$ is a proper pair of extremal classes satisfying ccc. If $c \in \mathbb{S}$, then there is a complementary maximal cube c^* of $\mathcal{D}^*$. Conversely, if c^* is a maximal cube of $\mathbb{S}^*$, then there is a complementary cube $c \in \mathbb{S}$. A maximal cube c^* of $\mathbb{S}^*$, cannot have two different complementary cubes in $\mathbb{S}$.*

Given Lemma 5.1, we will call $v = c \cap c^*$ the *special vertex* of c. We now note properties of v, assuming Lemma 5.1.

Corollary 5.2. *Assume that $(\mathcal{C}, \mathcal{D})$ is a proper pair of extremal classes satisfying ccc. Let $v = c \cap c^*$ where $c \in \mathbb{S}$ and $c^* \in \mathbb{S}^*$ are complementary maximal cubes. Then*

- *The set of vertices $v = c \cap c^*$, for all pairs of complementary maximal cubes $c \in \mathbb{S}$ and $c^* \in \mathbb{S}^*$, is $\mathcal{C} \setminus \mathcal{D}$.*
- *The mappings between c, c^*, v are bijections between $\mathbb{S}, \mathbb{S}^*$ and $\mathcal{C} \setminus \mathcal{D}$.*

Proof. Lemma 5.1 gives a bijection between $\mathbb{S}$ and $\mathbb{S}^*$, by the mapping from cubes c to complementary cubes c^*. Moreover by the Sandwich Lemma applied to extremal classes $\mathcal{C}, \mathcal{D}$, the cardinalities of $\mathcal{C} \setminus \mathcal{D}, \mathbb{S}^*$ are the same. Since the map from maximal cubes in $\mathbb{S}$ to special vertices is an injection by Lemma 5.1, it must be a bijection onto $\mathcal{C} \setminus \mathcal{D}$. $\qquad\square$

Proof of Lemma 5.1. Let $p : \{0, 1\}^n \to \{0, 1\}^d$ be the projection to the colours of $c \in \mathbb{S}$. Recall ccc for the pair $\mathcal{C}, \mathcal{D}$ says that for any cube of $\mathcal{C}$ which is not maximal, there is a cube of $\mathcal{D}$ with the same set of colours. So $p(\mathcal{D}) = \{0, 1\}^d \setminus \{w\}$ for some vertex w. For the only other case is that $p(\mathcal{D}) = \{0, 1\}^d$. But since $\mathcal{D}$ is extremal, this would imply $\mathcal{D}$ contained a cube with the same colours as c. This contradicts the assumption that c is maximal in $\mathcal{C}$, since for an extremal class, maximal cubes are unique for their sets of colours.

Define v, c^* by $v = p^{-1}(w) \cap c, c^* = p^{-1}(w)$ respectively. Clearly $c \cap c^* = v$. It remains to show that c^* is maximal in $\mathcal{D}^*$ and c^* cannot intersect some other cube of $\mathbb{S}$ in a single vertex.

Suppose that c^* was properly contained in a larger cube $\tilde{c}$ in $\mathcal{D}^*$. Then $p(\tilde{c})$ would properly contain $\{w\}$. Moreover $\tilde{c} = p^{-1}p(\tilde{c})$. But this contradicts our observation above that $p(\mathcal{D}) = \{0, 1|^d \setminus \{w\}$. So this completes the argument that c^* is maximal.

Next, assume that c^* is complementary to a second maximal cube $\hat{c} \in \mathbb{S}$. Consider the projection $\hat{p}$ of $\{0, 1\}^n$ to $\{0, 1\}^{\hat{d}}$ given by the $\hat{d}$ colours of $\hat{c}$. As above, $\hat{p}(\mathcal{D})$ consists of $\{0, 1\}^{\hat{d}}$ with a single vertex $\hat{w}$ removed. Moreover, $\hat{p}^{-1}(\hat{w})$ is a maximal cube c' of $\mathcal{D}^*$.

Now as $c^* \cap \hat{c} = u^*$ is a single vertex, $\hat{p}(c^*) = \hat{p}(u^*) = u$. As the cubes $c^*, \hat{c}$ are complementary, $c^* = \hat{p}^{-1}(u)$. But now we see that the colours of the anchor of c^* are both the colours of c and of $\hat{c}$. This gives a contradiction, since two maximal cubes of $\mathcal{C}$ cannot have the same set of colours. This completes the proof of Lemma 5.1. $\qquad\square$

Finally we show that the map from a special vertex v to the colours of the corresponding maximal cube c of $\mathbb{S}$ gives a representation mapping of $\mathcal{C} \setminus \mathcal{D}$ which also separates $\mathcal{C} \setminus \mathcal{D}$ from $\mathcal{D}$.

The second property follows immediately. For recall that if p is the projection given by the d colours of c, then $p(\mathcal{D}) = \{0, 1\}^d \setminus \{p(v)\}$. So the colours of c clearly separate v from all the vertices of $\mathcal{D}$. By this we mean that given any vertex $w \in D$, v, w differ on at least one coordinate from the colours of c.

Next suppose two vertices v, v' have the same values at all the coordinates corresponding to the colours of their corresponding cubes $c, c' \in \mathbb{S}$ of dimensions d, d' respectively. Project $\{0, 1\}^n$ to the union of the colours of c, c' by a mapping p''. The image of p'' is a cube $\{0, 1\}^{d''}$ of dimension $d'' \leq d + d'$. In this cube, $\mathcal{C}, \mathcal{D}$ map to extremal classes $p''(\mathcal{C}) = \mathcal{C}'', p''(\mathcal{D}) = \mathcal{D}''$.

We want to check that the pair of extremal classes $(\mathcal{C}'', \mathcal{D}'')$ satisfy ccc. But this follows easily. Choose any non maximal cube $\tilde{c} \in \mathcal{C}''$. As $\mathcal{C}$ is extremal, there is a cube c_0 in $\mathcal{C}$ which projects one-to-one to $\tilde{c}$ and therefore has the same colours as $\tilde{c}$. Since c_0 is not maximal, there is a cube c_1 in $\mathcal{D}$ with the same colours as c_0. Then $p''(c_1)$ is the required cube in $\mathcal{D}''$ with the same colours as $\tilde{c}$ as required.

Next, c, c' project one-to-one by p'' to cubes with colours spanning all of $\{0,1\}^{d''}$. Finally we claim that $p''(v) \neq p''(v')$. Since $p''(v), p''(v')$ are the special points for $p''(c), p''(c')$ in $\{0,1\}^{d''}$, they cannot coincide, by Lemma 5.1. But then $p''(v), p''(v')$ must have at least one different value of their coordinates at the colours of c, c' and hence the same is true for v, v'. So this gives a representation for $\mathcal{C} \setminus \mathcal{D}$, as we have verified the no-clashing condition.

6   Applications

Theorem 6.1. *Suppose that $\mathcal{C}$ is an extremal class of VC dimension d containing a maximum class $\mathcal{D}$ of VC dimension $d-1$. Then $\mathcal{C}$ has a d-size unlabelled compression scheme.*

Proof. This follows immediately by the construction in Section 5. We only have to check that there is a sequence of pairs of extremal classes starting at $\mathcal{C}, \mathcal{D}$, all satisfying ccc. But this is easy since any maximum class of VC dimension i is a union of a complete collection of maximal cubes of dimension i. Moreover any maximum class of VC dimension i has reductions which are products of maximum classes of VC dimension $i-1$ with $\{0,1\}$. $\qquad\square$

Theorem 6.2. *Suppose that $\mathcal{C}$ is an extremal class of VC dimension 2. Then $\mathcal{C}$ contains a proper extremal class $\mathcal{D}$ so that for every edge of $\mathcal{C}$, there is an edge of $\mathcal{D}$ of the same colour. Hence $\mathcal{C}$ has a compression scheme of size 2.*

Proof. Rubinstein and Rubinstein (2008) develops the idea of splitting classes by their reduction (associated with a particular projection direction): for extremal classes, splitting the reduction induces connected components that are themselves extremal. Choose a reduction $\mathcal{C}^1$ of $\mathcal{C}$ which splits off a component $\mathcal{E}$ with a smallest number of vertices, amongst all choices of reductions and complementary components. Let $\mathcal{D} = \mathcal{C} \setminus \mathcal{E}$. We claim that $\mathcal{D}$ is extremal and contains edges of each colour in $\mathcal{C}$.

The reason is that for any other reduction $\mathcal{C}^2$ of C, if $C_2 \cap \mathcal{E} \neq \emptyset$, then $C^2 \cap \mathcal{C}^1 \neq \emptyset$. For if $C^2 \cap \mathcal{C}^1 = \emptyset$, there is a smaller component which C^2 splits off from $\mathcal{E}$, which is a contradiction. But the condition $C^2 \cap \mathcal{C}^1 \neq \emptyset$ implies that $\mathcal{D}$ has edges of each colour of $\mathcal{C}$. Showing $\mathcal{D}$ is extremal is straightforward. $\qquad\square$

Remark 6.3. Chalopin et al. (2018) showed that VC-2 extremal classes have unlabelled 2-compression schemes. Theorem 6.2 gives an alternate proof using our criterion for extremal classes to have such schemes in Section 5.

7   Conclusion

Chalopin et al. (2018) constructed unlabelled compression schemes for maximum classes and extremal intersection-closed classes, while Moran and Warmuth (2016) developed labelled compression schemes for extremal classes. Here we have shown that intersection-closed classes can be embedded into extremal intersection-closed classes with an increase in VC dimension from d to $11d$ and so have $11d$-size unlabelled compression schemes. We simplify and extend the d-size unlabelled compression schemes for maximum classes due to Chalopin et al. (2018), to apply to extremal classes of VC dimension d which contain maximum classes of VC dimension $d-1$. We also give a sufficient condition for extremal classes to admit such compression schemes.

References

Richard P Anstee, Lajos Rónyai, and Attila Sali. Shattering news. *Graphs and Combinatorics*, 18(1): 59–73, 2002.

H. J. Bandelt, V. Chepoi, A. W. M. Dress, and J. H. Koolen. Combinatorics of lopsided sets. *European Journal of Combinatorics*, 27(5):669–689, 2006.

B. Bollobás and A. J. Radcliffe. Defect Sauer results. *Journal of Combinatorial Theory, Series (A)*, 72(2):189–208, 1995.

Jérémie Chalopin, Victor Chepoi, Shay Moran, and Manfred K. Warmuth. Unlabeled sample compression schemes and corner peelings for ample and maximum classes. *CoRR*, abs/1812.02099, 2018. URL `http://arxiv.org/abs/1812.02099`. updated 5 Jan 2022.

Rachel Cummings, Katrina Ligett, Kobbi Nissim, Aaron Roth, and Zhiwei Steven Wu. Adaptive learning with robust generalization guarantees. In *Conference on Learning Theory*, pages 772–814. PMLR, 2016.

A Dress. Towards a theory of holistic clustering. *DIMACS: Series in Discrete Mathematics and Theoretical Computer Science*, 37:271–289, 1997.

S. Floyd. Space-bounded learning and the Vapnik-Chervonenkis dimension. Technical Report TR-89-061, ICSI, UC Berkeley, 1989.

Sally Floyd and Manfred Warmuth. Sample compression, learnability, and the Vapnik-Chervonenkis dimension. *Machine learning*, 21(3):269–304, 1995.

D. Haussler, N. Littlestone, and M.K. Warmuth. Predicting $\{0, 1\}$ functions on randomly drawn points. *Information and Computation*, 115(2):284–293, 1994.

D. Kuzmin and M.K. Warmuth. Unlabeled compression schemes for maximum classes. *Journal of Machine Learning Research*, 8(Sep):2047–2081, 2007.

J. Langford. Tutorial on practical prediction theory for classification. *Journal of Machine Learning Research*, 6(Mar):273–306, 2005.

James Lawrence. Lopsided sets and orthant-intersection by convex sets. *Pacific Journal of Mathematics*, 104(1):155–173, 1983.

N. Littlestone and M.K. Warmuth. Relating data compression and learnability. Unpublished manuscript `http://www.cse.ucsc.edu/~manfred/pubs/lrnk-olivier.pdf`, 1986.

Shay Moran and Manfred K. Warmuth. Labeled compression schemes for extremal classes. In *International Conference on Algorithmic Learning Theory*, pages 34–49. Springer, 2016.

Shay Moran and Amir Yehudayoff. Sample compression schemes for VC classes. *Journal of the ACM (JACM)*, 63(3):1–10, 2016.

Alain Pajor. *Sous-espaces ℓ_1^n des espaces de Banach*. Travaux en Cours. Hermann, Paris, 1985.

Dömötör Pálvölgyi and Gábor Tardos. Unlabeled compression schemes exceeding the VC-dimension. *Discrete Applied Mathematics*, 276:102–107, 2020.

B. I. P. Rubinstein and J. H. Rubinstein. Geometric & topological representations of maximum classes with applications to sample compression. In *COLT'08*, pages 299–310, 2008.

B. I. P. Rubinstein and J. H. Rubinstein. A geometric approach to sample compression. *Journal of Machine Learning Research*, 13(Apr):1221–1261, 2012.

B. I. P. Rubinstein, P. L. Bartlett, and J. H. Rubinstein. Shifting: one-inclusion mistake bounds and sample compression. *Journal of Computer and System Sciences: Special Issue on Learning Theory 2006*, 75(1):37–59, January 2009.

J. Hyam Rubinstein, Benjamin I. P. Rubinstein, and Peter L. Bartlett. Bounding embeddings of VC classes into maximum classes. In V. Vovk, H. Papadopoulos, and A. Gammerman, editors, *Measures of Complexity: Festschrift of Alexey Chervonenkis*. Springer, Berlin, 2015. in press.

N. Sauer. On the density of families of sets. *Journal of Combinatorial Theory, Series A*, 13:145–147, 1972.

S. Shelah. A combinatorial problem; stability and order for models and theories in infinitary languages. *Pacific Journal of Mathematics*, 41(1):247–261, 1972.

V. N. Vapnik and A. Y. Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability and its Applications*, 16(2):264–280, 1971.

U. von Luxburg, O. Bousquet, and B. Schölkopf. A compression approach to support vector model selection. *Journal of Machine Learning Research*, 5:293–323, 2004.

M. K. Warmuth. Compressing to VC dimension many points. In *COLT'03*, 2003.

E. Welzl. Complete range spaces. Unpublished notes, 1987.