t-Divergence: A New Divergence Measure with Application to Robust Statistics & Clustering

Debolina Paul
Affiliate, Machine Learning
Research Group,
Indian Statistical Institute,
Kolkata, India

Saptarshi Chakraborty
Department of Statistics,
University of California,
Berkeley, USA

Swagatam Das
Electronics & Communication
Sciences Unit,
Indian Statistical Institute,
Kolkata, India

Abstract

This paper introduces the t-divergence, a novel divergence measure associated with the inverse tangent function. We investigate its intriguing consistent and outlier-robust features, particularly its quasi-metric properties and role in establishing weak convergence. Additionally, we showcase the efficacy of this divergence measure family in feature-weighted clustering for high-dimensional data.

1 Proposed *t*-Divergence

Practitioners often select divergence measures for their resilience against outliers. Employing less sensitive loss functions such as ℓ_1, Huber, or Geman-McClure typically guarantees this robustness. Nonetheless, these functions often lack smoothness, presenting challenges for derivative-based optimization methods. Our paper introduces the t-divergence, showcasing its efficacy in robust statistical estimation through comprehensive theoretical and experimental analysis. Formally, let Ω be the sample space and let $\mathcal{F}$ be a σ-algebra defined on it. Suppose $\mu : \mathcal{F} \to [0, \infty)$ be a measure defined on $(\Omega, \mathcal{F})$. Let D_μ be the set of all measures on $(\Omega, \mathcal{F})$, dominated by μ and $\int \frac{dP}{d\mu} \tan^{-1} \frac{dP}{d\mu} d\mu < \infty$, i.e. $D_\mu = \left\{ P : P \ll \mu \text{ and } \int \frac{dP}{d\mu} \tan^{-1} \frac{dP}{d\mu} d\mu < \infty \right\}$. Let, $p = \frac{dP}{d\mu}$ and $q = \frac{dQ}{d\mu}$. The t-divergence, $D : D_\mu \times D_\mu \to [0, \infty)$, between measures $P, Q \in D_\mu$, is defined as

$$D(P, Q) = \int (p(x) - q(x)) \tan^{-1}(p(x) - q(x)) d\mu(x)$$

Some intriguing properties of the proposed t-divergence are as follows:

1. $D(P, Q) \geq 0$. Moreover, $D(P, Q) = 0$ *iff* $p = q$, a.e. $[\mu]$.
2. $D(P, Q) < \infty$, for all $P, Q \in D_\mu$.
3. Let D_μ^p denote the set of all probability measures, dominated by μ. Unlike many other divergence measures, we observe that $0 \leq D(P, Q) < \infty$ for any $P, Q \in D_\mu^p$. This is because $D_\mu^p \subseteq D_\mu$.
4. The t-divergence is symmetric, i.e. $D(P, Q) = D(Q, P)$.
5. $D(P, Q) \leq \pi TV(P, Q)$, where $TV(P, Q)$ is the total variation distance between P and Q.
6. Suppose $\{P_n\}_{n \geq 1}$ be a sequence of probability measures in D_μ^p. Also let P be another probability measure, dominated by μ. Then $\lim_{n \to \infty} D(P_n, P) = 0$ implies $P_n \to P$ in distribution.
7. $D(\cdot, \cdot)$ is a near-metric (Burgin, 2017), with $\rho = 2$.

2 Applications

Application to Robust Statistical Inference Suppose $X_1, \ldots, X_n$ are i.i.d. according to the distribution G. Let $\mathcal{F} = \{F_\theta : \theta \in \Theta \subseteq \mathbb{R}^p\}$ be a family of distributions, indexed by θ. We assume that both $G, F_\theta \ll \mu$, for some dominating measure μ, for all $\theta \in \Theta$. Let $g = \frac{dG}{d\mu}$ and $f_\theta = \frac{dF_\theta}{d\mu}$ be respectively. We define the minimum t-functional as: $T(G) = \arg\min_{\theta \in \Theta} D(G, F_\theta)$. In the context of minimum divergence based estimation, one tries to minimise $D(G, F_\theta)$, w.r.t. θ, in order to obtain a point estimate of θ. Since in practice, the distribution G is unknown, one uses proxies (such as kernel density estimates or empirical cumulative distribution function) for G, based on the observed data $X_1, \ldots, X_n$. Let this estimate be $\hat{G}_n$, which has a density $\hat{g}_n$ w.r.t μ. The estimate for θ, based on the data is given by $\hat{\theta} = T(G_n) = \arg\min_{\theta \in \Theta} D(\hat{G}_n, F_\theta)$. We call this estimator as the minimum t-estimator. It can be shown that the t-estimator exists and is unique under mild regularity conditions (refer to Theorem 2). Additionally, the minimum t-estimator is robust under certain regularity conditions. This is done by deriving its influence function and showing that

its bounded. The influence function, $IF(\cdot)$ of a functional $T(\cdot)$ is defined though the following equation: $IF(y; T, G) = \lim_{\epsilon \downarrow 0} \frac{1}{\epsilon}(T((1 - \epsilon)G + \epsilon\delta_y) - T(G))$, where δ_y denotes the degenerate distribution, putting all its mass at y. Informally, we state the following theorem.

Theorem 1. *Under mild assumptions, the first order influence function* $IF(y; T, G)$ *for the minimum t-functional is given by* $\left(\sum_{x \in \mathcal{X}} \left[\frac{2f_{\theta_0}^2(x)u_{\theta_0}(x)u'_{\theta_0}(x)}{(1+(f_{\theta_0}(x)-g(x))^2)^2} + \rho(f_{\theta_0}(x) - g(x))f_{\theta_0}(x)(u_{\theta_0}^2(x) + u'_{\theta_0}(x)) \right] \right)^{-1} \left(\frac{2f_{\theta_0}(y)u_{\theta_0}(y)}{(1+(f_{\theta_0}(y)-g(y))^2)^2} - \sum_{x \in \mathcal{X}} \frac{g(x)f_{\theta_0}(x)u_{\theta_0}(x)}{(1+(f_{\theta_0}(x)-g(x))^2)^2} \right),$ *where* $\rho(x) = \tan^{-1}(x) + \frac{x}{1+x^2}$ *and* $u_\theta(x) = \frac{\partial}{\partial\theta} \log f_\theta(x).$

The first-order influence function for the minimum t-estimator remains bounded as shown in Theorem 1 when $f_\theta(y)u_\theta(y)$ is bounded across all $\theta \in \Theta$ and for all $y \in \mathcal{X}$. This condition is satisfied by exponential families and numerous commonly used distributions.

To demonstrate the robustness of the t-estimator, we conduct experiments with 100 datapoints which consist of $(1 - \epsilon)\%$ from $Binomial(50, \theta)$ and $\epsilon\%$ from $Uniform(40, \ldots, 50)$, with $\epsilon \in (0, 45)$ and true $\theta = 0.5$. We estimate θ under the binomial model using various methods, including maximum likelihood (MLE), median, minimum squared Hellinger estimate, minimum total variation estimate, and minimum t-estimate. We repeat this experiment 100 times and plot the average estimate for θ in Figure 1. The results highlight the vulnerability of MLE and the median to even small outlier fractions. Conversely, the minimum t-estimate demonstrates outlier robustness, performing comparably to the minimum Hellinger and minimum total variation estimates.

Figure 1: Comparison of different estimates of θ at $\theta = 0.5$ in terms of average point estimate under different levels of contamination.

Application to Clustering We use the t-divergence induced loss as opposed to the squarred error or Minkowski loss in the Weighted k-means algorithm to justtify its performance in practice. Our experimental results show that using the robust t-divergence induced loss improves the performance of Weighted k-means, even in a high-dimensional setting, where the number of features (p) far exceeds the number of observations (n). Given data points $\boldsymbol{x}_1, \ldots, \boldsymbol{x}_n \in \mathbb{R}^p$, the objective function is thus given by,

$$f(\boldsymbol{\Theta}, \boldsymbol{w}) = \sum_{i=1}^n \min_{1 \le j \le k} \sum_{l=1}^p w_l^\beta (x_{il} - \theta_{jl}) \tan^{-1}(x_{il} - \theta_{jl}), \quad \text{subject to } \sum_{l=1}^p w_l = 1 \quad (1)$$

A block coordinate descent algorithm is used to minimize the objective function equation 1 (detailed derivation given in the supplementary document), similar in spirit to Lloyd's k-means (Lloyd, 1982).

Table 1: Average ARI values for different peer algorithms on real data benchmarks ($+$ ($\approx$) denotes statistically significant (equivalent) results w.r.t. the best performing algorithm of that row; The last row indicates the average rank in terms of ARI).

Datasets	k-means	Wk-means	Minkowski	Sparse	Wk-means (Huber)	Wk-means (t)
Wine	$0.364^+(5)$	$0.561^+(4)$	$0.104^+(6)$	$0.806^\approx(2)$	$0.761^\approx(3)$	$\mathbf{0.830}(1)$
WBDC	$0.490^+(3)$	$0.013^+(6)$	$0.106^+(5)$	$0.491^+(2)$	$0.486^+(4)$	$\mathbf{0.730}(1)$
Lymphoma	$0.394^+(6)$	$0.768^+(4)$	$0.618^+(5)$	$0.848^+(2)$	$0.790^+(3)$	$\mathbf{0.947}(1)$
Leukemia	$0.683^+(3)$	$0.213^+(6)$	$0.401^+(5)$	$0.727^+(2)$	$0.581^+(4)$	$\mathbf{0.944}(1)$
Appendicitis	$0.229^+(4.5)$	$0.213^+(6)$	$0.229^+(4.5)$	$0.446^\approx(2)$	$0.251^+(3)$	$\mathbf{0.452}(1)$
Brain	$0.436^+(4)$	$0.432^+(5)$	$0.392^+(6)$	$0.446^+(3)$	$0.451^+(2)$	$\mathbf{0.534}(1)$
Colon	$0.016^+(5)$	$0.001^+(6)$	$0.014^+(4)$	$0.088^+(3)$	$0.102^+(2)$	$\mathbf{0.447}(1)$
Average Rank	4.36	5.43	5.07	2.29	3	$\mathbf{1}$

The study validates the efficacy of a weighted k-means algorithm using t-divergence induced loss on real-life datasets from Arizona State University and UCI Machine Learning Repository. The algorithm's performance was compared to classical k-means, Wk-means (Huang et al., 2005), MW-k-means (De Amorim & Mirkin, 2012), Sparse k-means (Witten & Tibshirani, 2010), and Wk-means with Huber loss using Adjusted Rand Index (ARI) as a performance indicator. Experiments, conducted involved running each algorithm 20 times on the same randomly chosen centroids until convergence. The weighted k-means with t-divergence induced loss showed enhanced performance over peer algorithms, as shown in Table 1, with statistical significance confirmed by Wilcoxon's signed-rank test at a 5% level. This robust approach notably improved the W-k-means algorithm's performance on benchmark datasets.

3 URM STATEMENT

This is to confirm that all the authors of this paper satisfy the URM criteria of the ICLR 2024 tiny papers track.

REFERENCES

Mark Burgin. *Semitopological Vector Spaces: Hypernorms, Hyperseminorms, and Operators*. CRC Press, 2017.

Renato Cordeiro De Amorim and Boris Mirkin. Minkowski metric, feature weighting and anomalous cluster initializing in k-means clustering. *Pattern Recognition*, 45(3):1061–1075, 2012.

Joshua Zhexue Huang, Michael K Ng, Hongqiang Rong, and Zichen Li. Automated variable weighting in k-means type clustering. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 27(5):657–668, 2005.

Stuart Lloyd. Least squares quantization in PCM. *IEEE Transactions on Information Theory*, 28(2):129–137, 1982.

Daniela M Witten and Robert Tibshirani. A framework for feature selection in clustering. *Journal of the American Statistical Association*, 105(490):713–726, 2010.

A PROOF OF PROPERTIES

Proof of Property 1. *Follows trivially as $\forall x, y \in \mathbb{R}$, $(x - y)\tan^{-1}(x - y) \geq 0$ and equality holds iff $x = y$.*

Proof of Property 2. *Let $p = \frac{dP}{d\mu}$ and $q = \frac{dQ}{d\mu}$. Since we know $(x - y)\tan^{-1}(x - y) \leq 2\big[(x - z)\tan^{-1}(x - z) + (z - y)\tan^{-1}(z - y)\big]$, $\forall x, y, z \in \mathbb{R}$, we observe that for all $x \in \Omega$, $0 \leq (p(x) - q(x))\tan^{-1}(p(x) - q(x)) \leq 2[p(x)\tan^{-1}p(x) + q(x)\tan^{-1}q(x)]$. Integrating w.r.t. μ, we get,*

$$D(P, Q) \leq 2\int (p\tan^{-1}p + q\tan^{-1}q)d\mu < \infty.$$

Proof of Property 3. *If $P \in D_\mu^p$. Let $p = \frac{dP}{d\mu}$. Then, $\int p\tan^{-1}pd\mu \leq \frac{\pi}{2}\int pd\mu = \frac{\pi}{2} < \infty$.*

Proof of Property 4. *Follows trivially since x and $\tan^{-1}x$ are both odd functions.*

Proof of Property 5. *We know that for all $z \in \mathbb{R}$, $|\tan^{-1}(z)| \leq \frac{\pi}{2}$. Thus, for all $z \in \mathbb{R}$, $z \leq z\tan^{-1}(z) \leq |\frac{\pi}{2}z|$. Thus, we have, $z\tan^{-1}(z) \leq \frac{\pi}{2}|z|$, for all $z \in \mathbb{R}$. Now, $D(P, Q) = \int (p - q)\tan^{-1}(p - q)d\mu \leq \int \frac{\pi}{2}|p - q|d\mu = \pi\frac{1}{2}\int |p - q|d\mu = \pi TV(P, Q)$.*

Proof of Property 6. *We will first show that $D(P_n, P) \to 0$ implies $TV(P_n, P) \to 0$. We fix $\epsilon > 0$. Thus, there exists $N_\epsilon \in \mathbb{N}$, such that $n \geq N_\epsilon$ implies $D(P_n, P) < \epsilon$. For any $\delta > 0$,*

$$\epsilon > \int (p_n - p)\tan^{-1}(p_n - p)d\mu$$

$$= \int_{|p_n - p| > \delta} (p_n - p)\tan^{-1}(p_n - p)d\mu + \int_{|p_n - p| < \delta} (p_n - p)\tan^{-1}(p_n - p)d\mu$$

$$\geq \tan^{-1}\delta \int_{|p_n - p| > \delta} |p_n - p|d\mu + \int_{|p_n - p| < \delta} (p_n - p)\tan^{-1}(p_n - p)d\mu$$

$$= \tan^{-1}\delta \int |p_n - p|d\mu + \int_{|p_n - p| \leq \delta} [(p_n - p)\tan^{-1}(p_n - p) - \tan^{-1}\delta|p_n - p|]d\mu$$

$$\geq \tan^{-1}\delta \int |p_n - p|d\mu - \int_{|p_n - p| \leq \delta} \tan^{-1}\delta|p_n - p|d\mu$$

$$\geq \tan^{-1}\delta \int |p_n - p|d\mu - \delta\tan^{-1}\delta.$$

Thus, $TV(P_n, P) < \frac{\epsilon}{\tan^{-1}\delta} + \delta$, for all $\delta > 0$. Thus, $TV(P_n, P) \le \inf_{\delta>0}\left(\frac{\epsilon}{\tan^{-1}\delta} + \delta\right)$, which can be made smaller that η, for any prefixed $\eta > 0$, if ϵ is chosen small enough. Thus, $TV(P_n, P) \to 0$ as $n \to \infty$. Now, let $g : \Omega \to \mathbb{R}$ be any bounded continuous function,

$$\left| \int g p_n d\mu - \int g p d\mu \right| \le \int |g||p_n - p|d\mu \le \sup_{x\in\Omega}|g(x)|TV(P_n, P) \to 0.$$

Thus, $P_n \xrightarrow{\mathcal{L}} P$, i.e. P_n converges to P in distribution.

Proof of Property 7. *The non-negativity, identity of indiscernibles and symmetry properties of $D(P,Q)$ have been showed before. What remains to show is that $D(P,Q) \le 2(D(P,Q)+D(R,Q))$ for all $P, Q, R \in D_\mu$. Let $p = \frac{dP}{d\mu}$, $q = \frac{dQ}{d\mu}$ and $r = \frac{dR}{d\mu}$. We note that $D(P,Q) = \int (p - q)\tan^{-1}(p-q)d\mu \le \int 2\left[(p-r)\tan^{-1}(p-r)+(r-q)\tan^{-1}(r-q)\right]d\mu = 2(D(P,Q)+D(R,Q))$.*

B EXISTENCE OF THE t-ESTIMATE

Theorem 2. *Let the parametric family $\mathcal{F}$ be identifiable and let Θ be a compact subset of $\mathbb{R}^p$. We also assume that $f_\theta(\cdot)$ is continuous a.e. $[\mu]$. Then the following holds:*

1. *For all $G \ll \mu$, $T(G)$ exists.*

2. *If $T(G)$ is unique, then $T(\cdot)$ is continuous at G, under the total variation topology, i.e. $T(G_n) \to T(G)$, whenever $\int |g_n - g|d\mu \to 0$. Here $g_n = \frac{dG_n}{d\mu}$.*

3. *$T(F_\theta) = \theta$ for all $\theta \in \Theta$.*

Proof. **Proof of part (1)**: Let $t_n \to t$ be a sequence of parameter values in Θ. Let $h(t) = D(G\|F_t)$. We observe that:

$$|D(G\|F_{t_n}) - D(G\|F_t)| = \left| \int \left[(g - f_{t_n})\tan^{-1}(g - f_{t_n}) - (g - f_t)\tan^{-1}(g - f_t)\right]d\mu \right|$$

$$\le \int \left|(g - f_{t_n})\tan^{-1}(g - f_{t_n}) - (g - f_t)\tan^{-1}(g - f_t)\right|d\mu. \quad (2)$$

We note that,

$$\left|(g - f_{t_n})\tan^{-1}(g - f_{t_n}) - (g - f_t)\tan^{-1}(g - f_t)\right| \le |(g - f_{t_n})\tan^{-1}(g - f_{t_n})| + |(g - f_t)\tan^{-1}(g - f_t)|$$

$$\le \frac{\pi}{2}\left[|(g - f_{t_n})| + |(g - f_t)|\right]$$

$$\le \frac{\pi}{2}\left[2g + f_{t_n} + f_t\right]. \quad (3)$$

We note that the LHS of equation equation 3 is μ-integrable and thus by simple application of Dominated Convergence Theorem (DCT) the RHS of equation 2 converges to 0 as $n \to \infty$. Hence $h(t)$ is continuous on Θ, which is compact. Hence $h(\cdot)$ attains its minimum on Θ.

Proof of part (2): Let $\{G_n\}_{n\ge 1}$ converges to G in total variation sense, i.e. $\int |g_n(x) - g(x)|d\mu(x) \to 0$, as $n \to \infty$. We define $h_n(t) = D(G_n\|F_t)$. We also assume that $\theta_n = T(G_n)$ and $\theta = T(G)$ are also defined uniquely. We observe the following,

$$|h_n(t) - h(t)| = \left| \int \left[(g_n - f_t)\tan^{-1}(g_n - f_t) - (g - f_t)\tan^{-1}(g - f_t)\right]d\mu \right| \quad (4)$$

$$= \int \left| \tan^{-1}(\xi_x) + \frac{\xi_x}{1 + \xi_x^2} \right| |g_n - g|d\mu$$

$$\le \left(\frac{\pi}{2} + 1\right) \int |g_n - g|d\mu$$

Equation equation 4 follows from applying first order Taylor's expansion on the function $x\tan^{-1}(x)$. Here ξ_x lies between $g_n(x) - f_t(x)$ and $(g(x) - f_t(x))$. From the above calculations we conclude that $\lim_{n\to\infty}\sup_{t\in\Theta}|h_n(t) - h(t)| = 0$. From the definition of θ_n and θ, we

observe that:

$$
\begin{aligned}
|h(\theta_n) - h(\theta)| &= h(\theta_n) - h(\theta) \\
&= (h(\theta_n) - h_n(\theta_n)) + (h_n(\theta_n) - h_n(\theta)) + (h_n(\theta) - h(\theta)) \\
&\leq (h(\theta_n) - h_n(\theta_n)) + (h_n(\theta) - h(\theta)) \\
&\leq 2 \sup_{t \in \Theta} |h_n(t) - h(t)| \to 0
\end{aligned}
$$
(5)

(6)

Equation equation 5 follows from the fact that, since θ_n is the minimiser if $h_n(\cdot)$, $h_n(\theta_n) \leq h_n(\theta)$. Thus, we get, $\lim_{n \to \infty} h(\theta_n) = h(\theta)$. We will now show that $\theta_n \to \theta$. We assume the contrary. Suppose $\theta_n \not\to \theta$. We note that $\{\theta_n\}_{n \geq 1}$ is a sequence in the compact set Θ. Thus it has a converging sub-sequence, say, $\{\theta_{n_l}\}_{l \geq 1}$, such that $\theta_{n_l} \to \theta_1$, where, $\theta_1 \neq \theta$. By the continuity of $h(\cdot)$, $h(\theta_{n_l}) \to h(\theta_1)$. This implies that $h(\theta) = h(\theta_1)$, since there cannot be two limit for the converging sequence $h(\theta_{n_l})$. Thus, $h(\theta) = h(\theta_1)$ gives us a contradiction, since, $T(G)$ is assumed to be unique. Thus, $\theta_n \to \theta$.

Proof of part (3): Since the parametric family $\mathcal{F}$ is identifiable, $D(F_\theta \| F_t) = 0$, only at the value $t = \theta$. Thus $T(F_\theta) = \theta$, uniquely. □

C PROOF OF THEOREM 1

We first derive the estimating equation for a minimum t-estimator. Let $\theta_0 = T(G)$ and θ_0 is an interior point of Θ, then θ_0 satisfies the following equation

$$
\left[\frac{\partial}{\partial \theta} \int (f_\theta(x) - g(x)) \tan^{-1}(f_\theta(x) - g(x)) d\mu(x) \right] \Bigg|_{\theta = \theta_0} = 0.
$$

Assuming the differentiability under the integral sign, we get,

$$
\left[\int \left[\tan^{-1}(f_\theta(x) - g(x)) + \frac{f_\theta(x) - g(x)}{1 + (f_\theta(x) - g(x))^2} \right] \frac{\partial f_\theta(x)}{\partial \theta} d\mu(x) \right] \Bigg|_{\theta = \theta_0} = 0.
$$

Thus, θ_0 satisfies the following equation.

$$
\int \rho(f_\theta(x) - g(x)) f_\theta(x) u_\theta(x) d\mu(x) = 0.
$$
(7)

Here $\rho(x) = \tan^{-1}(x) + \frac{x}{1+x^2}$ and $u_\theta(x) = \frac{\partial}{\partial \theta} \log f_\theta(x)$. Equation equation 7 gives the estimating equation for minimum t-estimator.

We will make the following technical assumptions.

A1 The support of F_θ is independent of θ.

A2 There exists $\eta > 0$ such that $T(G_\epsilon)$ is an interior point of Θ, for all $\epsilon < 0$ and for all $y \in \mathbb{R}$. Here $G_\epsilon = (1 - \epsilon)G + \epsilon \delta_y$.

A3 $T(\cdot)$ is Gateaux differentiable at G.

A4 The derivative on the Left hand side of equation 7 is permitted under the integral sign.

Let $G_\epsilon = (1 - \epsilon)G + \epsilon \Delta_y$. Here Δ_y denotes the degenerate distribution at y. We take μ to be the counting measure. Let $\theta_0 = T(G)$ and $\theta_\epsilon = T(G_\epsilon)$ be defined uniquely for all $\epsilon \geq 0$. The value of the influence function at y is given by $IF(y) = \frac{\partial \theta_\epsilon}{\partial \epsilon}\big|_{\epsilon=0}$. Observe that $g_\epsilon = (1 - \epsilon)g + \epsilon \delta_y$ is the density of G_ϵ w.r.t. μ. Here $\delta_y(x) = 1$ if $x = y$ and is 0, otherwise. Before we proceed, we observe that $\rho'(x) = \frac{2}{(1+x^2)^2}$. From the estimating equation equation 7, we observe that

$$
\sum_{x \in \mathcal{X}} \rho(f_{\theta_\epsilon}(x) - g_\epsilon(x)) f_{\theta_\epsilon}(x) u_{\theta_\epsilon}(x) = 0 \implies \sum_{x \in \mathcal{X}} \rho(f_{\theta_\epsilon}(x) - (1-\epsilon)g(x) - \epsilon \delta_y(x)) f_{\theta_\epsilon}(x) u_{\theta_\epsilon}(x) = 0
$$

Differentiating both sides w.r.t. ϵ and assuming that the derivative can be passed inside the summation, we get,

$$\sum_{x \in \mathcal{X}} \left[\frac{2}{(1 + (f_{\theta_\epsilon}(x) - g_\epsilon(x))^2)^2} (f'_{\theta_\epsilon}(x)\theta_\epsilon' + g(x) - \delta_y(x))f_{\theta_\epsilon}(x)u_{\theta_\epsilon}(x) \right.$$
$$\left. + \rho(f_{\theta_\epsilon}(x) - g_\epsilon(x))f'_{\theta_\epsilon}(x)\theta_\epsilon' u_{\theta_\epsilon}(x) + \rho(f_{\theta_\epsilon}(x) - g_\epsilon(x))f_{\theta_\epsilon}(x)u'_{\theta_\epsilon}(x)\theta_\epsilon' \right] = 0.$$

Substituting $\epsilon = 0$ in the above equation, we get,

$$\sum_{x \in \mathcal{X}} \left[\frac{2}{(1 + (f_{\theta_0}(x) - g(x))^2)^2} (f'_{\theta_0}(x)IF(y) + g(x) - \delta_y(x))f_{\theta_0}(x)u_{\theta_0}(x) \right.$$
$$\left. + \rho(f_{\theta_0}(x) - g(x))f'_{\theta_0}(x)IF(y)u_{\theta_0}(x) + \rho(f_{\theta_0}(x) - g(x))f_{\theta_0}(x)u'_{\theta_0}(x)IF(y) \right] = 0.$$

Thus,

$$IF(y) = \sum_{x \in \mathcal{X}} \left[\frac{2f'_{\theta_0}(x)f_{\theta_0}(x)u'_{\theta_0}(x)}{(1 + (f_{\theta_0}(x) - g(x))^2)^2} + \rho(f_{\theta_0}(x) - g(x))f'_{\theta_0}(x)u_{\theta_0}(x) + \rho(f_{\theta_0}(x) - g(x))f_{\theta_0}(x)u'_{\theta_0}(x) \right]$$
$$= \sum_{x \in \mathcal{X}} \frac{2(\delta_y(x) - g(x))f_{\theta_0}(x)u_{\theta_0}(x)}{(1 + (f_{\theta_0}(x) - g(x))^2)^2} = \frac{2f_{\theta_0}(y)u_{\theta_0}(y)}{(1 + (f_{\theta_0}(y) - g(y))^2)^2} - \sum_{x \in \mathcal{X}} \frac{g(x)f_{\theta_0}(x)u_{\theta_0}(x)}{(1 + (f_{\theta_0}(x) - g(x))^2)^2}.$$

simplifying the above equation, we get,

$$IF(y) = \frac{\frac{2f_{\theta_0}(y)u_{\theta_0}(y)}{(1+(f_{\theta_0}(y)-g(y))^2)^2} - \sum_{x \in \mathcal{X}} \frac{g(x)f_{\theta_0}(x)u_{\theta_0}(x)}{(1+(f_{\theta_0}(x)-g(x))^2)^2}}{\sum_{x \in \mathcal{X}} \frac{2f_{\theta_0}^2(x)u_{\theta_0}(x)u'_{\theta_0}(x)}{(1+(f_{\theta_0}(x)-g(x))^2)^2} + \rho(f_{\theta_0}(x) - g(x))f_{\theta_0}(x)(u_{\theta_0}^2(x) + u'_{\theta_0}(x))}.$$

D ALGORITHM FOR APPLICATION TO CLUSTERING

Algorithm 1 Weighted k-means with t-divergence induced loss

Input: $X \in \mathbb{R}^{n \times p}$, $\beta > 1$. **Output**: Cluster assignment matrix U, feature weight vector w.
repeat

    Step 1: Update U by $u_{ij}^{(t+1)} \leftarrow \begin{cases} 1 & \text{if } j = \arg\min_{1 \le j' \le k} \sum_{l=1}^{p} w_l^{(t)\beta}(x_{il} - \theta_{j'l}^{(t)}) \tan^{-1}(x_{il} - \theta_{j'l}^{(t)}), \\ 0 & \text{Otherwise.} \end{cases}$

    Step 2: Update Θ by taking $\theta_{jl}^{(t+1)} \leftarrow \arg\min_{\theta \in \mathbb{R}} \sum_{i=1}^{n} u_{ij}^{(t+1)}(x_{il} - \theta) \tan^{-1}(x_{il} - \theta)$ by the Newton-Raphson method.

    Setp 3: Update w by taking $w_l^{(t+1)} \leftarrow \frac{1/D_l^{(\beta-1)}}{\sum_{m=1}^{p} 1/D_m^{(\beta-1)}}$, where $D_l = \sum_{i=1}^{n} u_{ij}^{(t+1)}(x_{il} - \theta_{jl}^{(t+1)}) \tan^{-1}(x_{il} - \theta_{jl}^{(t+1)})$.
until objective equation 1 converges.