CONSIDERATIONS FOR DISTRIBUTION SHIFT ROBUSTNESS IN HEALTH

ABSTRACT

When analyzing robustness of predictive models under distribution shift, many works focus on tackling generalization in the presence of spurious correlations. In this case, one typically makes use of covariates or environment indicators to enforce independencies in learned models to guarantee generalization under various distribution shifts. In this work, we analyze a class of distribution shifts, where such independencies are not desirable, as there is a causal association between covariates and outcomes of interest. This case is common in the health space where covariates can be causally, as opposed to spuriously, related to outcomes of interest. We formalize this setting and relate it to common distribution shift settings from the literature. We theoretically show why standard supervised learning and invariant learning will not yield robust predictors in this case, while including the causal covariates into the prediction model can recover robustness. We demonstrate our theoretical findings in practice in experiments on synthetic data.

1 INTRODUCTION

In this work, we motivate how common assumptions in the domain generalization and invariant learning literature (Arjovsky et al., 2019; Ganin et al., 2016; Veitch et al., 2021) are violated in a set of broadly applicable problems, e.g. in healthcare. Invariant learning typically assumes a spurious or confounded association between outcome and covariates or auxiliary information. Hence, building a predictor that is invariant to the covariates or associated environment indicators can be shown to generalize better under distribution shift than standard empirical risk minimization. However, in many applications of interest for machine learning, e.g. in healthcare, there might not be only spurious associations between covariates and outcome, but also causal ones.

One illustrative example is body mass index (BMI), which is causally related to a host of conditions, e.g., left ventricular hypertrophy (LVH) (Lorell & Carabello, 2000). BMI is not "spurious" in the sense that it is merely associated with LVH, but can directly cause changes in left ventricular mass, which in turn can lead to LVH (Himeno et al., 1996). However, a shift in the prevalence of elevated BMI can shift the association between a signal — e.g., an electrocardiogram (ECG) — that is influenced by both BMI and LVH.

We formalize such a causal setting and show that it leads to regression in performance of machine learning models under distribution shift, that can not be mitigated with common invariant learning methods. Our contributions are the following:

- We motivate and formalize a class of problems where covariates, such as demographics or other auxiliary data causally influence the outcome of interest and explain the difference to the commonly considered confounded or spurious associations.

- For this class of problems we show theoretically and on simulated data, how distribution shifts along such causally influencing covariates cause discrepancies in performance that can not be mitigated with invariant learning methods designed for the commonly considered confounded setting.

(a) "Confounded graph" (b) "Direct causal graph" (this work).

Figure 1: Causal graphs considered in this work. (a) The "confounded graph" describes a spurious/confounded association between Y and V, and has been considered in the ML literature (Heinze-Deml & Meinshausen, 2021; Veitch et al., 2021; Makar et al., 2022; Puli et al., 2022). This setting requires that the marginal $P(Y)$ remains invariant across distribution shifts. (b) In the "direct causal graph" (this work), the shortcut variable V is a direct cause of the outcome Y, shifting the marginal $P(Y)$ when the intervention variable I_V shifts the marginal $P(V)$.

2 THEORY

Consider predicting outcome Y (e.g., health status) from features X (e.g., an ECG recording) in the presence of an auxiliary covariate V (e.g., age or BMI). One source of model brittleness can be "shortcuts", or features that are predictive in the training distribution, but not predictive under relevant distribution shifts (Arjovsky et al., 2019; Geirhos et al., 2020). To cope with such instability, one may try to remove the shortcuts during learning. One common approach to shortcut removal assumes a non-causal association between the potential shortcut V and the outcome to be predicted Y (Heinze-Deml & Meinshausen, 2021; Veitch et al., 2021; Makar et al., 2022; Puli et al., 2022), which, for instance, can arise due to a confounding covariate between V and Y. The goal is then to seek a predictor using only X that performs well across a range of distributions. For example, Makar et al. (2022) develop a risk invariant predictor across a family of related probability distributions motivated by the graph depicted in Figure 1a, that can be simplified for our analysis to

$$\mathcal{P}_{spur} = \{P_s(X \mid Y, V) \, P_s(Y) \, P_t(V \mid Y)\}, \tag{1}$$

for a source distribution denoted by s and shifted target distributions indexed by t. All target distributions in this family of distributions thus factor as $P_t(X, Y, V) = P_s(X \mid Y, V) \, P_s(Y) \, P_t(V \mid Y)$, i.e., they vary only in $P(V|Y)$ from the source distribution, while $P(X \mid Y, V)$ and $P(Y)$ remain unchanged. Notably, assuming that $P(Y)$ remains the same across all potential shifted distributions, can be an unrealistically strong assumption in applications like healthcare. For example, we would expect the prevalence of heart diseases (Y) to be higher in an older population (V).

Instead, in this work, we consider the scenario where the shortcut variable (e.g., age or BMI) is a direct causal parent of the outcome we wish to predict (e.g., myocardial infarction in an ECG), as depicted in Figure 1b. In this scenario, we wish to form good predictions for the family of distributions

$$\mathcal{P}_{cause} = \{P_s(X \mid Y, V) P_s(Y \mid V) P_t(V)\}. \tag{2}$$

That is, we allow for changing marginal distribution of $P(V)$, while holding the conditional distributions $P(Y \mid V)$ and $P(X \mid Y, V)$ fixed.

Using the notion of so-called stable sets from Pfister et al. (2021), one can derive from the graph in Figure 1b which sets of predictors are associated with the same conditional expectation across different interventions on V by checking which sets of covariates block all paths between I_V and Y. Hence, in our model, to block the path $I_V \rightarrow V \rightarrow Y$, the covariate V must be included in the set of predictors. The predictive distribution derived from the source that conditions on X and V is then invariant across the entire family, i.e., $P_s(Y \mid X, V) = P_t(Y \mid X, V)$, whereas the predictive distribution that only conditions on X is not invariant, i.e., $P_s(Y \mid X) \neq P_t(Y \mid X)$ in general. We formalize this in the following proposition (proof in Appendix A.1).

Proposition 1 *For any element $P_t \in \mathcal{P}_{cause}$ as defined in Eq. (2), it holds that $P_t(Y|X, V) = P_s(Y|X, V)$. Furthermore, for such a P_t, in general $P_t(Y|X) \neq P_s(Y|X)$.*

(a) "Extended confounded graph"   (b) "Extended direct causal graph" (this work).

Figure 2: Extended versions of causal graphs in Figure 1. (a) Graph considered in Makar et al. (2022) explicitly including invariant latent variable X^* (still leading to shifts in $\mathcal{P}_{spur}$). Here, X^* is a latent variable that describes variation in X caused by Y. Recovering the predictive signal $e(X) = X^*$ yields a predictor that is invariant across interventions I_V, but one that does not use information from V about Y. (b) Direct graph explicitly including X^* (still leading to shifts in $\mathcal{P}_{cause}$). In this setting, recovering X^* does not guarantee an invariant predictor across shifts due to I_V, demonstrated in Section 3

Hence, empirical risk minimization (ERM) using $\{V, X\}$ as predictors would yield a robust model with respect to $\mathcal{P}_{cause}$ while ERM using $\{X\}$ only would not.

Remark 1 *Even an invariant representation that is invariant to V and encodes only the information in X related to Y (e.g., X^* in Makar et al. (2022)) would suffer from a degradation in performance across the family $\mathcal{P}_{cause}$.*

We illustrate these findings with a simulation study in the next section.

3 EXPERIMENTS

To illustrate our findings above regarding the consequences of shifts in causally influencing covariates V, we set up a simulation from a simple example. To allow for analyses of the behaviour of invariant methods as well, we roll out the graphs from Figure 1 similar to Makar et al. (2022) by explicitly including an unobserved variable X^* (see Figure 2). Here, $X^* = e(X)$ for some function e is assumed to be a latent variable that only contains information about X that is related to Y and as such is invariant to V when conditioned on Y.

In this extended setting (2), we define our data generating process as

$$p(V = 1) = p \tag{3}$$
$$P(Y = 1 \mid V = 0) = .2 \tag{4}$$
$$P(Y = 1 \mid V = 1) = .9 \tag{5}$$
$$P(X \mid Y = y, V = v) = \mathcal{N}(\mu_{y,v}, 1) \tag{6}$$
$$P(X^* \mid Y = y) = \mathcal{N}(\mu_{y,0}, 1). \tag{7}$$

where $\mu_{0,0} = -2/3, \mu_{1,0} = 2/3, \mu_{0,1} = -.8$, and $\mu_{1,1,} = .8$. As such, it is constructed to allow to analyse shifts of the family $\mathcal{P}_{cause}$, where $P(V)$ can be shifted by varying p while $P(Y|V)$ and $P(X|Y, V)$ will remain the same.

In Figure 3, we compare the performance of the predictors $P_s(Y \mid X)$, $P_s(Y \mid X^*)$ and $P_s(Y \mid X, V)$ on distributions where the marginal $P(V)$ has been shifted, and the source distribution marginal is $P_s(V = 1) = .1$. Predictors are obtained in closed form, and performance metrics are calculated on a sample of size 20,000 drawn according to the source distribution. As discussed in Section 2, this shift induces a shift in $P(Y)$, $P(Y \mid X)$, *and* $P(Y \mid X^*)$, causing a degradation in performance of the predictors $P_s(Y \mid X)$ and $P_s(Y \mid X^*)$, but not of $P_s(Y \mid X, V)$, i.e. conditioning on V restores performance across distribution shifts.

(a) Accuracy across distribution shifts.

(b) AUCs across distribution shifts.

(c) General likelihood, $P(X \mid Y, V)$.

(d) Invariant likelihood, $P(X^* \mid Y)$.

Figure 3: Simulation study described in Section 3. Panel (a) compares the predictive accuracy of the models $P_s(Y \mid X)$, $P_s(Y \mid X^*)$ and $P_s(Y \mid X, V)$ as a function of the target marginal $P_t(V) = p$, with source $P_s(V) = .1$. Only $P_s(Y \mid X, V)$ does not degrade in accuracy as $P_t(V)$ shifts further away from $P_s(V)$. Panel (b) compares the predictive AUC of the same three models. Note that the predictor using X^*, $P_s(Y \mid X^*)$ does achieve invariance in AUC across shifts, but not accuracy (or likelihood), and cannot make use of information about Y from V. Panel (c) depicts the four likelihood models (one for each combination of Y and V) — note that $V = 1$ further separates the conditional distributions, making separation easier (hence the AUC goes up in Panel (b) as p increases). Panel (d) depicts a $P(X^*|Y, V)$, which is the same across values of V (unlike $P(X|Y, V)$ in Panel (c)). The overall key takeaway is the robustness of $P_s(Y \mid X, V)$, i.e. the model conditioning on both V and X versus the lack of robustness in models conditioning only on X or X^* in terms of predictive accuracy in Panel (a) (even when their AUC is robust across shifts, Panel (b)).

As a side note, note that the AUC performance of $P(Y \mid X^*)$ does not degrade, though a general risk (like accuracy or log-likelihood) does degrade. This is due to the fact that shifts in $P(V)$ only influence $P(Y \mid X^*)$ through the prevalence $P_t(Y) = \sum_{v'} P_s(Y \mid V = v') P_t(V = v')$, not through X^*. The AUC metric is invariant to prevalence, but general metrics like accuracy, log-likelihood, and calibration are sensitive to prevalence. Also note that the difference in AUC performance of $P(Y \mid X^*)$ and $P(Y \mid X)$ is due to the construction of the class conditional distributions depicted in Figures 3c and 3d.

Overall, the degradation (or robustness) of performance across shifts of the family $\mathcal{P}_{cause}$ is the main illustrative point to be observed in Figure 3 and this section.

4 DISCUSSION

Our theoretical findings show that for settings in which auxiliary covariates V causally influence the outcome of interest Y (rather than just being spuriously correlated to them), $P(Y|X, V)$ remains stable across shifts in $P(V)$, while $P(Y|X)$ in general does not. As such, regressing Y only on X to learn $P(Y|X)$ (or invariant derivations thereof) will lead to predictions that are not robust to such shifts, while regressing Y on X, V recovers the desired robustness, as we empirically demonstrate on simulated data. The canonical next step of our analysis is to demonstrate our findings on real world data in healthcare applications, where we would expect shifts of the family $\mathcal{P}_{cause}$ to occur in practice.

REFERENCES

Martin Arjovsky, Léon Bottou, Ishaan Gulrajani, and David Lopez-Paz. Invariant risk minimization. *arXiv preprint arXiv:1907.02893*, 2019.

Yaroslav Ganin, Evgeniya Ustinova, Hana Ajakan, Pascal Germain, Hugo Larochelle, François Laviolette, Mario Marchand, and Victor Lempitsky. Domain-adversarial training of neural networks. *The journal of machine learning research*, 17(1):2096–2030, 2016.

Robert Geirhos, Jörn-Henrik Jacobsen, Claudio Michaelis, Richard Zemel, Wieland Brendel, Matthias Bethge, and Felix A Wichmann. Shortcut learning in deep neural networks. *Nature Machine Intelligence*, 2(11):665–673, 2020.

Christina Heinze-Deml and Nicolai Meinshausen. Conditional variance penalties and domain shift robustness. *Machine Learning*, 110:303–348, 2021. doi: https://doi.org/10.1007/s10994-020-05924-1.

Etsuro Himeno, Kenji Nishino, Yoshiyuki Nakashima, Akio Kuroiwa, and Masaharu Ikeda. Weight reduction regresses left ventricular mass regardless of blood pressure level in obese subjects. *American heart journal*, 131(2):313–319, 1996.

Beverly H Lorell and Blase A Carabello. Left ventricular hypertrophy: pathogenesis, detection, and prognosis. *Circulation*, 102(4):470–479, 2000.

Maggie Makar, Ben Packer, Dan Moldovan, Davis Blalock, Yoni Halpern, and Alexander D'Amour. Causally motivated shortcut removal using auxiliary labels. In Gustau Camps-Valls, Francisco J. R. Ruiz, and Isabel Valera (eds.), *Proceedings of The 25th International Conference on Artificial Intelligence and Statistics*, volume 151 of *Proceedings of Machine Learning Research*, pp. 739–766. PMLR, 28–30 Mar 2022.

Niklas Pfister, Evan G. Williams, Jonas Peters, Ruedi Aebersold, and Peter Bühlmann. Stabilizing variable selection and regression. *The Annals of Applied Statistics*, 15(3):1220 – 1246, 2021. doi: 10.1214/21-AOAS1487.

Aahlad Manas Puli, Lily H Zhang, Eric Karl Oermann, and Rajesh Ranganath. Out-of-distribution generalization in the presence of nuisance-induced spurious correlations. In *International Conference on Learning Representations*, 2022.

Victor Veitch, Alexander D'Amour, Steve Yadlowsky, and Jacob Eisenstein. Counterfactual invariance to spurious correlations in text classification. In A. Beygelzimer, Y. Dauphin, P. Liang, and J. Wortman Vaughan (eds.), *Advances in Neural Information Processing Systems*, 2021.

A APPENDIX

A.1 PROOF OF PROPOSITION 1

First, we show that for any element P_t of this family, it holds that $P_t(Y|X,V) = P_s(Y|X,V)$. Remember that by the definition of $\mathcal{P}_{cause}$, we have that $P_t(X \mid Y,V) = P_s(X \mid Y,V)$ and $P_t(Y \mid V) = P_s(Y \mid V)$. From this, it quickly follows that

$$P_t(X|V) = \int P_t(X|Y,V)P_t(Y|V)dY \tag{8}$$

$$= \int P_s(X|Y,V)P_s(Y|V)dY \tag{9}$$

$$= P_s(X|V) \tag{10}$$

Then, using basic probability calculus, it follows

$$P_t(Y|X,V) = \frac{P_t(Y,X,V)}{P_t(X,V)} \tag{11}$$

$$= \frac{P_s(X|Y,V)P_t(Y|V)P_t(V)}{P_t(X|V)P_t(V)} \tag{12}$$

$$= \frac{P_s(X|Y,V)P_s(Y|V)}{P_t(X|V)} \tag{13}$$

$$= \frac{P_s(X|Y,V)P_s(Y|V)}{P_s(X|V)} \tag{14}$$

$$= \frac{P_s(X|Y,V)P_s(Y|V)P_s(V)}{P_s(X|V)P_s(V)} \tag{15}$$

$$= P_s(Y|X,V) \tag{16}$$

Next, we show that for such an element P_t of this family, in general $P_t(Y|X) \neq P_s(Y|X)$. Using the above result, this is indeed easy to see when marginalising over V:

$$P_t(Y|X) = \int P_t(Y|X,V)P_t(V|X)dV \tag{17}$$

$$= \int P_s(Y|X,V)P_t(V|X)dV \tag{18}$$

$$= \int P_s(Y|X,V)\frac{P_t(X|V)P_t(V)}{P_t(X)}dV \tag{19}$$

$$= \int P_s(Y|X,V)P_s(X|V)\frac{P_t(V)}{P_t(X)}dV \tag{20}$$

$$\tag{21}$$

Since in general $\frac{P_t(V)}{P_t(X)} \neq \frac{P_s(V)}{P_s(X)}$, this also implies that in general $P_t(Y|X) \neq P_s(Y|X)$.