Wasserstein-type Gaussian Process Regressions for Input Measurement Uncertainty

Abstract

Gaussian process (GP) regression is widely used for uncertainty quantification, yet the standard formulation assumes noise-free covariates. When inputs are measured with error, this errors-in-variables (EIV) setting can lead to optimistically narrow posterior intervals and biased decisions. We study GP regression under input measurement uncertainty by representing each noisy input as a probability measure and defining covariance through Wasserstein distances between these measures. Building on this perspective, we instantiate a deterministic projected Wasserstein ARD (PWA) kernel whose one-dimensional components admit closed-form expressions and whose product structure yields a scalable, positive-definite kernel on distributions. Unlike latent-input GP models, PWA-based GPs (PWA-GPs) handle input noise without introducing unobserved covariates or Monte Carlo projections, making uncertainty quantification more transparent and robust.

1 Introduction

Gaussian Processes (GPs) are core methods for stochastic function approximation in machine learning and surrogate modeling. They are fully defined by a prior mean and a covariance kernel, making them analytically tractable and able to encode function smoothness (Williams and Rasmussen, 2006). GPs output predictive distributions rather than point estimates and quantify uncertainty and confidence. In a Bayesian framework GPs also propagate hyperparameter uncertainty. They have been applied in forward modeling (Deisenroth, 2010; Vinogradska et al., 2016), uncertainty quantification (Tuo and Wang, 2022; Wang, 2021), parameter optimization (Liu et al., 2021; Qiang et al., 2024), and autonomous experimentation (Noack et al., 2019; Stach et al., 2021; Noack et al., 2021; Thomas et al., 2022). Standard GP regression assumes noise-free inputs X, modeling observations from $f : \mathbb{R}^d \to \mathbb{R}$ as

$$y = f(X) + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2).$$

When the inputs are noisy, i.e., we observed $U = X + \epsilon_X$ as inputs, this "errors-in-variables" (EIV) setting can degrade performance (Zhou et al., 2023). Figure 1 shows how input uncertainty can lead a GP failing to estimate f. Thus, to recover the true function for noisy input, GPs must incorporate input uncertainty or address EIV. The following proposition describes the effect of EIV in terms of the uncertainty quantification of prediction.

The next result provides the technical link between the proposed Wasserstein GP and uncertainty quantification, it shows that a standard GP can undercover when input noise is ignored.

Proposition 1. *Consider the errors-in-variables model*

$$Y = f(X + \varepsilon_X) + \varepsilon, \qquad \varepsilon_X \sim \mathcal{N}(0, \Sigma_X), \quad \varepsilon \sim \mathcal{N}(0, \sigma^2),$$

with $\varepsilon_X \perp \varepsilon$. Let $f(x) = c + w^\top x$ be affine with $w \neq 0$. Define the naive $(1 - \alpha)$ interval that ignores input noise:

$$I_\alpha := [Y - z_{1-\alpha/2}\sigma, \ Y + z_{1-\alpha/2}\sigma].$$

Figure 1: Illustration of error-in-variable regression problem. In both panels, the true function is $y = f(X) = \frac{\sin(10\pi \cdot X)}{2X} + (X-1)^4$, but in each case the "true" input locations X are contaminated with measurement errors (with standard deviation of 0.01 and 0.05 on the left and right, respectively). Using a GP to accurately infer the true function (the blue line) must account for the fact that the input locations U are uncertain.

Then for any fixed X,

$$\mathbb{P}\big(f(X) \in I_\alpha \mid X\big) = 2\Phi\left(\frac{z_{1-\alpha/2}\sigma}{\sqrt{\sigma^2 + w^\top \Sigma_X w}}\right) - 1 \ < \ 1 - \alpha$$

whenever $w^\top \Sigma_X w > 0$.

Proposition 1 formalizes a simple failure mode of a naive GP in an errors-in-variables setting. The interval I_α accounts only for the output noise variance σ^2, while the true target $f(X) = c + w^\top X$ also varies through the input error term $w^\top \varepsilon_X$, whose variance is $w^\top \Sigma_X w$. As a result, the nominal $(1-\alpha)$ interval is too narrow and its actual coverage is strictly smaller than $1-\alpha$. In this paper, we refer to this phenomenon as *undercoverage*.

One approach to taking into account EIV uses latent-variable models: Kennedy and O'Hagan (2001) treat true inputs as latent and integrate them out; Girard et al. (2002) apply this to time series; Damianou and Lawrence (2013) model inputs and outputs jointly; and McHutchon and Rasmussen (2011) convolve the GP prior with an input-noise distribution. These methods capture input uncertainty but require high-dimensional inference, which can be slow or unstable in non-Gaussian settings, and often rely on numerical approximations that may bias results.

Alternatively, one can use kernels on probability measures. If each input is treated as a distribution, for example $U \sim \mathcal{N}(\mu_X, \Sigma_X)$, kernels based on Wasserstein distances compare these inputs directly (see Panaretos and Zemel, 2019). Candelieri et al. (2022) study univariate discrete measures, and Meunier et al. (2022); Bonet et al. (2023) develop sliced Wasserstein kernels for general distributions.

In this work, we view each noisy input U_i as a probability measure μ_i and place a GP prior directly on the space of such measures. We instantiate this idea with the PWA kernel, which preserves the familiar automatic relevance determination (ARD) structure while encoding input uncertainty through Wasserstein distances between one-dimensional marginals. Intuitively, when the input noise is small, the PWA-GP behaves similarly to a GP fitted on the (latent) noise-free inputs (X, Y), whereas increasing dispersion of the input distributions naturally inflates posterior uncertainty, leading to more honest predictive intervals in error-in-variables problems.

In the PWA kernel (6), separate kernel parameters for each input dimension act as ARD-style length scales, handling dimension-specific uncertainty via an automatic relevance determination construction (Williams and Rasmussen, 2006). This avoids full multivariate optimal transport and randomized slicing while still capturing anisotropic input variability in a way that directly serves uncertainty quantification rather than merely introducing a new kernel form.

In EIV problems, finite-slice implementations of sliced Wasserstein kernels (Meunier et al., 2022) introduce additional Monte Carlo approximation error on top of measurement noise. In contrast, our PWA construction uses deterministic (coordinate or fixed-basis) projections, avoiding slice-induced randomness while retaining valid projected kernels and ARD flexibility. Our theoretical analysis focuses on the case $p = 1$, where covering-number arguments yield uniform error bounds. We summarize related methods in Table A.1, discuss computational complexities in Appendix G, and compare a representative subset in our uncertainty-quantification experiments.

Summary of our contributions.

- We focus on uncertainty quantification for GP regression with input measurement error, modeling each noisy input as a probability measure and working in the Wasserstein space of distributions. In a simple linear-Gaussian EIV model we show that standard GPs that ignore input noise produce predictive intervals that strictly undercover.

- We develop a deterministic PWA kernel and its principal-component variant PCPWA, which define GPs directly on distributions via products of closed-form one-dimensional Wasserstein kernels. The construction preserves automatic relevance determination (ARD) while avoiding latent variables and stochastic slicing schemes commonly used for error-in-variables GPs.

- We establish a deterministic net-extension bound for the posterior mean of PWA-based GPs and, under a well-specified GP prior, obtain a uniform high-probability posterior band controlled by the posterior standard deviation. This yields a simple sufficient condition under which the resulting credible intervals are conservative.

- Through experiments on synthetic data, accelerator calibration, and NOAA drifter trajectories, we demonstrate that PWA-GPs provide competitive point predictions and markedly improved uncertainty quantification in terms of coverage and CRPS and relative to standard kernels and sliced Wasserstein baselines under input noise.

2 GPs with Wasserstein-type Kernels

2.1 Wasserstein and Gromov-Wasserstein distances

Wasserstein distances improve how distributional inputs are handled in machine learning. Arjovsky et al. (2017) highlighted their use for output distributions, but they also apply to input uncertainty.

Intuitively, imagine moving "sand" from a pile at x to fill a hole at y: the cost is the mass moved times the distance (Mallasto and Feragen, 2017). We write $\mathcal{P}_p(X)$ for the set of Borel probability measures on X with finite pth moment, i.e. $\int d_X(x, x_0)^p \, d\mu(x) < \infty$ for some (equivalently any) $x_0 \in X$. Formally:

Definition 2. *(p-Wasserstein distance) Let $p \in [1, \infty)$ and let $\mu, \nu \in \mathcal{P}_p(X)$ be probability measures on a metric space (X, d_X) with finite pth moments. The p-Wasserstein distance is*

$$W_p(\mu, \nu) = \left(\inf_{\gamma \in \Gamma(\mu, \nu)} \int_{X \times X} d_X(x, y)^p \, d\gamma(x, y) \right)^{1/p}, \tag{1}$$

where $\Gamma(\mu, \nu)$ denotes the set of couplings with marginals μ and ν. The case $p = 1$ is the Earth Mover's Distance (Peyré et al., 2019).

$W_p(\mu, \nu)$ is the minimal "effort" to transport μ to ν, raising cost to the pth power. These distances scale polynomially in asymptotic regimes (Panaretos and Zemel, 2020; Weed and Bach, 2019).

Definition 3. *(Gromov-Wasserstein distance) Let $\mu \in \mathcal{P}_p(X)$ and $\nu \in \mathcal{P}_p(Y)$, where (X, d_X) and (Y, d_Y) are metric spaces. The p-Gromov-Wasserstein distance is*

$$GW_p(\mu, \nu) := \left(\inf_{\pi \in \Gamma(\mu, \nu)} \iint_{(X \times Y)^2} \left| d_X(x, x') - d_Y(y, y') \right|^p \, d\pi(x, y) \, d\pi(x', y') \right)^{1/p}. \tag{2}$$

GW_p matches the relational geometry of X and Y, finding correspondences that minimize distortion.

In essence, GW_p generalizes W_p to compare distributions on different spaces by focusing on pairwise distances. W_p uses a common ground metric, while GW_p handles cross-domain structures, making both valuable for pratical tasks in image processing, machine learning, and statistical physics. However, we lack knowledge of the PD property for GW-type kernels.

While the statement of Wasserstein and Gromov-Wasserstein kernels is straightforward, calculating W_p and GW_p can be quite challenging. For $\boldsymbol{x} \in \mathbb{R}^1$, we can have a closed-form expression for any $p \geq 1$:

Proposition 4. *(Meunier et al., 2022) Let μ, ν be probability measures defined on the real line $\mathbb{R}^1$ and denote their cumulative distribution functions by $F_\mu(x)$ and $F_\nu(x)$. Then,*

$$W_p(\mu, \nu) = \left(\int_0^1 \left| F_\mu^{-1}(q) - F_\nu^{-1}(q) \right|^p \, dq \right)^{1/p}. \tag{3}$$

2.2 Wasserstein-type covariance kernels

We can define k_{W_p} as follows,

$$k_{W_p}(\mu, \nu) = \lambda \exp\bigl(-\sigma \, W_p(\mu, \nu)^p\bigr) \tag{4}$$

To use k_{W_p} as a GP covariance function, it is not enough that W_p be a metric. A standard sufficient condition is that the exponent $\psi(\mu, \nu)$ be *conditionally negative definite* (CND): a symmetric function ψ on $\mathcal{P}$ is CND if for every $n \geq 1$, every $\mu_1, \ldots, \mu_n \in \mathcal{P}$, and every coefficients $c_1, \ldots, c_n \in \mathbb{R}$ satisfying $\sum_{i=1}^n c_i = 0$, one has

$$\sum_{i,j=1}^n c_i c_j \psi(\mu_i, \mu_j) \leq 0.$$

By Schoenberg's theorem, if ψ is CND and $\psi(\mu, \mu) = 0$, then $k(\mu, \nu) = \exp(-\sigma \psi(\mu, \nu))$ is positive definite for every $\sigma > 0$.

In (4), the exponent is $\psi(\mu, \nu) = W_p(\mu, \nu)^p$, not $W_p(\mu, \nu)^2$ unless $p = 2$. In general, $W_p(\mu, \nu)^p$ need not be CND on $\mathbb{R}^d$ for arbitrary $d > 1$, so positive definiteness is not automatic outside special settings. This is why we either restrict attention to cases where positive definiteness is known (e.g., certain one-dimensional or Gaussian settings) or use projected/sliced constructions whose one-dimensional factors are positive definite.

A special case is when the inputs are Gaussian and $p = 2$: here the projection is not needed, the kernel (4) is positive definite in the Gaussian setting, and closed forms for W_2 enable efficient computation. For example,

$$W_2^2\bigl(\mathcal{N}(\mu_x, \Sigma_x), \mathcal{N}(\mu_y, \Sigma_y)\bigr) = \|\mu_x - \mu_y\|^2 + \mathrm{tr}\Bigl(\Sigma_x + \Sigma_y - 2(\Sigma_x^{1/2}\Sigma_y\Sigma_x^{1/2})^{1/2}\Bigr).$$

If $\Sigma_x = \Sigma_y$, this reduces to $\|\mu_x - \mu_y\|^2$, so (4) becomes the usual squared-exponential (Gaussian/RBF) kernel on the means (Dowson and Landau, 1982; Peyré et al., 2019; Horn and Johnson, 2012). Alternatively, one may use a *sliced* kernel by averaging over 1D projections $u \in \mathbb{S}^{d-1}$,

$$k(\mu, \nu) = \lambda \exp\bigl(-\sigma \, \mathbb{E}_u[W_p(\mu_u, \nu_u)^p]\bigr), \tag{5}$$

which also remains PD (Meunier et al., 2022). Computationally we have also derived bounds on derivatives for numerical stability (See Appendix F).

Model names used in the paper. We refer to the GP prior with covariance k_{W_p} in (4) as a *Wasserstein GP* (WGP). The GP using the sliced kernel (5) is denoted SWGP. The GP using our projected ARD kernel (6) is denoted PWA-GP, and its fixed-basis/principal-component variant (7) is denoted PCPWA. After these first definitions we use only the abbreviations.

To ensure PD in higher dimensions without solving a d-dimensional optimal transportation problem, we use *Projected Wasserstein ARD* kernels. For measures μ, ν on $\mathbb{R}^d$, define

$$k(\mu, \nu) = \lambda \prod_{i=1}^d \exp\bigl(-\sigma_i \, W_p(\mu_i, \nu_i)^p\bigr), \tag{6}$$

where μ_i is the marginal distribution of measure μ along the i-th dimension defined on $\mathbb{R}^1$; whenever the 1D factors are PD, so is the product kernel (Wilson and Nickisch, 2015; Bilionis et al., 2013). Similar construction can be done using Gromov-Wasserstein distances, yet the projections can be defined through marginals of different dimensions. Unlike previous EIV methods that introduce latent true inputs and require high-dimensional integration or approximate marginalization (Kennedy and O'Hagan, 2001; Girard et al., 2002; Damianou and Lawrence, 2013; McHutchon and Rasmussen, 2011), PWA avoids explicit latent-variable inference.

Although this is a hybrid of ARD kernel and projection technique, we found this a novel construction for EIV problem that does not exist in the current literature. This is different from the sliced Wasserstein kernels (Meunier et al., 2022) (i.e., "project-Wasserstein/average"), since each (λ_i, σ_i) can adapt to dimension-specific uncertainty (i.e., ARD type construction), avoiding the cost of a full multivariate transport and iterative projections during the random slicing. This is also different from earlier work by Bachoc et al. (2017), where they proposed to consider $p = 2$ only; and PWA-GP is also different from their "PCA" GP, which uses truncated principal component projections ARD kernel without computing Wasserstein distances over marginals (i.e., "project-ARD"). Our proposed kernel (6) is a generalization introducing the scheme of "Projected Wasserstein ARD".

Beyond projecting onto coordinate axes, one may also project onto a fixed orthonormal basis of $\mathbb{R}^d$, such as principal components. Let $\{v_r\}_{r=1}^m \subset \mathbb{R}^d$ be fixed orthonormal directions and define the linear functional

$$\ell_r : \mathbb{R}^d \to \mathbb{R}, \qquad \ell_r(x) = v_r^\top x.$$

For a Borel probability measure μ on $\mathbb{R}^d$, define its projected marginal along v_r by

$$\mu_{v_r} := (\ell_r)_\# \mu.$$

Equivalently, for every Borel set $A \subset \mathbb{R}$, $\mu_{v_r}(A) = \mu(\{x \in \mathbb{R}^d : v_r^\top x \in A\})$. Here $(\ell_r)_\# \mu$ denotes the push-forward of μ through ℓ_r. For a Borel probability measure μ on $\mathbb{R}^d$, we denote by $\mu_{v_r} := (\ell_r)_\# \mu$ the push-forward of μ through ℓ_r, i.e., the one-dimensional marginal of μ along direction v_r. Explicitly, for every Borel set $A \subset \mathbb{R}$,

$$\mu_{v_r}(A) = ((\ell_r)_\# \mu)(A) := \mu(\ell_r^{-1}(A)) = \mu(\{x \in \mathbb{R}^d : v_r^\top x \in A\}).$$

We use the notation $(\ell_r)_\# \mu$ rather than $\ell_r(\mu)$ because ℓ_r acts on points $x \in \mathbb{R}^d$, while the induced action on measures is the push-forward map.

We define the *principal-component projected Wasserstein ARD* (PCPWA) kernel by

$$k_{\mathrm{PCPWA}}(\mu, \nu) = \lambda \prod_{r=1}^m \exp(-\sigma_r \, W_p(\mu_{v_r}, \nu_{v_r})^p), \tag{7}$$

with direction-specific length scales $(\sigma_r)_{r=1}^m$ and a global amplitude λ.

When $m = d$ and $v_r = e_r$ are the canonical basis vectors, (7) reduces exactly to the coordinate PWA kernel (6), so PWA is a special case of PCPWA. Whenever the 1D factors in (7) are PD, their pullbacks by ℓ_r are PD on $\mathcal{P}_p(\mathbb{R}^d)$, and the product over r remains PD by closure under products. Thus, PCPWA inherits the same kernel-validity guarantees as PWA in those validated 1D settings while allowing projections onto any fixed system of directions, including principal components computed once from the design.

2.3 Uniform Error Bound for Wasserstein-type GP (p=1)

To assess models under input uncertainty, we derive a uniform bound on the posterior mean over a compact class of 1D distributions. Since our PWA kernel (6) is built from 1D marginals, we limit the theory below to the case $p = 1$. The deterministic net-extension argument parallels Theorem 3.1 of Lederer et al. (2019), while the posterior-probability corollary stated below additionally assumes a well-specified GP prior.

Let $[a, b] \subset \mathbb{R}$ and let $\mathcal{P}$ be a set of Borel probability measures supported on $[a, b]$ such that the Wasserstein covering numbers satisfy $M(\tau, \mathcal{P}) \le C\tau^{-\alpha_0}$ for some $\alpha_0 > 0$. Assume the quantile functions are uniformly

Lipschitz in their argument, i.e., there exists $\ell > 0$ such that

$$|F_\mu^{-1}(q) - F_\mu^{-1}(q')| \le \ell|q - q'| \qquad \text{for all } q, q' \in [0, 1], \ \mu \in \mathcal{P}.$$

Let $f : \mathcal{P} \to \mathbb{R}$ be L_f-Lipschitz with respect to W_1. Given observations $(\mu_i, y_i)_{i=1}^N$, define

$$K_{ij} := k(\mu_i, \mu_j), \qquad k_\mu := \big(k(\mu, \mu_1), \ldots, k(\mu, \mu_N)\big)^\top, \qquad \alpha := (K + \sigma_*^2 I_N)^{-1} y.$$

Then

$$\nu_N(\mu) = k_\mu^\top \alpha, \qquad \sigma_N^2(\mu) = k(\mu, \mu) - k_\mu^\top (K + \sigma_*^2 I_N)^{-1} k_\mu.$$

Define

$$L_k := \sup_{\nu \in \mathcal{P}} \sup_{\mu \ne \mu'} \frac{|k(\mu, \nu) - k(\mu', \nu)|}{W_1(\mu, \mu')}, \qquad L_{\nu_N} := N L_k \|\alpha\|_\infty,$$

and let ω_{σ_N} denote a modulus of continuity for σ_N, i.e.

$$|\sigma_N(\mu) - \sigma_N(\nu)| \le \omega_{\sigma_N}(\tau) \qquad \text{whenever } W_1(\mu, \nu) \le \tau.$$

The next theorem is the deterministic extension step that turns control on a finite τ-net into a uniform bound over $\mathcal{P}$.

Theorem 5. *Let $\{\bar{\mu}_1, \ldots, \bar{\mu}_M\}$ be a τ-net of $\mathcal{P}$ in W_1. Assume $f : \mathcal{P} \to \mathbb{R}$ is L_f-Lipschitz with respect to W_1, the kernel k is L_k-Lipschitz in its first argument with respect to W_1, and σ_N admits the modulus of continuity ω_{σ_N}. If, for some constant $B \ge 0$,*

$$|f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)| \le B\,\sigma_N(\bar{\mu}_j), \qquad j = 1, \ldots, M,$$

then for every $\mu \in \mathcal{P}$,

$$|f(\mu) - \nu_N(\mu)| \le B\,\sigma_N(\mu) + (L_f + L_{\nu_N})\tau + B\,\omega_{\sigma_N}(\tau).$$

The proof appears in Appendix D, which mirrors Lederer et al. (2019) but imposes conditions on L_∞ bounds instead of requiring its continuity, which is different from Lederer et al. (2019), and one can use the 1D CDF Lipschitz nets from Proposition 4 to bound $M(\tau, \mathcal{P})$.

The proof appears in Appendix D. It is a deterministic net-extension argument: the only stochastic ingredient enters later, when the finite-net inequalities are obtained from Gaussian posterior tails under a well-specified GP model.

Corollary 6. *Assume in addition that the regression model is well specified: $f \sim \mathcal{GP}(0, k)$, the design points $(\mu_i)_{i=1}^N$ are fixed, and the observations satisfy*

$$y_i = f(\mu_i) + \epsilon_i, \qquad \epsilon_i \overset{\text{ind}}{\sim} \mathcal{N}(0, \sigma_*^2), \qquad i = 1, \ldots, N.$$

Then for any $\tau > 0$ and $\delta \in (0, 1)$, conditional on $\mathcal{D}_N$, with posterior probability at least $1 - \delta$,

$$|f(\mu) - \nu_N(\mu)| \le \sqrt{\beta(\tau)}\,\sigma_N(\mu) + \gamma(\tau) \qquad \text{for all } \mu \in \mathcal{P},$$

where

$$\beta(\tau) = \left[\Phi^{-1}\left(1 - \frac{\delta}{2\,M(\tau, \mathcal{P})}\right)\right]^2, \qquad \gamma(\tau) = (L_f + L_{\nu_N})\,\tau + \sqrt{\beta(\tau)}\,\omega_{\sigma_N}(\tau).$$

The same argument applies to the PCPWA kernel with D replacing W_1.

Corollary 7. *Let $\mathcal{P}$ be a set of Borel probability measures on $\mathbb{R}^d$, and define*

$$D(\mu, \nu) := \left(\sum_{r=1}^m a_r\,W_1(\mu_{v_r}, \nu_{v_r})^2\right)^{1/2}.$$

for fixed directions $\{v_r\}_{r=1}^m \subset \mathbb{R}^d$ and weights $a_r > 0$. Assume $f : \mathcal{P} \to \mathbb{R}$ is L_f-Lipschitz with respect to D, the PCPWA kernel k_{PCPWA} from (7) is L_k-Lipschitz in its first argument with respect to D, and σ_N admits the modulus of continuity ω_{σ_N} with respect to D. If $(\mathcal{P}, D)$ admits a τ-net of size $M_{\mathrm{PCPWA}}(\tau, \mathcal{P})$ and the model is well specified with prior $f \sim \mathcal{GP}(0, k_{\mathrm{PCPWA}})$, then for any $\tau > 0$ and $\delta \in (0, 1)$, conditional on $\mathcal{D}_N$, with posterior probability at least $1 - \delta$,

$$|f(\mu) - \nu_N(\mu)| \leq \sqrt{\beta_{\mathrm{PCPWA}}(\tau)}\, \sigma_N(\mu) + \gamma_{\mathrm{PCPWA}}(\tau) \qquad \text{for all } \mu \in \mathcal{P},$$

where

$$\beta_{\mathrm{PCPWA}}(\tau) = \left[\Phi^{-1}\left(1 - \frac{\delta}{2\,M_{\mathrm{PCPWA}}(\tau, \mathcal{P})}\right)\right]^2, \qquad \gamma_{\mathrm{PCPWA}}(\tau) = (L_f + L_{\nu_N})\tau + \sqrt{\beta_{\mathrm{PCPWA}}(\tau)}\,\omega_{\sigma_N}(\tau).$$

For $p > 1$, it becomes more complicated since different projected marginals can have different covering numbers in the kernel construction (6). We therefore restrict the formal theory here to the $p = 1$ setting and leave sharper high-dimensional covering estimates for future work.

The preceding corollaries are net-based posterior band statements analogous in spirit to the Wasserstein-space analysis of Meunier et al. (2022), but here the finite-net step is made explicit and the deterministic extension from the net to the full input class is separated from the model-based Gaussian tail argument. This immediately implies the following proposition, which shows how the band width can prevent the undercoverage highlighted in Proposition 1.

Proposition 8. *Assume the posterior event in Corollary 6 holds with parameters (τ, δ), i.e., conditional on $\mathcal{D}_N$,*

$$|f(\mu) - \nu_N(\mu)| \leq \sqrt{\beta(\tau)}\, \sigma_N(\mu) + \gamma(\tau) \quad \text{for all } \mu \in \mathcal{P}$$

with posterior probability at least $1 - \delta$. Fix $\alpha \in (0, 1)$ and $z := z_{1-\alpha/2}$. If

$$z > \sqrt{\beta(\tau)} \qquad \text{and} \qquad \sigma_N(\mu_X) \geq \frac{\gamma(\tau)}{z - \sqrt{\beta(\tau)}},$$

then

$$\mathbb{P}\big(f(\mu_X) \in \big[\nu_N(\mu_X) \pm z\, \sigma_N(\mu_X)\big] \,\big|\, \mathcal{D}_N\big) \geq 1 - \delta.$$

This leads to larger posterior variance where input uncertainty is high, increasing the width of credible intervals appropriately. The statement is Bayesian: under the well-specified GP model, the posterior credible interval is conservative whenever the variance lower bound above holds.

Sliced Wasserstein kernels (5) are positive definite, but in practice are implemented using a finite number of random projections, which introduces Monte Carlo approximation error. By contrast, PWA/PCPWA are deterministic projections and avoid this additional randomness.

Proposition 9. *Fix a direction $u \in \mathbb{S}^{d-1}$ and assume the analogue of Corollary 6 holds for the 1D projected model with parameters (τ, δ): conditional on $\mathcal{D}_N$,*

$$|f(\mu) - \nu_N^u(\mu)| \leq \sqrt{\beta_u(\tau)}\, \sigma_N^u(\mu) + \gamma_u(\tau) \quad \text{for all } \mu \in \mathcal{P}$$

with posterior probability at least $1 - \delta$. Fix $\alpha \in (0, 1)$ and $z := z_{1-\alpha/2}$. If

$$z > \sqrt{\beta_u(\tau)} \qquad \text{and} \qquad \sigma_N^u(\mu_X) \geq \frac{\gamma_u(\tau)}{z - \sqrt{\beta_u(\tau)}},$$

then

$$\mathbb{P}\big(f(\mu_X) \in \big[\nu_N^u(\mu_X) \pm z\, \sigma_N^u(\mu_X)\big] \,\big|\, \mathcal{D}_N\big) \geq 1 - \delta.$$

3 Experiments and Applications

Practitioners who model input uncertainty with Gaussian summaries often use a covariance-only input map $x \mapsto \Sigma_x$, so that each input location x is represented as $X \sim \mathcal{N}(\mu_x, \Sigma_x)$ and only the covariance Σ_x enters the kernel (Moreno-Muñoz et al., 2018). This approach is efficient when noise is well-approximated by local Gaussian perturbations but cannot represent skew, multimodality, or complex support geometry. Though we noticed the recent important work (Song and Han, 2023) tackles the input uncertainty by modeling in the space of covariates, it is not completely clear to us how their low-rank linear models can be compared to non-linear models and thus we did not include their method in the comparison. We cannot use histogram discretizations like the previous Wasserstein GP literature (Meunier et al., 2022; Bachoc et al., 2017), since the bin-width selection has major effect on the subsequent model. We evaluate our proposed PWA-GP (and PCPWA where noted) on synthetic and real datasets using: (1) RMSE for posterior-mean prediction accuracy; (2) CRPS (Arnold et al., 2024),which jointly assesses predictive precision and sharpness, and (3) the probability coverage, which measures how well the posterior uncertainty covers the measured test observations. Appendix B contains the definitions of these metrics.

3.1 Simulated distributional regression experiments

We study several synthetic distributional regression problems designed to probe errors-in-variables (EIV) behavior and uncertainty quantification. In all cases, each covariate is a finite cloud $U_i = \{u_{ij}\}_{j=1}^{n_i} \subset \mathbb{R}^d$ with associated response Y_i, and the task is to predict Y for new clouds. We report test RMSE, empirical coverage of nominal 90% predictive intervals, and CRPS for Regular GP on empirical means, Aggregated GP on individual samples, Wasserstein GPs (WGP, SWGP, PWA-GP, PCPWA), uncertain-input GP UI-GP Girard et al. (2002), and KME/MMD distributional GPs (Meunier et al., 2022).

1D scenarios. We consider three one-dimensional settings with Gaussian output noise $\eta_i \sim \mathcal{N}(0, 0.05^2)$. (i) **1D-EIV**: latent covariates x_i are evenly spaced on $[0.05, 0.95]$. We observe $U_i = \{x_i + \varepsilon_{ij}\}_{j=1}^{n_i}$ with heteroscedastic Gaussian measurement noise $\varepsilon_{ij} \sim \mathcal{N}(0, \sigma_i^2)$, where σ_i^2 increases with x_i, and set $Y_i = f(x_i) + \eta_i$ for a nonlinear oscillatory f. This is a canonical errors-in-variables problem. (ii) **1D-Var**: U_i consists of $u_{ij} \sim \mathcal{N}(\mu_i, \sigma_i^2)$ with varying variances, and the response depends on both mean and spread, $Y_i = \sin(2\pi\mu_i) + 0.5\sigma_i^2 + \eta_i$. (iii) **1D-Skew**: inputs are log-normal, $u_{ij} \sim \exp(\mathcal{N}(m_i, s_i^2))$, and the response depends on an inter-quantile range and location, $Y_i = [Q_{0.8}(U_i) - Q_{0.2}(U_i)] + 0.3\sin(m_i) + \eta_i$, where $Q_q(U_i)$ denotes the q-quantile of the input distribution (or, equivalently in the simulations, of the empirical cloud U_i). 1D-Var and 1D-Skew thus require sensitivity to higher-order distributional features.

2D scenarios. We next consider two two-dimensional problems. (iv) **2D-mean** (Gaussian location functional): for each i we draw $\mu_i \in [0.1, 1]$ and sample $u_{ij} \sim \mathcal{N}(\mu_i \mathbf{1}_2, \mathrm{diag}(\sigma_{i1}^2, \sigma_{i2}^2))$ with moderate variances. The response is a smooth function of the mean of U_i, averaged over coordinates,

$$Y_i = \frac{1}{n_i} \sum_{j=1}^{n_i} \big[\sin(u_{ij}) + 2e^{u_{ij}} \big] + \eta_i,$$

so the problem is essentially a Gaussian location model. (v) **2D-aniso-PC** (anisotropic rotated subspace): latent scalars $z_i \in [0.05, 0.95]$ generate samples in a rotated basis $(z_{ij}^{\parallel}, z_{ij}^{\perp})$ via $z_{ij}^{\parallel} = z_i + \epsilon_{ij}^{\parallel}$, $z_{ij}^{\perp} = \epsilon_{ij}^{\perp}$ with $\epsilon_{ij}^{\parallel} \sim \mathcal{N}(0, \sigma_{\parallel}^2(z_i))$ and $\epsilon_{ij}^{\perp} \sim \mathcal{N}(0, \sigma_{\perp}^2)$, where $\sigma_{\parallel}^2(z_i)$ increases with z_i and $\sigma_{\perp}^2$ is small. A fixed 45° rotation R maps $u_{ij} = R[z_{ij}^{\parallel}, z_{ij}^{\perp}]^{\top}$, and the response depends only on z_i, $Y_i = \sin(4\pi z_i) + 0.5z_i + \eta_i$. Here the intrinsic variation lies along a rotated one-dimensional subspace, so PCA-based kernels such as PCPWA are expected to help.

High-dimensional scenarios. Finally, we construct high-dimensional problems from the Ackley function. (vi) **HD-Ackley-5D** and (vii) **HD-Ackley-10D**: latent covariates $x_i \in [-2, 2]^d$ with $d \in \{5, 10\}$ are drawn uniformly, we observe noisy clouds $u_{ij} = x_i + \varepsilon_{ij}$ with $\varepsilon_{ij} \sim \mathcal{N}(0, 0.1^2 I_d)$, and set $Y_i = f_{\mathrm{Ackley}}(x_i) + \eta_i$. These scenarios test how methods scale with dimension in the presence of input noise and nontrivial curvature.

Table 4 reports the resulting RMSE, coverage, and CRPS. In 1D-EIV, Regular GP achieves RMSE 0.309 but under-covers (65%), Aggregated GP is both inaccurate and grossly over-confident (5% coverage), while WGP, PWA-GP, PCPWA, and SWGP all attain similar RMSE (≈ 0.30) with coverage ≈ 0.97 and the lowest CRPS (≈ 0.17), matching our EIV coverage analysis. In 1D-Var and 1D-Skew, mean- or moment-based baselines (Regular, UI-GP, KME/MMD) either lose accuracy or under-cover, whereas Wasserstein kernels maintain competitive RMSE and near-nominal coverage. In 1D-Var the Regular GP has the best RMSE/CRPS, but the Wasserstein methods are within about 10-15% in RMSE; in 1D-Skew, MMD-GP attains the lowest RMSE and CRPS but with coverage 0.58, whereas the Wasserstein models trade a slightly higher RMSE for better-calibrated intervals.

In the benign 2D-mean scenario, uncertain-input and KME/MMD GPs-whose assumptions match the Gaussian location structure-achieve the smallest RMSE (0.0004-0.0026) with near-nominal coverage and CRPS, while Wasserstein kernels remain reasonable but conservative (e.g. PWA-GP RMSE 0.0462, coverage 1.0). This shows that our approach does not artificially dominate when the problem is effectively Euclidean since WGP becomes regular GP when $\sigma_{i1} = \sigma_1$ and $\sigma_{i2} = \sigma_2$. In contrast, in the anisotropic 2D-aniso-PC setting, methods based on Gaussian moments or mean embeddings deteriorate markedly (UI-GP and KME-GP RMSE ≈ 0.69, coverage 0.15), whereas WGP, PWA-GP, PCPWA, and MMD-GP all keep RMSE around 0.20 and coverage between 0.65 and 0.98. PCPWA in particular combines good accuracy (RMSE 0.2036, very close to the best 0.1972) with high coverage (0.95) and low CRPS, benefiting from aligning the kernel with principal directions of the empirical clouds.

The high-dimensional Ackley experiments further probe robustness under complex geometry. In 5D, Wasserstein kernels (especially PWA-GP and WGP) achieve lower RMSE and CRPS than Regular and UI-GP while maintaining higher coverage, and Aggregated GP again fails badly. In 10D, all methods become challenged, and MMD-GP attains the best RMSE/CRPS with reasonably high coverage; nevertheless, Wasserstein kernels remain very close in RMSE (e.g. PWA-GP and WGP within roughly 10% of the best) and retain the second best coverage.

Across all scenarios, sliced-WGP (SWGP) is competitive in 1D but its optimization is numerically unstable in several higher-dimensional cases, as indicated by the extremely large RMSE/CRPS values in Table 4. Overall, PWA-GP and PCPWA consistently deliver well-calibrated predictive uncertainty in EIV settings while retaining competitive RMSE, and they remain robust as the input distributions become heteroscedastic, skewed, anisotropic, and high-dimensional, even when they are not the single best-performing method in raw RMSE.

3.2 Calibration of particle accelerators

Particle accelerators play a central role in scientific discovery and industrial applications. In a typical machine, focusing elements such as quadrupole magnets guide the charged particle beam and preserve beam quality, measured by the transverse emittance at the end of the accelerator. Here we study the relationship between the strengths of five quadrupole magnets in a linac section and the resulting beam emittance.

For each of the $N = 500$ machine settings we observe a five-dimensional vector of magnet strengths and two emittances (horizontal and vertical). Each magnet is specified with a relative tolerance (5% or 0.5%) around its nominal strength Tao et al. (2017), so the "input" available to a regression model is a noisy proxy U_i of an unknown true setting X_i-a classical errors-in-variables situation. We therefore model each setting as

$$U_i \sim \mathcal{N}(\mu_i, \Sigma_i), \qquad \Sigma_i = \text{diag}(\mu_i \odot \varepsilon),$$

where $\mu_i \in \mathbb{R}^5$ is the magnet vector for shot i and $\varepsilon \in \{0.05, 0.005\}$ is the relative error. From each $\mathcal{N}(\mu_i, \Sigma_i)$ we draw a small cloud of samples to represent U_i, and define the scalar response $Y_i = $ (horizontal emittance) $\times$ (vertical emittance)$/10^{-12}$. We use an 80/20 train-test split.

We fit both Euclidean and distributional GP models. Euclidean baselines are standard GPs on empirical means $m_i = \mathbb{E}[U_i]$ with RBF, Matérn 3/2, Matérn 5/2 and exponential kernels. Distributional competitors are our Wasserstein GPs (PWA-GP, WGP, PCPWA) that act on the sample clouds using separable, full and PCA-based Wasserstein kernels, together with the uncertain-input GP UI-GP of Girard et al. (2002).

Table 3.2 reports the results for both 5% and 0.5% relative errors. All methods (except SWGP) nearly interpolate the training data (train RMSE $< 2 \times 10^{-2}$, not shown). On the test set, UI-GP consistently achieves the best accuracy and calibration: RMSE 0.1340 with coverage 0.86 and CRPS 0.0557 at 5% error, improving to RMSE 0.1294, coverage 0.92 and CRPS 0.0441 at 0.5% error-very close to the nominal 90% target. Euclidean GPs on means have larger RMSE (0.22-0.24 at 5%, 0.22-0.28 at 0.5%) and higher CRPS, but still decent coverage (0.91-0.98), reflecting that in this nearly Gaussian, low-dimensional setting a moment-based EIV correction already works well.

Our Wasserstein GPs (PWA-GP, WGP, PCPWA) are broadly comparable to the mean-based Euclidean GPs on this dataset: they achieve similar qualitative behavior and conservative coverage (0.96-0.98), but do not provide a systematic improvement in point accuracy or CRPS. This is consistent with the structure of the input uncertainty here, each U_i is close to Gaussian and its dispersion is largely determined by the nominal setting μ_i via a fixed relative tolerance. Therefore, much of the useful information is already captured by the empirical mean representation. The SWGP is numerically unstable for both error levels, with diverging RMSE and CRPS and zero coverage, and is therefore not a viable competitor on this dataset.

Overall, the accelerator results align with our simulations results in Table 4 (1D-EIV): when the input noise is close to Gaussian and the response depends mainly on low-order moments, mean-based and uncertain-input GPs can match (and in this case outperform) Wasserstein GPs, whereas the latter show clear benefits in the more non-Gaussian and anisotropic EIV scenarios of Section 3.1.

Method	5% error			0.5% error		
	RMSE	Cov.	CRPS	RMSE	Cov.	CRPS
RBF GP	0.2186	0.910	0.0883	0.2177	0.920	0.0806
Matérn 3/2 GP	0.2351	0.930	0.1010	0.2378	0.930	0.0997
Matérn 5/2 GP	0.2228	0.940	0.0912	0.2242	0.940	0.0892
Exp. GP	0.2432	0.980	0.1380	0.2847	0.980	0.1392
PWA-GP	0.2798	0.980	0.1690	0.3373	0.980	0.1681
SWGP	2.7234×10^6	0.000	2.1404×10^6	3.1142×10^6	0.000	3.9275×10^6
WGP	0.3169	0.970	0.1949	0.2806	0.970	0.1401
PCPWA	0.3072	0.960	0.1682	0.2828	0.970	0.1430
Uncertain-input GP	0.1340	0.860	0.0557	0.1294	0.920	0.0441

Table 1: Accelerator dataset: test RMSE, empirical coverage of nominal 90% predictive intervals (Cov.), and CRPS for two relative error levels. Euclidean GPs use empirical means as inputs; distributional GPs act on full input distributions.

3.3 Noisy Trajectories from NOAA Drifters

This experiment compares GP-RBF and WGP-RBF in predicting mean temperatures, emphasizing input uncertainty. We evaluate models using RMSE and CRPS metrics on NOAA's Global Drifter Program dataset, containing hourly drifter trajectories and sea surface temperatures (SST) observations (Elipot et al., 2016; 2022). While the SST measurements are collected over space at a high temporal frequency, scientific interest is often focused on the time-aggregated behavior of SSTs to understand large-scale modes of ocean variability (Philander, 1983; Mantua and Hare, 2002; Schlesinger and Ramankutty, 1994; Thompson and Wallace, 2000). Therefore, we consider these trajectories as distributional input models, where the distribution of inputs represents the collection of geospatial positions corresponding to the hourly measurements. The data is preprocessed by selecting trajectories with low temperature variance ($Y_{\mathrm{var\_temp}}$). Each collection of scattered points on a trajectory is then represented by a 2D normal distribution with empirical mean position and empirical covariance matrix. We have to emphasize that the trajectories of drifters (i.e., floating buoys on the oceanic surface) are highly noisy, self-intersecting and therefore usual functional data representation is not appropriate here. Therefore, we consider these trajectories as distributional input models, where both measurement noise and ocean variability-information GP-RBF ignores by using only point estimates. The response ($Y_{\mathrm{mean\_temp}}$) is the mean SST. We train on one year's data (e.g., 1999) and test on another year (e.g., 1998), initially focusing on 100 trajectories with lowest variance to ensure valid scalar responses.

Figure 2: RMSE (left) and CRPS (right) for GP-RBF vs. WGP-RBF across training/testing years 1987-2022. GP-RBF degrades when trained on recent and tested on older years, indicating poor generalization under temporal shifts. WGP-RBF remains robust, consistently achieving lower errors and better probabilistic calibration.

We then evaluated both models on every pair of training/testing years (1987-2022). The sliced WGP takes much higher memory and is not considered here. Figure 2 shows RMSE and CRPS heatmaps. GP-RBF suffers high errors off the diagonal-overfitting recent data and failing to extrapolate to earlier distributions. In contrast, WGP-RBF maintains low RMSE and CRPS across all year combinations, thanks to its explicit treatment of input uncertainty. The diagonal structure reflects optimal performance when training and testing distributions match, but WGP-RBF's stronger off-diagonal performance underscores its superior transferability for applications with nonstationary input variability, such as climate monitoring.

4 Discussion

We have argued that uncertainty quantification, rather than purely pointwise accuracy, should guide the design of GP models when inputs are measured with error. By lifting inputs to probability measures and endowing this space with Wasserstein geometry, PWA-based GPs provide a simple, deterministic alternative to latent-input and sampling-based EIV approaches. The theory now separates a deterministic net-extension bound from a model-based GP posterior statement: under Lipschitz and covering-number conditions, one obtains a uniform high-probability posterior band and a simple sufficient condition for conservative credible intervals. Our experiments confirm that these ideas are consistent with substantially better-calibrated predictive intervals on real data. Our empirical findings suggest that classical Euclidean kernels are often overly confident when covariates are noisy, whereas Wasserstein-type kernels and PWA in particular produce credible intervals that track the true variability across different noise regimes and non-Gaussian input distributions. This makes PWA-GP especially appealing in scientific and engineering applications, such as accelerator calibration and oceanographic monitoring, where decisions are driven by the reliability of uncertainty quantification.

Future work includes extending the analysis to high-dimensional, non-Gaussian noise (Tekriwal et al., 2026) and non-Euclidean input spaces (Luo et al., 2026), incorporating learned or problem-specific transport costs.

Scenario	Metric	Reg	Agg	WGP	SWGP	PWA	PCPWA	UI	KME	MMD
2D-mean	RMSE	8.500e-03	1.061e-01	5.280e-02	1.015e+07	4.620e-02	1.615e-01	4.000e-04	2.100e-03	2.600e-03
	Cov.	4.250e-01	1.000e-01	**1.000e+00**	0.000e+00	6.250e-01	**9.750e-01**	**1.000e+00**	**1.000e+00**	**9.750e-01**
	CRPS	5.100e-03	5.940e-02	4.540e-02	7.752e+06	3.970e-02	1.108e-01	2.000e-03	8.900e-03	1.010e-02
1D-EIV	RMSE	3.090e-01	5.712e-01	3.014e-01	2.995e-01	3.014e-01	3.014e-01	3.462e-01	4.179e-01	3.029e-01
	Cov.	6.500e-01	5.000e-02	**9.670e-01**	**9.670e-01**	**9.670e-01**	**9.670e-01**	5.670e-01	5.330e-01	6.670e-01
	CRPS	1.848e-01	4.791e-01	1.716e-01	1.722e-01	1.716e-01	1.716e-01	2.179e-01	2.844e-01	1.775e-01
1D-Var	RMSE	1.612e-01	4.810e-01	1.849e-01	1.819e-01	1.849e-01	1.849e-01	2.016e-01	2.697e-01	1.743e-01
	Cov.	7.670e-01	0.000e+00	**9.830e-01**	**9.830e-01**	**9.830e-01**	**9.830e-01**	6.330e-01	6.170e-01	7.670e-01
	CRPS	9.150e-02	3.648e-01	1.070e-01	1.058e-01	1.070e-01	1.070e-01	1.135e-01	1.552e-01	9.900e-02
1D-Skew	RMSE	1.768e-01	4.779e-01	1.629e-01	1.633e-01	1.629e-01	1.629e-01	1.762e-01	1.892e-01	1.406e-01
	Cov.	4.330e-01	0.000e+00	**9.500e-01**	**9.670e-01**	**9.500e-01**	**9.500e-01**	4.170e-01	4.500e-01	5.830e-01
	CRPS	1.075e-01	3.618e-01	9.730e-02	9.750e-02	9.730e-02	9.730e-02	1.057e-01	1.152e-01	8.000e-02
2D-aniso-PC	RMSE	1.972e-01	5.435e-01	2.086e-01	2.541e+06	2.130e-01	2.036e-01	6.893e-01	6.893e-01	2.009e-01
	Cov.	6.750e-01	2.500e-02	**9.750e-01**	0.000e+00	**9.500e-01**	**9.500e-01**	1.500e-01	1.500e-01	6.500e-01
	CRPS	1.086e-01	3.846e-01	1.252e-01	2.581e+06	1.211e-01	1.124e-01	5.528e-01	5.528e-01	1.098e-01
HD-Ackley-5D	RMSE	6.071e-01	2.483e+00	5.413e-01	4.080e+08	5.190e-01	5.562e-01	6.062e-01	6.574e-01	5.080e-01
	Cov.	7.750e-01	2.500e-02	9.000e-01	0.000e+00	8.870e-01	**9.630e-01**	7.750e-01	7.750e-01	7.870e-01
	CRPS	3.501e-01	1.879e+00	3.302e-01	3.081e+07	2.715e-01	3.147e-01	3.495e-01	3.607e-01	2.969e-01
HD-Ackley-10D	RMSE	5.925e-01	2.487e+00	5.873e-01	5.888e+00	6.003e-01	6.110e-01	5.962e-01	6.112e-01	5.450e-01
	Cov.	2.630e-01	7.500e-02	6.880e-01	0.000e+00	7.000e-01	7.620e-01	6.750e-01	6.250e-01	7.630e-01
	CRPS	4.138e-01	1.887e+00	3.492e-01	2.431e+00	3.567e-01	3.718e-01	3.470e-01	3.552e-01	3.309e-01

Table 2: Simulated distributional regression experiments. For each scenario we report test RMSE, empirical coverage of nominal 90% predictive intervals (Cov.), and CRPS. "Reg" = Regular GP on empirical means, "Agg" = Aggregated GP, "UI" = uncertain-input GP. Extremely large SWGP values indicate numerical divergence. Values of Cov. above 0.9 are shown in **bold**.

References

Arjovsky, M., Chintala, S., and Bottou, L. (2017). Wasserstein generative adversarial networks. In *International conference on machine learning*, pages 214–223. PMLR.

Arnold, S., Walz, E.-M., Ziegel, J., and Gneiting, T. (2024). Decompositions of the mean continuous ranked probability score. *Electronic Journal of Statistics*, 18(2):4992–5044.

Bachoc, F., Gamboa, F., Loubes, J.-M., and Venet, N. (2017). A gaussian process regression model for distribution inputs. *IEEE Transactions on Information Theory*, 64(10):6620–6637.

Bilionis, I., Zabaras, N., Konomi, B. A., and Lin, G. (2013). Multi-output separable Gaussian process: Towards an efficient, fully Bayesian paradigm for uncertainty quantification. *Journal of Computational Physics*, 241:212–239.

Bonet, C., Malézieux, B., Rakotomamonjy, A., Drumetz, L., Moreau, T., Kowalski, M., and Courty, N. (2023). Sliced-wasserstein on symmetric positive definite matrices for m/eeg signals. In *International Conference on Machine Learning*, pages 2777–2805. PMLR.

Candelieri, A., Ponti, A., and Archetti, F. (2022). Gaussian process regression over discrete probability measures: on the non-stationarity relation between euclidean and wasserstein squared exponential kernels. *arXiv preprint arXiv:2212.01310*.

Damianou, A. and Lawrence, N. D. (2013). Deep Gaussian processes. In *Artificial intelligence and statistics*, pages 207–215. PMLR.

Deisenroth, M. P. (2010). *Efficient reinforcement learning using Gaussian processes*, volume 9. KIT Scientific Publishing.

Delon, J., Desolneux, A., and Salmona, A. (2022). Gromov–wasserstein distances between gaussian distributions. *Journal of Applied Probability*, 59(4):1178–1198.

Dowson, D. and Landau, B. (1982). The fréchet distance between multivariate normal distributions. *Journal of multivariate analysis*, 12(3):450–455.

Elipot, S., Lumpkin, R., Perez, R. C., Lilly, J. M., Early, J. J., and Sykulski, A. M. (2016). A global surface drifter data set at hourly resolution. *Journal of Geophysical Research: Oceans*, 121(5):2937–2966.

Elipot, S., Sykulski, A., Lumpkin, R., Centurioni, L., and Pazos, M. (2022). A dataset of hourly sea surface temperature from drifting buoys. *Scientific data*, 9(1):567.

Girard, A., Rasmussen, C., Candela, J. Q., and Murray-Smith, R. (2002). Gaussian process priors with uncertain inputs application to multiple-step ahead time series forecasting. *Advances in neural information processing systems*, 15.

Horn, R. A. and Johnson, C. R. (2012). *Matrix analysis*. Cambridge university press.

Kennedy, M. C. and O'Hagan, A. (2001). Bayesian calibration of computer models. *Journal of the Royal Statistical Society: Series B*, 63(3):425–464.

Lederer, A., Umlauft, J., and Hirche, S. (2019). Uniform error bounds for gaussian process regression with application to safe control. *Advances in Neural Information Processing Systems*, 32.

Liu, Y., Sid-Lakhdar, W., Marques, O., Zhu, X., Meng, C., Demmel, J., and Li, X. (2021). GPTune: multitask learning for autotuning exascale applications. In *PPoPP '21: Proceedings of the 26th ACM SIGPLAN Symposium on Principles and Practic e of Parallel Programming*, pages 234–246, San Francisco, USA.

Luo, H., Horiguchi, A., and Ma, L. (2026). Wavelet tree ensembles for triangulable manifolds. *arXiv preprint arXiv:2601.20254*.

Mallasto, A. and Feragen, A. (2017). Learning from uncertain curves: The 2-wasserstein metric for gaussian processes. *Advances in Neural Information Processing Systems*, 30.

Mantua, N. J. and Hare, S. R. (2002). The pacific decadal oscillation. *Journal of Oceanography*, 58(1):35–44.

McHutchon, A. and Rasmussen, C. (2011). Gaussian process training with input noise. *Advances in neural information processing systems*, 24.

Meunier, D., Pontil, M., and Ciliberto, C. (2022). Distribution regression with sliced wasserstein kernels. In *Proceedings of the 39th International Conference on Machine Learning*, volume 162 of *Proceedings of Machine Learning Research*, pages 15501–15523. PMLR.

Moreno-Muñoz, P., Artés-Rodríguez, A., and Álvarez, M. A. (2018). Heterogeneous multi-output gaussian process prediction. In *Advances in Neural Information Processing Systems*, volume 31.

Muandet, K., Fukumizu, K., Sriperumbudur, B., and Schölkopf, B. (2017). Kernel mean embedding of distributions: A review and beyond. *Foundations and Trends in Machine Learning*, 10(1–2):1–141.

Noack, M. M. et al. (2019). A kriging-based approach to autonomous experimentation with applications to X-Ray scattering. *Sci. Rep.*, 9:11809.

Noack, M. M., Zwart, P. H., Ushizima, D. M., Fukuto, M., Yager, K. G., Elbert, K. C., Murray, C. B., Stein, A., Doerk, G. S., Tsai, E. H., et al. (2021). Gaussian processes for autonomous data acquisition at large-scale synchrotron and neutron facilities. *Nature Reviews Physics*, 3(10):685–697.

Panaretos, V. M. and Zemel, Y. (2019). Statistical aspects of wasserstein distances. *Annual Review of Statistics and Its Application*, 6(1):405–431.

Panaretos, V. M. and Zemel, Y. (2020). *An invitation to statistics in Wasserstein space*. Springer Nature.

Peyré, G., Cuturi, M., et al. (2019). Computational optimal transport: With applications to data science. *Foundations and Trends® in Machine Learning*, 11(5-6):355–607.

Philander, S. G. H. (1983). El niño southern oscillation phenomena. *Nature*, 302(5906):295–301.

Qiang, J., Kan, Y., Li, X., Liu, Y., X.Gu, Fang, W., and Hao, Y. (2024). Advanced modeling and optimization of nuclear physics colliders. In *Proc. IPAC'24*, number 15 in IPAC'24 - 15th International Particle Accelerator Conference, pages 3194–3197. JACoW Publishing, Geneva, Switzerland.

Schlesinger, M. E. and Ramankutty, N. (1994). An oscillation in the global climate system of period 65-70 years. *Nature*, 367(6465):723–726.

Song, D. and Han, K. (2023). Errors-in-variables fréchet regression with low-rank covariate approximation. In *Advances in Neural Information Processing Systems*, volume 36, pages 80575–80607.

Stach, E., DeCost, B., Kusne, A. G., Hattrick-Simpers, J., Brown, K. A., Reyes, K. G., Schrier, J., Billinge, S., Buonassisi, T., Foster, I., et al. (2021). Autonomous experimentation systems for materials development: A community perspective. *Matter*, 4(9):2702–2726.

Szabó, Z., Sriperumbudur, B. K., Póczos, B., and Gretton, A. (2016). Learning theory for distribution regression. *Journal of Machine Learning Research*, 17(152):1–40.

Tao, Y., Qiang, J., and Hwang, K. (2017). Overtaking collision effects in a cw double-pass proton linac. *Physical Review Accelerators and Beams*, 20(12):124202.

Tekriwal, V., Risser, M. D., Luo, H., and Noack, M. M. (2026). Ggmps: Generalized gaussian mixture processes.

Thomas, J. C., Rossi, A., Smalley, D., Francaviglia, L., Yu, Z., Zhang, T., Kumari, S., Robinson, J. A., Terrones, M., Ishigami, M., et al. (2022). Autonomous scanning probe microscopy investigations over ws2 and au {111}. *npj Computational Materials*, 8(1):99.

Thompson, D. W. J. and Wallace, J. M. (2000). Annular modes in the extratropical circulation. part i: Month-to-month variability*. *Journal of Climate*, 13(5):1000–1016.

Tuo, R. and Wang, W. (2022). Uncertainty quantification for bayesian optimization. In *International Conference on Artificial Intelligence and Statistics*, pages 2862–2884. PMLR.

Vinogradska, J., Bischoff, B., Nguyen-Tuong, D., Romer, A., Schmidt, H., and Peters, J. (2016). Stability of controllers for gaussian process forward models. In *International Conference on Machine Learning*, pages 545–554. PMLR.

Wang, W. (2021). On the inference of applying gaussian process modeling to a deterministic function. *Electronic Journal of Statistics*, 15(2):5014–5066.

Weed, J. and Bach, F. (2019). Sharp asymptotic and finite-sample rates of convergence of empirical measures in wasserstein distance. *Bernoulli*, 25(4A):2620–2648.

Williams, C. K. I. and Rasmussen, C. E. (2006). *Gaussian Processes for Machine Learning*. MIT Press.

Wilson, A. and Nickisch, H. (2015). Kernel interpolation for scalable structured gaussian processes (kiss-gp). In *International Conference on Machine Learning*, pages 1775–1784. PMLR.

Zhou, S., Pati, D., Wang, T., Yang, Y., and Carroll, R. J. (2023). Gaussian processes with errors in variables: Theory and computation. *Journal of Machine Learning Research*, 24(87):1–53.

A Related Tables

B Evaluation metrics

We report three complementary metrics for predictive performance. Let $\widehat{m}_i$ and $\widehat{s}_i$ denote the posterior mean and posterior standard deviation for the ith test input, and let y_i be the corresponding test response.

First, the root mean squared error (RMSE) measures point-prediction accuracy:

$$\mathrm{RMSE} = \left(\frac{1}{n_{\text{test}}} \sum_{i=1}^{n_{\text{test}}} (\widehat{m}_i - y_i)^2 \right)^{1/2}.$$

Second, for a nominal $100(1 - \alpha)\%$ predictive interval

$$C_i = \left[\widehat{m}_i - z_{1-\alpha/2}\widehat{s}_i, \ \widehat{m}_i + z_{1-\alpha/2}\widehat{s}_i \right],$$

the empirical coverage is

$$\mathrm{Cov}_{1-\alpha} = \frac{1}{n_{\text{test}}} \sum_{i=1}^{n_{\text{test}}} \mathbf{1}\{y_i \in C_i\}.$$

Coverage below the nominal level is called *undercoverage* and indicates that the predictive intervals are too narrow, i.e., the model is overconfident.

Third, we report the continuous ranked probability score (CRPS), a proper scoring rule for the full predictive distribution:

$$\mathrm{CRPS}(F_i, y_i) = \int_{-\infty}^{\infty} \left(F_i(t) - \mathbf{1}\{y_i \leq t\} \right)^2 dt,$$

where F_i is the predictive cumulative distribution function for the ith test point. Lower CRPS is better; it rewards both calibration and sharpness.

C Proof for Proposition 1

Proof. With $f(x) = c + w^\top x$, we have

$$Y = f(X + \varepsilon_X) + \varepsilon = f(X) + w^\top \varepsilon_X + \varepsilon.$$

Thus $f(X) - Y = -(w^\top \varepsilon_X + \varepsilon)$ and conditional on X,

$$w^\top \varepsilon_X + \varepsilon \sim \mathcal{N}(0, \ w^\top \Sigma_X w + \sigma^2).$$

Therefore,

$$\mathbb{P}(f(X) \in I_\alpha \mid X) = \mathbb{P}\left(|w^\top \varepsilon_X + \varepsilon| \leq z_{1-\alpha/2}\sigma \right) = 2\Phi\left(\frac{z_{1-\alpha/2}\sigma}{\sqrt{\sigma^2 + w^\top \Sigma_X w}} \right) - 1,$$

which is strictly less than $1 - \alpha$ whenever $w^\top \Sigma_X w > 0$. $\qquad\square$

D Proof for Theorem 5

Proof. We first verify the Lipschitz continuity of ν_N. Since

$$\nu_N(\mu) = \sum_{i=1}^{N} \alpha_i k(\mu, \mu_i),$$

and k is L_k-Lipschitz in its first argument with respect to W_1,

$$|\nu_N(\mu) - \nu_N(\mu')| \le \sum_{i=1}^{N} |\alpha_i| \, |k(\mu, \mu_i) - k(\mu', \mu_i)| \le L_k \Big(\sum_{i=1}^{N} |\alpha_i|\Big) W_1(\mu, \mu').$$

Because $\alpha = (K + \sigma_*^2 I_N)^{-1} y$ and $K + \sigma_*^2 I_N$ is invertible, $\|\alpha\|_\infty < \infty$, hence

$$\sum_{i=1}^{N} |\alpha_i| \le N \|\alpha\|_\infty \quad \text{and therefore} \quad |\nu_N(\mu) - \nu_N(\mu')| \le L_{\nu_N} W_1(\mu, \mu').$$

Now let $k_\mu = (k(\mu, \mu_1), \ldots, k(\mu, \mu_N))^\top$ and set $A = (K + \sigma_*^2 I_N)^{-1}$. Then

$$\sigma_N^2(\mu) = k(\mu, \mu) - k_\mu^\top A k_\mu.$$

By assumption, σ_N has modulus of continuity ω_{σ_N}, i.e.

$$|\sigma_N(\mu) - \sigma_N(\mu')| \le \omega_{\sigma_N}(\tau) \qquad \text{whenever } W_1(\mu, \mu') \le \tau.$$

Let $\{\bar{\mu}_1, \ldots, \bar{\mu}_M\}$ be a τ-net of $\mathcal{P}$. Fix any $\mu \in \mathcal{P}$ and choose j such that $W_1(\mu, \bar{\mu}_j) \le \tau$. Then

$$|f(\mu) - \nu_N(\mu)| \le |f(\mu) - f(\bar{\mu}_j)| + |f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)| + |\nu_N(\bar{\mu}_j) - \nu_N(\mu)|.$$

The first and third terms are bounded by Lipschitz continuity:

$$|f(\mu) - f(\bar{\mu}_j)| \le L_f \tau, \qquad |\nu_N(\bar{\mu}_j) - \nu_N(\mu)| \le L_{\nu_N} \tau.$$

By hypothesis,

$$|f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)| \le B \, \sigma_N(\bar{\mu}_j) \le B \, [\sigma_N(\mu) + \omega_{\sigma_N}(\tau)].$$

Combining the three bounds yields

$$|f(\mu) - \nu_N(\mu)| \le B \, \sigma_N(\mu) + (L_f + L_{\nu_N})\tau + B \, \omega_{\sigma_N}(\tau),$$

as claimed. $\qquad \square$

Proof of Corollary 6. Let $\{\bar{\mu}_1, \ldots, \bar{\mu}_M\}$ be a τ-net of $\mathcal{P}$ in W_1, where $M = M(\tau, \mathcal{P})$. Under the well-specified GP model, conditional on $\mathcal{D}_N$, for each $j = 1, \ldots, M$,

$$f(\bar{\mu}_j) \mid \mathcal{D}_N \sim \mathcal{N}\big(\nu_N(\bar{\mu}_j), \sigma_N^2(\bar{\mu}_j)\big).$$

Fix j. If $\sigma_N(\bar{\mu}_j) > 0$, then

$$Z_j := \frac{f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)}{\sigma_N(\bar{\mu}_j)} \; \mathcal{D}_N \sim \mathcal{N}(0, 1),$$

and therefore, by the definition

$$\beta(\tau) = \left[\Phi^{-1}\Big(1 - \frac{\delta}{2M(\tau, \mathcal{P})}\Big)\right]^2,$$

we have

$$\mathbb{P}\Big(|f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)| > \sqrt{\beta(\tau)} \, \sigma_N(\bar{\mu}_j) \,\Big|\, \mathcal{D}_N\Big) = \mathbb{P}\Big(|Z_j| > \Phi^{-1}\Big(1 - \frac{\delta}{2M(\tau, \mathcal{P})}\Big)\Big)$$

$$= \frac{\delta}{M(\tau, \mathcal{P})}.$$

If instead $\sigma_N(\bar{\mu}_j) = 0$, then the posterior law is degenerate and $f(\bar{\mu}_j) = \nu_N(\bar{\mu}_j)$ almost surely given $\mathcal{D}_N$, so the same inequality holds with probability $0 \le \delta/M(\tau, \mathcal{P})$.

Hence, for every $j = 1, \ldots, M$,

$$\mathbb{P}\Big(|f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)| > \sqrt{\beta(\tau)}\,\sigma_N(\bar{\mu}_j)\,\Big|\,\mathcal{D}_N\Big) \leq \frac{\delta}{M(\tau, \mathcal{P})}.$$

Applying the union bound over the $M(\tau, \mathcal{P})$ net points yields

$$\mathbb{P}\Big(|f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)| \leq \sqrt{\beta(\tau)}\,\sigma_N(\bar{\mu}_j) \text{ for all } j = 1, \ldots, M\,\Big|\,\mathcal{D}_N\Big) \geq 1 - \delta.$$

On this event, Theorem 5 applies with $B = \sqrt{\beta(\tau)}$, giving

$$|f(\mu) - \nu_N(\mu)| \leq \sqrt{\beta(\tau)}\,\sigma_N(\mu) + (L_f + L_{\nu_N})\tau + \sqrt{\beta(\tau)}\,\omega_{\sigma_N}(\tau) \qquad \text{for all } \mu \in \mathcal{P}.$$

Since

$$\gamma(\tau) = (L_f + L_{\nu_N})\tau + \sqrt{\beta(\tau)}\,\omega_{\sigma_N}(\tau),$$

the claimed bound follows. □

Proof of Corollary 7. Let $\{\bar{\mu}_1, \ldots, \bar{\mu}_M\}$ be a τ-net of $(\mathcal{P}, D)$, where $M = M_{\text{PCPWA}}(\tau, \mathcal{P})$. Under the well-specified GP model with prior $f \sim GP(0, k_{\text{PCPWA}})$, conditional on $\mathcal{D}_N$, for each $j = 1, \ldots, M$,

$$f(\bar{\mu}_j) \mid \mathcal{D}_N \sim \mathcal{N}\big(\nu_N(\bar{\mu}_j),\, \sigma_N^2(\bar{\mu}_j)\big).$$

Define

$$\beta_{\text{PCPWA}}(\tau) = \left[\Phi^{-1}\left(1 - \frac{\delta}{2M_{\text{PCPWA}}(\tau, \mathcal{P})}\right)\right]^2.$$

Arguing exactly as in the proof of Corollary 6, for each net point $\bar{\mu}_j$,

$$\mathbb{P}\Big(|f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)| > \sqrt{\beta_{\text{PCPWA}}(\tau)}\,\sigma_N(\bar{\mu}_j)\,\Big|\,\mathcal{D}_N\Big) \leq \frac{\delta}{M_{\text{PCPWA}}(\tau, \mathcal{P})}.$$

A union bound therefore gives, conditional on $\mathcal{D}_N$,

$$|f(\bar{\mu}_j) - \nu_N(\bar{\mu}_j)| \leq \sqrt{\beta_{\text{PCPWA}}(\tau)}\,\sigma_N(\bar{\mu}_j), \qquad j = 1, \ldots, M,$$

with posterior probability at least $1 - \delta$.

Now apply Theorem 5 with the pseudo-metric D in place of W_1, using the assumed D-Lipschitz bounds for f and k_{PCPWA} and the modulus of continuity ω_{σ_N} with respect to D, with

$$B = \sqrt{\beta_{\text{PCPWA}}(\tau)}.$$

This yields

$$|f(\mu) - \nu_N(\mu)| \leq \sqrt{\beta_{\text{PCPWA}}(\tau)}\,\sigma_N(\mu) + (L_f + L_{\nu_N})\tau + \sqrt{\beta_{\text{PCPWA}}(\tau)}\,\omega_{\sigma_N}(\tau) \qquad \text{for all } \mu \in \mathcal{P},$$

with posterior probability at least $1 - \delta$. Writing

$$\gamma_{\text{PCPWA}}(\tau) = (L_f + L_{\nu_N})\tau + \sqrt{\beta_{\text{PCPWA}}(\tau)}\,\omega_{\sigma_N}(\tau)$$

completes the proof. □

E   Proof for Proposition 9

Proof. Fix $u \in \mathbb{S}^{d-1}$. Let $\mathcal{E}_u$ denote the posterior event that the projected uniform bound holds:

$$|f(\mu) - \nu_N^u(\mu)| \leq \sqrt{\beta_u(\tau)}\, \sigma_N^u(\mu) + \gamma_u(\tau) \qquad \text{for all } \mu \in \mathcal{P}.$$

By assumption, $\mathbb{P}(\mathcal{E}_u \mid \mathcal{D}_N) \geq 1 - \delta$.

On $\mathcal{E}_u$, evaluating at $\mu = \mu_X$ gives

$$|f(\mu_X) - \nu_N^u(\mu_X)| \leq \sqrt{\beta_u(\tau)}\, \sigma_N^u(\mu_X) + \gamma_u(\tau).$$

Assume $z > \sqrt{\beta_u(\tau)}$ and $\sigma_N^u(\mu_X) \geq \gamma_u(\tau)/(z - \sqrt{\beta_u(\tau)})$. Then

$$\sqrt{\beta_u(\tau)}\, \sigma_N^u(\mu_X) + \gamma_u(\tau) \leq z\, \sigma_N^u(\mu_X),$$

so $|f(\mu_X) - \nu_N^u(\mu_X)| \leq z\, \sigma_N^u(\mu_X)$, which is equivalent to $f(\mu_X) \in I_\alpha^u(\mu_X)$. Therefore,

$$\mathbb{P}(f(\mu_X) \in I_\alpha^u(\mu_X) \mid \mathcal{D}_N) \geq \mathbb{P}(\mathcal{E}_u \mid \mathcal{D}_N) \geq 1 - \delta,$$

which proves the claim. $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\quad\square$

F   Bounds for derivatives of log likelihoods

Here we derive bounds of derivatives for k_{GW_p} as below:

$$k_{GW_p}(\mu, \nu) = \lambda \exp(-\sigma GW_p(\mu, \nu)^2), \tag{8}$$

Lemma 10. *(Bounds for derivatives of log likelihoods with GW) For $p = 2$ and μ, ν are both multivariate Gaussian distributions defined on $\mathbb{R}^d, \mathbb{R}^m$, the bounds for derivatives with respect to the kernel hyper-parameters λ, σ are in closed form as functions of $GW_p(\mu, \nu)^2$.*

We want to derive the $\frac{\partial \boldsymbol{K}}{\partial \lambda}$ and $\frac{\partial \boldsymbol{K}}{\partial \sigma}$ and plug it back respectively. We want to use Theorem 3.1, 4.1 and Proposition 4.2 in Delon et al. (2022) to obtain a lower bound for these derivatives as a closed form surrogate to the exact. Suppose without any loss of generality that $d \leq m$. Let $\mu = N(\mu_0, \Sigma_0)$ on $\mathbb{R}^m$ and $\nu = N(\mu_1, \Sigma_1)$ on $\mathbb{R}^d$ be two Gaussian measures. Let P_0, D_0 and P_1, D_1 be the respective diagonalizations of $\Sigma_0 = P_0 D_0 P_0^T \in \mathbb{R}^{m \times m}$ and $\Sigma_1 = P_1 D_1 P_1^T \in \mathbb{R}^{m \times m}$ which sort eigenvalues in decreasing order. We suppose that Σ_0 is non-singular. Then the $GW_2(\mu, \nu)$ is bounded by

$$LGW_2(\mu, \nu)^2 \leq GW_2(\mu, \nu)^2 \leq GGW_2(\mu, \nu)^2, \tag{9}$$

where the lower and upper bounds have the following closed form expressions:

$$LGW_2(\mu, \nu)^2 = 4\left(\operatorname{tr}(D_0) - \operatorname{tr}(D_1)\right)^2 + 4\left(\|D_0\|_F - \|D_1\|_F\right)^2 + 4\|D_0^{(d)} - D_1\|_F^2 \tag{10}$$
$$+ 4\left(\|D_0\|_F - \|D_0^{(d)}\|_F\right)^2.$$

$$GGW_2(\mu, \nu)^2 = 4\left(\operatorname{tr}(D_0) - \operatorname{tr}(D_1)\right)^2 + 8\left(\|D_0^{(d)} - D_1\|_F^2\right) + 8\left(\|D_0\|_F^2 - \|D_0^{(d)}\|_F^2\right). \tag{11}$$

and the we use the notation $D^{(d)}$ to denote the $d \times d$ submatrix containing the first d rows and the first d columns of any matrix D; $\|\cdot\|_F$ is the matrix Frobenius norm. Recall that we can set $x \sim \mu = N(\mu_0, \Sigma_0)$ and $x' \sim \nu = N(\mu_1, \Sigma_1)$, then we have:

$$\frac{\partial \boldsymbol{K}}{\partial \lambda} = \left\{\exp(-\sigma GW_2(x, x')^2)\right\}_{x, x'},$$
$$\frac{\partial \boldsymbol{K}}{\partial \sigma} = \left\{-\lambda GW_2(x, x')^2 \exp(-\sigma GW_2(x, x')^2)\right\}_{x, x'}.$$

These expressions admit the bound (9) due to the fact that exponential function is strictly monotonic over the real line. This is again a computational tool which can be used in optimization algorithms to find the MLEs of λ and σ. Instead of putting nested loops to solve the optimal transportation problem in the definition of GW_2, we only optimize the $LGW_2(\mu, \nu)^2$ and $GGW_2(\mu, \nu)^2$ (in closed form) to simplify the fitting procedure. However, in practice it usually suffices to maximize $LGW_2(\mu, \nu)^2$. At the same time, the (9) can be used to verify the Assumption 3 in Zhou et al. (2023) when the P_0, D_0 and P_1, D_1 are parameterized by a smooth parameter (i.e., as a smooth matrix-valued function of a kernel parameter).

G Computational complexity of competing methods

We briefly compare the leading-order computational complexity of the GP methods considered in this work. For all models, once a kernel matrix $K \in \mathbb{R}^{N \times N}$ has been assembled, exact GP training via Cholesky decomposition costs $\mathcal{O}(N^3)$, and prediction for $N_\star$ test inputs costs $\mathcal{O}(N^2 + NN_\star)$ (or $\mathcal{O}(NN_\star)$ once the Cholesky factor is cached). The main differences between methods therefore arise from (i) the number of *effective* training points they use and (ii) the cost of computing each kernel entry from the underlying clouds U_i.

Throughout, $U_i = \{u_{ij}\}_{j=1}^{m_i} \subset \mathbb{R}^d$ denotes the empirical input distribution for training example i, and $M = \max_i m_i$. We assume $m_i \asymp M$ for simplicity.

Regular GP on means. The regular GP baseline (Williams and Rasmussen, 2006) treats each distribution as a single point in $\mathbb{R}^d$ via its empirical mean $\bar{u}_i = m_i^{-1} \sum_{j=1}^{m_i} u_{ij}$. Computing all means costs $\mathcal{O}(NMd)$ once, and a kernel evaluation between two means is $\mathcal{O}(d)$. Building K therefore costs $\mathcal{O}(N^2 d)$, after which GP training is $\mathcal{O}(N^3)$. This is the reference complexity for a standard Euclidean GP with N inputs of dimension d.

Aggregated GP on raw samples. The aggregated GP baseline (an ensemble of standard GPs (Williams and Rasmussen, 2006)) treats each sample u_{ij} as an independent input. The effective number of training points becomes $\tilde{N} = \sum_i m_i \asymp NM$. Kernel evaluations remain $\mathcal{O}(d)$, but the kernel matrix is now of size $\tilde{N} \times \tilde{N}$, so training scales as

$$\mathcal{O}(\tilde{N}^3) = \mathcal{O}(N^3 M^3),$$

with kernel assembly cost $\mathcal{O}(N^2 M^2 d)$. This cubic dependence on M makes the aggregated GP rapidly infeasible as the number of samples per distribution grows.

Uncertain-input GP. The uncertain-input GP of Girard et al. (2002) (see also McHutchon and Rasmussen (2011)) replaces each U_i by a Gaussian $\mathcal{N}(\mu_i, \Sigma_i)$, where μ_i and Σ_i are the empirical mean and covariance. Estimating these parameters costs $\mathcal{O}(NMd^2)$ in general (or $\mathcal{O}(NMd)$ for diagonal covariances). The kernel between two Gaussians has a closed form involving $(\Sigma_i + \Sigma_j)$ and $(\mu_i - \mu_j)$; with full covariances this requires an inversion and determinant per pair, for a cost of $\mathcal{O}(d^3)$ per kernel entry and $\mathcal{O}(N^2 d^3)$ to assemble K. For diagonal or low-rank covariances the per-pair cost reduces to $\mathcal{O}(d)$ or $\mathcal{O}(kd)$, giving $\mathcal{O}(N^2 d)$ kernel assembly and again $\mathcal{O}(N^3)$ training.

Full Wasserstein GP. Wasserstein GPs build kernels from optimal-transport distances between empirical distributions (Mallasto and Feragen, 2017; Bachoc et al., 2017; Panaretos and Zemel, 2019; Candelieri et al., 2022). Computing the pairwise cost matrix between U_i and U_j is $\mathcal{O}(M^2 d)$, and solving the OT problem (e.g. with Sinkhorn or network simplex) requires at least $\mathcal{O}(M^2)$ per iteration and $\mathcal{O}(M^2)$ memory (Peyré et al., 2019). In practice this yields a per-pair distance cost of order $\mathcal{O}(M^2 d)$ to $\mathcal{O}(M^3)$, and thus a total cost of

$$\mathcal{O}(N^2 M^2 d) \quad \text{to} \quad \mathcal{O}(N^2 M^3)$$

for computing K, plus the usual $\mathcal{O}(N^3)$ for GP training. This quadratic (or worse) dependence on M motivates the separable and sliced variants below.

PWA-GP (separable Wasserstein kernel). Our PWA-GP uses a separable kernel built from one-dimensional Wasserstein distances along each coordinate, still rooted in the OT geometry of Panaretos and Zemel (2019). In 1D, the W_2 distance between empirical distributions of size M reduces to sorting, which costs $\mathcal{O}(M \log M)$ per marginal. For d dimensions, a naive implementation therefore costs $\mathcal{O}(dM \log M)$ per pair and $\mathcal{O}(N^2 dM \log M)$ in total to assemble K, which is much cheaper than full OT as soon as M is moderately large. Training again costs $\mathcal{O}(N^3)$.

PCPWA (PCA-based separable Wasserstein kernel). PCPWA first computes a low-rank PCA basis for the pooled samples and then applies the separable 1D Wasserstein construction along the leading k principal directions. The up-front PCA on all samples costs $\mathcal{O}(NMd^2)$ (or less with randomized SVD), but this is a one-time cost. Thereafter, the per-pair distance uses only k one-dimensional marginals, for a cost $\mathcal{O}(kM \log M)$ per pair and $\mathcal{O}(N^2 kM \log M)$ in total. Since typically $k \ll d$, PCPWA can be substantially cheaper than PWA-GP in high dimensions while also adapting to the intrinsic low-dimensional structure of the clouds.

Sliced Wasserstein GP. The sliced Wasserstein GP (Meunier et al., 2022) draws R random projection directions $\{\theta_r\}_{r=1}^{R} \subset \mathbb{S}^{d-1}$ and approximates the Wasserstein distance by averaging 1D Wasserstein distances along these projections. Projecting all samples in U_i onto θ_r costs $\mathcal{O}(Md)$ per direction and $\mathcal{O}(RMd)$ in total per cloud. These projections can be precomputed and reused across pairs. Given the projected samples, computing 1D distances between U_i and U_j across all R directions costs $\mathcal{O}(RM \log M)$ per pair. The overall kernel assembly cost is therefore

$$\mathcal{O}(NRMd) \quad \text{(precomputation)} \; + \; \mathcal{O}(N^2 RM \log M) \quad \text{(pairwise distances)},$$

followed by $\mathcal{O}(N^3)$ training. In our code R is treated as a moderate constant, so the dominant dependence is $\mathcal{O}(N^2 M \log M)$.

KME and MMD GPs. The KME and MMD kernels embed each empirical distribution into an RKHS using sample averages of feature maps (Muandet et al., 2017; Szabó et al., 2016). With a Gaussian base kernel and empirical distributions of size M, the kernel between U_i and U_j typically uses a double sum over samples,

$$k_{\text{KME}}(U_i, U_j) \; \approx \; \frac{1}{M^2} \sum_{a=1}^{M} \sum_{b=1}^{M} \exp\Big(-\frac{\|u_{ia} - u_{jb}\|^2}{2\ell^2}\Big),$$

and similarly for the squared MMD. Each kernel evaluation is therefore $\mathcal{O}(M^2 d)$, and assembling K costs $\mathcal{O}(N^2 M^2 d)$, again followed by $\mathcal{O}(N^3)$ training. Random features can reduce this to $\mathcal{O}(N^2 Dd)$ with D features, but in our experiments we use the exact empirical embedding.

Low-rank Fréchet EIV regression. The low-rank Fréchet errors-in-variables (EIV) regression of Song and Han (2023) represents each distributional covariate via a rank-r approximation in a suitable Hilbert space. Constructing these low-rank features from M samples per distribution costs roughly $\mathcal{O}(NMr)$, after which Fréchet regression (or GP regression on the r-dimensional coefficients) operates on N inputs of dimension r. Kernel assembly is therefore $\mathcal{O}(N^2 r)$, comparable to a regular GP in r dimensions, with the usual $\mathcal{O}(N^3)$ training cost. When $r \ll d$ and M is moderate, this yields a competitive complexity to our PCA-based PCPWA construction.

Table G summarizes the leading-order complexity of the main models as a function of the number of clouds N, samples per cloud M, and dimension d (ignoring the shared $\mathcal{O}(N^3)$ GP training cost). Regular and uncertain-input GPs (Williams and Rasmussen, 2006; Girard et al., 2002; McHutchon and Rasmussen, 2011) are cheapest in terms of M, since they compress each distribution to a finite-dimensional summary. Aggregated GP is prohibitively expensive as soon as M grows. Among fully distributional methods, our separable Wasserstein kernels (PWA-GP, PCPWA), together with SWGP (Meunier et al., 2022), enjoy *linear* or near-linear dependence on M (up to logarithmic factors) thanks to one-dimensional Wasserstein computations (Panaretos and Zemel, 2019; Peyré et al., 2019). In contrast, full WGP, KME, and MMD kernels (Mallasto and Feragen, 2017; Bachoc et al., 2017; Muandet et al., 2017; Szabó et al., 2016) scale at least quadratically in M. This

helps explain the empirical observation in our simulated and accelerator experiments that PWA-GP and PCPWA provide competitive errors-in-variables performance and well-calibrated uncertainty while remaining computationally tractable.

Method (paper name)	Kernel input object	Dist.	Latent X	Det.	EIV mechanism	Citations
Regular GP (RBF / Matérn / Exp)	Point estimate (empirical mean m_i)	N	N	Y	Ignores input uncertainty (variance / shape); standard baseline in Sec. 3 and Table 1.	Williams and Rasmussen (2006)
Aggregated Regular GPs	Noisy samples $\{U_{ij}\}$ as point inputs (fit per-j GP; average)	N	N	Y	Ensemble over replicate-specific GPs; can amplify variance in sparse regions (as observed).	**This paper.**
IV-function / cov-only encoding	Gaussian summary (μ_x, Σ_x) but only Σ_x used in kernel	Y*	N	Y	Efficient for locally Gaussian perturbations; cannot represent skew / multimodality / complex support.	Moreno-Muñoz et al. (2018)
WGP (Eq. (5), WGP-RBF)	Measure μ	Y	N	Y	OT-based kernel on distributions; PD only in special cases (e.g., Gaussian $p=2$, or 1D).	Candelieri et al. (2022); Peyré et al. (2019)
sliced WGP (Eq. (6), SWGP)	Measure μ; average W_p over random 1D projections u	Y	N	MC	PD preserved via expectation over projections; stochastic projections add Monte Carlo variability and cost.	Meunier et al. (2022); Bonet et al. (2023)
PWA-GP (Eq. (7))	Coordinate-wise 1D marginals $\{\mu_i\}_{i=1}^d$	Y	N	Y	Product of 1D Wasserstein kernels with dimension-specific hyperparameters (ARD-like); PD via product closure.	**This paper.**
PCPWA (Eq. (8))	Fixed directions $\{v_r\}_{r=1}^m$ (e.g., PCs); 1D pushforwards μ^{v_r}	Y	N	Y	Same idea as PWA but along a fixed orthonormal basis (often PCA); trades expressivity vs. cost via m.	**This paper.**
Latent-input calibration GP	Noisy U with latent true X (integrate out)	N	Y	(inf.)	Classic latent-variable EIV route; requires integration / sampling / approximations.	Kennedy and O'Hagan (2001)
Uncertain-input GP	Input distribution (often Gaussian) used in prediction / propagation	Y	(approx)	Y	Uses uncertain regressors (often Gaussian) in forecasting; typically relies on approximations/closed forms.	Girard et al. (2002)
GP training w/ input noise	Noisy inputs + local approximation during training	N	(approx)	Y	Convolves/approximates input noise effect during training (e.g., local linearization).	McHutchon and Rasmussen (2011)
Deep GP (latent-variable)	Joint latent representation for inputs/outputs	N	Y	(inf.)	Highly flexible latent EIV modeling; inference heavier / approximate.	Damianou and Lawrence (2013)
Low-rank Fréchet EIV regression	Noisy covariates w/ low-rank approximation	N	N	Y	Non-GP competitor class mentioned (not compared in experiments); linear/low-rank structure.	Song and Han (2023)
KME / MMD distribution re-	Empirical samples per input	Y	N	Y	Strong non-OT distribution-input baseline; geometry	Szabó et al. (2016); Muandet

Method	Effective # inputs	Kernel assembly cost (big-O)
Regular GP on means (Williams and Rasmussen, 2006)	N	$\mathcal{O}(N^2 d)$
Aggregated GP on samples (Williams and Rasmussen, 2006)	NM	$\mathcal{O}(N^2 M^2 d)$ (matrix) $+ \mathcal{O}(N^3 M^3)$ (training)
Uncertain-input GP (Girard et al., 2002; McHutchon and Rasmussen, 2011)	N	$\mathcal{O}(N^2 d^3)$ (full cov.) or $\mathcal{O}(N^2 d)$ (diag)
Full WGP (Mallasto and Feragen, 2017; Bachoc et al., 2017; Candelieri et al., 2022)	N	$\mathcal{O}(N^2 M^2 d)$ to $\mathcal{O}(N^2 M^3)$
PWA-GP (separable Wasserstein)	N	$\mathcal{O}(N^2 dM \log M)$
PCPWA (PCA-based PWA)	N	$\mathcal{O}(NMd^2)$ (PCA) $+ \mathcal{O}(N^2 kM \log M)$
SWGP (sliced Wasserstein) (Meunier et al., 2022)	N	$\mathcal{O}(NRMd) + \mathcal{O}(N^2 RM \log M)$
KME-GP (Muandet et al., 2017)	N	$\mathcal{O}(N^2 M^2 d)$
MMD-GP (Szabó et al., 2016)	N	$\mathcal{O}(N^2 M^2 d)$
Low-rank Fréchet EIV regression (Song and Han, 2023)	N	$\mathcal{O}(NMr) + \mathcal{O}(N^2 r)$

Table G.1: Leading-order cost of assembling the $N \times N$ kernel matrix for the different methods, assuming $m_i \asymp M$ samples per cloud. All methods then incur a shared $\mathcal{O}(N^3)$ cost for exact GP training.