Reinforcement Learning
with Non-Exponential Discounting

Abstract

Commonly in reinforcement learning (RL), rewards are discounted over time using an exponential function to model time preference, thereby bounding the expected long-term reward. In contrast, in economics and psychology, it has been shown that humans often adopt a hyperbolic discounting scheme, which is optimal when a specific task termination time distribution is assumed. In this work, we propose a theory for continuous-time reinforcement learning generalized to arbitrary discount functions. This formulation covers the case in which there is a random termination time. We derive a Hamilton–Jacobi–Bellman (HJB) equation characterizing the optimal policy and describe how it can be solved using a collocation method, which uses deep learning for function approximation. Further, we show how the inverse RL problem can be approached, in which one tries to recover properties of the discount function given decision data. We validate the applicability of our proposed approach on two simulated problems. Our approach opens the way for the analysis of human discounting in sequential decision-making tasks.

1 Introduction

An often observed phenomenon in humans and animals is that they prefer rewards rather sooner than later [1]. It comes with no surprise that an animal searching for food aims to find it as soon as possible and employees working in a company prefer to be paid after each month instead of after each year. In behavioral experiments, it was observed that people are even willing to pay a price to receive rewards earlier [2]. It can be concluded that rewards become of less value in the future, a phenomenon which is known as discounting [2].

Modelling and experimentally inferring discounting functions, which describe how values are discounted over time, already has a long history in economics and psychology, where many different functional forms have been proposed [3, 4, 5]. While from an economic perspective, a fixed interest rate seems reasonable, leading to an exponential discount function, human behavior is oftentimes better fitted by a hyperbolic curve [6]. The reason for this is that human decisions are typically not consistent regarding shifting rewards in time, a phenomenon named preference reversal: We might find that a subject prefers a smaller reward on the same day over a larger reward one day after. However, when given the choice between the same smaller reward in 365 days, compared to the same larger reward in 366 days, the subject is more likely to be willing to wait the one day more for the larger reward.

While some literature branded the observed discounting behavior as not being rational [7], there has been an increasing amount of work identifying circumstances under which this behavior is indeed

optimal [8, 9, 10]. One widely-adopted theory rationalizing hyperbolic discounting is to assume a constant risk for the reward to become unavailable with an uncertainty about the risk [11]. Under this condition, one should adapt the preference over time, as with time the expected risk decreases.

Discounting is also widely applied in the field of reinforcement learning (RL) and optimal control [12, 13, 14]. First, when modelling infinite horizon time objectives, a discount function is needed to make the expected long-term reward objective well-defined, as otherwise it would become infinite. Second, for autonomous agents, it also makes sense to model a preference for earlier rewards in order to find shortest paths to save time and energy. Third, the discount function can be interpreted as the probability of termination inducing a specific end time distribution [12].

Despite these obvious connections, discounting models of psychology and reinforcement learning have remained mostly independent except for few exceptions [15, 16]. Generalizing reinforcement learning to a broader range of discount functions would enable solutions for applications in which the end time follows a specific distribution. On the other hand, methods for determining optimal decisions in sequential decision making tasks with general end time distributions would provide tools which can help to explain human decision making under uncertainty.

In this work, we present a theory for reinforcement learning in continuous time based on non-exponential discount functions. First, we investigate the conditions under which the objective of maximizing the long-term reward formulated with hyperbolic discounting is well-defined. Then, we derive a Hamilton–Jacobi–Bellman (HJB)-type equation for a general discount function and describe how to solve it to obtain the optimal policy. Further, we provide an approach to tackle the inverse reinforcement learning (IRL) problem, to estimate parameters of the discount function given decision data. Finally, we show the applicability of our proposed method on two simulated problems.

2 Related work

Optimal control in continuous time and space has a long history with many classical works [17, 18, 19, 20]. Continuous-time reinforcement learning formulations have been regarded [14] and various solution methods have been proposed [21, 22]. Solution approaches that solve the HJB equation directly, include linearization techniques [23, 24], path integral formulations [25, 12], and collocation-based methods [26]. In the recent years, it became increasingly popular to use neural networks as a function approximation to solve the HJB equation [27, 28, 29, 30, 31].

Non-exponential discounting was considered in literature in economics, psychology, and neuroscience [4, 32, 2]. For humans and animals, preference reversal behavior was found [33, 34, 5] and different functional forms for the discount function have been proposed [3, 35]. In some work, methods were developed to efficiently estimate the discount function for non-sequential decision data [36, 37]. Further, there has been research aiming to find rational explanations for the encountered discounting behavior [8, 9, 10], e.g., by assuming uncertainty about a constant hazard [11].

Decision processes with non-exponential time distributions have been considered in the field of semi-Markov Decision Processes (SMDPs) [38, 39, 40, 41], where transition times between discrete states can follow arbitrary distributions. MDPs with quasi-hyperbolic discounting, for which all future rewards are additionally discounted by a constant factor, have been addressed in [42, 43, 44]. They can be used to model preference reversal but are limited to the specific form assumed.

More closely related to our work, Fedus et al. [16] presented an approximate method for solving MDPs with hyperbolic discounting. They approached the problem by solving the corresponding exponentially-discounted problem for many different discount factors and combining the results. Part of the approximation, however, is that the value function and policy are static, failing to model preference reversals over time. Another line of work considers MDPs with discount factors that are coupled to the value function to imitate hyperbolic discounting behavior [15, 45]. As time is only considered indirectly through the magnitude of the value function, these approaches cannot be used for finding the solution to a given specific discount function.

Inverse reinforcement learning approaches have been mainly used to learn reward functions given data [46, 47, 48, 49]. These methods were also applied to learning properties of human behavior [50, 51, 52]. Other inverse approaches focused on learning dynamics models [53] or learning rules [54] from sequential decision-making data.

3 Background

3.1 Survival analysis

In survival analysis [55], one is interested in the duration until events occur. In its classical form one considers a single event, for which the duration can be described as a continuous random variable T with cumulative distribution function $F(t) = P(T \leq t)$ and probability density function $f(t)$, where $t \in \mathbb{R}_0^+$ denotes the elapsed time. In survival analysis literature, $F(t)$ is known as the failure function and the survival function is defined as $S(t) = 1 - F(t) = P(T > t)$. The survival function is monotonically decreasing and has the properties $S(0) = 1$ and $\lim_{t \to \infty} S(t) = 0$. By the conditioning rule, one finds $P(T > t_1 \mid T > t_0) = S(t_1)/S(t_0)$ if $t_1 > t_0$. The hazard rate is defined as $\alpha(t) = \lim_{\Delta t \to 0} \frac{1}{\Delta t} P(t \leq T < t + \Delta t \mid t \leq T)$, yielding the relations

$$\alpha(t) = \lim_{\Delta t \to 0} \frac{1}{\Delta t} \frac{S(t) - S(t + \Delta t)}{S(t)} = -\frac{S'(t)}{S(t)} \quad \text{and} \quad S(t) = \exp\left(-\int_0^t \alpha(\tau) \, d\tau\right). \quad (1)$$

For a constant hazard rate $\alpha(t) = \lambda$, one finds $S(t) = \exp(-\lambda t)$, which can be shown to be the unique memory-less survival function [55], i.e., $P(T > t + \Delta t \mid T > t) = P(T > \Delta t)$ for all $t, \Delta t \in \mathbb{R}_0^+$.

3.2 Discounting and preference reversal

We consider the setting in which a subject collects rewards and shows a form of time preference, i.e., rewards are desired rather sooner than later. We model the value of a reward $r \in \mathbb{R}$ as a function $L : \mathbb{R} \times \mathbb{R}_0^+ \to \mathbb{R}$ with $L(r,t) = S(t) \cdot r$, where $S(t)$ is the discount function decreasing with time $t \in \mathbb{R}_0^+$. With the convention that $S(t_0) = 1$ and $\lim_{t \to \infty} S(t) = 0$, we can regard $S(t)$ as the survival function (Section 3.1), with the interpretation that the reward becomes unavailable with hazard rate $\alpha(t)$ [11].

When assuming a constant hazard rate $\alpha(t) = \lambda$, we say that the subject discounts exponentially, as $S(t) = \exp(-\lambda t)$. By the memory-less property, we have that if the subject prefers reward r_1 after t_1 over r_2 after t_2, she or he would remain consistent with the election if we presented the choice again later in time. On the other hand, if we assume a constant but unknown hazard rate λ with belief $p(\lambda) = \mathrm{Gamma}(\lambda; \alpha_0, \beta_0)$, we obtain a hyperbolic form for the expected survival function:

$$S(t; \alpha_0, \beta_0) = \int_\lambda \exp(-\lambda t) \, p(\lambda) \, d\lambda = \frac{1}{(\frac{t}{\beta_0} + 1)^{\alpha_0}} \quad (2)$$

The posterior belief over λ at a later point in time can be derived using Bayes rule and is given by $p(\lambda \mid t) = \mathrm{Gamma}(\lambda; \alpha_0, \beta_0 + t)$. The expected hazard rate $\alpha(t)$ is given by the posterior mean,

$$\alpha(t) = \int_\lambda \lambda \, p(\lambda \mid t) \, d\lambda = \frac{\alpha_0}{\beta_0 + t}. \quad (3)$$

For discount functions other than the exponential, such as the hyperbolic discount function in Eq. (2), the hazard rate varies over time and preferences among options may change.

3.3 Optimal control

In stochastic optimal control [56], we consider a Markovian system with continuous state $\mathbf{x}(t) \in \mathbb{R}^n$ evolving according to the stochastic differential equation (SDE) $d\mathbf{X}(t) = f(\mathbf{X}(t), \mathbf{u}(t)) \, dt + G(\mathbf{X}(t), \mathbf{u}(t)) \, d\mathbf{W}(t)$ where $t \in \mathbb{R}_0^+$ denotes time, $f : \mathcal{X} \times \mathcal{U} \to \mathcal{X}$ the drift function, $G : \mathcal{X} \times \mathcal{U} \to \mathcal{X} \times \mathbb{R}^m$ the dispersion matrix, and $\mathbf{W}(t) \in \mathbb{R}^m$ m-dimensional Brownian motion. The goal of

optimal control is to determine the control inputs $\mathbf{u}(t) \in \mathcal{U}$ given the current state $\mathbf{x}$ at time t, in order to maximize the expected long term discounted reward with reward function $R : \mathcal{X} \times \mathcal{U} \times \mathbb{R}_0^+ \to \mathbb{R}$. The solution is characterized by the optimal value function,

$$V^*(\mathbf{x}) = \max_{\mathbf{u}_{[t,\infty)}} \mathbb{E}\left[\int_t^\infty \exp(-\lambda(\tau - t))\, R(\mathbf{X}(\tau), \mathbf{u}(\tau))\, \mathrm{d}\tau \,\Big|\, \mathbf{X}(t) = \mathbf{x}\right], \tag{4}$$

where maximization is carried out over all trajectories $\mathbf{u}_{[t,\infty)} := \{\mathbf{u}(\tau)\}_{\tau \in [t,\infty]}$. The quantity $\exp(-\lambda t)$ is the discount factor, which ensures convergence of the integral and models a preference for earlier rewards. λ can be interpreted in terms of survival analysis as the hazard rate for termination (cf. Section 3.1). According to the principle of optimality, the stochastic Hamilton-Jacobi-Bellman (HJB) equation, given by

$$\lambda V^*(\mathbf{x}) = \max_{\mathbf{u} \in \mathcal{U}} \left[R(\mathbf{x}, u) + V_\mathbf{x}^*(\mathbf{x})^T f(\mathbf{x}, \mathbf{u}) + \frac{1}{2} V_{\mathbf{xx}}^*(\mathbf{x}) G(\mathbf{x}, \mathbf{u}) G(\mathbf{x}, \mathbf{u})^T \right],$$

provides a condition for the optimal value function. Here, partial derivatives are denoted by the index notation, i.e., $V_\mathbf{x}$ denotes the partial derivative of V w.r.t. $\mathbf{x}$ and $V_{\mathbf{xx}}$ the respective Hessian. Note that the optimal value function depends on time only indirectly through the state $\mathbf{x}$ due to the Markov property and the memory-less discount function. This dependence also applies for the optimal policy $\pi^* : \mathcal{X} \to \mathcal{U}$, given by the maximizer of the right-hand side of the HJB equation.

4 Reinforcement learning with general discount function

We consider a system as in stochastic optimal control (cf. Section 3.3) with continuous state space $\mathbb{R}^n$ and finite set of controls $\mathcal{U}$. Instead of an exponential discount function, we allow for general survival functions $S(t)$ based on a time-dependent hazard rate $\alpha(t)$. The objective function measuring the total expected discounted reward is given by

$$J\left(\mathbf{u}_{[0,\infty)}\right) = \mathbb{E}\left[\int_0^\infty S(\tau)\, R(\mathbf{X}(\tau), \mathbf{u}(\tau))\, \mathrm{d}\tau\right]. \tag{5}$$

Analogous to Section 3.3, we define the expected reward-to-go as the value function

$$V^*(\mathbf{x}, t) = \max_{\mathbf{u}_{[t,\infty)}} \mathbb{E}\left[\int_t^\infty \frac{S(\tau)}{S(t)} R(\mathbf{X}(\tau), \mathbf{u}(\tau))\, \mathrm{d}\tau \,\Big|\, \mathbf{X}(t) = \mathbf{x}\right], \tag{6}$$

where $S(\tau)/S(t)$ is the probability of survival until time τ, conditioned on the fact that one already has survived until time t (cf. Section 3.1). In contrast to Eq. (4), the value function becomes time-dependent through the general survival function and also the optimal policy depends on time.

4.1 Convergence of the value function for hyperbolic discounting

The objective as stated in Eq. (5) is not well-defined for a general model and survival function, as the integral may diverge. For the case of a hyperbolic discount function as in Eq. (2), we find the following theorem:

Theorem 1. *If the reward function $R(\mathbf{x}, \mathbf{u})$ is bounded above for all $(\mathbf{x}, \mathbf{u}) \in \mathcal{X} \times \mathcal{U}$, and $\alpha_0 > 1$, the value function defined in equation Eq. (6) is well-defined. If $R(\mathbf{x}, \mathbf{u})$ is bounded below for all $(\mathbf{x}, \mathbf{u}) \in \mathcal{X} \times \mathcal{U}$, and $\alpha_0 \leq 1$, the value function diverges.*

Proof. See Appendix A.

In the examples considered later, we will assume a bounded reward function and a hyperbolic discount function with $\alpha_0 > 1$, for which Eq. (5) and Eq. (6) are well-defined.

4.2 HJB equation for a general discount function

In the following, we give a brief overview about the derivation of the HJB equation for a general discount function. A more detailed derivation is provided in Appendix B. First, we split the integral in Eq. (6) into two terms such that we obtain a recursive formulation of the value function:

$$
V^*(\mathbf{x}, t) = \max_{\mathbf{u}_{[t, t+\Delta t]}} \mathbb{E} \left[\int_t^{t+\Delta t} \frac{S(\tau)}{S(t)} R(\mathbf{X}(\tau), \mathbf{u}(\tau), \tau) \, \mathrm{d}\tau \right.
$$
$$
\left. + \frac{S(t + \Delta t)}{S(t)} V^*(\mathbf{X}(t + \Delta t), t + \Delta t) \, \Big| \, \mathbf{X}(t) = \mathbf{x} \right]
$$

For the second term in the expectation, we apply a Taylor expansion and Itô's formula [56] and obtain

$$
V^*(\mathbf{X}(t + \Delta t), t + \Delta t) = V^*(\mathbf{X}(t), t) + \int_t^{t+\Delta t} V_{\mathbf{x}}^*(\mathbf{X}(\tau), \tau) \, f(\mathbf{X}(\tau), \mathbf{u}(\tau), \tau) \, \mathrm{d}\tau
$$
$$
+ \int_t^{t+\Delta t} V_{\mathbf{x}}^*(\mathbf{X}(\tau), \tau) \, G(\mathbf{X}(\tau), \mathbf{u}(\tau), \tau) \, \mathrm{d}\mathbf{W}(\tau) + \int_t^{t+\Delta t} V_t^*(\mathbf{X}(\tau), \tau) \, \mathrm{d}\tau
$$
$$
+ \int_t^{t+\Delta t} \frac{1}{2} \, \mathrm{tr} \left\{ V_{\mathbf{xx}}^*(\mathbf{X}(\tau), \tau) \, G(\mathbf{X}(\tau), \mathbf{u}(\tau), \tau) \, G(\mathbf{X}(\tau), \mathbf{u}(\tau), \tau)^T \right\} \mathrm{d}\tau + o(\Delta t).
$$

Plugging in this result into the equation above, dividing both sides by Δt, and taking the limit $\Delta t \to 0$, as well as calculating the expectation w.r.t. $\mathbf{W}(t)$ leads to the desired HJB equation

$$
\alpha(t) V^*(\mathbf{x}, t) = \max_{\mathbf{u}} \left\{ R(\mathbf{x}, \mathbf{u}, t) + V_t^*(\mathbf{x}, t) + V_{\mathbf{x}}^*(\mathbf{x}, t) \, f(\mathbf{x}, \mathbf{u}, t) \right.
$$
$$
\left. + \frac{1}{2} \, \mathrm{tr} \left(V_{\mathbf{xx}}^*(\mathbf{x}, t) \, G(\mathbf{x}, \mathbf{u}, t) \, G(\mathbf{x}, \mathbf{u}, t)^T \right) \right\},
\tag{7}
$$

where $\alpha(t)$ can be recognized to be the hazard rate corresponding to the survival function $S(t)$. We define the r.h.s. of the HJB equation in Eq. (7) without the maximization w.r.t. the action as

$$
Q(\mathbf{x}, \mathbf{u}, t) := R(\mathbf{x}, \mathbf{u}, t) + V_t^*(\mathbf{x}, t) + V_{\mathbf{x}}^*(\mathbf{x}, t) \, f(\mathbf{x}, \mathbf{u}, t)
$$
$$
+ \frac{1}{2} \, \mathrm{tr} \left(V_{\mathbf{xx}}^*(\mathbf{x}, t) \, G(\mathbf{x}, \mathbf{u}, t) \, G(\mathbf{x}, \mathbf{u}, t)^T \right),
$$

so that the optimal policy is given by $\pi^*(\mathbf{x}, t) = \arg\max_{\mathbf{u}} Q(\mathbf{x}, \mathbf{u}, t)$. Later on, we will consider hyperbolic discounting, for which the hazard rate $\alpha(t)$ is given by Eq. (3), i.e., $\alpha(t) = \alpha_0 / (\beta_0 + t)$.

4.3 Solving the HJB equation

In order to solve the HJB equation in Eq. (7), which is a PDE, we apply a collocation-based method [26, 27, 29]. To do so, we first reformulate the HJB equation as

$$
E(V, \mathbf{x}, t) := -\alpha(t) V(\mathbf{x}, t) + \max_u \left[R(\mathbf{x}, \mathbf{u}, t) + V_t(\mathbf{x}, t) + V_{\mathbf{x}}(\mathbf{x}, t) \, f(\mathbf{x}, \mathbf{u}, t) \right.
$$
$$
\left. + \frac{1}{2} \, \mathrm{tr} \left(V_{\mathbf{xx}}(\mathbf{x}, t) \, G(\mathbf{x}, \mathbf{u}, t) \, G(\mathbf{x}, \mathbf{u}, t)^T \right) \right] = 0,
\tag{8}
$$

and use a function approximator $V^\psi(\mathbf{x}, t)$ for $V^*(\mathbf{x}, t)$. The parameters ψ of the approximator can be determined by sampling random states $\hat{\mathbf{x}}_i$ and time points $\hat{t}_i$ and minimizing $\sum_i E(V^\psi, \hat{\mathbf{x}}_i, \hat{t}_i)^2$ w.r.t. ψ. Calculating the derivatives $V_{\mathbf{x}}^\psi(\mathbf{x}, t)$, $V_t^\psi(\mathbf{x}, t)$, $V_{\mathbf{xx}}^\psi(\mathbf{x}, t)$ and differentiating the objective function w.r.t. ψ is straightforward via automatic differentiation when choosing a neural network as function approximator [57]. As t is not bounded, we need to choose a reparametrization of t which maps all t to the interval $[0, 1)$ before feeding the values into the network. More details about the implemented parametrization and application of the collocation method is provided in Appendix C. The complete algorithm to learn the value function and policy is provided in Algorithm 1.

Algorithm 1: Computation of the optimal value function and policy for non-exp. discounting

---

Result: Optimal value function $V^\psi(\mathbf{x}, t)$, optimal policy $\pi^\psi(\mathbf{x}, t)$
Input: Parameters $\boldsymbol{\theta}$ of the discount function, system model, number of iterations K
Initialize parameters ψ of neural network for modelling $V^\psi(\mathbf{x}, t)$;
for $k \leftarrow 0$ **to** $K - 1$ **do**
    Sample a batch of states and time points $(\hat{\mathbf{x}}, \hat{t})_{i=\{1,...,N\}}$;
    Push $(\hat{\mathbf{x}}, \hat{t})_{i=\{1,...,N\}}$ through the network to obtain $\hat{V}^\psi_{i=\{1,...,N\}}$;
    Use back-propagation to compute $\hat{V}^\psi_{\mathbf{x}i}, \hat{V}^\psi_{ti}, \hat{V}^\psi_{\mathbf{xx}i}$;
    Evaluate $E(\hat{V}^\psi_i, \hat{\mathbf{x}}_i, \hat{t}_i)$ in Eq. (8) and determine maximizing action $\hat{\mathbf{u}}_i$ for $i = 1, \ldots, N$;
    Use back-propagation to compute the gradient of $\sum_i E(\hat{V}^\psi_i, \hat{\mathbf{x}}_i, \hat{t}_i)^2$ w.r.t. ψ ;
    Update ψ using the gradient ;
end
return $V^\psi(\mathbf{x}, t), \pi^\psi(\mathbf{x}, t) = \arg\max_{\mathbf{u}} Q(\mathbf{x}, \mathbf{u}, t)$

4.4 Inverse reinforcement learning for inferring the discount function

When analysing human behavior, one might be interested in learning the underlying discount function that led to some behavioral data. In contrast to standard inverse reinforcement learning settings, where the goal is to learn the underlying reward function R, here we assume that the reward function is given and the discount function $S(t)$ is unknown and needs to be inferred.

As given data, we assume the states and time points at which the subject switches from one action to another one, i.e., $\mathcal{D} = \{\mathbf{x}, \mathbf{u}^-, \mathbf{u}^+, t\}_{i=1...N}$, describing that in state $\mathbf{x}$ at time t the action u^- is switched to u^+. The observed decision maker is assumed to use the optimal policy $\pi(\mathbf{x}, t) = \arg\max_{\mathbf{u}} Q(\mathbf{x}, \mathbf{u}, t)$. Shortly before and after switching time t, we have $Q(\mathbf{x}(t-\Delta t), \mathbf{u}^-, t-\Delta t) > Q(\mathbf{x}(t-\Delta t), \mathbf{u}^+, t-\Delta t)$ and $Q(\mathbf{x}(t+\Delta t), \mathbf{u}^-, t+\Delta t) < Q(\mathbf{x}(t+\Delta t), \mathbf{u}^+, t+\Delta t)$, respectively, indicating that before t action $\mathbf{u}^-$ is preferred over $\mathbf{u}^+$ and vice versa afterwards. By letting $\Delta t \to 0$, one finds $Q(\mathbf{x}(t), \mathbf{u}^-, t) = Q(\mathbf{x}(t), \mathbf{u}^+, t)$. A sensible objective for the inverse problem is therefore to minimize $\sum_i F(\mathbf{x}_i, \mathbf{u}_i^-, \mathbf{u}_i^+, t_i)^2$, defined as the squared difference of both Q values, i.e.,

$$
\begin{aligned}
F(\mathbf{x}, \mathbf{u}^-, \mathbf{u}^+, t) &= \left\{ Q(\mathbf{x}, \mathbf{u}^-, t) - Q(\mathbf{x}, \mathbf{u}^+, t_i) \right\}^2 \\
&= \big\{ R(\mathbf{x}, \mathbf{u}^+) - R(\mathbf{x}, \mathbf{u}^-) + V_{\mathbf{x}}(\mathbf{x}, t) \left(f(\mathbf{x}, \mathbf{u}^+) - f(\mathbf{x}, \mathbf{u}^-) \right) \\
&\quad + \frac{1}{2} \operatorname{tr} \left(V_{\mathbf{xx}}(\mathbf{x}, t) \left(G(\mathbf{x}, \mathbf{u}^+, t) G(\mathbf{x}, \mathbf{u}^+, t)^T - G(\mathbf{x}, \mathbf{u}^-, t) G(\mathbf{x}, \mathbf{u}^-, t)^T \right) \right) \big\}^2,
\end{aligned}
\tag{9}
$$

w.r.t. the parameters $\boldsymbol{\theta}$ of the discount function. The optimization needs to take Eq. (8) as a constraint into account to ensure that the HJB equation is fulfilled. Note that Eq. (9) depends indirectly on the parameters $\boldsymbol{\theta}$ through the definition of the value function. However, the objective function F could also directly depend on $\boldsymbol{\theta}$, we consider the general case in the following. In principle, the state at which the switching occurs, also depends on $\boldsymbol{\theta}$. Nevertheless, we will assume that the parameters have only minor influence on the states of switching, so that terms including the variation of $\mathbf{x}$ w.r.t. $\boldsymbol{\theta}$ can be neglected.

Gradient computation For the minimization, it is useful to determine the gradient of the objective function F w.r.t. $\boldsymbol{\theta}$, which can be formulated using the chain rule for total derivatives as

$$
\begin{aligned}
\frac{\mathrm{d}F}{\mathrm{d}\boldsymbol{\theta}} &= F_{\boldsymbol{\theta}} + F_V V_{\boldsymbol{\theta}} + F_{V_{\mathbf{x}}} (V_{\mathbf{x}})_{\boldsymbol{\theta}} + F_{V_{\mathbf{xx}}} (V_{\mathbf{xx}})_{\boldsymbol{\theta}} \\
&= F_{\boldsymbol{\theta}} + F_V V_{\boldsymbol{\theta}} + F_{V_{\mathbf{x}}} (V_{\boldsymbol{\theta}})_{\mathbf{x}} + F_{V_{\mathbf{xx}}} (V_{\boldsymbol{\theta}})_{\mathbf{xx}}.
\end{aligned}
\tag{10}
$$

By switching the order of the partial derivatives in the last step, we have obtained an expression for the gradient depending on $V_{\boldsymbol{\theta}}$. The partial derivative $V_{\boldsymbol{\theta}}$ measures the influence of the discounting parameters on the value function and is not straightforward to evaluate. We can get a PDE for this quantity by observing that the constraint $E(V^*, \mathbf{x}, t)$ in Eq. (8) is zero everywhere for the optimal

Figure 1: **Results for the investment problem. A** Learned value function for account balances and time points for hyperbolic discounting. **B** Learned Q function for both actions over time (median and quantiles for 10 runs). **C** Learned policy showing preference reversals for all account balances except for $b = 1$. **D** Simulation with account balance (top), interest rate (middle), and action (bottom) over time. **E** Learned value function for an exponential discount function, which is constant over time. **G** Learned Q function for an exponential discount function. **G** Gradients obtained for parameter pairs $[\alpha_0, \beta_0]$ of the discount function given simulated data on a 11×11 grid. The parameters used to generate the simulated trajectories are indicated by a red cross. **H** Values of the objective function F (Eq. (9)) for parameter pairs on a 11×11 grid when applying the IRL method.

value function V^* and therefore its derivative also needs to be zero. This method is known as the forward sensitivity method [58] and provides an additional PDE for the desired gradient $V_{\boldsymbol{\theta}}$:

$$
\begin{aligned}
0 = \frac{\mathrm{d}E}{\mathrm{d}\boldsymbol{\theta}} &= E_V V_{\boldsymbol{\theta}} + E_{V_{\mathbf{x}}} (V_{\mathbf{x}})_{\boldsymbol{\theta}} + E_{V_{\mathbf{xx}}} (V_{\mathbf{xx}})_{\boldsymbol{\theta}} \\
&= E_V V_{\boldsymbol{\theta}} + E_{V_{\mathbf{x}}} (V_{\boldsymbol{\theta}})_{\mathbf{x}} + E_{V_{\mathbf{xx}}} (V_{\boldsymbol{\theta}})_{\mathbf{xx}} =: H(V, V_{\boldsymbol{\theta}}, \mathbf{x}, t)
\end{aligned}
\tag{11}
$$

To solve the PDE and obtain $V_{\boldsymbol{\theta}}$, we can use the same procedure as in Section 4.3. To obtain the gradient, first one has to solve the HJB equation Eq. (7) to obtain V^ψ, then Eq. (11) to determine $V_{\boldsymbol{\theta}}^\phi$. Afterwards, Eq. (10) gives the gradient to update the parameters $\boldsymbol{\theta}$ of the discount function. The quantities $V_{\boldsymbol{\theta}}, (V_{\boldsymbol{\theta}})_{\mathbf{x}}, (V_{\boldsymbol{\theta}})_{\mathbf{xx}}$ for the derivative be computed via automatic differentiation as in Section 4.3. The complete algorithm for computing the gradient is presented as Algorithm 2 in Appendix D.

5 Experiments

We tested our derived method on two simulated problems with a random termination time following a hyperbolic survival function. First, we solved the HJB equation in Eq. (7) using the collocation method (Algorithm 1) and computed the optimal policy. Then, for applying the inverse reinforcement learning method, we evaluated the objective function in Eq. (9) and computed gradients for different parameter sets $\boldsymbol{\theta} = [\alpha_0, \beta_0]$ of the hyperbolic discount function in Eq. (2) using Algorithm 2. For generating data, we randomly sampled starting states and determined the time points at which subjects would switch their action. Afterwards, the determined time points were distorted by Gaussian noise.

All proposed methods were implemented in Python using the PyTorch [57] framework. The used hyperparameters are presented in Appendix F. As tasks, we considered an investment problem and a problem to control a point on a line. In the following we provide a brief overview over the considered problems, more details can be found in Appendix E.

Figure 2: **Results for the line problem. A** Learned value function for hyperbolic discount function. **B** Learned Q function for each action over time (median and quantiles for 10 runs) **C** Learned policy showing preference reversals. **D** Simulation showing the state (top) and action (bottom) over time. **E** Learned value function for an exponential discount function. **G** Learned Q function for an exponential discount function. **G** Gradients obtained for parameter pairs $[\alpha_0, \beta_0]$ on a 15×15 grid given simulated data. The parameter used to generate the simulated trajectories is indicated by a red cross. **H** Values of the objective function of the IRL methods for parameter pairs on a 15×15 grid.

5.1   Experimental Tasks

In the investment problem, we model a subject having to decide whether to invest her or his income to the bank account leading to future interests (rewards), or to spend the money for immediate reward. We model the state as the current balance of the bank account as well as the current interest rate. When the money is spent (action *spend*), the subject receives rewards with rate 0.1 but the balance of the account remains unchanged. When the income is invested (action *invest*), the balance of the bank account increases with rate 0.1 but there is no additional reward. In both cases the subject receives rewards through interests, being proportional to the current balance on the bank account. We assume that the interest rate varies over time following a Gaussian diffusion model. To keep the state bounded, we model a maximum balance for the account.

In the line problem, the task is to control a point along a line. The state consists of the current position of the point. Possible actions are *left* and *right*, which move the point to the respective direction, as well as *stay*, which does not move the point at all. When moving the point, there is a Gaussian diffusion on the position and one has to pay a small action cost (negative reward). We model a state dependent reward modelling a high reward for distant states on one side and low reward for close states on the other side. A more detailed description of the functional form is provided in Appendix E.

5.2   Results

The proposed method in Algorithm 1 produces plausible value functions for the considered problems. Figure 1 A shows the learned value function for the investment problem. The values are increasing with account balance (b), as expected rewards through interests become higher. Further in time, the value also increases, as the hazard rate can be assumed to be lower and one is expected to collect rewards for a longer time. Figure 1 B and C shows the learned Q function and policy, respectively. While it is advantageous to spend the income in the beginning, preference reversal occurs when the risk of termination is assumed to relatively low, and investing becomes more attractive. The simulation in Fig. 1 D also reflects this behavior. For comparison, when assuming exponential discounting (Fig. 1 E and F), the value and Q function do not show preference reversal and remain constant over time. With regard to the inverse approach, Fig. 1 G shows the computed gradients of the IRL objective function F in Eq. (9) for different parameter pairs. One can observe that the

gradients mostly point into the direction of the parameters used for simulation, indicated by a red cross. Figure 1 H shows the evaluated IRL objective function F for the parameter pairs, correctly displaying low values in the area close to the true parameters.

For the line problem, the computed value function is depicted in Fig. 2 A. Also in this task, the learned value increases over time and is high in areas close to highly rewarding states. Figure 2 B and C shows the learned Q function and policy respectively. In the beginning when the risk could be potentially high, if not close enough to the large reward, it is best to move right to collect the smaller sooner reward. With time when the risk of termination decreases, it becomes increasingly advantageous to move left to collect the larger later reward. In between, there is a time span, during which it is best to stay: The moving cost is too high for moving right and shortly turning afterwards, but the risk is not low enough for being worth it to move left to the higher reward. The simulation in Fig. 2 D presents a sampled trajectory, showing that the subject first moves right until the maximum smaller sooner reward at $x = 0.5$ is reached and stays there for some time. After the risk has decreased over time, the simulated subject finally moves right to collect the larger later reward – her or his preferences have reversed. Like for the investment problem, the value and Q function (Fig. 2 E and F) do not model any time-varying behavior. With regard to the IRL problem, the gradients and function values of F of Eq. (9) in Fig. 2 G and H also lead closely to parameter values used for the simulations.

6 Conclusion

In this work, we have proposed a method for reinforcement learning with non-exponential discount functions. The approach can be used to solve decision-making problems with an arbitrary end-time distribution and to model human discounting behavior. First, we have shown the conditions for which the problem is well-defined when using a hyperbolic discount function. Then, we derived a HJB-type equation providing conditions for the optimal time-dependent value function. We presented how the obtained PDE can be approximately solved using a collocation method, leading to the optimal policy. Further, we introduced an approach for the inverse problem, in which the discount function needs to be inferred given behavioral data. The application of our methods on two simulated problems led to plausible solutions, opening the way for further applications such as the use in human experiments.

Limitations and future work   In our proposed methods, we assume a finite action space. While this assumption applies to many behavioral experiments, it can be a limitation for the application to classical optimal control. To extend the method to continuous control, one has to determine how the maximization problem in the HJB equation is solved. For strictly convex action costs, there have been approaches proposed that efficiently solve the maximization in the HJB equation [29], under some conditions even in closed-form [27]. Another requirement of our method is that the model needs to be known. Model-free approaches, such as TD-error learning for general discount functions could be explored, in line with work such as [15]. Further, when approaching problems with large state spaces, the proposed collocation method is likely to converge slowly. For these cases, adapted collocation methods [28] or advantage updating approaches [22] could be considered instead. To apply our method to more advanced problems in financial engineering, such as modeling stocks with discontinuous returns [59], one has to consider general jump-diffusion processes. While an extension to these models is straightforward from a theoretical point, we here left it out to keep the focus on the handling of the discount function. Regarding our inverse reinforcement learning approach, we have assumed the states and time points of the action switches to be given. While this assumption is reasonable for many human behavior experiments, it might be interesting to learn discount functions for given discretized trajectories instead. Also, incorporating an extended timing model of human decisions [60] instead of fixed-variance Gaussian diffusion would be an interesting extension.

In the future, we are planning to apply our proposed methods in human experiments to get new insights to human discounting behavior. Characterizing individual human subjects by analyzing their behavior comes with the risk to be used with negative social impact. This matter can be counteracted by collecting only anonymized data for the application of our method.

References

[1] Ted O'Donoghue and Matthew Rabin. The economics of immediate gratification. *Journal of Behavioral Decision Making*, 13(2):233–250, 2000.

[2] Shane Frederick, George Loewenstein, and Ted O'donoghue. Time discounting and time preference: A critical review. *Journal of Economic Literature*, 40(2):351–401, 2002.

[3] Todd L McKerchar, Leonard Green, Joel Myerson, T Stephen Pickford, Jade C Hill, and Steven C Stout. A comparison of four models of delay discounting in humans. *Behavioural Processes*, 81(2):256–259, 2009.

[4] Robert Henry Strotz. Myopia and inconsistency in dynamic utility maximization. *The Review of Economic Studies*, 23(3):165–180, 1955.

[5] Richard Thaler. Some empirical evidence on dynamic inconsistency. *Economics Letters*, 8(3):201–207, 1981.

[6] James E Mazur. An adjusting procedure for studying delayed reinforcement. *Quantitative Analyses of Behavior*, 5:55–73, 1987.

[7] Irving Fisher. *Theory of interest: as determined by impatience to spend income and opportunity to invest it*. Augustusm Kelly Publishers, 1930.

[8] Partha Dasgupta and Eric Maskin. Uncertainty and hyperbolic discounting. *American Economic Review*, 95(4):1290–1299, 2005.

[9] Taiki Takahashi. Loss of self-control in intertemporal choice may be attributable to logarithmic time-perception. *Medical Hypotheses*, 65(4):691–693, 2005.

[10] Debajyoti Ray and Peter Bossaerts. Positive temporal dependence of the biological clock implies hyperbolic discounting. *Frontiers in Neuroscience*, 5:2, 2011.

[11] Peter D Sozou. On hyperbolic discounting and uncertain hazard rates. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 265(1409):2015–2020, 1998.

[12] Richard S Sutton and Andrew G Barto. *Reinforcement learning: An introduction*. MIT Press, 2018.

[13] Dimitri Bertsekas. *Reinforcement learning and optimal control*. Athena Scientific, 2019.

[14] Kenji Doya. Reinforcement learning in continuous time and space. *Neural Computation*, 12(1):219–245, 2000.

[15] William H Alexander and Joshua W Brown. Hyperbolically discounted temporal difference learning. *Neural Computation*, 22(6):1511–1527, 2010.

[16] William Fedus, Carles Gelada, Yoshua Bengio, Marc G Bellemare, and Hugo Larochelle. Hyperbolic discounting and learning over multiple horizons. *arXiv preprint arXiv:1902.06865*, 2019.

[17] Ruslan L Stratonovich. Conditional markov processes and their application to the theory of optimal control. 1968.

[18] Harold J Kushner and Paul G Dupuis. *Numerical methods for stochastic control problems in continuous time*, volume 24. Springer Science & Business Media, 2001.

[19] Wendell H Fleming and Halil Mete Soner. *Controlled Markov processes and viscosity solutions*, volume 25. Springer Science & Business Media, 2006.

[20] Lev Semenovich Pontryagin. *Mathematical theory of optimal processes*. CRC Press, 1987.

[21] Kyriakos G Vamvoudakis and Frank L Lewis. Online actor–critic algorithm to solve the continuous-time infinite horizon optimal control problem. *Automatica*, 46(5):878–888, 2010.

[22] Leemon C Baird. Reinforcement learning in continuous time: Advantage updating. In *IEEE International Conference on Neural Networks*, volume 4, pages 2448–2453. IEEE, 1994.

[23] David H Jacobson. New second-order and first-order algorithms for determining optimal control: A differential dynamic programming approach. *Journal of Optimization Theory and Applications*, 2(6): 411–440, 1968.

[24] Yuval Tassa, Tom Erez, and William Smart. Receding horizon differential dynamic programming. In *Advances in Neural Information Processing Systems*, volume 20. Curran Associates, Inc., 2007.

[25] Hilbert J Kappen. Path integrals and symmetry breaking for optimal control theory. *Journal of Statistical Mechanics: Theory and Experiment*, 2005(11):P11011, 2005.

[26] Alex Simpkins and Emanuel Todorov. Practical numerical methods for stochastic optimal control of biological systems in continuous time and space. In *IEEE Symposium on Adaptive Dynamic Programming and Reinforcement Learning*, pages 212–218. IEEE, 2009.

[27] Yuval Tassa and Tom Erez. Least squares solutions of the HJB equation with neural network value-function approximators. *IEEE Transactions on Neural Networks*, 18(4):1031–1041, 2007.

[28] Justin Sirignano and Konstantinos Spiliopoulos. DGM: A deep learning algorithm for solving partial differential equations. *Journal of Computational Physics*, 375:1339–1364, 2018.

[29] Michael Lutter, Boris Belousov, Kim Listmann, Debora Clever, and Jan Peters. HJB optimal feedback control with deep differential value functions and action constraints. In *Conference on Robot Learning*, pages 640–650. PMLR, 2020.

[30] Jiequn Han, Arnulf Jentzen, and E Weinan. Solving high-dimensional partial differential equations using deep learning. *Proceedings of the National Academy of Sciences*, 115(34):8505–8510, 2018.

[31] Bastian Alt, Matthias Schultheis, and Heinz Koeppl. POMDPs in continuous time and discrete spaces. In *Advances in Neural Information Processing Systems*, volume 33, pages 13151–13162. Curran Associates, Inc., 2020.

[32] Richard H Thaler and Hersh M Shefrin. An economic theory of self-control. *Journal of Political Economy*, 89(2):392–406, 1981.

[33] George Ainslie and Richard J Herrnstein. Preference reversal and delayed reinforcement. *Animal Learning & Behavior*, 9(4):476–482, 1981.

[34] Leonard Green, Nathanael Fristoe, and Joel Myerson. Temporal discounting and preference reversals in choice between delayed outcomes. *Psychonomic Bulletin & Review*, 1(3):383–389, 1994.

[35] Steffen Andersen, Glenn W Harrison, Morten I Lau, and E Elisabet Rutström. Discounting behavior: A reconsideration. *European Economic Review*, 71:15–33, 2014.

[36] Daniel R Cavagnaro, Gabriel J Aranovich, Samuel M McClure, Mark A Pitt, and Jay I Myung. On the functional form of temporal discounting: An optimized adaptive test. *Journal of Risk and Uncertainty*, 52 (3):233–254, 2016.

[37] Jorge Chang, Jiseob Kim, Byoung-Tak Zhang, Mark A Pitt, and Jay I Myung. Modeling delay discounting using gaussian process with active learning. In *CogSci*, pages 1479–1485, 2019.

[38] Ronald A Howard. Semi-markovian decision processes. *Bulletin of the International Statistical Institute*, 40(2):625–652, 1963.

[39] Vo S Korolyuk, SM Brodi, and AF Turbin. Semi-markov processes and their applications. *Journal of Soviet Mathematics*, 4(3):244–280, 1975.

[40] Sheldon M Ross. Average cost semi-markov decision processes. *Journal of Applied Probability*, 7(3): 649–656, 1970.

[41] Steven Bradtke and Michael Duff. Reinforcement learning methods for continuous-time markov decision problems. In *Advances in Neural Information Processing Systems*, volume 7. MIT Press, 1994.

[42] Anna Jaśkiewicz and Andrzej S Nowak. Markov decision processes with quasi-hyperbolic discounting. *Finance and Stochastics*, 25(2):189–229, 2021.

[43] Tomas Bjork and Agatha Murgoci. A general theory of markovian time inconsistent stochastic control problems. *Available at SSRN*, 2010.

[44] Tomas Björk and Agatha Murgoci. A theory of markovian time-inconsistent stochastic control in discrete time. *Finance and Stochastics*, 18(3):545–592, 2014.

[45] Hado van Hasselt, John Quan, Matteo Hessel, Zhongwen Xu, Diana Borsa, and André Barreto. General non-linear bellman equations. *arXiv preprint arXiv:1907.03687*, 2019.

[46] Andrew Y Ng, Stuart J Russell, et al. Algorithms for inverse reinforcement learning. In *International Conference on Machine Learning*, volume 1, 2000.

[47] Pieter Abbeel and Andrew Y Ng. Apprenticeship learning via inverse reinforcement learning. In *International Conference on Machine Learning*, volume 5, 2004.

[48] Brian D Ziebart, Andrew L Maas, J Andrew Bagnell, Anind K Dey, et al. Maximum entropy inverse reinforcement learning. In *AAAI*, volume 8, pages 1433–1438, 2008.

[49] Divyansh Garg, Shuvam Chakraborty, Chris Cundy, Jiaming Song, and Stefano Ermon. IQ-Learn: Inverse soft-q learning for imitation. In *Advances in Neural Information Processing Systems*, volume 34, pages 4028–4039. Curran Associates, Inc., 2021.

[50] Katja Mombaur, Anh Truong, and Jean-Paul Laumond. From human to humanoid locomotion—an inverse optimal control approach. *Autonomous Robots*, 28(3):369–383, 2010.

[51] Katharina Muelling, Abdeslam Boularias, Betty Mohler, Bernhard Schölkopf, and Jan Peters. Learning strategies in table tennis using inverse reinforcement learning. *Biological Cybernetics*, 108(5):603–619, 2014.

[52] Matthias Schultheis, Dominik Straub, and Constantin A Rothkopf. Inverse optimal control adapted to the noise characteristics of the human sensorimotor system. In *Advances in Neural Information Processing Systems*, volume 34, pages 9429–9442. Curran Associates, Inc., 2021.

[53] Matthew Golub, Steven Chase, and Byron Yu. Learning an internal dynamics model from control demonstration. In *International Conference on Machine Learning*, pages 606–614. PMLR, 2013.

[54] Zoe Ashwood, Nicholas A. Roy, Ji Hyun Bak, and Jonathan W Pillow. Inferring learning rules from animal decision-making. In *Advances in Neural Information Processing Systems*, volume 33, pages 3442–3453. Curran Associates, Inc., 2020.

[55] Odd Aalen, Ornulf Borgan, and Hakon Gjessing. *Survival and event history analysis: a process point of view*. Springer Science & Business Media, 2008.

[56] Floyd B Hanson. *Applied stochastic processes and control for jump-diffusions: modeling, analysis and computation*. SIAM, 2007.

[57] Adam Paszke, Sam Gross, Francisco Massa, Adam Lerer, James Bradbury, Gregory Chanan, Trevor Killeen, Zeming Lin, Natalia Gimelshein, Luca Antiga, Alban Desmaison, Andreas Kopf, Edward Yang, Zachary DeVito, Martin Raison, Alykhan Tejani, Sasank Chilamkurthy, Benoit Steiner, Lu Fang, Junjie Bai, and Soumith Chintala. Pytorch: An imperative style, high-performance deep learning library. In *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019.

[58] Herschel Rabitz, Mark Kramer, and D Dacol. Sensitivity analysis in chemical kinetics. *Annual Review of Physical Chemistry*, 34(1):419–461, 1983.

[59] Robert C Merton. Option pricing when underlying stock returns are discontinuous. *Journal of Financial Economics*, 3(1-2):125–144, 1976.

[60] Vijay Mohan K Namboodiri, Stefan Mihalas, Tanya Marton, and Marshall Gilmer Hussain Shuler. A general theory of intertemporal decision-making and the perception of time. *Frontiers in Behavioral Neuroscience*, 8:61, 2014.